



UNIVERSITI PUTRA MALAYSIA

***EFFECTS OF *Borreria latifolia* (Aubl.) K.SCHUM. AND *Rosmarinus officinalis*
L. AS FEED ADDITIVES ON PERFORMANCE AND MEAT QUALITY OF
FINISHING VILLAGE CHICKEN***

ALI ABDUL AMEER

FP 2016 63



EFFECTS OF *Borreria latifolia* (Aubl.) K.SCHUM. AND *Rosmarinus officinalis* L. AS FEED ADDITIVES ON PERFORMANCE AND MEAT QUALITY OF FINISHING VILLAGE CHICKEN

By

ALI ABDUL AMEER

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

October 2016



© COPYRIGHT UPM

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATIONS

To My Beloved Mother And Father

And

To My Family With Love And Respect



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

EFFECTS OF *Borreria latifolia* (Aubl.) K.SCHUM. AND *Rosmarinus officinalis* L.AS FEED ADDITIVES ON PERFORMANCE AND MEAT QUALITY OF FINISHING VILLAGE CHICKEN

By

ALI ABDUL AMEER

October 2016

Chairman : Associate Professor Azhar Kasim, PhD
Faculty : Agriculture

The poultry industry would extremely appreciate natural antioxidants that could replace synthetic ones and satisfy consumer demands for food products without residues from substances that can possibly hurt human health. Besides health decay, lipid oxidation is also a major cause of meat quality deterioration, affecting color, flavor, texture, and nutritional value.

The objective of this study was to investigate the impact of dietary supplementation of *Borreria latifolia* and *Rosmarinus officinalis* as natural antioxidant on performance and meat quality of finishing village chicken. Two experiments were conducted, the first experiment was to confirm the antioxidant activity of BL through a comparison with *Rosmarinus officinalis* (RO) as a reference in addition to evaluate their effect on performance and meat quality. Three treatments were employed: T1 (control basal diet without supplementation), T2 (basal diet with 1% of BL), T3 (basal diet with 1% of RO). There was a significant difference ($p < 0.05$) between dietary treatments in cooking loss value. Cooking loss value was higher in T1 compared to the T2 and T3. Significantly ($p < 0.05$), pH value was affected by dietary treatments with T2 and T3 having higher pH value compared to T1. Additionally, significant effect of dietary treatments was recorded during sensory evaluation test whereas T2, T3 had a higher score in overall acceptability compared to control with T2 higher ($p < 0.05$) than T3. Moreover, significant effect ($p < 0.05$) of dietary treatment on lipid oxidation value was observed. T2 and T3 had a lower malondialdehyde (MDA) concentrations compared to T1. Additionally, there was significant ($p < 0.05$) difference between males and female detected in final body weight, weight gain, feed intake, carcass weight, breast and thigh weight, wing weight, neck and loin weight. Moreover, the sex effect was significant ($p < 0.05$) on MUFA, PUFA, PUFA:SFA ratio, n6:n3 ratio, pH value, yellowness (b^*) and lipid oxidation. In addition, significant ($p < 0.05$) increase in tenderness, pH, and lipid oxidation through post mortem aging period (1, 3, 5 d) were recorded, while cooking loss (CL) and redness (a^*) were decreased. Significant interactions between herb and

sex on pH parameter, and herb by sex, herb by post-mortem period, sex by post-mortem period, and herb by sex by post-mortem period interactions in regards with lipid oxidation were observed.

The second experiment was to evaluate the effect of three different levels of BL on meat quality and growth performance. Four treatments applied were: T1 (control basal diet without supplementation), T2 (basal diet with 1.5% of BL), T3 (basal diet with 2% of BL), and T4 (basal diet with 2.5% of BL). There was a significant ($p < 0.05$) effect of the dietary treatments on redness (a^*) value. T4 showed higher value than T1, T2, and T3. Furthermore, T4 produced higher ($p < 0.05$) scores than T1, T2, and T3 in both overall acceptability and texture through sensory evaluation test. With regard to lipid oxidation test, T4 showed the lowest malondialdehyde concentration whereas T1 had the highest concentration of MDA. Meanwhile, T2 and T3 had a higher concentration of MDA compared to T4. Furthermore, there was significant ($p < 0.05$) difference between males and females in MUFA, PUFA, PUFA:SFA ratio, n6:n3 ratio, yellowness (b^*) and lipid oxidation. In addition, significant ($p < 0.05$) effect of sex on final body weight, weight gain, feed intake, carcass weight, breast and thigh weight, wings weight, neck and loin weight, and liver weight was recorded. The obtained result of the current studies showed significant ($p < 0.05$) decreases in redness (a^*) and cooking loss (CL) during post mortem aging period while pH value, tenderness, and lipid oxidation were elevated. Additionally, there were significant interactions between herb and sex, herb and aging period, sex and aging period, and the herb by sex by aging period interaction on lipid oxidation as well as a significant interaction between herb and sex on pH was noticed in the study.

Generally, the present study revealed that dietary supplementation of 2.5% BL improved oxidative stability which could have extended meat shelf life and may enhance meat acceptability including a partial improvement in meat color. While, the enhancement of growth performance was not revealed.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KESAN PEMAKANAN *BORRERIA LATIFOLIA* KE ATAS PRESTASI
PERTUMBUHAN, CIRI-CIRI BANGKAI DAN KUALITI DAGING DALAM
AYAM KAMPUNG**

Oleh

ALI ABDUL AMEER

Oktober 2016

Pengerusi : Profesor Madya Azhar Kasim, PhD
Fakulti : Pertanian

Potensi keupayaan anti oksidan bagi *Borreria latifolia* dikaji dalam kajian ini. Dua eksperimen telah dijalankan untuk mengkaji kesan daripada suplemen pemakanan *Borreria latifolia* (BL) pada prestasi pertumbuhan, ciri-ciri bangkai, dan kualiti daging ayam kampung. Dua eksperimen telah dijalankan, percubaan pertama adalah untuk mengesahkan aktiviti antioksidan BL melalui perbandingan dengan *Officinalis Rosmarinus* (RO) sebagai rujukan di samping menilai kesannya pada prestasi dan kualiti daging. Tiga rawatan adalah digunakan: T1 (mengawal diet basal tanpa suplemen), T2 (diet basal dengan 1 % daripada BL), T3 (diet basal dengan 1 % daripada RO). Terdapat perbezaan yang signifikan ($p < 0.05$) antara rawatan pemakanan dalam nilai penurunan memasak. Penurunan nilai memasak adalah lebih tinggi pada T1 berbanding dengan T2 dan T3. Secara ketaranya ($p < 0.05$), nilai pH terjejas oleh rawatan diet dengan T2 dan T3 yang mempunyai nilai pH yang lebih tinggi berbanding dengan T1. Selain itu, kesan yang ketara bagi rawatan pemakanan dicatatkan semasa ujian penilaian deria manakala T2, T3 mempunyai skor yang lebih tinggi dalam penerimaan keseluruhan berbanding untuk mengawal dengan T2 lebih tinggi ($p < 0.05$) berbanding T3. Selain itu, kesan yang signifikan ($p < 0.05$) rawatan pemakanan nilai pengoksidaan lipid diperhatikan. T2 dan T3 mempunyai kepekatan MDA yang lebih rendah berbanding dengan T1. Selain itu, terdapat perbezaan signifikan ($p < 0.05$) antara jantan dan betina dikesan dalam berat akhir badan, berat badan, pengambilan makanan, berat bangkai, payudara dan berat paha, berat sayap, leher dan berat pinggang. Selain itu, kesan seks itu yang signifikan ($p < 0.05$) pada MUFA, PUFA, PUFA: Nisbah SFA, n6: Nisbah n3, nilai pH, kekuningan (b *) dan lipid pengoksidaan. Di samping itu, peningkatan ketara ($p < 0.05$) dalam kelembutan, pH, dan lipid pengoksidaan melalui bedah siasat penuaan period (1, 3, 5 d) telah direkodkan penuaan, semasa penurunan memasak (CL) dan kemerahan (* a) telah menurun. Interaksi yang signifikan antara herba dan seks di parameter pH, dan di antara herba dan seks, herba dan penuaan tempoh, seks dan penuaan tempoh, dan herba mengikut jantina oleh penuaan interaksi tempoh dalam hal dengan pengoksidaan lipid diperhatikan. Percubaan kedua adalah untuk mengkaji kesan tiga

tahap yang berbeza BL kepada kualiti daging dan prestasi pertumbuhan. Empat rawatan digunakan iaitu: T1 (mengawal diet basal tanpa suplemen), T2 (diet basal dengan 1.5 % daripada BL), T3 (diet basal dengan 2 % daripada BL) dan T4 (diet basal dengan 2.5 % daripada BL). Terdapat perbezaan yang bererti ($p < 0.05$) kesan rawatan pemakanan pada kemerahan (a *) nilai. T4 menunjukkan nilai lebih tinggi daripada T1, T2 dan T3. Tambahan pula, T4 dihasilkan ($p < 0.05$) pada skor yang lebih tinggi daripada T1, T2, T3 dan dalam kedua-dua penerimaan keseluruhan dan tekstur melalui ujian penilaian deria.

Berhubung dengan ujian lipid pengoksidaan, T4 menunjukkan kepekatan MDA yang paling rendah manakala T1 mempunyai kepekatan tertinggi MDA. Sementara itu, T2 dan T3 mempunyai kepekatan yang lebih tinggi MDA berbanding T4. Tambahan pula, ada perbezaan signifikan ($p < 0.05$) antara jantan dan betina dalam MUFA, PUFA, PUFA: Nisbah SFA, n6: Nisbah n3, kekuningan (* b) dan lipid pengoksidaan. Di samping itu, ketara kesan ($p < 0.05$) seks pada berat akhir badan, berat badan, pengambilan makanan, berat bangkai, payudara dan berat paha, berat sayap, leher dan berat badan pinggang, dan berat hati direkodkan. Keputusan yang diperolehi daripada kajian semasa menunjukkan signifikan ($p < 0.05$) mengurangkan kemerahan (* a) dan kehilangan memasak (CL) semasa bedah siasat tempoh penuaan manakala nilai pH, kelembutan, dan pengoksidaan lipid dinaikkan. Selain itu, terdapat interaksi yang signifikan antara herba dan seks, herba dan penuaan tempoh, seks dan penuaan tempoh, dan herba mengikut jantina oleh penuaan tempoh interaksi pada pengoksidaan lipid serta interaksi yang signifikan antara herba dan hubungan seks pada pH telah disedari dalam kajian ini. Secara umumnya, kajian ini mendedahkan bahawa suplemen pemakanan sebanyak 2.5% BL bertambah baik kestabilan oksidatif yang boleh dilanjutkan hidup daging rak dan boleh meningkatkan daging penerimaan termasuk peningkatan separa dalam warna daging. Sementara itu, peningkatan prestasi pertumbuhan tidak didedahkan.

ACKNOWLEDGEMENTS

My sincere thanks to the Almighty Allah for giving me the health and patience to complete this work.

I'm indebted to my wonderful supervisor, Associate Professor Dr. Azhar Kasim. It is difficult to find suitable words to show my deepest thanks and gratitude, for his guidance, patience, willingness to help, kindness and valuable advice. Special thanks for his endless support during the preparation of this thesis.

I am also thankful to the members of my supervisory committee, Prof. Dr Yazid Abdul Manap and Dr. Awis Qurni Bin Sazili for their suggestions and support during my study.

My sincere thanks and appreciation to my friends and my lab mates, the academic and supporting staff of Department of Animal Science, Faculty of Agriculture, and Poultry Unit for their technical assistance, kind cooperation and help during the field study at UPM.

I would like to express my appreciation to my loving, caring mother and father who advised me and supported me in my pursuit for my higher education and expressed understanding and consideration towards me. My deepest gratitude is extended to my brothers and sisters for their encouragement and support. I would like to extend my special sincere thanks to my bigger brother Ahmad Abdul Ameer for his advice and support.

I'm grateful to my true and actual souls mate my beloved wife Hawraa Hussein, for her patience, encouragement and support, and for caring for me and my beloved daughter. To my beloved daughter, Fatimah who is my life and make me a better father and man. You are an inspiration to me on a daily basis to be more than what I was the day before.


I certify that a Thesis Examination Committee has met on 13 October 2016 to conduct the final examination of Ali Abdulameer Abbood on his thesis entitled "Effects of *Borreria latifolia* (Aubl.) K.Schum. and *Rosmarinus officinalis* L. as Feed Additives on Performance and Meat Quality of Finishing Village Chicken" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Dahlan bin Ismail, PhD
Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Loh Teck Chwen, PhD
Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Wan Zahari Mohamed, PhD
Professor
Universiti Malaysia Kelantan
Malaysia
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 27 December 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Azhar Kasim, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Yazid Abdul Manap, PhD

Professor
Faculty of Food Science
Universiti Putra Malaysia
(Member)

Awis Qurni Sazili, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced
- the thesis has not been submitted previously or concurrently for any other degree at any institution
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and deputy vice-chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar paper, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No.: Ali Abdul Ameer, GS39697

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xv
LIST OF ABBRIVIATIONS	xvi
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	3
2.1 Village chicken in Malaysia	3
2.2 Medicinal herbs in chicken production	4
2.3 <i>Borreria latifolia</i>	4
2.4 <i>Rosmarinas officinalis</i>	5
2.5 Influence of herbs on growth performance	6
2.6 Chicken meat quality	7
2.6.1 Effect of color on meat quality	8
2.6.2 Water-holding capacity and meat quality	8
2.6.3 Effect of breast muscle pH on meat quality	9
2.6.4 Tenderness and meat quality	9
2.6.5 Influence of fatty acids on meat quality	10
2.6.6 Effect of lipid oxidation on meat quality	10
2.6.7 Effect of herbs on lipid oxidation	10
3 GENERAL MATERIALS AND METHODS	12
3.1 Preparation of herbs	12
3.2 Bird husbandry	12
3.3 Slaughtering and sample preparation	15
3.4 Proximate analysis	15
3.5 Determination of total phenolic compounds	15
3.6 PH measurement	16
3.7 Color measurement	16
3.8 Water holding capacity	16
3.8.1 Drip loss	16
3.8.2 Cooking loss	17
3.9 Meat tenderness	17
3.10 Fatty acid profile	18
3.11 Lipid oxidation	18
3.12 Sensory evaluation	19
3.13 Statistical analysis	19

4	CROSS COMPARISON BETWEEN DIETARY <i>BORRERIA LATIFOLIA</i> AND <i>ROSMARINUS OFFICINALIS</i> AND THEIR EFFECTS ON GROWTH PERFORMANCES, CARCASS CHARACTERISTICS AND MEAT QUALITY OF UPM-BRED VILLAGE CHICKEN	20
4.1	Introduction	20
4.2	Objectives	20
4.3	Materials and Methods	20
4.3.1	Treatments	20
4.3.2	Bird husbandry	21
4.3.3	Sampling and slaughtering procedure	21
4.3.4	Proximate analysis	21
4.3.5	Total phenolic compounds	21
4.3.6	PH measurement	21
4.3.7	Color measurement	21
4.3.8	Water holding capacity	22
4.3.9	Shear force value	22
4.3.10	Fatty acid profile	22
4.3.11	Lipid oxidation	22
4.3.12	Sensory evaluation	22
4.3.13	Statistical analysis	22
4.4	Results	22
4.4.1	Growth performance	22
4.4.2	Carcass characteristics	23
4.4.3	Chemical composition of <i>pectoralis</i> muscle	24
4.4.4	PH values	25
4.4.5	Color	27
4.4.6	Water holding capacity	27
4.4.7	Tenderness	28
4.4.8	Fatty acid profile	28
4.4.9	Lipid oxidation	30
4.4.10	Sensory evaluation	32
4.5	Discussion	32
4.5.1	Effect of BL & RO on growth performance	32
4.5.2	Effect of BL & RO on carcass characteristics	33
4.5.3	Effect of BL & RO on meat composition	34
4.5.4	Effect of BL & RO on pH values	34
4.5.5	Effect of BL & RO on meat color	34
4.5.6	Effect of BL & RO on WHC	35
	4.5.6.1 Cooking loss	35
	4.5.6.2 Drip loss	36
4.5.7	Effect of BL & RO on shear force value	36
4.5.8	Effect of BL & RO on fatty acid profile	37
4.5.9	Effect of BL & RO on lipid oxidation	37
4.5.10	Effect of BL & RO on sensory Evaluation	38
4.6	Conclusion	39

5	THE EFFECT OF THREE DIETARY LEVELS OF <i>BORRERIA LATIFOLIA</i> ON GROWTH PERFORMANCE AND MEAT QUALITY IN CROSSBRED VILLAGE CHICKEN	40
5.1	Introduction	40
5.2	Objectives	40
5.3	Materials and Methods	40
5.4	Results	41
5.4.1	Growth performance	41
5.4.2	Carcass characteristics	42
5.4.3	Chemical composition of <i>pectoralis</i> muscle	43
5.4.4	PH values	44
5.4.5	Meat color	45
5.4.6	Water holding capacity	46
5.4.7	Shear force value	46
5.4.8	Fatty acid profile	46
5.4.9	Lipid oxidation	48
5.4.10	Sensory evaluation	49
5.5	Discussion	49
5.5.1	Effect of BL on growth performance	49
5.5.2	Effect of BL on carcass characteristics	50
5.5.3	Effect of BL on chemical composition of meat	51
5.5.4	Effect of BL on pH values	51
5.5.5	Effect of BL on color	52
5.5.6	Effect of BL on WHC	52
5.5.6.1	Effect of BL on drip loss	52
5.5.6.2	Effect of BL on cooking loss	53
5.5.7	Effect of BL on shear force	53
5.5.8	Effect of BL on fatty acid profile	54
5.5.9	Effect of BL on lipid oxidation	54
5.5.10	Effect of BL on sensory evaluation	55
5.6	Conclusion	56
6	GENERAL DISCUSSION	57
7	CONCLUSIONS AND RECOMMENDATION FOR FUTURE WORK	65
	REFERENCES	66
	APPENDICES	86
	BIODATA OF STUDENT	90
	LIST OF PUBLICATIONS	91

LIST OF TABLES

Table		Page
3.1	Chemical composition and total phenolic compounds (TPH) of the dietary treatments, <i>Borreria latifolia</i> and <i>Rosmarinus officinalis</i>	13
4.1	Dietary supplementation of <i>Borreria latifolia</i> (BL) and <i>Rosmarinus officinalis</i> (RO) to the diet of village chicken	21
4.2	Effect of bird's sex and dietary supplementations of <i>Borreria latifolia</i> (BL) and <i>Rosmarinus officinalis</i> (RO) on body live weight, weight gain, feed conversion ratio, and feed intake in UPM-bred village chicken	23
4.3	Effect of bird's sex and dietary supplementations of <i>Borreria latifolia</i> (BL) and <i>Rosmarinus officinalis</i> (RO) on the weights of carcass, breast, thigh, wings, neck and loin and dressing percentage in UPM-bred village chicken	24
4.4	Effect of bird's sex and dietary supplementations of <i>Borreria latifolia</i> (BL) and <i>Rosmarinus officinalis</i> (RO) on the weights of heart, liver, spleen, and crop in UPM-bred village chicken.	24
4.5	Effect of bird's sex and dietary supplementations of <i>Borreria latifolia</i> (BL) and <i>Rosmarinus officinalis</i> (RO) on meat composition of <i>Pectoralis major</i> muscle in UPM-bred village chicken	25
4.6	Effect of bird's sex, post-mortem period and dietary supplementations of <i>Borreria latifolia</i> (BL) and <i>Rosmarinus officinalis</i> (RO) on the means of pH, lightness, redness, yellowness, cooking loss, drip loss and shear force of <i>Pectoralis major</i> muscle in UPM-bred village chicken	26
4.7	Effect of bird's sex and dietary supplementations of <i>Borreria latifolia</i> (BL) and <i>Rosmarinus officinalis</i> (RO) on fatty acid composition of <i>Pectoralis major</i> muscle in UPM-bred village chicken	29
4.8	Lipid oxidation values of <i>Pectoralis major</i> muscle during post-mortem aging periods in males and females of UPM-bred village chicken fed dietary treatments of <i>Borreria latifolia</i> (BL) and <i>Rosmarinus officinalis</i> (RO)	31
4.9	Effect of bird's sex and dietary supplementations of <i>Borreria latifolia</i> (BL) and <i>Rosmarinus officinalis</i> (RO) on sensory evaluation scores of <i>Pectoralis major</i> muscle in UPM-bred village chicken	32

5.1	Dietary supplementation of three levels of <i>Borreria latifolia</i> to the diet of village chicken	41
5.2	The effect of bird's sex and feeding different levels of <i>Borreria latifolia</i> (BL) on body live weight, weight gain, feed conversion ratio, and feed intake in crossbred village chicken	42
5.3	The effect of bird's sex and feeding different levels of <i>Borreria latifolia</i> (BL) on the dressing percentage and weights of carcass, breast, thigh, wings, neck and loin in crossbred village chicken.	43
5.4	The effect of feeding different levels of <i>Borreria latifolia</i> (BL) and bird's sex on weights of heart, liver, spleen, and crop in crossbred village chicken	43
5.5	The effect of bird's sex and dietary supplementations of different levels of <i>Borreria latifolia</i> (BL) on meat composition of Pectoralis major muscle in crossbred village chicken	44
5.6	The effect of feeding different levels of <i>Borreria latifolia</i> (BL), bird's sex and post-mortem period on the means of pH value, lightness, redness, yellowness, cooking loss, drip loss and shear force of Pectoralis major muscle in crossbred village chicken	45
5.7	Effect of feeding different levels of <i>Borreria latifolia</i> and bird's sex on fatty acid composition (mg/100 g) of pectoralis major muscle in crossbred village chicken	47
5.8	Effect of post-mortem period, bird's sex and different levels of <i>Borreria latifolia</i> (BL) on lipid oxidation values of <i>Pectoralis</i> major muscle in crossbred village chicken	48
5.9	Effect of feeding <i>Borreria latifolia</i> (BL) and bird's sex on sensory scores of <i>Pectoralis</i> major muscle samples in crossbred village chicken	49


LIST OF FIGURES

Figure		Page
2.1	<i>Borreria latifolia</i> (BL)	5
2.2	<i>Rosmarinus officinalis</i>	5
3.1	Methodology of the research	14
3.2	<i>Pectoralis</i> muscle of village	15
4.1	Sex by herb interaction in pH value for males and females in <i>Pectoralis</i> muscle of village chicken.	27
4.2	Herb by sex by period interaction ($p < 0.05$) in lipid oxidation test for males	30
4.3	Herb by sex by period interaction ($p < 0.05$) in lipid oxidation test for females	30

LIST OF ABBREVIATIONS

a*	Redness
b*	Yellowness
BL	<i>Borreria latifolia</i>
°C	Celsius degrees
CL	Cooking loss
CP	Crude protein
cm ²	Square centimeter
d	Day
ddH ₂ O	Deionized distilled water
DL	Drip loss
DM	Dry matter
EE	Ether extract
FA	Fatty acid
FCR	Feed conversion ratio
g	Gram
GC	Gas chromatography
HCL	Hydrochloric acid
h	Hour
H ₂ SO ₄	Sulfuric acid
kg	kilogram
KOH	Potassium hydroxide
L	Liter
L*	Lightness
MDA	Malondialdehyde
mg	Milligram
min	Minute
ml	Milliliter
MUFA	Monounsaturated fatty acids
OM	Organic matter
PBS	Phosphate buffered saline
PSE	Pale Soft Executive
PUSFAs	Polyunsaturated fatty acids
PUFA:SFA ratio	Total PUFA to Total SFA ratio
RCBD	Randomized completely block design
RO	<i>Rosmarinus officinalis</i>
SE	Standard error

Se	Second
SFA	Saturated fatty acids
TBA	Thiobarbituric acid
TBARS	Thiobarbituric acid reactive substances
TCA	Trichloroacetic acid
USFA	Unsaturated fatty acid
USFS:SFA ratio	Total USFA to Total SFA ratio
μl	Micro liter
μM	Micromole
VSF	Volodkovitch Shear Force
v/v	Volume per volume
WHC	Water holding capacity



CHAPTER 1

GENERAL INTRODUCTION

Traditionally, herbs have been used for centuries in many civilizations to treat certain medical conditions due to the efficiency and availability of these herbs (Schulz *et al.*, 2001). Primitive man had been impressed by the diversity of plants that could be obtained, which provided food, clothing and medicine. The attention given on the use of medicinal herbs was the result of years of struggle against sicknesses which drove man to seek medication in tree barks, seeds and different parts of plants. Herbs have been the source of treatment and prophylaxis (Kelly, 2009). In addition, substances from herbs offer great promise to the pharmaceutical industry for the production of medication to treat coronary illness, hypertension, torment, asthma, and various other medical conditions.

In poultry production, the utilization of intricate intensive poultry production systems creates the revelation and prevalence of using various herbs as feed additives. The principle goal of adding herbs is to boost animal performance and production quality. Herbs and their extracts are included in chicken feed to improve the utilization of feed nutrients and lead to better feed efficiency and higher production. Several herbs and spices contain synthetic substances, like polyphenols, quinines, and polypeptides. Some of them show remedial qualities, such as antioxidant activities (Li, 1993).

Poultry meat is particularly prone to oxidative deterioration due to high concentration of polyunsaturated fatty acids (PUFA; Igene and Pearson, 1979). There are many studies showing an improvement in the oxidative stability of tissue after feeding poultry with antioxidant compounds added into the diet (Botsoglou *et al.*, 2002; Lee *et al.*, 2004; Govaris *et al.*, 2005). There is a trend to search for compounds that may allow a shift from synthetic to natural antioxidants (Sheehy *et al.*, 1995; Yanishlieva, 2001; Botsoglou *et al.*, 2002). This trend is justified because a carcinogenic potential from the use of synthetic antioxidants has been suspected (Madavi & Salunkhe, 1995).

Plant phenols show *in vitro* antioxidant activity, acting to restrain lipid peroxidation as chain-breaking peroxy radical scavengers (Proestos *et al.*, 2006; Miguel, 2010). Along these lines, they can assume a defensive part aimed at unsaturated lipids in food against oxidative damage (Couladis *et al.*, 2003). Oxidative deterioration is the major non-microbial element that causes value crumbling in foods (Descalzo *et al.*, 2005). Throughout the handling, processing and storage of fresh meat, lipid oxidation is associated with discoloration, drip losses, off-flavor and off-odor development in addition to the ability to form toxic compounds (Gray *et al.*, 1996).

A few *in vivo* studies have been directed to test natural substances capable of enhancing the stability of lipid in meat. It has been shown that dietary supplementation with natural antioxidants, for example, rosemary (*Rosmarinus officinalis*), sage (*Salvia officinalis*) and oregano (*Origanum vulgare*) derivatives may improve antioxidant defenses and meat shelf life (Lopez-Bote *et al.*, 1998, Florou-Paneri *et al.*, 2006). Among many kinds of herbs, *Borreria latifolia* is an annual herb and a member of the Rubiaceae family, which develops as a prevailing weed on waste ranges and/or agricultural farms. It originated from tropical

America and was ordinarily dispersed in India, Southeast Asia, and Malaysia. *Borreria alata*, *Borreria scaberrima* bold, *Spermacoce latifolia* Aubl, are synonyms for *Borreria latifolia*.

In the present study, addition of *Borreria latifolia* and *Rosmarinus officinalis* as feed additives was to improve meat quality and evaluating the performance at finishing stage of village chicken. However, there is a dearth of information and published scientific data on the effects of *Borreria latifolia* as feed supplementation for growth performance and meat quality in chicken diets. Therefore, the present study was carried out to determine the effect of adding *Borreria latifolia* to the finisher diets on growth performances, carcass characteristics and meat quality. Hence, this study is aimed at achieving these objectives:

1. To determine the capability of *Borreria latifolia* as a natural antioxidant.
2. To examine the impact of feeding *Borreria latifolia* on the performance of village chicken.
3. To evaluate the effect of feeding *Borreria latifolia* on chicken meat parameters.

REFERENCES

- Aberra, M., Tegene, N., (2011). Phenotypic and morphological characterization of indigenous chicken populations in southern region of Ethiopia. *Animal Genetic Resources*, 49, 19–31.
- Acamovic T. and Brooker J.D. (2005). Biochemistry of plant secondary metabolites and their effects in animals. *Proceedings of the Nutrition Society*, 64(3), p.403.
- Ahmad R, Ali A.M., Israf D.A., Ismail N.H., Shaari K. and Lajis N.H. (2005). Antioxidant, radical-scavenging, anti-inflammatory, cytotoxic and antibacterial activities of methanolic extracts of some Hedyotis species. *Life Science*, 76: 1953-1964.
- Ahn, D.U., Wolfe, F.H. and im, J. (1995). Dietary α -Linolenic Acid and Mixed Tocopherols, and Packaging Influences on Lipid Stability in Broiler Chicken Breast and Leg Muscle. *Journal of Food Science*, 60(5), pp.1013-1018.
- Aji, S.B., Ignatuius, K., Ado, A.Y., Nuhu, J.B. and Abdulkarim, A. (2011). Effect of feeding onion (*Allium cepa*) and garlic (*Allium sativum*) on some performance characteristics of broiler chickens. *Research Journal of Poultry Science*, 4: 22-27.
- Alçiçek, A, Bozkurt, M. and Çabuk, M. (2003). The effects of an essential oil combination derived from selected herbs growing wild in Turkey on broiler performance. *South African Journal of Animal Science*, 33: 89–94.
- Alçiçek, A., Bozkurt, M. and Çabuk, M. (2004). The effect of a mixture of herbal essential oils, an organic acid or a probiotic on broiler performance. *South African Society for Animal Science*. 34 (4). pp: 217-222
- Al-Harthi, M.A. (2002). Performance and carcass characteristics of broiler chicks as affected by different dietary types and levels of herbs and spices as non classical growth promoters. *Egypt. Poult. Sci*, 22(1), pp.325-343.
- Al-Kassie, G.A.M. (2009). Influence of two plant extracts derived from thyme and cinnamon on broiler performance. *Pakistan Veterinary Journal*, 29: 1-5.
- Allen, C.D., Fletcher, D.L., Northcutt, J.K. and Russell, S.M. (1998). The relationship of broiler breast color to meat quality and shelf-life. *Poultry Science*, 77(2), pp.361-366.
- Allen, C.D., Russell, S.M., and Fletcher, D.L. (1997). The relationship of broiler breast meat color and pH to shelf life and odor development. *Poultry Science*, 76:1042–1046
- Allinson, I.B., Ekunseitan, D.A., Ayoola, A.A., Ogunade, I.M., and Njoku, C.P. (2013). Effect of herbal supplement on growth response and faecal egg counts of cockerels. *Online Journal of Animal Feed Research*, 3: 68-73.
- Altinier, G., Sosa, S., Aquino, R.P., Mencherini, T., Loggia, R.D. and Tubaro, A. (2007). Characterization of topical anti-inflammatory compounds in *Rosmarinus officinalis* L. *Journal of Agricultural and Food Chemistry*, 55(5), pp.1718-1723.

- Alvarado, C.Z., and Sams, A.R. (2004). Early postmortem injection and tumble marination effects on broiler breast meat tenderness. *Poultry Science*, 83(6), 1035–1038.
- AMSA. 2012. AMSA Meat Color Measurement Guidelines. *American Meat Science Association*, Illinois, USA.
- AOAC. (1984). Official Methods of Analysis, 14th edn. Association of Official Analytical Chemists, Arlington, VA.
- Ayeleso, A.O., Oguntibeju, O.O. and Brooks, N.L. (2014). In Vitro Study on the Antioxidant Potentials of the Leaves and Fruits of *Nauclea latifolia*. *The Scientific World Journal*, 2014.
- Bakhiet, A.O. and Adam, S.E.I. (1995). Therapeutic Utility, Constituents and Toxicity of Some Medicinal Plants: A Review. *Veterinary Human Toxicol*, 37: 255-258.
- Bampidis, V.A., Christodoulou, V., Florou-Paneri, P., Christaki, E., Chatzopoulou, P.S., Tsiligianni, T. and Spais, A.B. (2005). Effect of dietary dried oregano leaves on growth performance, carcass characteristics and serum cholesterol of female early maturing turkeys. *British Poultry Science*, 46, 595–601.
- Basmacıoğlu, H., okuşoğlu, Ö. and Ergül, M. (2004). "The effect of oregano and rosemary essential oils or alpha-tocopheryl acetate on performance and lipid oxidation of meat enriched with n- PUFA's in broilers." *South African Journal of Animal Science*, 34.3: 197.
- Battula, V., Schilling, M.W., Vizzier-Thaxton, Y., Behrends, J.M., Williams, J.B. and Schmidt, T.B. (2008). The effects of low-atmosphere stunning and deboning time on broiler breast meat quality. *Poultry Science*, 87:1202-1210.
- Bendall, J.R. (1975). "Cold-contraction and atp-turnover in the red and white musculature of the pig, post mortem." *Journal of the Science of Food and Agriculture*, 26. (1) 55-71.
- Benkeblia, N. (2004). Antimicrobial activity of essential oil extracts of various onions (*Allium cepa*) and garlic (*Allium sativum*). *LWT-Food Science and Technology*, 37: 263–268.
- Biswas, A.H. and Wakita, M. (2001). Effect of Dietary Japanese Green Tea Powder Supplementation on Feed Utilization and Carcass Profiles in Broilers. *Journal of Poultry Science*, 38(1), pp.50-57.
- Bogosavljević-Bošković, Pavlovski, Z., Petrović, M.D., Dosković, V. and Rakonjac, S. (2015). Broiler meat quality: Proteins and lipids of muscle tissue. *African Journal of Biotechnology*, 9(54), pp.9177-9182.
- Bölükbaşı, Ş.C., Erhan, M.K. and Özkan, A. (2006). "Effect of dietary thyme oil and vitamin E on growth, lipid oxidation, meat fatty acid composition and serum lipoproteins of broilers." *South African Journal of Animal Science*, 36.(3) 189-196.
- Botsoglou, E., Govaris, A., Fletouris, D. and Iliadis, S. (2013). Olive leaves (*Olea europea* L.) and α -tocopheryl acetate as feed antioxidants for improving the oxidative stability of α -linolenic acid-enriched eggs. *Journal of animal physiology and animal nutrition*, 97(4), pp.740-753.

- Botsoglou, N., Florou-Paneri, E. Botsoglou, P., Dotas, V., Giannenas, I., Koidis, A. and Mitrakos, P. (2005). The effect of feeding rosemary, oregano, saffron and α -tocopheryl acetate on hen performance and oxidative stability of eggs. *South African Journal of Animal Science*, 35: 143-151.
- Botsoglou, N.A.P. Florou-Paneri, E. Christaki, D.J. Fletouris and A.B. Spais, (2002). Effect of dietary oregano essential oil on performance of chickens and on iron-induced lipid oxidation of breast, thigh and abdominal fat tissues. *British Poultry Science*, 43: 223-230.
- Botsoglou, N.A., Christaki, E., Fletouris, D.J., Florou-Paneri, P. and Spais A.B. (2002). The effect of dietary oregano essential oil on lipid oxidation in raw and cooked chicken during refrigerated storage. *Meat Science*, 62:259–265.
- Bragagnolo, N. (2001), December. Aspectos comparativos entre carnes segundo a composição de ácidos graxos e teor de colesterol. In Conferência Internacional Virtual sobre Qualidade de Carne Suína (Vol. 2, pp. 393-402). Concórdia: Embrapa Suínos e Aves.
- Bravo, L. (1998). Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutrition Reviews*, 56, 317–333.
- Brenes, A., Viveros, A., Goñi, I., Centeno, C., Saura-Calixto, F. and Arijia, I. (2010). Effect of grape seed extract on growth performance, protein and polyphenol digestibilities, and antioxidant activity in chickens. *Spanish Journal of Agricultural Research*, 8(2), pp.326-335.
- Byarugaba, D. (2007). "The structure and importance of the commercial and village based poultry systems in Uganda." Kampala, FAO. (Consultancy report). Available at: apfuganda.ning.com/forum/attachment/download.
- Cabuk, M., Bozkurt, M., Alcicek, A., Akbağ, Y. and Küçükyılmaz, K. (2006). Effect of a herbal essential oil mixture on growth and internal organ weight of broilers from young and old breeder flocks. *South African Journal of Animal Science*, 36(2), pp.135-141.
- Campo, J.D. and Amiot, M.J. (2000). Antimicrobial effect of rosemary extracts. *Journal of Food Protection*, 63(10), pp.1359-1368.
- Candan, F., Unlu, M., Tepe, B., Daferera, D., Polissiou, M., Sokmen, A. and Akpulat, A. (2003). Antioxidant and antimicrobial activity of the essential oil and methanol extracts of *Achillea millefolium* subsp. *millefolium* Afan. (Asteraceae). *Journal of Ethnopharmacology*, 87: 215–220.
- Cañeque V. La canal de cordeiro. (1989). In: Producción de Carne de Cordeiro. (1989). México, MEX. México. 367-436.
- Cavitt, L.C., Meullenet, J.F., Gandhapuneni, R.K., Youm, G.W. and Owens, C.M. (2005). Rigor development and meat quality of large and small broilers and the use of Allo-Kramer shear, needle puncture, and razor blade shear to measure texture. *Poultry Science*, 84(1), pp.113-118.
- Chen, A., Xu, J., Yang, C. and Hong, Q. (2009). Effects of cinnamon extracts on growth performance and excreta urease activity and nitrogen loss in broilers. In *Livestock Environment VIII*, 31 August–4 September 2008, Iguassu Falls, Brazil (p. 46). *American Society of Agricultural and Biological Engineers*.

- Chen, X.D., Ma, Q.G., Tang, M.Y. and Ji, C. (2007). Development of breast muscle and meat quality in Arbor Acres broilers, Jingxing 100 crossbred chickens and Beijing fatty chickens. *Meat science*, 77(2), pp.220-227.
- Chen, Y.J., Kim, I.H., Cho, J.H., Yoo, J.S., Wang, Q., Wang, Y. and Huang, Y. (2008). Evaluation of dietary L-carnitine or garlic powder on growth performance, dry matter and nitrogen digestibilities, blood profiles and meat quality in finishing pigs. *Animal feed science and technology*, 141(1), pp.141-152.
- Choudhury, K.D., Choudhury, M.D. and Baruah, M. (2012). Antibacterial activity of some plants belonging to the family Rubiaceae: a review. *World Journal of Pharmacy and Pharmaceutical Sciences*, 1(3), pp.1170-1192.
- Chouliara, E., Karatapanis, A., Savvaidis, I.N. and Kontominas, M.G. (2007). Combined effect of oregano essential oil and modified atmosphere packaging on shelf-life extension of fresh chicken breast meat, stored at 4 C. *Food Microbiology*, 24(6), pp.607-617.
- Christaki, E. (2012). Naturally derived antioxidants in poultry nutrition. *Research Journal of Biotechnology*, 7(3).
- Coleman, R.A., and Korver, D.R. (2004). "Amino acid requirements of broilers: relationships with growth and meat quality." Proceedings of the 16th Australian Poultry Science Symposium, Sydney, New South Wales, Australia, 9-11 February. *Poultry Research Foundation*.
- Conserva, L.M. and Jesu Costa Ferreira, J. (2012). Borreria and Spermaceoce species (Rubiaceae): A review of their ethnomedicinal properties, chemical constituents, and biological activities. *Pharmacognosy reviews*, 6(11), p.46.
- Corzo, A., Schilling, M.W., Loar, R.E., Jackson, V., Kin, S. and Radhakrishnan, V. (2009). The effects of feeding distillers dried grains with solubles on broiler meat quality. *Poultry Science*, 88(2), pp.432-439.
- Cosgrove, J.P., Church, D.F., and Pryor, W.A. (1987). The kinetics of the autoxidation of polyunsaturated fatty-acids. *Lipids*, 22, 299-304.
- Couladis, M., Tzakou, O., Verykokidou, E. and Harvala, C. (2003). Screening of some Greek aromatic plants for antioxidant activity. *Phytotherapy research*, 17(2), pp.194-195.
- Cross D.E., Acamovic T., Deans S.G., Cdevitt R.M. (2002). The effects of dietary inclusion of herbs and their volatile oils on the performance of growing chickens. *British Poultry Science*, 43, S33-S35.
- Cross D.E., McDevitt R.M., Hillman K., Acamovic T., (2007). The effect of herbs and their associated essential oils on performance, dietary digestibility and gut microflora in chickens from 7 to 28 days of age. *British Poultry Science*, 48, 496-506.
- Cuppett, Susan L., and Clifford A. Hall. (1998). "Antioxidant activity of the Labiatae." *Advances in food and nutrition research* 42: 245-272.
- De Marchi, M., Cassandro, M., Lunardi, E., Baldan, G. and Siegel, P.B. (2005). Carcass characteristics and qualitative meat traits of the Padovana breed of chicken. *International Journal of Poultry Science*, 4(4), pp.233-238.

- Debersac, P., Vernevaut, M.F., Amiot, M.J., Suschetet, M. and Siess, M.H. (2001). Effects of a water-soluble extract of rosemary and its purified component rosmarinic acid on xenobiotic-metabolizing enzymes in rat liver. *Food and Chemical Toxicology*, 39(2), pp.109-117.
- Demir E., Sarica S., Ozcan M.A., Suicmez M. (2003). The use of natural feed additives as alternatives for an Antibiotic growth promoter in broiler diets. *British Poultry Science*, 44, S44–S45.
- Denbow, M. D. (2003). Factors Affecting Meat Quality. <http://www.poultryscience.org/pba/1952-2003/2000/2000%20Denbow.pdf>. Accessed on February 7th, 2010.
- Denli, M., Okan, F. and Uluocak, A.N. (2004). "Effect of dietary supplementation of herb essential oils on the growth performance, carcass and intestinal characteristics of quail (*Coturnix coturnix japonica*)." *South African Journal of Animal Science* 34: 174-179.
- Descalzo, A.M., Insani, E.M., Biolatto, A., Sancho, A.M., Garcia, P.T., Pensel, N.A. and Josifovich, J.A. (2005). Influence of pasture or grain-based diets supplemented with vitamin E on antioxidant/oxidative balance of Argentine beef. *Meat Science*, 70(1), pp.35-44.
- Dickens, J.A. and Lyon, C.E. (1995). The effects of electric stimulation and extended chilling times on the biochemical reactions and texture of cooked broiler breast meat. *Poultry Science*, 74(12), pp.2035-2040.
- Doley, S., Gupta, J.J. and Reddy, P.B. (2009). Effect of supplementation of ginger, garlic and turmeric in broiler chicken. *Indian Veterinary Journal*, 86(6), pp.644-645.
- Dorman, H.J.D., Deans, S.G., Noble, R.C., Surai, P. (1995). Evaluation in vitro of plant essential oils as natural antioxidants. *Journal Essential Oil Research*, 7, 645– 651.
- Dransfield, E. and Sosnicki, A. A. (1999). Relationship between muscle growth and poultry meat quality. *Poultry Science*, 78(5), pp.743-746.
- Durrani, F.R., Ismail, M., Sultan, A., Suhail, S.M., Chand, N. and Durrani, Z. (2006). Effect of different levels of feed added turmeric (*Curcuma longa*) on the performance of broiler chicks. *Journal of Agricultural and Biological Science*, 1(2), pp.9-11.
- Ebrahimi, M., Rajion, M.A., Goh, Y.M. and Sazili, A.Q. (2012). Impact of different inclusion levels of oil palm (*Elaeis guineensis* Jacq.) fronds on fatty acid profiles of goat muscles. *Journal of Animal Physiology and Animal Nutrition*, 96(6), pp.962-969.
- Edwards, H.M. and Denman, F. (1975). Carcass composition studies 2. Influences of breed, sex and diet on gross composition of the carcass and fatty acid composition of the adipose tissue. *Poultry Science*, 54(4), pp.1230-1238.
- El-Deek, A.A., Attia, Y.A., Maysa, M., Hannfy, M. (2002). Effect of anise (*Pimpinella anisum*), ginger (*Zingiber officinale* Roscoe) and fennel (*Foeniculum vulgare*) and their mixture on performance of broilers *Archiv für Geflügelkunde*, 67(2), pp.92-96.

- Engberg, R.M., Jensen, B.B. and Hojberg, O., (2007), August. Plant of the Juglandaceae family as alternative to antibiotic growth promoters in broiler production. In 16th European Symposium on Poultry Nutrition, Strasbourg, France.
- Engku Azahan, E.A., Yeong, S.W. and Noraziah, M. (1990), October. Intensive rearing of kampung chickens for meat production. In Proc. 2nd Conf. Veterinary Assoc. Malaysia (pp. 6-10).
- Fadlalla, I.M.T., Mohammed, B.H. and Bakhiet, A.O. (2010). Effect of feeding garlic on the performance and immunity of broilers. *Asian Journal of Poultry Science*, 4(4), pp.182-189.
- Falowo, A.B., Fayemi, P.O. and Muchenje, V. (2014). Natural antioxidants against lipid-protein oxidative deterioration in meat and meat products: A review. *Food Research International*, 64, pp.171-181.
- Fellenberg, M.A. and Speisky, H. (2006). Antioxidants: their effects on broiler oxidative stress and its meat oxidative stability. *World's Poultry Science Journal*, 62(01), pp.53-70.
- Fletcher, D.L. (1992). Methodology for achieving pigment specifications. *Poultry science*, 71(4), pp.733-743.
- Fletcher, D.L. (2002). Poultry meat quality. *World's Poultry Science Journal*, 58 (02), pp.131-145.
- Fletcher, D.L. (1999). Broiler breast meat color variation, pH, and texture. *Poultry Science*, 78:1323-1327.
- Florou-Paneri, P., Giannenas, I., Christaki, E., Govaris, A. and Botsoglou, N. (2006). Performance of chickens and oxidative stability of the produced meat as affected by feed supplementation with oregano, vitamin C, vitamin E and their combinations. *Archiv fur Geflugelkunde*, 70(5), pp.232-239.
- Folch, J., Lees, M. and Sloane-Stanley, G.H. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *The Journal Biological chemistry*, 226(1), pp.497-509.
- Freitas, R.D., Fonseca, J.B., Soares, R.D.T.R.N., Rostagno, H.S. and Soares, P.R. (2001). Utilization of garlic (*Allium sativum* L.) as growth promoter of broilers. *Revista Brasileira de Zootecnia*, 30(3), pp.761-765.
- Froning, G.W. (1995). Color of poultry meat. *Poultry and avian biology reviews*.
- Ghazalah, A.A. and Ali, A.M. (2008). Rosemary leaves as a dietary supplement for growth in broiler chickens. *International Journal of Poultry Science*, 7(3), pp.234-239.
- Gill, C. (2000). Botanical feed additives. *Feed. Int*, 21, pp.14-17.
- Gollnisch, K., Halle, I., Schubert, R., Flachowsky, G., Jahreis, G. and Bitsch, R. (2001). Effects of essential oils and herbs in animal nutrition. In Vitamine und Zusatzstoffe in der Ernährung von Mensch und Tier. 8. Symposium, 26. and 27. September, 2001, Jena/Thüringen, Germany. (pp. 197-204). Friedrich-Schiller-Universität.

- Gopi, M., Karthik, K., Manjunathachar, H.V., Tamilmahan, P., Kesavan, M., Dashprakash, M., Balaraju, B.L. and Purushothaman, M.R., (2014). Essential oils as a feed additive in poultry nutrition. *Advances in Animal and Veterinary Science*, 2(1), pp.1-7.
- Govaris, A., Botsoglou, E., Florou-Paneri, P., Moulas, A. and Papageorgiou, G. (2005). Dietary supplementation of oregano essential oil and-tocopheryl acetate on microbial growth and lipid oxidation of turkey breast fillets during storage. *International Journal of Poultry Science*, 4(12), pp.969-975.
- Govaris, A., Botsoglou, E., Moulas, A. and Botsoglou, N. (2010). Effect of dietary olive leaves and rosemary on microbial growth and lipid oxidation of turkey breast during refrigerated storage. *South African Journal of Animal Science* 40: 145-155.
- Gray, J.I., Gomaa, E.A. and Buckley, D.J. (1996). Oxidative quality and shelf life of meats. *Meat science*, 43, pp.111-123.
- Grela, E.R. (2000). Influence of herb supplements in pig feeding on carcass traits and some organoleptic and chemical parameters of meat. *Roczniki Naukowe Zootechniki*, (Supplement z. 6), pp.167-171.
- Griggs, J.P. and Jacob, J.P. (2005). Alternatives to antibiotics for organic poultry production. *The Journal of Applied Poultry Research*, 14(4), pp.750-756.
- Groom, G.M. (1990). Factors affecting poultry meat quality. *L'Aviculture en Méditerranée. Opt Mediterr Ser A*, 7, pp.205-210.
- Guo, F., wakkel R.P.K. and Versteegen, M.W.A. (2000). The use of Chinese herbs as alternative for growth promoters in broiler diets. Proc. of XII World's Poultry. Cong., 20-24 Aug., 2000, Montreal, Canada.
- Halima, H., Neser, F.W.C., van Marle-Koster, E. and de Kock, A. (2007). Phenotypic variation of native chicken populations in northwest Ethiopia. *Tropical animal health and production*, 39(7), pp.507-513.
- Hascik, P., Elimam, I.O.E., Garlik, J., Kacaniova, M., Cubon, J., Bobko, M., Vavrisinova, K. and Arpasova, H. (2013). The effect of bee pollen as dietary supplement on meat chemical composition for broiler Ross 308. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis* (Czech Republic).
- Hassan W.A. and Adamu U.A. (1997). Pigeon genetic resources in semi-arid zone of Nigeria: Initial results from characterization studies. Paper submitted at INFPD Workshop and General Meeting, 9-13 December 1997, M'Bour, Senegal.
- Havenstein, G.B., Ferket, P.R. and Qureshi, M.A. (2003). Carcass composition and yield of 1957 versus 2001 broilers when fed representative 1957 and 2001 broiler diets. *Poultry Science*, 82(10), pp.1509-1518.
- Heim, K.E., Tagliaferro, A.R. and Bobilya, D.J. (2002). Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *The Journal of Nutritional Biochemistry*, 13(10), pp.572-584.

- Hernandez, F., Madrid, J., Garcia, V., Orengo, J. and Megias, M.D. (2004). Influence of two plant extracts on broilers performance, digestibility, and digestive organ size. *Poultry Science*, 83(2), pp.169-174.
- Hertrampf J.W. (2001). Alternative antibacterial performance promoters. *Poultry International*, 40:50–52.
- Ho, C.T., Wang, M., Wei, G.J., Huang, T.C. and Huang, M.T. (2000). Chemistry and antioxidative factors in rosemary and sage. *Biofactors*, 13(1-4), pp.161- 166.
- Holcman A., Vadnjal R., Žlender B., tibilj V. (2003): Chemical composition of chicken meat from free range and extensive indoor rearing. *Arch. Geflügelk.*, 67, 120–124].
- Honikel, K.O. (1998). Reference methods for the assessment of physical characteristics of meat. *Meat Science*, 49(4), 447-457.
- Huff, E.J. and Parrish, F.C. (1993). Bovine longissimus muscle tenderness as affected by postmortem aging time, animal age and sex. *Journal of Food Science*, 58(4), pp.713-716.
- Hughes, R.J., Brooker, J.D. and Smyl, C. (2005). Growth rate of broiler chickens given condensed tannins extracted from grape seed. *In Australian Poultry Science Symposium* (Vol. 17, pp. 65-68).
- Hunt, M.C. (1980). Meat color measurements. Meat Conference of the American Meat Science Assoc. Purdue University, Lafayette, Ind. 41-46.
- Igene, J.O. and Pearson, A.M. (1979). *Journal of Food Science*, 44, 1285.
- Isidahomen, C.E., Ilori B.M. and Akano, K. (2012). Genetic and Sex Differences in Carcass Traits of Nigerian Indigenous Chickens. *Journal of Animal Science Advances*. 2: 637-648.
- Ismail, N.H., Ali, A.M., Aimi, N., Kitajima, M., Takayama, H. and Lajis, N.H. (1997). Anthraquinones from *Morinda elliptica*. *Phytochemistry*, 45(8), pp.1723-1725.
- Jamroz D, Kamel C. Wiliczekiewicz A. Wertelecki T. Orda J. and Skorupinska J. (2005). Use of active substances of plant origin in chicken diets based on maize and locally grown cereals. *British Poultry Science*, 46: 485–493.
- Jang, A., Liu, X.D., Shin, M.H., Lee, B.D., Lee, S.K., Lee, J.H. and Jo, C. (2008). Antioxidative potential of raw breast meat from broiler chicks fed a dietary medicinal herb extract mix. *Poultry Science*, 87(11), pp.2382-2389.
- Janz, J.A.M., Morel, P.C.H., Wilkinson, B.H.P. and Purchas, R.W. (2007). Preliminary investigation of the effects of low-level dietary inclusion of fragrant essential oils and oleoresins on pig performance and pork quality. *Meat Science*, 75(2), pp.350-355.
- Jaturasitha, S., Srikanchai, T., Kreuzer, M. and Wicke, M. (2008). Differences in carcass and meat characteristics between chicken indigenous to northern Thailand (Black-boned and Thai native) and imported extensive breeds (Bresse and Rhode Island Red). *Poultry Science*, 87(1), pp.160-169.

- Jensen, C., Skibsted, L.H. and Bertelsen, G. (1998). Oxidative stability of frozen-stored raw pork chops, chill-stored pre-frozen raw pork chops, and frozen-stored pre-cooked sausages in relation to dietary CuSO₄, rapeseed oil and Vitamin E. *Zeitschrift für Lebensmitteluntersuchung und -Forschung A*, 207 (5), pp. 363-368.
- Jensen, C., Skibsted, L.H., Jakobsen, K. and Bertelsen, G. (1995). Supplementation of broiler diets with all-rac-alpha or a mixture of natural source RRR-alpha-, gamma-, delta-tocopheryl acetate. 2. Effect on the oxidative stability of raw and precooked broiler meat products. *Poultry Science*, 74(12), pp.2048-2056.
- Kähkönen, M.P., Hopia, A.I., Vuorela, H.J., Rauha, J.P., Pihlaja, K., Kujala, T.S. and Heinonen, M. (1999). Antioxidant activity of plant extracts containing phenolic compounds. *Journal of Agricultural and Food Chemistry*, 47(10), pp.3954-3962.
- Kannan, G., Kouakou, B. and Gelaye, S. (2001). Color changes reflecting myoglobin and lipid oxidation in chevon cuts during refrigerated display. *Small Ruminant Research*, 42(1), pp.67-74.
- Kanner, J. (1994). Oxidative processes in meat and meat products: quality implications. *Meat Science*, 36(1), pp.169-189.
- Karpinska, M., Borowski, J. and Danowska-Oziewicz, M. (2000). Antioxidative activity of rosemary extract in lipid fraction of minced meat balls during storage in a freezer. *Food/Nahrung*, 44(1), pp.38-41.
- Kelly-Pieper, K., Patil, S.P., Busse, P., Yang, N., Sampson, H., Li, X.M., Wisnivesky, J.P. and Kattan, M. (2009). Safety and Tolerability of an Antiasthma Herbal Formula (A HMI™) in adult subjects with asthma: a randomized, double-blinded, placebo-controlled, dose-escalation Phase I study. *The Journal of Alternative and Complementary Medicine*, 15(7), pp.735-743.
- Kgwatalala, P.M., Bolowe, A.M., Thutwa, K. and Nsoso, S.J. (2013). Carcass traits of the naked-neck, dwarf and normal strains of indigenous Tswana chickens under an intensive management system. *Breast*, 62, pp.0-91.
- Khan, R.U., Nikousefat, Z., Tufarelli, V., Naz, S., Javdani, M. and Laudadio, V. (2012). Garlic (*Allium sativum*) supplementation in poultry diets: effect on production and physiology. *World's Poultry Science Journal*, 68(03), pp.417-424.
- Kidd, M.T., Corzo, A., Hoehler, D., Miller, E.R. and Dozier, W.A. (2005). Broiler responsiveness (Ross x 708) to diets varying in amino acid density. *Poultry Science*, 84(9), pp.1389-1396.
- Kim, Y.J., Jin, S.K. and Yang, H.S. (2009). Effect of dietary garlic bulb and husk on the physicochemical properties of chicken meat. *Poultry Science*, 88(2), pp.398-405.
- Kołodziej-Skalska, A., Rybarczyk, A., Matysiak, B., Jacyno, E., Pietruszka, A., and Kawęcka, M. (2011). Effect of dietary plant extracts mixture on pork meat quality. *Acta Agriculturae Scandinavica, Section A—Animal Science*, 6, 80–85.

- Kotula, K.L. and Wang, Y. (1994). Characterization of broiler meat quality factors as influence by feed withdrawal time. *The Journal of Applied Poultry Research*, 3(2), pp.103-110.
- Kumar, M., Choudhary, R.S. and Vaishnav, J.K. (2005). Effect of supplemental prebiotic, probiotic and turmeric in diet on the performance of broiler chicks during summer. *Indian Journal of Poultry Science*, 40(2), pp.137-141.
- Kwiecien, M. and Mieczan, W. (2009). Effect of addition of herbs on body weight and assessment of physical and chemical alterations in the tibia bones of broiler chickens. *Journal of Elementology*, 14, 705-715.
- Langhout, P. (2000). New additives for broiler chickens. *World Poultry-Elsevier*, 16(3), 22-27.
- Lawrie, R. A. (2005). *Ciência da Carne*. 6 ed. Porto Alegre: Artmed, 384 p.
- Le Bihan-Duval, E., Millet, N. and Remignon, H. (1999). Broiler meat quality: effect of selection for increased carcass quality and estimates of genetic parameters. *Poultry Science*, 78(6), pp.822-826.
- Ledward, D.A. (1985). Post-slaughter influences on the formation of metmyoglobin in beef muscles. *Meat Science*, 15(3), 149-171.
- Lee K.W., Everts, H., Kappert, H.J., Frehner, M., Losa, R., Beynen, A.C. (2003). Effects of dietary essential oil components on growth performance, digestive enzymes and lipid metabolism in female broiler chickens. *British Poultry Science*, 44, 450–457.
- Lee, W.K., Everts, H. and Beynen, A.C. (2004). Essential oils in broiler nutrition. *Internatinal Journal of Poultry Science*, 3:738–752.
- Lestingi, A., Laudadio, V., Marsico, G., Caroli, A. and Vicenti, A. (2004). Effects of dietary safflower oil on growth performance, carcass characteristics and meat chemical and fatty acid composition of wild duck (*Anas platyrhynchos*). *Journal of Food, Agriculture and Environment*, 2, pp.47-50.
- Li, T.S. (1993), July. Evaluation of chemical and non-chemical treatments for the control of ginseng replant disease. In III *International Symposium on Replant Problems*, 363 (pp. 141-146).
- Liu, Y., Lyon, B.G., Windham, W.R., Lyon, C.E. and Savage, E.M. (2004). Principal component analysis of physical, color, and sensory characteristics of chicken breasts deboned at two, four, six, and twenty-four hours postmortem. *Poultry Science*, 83(1), pp.101-108.
- Lo, A.H., Liang, Y.C., Lin-Shiau, S.Y., Ho, C.T. and Lin, J.K. (2002). Carnosol, an antioxidant in rosemary, suppresses inducible nitric oxide synthase through down-regulating nuclear factor- κ B in mouse macrophages. *Carcinogenesis*, 23(6), pp.983-991.
- Lonergan, S.M., Deeb, N., Fedler, C.A. and Lamont, S.J. (2003). Breast meat quality and composition in unique chicken populations. *Poultry Science*, 82(12), pp.1990-1994.
- López, K.P., Schilling, M.W. and Corzo, A. (2011). Broiler genetic strain and sex effects on meat characteristics. *Poultry Science*, 90(5), pp.1105-1111.

- Lopez-Bote, C.J., Gray, J.I., Gomaa, E.A. and Flegal, C.J. (1998). Effect of dietary administration of oil extracts from rosemary and sage on lipid oxidation in broiler meat. *British Poultry Science*, 39(2), pp.235-240.
- Lovland, A. and Kaldhusdal, M., (2001). Severely impaired production performance in broiler flocks with high incidence of *Clostridium perfringens*-associated hepatitis. *Avian Pathology*, 30(1), pp.73-81.
- Luna, A., Labaque, M.C., Zygadlo, J.A. and Marin, R.H. (2010). Effects of thymol and carvacrol feed supplementation on lipid oxidation in broiler meat. *Poultry Science*, 89(2), pp.366-370.
- Lyon, C.E. and Wilson, R.L. (1986). Effects of sex, rigor condition, and heating method on yield and objective texture of broiler breast meat. *Poultry Science*, 65(5), pp.907-914.
- Lyon, C.E., Lyon, B.G., Papa, C.M. and Robach, M.C. (1992). Broiler tenderness: Effects of postchill deboning time and fillet holding time. *The Journal of Applied Poultry Research*, 1(1), pp.27-32.
- Lyon, C.E., Hamm, D., Hudspeth, J.P. and Benoff, F.H. (1983), January. Effects of age and sex of the bird on thaw and cook losses and texture of broiler breast meat. In poultry science (Vol. 62, No. 7, pp. 1459-1459). 1111 NORTH DUNLAP AVE, SAVOY, IL 61874: *Poultry Science Assoc Inc*.
- Madavi, D.L. and Salunkhe, D.K. (1995). Toxicological aspects of food antioxidants. In D. L. Madavi, S. S. Deshpande, & D. K. Salunkhe, Food antioxidants (pp. 267). New York: Marcel Dekker Inc
- Mahan, L.K. and Escott-Stump, S. (2005). Krause, alimentos, nutrição and dietoterapia. Editora Roca.
- Mahmoud, El-Sayed, R., Ibrahim, Doaa. and Badawi, El-Sayed, M. (2013). Effect of supplementation of broiler diet with guava leaves and /or olive oil on growth, meat composition, blood metabolites and immune response. *Benha Veterinary Medical Journal*. Vol.25, No. 2:23-32.
- Maki, A.A. and Froning, G.W. (1987). Effect on the quality characteristics of turkey breast muscle of tumbling whole carcasses in the presence of salt and phosphate. *Poultry Science*, 66(7), pp.1180-1183.
- Makkar, H.P., Blümmel, M., Borowy, N.K. and Becker, K. (1993). Gravimetric determination of tannins and their correlations with chemical and protein precipitation methods. *Journal of the Science of Food and Agriculture*, 61(2), pp.161-165.
- Manzanilla, E.G., Baucells, F., Kamel, C., Morales, J., Perez, J.F. and Gasa, J. (2001). Effects of Plant Extracts on the Performance and Lower Gut Microflora of Early Weaned Piglets. *Journal of Animal Science*, Suppl., 1: 47.
- Mathis, G.F., Hofacre, C. and Scicutella, N. (2007). Performance improvement with a feed added coated blend of essential oils, a coated blend of organic and inorganic acids with essential oils, or virginiamycin in broilers challenged with *Clostridium perfringens*. In *World Poultry Science Association, Proceedings of the 16th European Symposium on Poultry Nutrition*,

Strasbourg, France, 26-30 August, 2007. *World's Poultry Science Association* (WPSA).

- McKenna, D.R., Mies, P.D., Baird, B.E., Pfeiffer, K.D., Ellebracht, J.W. and Savell, J.W. (2005). Biochemical and physical factors affecting discoloration characteristics of 19 bovine muscles. *Meat Science*, 70(4), 665-682.
- McNeal, W.D., and Fletcher, D.L. (2003). Effects of high frequency electrical stunning and decapitation on early rigor development and meat quality of broiler breast meat. *Poultry Science*, 82:1352–1355.
- Mehaffey, J.M., Pradhan, S.P. Meullenet, J.F. Emmert, J.L. Mckee, S.R. and Owens, C.M. (2006). Meat quality evaluation of minimally aged broiler breast fillets from five commercial genetic strains. *Poultry Science*, 85:902–908.
- Meilgaard, M.C., Civille, G.V. and Carr, B.T. (2007). Introduction to sensory techniques. *Sensory Evaluation Techniques*, pp.1-5.
- Miguel, M.G. (2010). Antioxidant activity of medicinal and aromatic plants. A review. *Flavour and Fragrance Journal*, 25(5), pp.291-312.
- Mikulski, D., Jankowski, J., Zdunczyk, Z., Juskiewicz, J. and Slominski, B.A. (2012). The effect of different dietary levels of rapeseed meal on growth performance, carcass traits, and meat quality in turkeys. *Poultry Science*, 91(1), pp.215-223.
- Min D.B., and Boff, J.M. (2003). Crude fat analysis. In: Nielsen SS, editor. *Food analysis*. 3rd ed. New York, New York: Kluwer Academic. P 113-129.
- Mirzaei-Aghsaghali, A. (2012). Importance of medical herbs in animal feeding: A review. *Ann. Biol. Re*, 3, pp.918-923.
- Missohou A, Sow R.S. and Ngwe-Assoumou C. (1997). Caractéristiques morphobiométriques de la poule du Sénégal. Paper presented at INFPD Workshop and General Meeting, 9- 13 December 1997, M'Bour, Senegal.
- Mottaghitlab, M. (2000). "Beneficial effects of garlic (*Allium sativum*) as a growth promoter for broilers and their economic performance." *Proc. of XXI World's Poult. Cong*, 20-24.
- Musa, H.H., Chen, G.H., Cheng, J.H., Li, B.C. and Mekki, D.M. (2006). Study on carcass characteristics of chicken breeds raised under the intensive condition. *International Journal of Poultry Science*, 5(6), pp.530-533.
- Musa, H.H., Chen, G.H., Cheng, J.H., Shuipe, E.S. and Bao, W.B. (2006). Breed and sex effect on meat quality of chicken. *International Journal of Poultry Science*, 5(6), pp.566-568.
- Nakatani, N. (1994). Chemistry of antioxidants from Labiatae herbs. In ACS symposium series (USA). National Research Council, 1988. Designing Food-Animal product options in the market place. Existing technological options and future research needs. P.115. National Academy Press, Washington.D.C.
- Naveena, B.M. and Mendiratta, S.K. (2004). The tenderization of buffalo meat using ginger extract. *Journal Muscle Foods*, 15:235–239.
- Newman, K.E. and Devegowda, G. (1999). Merging modern agriculture with the

- herbal revolution: possibilities for livestock production, what we do and do not know. In Proc. of Alltech's 15th Annu. Symp, Biotechn. In the Feed Industry. Alltech Technical Publications, Nottingham University Press. Nicholasville, KY (pp. 301-306).
- Ngoka, D.A., Froning, G.W., Lowry, S.R. and Babji, A.S. (1982). Effects of sex, age, preslaughter factors, and holding conditions on the quality characteristics and chemical composition of turkey breast muscles. *Poultry Science*, 61(10), pp.1996-2003.
- Ngoupayou, J.D. (1990), October. Country report on small-holder rural poultry production in Cameroon. In CTA seminar proceedings (Vol. 2, pp. 39-47).
- Nielsen, S.E., Young, J.F., Daneshvar, B., Lauridsen, S.T., Knuthsen, P., Sandström, B. and Dragsted, L.O. (1999). Effect of parsley (*Petroselinum crispum*) intake on urinary apigenin excretion, blood antioxidant enzymes and biomarkers for oxidative stress in human subjects. *British Journal of Nutrition*, 81(06), pp.447-455.
- Nigussie, D. (2011). Breeding programs for indigenous chicken in Ethiopia, Analysis of diversity in production systems and chicken populations. PhD Thesis submitted in fulfillment of the requirements for the degree of doctor at Wageningen University Netherlands, 148.
- Niu, Z., Shi, J., Liu, F., Wang, X., Gao, C. and Yao, L. (2009). Effects of dietary energy and protein on growth performance and carcass quality of broilers during starter phase. *International Journal of Poultry Science*, 8(5), pp.508-511.
- Nkukwana, T.T., Muchenje, V., Pieterse, E., Masika, P.J., Mabusela, T.P., Hoffman, L.C. and Dzama, K. (2014). Effect of Moringa oleifera leaf meal on growth performance, apparent digestibility, digestive organ size and carcass yield in broiler chickens. *Livestock Science*, 161, pp.139-146.
- Northcutt, J.K. (2009). Factors Affecting Poultry Meat Quality. University's Bulletin 1157. <http://pubs.caes.uga.edu/caespubs/pubcd/B1157/B1157.htm>. Accessed on March 20th 2009.
- Northcutt, J.K., Buhr, R.J., Young, L.L., Lyon, C.E. and Ware, G.O. (2001). Influence of age and postchill carcass aging duration on chicken breast fillet quality. *Poultry science*, 80(6), pp.808-812.
- Nuernberg, K., Slamecka, J., Mojto, J., Gasparik, J. and Nuernberg, G. (2011). Muscle fat composition of pheasants (*Phasianus colchicus*), wild ducks (*Anas platyrhynchos*) and black coots (*Fulica atra*). *European Journal of Wildlife Research*, 57(4), pp.795-803.
- Nyachoti, C.M., Atkinson, J.L. and Leeson, S. (1997). Sorghum tannins: a review. *World's Poultry Science Journal*, 53(01), pp.5-21.
- Ocak, N., Erener, G., Burak A.k.F., Sungu, M., Altop, A. and Ozmen, A. (2008). Performance of broilers fed diets supplemented with dry peppermint (*Mentha piperita* L.) or thyme (*Thymus vulgaris* L.) leaves as growth promoter source. *Czech Journal of Animal Science*, 53(4), p.169.

- O'Neill, L.M., Galvin, K., Morrissey, P.A. and Buckley, D.J. (1998). Comparison of effects of dietary olive oil, tallow and vitamin E on the quality of broiler meat and meat products. *British Poultry Science*, 39(3), pp.365-371.
- Onibi, G.E., Adebisi, O.E., Fajemisin, A.N. and Adetunji, A.V. (2009). Response of broiler chickens in terms of performance and meat quality to garlic (*Allium sativum*) supplementation. *African Journal of Agricultural Research*, 4(5), pp.511-517.
- Pai, S.T. and Platt, M.W. (1995). Antifungal effects of *Allium sativum* (garlic) extract against the *Aspergillus* species involved in otomycosis. *Letters in Applied Microbiology*, 20(1), pp.14-18.
- Papa, C., LYON, C. and Wilson, R. (1992). Moisture retention and textural properties of ground chicken meat as affected by sodium tripolyphosphate, ionic strength and pH. *Journal of Food Science*, 57(6), pp.1291-1293.
- Papageorgiou, G., Botsoglou, N., Govaris, A., Giannenas, I., Iliadis, S. and Botsoglou, E. (2003). Effect of dietary oregano oil and α -tocopheryl acetate supplementation on iron-induced lipid oxidation of turkey breast, thigh, liver and heart tissues. *Journal of Animal Physiology and Animal Nutrition*, 87(9-10), pp.324-335.
- Parthasarathy, S., Bin Azizi, J., Ramanathan, S., Ismail, S., Sasidharan, S., Said, M. I. M. and Mansor, S. M. (2009). Evaluation of antioxidant and antibacterial activities of aqueous, methanolic and alkaloid extracts from *Mitragyna speciosa* (Rubiaceae family) leaves. *Molecules*, 14(10), pp.3964-3974.
- Paschma, J., and Wawrzynski, M. (2003). Effect of dietary herb supplement for pigs on growth parameters, slaughter traits and dietetic value of pork. *Roczniki Naukowe Zootechniki*, 30(1), 79–88.
- Peng, Y., Yuan, J., Liu, F., Ye, J. (2005). Determination of active components in rosemary by capillary electrophoresis with electrochemical detection. *Journal of Pharmaceutical and Biomedical Analysis* 39, 431–437
- Pilarski, R., Zieliński, H., Ciesiołka, D. and Gulewicz, K. (2006). Antioxidant activity of ethanolic and aqueous extracts of *Uncaria tomentosa* (Willd.) DC. *Journal of Ethnopharmacology*, 104(1), pp.18-23.
- Poole, G.H., Lyon, C. E., Buhr, R J., Young, L.L., Alley, A., Hess, J.B., Bilgili, S.F. and Northcutt, J.K. (1999). Evaluation of age, gender, strain, and diet on the cooked yield and shear values of broiler breast fillets. *The Journal of Applied Poultry Research*, 8(2), pp.170-176.
- Proestos, C., Boziaris, I.S., Nychas, G.J. and Komaitis, M. (2006). Analysis of flavonoids and phenolic acids in Greek aromatic plants: Investigation of their antioxidant capacity and antimicrobial activity. *Food Chemistry*, 95(4), pp.664-671.
- Qiao, M., Fletcher, D.L., Smith, D.P. and Northcutt, J.K. (2001). The effect of broiler breast meat color on pH, moisture, water-holding capacity, and emulsification capacity. *Poultry Science*, 80(5), pp.676-680.
- Raes, K., De Smet, S. and Demeyer, D. (2004). Effect of dietary fatty acids on incorporation of long chain polyunsaturated fatty acids and conjugated

- linoleic acid in lamb, beef and pork meat: a review. *Animal Feed Science and Technology*, 113(1-4), 199-221.
- Rahal, A., Kumar, A., Singh, V., Yadav, B., Tiwari, R., Chakraborty, S. and Dhama, K. (2014). Oxidative stress, prooxidants, and antioxidants: the interplay. *BioMed research international*, 2014.
- Rahayu, H.I., Zulkifli, I., Vidyadaran, M.K., Alimon, A.R. and Babjee, S.A. (2008). Carcass variables and chemical composition of commercial broiler chickens and the red jungle fowl. *Asian-Australasian Journal of Animal Sciences*, 21(9), pp.1376-1382.
- Rahim, A., Aghazadeh, A.M. and Daneshyar, M. (2011). Growth performance and some carcass characteristics in broiler chickens supplemented with Thymus extract (*Thymus vulgaris*) in drinking water. *Journal of American Science*, 7(11), pp.400-405.
- Rahimi, S., Teymori Zadeh, Z., Torshizi, K., Omidbaigi, R. and Rokni, H. (2011). Effect of the three herbal extracts on growth performance, immune system, blood factors and intestinal selected bacterial population in broiler chickens. *Journal of Agricultural Science and Technology*, 13, pp.527-539.
- Rajion, M.A., McLean, J.G. and Cahill, R.N. (1985). Essential fatty acids in the fetal and newborn lamb. *Australian Journal of Biological Sciences*, 38(1), pp.33-40.
- Rice-Evans, C.A., Miller, N.J., Bolwell, P.G., Bramley, P.M. and Pridham, J.B. (1995). The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radical Research*, 22, 375-383.
- Rios, J.L. and Recio, M.C. (2005). Medicinal plants and antimicrobial activity. *Journal of ethnopharmacology*, 100(1), pp.80-84.
- Rondelli, S., Martinez, O. and Garcia, P.T. (2003). Sex effect on productive parameters, carcass and body fat composition of two commercial broilers lines. *Revista Brasileira de Ciência Avícola*, 5(3), pp.169-173.
- Ruberto, G., Baratta, M.T., Sari, M. and Kaâbeche, M. (2002). Chemical composition and antioxidant activity of essential oils from Algerian *Origanum glandulosum* Desf. *Flavour and Fragrance Journal*, 17(4), pp.251-254.
- Sacchetti, G., Maietti, S., Muzzoli, M., Scaglianti, M., Manfredini, S., Radice, M. and Bruni, R. (2005). Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food chemistry*, 91(4), pp.621-632.
- Sahin, K., Orhan, C., Tuzcu, M., Ali, S., Sahin, N. and Hayirli, A. (2010). Epigallocatechin-3-gallate prevents lipid peroxidation and enhances antioxidant defense system via modulating hepatic nuclear transcription factors in heat-stressed quails. *Poultry Science*, 89:2251-2258.
- Sang-Oh, P., Chae-Min, R., Byung-Sung, P. and Jong, H. (2013). The meat quality and growth performance in broiler chickens fed diet with cinnamon powder. *Journal of Environmental Biology*, 34(1), p.127.

- Sarica, S., Ciftci, A., Demir, E., Kilinc, K. and Yildirim, Y. (2005). Use of an antibiotic growth promoter and two herbal natural feed additives with and without exogenous enzymes in wheat based broiler diets. *South African Journal of Animal Science*, 35(1), pp.61-72.
- Sarker, M.S.K., Yim, K.J., Ko, S.Y., Uganbayar, D., Kim, G.M., Bae, I.H., Oh, J. I., Yee, S.T., and Yang, C.J. (2010). Green tea level on growth performance and meat quality in finishing pigs. *Pakistan Journal of Nutrition*, 9, 10–14.
- Sartowska, K.E., Korwin-Kossakowska, A., Polawska, E., Lipinska, P. and Sender, G. (2014). Sex-related differences in the nutritional value of Japanese quail meat. *International Journal of Food Science & Technology*, 49(12), pp.2635-2642.
- SAS. (2007). User's Guide. 9.4 ed. In SAS Inst. Inc, Cary, NC. USA.
- Savell, J.W., Cross, H.R., Francis, J.J., Wise, J.W., Hale, D.S., Wilkes, D.L. and Smith, G.C. (1989). National consumer retail beef study: Interaction of trim level, price and grade on consumer acceptance of beef steaks and roasts. *Journal of Food Quality*, 12(4), pp.251-274.
- Sazili, A.Q., Parr, T., Sensky, P.L., Jones, S.W., Bardsley, R.G. and Buttery, P.J. (2005). The relationship between slow and fast myosin heavy chain content, calpastatin and meat tenderness in different ovine skeletal muscles. *Meat Science*, 69(1), 17-25
- Schilling, M. W., Schilling, J.K., Claus, J. R., Marriott, N.G., Duncan, S.E. and Wang, H. (2003). Instrumental texture assessment and consumer acceptability of cooked broiler breasts evaluated using a geometrically uniform-shaped sample. *Journal of Muscle Foods* 14:11-23.
- Schilling, M.W., Radhakrishnan, V., Thaxton, Y.V., Christensen, K., Thaxton, J.P. and Jackson, V. (2008). The effects of broiler catching method on breast meat quality. *Meat Science*, 79(1), pp.163-171.
- Schneider, B.L., Renema, R.A., Betti, M., Carney, V.L. and Zuidhof, M.J. (2012). Effect of holding temperature, shackling, sex, and age on broiler breast meat quality. *Poultry Science*, 91(2), pp.468-477.
- Schultz, V., Hänsel, R. and Tyler, V.E. (2001). Rational phytotherapy: a physician's guide to herbal medicine. Psychology Press.
- Sebastian, S., Touchburn, S.P., Chavez, E.R. and Lague, P.C. (1996). The effects of supplemental microbial phytase on the performance and utilization of dietary calcium, phosphorus, copper, and zinc in broiler chickens fed corn-soybean diets. *Poultry Science*, 75(6), pp.729-736.
- Shafey, T.M., Al-Mufarej, S., Shalaby, M.I. and Jarelnabi, A.J. (2001). The effect of feeding mannan-oligosaccharides (Bio-Mos) on the performance of meat chickens under two different vaccination programs. *Asian Australasian Journal of Animal Sciences*, 14(4), pp.559-563.
- Sheehy, P.J.A., Morrissey, P.A. and Buckley, D.J. (1995). Advances in research and application of vitamin E as an antioxidant for poultry meat. Pages 425–433 in Poultry Meat Quality. Proc. XII Eur. Symp. Qual. Poult. Meat, Zaragoza, Spain. R. Cepero Briz, ed.

- Simitzis, P.E., Deligeorgis, S.G., Bizelis, J.A., Dardamani, A., Theodosiou, I. and Fegeros, K. (2008). Effect of dietary oregano oil supplementation on lamb meat characteristics. *Meat Science*, 79(2), pp.217-223.
- Sirri, F., Petracci, M., Bianchi, M. and Meluzzi, A. (2010). Survey of skin pigmentation of yellow-skinned broiler chickens. *Poultry science*, 89(7), pp.1556-1561.
- Skrabka-Blotnicka, T., Rosin'ski, A., Przysie-Zzna, E., Woloszyn, J. And Elminowska-Wenda, G. (1997). Effect of Dietary Formulation Supplemented with Herbal Mixture on Goose Breast Muscle Quality. Report I. The Effect on the Chemical Composition. *Archiv fu'r Geflu'gelkunde*, 61(3): 135-138.
- Smet, K., Raes, K., Huyghebaert, G., Haak, L., Arnouts, S. and De Smet, S. (2008). Lipid and protein oxidation of broiler meat as influenced by dietary natural antioxidant supplementation. *Poultry Science*, 87(8), pp.1682-1688.
- Smith, S. B., Smith, D.R. and Lunt, D.K. (2004). Chemical and physical characteristics of meat: adipose tissue. *Encyclopedia of Meat Sciences*, pp.225-238.
- Solomon, M.B., Laack, R.L.J.M. and Eastridge, J.S. (1998). Biophysical basis of pale, soft, exudative (PSE) pork and poultry muscle: A review. *Journal of Muscle Foods*, 9(1), pp.1-11.
- Soltan, M.A., Shewita, R.S. and El-Katcha, M.I. (2008). Effect of dietary anise seeds supplementation on growth performance, immune response, carcass traits and some blood parameters of broiler chickens. *International Journal of Poultry Science*, 7(11), pp.1078-1088.
- Song, J.H. and Miyazawa, T. (2001). Enhanced level of n-3 fatty acid in membrane phospholipids induces lipid peroxidation in rats fed dietary docosahexaenoic acid oil. *Atherosclerosis*, 155(1), pp.9-18.
- Souri, H., Khatibjoo, A., Taherpoor, K., Hassan Abadi, A., Fattahnia, F. and Askari, M. (2015). Effect of *Thymus vulgaris* and *Satureja khuzestanica* ethanolic extracts on broiler chickens' performance and immune response. *Iranian Journal of Applied Animal Science*, 5(2), pp.437-446.
- Souza, P.A., Kodawara, L.M., Pelicano, E.R.L., Souza, H.B.A., Oba, A., Leonel, F. R., Norkus, E.A. and Lima, T.M.A. (2005). Effect of deboning time on the quality of broiler breast meat (*Pectoralis major*). *Revista Brasileira de Ciênciã Avícola*, 7(2), pp.123-128.
- Standard Malaysia. (2009). Halal food-production, preparation, handling and storage- general guidelines (second revision). Department of Standard Malaysia, MS1500:2009.
- Swennen, Q., Everaert, N., Debonne, M., Verbaeys, I., Careghi, C., Tona, K., Janssens, G.P.J., Decuyper, E., Bruggeman, V. and Buyse, J. (2010). Effect of macronutrient ratio of the pre-starter diet on broiler performance and intermediary metabolism. *Journal of Animal Physiology and Animal Nutrition*, 94(3), pp.375-384.
- Symeon, G.K., Athanasiou, A., Lykos, N., Charismiadou, M.A., Goliomytis, M., Demiris, N., Ayoutanti, A., Simitzis, P.E. and Deligeorgis, S. G. (2014). The

- Effects of Dietary Cinnamon (*Cinnamomum Zeylanicum*) Oil Supplementation on Broiler Feeding Behaviour, Growth Performance, Carcass Traits and Meat Quality Characteristics. *Annals of Animal Science*, 14(4), pp.883-895.
- Takayama, H., Kurihara, M., Kitajima, M., Said, I. M. and Aimi, N. (1998). New indole alkaloids from the leaves of Malaysian *Mitragyna speciosa*. *Tetrahedron*, 54(29), pp.8433-8440.
- Tavárez, M.A., Boler, D.D., Bess, K.N., Zhao, J., Yan, F., Dilger, A.C., McKeith, F. K. and Killefer, J. (2011). Effect of antioxidant inclusion and oil quality on broiler performance, meat quality, and lipid oxidation. *Poultry science*, 90(4), pp.922-930.
- Teixeira, A., Pereira, E. and Rodrigues, E.S. (2011). Goat meat quality. Effects of salt airdrying and ageing processes. *Small Ruminant Research*, 98(1), 55-58.
- Thielke S, Lhafi, S.K. and Kühne M. (2005). Effects of Aging Prior to Freezing on Poultry Meat Tenderness. *Poultry Science*, 84:607–612.
- Tjitrosoedirdjo S.S. (1992). *Borreria latifolia* (Aubl.) K. Sch. Weed Info Sheet No. 13. Bogor, Indonesia: The Southeast Asian Weed Information Centre (SEAWIC).
- Toghyani, M., Toghyani, M., Gheisari, A., Ghalamkari, G. and Mohammadrezaei, M. (2010). Growth performance, serum biochemistry and blood hematology of broiler chicks fed different levels of black seed (*Nigella sativa*) and peppermint (*Mentha piperita*). *Livestock Science*, 129(1), pp.173-178.
- Tollba, A.A.H. and M. Hassan, S.H. (2003). Using some natural additives to improve physiological and productive performance of broiler chicks under high temperature condition. 2. Black cumin (*Nigella sativa*) or garlic (*Allium sativum*). *Poultry Science*, 23: 327-340.
- Tsinas, A.C., Giannakopoulos, C.G., Papasteriades, A., Alexopoulos, C., Mavromatis, J. and Kyriakis, S.C. (1998), July. Use of origanum essential oils as growth promoter in pigs. In Proceedings of 15th IPVS Congress, Vol. 3, p. 221.
- Tucker, L. (2002). Botanical broilers: Plant extracts to maintain poultry performance. *Feed International*, 23, pp.26-29.
- Tudor, G.D., Coupar, F.J. and Pethick, D.W. (1996). Effect of silage diets on glycogen concentration in the muscle of yearling cattle. In proceedings-australian society of animal production (Vol. 21, pp. 451-451). *Australian Society of Animal Production*.
- Van Laack, R.L.J.M., Liu, C.H., Smith, M.O. and Loveday, H.D. (2000). Characteristics of Pale, Soft, Exudative broiler breast meat. *Poultry Science*, 79(7), pp.1057-1061.
- Vignola, G., Lambertini, L., Mazzone, G., Giammarco, M., Tassinari, M., Martelli, G. and Bertin, G. (2009). Effects of selenium source and level of supplementation on the performance and meat quality of lambs. *Meat Science*, 81, 678–685.

- Wang, R., Li, D. and Bourne, S. (1998). Can 2000 years of herbal medicine history help us solve problems in the year 2000. *In alltechs annual symposium* (Vol. 14, pp. 168-184).
- Wang, Y.X., Zhan, X.A., Yuan, D., Zhang, X.W. and Wu, R.J. (2011). Effects of selenomethionine and sodium selenite supplementation on meat quality, selenium distribution and antioxidant status in broilers. *Czech Journal of Animal Science*, 56(7), pp.305-313.
- Weber, G.M. and Antipatis, C. (2001). Pork meat quality and dietary vitamin E. In *International Virtual Conference on Pork Quality* (Vol. 2, pp. 122-128).
- Wei, A. and Shibamoto, T. (2007). Antioxidant activities and volatile constituents of various essential oils. *Journal of agricultural and food chemistry*, 55(5), pp.1737-1742.
- Wenk, C. (2000). Herbs, spices and botanicals: 'Old fashioned' or the new feed additives for tomorrow's feed formulations? Concepts for their successful use in biotechnology in the feed industry. Proceedings of Alltech's 16th Annual Symposium (Lyons, TP and Jacques, KA (Eds.). *Nottingham Press, Nottingham: 79-95*.
- Wheeler, B.R., Matthews, N.S., Miller, R.K. and Sams, A.R. (1999). A halothane test to detect turkeys prone to developing pale, soft, and exudative meat. *Poultry science*, 78(11), pp.1634-1638.
- Williams, P. and Losa, R. (2001). The use of essential oils and their compounds in poultry nutrition. *World Poultry*, 17(4), pp.14-15.
- Wojdyło, A., Oszmiański, J. and Czemerys, R. (2007). Antioxidant activity and phenolic compounds in 32 selected herbs. *Food chemistry*, 105(3), pp.940-949.
- Wood, J.D., Enser, M., Fisher, A.V., Nute, G.R., Sheard, P.R., Richardson, R.I., Hughes, S.I. and Whittington, F.M. (2008). Fat deposition, fatty acid composition and meat quality: A review. *Meat Science*, 78(4), 343-358.
- Wulf, D.M., Morgan, J.B., Sanders, S.K., Tatum, J.D., Smith, G.C. & Williams, S. (1995). Effects of dietary supplementation of vitamin E on storage and caselife properties of lamb retail cuts. *Journal of Animal Science*, 73(2), 399-405.
- Yang, C.C., and Chen, T.C. (1993). Effects of refrigerated storage, pH adjustment, and marinade on color of raw and microwave cooked chicken meat. *Poultry Science*, 72:355-362.
- Yanishlieva, N.V., Marinova, E.M., Gordon, M.H. and Raneva, V.G. (1999). Antioxidant activity and mechanism of action of thymol and carvacrol in two lipid systems. *Food Chemistry*, 64(1), pp.59-66.
- Yanishlieva-Maslarova, N.V., Pokorny, J., Yanishlieva, N. and Gordon, M. (2001). Inhibiting oxidation. Antioxidants in food: Practical applications, pp.22-70.
- Yates, J.D., Brunson, C.C. and Webb, J.E. (1976). Relationships of certain biochemical, physical, and quality characteristics of broiler muscles. *Poultry Science*, 55(1), pp.369-378.

- Yin, M.C. and Cheng, W.S. (1998). Antioxidant activity of several Allium members. *Journal Agriculture and Food Chemistry*, 46: 4097–4101.
- Young, J.F., Stagsted, J., Jensen, S.K., Karlsson, A.H. and Henckel, P. (2003). Ascorbic acid, a-tocopherol and oregano supplements reduce stress-induced deterioration of chicken meat quality. *Poultry Science*, 82, 1343–1351.
- Young, L.L., Lyon, C.E., Searcy, G.K. and Wilson, R.L. (1987). Influence of Sodium Tripolyphosphate and Sodium Chloride on Moisture-Retention and Textural Characteristics of Chicken Breast Meat Patties. *Journal of Food Science*, 52(3), pp.571-574.
- Zhang K.Y., Yan F., Keen C.A. and Waldroup P.W. (2005). Evaluation of microencapsulated essential oils and organic acids in diets for broiler chickens. *International Journal of Poultry Science*, 4: 612–619.
- Zhang, G.F., Yang, Z.B., Wang, Y., Yang, W.R., Jiang, S.Z. and Gai, G.S. (2009). Effects of ginger root (*Zingiber officinale*) processed to different particle sizes on growth performance, antioxidant status, and serum metabolites of broiler chickens. *Poultry science*, 88(10), pp.2159-2166.
- Zollitsch, W., Knaus, W., Aichinger, F. and Lettner, F. (1997). Effects of different dietary fat sources on performance and carcass characteristics of broilers. *Animal Feed Science and Technology*, 66(1), pp.63-73.