



UNIVERSITI PUTRA MALAYSIA

***EXPRESSION PROFILING OF GENES RELATED TO ENDOTHELIAL
CELL FUNCTION IN PRE-DIABETES AND TYPE 2 DIABETES SUBJECTS***

SARA MORADIPOOR

FPSK(M) 2016 5



**EXPRESSION PROFILING OF GENES RELATED TO ENDOTHELIAL CELL
FUNCTION IN PRE-DIABETES AND TYPE 2 DIABETES SUBJECTS**

By

SARA MORADIPOOR

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

March 2016



© COPYRIGHT UPM

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Dedicated to:

my beloved parents and siblings,

*for their love, endless support, encouragement and for being a source of motivation
and strength during moments of despair and discouragement.*



© COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of science

EXPRESSION PROFILING OF GENES RELATED TO ENDOTHELIAL CELL FUNCTION IN PRE-DIABETES AND TYPE 2 DIABETES SUBJECTS

By

SARA MORADIPOOR

March 2016

Chairman : Prof. Patimah Ismail, PhD
Faculty : Medicine and Health Sciences

Type 2 diabetes mellitus (DM2) is a major chronic disease with high morbidity, mortality, and economic burden on public health. The term “endothelial dysfunction” refers to the inability of the endothelium to properly maintain vascular homeostasis. Substantial clinical and experimental evidence suggests that endothelial dysfunction occurs in T2DM patient, in both the resistance and conduit vessels of the peripheral circulation as well as in the coronary circulation. Endothelial dysfunction could be the mechanism in explaining the strong association observed between atherosclerotic cardiovascular disease (CVD) and type 2 diabetes mellitus. It has also been shown that patients with prediabetic conditions, such as impaired fasting glucose and impaired glucose tolerance, are at increased risk of cardiovascular disease as well. The first step of the adverse sequence of events that leads to the atherosclerotic process is thought to be endothelial dysfunction. Many genes have already been reported to be related to endothelial dysfunction in type 2 diabetes and prediabetic patients [e.g. endothelial nitric oxide synthase (eNOS)], but the identification of more genes is always a field of further investigation because of the potential finding of new targets for prevention of CVD in type 2 diabetes patients. The aim of this study was to evaluate the expression of a set of genes in peripheral blood associated with endothelial dysfunction in patients with type 2 diabetes mellitus and in those with impaired fasting glycaemia and impaired glucose tolerance (prediabetic patients), trying to find out the relationship between expression of this set of genes and these two pathological conditions. 45 subjects (22 men, 23 women), with a mean age of 48.9 ± 5.71 years, were included in the study. We have recruited the participants after evaluation of fasting glucose and glucose tolerance and HbA1c. According to the results of fasting glucose, glucose tolerance and HbA1c, the participants were divided into three age-matched groups: group 1: diabetes, defined as fasting plasma glucose ≥ 126 mg/dl or 2-h plasma glucose ≥ 200 mg/dl and HbA1c ≥ 6.5 (n=15), group 2: Pre-diabetes, defined as fasting plasma glucose between 110 to 125 mg/dL or IGT 2h post-oral glucose load (75g) between 140mg/dl and 200mg/d and HbA1c between 5.7% to 6.4% (n=15), and group 3 (control group): healthy individuals with normal fasting plasma glucose and normal glucose tolerance (n=15). In patients with type 2 diabetes, we found 59 genes with increased expression. Decreased expression was observed for 4 genes. In pre-diabetes

patients, 2 genes were seen to be significantly down-regulated and 49 genes were significantly up-regulated. Our results indicate that diabetic and pre-diabetic condition may contribute to endothelial dysfunction by disrupting the expression of genes involved in the regulation of endothelial function.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PEMPROFILAN EKSPRESI GEN YANG BERKAITAN DENGAN FUNGSI
SEL ENDOTHELIAL DALAM SAMPEL PRE-DIABETES DAN DIABETES
JENIS 2**

Oleh

SARA MORADIPOOR

Mac 2016

Pengerusi : Prof. Patimah Ismail, PhD, PhD
Fakulti : Perubatan dan Sains Kesihatan

Kencing manis jenis 2 (DM2) merupakan penyakit kronik utama dengan morbiditi, mortaliti, dan beban ekonomi yang tinggi ke atas kesihatan awam. Istilah "disfungsi endotelial" merujuk kepada ketidakupayaan endotelium untuk mengekalkan homeostasis vaskular dengan betul. Bukti klinikal dan eksperimen yang cukup banyak menunjukkan bahawa disfungsi endotelial berlaku kepada pesakit T2DM, terdapat rintangan di dalam kedua-dua saluran pembuluh peredaran periferi dan juga dalam peredaran koronari. Disfungsi endotelial diperhatikan boleh menjadi mekanisme yang menjelaskan perkaitan yang kuat di antara penyakit aterosklerotik kardiovaskular (CVD) dan kencing manis jenis 2. Penyakit ini juga telah menunjukkan bahawa pesakit yang menghidap kencing manis peringkat awal, seperti glukosa berlarap terjejas dan toleransi glukosa terjejas, turut mempunyai risiko peningkatan penyakit kardiovaskular. Langkah awal urutan perkembangan yang berlawanan membawa kepada proses aterosklerotik yang dianggap sebagai disfungsi endotelial. Banyak gen yang telah dilaporkan berkaitan dengan disfungsi endotelial bagi pesakit kencing manis jenis 2 dan pesakit kencing manis peringkat awal [contohnya Endotelial Nitrik Oksida Sintesis (eNOS)], tetapi bagi mengenal pasti lebih banyak gen, kajian lanjut di lapangan sering dijalankan kerana potensi dapatan sasaran baru untuk pencegahan penyakit CVD bagi pesakit kencing manis jenis 2. Tujuan kajian ini adalah untuk menilai ekspresi satu set gen dalam pereferi darah yang berkait dengan disfungsi endotelial pada pesakit kencing manis jenis 2 dan pada pesakit yang mengalami glycaemia berlarap terjejas dan toleransi glukosa terjejas (pesakit kencing manis peringkat awal), dan berusaha untuk mengetahui hubungan antara ekspresi set gen ini dan kedua-dua keadaan patologi tersebut. Seramai 45 pesakit (22 lelaki, 23 wanita), umur purata 48.9 ± 5.71 tahun, telah dilibatkan dalam kajian ini. Kami memilih peserta selepas penilaian glukosa berlarap dan toleransi glukosa dan HbA1c. Berdasarkan kepada hasil penilaian glukosa berlarap, toleransi glukosa dan HbA1c, para peserta dibahagikan kepada tiga kumpulan umur yang dipadankan: Kumpulan 1: kencing manis, ditakrifkan sebagai plasma glukosa berlarap ≥ 126 mg/dl atau 2-h plasma glukosa ≥ 200 mg/dl dan HbA1c ≥ 6.5 (n = 15), kumpulan 2: Pesakit kencing manis peringkat awal, ditakrifkan sebagai plasma glukosa berlarap antara 110 hingga 125mg/dl atau IGT 2h beban glukosa bedah oral (75g) antara 140mg/dl dan 200mg/d

dan HbA1c antara 5.7% hingga 6.4% ($n = 15$), dan kumpulan 3 (kumpulan kawalan): individu yang sihat memiliki plasma glukosa berlapar normal dan toleransi glukosa normal ($n = 15$). Pada pesakit kencing manis jenis 2, kami mendapati 59 gen dengan ekspresi meningkat. Ekspresi menurun diperhatikan untuk 4 gen. Bagi pesakit-pesakit kencing manis peringkat awal, kami mendapati 49 gen dengan ekspresi meningkat. Ekspresi menurun diperhatikan untuk 2 gen. Keputusan kami menunjukkan bahawa keadaan kencing manis dan kencing manis peringkat awal boleh menyebabkan disfungsi endotelial melalui gangguan ekspresi gen yang terlibat dalam peraturan fungsi endotelial.

ACKNOWLEDGEMENTS

I would like to express my sincere gratitude to my supervisor Prof. Dr. Patimah Binti Ismail, whose expertise, understanding, and patience, added considerably to my graduate experience. Her guidance helped me in all the time of research and writing of this thesis. I could not have imagined having a better advisor and mentor for my master study.

I would like to thank the other member of my committee, Dr. Wan Aliaa Binti Wan Sulaiman for the assistance she provided at all levels of the research project. A very special thanks goes out to Dr. Ali Etemad for all the instances in which his assistance helped me along the way.

I thank my friends in the Molecular Biology and Bioinformatics Laboratory, for our exchanges of knowledge, skills, and venting of frustration during my master program, which helped enrich the experience.

I would like to express my sincere gratefulness to my dear father and mother which without their endless love, moral and financial support and encouragement, I would have never been able to complete this important step in my life. Last but not least, I would like to deliver a bouquet of thanks to my best friend Sina for his endless support and kindness.

I certify that a Thesis Examination Committee has met on 09 March 2016 to conduct the final examination of Sara Moradipoor on her thesis entitled "Expression Profiling of Genes Related to Endothelial Cell Function in Pre-Diabetes and Type 2 Diabetes Subjects" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Fauziah bt Othman, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Norshariza binti Nordin, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Internal Examiner)

Srikumar Chakravarthi, PhD

Associate Professor
Graduate School of Medicine
Perdana University Mardi Campus
(External Examiner)



ZULKARNAIN ZAINAL, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 June 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Patimah Binti Ismail, PhD

Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Wan Aliaa Binti Wan Sulaiman

Lecturer
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Member)

BUJANG KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Sara Moradipoor, GS37982

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of
Chairman of
Supervisory
Committee:

Signature: _____

Name of
Member of
Supervisory
Committee:

TABLE OF CONTENTS

		Page
ABSTRACT		i
ABSTRAK		iii
ACKNOWLEDGEMENTS		v
APPROVAL		vi
DECLARATION		viii
LIST OF TABLES		xii
LIST OF FIGURES		xiii
LIST OF APPENDICES		xiv
LIST OF ABBREVIATIONS		xv
CHAPTER		
1	INTRODUCTION	1
	1.1 Research Background	1
	1.2 Statements of Problem	2
	1.3 Hypothesis	2
	1.4 Objectives	3
2	LITERATURE REVIEW	4
	2.1 Definition of Type 2 Diabetes & Pre-diabetes	4
	2.2 Diagnosis	4
	2.3 Pathophysiology	5
	2.3.1 Insulin Resistance	5
	2.3.2 Impaired Insulin Secretion	6
	2.3.3 Hepatic Glucose Production	6
	2.4 Etiology	6
	2.4.1 Genetic Factors	7
	2.4.2 Environmental Factors	7
	2.5 Prevalence	7
	2.6 Endothelial Function	9
	2.7 Endothelial Dysfunction	10
	2.8 Endothelial Dysfunction in Diabetes Mellitus	10
	2.9 Cellular and Molecular Mechanisms of Endothelial Dysfunction in Diabetes	11
	2.9.1 Hyperglycemia	11
	2.9.2 Insulin Resistance	15
	2.9.3 Oxidative Stress	18
	2.9.4 Pro-inflammatory Activation of Endothelium	20
	2.10 RT ² Profiler™ PCR Arrays	21
3	METHODOLOGY	22
	3.1 Study Design	22
	3.2 Ethical Approval	22
	3.3 Methodology flow chart	23
	3.4 Sampling	24
	3.4.1 Study Population	24
	3.4.2 Study Subject	24
	3.4.3 Clinical Measurements	25
	3.4.4 Sampling Location	25

	3.4.5	Sampling Method	26
3.5	RNA		26
	3.5.1	RNA Extraction and Purification	26
	3.5.2	RNA Quantification	27
3.6	cDNA Preparation		27
3.7	RT ² Profiler™ PCR Arrays		28
3.8	Data Analysis		33
4	RESULTS		34
4.1	Subjects		34
	4.1.1	Clinical Measurements	34
	4.1.2	Blood Glucose Level	35
	4.1.3	Chronic Disease	35
4.2	RNA		35
	4.2.1	RNA Quantification and Purity	35
4.3	Expression of Endothelial Cell Biology Genes		36
	4.3.1	Expression of Endothelial Cell Biology Genes in Type 2 Diabetic Group	36
	4.3.2	Expression of Endothelial Cell Biology Genes in Pre-diabetic Group	44
	4.3.3	Direct Comparison of Gene Expression Levels between Group 1 and Group 2	51
5	DISCUSSION		54
5.1	Subjects		54
5.2	Gene Expression Study		54
6	SUMMARY, CONCLUSION, LIMITATION AND RECOMMENDATIONS FOR FUTURE RESEARCH		60
	REFERENCES		62
	APPENDICES		77
	BIODATA OF STUDENT		92
	LIST OF PUBLICATIONS		93

LIST OF TABLES

Table		Page
2.1	Criteria for the diagnosis of diabetes mellitus and pre-diabetes	5
3.1	Subjects' selected criteria for group with type 2 diabetes	24
3.2	Subjects' selected criteria for group with pre-diabetes	25
3.3	Reverse-transcription mix	27
3.4	RT ² Profiler™ PCR Array human endothelial cell biology gene table	30
3.5	PCR components mix	32
3.6	Cycling program	32
3.7	C _T value for The RT2 Profiler™ PCR Arrays control elements	33
4.1	Clinical characteristics of the participants in the different groups	34
4.2	Average of subjects' blood glucose level	35
4.3	Genes with altered expression in group 1	36
4.4	Average Delta (C _T) of genes with altered expression in group 1	41
4.5	Genes with up- and down-regulation, comparing men and women in group 1	43
4.6	Genes with altered expression in group 2	44
4.7	Average Delta (C _T) of genes with altered expression in group 2	49
4.8	Genes with up- and down-regulation, comparing men and women in group 2	51
4.9	Direct comparison of gene expression levels between group 1 and 2	52
5.1	Functional gene grouping	55

LIST OF FIGURES

Figure		Page
2.1	Hyperglycemia leading to endothelial dysfunction	12
2.2	Insulin resistance leading to endothelial dysfunction	16
3.1	Methodology flow chart	23
3.2	RT ² Profiler™ PCR Array format R	29
4.1	Fold regulation in gene expression in T2DM compared to healthy individual	39
4.2	Relative expression comparison for 84 endothelial cells-related gene between type 2 diabetes patients and healthy control	40
4.3	Genes with altered expression comparing men and women in group 1	44
4.4	Genes with altered expression more than 3 fold in group 2	47
4.5	Relative expression comparison for 84 endothelial cells-related gene between pre-diabetes patients and healthy control	48
4.6	Genes with altered expression comparing men and women in group 2	51
4.7	Genes with altered expression more than 3 folds, comparing pre-diabetic group and T2DM group	53

LIST OF APPENDICES

Appendix		Page
1	Ethic approval	77
2	Ethic approval	78
3	Questionnaire form	82
4	Consent form	84
5	Expression levels of all 84 genes evaluated by RT ² Profiler™ PCR Array format R	86



LIST OF ABBREVIATIONS

ADA	American diabetes association
AGE	Advanced glycation end products
AKT	Protein kinase B
ALOX5	Arachidonate 5-lipoxygenase
ANXA5	Annexin A5
AR	Aldose reductase
B2M	Beta-2 microglobulin
BH4	Tetrahydrobiopterin
BMI	Body mass index
°C	Centigrade
Ca	Calcium
CASP1	Caspase-1
CCL2	Chemokine (C-C motif) ligand 2
cDNA	Complementary Deoxyribonucleic acid
CI	confidence interval
COPD	Chronic obstructive pulmonary disease
COX-2	Cyclooxygenase-2
CRP	C-reactive protein
C _T	threshold cycle
CVD	Cardio vascular disease
DAG	Diacylglycerol
DM	Diabetes mellitus
DNA	Deoxyribonucleic acid
EC	Endothelial cell
ECM	Extracellular matrix
EDCFs	Endothelium-derived constricting factors
EDRFs	Endothelium-derived relaxing factors
eNOS	Endothelial nitric oxide synthase
ERK	Extracellular signal-regulated kinases
ET-1	Endothelin-1
F-6-P	Fructose 6-phosphate
FC	Fold change
FFA	Free fatty acids
FPG	Fasting plasma glucose
GFAT	Glucosamine-fructose-6-phosphate aminotransferase
GFPT	Glucosamine-fructose-6-phosphate aminotransferase
GlcNAc	N-acetylglucosamine
GlucN-6-P	Glucosamine-6-phosphate
GlucNAc-1-P	N-acetylglucosamine-6-phosphate
GluN	Glutamine
GLUT1	Glucose transporter 1
HAEC	Human aortic endothelial cell
HbA1c	Glycated haemoglobin
HBP	Hexosamine biosynthetic pathway
HDL	High-density lipoprotein

HG	High glucose
HGEC	Human glomerular endothelial cell
HNF-1 α	Hepatocyte nuclear factor 1 alpha
HUVEC	Human umbilical vein endothelial cells
ICAM	Intercellular adhesion molecule
IFG	Impaired fasting glucose
Ig	Immunoglobulin
IGT	Impaired glucose tolerance
IL-1B	Interleukin 1, beta
IL-1Ra	Interleukin-1 receptor antagonist
IL-6	Interleukin 6
IL7	Interleukin-7
IMR	Institute for Medical Research
iNOS	Inducible nitric oxide synthase
IR	Insulin resistance
IRS	Insulin receptor substrate
ITGA5	Integrin alpha-5
ITGAV	Integrin, alpha V
Kg	Kilogram
L	Litter
LAM	Leukocyte adhesion molecule
LKB1	Liver kinase B1
MAPK	Mitogen activated protein kinase
mg	Milligram
min	Minute
mmol	Millimole
MMP	Matrix metalloproteinase
MREC	Medical Research Ethic Committee
mRNA	Messenger Ribonucleic acid
NADH	Nicotinamide adenine dinucleotide
NADPH	Nicotinamide adenine dinucleotide phosphate
NF- κ B	Nuclear factor-kappa B
ng	Nanogram
NIDDM	Noninsulin-dependent diabetes mellitus
nm	Nanometer
NO	Nitric oxide
NOS3	Nitric oxide synthase 3
nx	plate number of test group
ny	plate number of control group
O ²⁻	superoxide anion
O-GlcNAc	O- linked- N-acetylglucosamine
OGT	O-linked N-acetylglucosamine transferase
OGTT	Oral glucose tolerance test
ONOO ⁻	Peroxynitrite anion
PAI-1	Plasminogen activator inhibitor-1
PBS	Phosphate-buffered saline
PDK-1	pyruvate dehydrogenase kinase, isozyme 1
PECAM1	Platelet endothelial cell adhesion molecule
PGH2	Prostaglandin H2
PI3K	Phosphoinositide 3 kinase

PIP3	Phosphatidylinositol (3,4,5)-trisphosphate
PKC	Protein kinase C
PKC- β	Protein kinase C beta
PLAU	Urokinase-type plasminogen activator
PLG	plasminogen
PPAR- γ	Peroxisome proliferator-active receptor gamma
PTEN	phosphatase and tensin homolog
RAGE	Advanced glycation end products receptor
RCF	Relative centrifugal force
RNA	Ribonucleic acid
ROS	Reactive oxygen species
RPLP0	Ribosomal protein, large, P0
RT-PCR	Reverse transcription polymerase chain reaction
SD	Standard deviation
SDH	Sorbitol dehydrogenase
SELL	Selectin L
SELPLG	Selectin P ligand 1
SMC	Smooth muscle cell
SP1	Specificity protein 1
T2DM	Type 2 diabetes mellitus
TGFB1	Transforming growth factor beta-1
TIMP1	Tissue inhibitor of metalloproteinase
TNF- α	Tumor necrosis factor alpha
TXA2	Thromboxane A2
UCP	Uncoupling protein
UDP-GalNAc	Uridine diphosphate N-acetylgalactosamine
UDP-GlucNAc	Uridine diphosphate N-acetylglucosamine
uPA	Urokinase-type plasminogen activator
UPM	University Putra Malaysia
USA	United states of america
VCAM	Vascular cell adhesion molecule
VEGF	Vascular endothelial growth factor
vWF	von Willibrand's factor
WHR	Waist-hip ratio
μ	Micro
σ_x	standard deviation of Delta Ct of test group
σ_y	standard deviation of Delta(Ct) of control group



© COPYRIGHT UPM

CHAPTER 1

INTRODUCTION

1.1 Research Background

Diabetes is a group of metabolic diseases that are characterized by glucose level, chronically elevated above the normal range. Undiagnosed, or poorly controlled, it can lead to complications such as blindness, renal insufficiency, stroke, and amputation of extremities. Diabetes is associated with a decrease in life expectancy (Malecki, 2005). There are several distinct types of diabetes, but two most common types are type 1 diabetes and type 2 diabetes (T2DM), of which type 2 diabetes constitutes more than 90% of cases of diabetes. The worldwide prevalence of diabetes has been dramatically increasing, particularly in developing countries like India. Current estimates of diabetes suggest that 285 million of the world's adult population (6.4%) is suffering from diabetes. India has 50.8 million and China has 43.2 million diabetics. The largest age group involved is 40–59 years. By 2030 it is projected that 438 million of the world's adult population (7.8%) will be diabetics with major chunk from India and China. The largest age group affected will be between 60 and 79 years. T2DM is caused by complex interactions between adverse environmental factors and certain genetic factors. Several risk factors, including common, easily measurable phenotypic features (such as obesity, hypertension, low HDL cholesterol levels, elevated triglyceride levels and impaired fasting glucose) as well as parental history of diabetes are already used efficiently to predict the development of T2DM (Kota, Meher, Jammula, Kota, & Modi, 2012). According to the American Diabetes Association (ADA), the term 'pre-diabetes' is defined as a metabolic clinical condition able to predispose affected individual to a future development of diabetes (American Diabetes Association, 2014). The endothelium, which is originally considered to be simply a physical barrier between the blood and vascular wall, is now recognized as the most important component of normal vascular homeostasis, because it serves to maintain the anticoagulant, antiplatelet, and fibrinolytic phenotypes of vascular cells (J. Xu & Zou, 2009). The term endothelial dysfunction refers to a condition in which the endothelium loses its physiological properties: the tendency to promote vasodilation, fibrinolysis and antiaggregation (Avogaro, Fadini, Gallo, Pagnin, & de Kreutzenberg, 2006). Endothelial dysfunction plays a critical role in the development of diabetic vasculopathy, which is associated with reduced NO bioavailability resulting from ROS overproduction, lipid peroxidation, and increased generation of adhesion molecules (W. T. Wong, Wong, Tian, & Huang, 2010). Several evidences indicate that pre-diabetes conditions may be associated with an increased cardiovascular risk profile of individuals (Ciccone et al., 2014). Ying Su et al. considered 30 patients with IFG, 38 with IGT, 46 with normal glucose tolerance, and 44 with T2DM and they confirmed that endothelial dysfunction, an early marker of macrovascular disease, is present in subjects with pre-diabetes, indicating endothelial damage in these stages (Su et al., 2008).

1.2 Statements of Problem

The prevalence of diabetes, constituted chiefly by type 2 diabetes (T2D), is a global public health threat. Asian countries contribute to more than 60% of the world's diabetic population as the prevalence of diabetes is increasing in these countries. The prevalence among adults aged 20-70 years is expected to rise from 285 million in 2010 to 438 million by the year 2030. While T2D poses a huge economic burden to all nations, developing countries bear the highest burden since more than 80% of cases occur in these countries. Prevalence estimates of diabetes and impaired glucose tolerance (IGT) are high in all Asian countries and are expected to increase further in the next two decades. The present trend indicates that more than 60% of the world's diabetic population will be in Asia (Guo, He, Zhang, & Walton, 2012). Malaysia has one of the world's highest number of diabetes cases among its population with 2.6 million registered patients (Kearney, Whelton, Reynolds, Whelton, & He, 2004). The total estimated cost of diagnosed diabetes in 2012 is \$245 billion, including \$176 billion in direct medical costs and \$69 billion in reduced productivity. This figure represents a 41 percent increase over a five-year period (\$174 billion in 2007 dollars) (American Diabetes Association, 2013). Micro- and macro-vascular dysfunctions are the major causes of morbidity and mortality in patients with diabetes mellitus (W. T. Wong et al., 2010). Type 2 diabetes is characterized by a two- to fourfold increased risk of cardiovascular disease. It has also been shown that patients with pre-diabetic conditions, such as impaired fasting glucose and impaired glucose tolerance, are at increased risk of cardiovascular disease as well (Kirpichnikov & Sowers, 2001). Pre-diabetic condition is a further expression of incipient atherosclerosis development. The synergy between a systemic inflammatory condition and the presence of high blood glucose concentrations are able to impair vascular endothelium in its function, thus predisposing to atherosclerotic lesions (Ciccione et al., 2014). The first step of the adverse sequence of events that leads to the atherosclerotic process is thought to be endothelial dysfunction (Avogaro et al., 2006).

According to the above studies, diabetes and pre-diabetes are costly diseases with high prevalence worldwide, mostly Asian countries including Malaysia. It has also been shown that patients with type 2 diabetes and patients with pre-diabetes are at increased risk of cardiovascular disease, and endothelial dysfunction is thought to be the most important event that leads to the atherosclerotic process and cardiovascular disease. Therefore, research on expression of endothelial cells-related genes in type 2 diabetes and pre-diabetes patients seems to be important in prevention of atherosclerosis and cardiovascular disease in these metabolic disorders.

1.3 Hypothesis:

H1: The rate of expression of genes regulating the endothelial cells function, is related to endothelial dysfunction in type 2 diabetes patients.

H2: The amount of expression of genes regulating the endothelial cells function, is related to endothelial dysfunction in pre-diabetic condition.

1.4 Objectives:

General Objective:

To determine the expression of genes related to endothelial cells function in peripheral blood of patients with type 2 diabetes, and pre-diabetic patients.

Specific Objectives:

1. To compare the expression level of seven groups of genes – angiogenesis, vasoconstriction & vasodilation, inflammatory response, apoptosis, cell adhesion, coagulation, platelet activation – which are related to endothelial cells function, between type 2 diabetes patients and control.
2. To compare the expression level of seven groups of genes – angiogenesis, vasoconstriction & vasodilation, inflammatory response, apoptosis, cell adhesion, coagulation, platelet activation – which are related to endothelial cells function, between pre-diabetes patients and control.

REFERENCES

- Addabbo, F., Montagnani, M., & Goligorsky, M. S. (2009). Mitochondria and Reactive Oxygen Species. *Hypertension*, 53(6), 885-892. doi: 10.1161/HYPERTENSIONAHA.109.130054
- Adeghate, E., Schattner, P., & Dunn, E. (2006). An update on the etiology and epidemiology of diabetes mellitus. *Ann N Y Acad Sci*, 1084, 1-29. doi: 10.1196/annals.1372.029
- Alexander, J. S., & Elrod, J. W. (2002). Extracellular matrix, junctional integrity and matrix metalloproteinase interactions in endothelial permeability regulation. *J Anat*, 200(6), 561-574.
- Ali, O., Tan, T. T., Sakinah, O., Khalid, B. A., Wu, L. L., & Ng, M. L. (1993). Prevalence of NIDDM and impaired glucose tolerance in aborigines and Malays in Malaysia and their relationship to sociodemographic, health, and nutritional factors. *Diabetes Care*, 16(1), 68-75.
- Alsaigh, T., Pocock, E. S., Bergan, J. J., & Schmid-Schonbein, G. W. (2011). Acute venous occlusion enhances matrix metalloprotease activity: Implications on endothelial dysfunction. *Microvasc Res*, 81(1), 108-116. doi: 10.1016/j.mvr.2010.09.010
- American Diabetes Association. (2013). Economic Costs of Diabetes in the U.S. in 2012. *Diabetes Care*, 36(4), 1033-1046. doi: 10.2337/dc12-2625
- American Diabetes Association. (2014). Diagnosis and classification of diabetes mellitus. *Diabetes Care*, 37 Suppl 1, S81-90. doi: 10.2337/dc14-S081
- Arikawa, E., Quellhorst, G., Han, Y., Pan, H., & Yang, J. (2011). RT2 Profiler™ PCR Arrays: Pathway-focused gene expression profiling with qRT-PCR. *SA Biosciences Technical Article*, 11.
- Aso, Y., Fujiwara, Y., Tayama, K., Takebayashi, K., Inukai, T., & Takemura, Y. (2000). Relationship between soluble thrombomodulin in plasma and coagulation or fibrinolysis in type 2 diabetes. *Clin Chim Acta*, 301(1-2), 135-145.
- Avogaro, A., Fadini, G. P., Gallo, A., Pagnin, E., & de Kreutzenberg, S. (2006). Endothelial dysfunction in type 2 diabetes mellitus. *Nutr Metab Cardiovasc Dis*, 16 Suppl 1, S39-45. doi: 10.1016/j.numecd.2005.10.015
- Bagi, Z., Erdei, N., Toth, A., Li, W., Hintze, T. H., Koller, A., & Kaley, G. (2005). Type 2 diabetic mice have increased arteriolar tone and blood pressure: enhanced release of COX-2-derived constrictor prostaglandins. *Arterioscler Thromb Vasc Biol*, 25(8), 1610-1616. doi: 10.1161/01.ATV.0000172688.26838.9f
- Bakker, W., Eringa, E. C., Sipkema, P., & van Hinsbergh, V. W. (2009). Endothelial dysfunction and diabetes: roles of hyperglycemia, impaired insulin signaling and obesity. *Cell Tissue Res*, 335(1), 165-189. doi: 10.1007/s00441-008-0685-6
- Balletshofer, B. M., Rittig, K., Enderle, M. D., Volk, A., Maerker, E., Jacob, S., . . . Haring, H. U. (2000). Endothelial dysfunction is detectable in young normotensive first-degree relatives of subjects with type 2 diabetes in association with insulin resistance. *Circulation*, 101(15), 1780-1784.
- Banerjee, M., & Vats, P. (2014). Reactive metabolites and antioxidant gene polymorphisms in type 2 diabetes mellitus. *Indian Journal of Human Genetics*, 20(1), 10-19. doi: 10.4103/0971-6866.132747
- Bansal, S., Chawla, D., Siddarth, M., Banerjee, B. D., Madhu, S. V., & Tripathi, A. K. (2013). A study on serum advanced glycation end products and its association

- with oxidative stress and paraoxonase activity in type 2 diabetic patients with vascular complications. *Clinical Biochemistry*, 46(1–2), 109-114. doi: <http://dx.doi.org/10.1016/j.clinbiochem.2012.10.019>
- Bayraktutan, U. (2002). Free radicals, diabetes and endothelial dysfunction. *Diabetes, Obesity and Metabolism*, 4(4), 224-238. doi: 10.1046/j.1463-1326.2002.00184.x
- Beckman, J. A., Creager, M. A., & Libby, P. (2002). Diabetes and atherosclerosis: epidemiology, pathophysiology, and management. *Jama*, 287(19), 2570-2581.
- Beckman, J. A., Goldfine, A. B., Gordon, M. B., Garrett, L. A., & Creager, M. A. (2002). Inhibition of protein kinase C β prevents impaired endothelium-dependent vasodilation caused by hyperglycemia in humans. *Circ Res*, 90(1), 107-111.
- Bonora, E., Kiechl, S., Willeit, J., Oberhollenzer, F., Egger, G., Targher, G., . . . Muggeo, M. (1998). Prevalence of insulin resistance in metabolic disorders: the Bruneck Study. *Diabetes*, 47(10), 1643-1649.
- Brizzi, M. F., Garbarino, G., Rossi, P. R., Pagliardi, G. L., Arduino, C., Avanzi, G. C., & Pegoraro, L. (1993). Interleukin 3 stimulates proliferation and triggers endothelial-leukocyte adhesion molecule 1 gene activation of human endothelial cells. *J Clin Invest*, 91(6), 2887-2892. doi: 10.1172/jci116534
- Brownlee, M. (2001). Biochemistry and molecular cell biology of diabetic complications. *Nature*, 414(6865), 813-820.
- Bullard, K. M., Saydah, S. H., Imperatore, G., Cowie, C. C., Gregg, E. W., Geiss, L. S., . . . Caspersen, C. J. (2013). Secular changes in U.S. Prediabetes prevalence defined by hemoglobin A1c and fasting plasma glucose: National Health and Nutrition Examination Surveys, 1999-2010. *Diabetes Care*, 36(8), 2286-2293. doi: 10.2337/dc12-2563
- Burtea, C., Laurent, S., Murariu, O., Rattat, D., Toubeau, G., Verbruggen, A., . . . Muller, R. N. (2008). Molecular imaging of $\alpha v\beta 3$ integrin expression in atherosclerotic plaques with a mimetic of RGD peptide grafted to Gd-DTPA. *Cardiovascular research*.
- Buse, M. G. (2006). Hexosamines, insulin resistance and the complications of diabetes: current status. *American journal of physiology. Endocrinology and metabolism*, 290(1), E1-E8. doi: 10.1152/ajpendo.00329.2005
- Caballero, A. E., Arora, S., Saouaf, R., Lim, S. C., Smakowski, P., Park, J. Y., . . . Veves, A. (1999). Microvascular and macrovascular reactivity is reduced in subjects at risk for type 2 diabetes. *Diabetes*, 48(9), 1856-1862.
- Cade, W. T. (2008). Diabetes-related microvascular and macrovascular diseases in the physical therapy setting. *Phys Ther*, 88(11), 1322-1335. doi: 10.2522/ptj.20080008
- Calles-Escandon, J., & Cipolla, M. (2001). Diabetes and endothelial dysfunction: a clinical perspective. *Endocr Rev*, 22(1), 36-52. doi: 10.1210/edrv.22.1.0417
- Cameron, N. E., Cotter, M. A., Robertson, S., & Cox, D. (1992). Muscle and nerve dysfunction in rats with experimental galactosaemia. *Exp Physiol*, 77(1), 89-108.
- Candido, R., Bernardi, S., & Allen, T. J. (2009). Linking diabetes and atherosclerosis.
- Canivell, S., & Gomis, R. (2014). Diagnosis and classification of autoimmune diabetes mellitus. *Autoimmun Rev*, 13(4-5), 403-407. doi: 10.1016/j.autrev.2014.01.020
- Ceriello, A. (2003). New insights on oxidative stress and diabetic complications may lead to a "causal" antioxidant therapy. *Diabetes Care*, 26(5), 1589-1596.
- Ceriello, A. (2010). Point: postprandial glucose levels are a clinically important treatment target. *Diabetes Care*, 33(8), 1905-1907. doi: 10.2337/dc10-0634

- Chakravarthy, U., Hayes, R. G., Stitt, A. W., McAuley, E., & Archer, D. B. (1998). Constitutive nitric oxide synthase expression in retinal vascular endothelial cells is suppressed by high glucose and advanced glycation end products. *Diabetes*, *47*(6), 945-952.
- Chan, J. C., Malik, V., Jia, W., Kadowaki, T., Yajnik, C. S., Yoon, K. H., & Hu, F. B. (2009). Diabetes in Asia: epidemiology, risk factors, and pathophysiology. *Jama*, *301*(20), 2129-2140. doi: 10.1001/jama.2009.726
- Chawla, D., Bansal, S., Banerjee, B. D., Madhu, S. V., Kalra, O. P., & Tripathi, A. K. (2014). Role of advanced glycation end product (AGE)-induced receptor (RAGE) expression in diabetic vascular complications. *Microvascular Research*, *95*(0), 1-6. doi: <http://dx.doi.org/10.1016/j.mvr.2014.06.010>
- Chen, S., Khan, Z. A., Cukiernik, M., & Chakrabarti, S. (2003). Differential activation of NF-kappa B and AP-1 in increased fibronectin synthesis in target organs of diabetic complications. *Am J Physiol Endocrinol Metab*, *284*(6), E1089-1097. doi: 10.1152/ajpendo.00540.2002
- Chen, Y. Q., Su, M., Walia, R. R., Hao, Q., Covington, J. W., & Vaughan, D. E. (1998). Sp1 sites mediate activation of the plasminogen activator inhibitor-1 promoter by glucose in vascular smooth muscle cells. *J Biol Chem*, *273*(14), 8225-8231.
- Ciccone, M. M., Scicchitano, P., Cameli, M., Cecere, A., Cortese, F., Dentamaro, I., . . . Modesti, P. A. (2014). Endothelial function in pre-diabetes, diabetes and diabetic cardiomyopathy: a review. *J Diabetes Metab*, *5*(364), 2.
- Cifarelli, V., Geng, X., Styche, A., Lakomy, R., Trucco, M., & Luppi, P. (2011). C-peptide reduces high-glucose-induced apoptosis of endothelial cells and decreases NAD(P)H-oxidase reactive oxygen species generation in human aortic endothelial cells. *Diabetologia*, *54*(10), 2702-2712. doi: 10.1007/s00125-011-2251-0
- Cipollone, F., Mezzetti, A., Fazio, M. L., Cuccurullo, C., Iezzi, A., Uchino, S., . . . Stafforini, D. M. (2005). Association between 5-lipoxygenase expression and plaque instability in humans. *Arterioscler Thromb Vasc Biol*, *25*(8), 1665-1670. doi: 10.1161/01.ATV.0000172632.96987.2d
- Cosentino, F., Eto, M., De Paolis, P., van der Loo, B., Bachschmid, M., Ullrich, V., . . . Luscher, T. F. (2003). High glucose causes upregulation of cyclooxygenase-2 and alters prostanoid profile in human endothelial cells: role of protein kinase C and reactive oxygen species. *Circulation*, *107*(7), 1017-1023.
- Cotter, M. A., Cameron, N. E., & Robertson, S. (1992). Polyol pathway-mediated changes in cardiac muscle contractile properties: studies in streptozotocin-diabetic and galactose-fed rats. *Exp Physiol*, *77*(6), 829-838.
- Cotter, M. A., Jack, A. M., & Cameron, N. E. (2002). Effects of the protein kinase C beta inhibitor LY333531 on neural and vascular function in rats with streptozotocin-induced diabetes. *Clin Sci (Lond)*, *103*(3), 311-321. doi: 10.1042/
- Cusi, K., Maezono, K., Osman, A., Pendergrass, M., Patti, M. E., Pratipanawatr, T., . . . Mandarino, L. J. (2000). Insulin resistance differentially affects the PI 3-kinase- and MAP kinase-mediated signaling in human muscle. *Journal of Clinical Investigation*, *105*(3), 311-320.
- D'Adamo, E., & Caprio, S. (2011). Type 2 diabetes in youth: epidemiology and pathophysiology. *Diabetes Care*, *34 Suppl 2*, S161-165. doi: 10.2337/dc11-s212

- Dagher, Z., Park, Y. S., Asnaghi, V., Hoehn, T., Gerhardinger, C., & Lorenzi, M. (2004). Studies of rat and human retinas predict a role for the polyol pathway in human diabetic retinopathy. *Diabetes*, *53*(9), 2404-2411.
- de Carvalho Vidigal, F., Guedes Cocate, P., Goncalves Pereira, L., & de Cassia Goncalves Alfenas, R. (2012). The role of hyperglycemia in the induction of oxidative stress and inflammatory process. *Nutr Hosp*, *27*(5), 1391-1398. doi: 10.3305/nh.2012.27.5.5917
- de la Sierra, A., & Larrousse, M. (2010). Endothelial dysfunction is associated with increased levels of biomarkers in essential hypertension. *J Hum Hypertens*, *24*(6), 373-379.
- De Vriese, A. S., Verbeuren, T. J., Van de Voorde, J., Lameire, N. H., & Vanhoutte, P. M. (2000). Endothelial dysfunction in diabetes. *British Journal of Pharmacology*, *130*(5), 963-974. doi: 10.1038/sj.bjp.0703393
- Death, A. K., Fisher, E. J., McGrath, K. C. Y., & Yue, D. K. (2003). High glucose alters matrix metalloproteinase expression in two key vascular cells: potential impact on atherosclerosis in diabetes. *Atherosclerosis*, *168*(2), 263-269. doi: [http://dx.doi.org/10.1016/S0021-9150\(03\)00140-0](http://dx.doi.org/10.1016/S0021-9150(03)00140-0)
- DeFronzo, R. A. (2004). Pathogenesis of type 2 diabetes mellitus. *Med Clin North Am*, *88*(4), 787-835, ix. doi: 10.1016/j.mcna.2004.04.013
- Derosa, G., D'Angelo, A., Tinelli, C., Devangelio, E., Consoli, A., Miccoli, R., . . . Cicero, A. F. (2007). Evaluation of metalloproteinase 2 and 9 levels and their inhibitors in diabetic and healthy subjects. *Diabetes Metab*, *33*(2), 129-134. doi: 10.1016/j.diabet.2006.11.008
- Dimaio, T. A., Wang, S., Huang, Q., Scheef, E. A., Sorenson, C. M., & Sheibani, N. (2008). Attenuation of retinal vascular development and neovascularization in PECAM-1-deficient mice. *Dev Biol*, *315*(1), 72-88. doi: 10.1016/j.ydbio.2007.12.008
- Ding, H., Aljofan, M., & Triggle, C. R. (2007). Oxidative stress and increased eNOS and NADPH oxidase expression in mouse microvessel endothelial cells. *Journal of cellular physiology*, *212*(3), 682-689. doi: 10.1002/jcp.21063
- Diogo, C. V., Suski, J. M., Lebieczinska, M., Karkucinska-Wieckowska, A., Wojtala, A., Pronicki, M., . . . Wieckowski, M. R. (2013). Cardiac mitochondrial dysfunction during hyperglycemia--the role of oxidative stress and p66Shc signaling. *Int J Biochem Cell Biol*, *45*(1), 114-122. doi: 10.1016/j.biocel.2012.07.004
- Dosi, R., Acharya, D., & Patell, R. (2012). Endothelial dysfunction in a cohort of Indian patients with type-2 diabetes mellitus. *Journal, Indian Academy of Clinical Medicine*, *12*(3), 206-209.
- Drzewoski, J., Sliwinska, A., Przybyłowska, K., Sliwinski, T., Kasznicki, J., Zurawska-Klis, M., . . . Majsterek, I. (2008). Gene polymorphisms and antigen levels of matrix metalloproteinase-1 in type 2 diabetes mellitus coexisting with coronary heart disease. *Kardiol Pol*, *66*(10), 1042-1048; discussion 1049.
- Du, X. L., Edelstein, D., Dimmeler, S., Ju, Q., Sui, C., & Brownlee, M. (2001). Hyperglycemia inhibits endothelial nitric oxide synthase activity by posttranslational modification at the Akt site. *J Clin Invest*, *108*(9), 1341-1348. doi: 10.1172/jci11235
- Du, X. L., Edelstein, D., Rossetti, L., Fantus, I. G., Goldberg, H., Ziyadeh, F., . . . Brownlee, M. (2000). Hyperglycemia-induced mitochondrial superoxide overproduction activates the hexosamine pathway and induces plasminogen activator inhibitor-1 expression by increasing Sp1 glycosylation. *Proc Natl Acad Sci U S A*, *97*(22), 12222-12226. doi: 10.1073/pnas.97.22.12222

- Eriksson, J. W. (2007). Metabolic stress in insulin's target cells leads to ROS accumulation - a hypothetical common pathway causing insulin resistance. *FEBS Lett*, 581(19), 3734-3742. doi: 10.1016/j.febslet.2007.06.044
- Evcimen, N. D., & King, G. L. (2007). The role of protein kinase C activation and the vascular complications of diabetes. *Pharmacological research*, 55(6), 498-510.
- Ewing, M. M., de Vries, M. R., Nordzell, M., Pettersson, K., de Boer, H. C., van Zonneveld, A. J., . . . Quax, P. H. (2011). Annexin A5 therapy attenuates vascular inflammation and remodeling and improves endothelial function in mice. *Arterioscler Thromb Vasc Biol*, 31(1), 95-101. doi: 10.1161/atvbaha.110.216747
- Faerch, K., & Vistisen, D. (2014). Prevalence of diabetes in Chinese adults. *Jama*, 311(2), 200. doi: 10.1001/jama.2013.283127
- Falkenberg, M., Tom, C., DeYoung, M. B., Wen, S., Linnemann, R., & Dichek, D. A. (2002). Increased expression of urokinase during atherosclerotic lesion development causes arterial constriction and lumen loss, and accelerates lesion growth. *Proceedings of the National Academy of Sciences*, 99(16), 10665-10670.
- Fantus, I. G., Goldberg, H., Whiteside, C., & Topic, D. (2006). The Hexosamine Biosynthesis Pathway. In P. Cortes & C. Mogensen (Eds.), *The Diabetic Kidney* (pp. 117-133): Humana Press.
- Federici, M., Menghini, R., Mauriello, A., Hribal, M. L., Ferrelli, F., Lauro, D., . . . Lauro, R. (2002). Insulin-dependent activation of endothelial nitric oxide synthase is impaired by O-linked glycosylation modification of signaling proteins in human coronary endothelial cells. *Circulation*, 106(4), 466-472.
- Federici, M., Pandolfi, A., De Filippis, E. A., Pellegrini, G., Menghini, R., Lauro, D., . . . Consoli, A. (2004). G972R IRS-1 variant impairs insulin regulation of endothelial nitric oxide synthase in cultured human endothelial cells. *Circulation*, 109(3), 399-405. doi: 10.1161/01.cir.0000109498.77895.6f
- Félétou, M. (2011). The endothelium: Part 1: multiple functions of the endothelial cells—focus on endothelium-derived vasoactive mediators.
- Fernández-Velasco, M., Ruiz-Hurtado, G., Gómez, A. M., & Rueda, A. (2014). Ca²⁺-handling alterations and vascular dysfunction in diabetes. *Cell Calcium*, 56(5), 397-407. doi: <http://dx.doi.org/10.1016/j.ceca.2014.08.007>
- Folsom, A. R., Aleksic, N., Park, E., Salomaa, V., Juneja, H., & Wu, K. K. (2001). Prospective study of fibrinolytic factors and incident coronary heart disease: the Atherosclerosis Risk in Communities (ARIC) Study. *Arterioscler Thromb Vasc Biol*, 21(4), 611-617.
- Forster, H. G., ter Wee, P. M., Hohman, T. C., & Epstein, M. (1996). Impairment of afferent arteriolar myogenic responsiveness in the galactose-fed rat is prevented by tolrestat. *Diabetologia*, 39(8), 907-914.
- Gerich, J. E. (2002). Is reduced first-phase insulin release the earliest detectable abnormality in individuals destined to develop type 2 diabetes? *Diabetes*, 51 Suppl 1, S117-121.
- Giacco, F., & Brownlee, M. (2010). Oxidative stress and diabetic complications. *Circ Res*, 107(9), 1058-1070. doi: 10.1161/CIRCRESAHA.110.223545
- Gogg, S., Smith, U., & Jansson, P. A. (2009). Increased MAPK activation and impaired insulin signaling in subcutaneous microvascular endothelial cells in type 2 diabetes: the role of endothelin-1. *Diabetes*, 58(10), 2238-2245. doi: 10.2337/db08-0961

- Gradinaru, D., Borsa, C., Ionescu, C., & Margina, D. (2013). Advanced oxidative and glycoxidative protein damage markers in the elderly with type 2 diabetes. *Journal of proteomics*, 92, 313-322. doi: 10.1016/j.jprot.2013.03.034
- Grassi, D., Desideri, G., Necozione, S., Ruggieri, F., Blumberg, J. B., Stornello, M., & Ferri, C. (2012). Protective effects of flavanol-rich dark chocolate on endothelial function and wave reflection during acute hyperglycemia. *Hypertension*, 60(3), 827-832. doi: 10.1161/hypertensionaha.112.193995
- Guariguata, L., Whiting, D. R., Hambleton, I., Beagley, J., Linnenkamp, U., & Shaw, J. E. (2014). Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes Res Clin Pract*, 103(2), 137-149. doi: <http://dx.doi.org/10.1016/j.diabres.2013.11.002>
- Guo, F., He, D., Zhang, W., & Walton, R. G. (2012). Trends in prevalence, awareness, management, and control of hypertension among United States adults, 1999 to 2010. *J Am Coll Cardiol*, 60(7), 599-606. doi: 10.1016/j.jacc.2012.04.026
- Guzik, T. J., Mussa, S., Gastaldi, D., Sadowski, J., Ratnatunga, C., Pillai, R., & Channon, K. M. (2002). Mechanisms of increased vascular superoxide production in human diabetes mellitus: role of NAD(P)H oxidase and endothelial nitric oxide synthase. *Circulation*, 105(14), 1656-1662.
- Hadi, H. A. R., & Suwaidi, J. A. (2007). Endothelial dysfunction in diabetes mellitus. *Vascular Health and Risk Management*, 3(6), 853-876.
- Haffner, S. M., Lehto, S., Ronnema, T., Pyorala, K., & Laakso, M. (1998). Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med*, 339(4), 229-234. doi: 10.1056/nejm199807233390404
- Hamburg, N. M., Larson, M. G., Vita, J. A., Vasan, R. S., Keyes, M. J., Widlansky, M. E., . . . Meigs, J. B. (2008). Metabolic syndrome, insulin resistance, and brachial artery vasodilator function in Framingham Offspring participants without clinical evidence of cardiovascular disease. *Am J Cardiol*, 101(1), 82-88.
- Hamilton, S. J., Chew, G. T., & Watts, G. F. (2007). Therapeutic regulation of endothelial dysfunction in type 2 diabetes mellitus. *Diab Vasc Dis Res*, 4(2), 89-102. doi: 10.3132/dvdr.2007.026
- Hart, G. W., Housley, M. P., & Slawson, C. (2007). Cycling of O-linked [beta]-N-acetylglucosamine on nucleocytoplasmic proteins. *Nature*, 446(7139), 1017-1022.
- Henquin, J. C., Ishiyama, N., Nenquin, M., Ravier, M. A., & Jonas, J. C. (2002). Signals and pools underlying biphasic insulin secretion. *Diabetes*, 51 Suppl 1, S60-67.
- Henriksen, J. E., Alford, F., Handberg, A., Vaag, A., Ward, G. M., Kalfas, A., & Beck-Nielsen, H. (1994). Increased glucose effectiveness in normoglycemic but insulin-resistant relatives of patients with non-insulin-dependent diabetes mellitus. A novel compensatory mechanism. *J Clin Invest*, 94(3), 1196-1204. doi: 10.1172/jci117436
- Herman, W. H., & Zimmet, P. (2012). Type 2 diabetes: an epidemic requiring global attention and urgent action. *Diabetes Care*, 35(5), 943-944. doi: 10.2337/dc12-0298
- Hink, U., Li, H., Mollnau, H., Oelze, M., Matheis, E., Hartmann, M., . . . Munzel, T. (2001). Mechanisms underlying endothelial dysfunction in diabetes mellitus. *Circ Res*, 88(2), E14-22.

- Hirose, A., Tanikawa, T., Mori, H., Okada, Y., & Tanaka, Y. (2010). Advanced glycation end products increase endothelial permeability through the RAGE/Rho signaling pathway. *FEBS Letters*, *584*(1), 61-66. doi: <http://dx.doi.org/10.1016/j.febslet.2009.11.082>
- Hoshiyama, M., Li, B., Yao, J., Harada, T., Morioka, T., & Oite, T. (2003). Effect of high glucose on nitric oxide production and endothelial nitric oxide synthase protein expression in human glomerular endothelial cells. *Nephron Exp Nephrol*, *95*(2), e62-68. doi: 73673
- Huang, A. L., & Vita, J. A. (2006). Effects of systemic inflammation on endothelium-dependent vasodilation. *Trends Cardiovasc Med*, *16*(1), 15-20. doi: 10.1016/j.tcm.2005.10.002
- Huang, Y., Liu, X. Q., Rall, S. C., Jr., Taylor, J. M., von Eckardstein, A., Assmann, G., & Mahley, R. W. (1998). Overexpression and accumulation of apolipoprotein E as a cause of hypertriglyceridemia. *J Biol Chem*, *273*(41), 26388-26393.
- Hussain, A., Claussen, B., Ramachandran, A., & Williams, R. (2007). Prevention of type 2 diabetes: a review. *Diabetes Res Clin Pract*, *76*(3), 317-326. doi: 10.1016/j.diabres.2006.09.020
- Inoguchi, T., Battan, R., Handler, E., Sportsman, J. R., Heath, W., & King, G. L. (1992). Preferential elevation of protein kinase C isoform beta II and diacylglycerol levels in the aorta and heart of diabetic rats: differential reversibility to glycemic control by islet cell transplantation. *Proc Natl Acad Sci U S A*, *89*(22), 11059-11063.
- Inoguchi, T., Li, P., Umeda, F., Yu, H. Y., Kakimoto, M., Imamura, M., . . . Nawata, H. (2000). High glucose level and free fatty acid stimulate reactive oxygen species production through protein kinase C--dependent activation of NAD(P)H oxidase in cultured vascular cells. *Diabetes*, *49*(11), 1939-1945.
- Kanters, S. D., Banga, J. D., Algra, A., Frijns, R. C., Beutler, J. J., & Fijnheer, R. (2001). Plasma levels of cellular fibronectin in diabetes. *Diabetes Care*, *24*(2), 323-327.
- Kassab, A., & Piwowar, A. (2012). Cell oxidant stress delivery and cell dysfunction onset in type 2 diabetes. *Biochimie*, *94*(9), 1837-1848. doi: <http://dx.doi.org/10.1016/j.biochi.2012.01.020>
- Kearney, P. M., Whelton, M., Reynolds, K., Whelton, P. K., & He, J. (2004). Worldwide prevalence of hypertension: a systematic review. *J Hypertens*, *22*(1), 11-19.
- Kenichi, M., Masanobu, M., Takehiko, K., Shoko, T., Akira, F., Katsushige, A., . . . Shigeru, K. (2004). Renal synthesis of urokinase type-plasminogen activator, its receptor, and plasminogen activator inhibitor-1 in diabetic nephropathy in rats: modulation by angiotensin-converting-enzyme inhibitor. *J Lab Clin Med*, *144*(2), 69-77. doi: 10.1016/j.lab.2004.04.002
- Khew-Goodall, Y., Butcher, C. M., Litwin, M. S., Newlands, S., Korpelainen, E. I., Noack, L. M., . . . Vadas, M. A. (1996). Chronic expression of P-selectin on endothelial cells stimulated by the T-cell cytokine, interleukin-3. *Blood*, *87*(4), 1432-1438.
- Kienast, J., Padro, T., Steins, M., Li, C. X., Schmid, K. W., Hammel, D., . . . van de Loo, J. C. (1998). Relation of urokinase-type plasminogen activator expression to presence and severity of atherosclerotic lesions in human coronary arteries. *Thromb Haemost*, *79*(3), 579-586.
- Kikuchi, Y., Imakiire, T., Hyodo, T., Kushiyama, T., Higashi, K., Hyodo, N., . . . Miura, S. (2005). Advanced glycation end-product induces fractalkine gene

- upregulation in normal rat glomeruli. *Nephrol Dial Transplant*, 20(12), 2690-2696. doi: 10.1093/ndt/gfi232
- Kim, J. A., Montagnani, M., Koh, K. K., & Quon, M. J. (2006). Reciprocal relationships between insulin resistance and endothelial dysfunction: molecular and pathophysiological mechanisms. *Circulation*, 113(15), 1888-1904. doi: 10.1161/circulationaha.105.563213
- Kirpichnikov, D., & Sowers, J. R. (2001). Diabetes mellitus and diabetes-associated vascular disease. *Trends Endocrinol Metab*, 12(5), 225-230.
- Kizub, I. V., Klymenko, K. I., & Soloviev, A. I. (2014). Protein kinase C in enhanced vascular tone in diabetes mellitus. *Int J Cardiol*, 174(2), 230-242. doi: 10.1016/j.ijcard.2014.04.117
- Kohei, K. (2010). Pathophysiology of type 2 diabetes and its treatment policy. *Jpn Med Assoc*, 53, 41-46.
- Kolluru, G. K., Bir, S. C., & Kevil, C. G. (2012). Endothelial dysfunction and diabetes: effects on angiogenesis, vascular remodeling, and wound healing. *International journal of vascular medicine*, 2012.
- Korpelainen, E. I., Gamble, J. R., Smith, W. B., Goodall, G. J., Qiyu, S., Woodcock, J. M., . . . Lopez, A. F. (1993). The receptor for interleukin 3 is selectively induced in human endothelial cells by tumor necrosis factor alpha and potentiates interleukin 8 secretion and neutrophil transmigration. *Proc Natl Acad Sci U S A*, 90(23), 11137-11141.
- Korshunov, S. S., Skulachev, V. P., & Starkov, A. A. (1997). High protonic potential actuates a mechanism of production of reactive oxygen species in mitochondria. *FEBS Letters*, 416(1), 15-18.
- Kota, S. K., Meher, L. K., Jammula, S., Kota, S. K., & Modi, K. D. (2012). Genetics of type 2 diabetes mellitus and other specific types of diabetes; its role in treatment modalities. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 6(1), 54-58. doi: <http://dx.doi.org/10.1016/j.dsx.2012.05.014>
- Kouroedov, A., Eto, M., Joch, H., Volpe, M., Lüscher, T. F., & Cosentino, F. (2004). Selective inhibition of protein kinase C β 2 prevents acute effects of high glucose on vascular cell adhesion molecule-1 expression in human endothelial cells. *Circulation*, 110(1), 91-96.
- Kowluru, R. A. (2010). Role of matrix metalloproteinase-9 in the development of diabetic retinopathy and its regulation by H-Ras. *Invest Ophthalmol Vis Sci*, 51(8), 4320-4326. doi: 10.1167/iovs.09-4851
- Kuboki, K., Jiang, Z. Y., Takahara, N., Ha, S. W., Igarashi, M., Yamauchi, T., . . . King, G. L. (2000). Regulation of endothelial constitutive nitric oxide synthase gene expression in endothelial cells and in vivo : a specific vascular action of insulin. *Circulation*, 101(6), 676-681.
- Kumar, A., Bharti, S. K., & Kumar, A. (2014). Type 2 Diabetes Mellitus: The Concerned Complications and Target Organs. *Apollo Medicine*, 11(3), 161-166. doi: <http://dx.doi.org/10.1016/j.apme.2014.01.009>
- Laughlin, M. H., Newcomer, S. C., & Bender, S. B. (2008). Importance of hemodynamic forces as signals for exercise-induced changes in endothelial cell phenotype. *Journal of Applied Physiology*, 104(3), 588-600.
- Leiva, E., Mujica, V., Orrego, R., Prieto, M., & Arredondo, M. (2005). Apolipoprotein E polymorphism in type 2 diabetic patients of Talca, Chile. *Diabetes Res Clin Pract*, 68(3), 244-249. doi: 10.1016/j.diabetes.2004.09.017
- Li, L., Crockett, E., Wang, D. H., Galligan, J. J., Fink, G. D., & Chen, A. F. (2002). Gene transfer of endothelial NO synthase and manganese superoxide dismutase on arterial vascular cell adhesion molecule-1 expression and

- superoxide production in deoxycorticosterone acetate-salt hypertension. *Arterioscler Thromb Vasc Biol*, 22(2), 249-255.
- Lu, X., Bean, J. S., Kassab, G. S., & Rekhter, M. D. (2011). Protein kinase C inhibition ameliorates functional endothelial insulin resistance and vascular smooth muscle cell hypersensitivity to insulin in diabetic hypertensive rats. *Cardiovasc Diabetol*, 10, 48. doi: 10.1186/1475-2840-10-48
- Ma, H., Zhang, H. F., Yu, L., Zhang, Q. J., Li, J., Huo, J. H., . . . Gao, F. (2006). Vasculoprotective effect of insulin in the ischemic/reperfused canine heart: role of Akt-stimulated NO production. *Cardiovasc Res*, 69(1), 57-65. doi: 10.1016/j.cardiores.2005.08.019
- Mainous, A. G., 3rd, Tanner, R. J., Baker, R., Zayas, C. E., & Harle, C. A. (2014). Prevalence of prediabetes in England from 2003 to 2011: population-based, cross-sectional study. *BMJ Open*, 4(6), e005002. doi: 10.1136/bmjopen-2014-005002
- Malecki, M. T. (2005). Genetics of type 2 diabetes mellitus. *Diabetes Res Clin Pract*, 68, Supplement 1(0), S10-S21. doi: <http://dx.doi.org/10.1016/j.diabres.2005.03.003>
- Matsumoto, T., Ishida, K., Nakayama, N., Kobayashi, T., & Kamata, K. (2009). Involvement of NO and MEK/ERK pathway in enhancement of endothelin-1-induced mesenteric artery contraction in later-stage type 2 diabetic Goto-Kakizaki rat. *Am J Physiol Heart Circ Physiol*, 296(5), H1388-1397. doi: 10.1152/ajpheart.00043.2009
- McVeigh, G. E., Brennan, G. M., Johnston, G. D., McDermott, B. J., McGrath, L. T., Henry, W. R., . . . Hayes, J. R. (1992). Impaired endothelium-dependent and independent vasodilation in patients with type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia*, 35(8), 771-776.
- Mehta, N. N., Sheetz, M., Price, K., Comiskey, L., Amrutia, S., Iqbal, N., . . . Reilly, M. P. (2009). Selective PKC beta inhibition with ruboxistaurin and endothelial function in type-2 diabetes mellitus. *Cardiovasc Drugs Ther*, 23(1), 17-24. doi: 10.1007/s10557-008-6144-5
- Michiels, C. (2003). Endothelial cell functions. *Journal of cellular physiology*, 196(3), 430-443.
- Mizutani, M., Kern, T. S., & Lorenzi, M. (1996). Accelerated death of retinal microvascular cells in human and experimental diabetic retinopathy. *J Clin Invest*, 97(12), 2883-2890. doi: 10.1172/jci118746
- Moore, R., Hawley, A., Sigler, R., Farris, D., Wroblewski, S., Ramacciotti, E., & Myers, D. (2009). Tissue Inhibitor of Metalloproteinase-1 is an Early Marker of Acute Endothelial Dysfunction in a Rodent Model of Venous Oxidative Injury. *Annals of Vascular Surgery*, 23(4), 498-505. doi: <http://dx.doi.org/10.1016/j.avsg.2009.03.005>
- Morigi, M., Angioletti, S., Imberti, B., Donadelli, R., Micheletti, G., Figliuzzi, M., . . . Remuzzi, G. (1998). Leukocyte-endothelial interaction is augmented by high glucose concentrations and hyperglycemia in a NF-kB-dependent fashion. *Journal of Clinical Investigation*, 101(9), 1905-1915.
- Movva, L. R., Ho, D. K., Corbet, E. F., & Leung, W. K. (2014). Type-2 diabetes mellitus, metabolic control, serum inflammatory factors, lifestyle, and periodontal status. *Journal of Dental Sciences*, 9(1), 1-9.
- Muniyappa, R., & Sowers, J. R. (2013). Role of insulin resistance in endothelial dysfunction. *Rev Endocr Metab Disord*, 14(1), 5-12. doi: 10.1007/s11154-012-9229-1

- Nangle, M. R., Cotter, M. A., & Cameron, N. E. (2003). Protein kinase C beta inhibition and aorta and corpus cavernosum function in streptozotocin-diabetic mice. *Eur J Pharmacol*, 475(1-3), 99-106.
- Ning, A., Cui, J., Maberley, D., Ma, P., & Matsubara, J. (2008). Expression of integrins in human proliferative diabetic retinopathy membranes. *Canadian Journal of Ophthalmology/Journal Canadien d'Ophthalmologie*, 43(6), 683-688.
- Nishikawa, T., Edelstein, D., Du, X. L., Yamagishi, S., Matsumura, T., Kaneda, Y., . . . Brownlee, M. (2000). Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. *Nature*, 404(6779), 787-790. doi: 10.1038/35008121
- Nolan, C. J., Damm, P., & Prentki, M. (2011). Type 2 diabetes across generations: from pathophysiology to prevention and management. *Lancet*, 378(9786), 169-181. doi: 10.1016/s0140-6736(11)60614-4
- Nomiyama, T., Igarashi, Y., Taka, H., Mineki, R., Uchida, T., Ogihara, T., . . . Watada, H. (2004). Reduction of insulin-stimulated glucose uptake by peroxynitrite is concurrent with tyrosine nitration of insulin receptor substrate-1. *Biochemical and Biophysical Research Communications*, 320(3), 639-647. doi: <http://dx.doi.org/10.1016/j.bbrc.2004.06.019>
- Okon, E. B., Chung, A. W., Rauniyar, P., Padilla, E., Tejerina, T., McManus, B. M., . . . van Breemen, C. (2005). Compromised arterial function in human type 2 diabetic patients. *Diabetes*, 54(8), 2415-2423.
- Okouchi, M., Okayama, N., Imai, S., Omi, H., Shimizu, M., Fukutomi, T., & Itoh, M. (2002). High insulin enhances neutrophil transendothelial migration through increasing surface expression of platelet endothelial cell adhesion molecule-1 via activation of mitogen activated protein kinase. *Diabetologia*, 45(10), 1449-1456. doi: 10.1007/s00125-002-0902-x
- Olukman, M., Orhan, C. E., Celenk, F. G., & Ulker, S. (2010). Apocynin restores endothelial dysfunction in streptozotocin diabetic rats through regulation of nitric oxide synthase and NADPH oxidase expressions. *J Diabetes Complications*, 24(6), 415-423. doi: 10.1016/j.jdiacomp.2010.02.001
- Park, K., Gross, M., Lee, D.-H., Holvoet, P., Himes, J. H., Shikany, J. M., & Jacobs, D. R. (2009). Oxidative Stress and Insulin Resistance: The Coronary Artery Risk Development in Young Adults study. *Diabetes Care*, 32(7), 1302-1307. doi: 10.2337/dc09-0259
- Peppas, M., & Vlassara, H. (2005). Advanced glycation end products and diabetic complications: a general overview. *Hormones (Athens)*, 4(1), 28-37.
- Perticone, F., Sciacqua, A., Scozzafava, A., Ventura, G., Laratta, E., Pujia, A., . . . Sesti, G. (2004). Impaired endothelial function in never-treated hypertensive subjects carrying the Arg972 polymorphism in the insulin receptor substrate-1 gene. *J Clin Endocrinol Metab*, 89(7), 3606-3609. doi: 10.1210/jc.2003-032161
- Pfeiffer, A., Middelberg-Bisping, K., Drewes, C., & Schatz, H. (1996). Elevated plasma levels of transforming growth factor-beta 1 in NIDDM. *Diabetes Care*, 19(10), 1113-1117.
- Pieper, G. M. (1997). Activation of nuclear factor- κ B in cultured endothelial cells by increased glucose concentration: prevention by calphostin C. *J Cardiovasc Pharmacol*, 30(4), 528-532.
- Quagliaro, L., Piconi, L., Assaloni, R., Martinelli, L., Motz, E., & Ceriello, A. (2003). Intermittent high glucose enhances apoptosis related to oxidative stress in human umbilical vein endothelial cells the role of protein kinase C and NAD(P)H-oxidase activation. *Diabetes*, 52(11), 2795-2804.

- Rajecki, M., Pajunen, P., Jousilahti, P., Rasi, V., Vahtera, E., & Salomaa, V. (2005). Hemostatic factors as predictors of stroke and cardiovascular diseases: the FINRISK '92 Hemostasis Study. *Blood Coagul Fibrinolysis*, *16*(2), 119-124.
- Ramachandran, A., Ma, R. C. W., & Snehalatha, C. (2010). Diabetes in Asia. *The Lancet*, *375*(9712), 408-418.
- Rampal, S., Rampal, L., Rahmat, R., Zain, A. M., Yap, Y. G., Mohamed, M., & Taha, M. (2010). Variation in the prevalence, awareness, and control of diabetes in a multiethnic population: a nationwide population study in Malaysia. *Asia Pac J Public Health*, *22*(2), 194-202. doi: 10.1177/1010539509334816
- Randriamboavonjy, V., Pistrosch, F., Bolck, B., Schwinger, R. H., Dixit, M., Badenhoop, K., . . . Fleming, I. (2008). Platelet sarcoplasmic endoplasmic reticulum Ca²⁺-ATPase and mu-calpain activity are altered in type 2 diabetes mellitus and restored by rosiglitazone. *Circulation*, *117*(1), 52-60. doi: 10.1161/circulationaha.107.719807
- Rask-Madsen, C., & King, G. L. (2007). Mechanisms of disease: endothelial dysfunction in insulin resistance and diabetes. *Nature Clinical Practice Endocrinology & Metabolism*, *3*(1), 46-56.
- Rask-Madsen, C., Li, Q., Freund, B., Feather, D., Abramov, R., Wu, I. H., . . . King, G. L. (2010). Loss of insulin signaling in vascular endothelial cells accelerates atherosclerosis in apolipoprotein E null mice. *Cell metabolism*, *11*(5), 379-389. doi: 10.1016/j.cmet.2010.03.013
- Rattan, V., Shen, Y., Sultana, C., Kumar, D., & Kalra, V. K. (1996). Glucose-induced transmigration of monocytes is linked to phosphorylation of PECAM-1 in cultured endothelial cells. *Am J Physiol*, *271*(4 Pt 1), E711-717.
- Reynolds, A. R., Reynolds, L. E., Nagel, T. E., Lively, J. C., Robinson, S. D., Hicklin, D. J., . . . Hodivala-Dilke, K. M. (2004). Elevated Flk1 (vascular endothelial growth factor receptor 2) signaling mediates enhanced angiogenesis in β 3-integrin-deficient mice. *Cancer Research*, *64*(23), 8643-8650.
- Roberts, A. C., & Porter, K. E. (2013). Cellular and molecular mechanisms of endothelial dysfunction in diabetes. *Diab Vasc Dis Res*, *10*(6), 472-482. doi: 10.1177/1479164113500680
- Rohwedder, I., Montanez, E., Beckmann, K., Bengtsson, E., Duner, P., Nilsson, J., . . . Fassler, R. (2012). Plasma fibronectin deficiency impedes atherosclerosis progression and fibrous cap formation. *EMBO Mol Med*, *4*(7), 564-576. doi: 10.1002/emmm.201200237
- Rolo, A. P., & Palmeira, C. M. (2006). Diabetes and mitochondrial function: Role of hyperglycemia and oxidative stress. *Toxicology and Applied Pharmacology*, *212*(2), 167-178. doi: <http://dx.doi.org/10.1016/j.taap.2006.01.003>
- Romao, I., & Roth, J. (2008). Genetic and Environmental Interactions in Obesity and Type 2 Diabetes. *Journal of the American Dietetic Association*, *108*(4), S24-S28. doi: 10.1016/j.jada.2008.01.022
- Roth, T., Podestá, F., Stepp, M. A., Boeri, D., & Lorenzi, M. (1993). Integrin overexpression induced by high glucose and by human diabetes: potential pathway to cell dysfunction in diabetic microangiopathy. *Proc Natl Acad Sci U S A*, *90*(20), 9640-9644.
- Ruberg, F. L., Leopold, J. A., & Loscalzo, J. (2002). Atherothrombosis: plaque instability and thrombogenesis. *Prog Cardiovasc Dis*, *44*(5), 381-394.
- Schafer, A., Schulz, C., Fraccarollo, D., Tas, P., Leutke, M., Eigenthaler, M., . . . Bauersachs, J. (2007). The CX3C chemokine fractalkine induces vascular dysfunction by generation of superoxide anions. *Arterioscler Thromb Vasc Biol*, *27*(1), 55-62. doi: 10.1161/01.atv.0000251535.30191.60

- Schalkwijk, C. G., & Stehouwer, C. D. (2005). Vascular complications in diabetes mellitus: the role of endothelial dysfunction. *Clin Sci (Lond)*, *109*(2), 143-159. doi: 10.1042/cs20050025
- Schulze, P. C., Yoshioka, J., Takahashi, T., He, Z., King, G. L., & Lee, R. T. (2004). Hyperglycemia promotes oxidative stress through inhibition of thioredoxin function by thioredoxin-interacting protein. *J Biol Chem*, *279*(29), 30369-30374. doi: 10.1074/jbc.M400549200
- Selvaraju, V., Joshi, M., Suresh, S., Sanchez, J. A., Maulik, N., & Maulik, G. (2012). Diabetes, oxidative stress, molecular mechanism, and cardiovascular disease--an overview. *Toxicol Mech Methods*, *22*(5), 330-335. doi: 10.3109/15376516.2012.666648
- Sena, C. M., Pereira, A. M., & Seica, R. (2013). Endothelial dysfunction — A major mediator of diabetic vascular disease. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, *1832*(12), 2216-2231. doi: <http://dx.doi.org/10.1016/j.bbadis.2013.08.006>
- Sharma, V., & PL, S. (2013). Role of Different Molecular Pathways in the Development of Diabetes-Induced Nephropathy. *J Diabetes Metab*, *S9*(004). doi: 10.4172/2155-6156.S9-004
- Sigala, F., Vourliotakis, G., Georgopoulos, S., Kavantzias, N., Papalambros, E., Agapitos, M., & Bastounis, E. (2003). Vascular endothelial cadherin expression in human carotid atherosclerotic plaque and its relationship with plaque morphology and clinical data. *Eur J Vasc Endovasc Surg*, *26*(5), 523-528.
- Simpson, R. W., Shaw, J. E., & Zimmet, P. Z. (2003). The prevention of type 2 diabetes--lifestyle change or pharmacotherapy? A challenge for the 21st century. *Diabetes Res Clin Pract*, *59*(3), 165-180.
- Solowiej, A., Biswas, P., Graesser, D., & Madri, J. A. (2003). Lack of platelet endothelial cell adhesion molecule-1 attenuates foreign body inflammation because of decreased angiogenesis. *Am J Pathol*, *162*(3), 953-962. doi: 10.1016/s0002-9440(10)63890-4
- Song, K. H., Park, J., Park, J. H., Natarajan, R., & Ha, H. (2013). Fractalkine and its receptor mediate extracellular matrix accumulation in diabetic nephropathy in mice. *Diabetologia*, *56*(7), 1661-1669. doi: 10.1007/s00125-013-2907-z
- Song, P., Wu, Y., Xu, J., Xie, Z., Dong, Y., Zhang, M., & Zou, M. H. (2007). Reactive nitrogen species induced by hyperglycemia suppresses Akt signaling and triggers apoptosis by upregulating phosphatase PTEN (phosphatase and tensin homologue deleted on chromosome 10) in an LKB1-dependent manner. *Circulation*, *116*(14), 1585-1595. doi: 10.1161/circulationaha.107.716498
- Srinivasan, V., Sandhya, N., Sampathkumar, R., Farooq, S., Mohan, V., & Balasubramanyam, M. (2007). Glutamine fructose-6-phosphate amidotransferase (GFAT) gene expression and activity in patients with type 2 diabetes: inter-relationships with hyperglycaemia and oxidative stress. *Clin Biochem*, *40*(13-14), 952-957. doi: 10.1016/j.clinbiochem.2007.05.002
- Steinberg, H. O., Chaker, H., Leaming, R., Johnson, A., Brechtel, G., & Baron, A. D. (1996). Obesity/insulin resistance is associated with endothelial dysfunction. Implications for the syndrome of insulin resistance. *J Clin Invest*, *97*(11), 2601-2610. doi: 10.1172/jci118709
- Stevens, H. Y., Melchior, B., Bell, K. S., Yun, S., Yeh, J. C., & Frangos, J. A. (2008). PECAM-1 is a critical mediator of atherosclerosis. *Dis Model Mech*, *1*(2-3), 175-181; discussion 179. doi: 10.1242/dmm.000547

- Stoynev, N., Dimova, I., Rukova, B., Hadjidekova, S., Nikolova, D., Toncheva, D., & Tankova, T. (2014). Gene expression in peripheral blood of patients with hypertension and patients with type 2 diabetes. *J Cardiovasc Med (Hagerstown)*, *15*(9), 702-709. doi: 10.2459/JCM.0b013e32835dbcc8
- Su, Y., Liu, X.-M., Sun, Y.-M., Wang, Y.-Y., Luan, Y., & Wu, Y. (2008). Endothelial Dysfunction in Impaired Fasting Glycemia, Impaired Glucose Tolerance, and Type 2 Diabetes Mellitus. *The American Journal of Cardiology*, *102*(4), 497-498. doi: <http://dx.doi.org/10.1016/j.amjcard.2008.03.087>
- Tabit, C. E., Chung, W. B., Hamburg, N. M., & Vita, J. A. (2010). Endothelial dysfunction in diabetes mellitus: molecular mechanisms and clinical implications. *Rev Endocr Metab Disord*, *11*(1), 61-74. doi: 10.1007/s11154-010-9134-4
- Takamura, T., Sakurai, M., Ota, T., Ando, H., Honda, M., & Kaneko, S. (2004). Genes for systemic vascular complications are differentially expressed in the livers of type 2 diabetic patients. *Diabetologia*, *47*(4), 638-647. doi: 10.1007/s00125-004-1366-y
- Tan, K. C., Chow, W. S., Ai, V. H., Metz, C., Bucala, R., & Lam, K. S. (2002). Advanced glycation end products and endothelial dysfunction in type 2 diabetes. *Diabetes Care*, *25*(6), 1055-1059.
- Taye, A., Saad, A. H., Kumar, A. H., & Morawietz, H. (2010). Effect of apocynin on NADPH oxidase-mediated oxidative stress-LOX-1-eNOS pathway in human endothelial cells exposed to high glucose. *Eur J Pharmacol*, *627*(1-3), 42-48. doi: 10.1016/j.ejphar.2009.10.045
- Tesauro, M., & Cardillo, C. (2011). Obesity, blood vessels and metabolic syndrome. *Acta Physiol (Oxf)*, *203*(1), 279-286. doi: 10.1111/j.1748-1716.2011.02290.x
- Tesauro, M., Rizza, S., Iantorno, M., Campia, U., Cardillo, C., Lauro, D., . . . Quon, M. J. (2007). Vascular, metabolic, and inflammatory abnormalities in normoglycemic offspring of patients with type 2 diabetes mellitus. *Metabolism*, *56*(3), 413-419. doi: 10.1016/j.metabol.2006.10.026
- Thum, T., Haverich, A., & Borlak, J. (2000). Cellular dedifferentiation of endothelium is linked to activation and silencing of certain nuclear transcription factors: implications for endothelial dysfunction and vascular biology. *Faseb j*, *14*(5), 740-751.
- Ting, H. H., Timimi, F. K., Boles, K. S., Creager, S. J., Ganz, P., & Creager, M. A. (1996). Vitamin C improves endothelium-dependent vasodilation in patients with non-insulin-dependent diabetes mellitus. *Journal of Clinical Investigation*, *97*(1), 22.
- Tiwari, B. K., Pandey, K. B., Abidi, A. B., & Rizvi, S. I. (2013). Markers of Oxidative Stress during Diabetes Mellitus. *Journal of Biomarkers*, *2013*, 8. doi: 10.1155/2013/378790
- Tusié Luna, & Teresa, M. (2005). Genes and Type 2 Diabetes Mellitus. *Archives of Medical Research*, *36*(3), 210-222. doi: <http://dx.doi.org/10.1016/j.arcmed.2005.03.004>
- van den Oever, I. A., Raterman, H. G., Nurmohamed, M. T., & Simsek, S. (2010). Endothelial dysfunction, inflammation, and apoptosis in diabetes mellitus. *Mediators Inflamm*, *2010*, 792393. doi: 10.1155/2010/792393
- van Haeften, T. W. (2002). Early disturbances in insulin secretion in the development of type 2 diabetes mellitus. *Mol Cell Endocrinol*, *197*(1-2), 197-204.
- von der Thusen, J. H., Kuiper, J., van Berkel, T. J., & Biessen, E. A. (2003). Interleukins in atherosclerosis: molecular pathways and therapeutic potential. *Pharmacol Rev*, *55*(1), 133-166. doi: 10.1124/pr.55.1.5

- Wang, Y., Cheng, K. K., Lam, K. S., Wu, D., Wang, Y., Huang, Y., . . . Xu, A. (2011). APPL1 counteracts obesity-induced vascular insulin resistance and endothelial dysfunction by modulating the endothelial production of nitric oxide and endothelin-1 in mice. *Diabetes*, *60*(11), 3044-3054. doi: 10.2337/db11-0666
- Weisensee, D., Bereiter-Hahn, J., Schoeppe, W., & Low-Friedrich, I. (1993). Effects of cytokines on the contractility of cultured cardiac myocytes. *Int J Immunopharmacol*, *15*(5), 581-587.
- Williams, S. B., Cusco, J. A., Roddy, M.-A., Johnstone, M. T., & Creager, M. A. (1996). Impaired nitric oxide-mediated vasodilation in patients with non-insulin-dependent diabetes mellitus. *J Am Coll Cardiol*, *27*(3), 567-574.
- Williamson, J. R., Chang, K., Frangos, M., Hasan, K. S., Ido, Y., Kawamura, T., . . . Tilton, R. G. (1993). Hyperglycemic pseudohypoxia and diabetic complications. *Diabetes*, *42*(6), 801-813.
- Wojta, J., Hoover, R. L., & Daniel, T. O. (1989). Vascular origin determines plasminogen activator expression in human endothelial cells. Renal endothelial cells produce large amounts of single chain urokinase type plasminogen activator. *J Biol Chem*, *264*(5), 2846-2852.
- Wong, B. W., Wong, D., & McManus, B. M. (2002). Characterization of fractalkine (CX3CL1) and CX3CR1 in human coronary arteries with native atherosclerosis, diabetes mellitus, and transplant vascular disease. *Cardiovasc Pathol*, *11*(6), 332-338.
- Wong, W. T., Wong, S. L., Tian, X. Y., & Huang, Y. (2010). Endothelial dysfunction: the common consequence in diabetes and hypertension. *J Cardiovasc Pharmacol*, *55*(4), 300-307.
- Xu, J., & Zou, M.-H. (2009). Molecular Insights and Therapeutic Targets for Diabetic Endothelial Dysfunction. *Circulation*, *120*(13), 1266-1286. doi: 10.1161/CIRCULATIONAHA.108.835223
- Xu, Y., Wang, S., Feng, L., Zhu, Q., Xiang, P., & He, B. (2010). Blockade of PKC-beta protects HUVEC from advanced glycation end products induced inflammation. *International immunopharmacology*, *10*(12), 1552-1559. doi: <http://dx.doi.org/10.1016/j.intimp.2010.09.006>
- Yaghoubi, A. R., Kargar, F., & Khaki-Khatibi, F. (2016). Association of Fibronectin, Leptin and LDL-Oxide Serum Levels With Coronary Artery Disease in Non-smoker and Non-diabetic Patients. *Crescent Journal of Medical and Biological Sciences*, *3*(1), 23-27.
- Yang, Z., & Li, J.-C. (2008). Stimulation of endothelin-1 gene expression by insulin via phosphoinositide-3 kinase-glycogen synthase kinase-3 β signaling in endothelial cells. *Life Sciences*, *82*(9-10), 512-518. doi: <http://dx.doi.org/10.1016/j.lfs.2007.12.005>
- Yener, S., Comlekci, A., Akinci, B., Akan, P., Demir, T., Bayraktar, F., & Yesil, S. (2008). Serum transforming growth factor-beta 1 levels in normoalbuminuric and normotensive patients with type 2 diabetes. Effect of metformin and rosiglitazone. *Hormones (Athens)*, *7*(1), 70-76.
- Yildirim, N., Sahin, A., Erol, N., Kara, S., Uslu, S., & Topbas, S. (2008). The relationship between plasma MMP-9 and TIMP-2 levels and intraocular pressure elevation in diabetic patients after intravitreal triamcinolone injection. *J Glaucoma*, *17*(4), 253-256. doi: 10.1097/IJG.0b013e31815c3a07
- Yoon, K. H., Lee, J. H., Kim, J. W., Cho, J. H., Choi, Y. H., Ko, S. H., . . . Son, H. Y. (2006). Epidemic obesity and type 2 diabetes in Asia. *Lancet*, *368*(9548), 1681-1688. doi: 10.1016/s0140-6736(06)69703-1

- Yu, Q., Gao, F., & Ma, X. L. (2011). Insulin says NO to cardiovascular disease. *Cardiovasc Res*, 89(3), 516-524. doi: 10.1093/cvr/cvq349
- Zaghloul, A., Al-Bukhari, T. A., Al-Pakistani, H. A., Shalaby, M., Halawani, S. H., Bajuaifer, N., & Teama, S. H. (2014). Soluble endothelial protein C receptor and high sensitivity C reactive protein levels as markers of endothelial dysfunction in patients with type 1 and type 2 diabetes mellitus: Their role in the prediction of vascular complications. *Diabetes Res Clin Pract*, 106(3), 597-604. doi: 10.1016/j.diabres.2014.09.007
- Zeng, G., Nystrom, F. H., Ravichandran, L. V., Cong, L. N., Kirby, M., Mostowski, H., & Quon, M. J. (2000). Roles for insulin receptor, PI3-kinase, and Akt in insulin-signaling pathways related to production of nitric oxide in human vascular endothelial cells. *Circulation*, 101(13), 1539-1545.
- Zerón, H. M., Páez, A., Zapata, E., Massó, F., & Montaña, L. F. (2010). HUVECs from newborns with a strong family history of diabetes show increased apoptosis by flow cytometry with annexin V. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 4(3), 168-173. doi: <http://dx.doi.org/10.1016/j.dsx.2010.05.013>
- Zhang, H., Dellsperger, K., & Zhang, C. (2011). The link between metabolic abnormalities and endothelial dysfunction in type 2 diabetes: an update. *Basic Research in Cardiology*, 107(1), 1-11. doi: 10.1007/s00395-011-0237-1
- Zhang, J., Patel, J. M., Li, Y. D., & Block, E. R. (1997). Proinflammatory cytokines downregulate gene expression and activity of constitutive nitric oxide synthase in porcine pulmonary artery endothelial cells. *Res Commun Mol Pathol Pharmacol*, 96(1), 71-87.
- Zhang, L., Gong, D., Li, S., & Zhou, X. (2012). Meta-analysis of the effects of statin therapy on endothelial function in patients with diabetes mellitus. *Atherosclerosis*, 223(1), 78-85. doi: 10.1016/j.atherosclerosis.2012.01.031
- Zhou, Y. J., Wang, J. H., Li, L., Yang, H. W., Wen de, L., & He, Q. C. (2007). Expanding expression of the 5-lipoxygenase/leukotriene B4 pathway in atherosclerotic lesions of diabetic patients promotes plaque instability. *Biochem Biophys Res Commun*, 363(1), 30-36. doi: 10.1016/j.bbrc.2007.08.134
- Zumbach, M., Hofmann, M., Borcea, V., Luther, T., Kotzsch, M., Muller, M., . . . Nawroth, P. P. (1997). Tissue factor antigen is elevated in patients with microvascular complications of diabetes mellitus. *Exp Clin Endocrinol Diabetes*, 105(4), 206-212. doi: 10.1055/s-0029-1211753