



UNIVERSITI PUTRA MALAYSIA

***FACTORS ASSOCIATED WITH BODY WEIGHT STATUS AMONG
MALAYSIAN ADOLESCENTS LIVING IN DAY-SCHOOL HOSTELS***

LAI SOKE CHING

FPSK(M) 2016 6



**FACTORS ASSOCIATED WITH BODY WEIGHT STATUS AMONG
MALAYSIAN ADOLESCENTS LIVING IN DAY-SCHOOL HOSTELS**

By

LAI SOKE CHING

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of
Master of Science**

January 2016

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Degree of Master of Science

**FACTORS ASSOCIATED WITH BODY WEIGHT STATUS AMONG
MALAYSIAN ADOLESCENTS LIVING IN DAY-SCHOOL HOSTELS**

By

LAI SOKE CHING

January 2016

Chair : Chin Yit Siew, PhD
Faculty : Medicine and Health Sciences

The lives of adolescents in school hostels are well-managed with respect to their daily food intake and physical activity. They are likely to be physically active and are assumed to consume nutritious food. However, little is known about their body weight status and its associated factors. Thus, this study determined the association between socio-demographic characteristics, dietary practice, physical activity, knowledge, attitude and practice on healthy lifestyle, Transtheoretical Model of behavioral change on healthy diet and exercise, and body image perception with body weight status among Malaysian adolescents living in day-school hostels.

A total of 4189 school-going adolescents (aged 13 years) were involved in this study. Information on socio-demographic characteristics, dietary practice, physical activity, knowledge, attitude and practice on healthy lifestyle, Transtheoretical Model of behavioral change on healthy diet and exercise and body image were collected using a self-administered questionnaire. Body weight and height were measured while BMI-for-age (z-score) was determined using WHO Growth Reference. One day 24-hour dietary and physical activity recalls were obtained by face-to-face interview. Multivariate general linear regression analysis was conducted.

Majority of the respondents were females (59.5%), Malays (70.5%), from rural area (70.2%) and were achieved pubertal status (65.9%). The prevalence of overweight and obesity [Both sexes: 23.3% (95% Confidence Interval [CI]: 22.1, 24.7); Males: 21.7% (95% CI: 19.8, 23.8); Females: 24.5% (95% CI: 22.8, 26.2)] was about six times higher than the prevalence of thinness [Both Sexes: 4.3% (95% CI: 3.7, 5.0); Males: 5.6% (95% CI: 4.5, 6.9); Females: 3.4% (95% CI: 2.7, 4.2)] among the respondents. The mean total daily energy intake was 2070 Kcal (95% CI: 2045, 2097) and the mean total daily energy expenditure was 1965 Kcal (95% CI: 1952, 1978). The mean score for knowledge and attitude were 18.6 (95% CI: 18.4, 18.7) and 98.9 (95% CI: 98.4, 99.2), respectively. Self-efficacy and perceived benefits on healthy diet had mean score of 15.8 (95% CI: 15.7, 15.9) and 27.8 (95% CI: 27.6, 27.9), respectively. Slightly less than half of the respondents (48.9%, 95% CI: 47.4, 50.5) had correct perception on body weight, about

one in three respondents (32.4%, 95% CI: 31.0, 33.9) satisfied with their body size and two in three respondents (68.2%, 95% CI: 66.7, 69.6) had correct perception on healthy body size.

Bivariate analysis indicated that sex, area, pubertal status, energy expenditure per kilogram body weight, knowledge and attitude on healthy lifestyle, self-efficacy and perceived benefits on exercise, body weight status perception, body size satisfaction and healthy body size perception were significantly associated with BMI-for-age of the respondents. No associations were found between ethnicity, dietary practice, physical activity level, practice on healthy lifestyle, stages of healthy diet behavioral change, stages of exercise behavioral change, self-efficacy on exercise, perceived barriers for both healthy diet and exercise, and perceived benefits for exercise with BMI-for-age of the respondents. In multivariate general linear regression model, being female ($\beta= 0.620$) having achieved pubertal status ($\beta= 0.408$), positive attitude on healthy diet ($\beta= 0.005$), better self-efficacy on healthy diet ($\beta= -0.014$), lower energy expenditure per kilogram body weight ($\beta= -0.075$), incorrect body weight status perception ($\beta= 0.177$), dissatisfaction with body size ($\beta= 0.144$), and incorrect healthy body size perception ($\beta= 0.113$) significantly contributed to the high BMI-for-age of the respondents at $p<0.05$ level of significance explaining 34.7% of the variance in BMI-for-age ($R^2=0.347$, $F=119.512$, $p<0.001$).

Dual-form of malnutrition co-exists with overweight and obesity were more prevalent than thinness. Future healthy body weight intervention programmes should emphasize on sex-specific, attitude on healthy lifestyle, self-efficacy on healthy diet and body image approach among these adolescents living in day-school hostels.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

**FAKTOR BERKAITAN DENGAN STATUS BERAT BADAN DALAM
KALANGAN REMAJA MALAYSIA YANG TINGGAL DI ASRAMA
SEKOLAH HARIAN**

Oleh

LAI SOKE CHING

Januari 2016

Pengerusi : Chin Yit Siew, PhD
Fakulti : Perubatan dan Sains Kesihatan

Kehidupan remaja di asrama sekolah adalah diurus dengan baik dari segi pengambilan makanan harian dan aktiviti fizikal. Mereka diandaikan aktif secara fizikal dan mengambil makanan berkhasiat. Walau bagaimanapun, tidak banyak yang diketahui mengenai status berat badan dan faktor yang berkaitan dengannya. Oleh itu, kajian ini bertujuan untuk menentukan perkaitan antara ciri socio-demografi, pengambilan makanan, tahap aktiviti fizikal, pengetahuan, sikap dan amalan gaya hidup sihat, Model Transtheoretical perubahan tingkah laku ke arah pemakanan yang sihat dan senaman, persepsi imej tubuh, dan status berat badan dalam kalangan remaja Malaysia yang tinggal di asrama sekolah harian.

Sejumlah 4189 remaja yang masih bersekolah (berumur 13 tahun) terlibat dalam kajian ini. Maklumat yang merangkumi ciri sosio-demografi, pengambilan pemakanan, aktiviti fizikal, pengetahuan, sikap dan amalan gaya hidup sihat, Model Transheoretical perubahan tingkah laku ke arah pemakanan yang sihat dan senaman, dan imej tubuh telah diambil daripada responden melalui borang soal-selidik. Berat badan dan tinggi telah diukur. BMI-untuk-umur (z-skor) telah ditentukan berdasarkan Rujukan Pertumbuhan WHO. Ingatan diet 24-jam dan ingatan aktiviti fizikal 24-jam telah dijalankan melalui temuduga muka-ke-muka. Analisis regresi linear berganda telah dijalankan.

Majoriti responden adalah perempuan (59.5%), Melayu (70.5%), dari kawasan luar bandar (70.2%) dan mencapai akil baligh (65.9%). Prevalens untuk berlebihan berat badan dan obesiti [Kedua-dua jantina: 23.3% (95% *Confidence Interval* [CI]: 22.1, 24.7); Lelaki: 21.7% (95% CI: 19.8, 23.8); Perempuan: 24.5% (95% CI: 22.8, 26.2)] adalah hampir enam kali lebih tinggi daripada prevalens kekurangan berat badan [Kedua-dua jantina: 4.3% (95% CI: 3.7, 5.0); Lelaki: 5.6% (95% CI: 4.5, 6.9); Perempuan: 3.4% (95% CI: 2.7, 4.2)] dalam kalangan responden. Min pengambilan tenaga sehari adalah 2070 Kcal (95% CI: 2045, 2097) dan min penggunaan tenaga sehari adalah 1965 Kcal (95% CI: 1952, 1978). Min skor pengetahuan dan sikap masing-masing adalah 18.6 (95% CI:

18.4, 18.7) dan 98.9 (95% CI: 98.4, 99.2). Keberkesanan diri dan manfaat yang dirasakan pada pemakanan yang sihat masing-masing mempunyai min skor 15.8 (95% CI: 15.7, 15.9) dan 27.8 (95% CI: 27.6, 27.9). Hampir separuh daripada responden (48.9%, 95% CI: 47.4, 50.5) mempunyai persepsi yang betul terhadap status berat badan, iaitu satu daripada tiga responden (32.4%, 95% CI: 31.0, 33.9) berpuas hati dengan saiz tubuh mereka dan dua daripada tiga responden (68.2%, 95% CI: 66.7, 69.6) mempunyai persepsi yang betul terhadap saiz tubuh yang sihat.

Analisis bivariate menunjukkan jantina, kawasan, status akil baligh, penggunaan tenaga per kilogram berat badan, pengetahuan dan sikap ke arah hidup sihat, keberkesanan diri dan manfaat yang dirasakan daripada senaman, persepsi terhadap status berat badan, kepuasan terhadap saiz tubuh dan persepsi terhadap saiz tubuh yang sihat mempunyai perkaitan yang signifikan dengan BMI-untuk-umur responden. Tiada perkaitan didapati antara etnik, amalan pemakanan, tahap aktiviti fizikal, amalan cara hidup sihat, peringkat perubahan tingkah laku pemakanan yang sihat, peringkat-peringkat perubahan tingkah laku senaman, keberkesanan diri terhadap senaman, halangan yang dirasakan terhadap pemakanan yang sihat dan senaman, dan manfaat yang dirasakan terhadap senaman dengan BMI-untuk-umur responden.

Analisis regresi linear berganda model menunjukkan faktor seperti perempuan ($\beta = 0.620$), mencapai akil baligh ($\beta = 0.408$), sikap yang positif terhadap pemakanan yang sihat ($\beta = 0.005$), keberkesanan diri yang tinggi terhadap pemakanan yang sihat ($\beta = -0.014$), penggunaan tenaga per kilogram berat badan yang rendah ($\beta = -0.075$), persepsi terhadap status berat badan yang salah ($\beta = 0.177$), ketidakpuasan terhadap saiz tubuh ($\beta = 0.144$), persepsi terhadap saiz tubuh yang sihat yang salah ($\beta = 0.113$) menyumbang secara signifikan terhadap BMI-untuk-umur responden yang tinggi pada tahap keyakinan $p < 0.05$ menjelaskan 34.7% daripada variasi dalam BMI-untuk-umur ($R^2 = 0.347$, $F = 119.512$, $p < 0.001$).

Dwi-bentuk malnutrisi telah wujud di mana prevalens berlebihan berat badan dan obes lebih tinggi berbanding dengan kekurangan berat badan. Program intervensi berkaitan berat badan yang sihat pada masa depan harus memberi perhatian terhadap pengkhususan kepada jantina tertentu, sikap terhadap gaya hidup sihat, keberkesanan diri terhadap pemakanan yang sihat dan imej tubuh dalam kalangan remaja yang tinggal di sekolah asrama harian.

ACKNOWLEDGEMENT

This dissertation would not have been possible without the support and assistance of several individuals who contributed that in one way or another contributed and extended their valuable assistance in the preparation and completion of this study.

First and foremost, I would like to extend my gratitude and appreciation to my supervisor, Dr. Chin Yit Siew for her inspiration, guidance and advice throughout the study. I am deeply appreciated and grateful for her constant encouragement and support at various stages of my study.

I am thankful to Assoc. Prof. Dr. Chan Yoke Mun and Assoc. Prof. Dr. Mohd Nasir Mohd Taib for their advice, valuable comments and timely suggestions that enabled me to complete this dissertation.

I would like to express my sincere thanks to Chong Lin Siew and Lim Shiang Cheng for their encouragement and being a good companion throughout my Master degree, and research team that giving their full support and huge contribution to this study. Without their support and commitment, this study would not have come this far.

This study will not be completed without full commitment given by Ministry of Education in conducting this study at the selected school nationwide. More specifically to all the State of Department Education, school principals, teachers, students and parents for their excellent cooperation and commitment throughout the study.

Lastly, I would like to take this opportunity to record my appreciation to my family members, especially my parents, for their love, patience and unending support throughout the process.

I certify that a Thesis Examination Committee has met on 21 January 2016 to conduct the final examination of Lai Soke Ching on her thesis entitled "Factors Associated with Body Weight Status among Malaysian Adolescents Living in Day-School Hostels" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Loh Su Peng, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Rosita binti Jamaluddin, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Nik Shanita binti Safii, PhD

Associate Professor
Universiti Kebangsaan Malaysia
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 24 March 2016 .

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Chin Yit Siew, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Chan Yoke Mun, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Mohd Nasir Mohd Taib, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

BUJANG KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property form the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in this thesis, and scholarly integrity was upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____

Date: _____

Name and Matric No.: Lai Soke Ching, GS 35385

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to

Signature: _____

Name of

Chairman of

Supervisory

Committee: Dr. Chin Yit Siew

Signature: _____

Name of

Member of

Supervisory

Committee: Assoc. Prof. Dr. Mohd Nasir Mohd Taib

Signature: _____

Name of

Member of

Supervisory

Committee: Assoc. Prof. Dr. Chan Yoke Mun

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xv
LIST OF APPENDICES	xvi
LIST OF ABBREVIATIONS	xvii
 CHAPTER	
1 INTRODUCTION	1
1.1 Background of the Study	1
1.2 Problem Statement	2
1.3 Significance of the Study	4
1.4 Objectives of the Study	5
1.4.1 General Objective	5
1.4.2 Specific Objectives	5
1.5 Null Hypothesis	6
1.6 Conceptual Framework	6
 2 LITERATURE REVIEW	 9
2.1 Adolescence	9
2.2 General Problems in Adolescents	9
2.3 Body Weight Status	10
2.3.1 Definition of Body Weight Status	10
2.3.2 Body Weight Status - Worldwide	10
2.3.3 Body Weight Status - Malaysia	11
2.3.4 Consequences of Malnutrition	13
2.4 Association between Socio-demographic Factors and Overweight and Obesity	13
2.4.1 Sex	13
2.4.2 Ethnicity	14
2.4.3 Area	14
2.4.4 Pubertal Status	15
2.5 Association between Dietary Practice Factors and Overweight and Obesity	16
2.5.1 Energy and Macronutrient Intakes	16
2.5.2 Eating Behavior	17
2.6 Association between Energy Expenditure and Physical Activity Level Factors and Overweight and Obesity	18
2.7 Association between Knowledge, Attitude and Practice on Healthy Lifestyle Factors and Overweight and Obesity	19
2.7.1 Knowledge on Healthy Lifestyle	20
2.7.2 Attitude on Healthy Lifestyle	21

2.7.3	Practice on Healthy Lifestyle	21
2.8	Association between Transtheoretical Model of Behavioral Change Factors and Overweight and Obesity	22
2.8.1	Stages of Dietary Behavioral Change	23
2.8.2	Stages of Exercise Behavioral Change	24
2.8.3	Self-efficacy on Healthy Diet	25
2.8.4	Self-efficacy on Exercise	26
2.8.5	Decisional Balance on Healthy Diet	27
2.8.6	Decisional Balance on Exercise	28
2.9	Association between Body Image Factors and Overweight and Obesity	28
2.10	Multifactorial Model of Body Weight Status	30
3	METHODOLOGY	32
3.1	Research Background and Study Design	32
3.2	Study Location and Subjects	32
3.3	Study Inclusion Criteria and Exclusion Criteria	33
3.4	Sample Size Calculation	33
3.5	Sampling and Sample Design	35
3.6	Translation of Questionnaire	37
3.7	Research Instruments	37
3.7.1	Anthropometric Measurements	37
3.7.2	Self-administered Questionnaire	37
3.7.3	24-hour Dietary Recall	44
3.7.4	24-hour Physical Activity Recall	45
3.8	Ethical Approvals	46
3.9	Data Collection	47
3.10	Data Analysis	47
4	RESULTS	49
4.1	Socio-demographic Characteristics	49
4.2	Body Weight Status	51
4.3	Dietary Practices	53
4.3.1	Energy and Nutrient Intakes	53
4.3.2	Pattern of Meal Consumption and Meal Skipping Behaviors	57
4.4	Energy Expenditure	59
4.4.1	Physical Activity Level (PAL)	60
4.4.2	Energy Balance	61
4.5	Knowledge, Attitude and Practice on Healthy Lifestyle	62
4.6	Transtheoretical Model of Behavioral Change	71
4.6.1	Stages of Change	71
4.6.2	Self-efficacy	73
4.6.3	Decisional Balance	74
4.7	Body Image	77
4.7.1	Body Weight Status Perception	77
4.7.2	Body Size Perception	79
4.8	Association between Socio-demographic Characteristics and BMI-for-age	82
4.9	Association between Dietary Practice and BMI -for-age	83

4.10	Association between Physical Activity and BMI -for-age	84
4.11	Association between Knowledge, Attitude and Practice (KAP) on Healthy Lifestyle Factors and BMI -for-age	84
4.12	Association between Transtheoretical Model Behavioral Change and BMI -for-age	85
4.13	Association between Body Image and BMI -for-age	86
4.14	Contribution of Socio-demographic, Knowledge, Attitude and Practice (KAP) on Healthy Lifestyle Factors, Behavioral Change, Dietary Practice, Physical Activity and Body Image Factors towards BMI-for-age (z-score)	87
5	DISCUSSIONS	89
5.1	Body Weight Status	89
5.2	Association between Socio-demographic Factors and BMI-for-age	90
5.3	Association between Dietary Practice Factors and BMI-for-age	91
5.4	Association between Physical Activity Factors and the BMI-for-age	91
5.5	Association between Knowledge, Attitude and Practice on Healthy Lifestyle Factors and the BMI-for-age	92
5.6	Association between Transtheoretical Model Behavioral Change and the BMI-for-age	93
5.7	Association between Body Image Factors and the BMI-for-age	95
5.8	Contribution of Factors towards the BMI-for-age	96
6	CONCLUSION AND RECOMMENDATIONS	98
6.1	Conclusion	98
6.2	Limitations	98
6.3	Recommendations	98
	BIBLIOGRAPHY	100
	APPENDICES	121
	BIODATA OF STUDENT	168
	LIST OF PUBLICATIONS	169

LIST OF TABLES

Table		Page
2.1	Worldwide trends of overweight and obesity among adolescents aged below 18 years	12
3.1	Sample size calculation for prevalence study	33
3.2	Sample size calculation for multiple linear regression analysis using G* Power 3.1.9.2	34
3.3	Classification of BMI-for-age	37
3.4	Classification of stages of healthy diet and exercise behavioral change	39
3.5	Classification of body weight status perception	42
3.6	Classification of body size perception	44
3.7	Classification of accuracy in reporting dietary energy intake	45
3.8	Classification of physical activity level	46
4.1	Distribution of respondents by socio-demographic characteristics	49
4.2	Mean body weight, height, BMI and BMI-for-age of respondents	52
4.3	Energy and nutrient intakes among respondents	53
4.4	Distribution of body weight status respondents by energy intake	54
4.5	Distribution of respondents according to nutrient adequacy	54
4.6	Distribution of respondents by under-reporting, acceptable-reporting, and over-reporting of energy intake	56
4.7	Distribution of body weight status respondents by under-reporting, acceptable-reporting, and over-reporting of energy intake	57
4.8	Distribution of respondents according to pattern of meal consumption	58
4.9	Distribution of respondents by meal skipping behaviors	58
4.10	Distribution of respondents by snacking behaviors	59
4.11	Mean energy expenditure, physical activity level, and BMR of respondents	59
4.12	Distribution of body weight status respondents by mean energy expenditure, physical activity level, and BMR of respondents	60
4.13	Distribution of respondents by physical activity level	61
4.14	Distribution of body weight status respondents by physical activity level	61
4.15	Distribution of respondents by energy balance	62
4.16	Distribution of respondents by knowledge on healthy lifestyle	62
4.17	Distribution of respondents by attitude on healthy lifestyle	65
4.18	Distribution of respondents by practice on healthy lifestyle	69
4.19	Distribution of current stages of change for healthy diet and exercise	72
4.20	Distribution of respondents according to self-efficacy for healthy diet	73
4.21	Distribution of respondents according to self-efficacy for exercise	74
4.22	Distribution of respondents according to perceived benefits and perceived barriers for healthy diet	75
4.23	Distribution of respondents according to perceived benefits and perceived barriers for exercise	76

4.24	Distribution of respondents according to body weight status perception	78
4.25	Distribution of respondents by correct- and incorrect-body weight status perception	78
4.26	Distribution of respondents according to perception of the current body size, perception of healthy body size based on Contour Drawing Rating Scale (Thompson and Gray, 1995)	80
4.27	Mean scores of body size discrepancy score	81
4.28	Distribution of respondents by body shape perception	82
4.29	Association between socio-demographic characteristics and BMI-for-age	83
4.30	Association between dietary practice and BMI -for-age	83
4.31	Association between energy intake and macronutrients intake and BMI-for-age	84
4.32	Association between physical activity factors and BMI -for-age	84
4.33	Association between knowledge, attitude and practice (KAP) on healthy lifestyle factors and BMI -for-age	85
4.34	Association between stage of changes and BMI -for-age	85
4.35	Association between self-efficacy, decisional balance and BMI-for-age	86
4.36	Association between body image and BMI -for-age	86
4.37	General Linear Model for BMI-for-age	88

LIST OF FIGURES

Table		Page
1.1	Conceptual framework	8
3.1	Multistage Stratified Cluster Sampling Design	36
3.2	Series figure of “Contour Drawing Rating Scale” for female (Thompson & Gray, 1995)	43
3.3	Series figure of “Contour Drawing Rating Scale” for male (Thompson & Gray, 1995)	43
4.1	Distribution of respondents by pubertal status	51
4.2	Distribution of respondents by body weight status	52



LIST OF APPENDICES

Appendix		Page
A	Information Sheet	121
B	Consent Form	124
C	Self-administered Questionnaire	127
D	24-Hour Dietary Recall Form	146
E	24-Hour Physical Activity Recall Form	148
F	Ethics Approval	150
G	Approval Letter (MoE)	153
H	Approval Letter (State of Education Department)	155



LIST OF ABBREVIATIONS

BMI	Body Mass Index
BMR	Basal Metabolic Rate
CDC	Centers for Disease Control and Prevention
CI	Confidence Interval
EI	Energy Intake
EE	Energy Expenditure
FAO	Food and Agriculture Organization
IOTF	International Obesity Task Force
IPH	Institute of Public Health
KAP-HLQ	Knowledge, Attitude, and Practice on Healthy Lifestyle Questionnaire
MET	Metabolic Equivalent
MoE	Ministry of Education
MSNS	Malaysian School-Based Nutrition Survey
MVPA	Moderate to Vigorous Physical Activity
NCCFN	National Coordinating Committee on Food and Nutrition
NHANES	National Health and Nutrition Examination Survey
NHMS	National Health and Morbidity Survey
PAL	Physical Activity Level
RM	Ringgit Malaysia
RMR	Resting Metabolic Rate
RNI	Recommended Nutrient Intakes for Malaysians
SOC	Stages of Change
SPSS	Statistical Packages for Social Sciences
TDEE	Total Daily Energy Expenditure
TTM	Transtheoretical Model
UNU	United Nations University
UPM	Universiti Putra Malaysia
WGOC	Working Group on Obesity in China
WHO	World Health Organization
RM	Ringgit Malaysia

CHAPTER 1

INTRODUCTION

1.1 Background of Study

World Health Organization [WHO] (2000) defined obesity as abnormal and excessive fat accumulation in the body. Obesity may impair health and is among the most common non-communicable diseases that causes morbidity and mortality throughout the world. The transitions in economic development and urbanization resulted in lifestyle changes, with increased consumption of energy-dense food and lack of physical activities (Popkin, 2001). This concomitant transition in nutrition has contributed to nutritional problem in which thinness and obesity co-exist. Globally, the rate of obesity is increasing dramatically and extended from developed countries to developing countries.

Previous studies conducted in developed countries had shown the high prevalence in overweight and obesity (Ogden, Carroll, Kit, & Flegal, 2012b, 2014; Schönbeck et al., 2011). For instance, a study conducted among Dutch children and adolescents showed increasing trend in which the prevalence have increased two to three fold in overweight and four to six fold in obesity since 1980 compared to the year 2009 (Schönbeck et al., 2011) according to the International Obesity Task Force (IOTF) Classification. Another study conducted in the United States revealed that about two in five youth (16.9%) aged 2 to 19 years old were obese (Ogden et al., 2014), which is based on the Centers for Disease Control and Prevention (CDC) BMI-for-age growth charts. However, the data remained unchanged compared with that in 2009-2010 (16.9%) (Ogden et al., 2012b), despite the high prevalence in obesity.

On the other hand, studies on Asian countries reported that the prevalence of overweight and obesity were more prevalent than thinness using different cut-offs. For example, in Vietnam, Tang et al. (2010) used the IOTF Classification to categorize body mass index according to age and found that the prevalence of overweight (14.7%) and obesity (18.9%) among adolescents aged 6-19 years was higher than the prevalence of thinness (7.6%). China, as one of the world's fastest growing economy (Gupta & Wang, 2009), reported an escalating epidemic of overweight and obesity among children and adolescents aged 7-18 years from 1985 to 2010 (male: 20.2% to 33.2%; female: 2.3% to 19.1%) by using the Working Group on Obesity in China (WGOC) standard (Zhang & Wang, 2012). In India, Gupta et al. (2011), also using the IOTF Classification, found that there has been a decline in the prevalence of thinness (11.3% to 3.9%) accompanied by an increase in the prevalence of obesity (9.8% to 11.7%) from 2006 to 2009 among urbanite adolescents aged 14-17 years in New Delhi (North India) reflecting the effects of nutrition transition (Gupta et al., 2011).

Malaysia is also currently undergoing an active epidemiological and nutrition transition with a rapid increase in adolescents overweight and obesity. Data from the National

Health and Morbidity Survey (NHMS) III in 2006 (Institute for Public Health [IPH], 2008), which was based on the CDC 2000 reference (weight-for-age), reported that the prevalence of thinness (weight-for-age $< -2SD$) and overweight (weight-for-age $> +2SD$) among Malaysian adolescents aged less than 18 years were 13.2% and 5.4%, respectively (IPH, 2008). Subsequently, the National Health and Morbidity Survey (NHMS) IV in 2011 (IPH, 2011) has revealed that the prevalence of thinness (BMI-for-age $< -2SD$) and obesity (BMI-for-age $> +2SD$) among Malaysian adolescents aged less than 18 years were 12.2% and 6.1%, respectively (CDC 2000 reference) (IPH, 2011). The latest Malaysia School-Based Nutrition Survey 2012 focused on school-going adolescents aged 10 to 17 years reported that the prevalence of thinness (BMI-for-age $< -2SD$) and obesity (BMI-for-age $> +2SD$) were 7.4% and 12.3%, respectively. As such, the co-existence of thinness, overweight and obesity prevalent are of public health concern, particularly because obesity is a forewarning risk factor of non-communicable diseases (IPH, 2013).

Previous studies found that both under- and over-nutrition during adolescence have important short- and long-term implications. Over-nutrition among adolescents is associated with several adverse outcomes in adulthood including cardiovascular disease (Alp et al., 2014), metabolic syndrome (Steinberger & Kelly, 2014), and psychological outcomes (Brixval, Rayce, Rasmussen, Holstein, & Due, 2012). Additionally, under-nutrition during adolescence has been linked to psychiatric disorders (Bühren et al., 2013), delayed growth and pubertal development (Ferrar & Olds, 2010).

It is also noteworthy that adolescent obesity may persist into adulthood and increases the likelihood of adult morbidity and mortality (Dietz, 1998). Regardless of the reference cut-off points used for weight status, the increasing epidemic of overweight and obesity among adolescents is of worrying. Treatment of these epidemics in adolescents is challenging, and prevention is the only feasible option to curb this public health problem. Thus, a better knowledge of the body weight status and the risk factors associated with them is a pre-requisite for setting up health promotion programs.

1.2 Problem Statement

Adolescence is a period which involves dramatic physical, psychological, and cognitive changes. In order to achieve optimal growth and development during adolescence, the nutritional requirements during this period are the highest across the life span (National Coordinating Committee on Food and Nutrition [NCCFN], 2010). However, immature cognitive abilities, increasing autonomy in making decisions, and susceptibility to environmental influences during this stage may contribute to unhealthy nutritional and lifestyle choices among adolescents (Stang & Story, 2005) which in turn, may become risk factors for thinness, overweight and obesity (De-Jong et al., 2013; Law, Mohd Nasir, & Hazizi, 2013). The thinness, overweight, and obesity generation is not only a serious and costly health condition but such health complication will also affect the growth and development of the country.

Previous local national studies revealed that thinness, overweight, and obesity coexist among adolescents in Malaysia (IPH, 2011; IPH, 2013). Meanwhile, the latest nationwide studies, focused on school-going adolescents, reported that the problem of overweight and obesity was more prevalent than the problem of thinness among school-going adolescents (IPH, 2013; Poh et al., 2013). However, most of the local nationwide studies have been conducted among school going adolescents (Baharudin et al., 2013; IPH, 2008; IPH, 2011), not adolescents attending and living in day-school hostels.

In fact, there are two types of government-financed residential schools in Malaysia, which are fully residential school (*Sekolah Berasrama Penuh*) and day-school facilitated with school-hostel (*Sekolah Berasrama Harian*) (Ministry of Education Malaysia [MoE], 2012). Fully residential school is established to nurture outstanding students to excel in academics as well as co-curriculum and students are required to live in school hostels provided. On the other hand, day-schools facilitated with school hostel is established to provide conducive and safe accommodation for less privileged students who live too far from school to commute daily. Although they moved away from home to stay in a different environment which is in school hostels, studies have yet to focus on this group of adolescents.

According to the MoE (2012), meals are standardized for all these adolescents living in school hostels and extra-curriculum activities are included in their daily schedule. Since they are likely to consume nutritious food and active in physical activity, it is hypothesized that the prevalence of overweight and obesity would be low among this group of adolescents. However, the current trend of under- and over-nutrition among adolescents is prevalent in Malaysia, and studies on nutritional status among adolescents living in day-school hostels are limited (Baharudin et al., 2014; Poh et al., 2013). Therefore, it is timely to determine whether overweight and obesity are prevalent among adolescents living in day-school hostels.

Overweight and obesity among adolescents are a multifactorial disease. Many previous studies have shown that the prevalence of overweight and obesity among adolescents appears to be associated with sociodemographic characteristics (Baharudin et al., 2013; Khambalia, Lim, Gill, & Bulgiba, 2012). Previous studies also reported that excess energy intake (Soo, Wan, Abdul, & Lee, 2011) and low physical activity level (Su et al., 2014; Teo, Nurul-Fadhilah, Aziz, Hills, & Foo, 2014) were significantly associated with overweight and obesity. Obese adolescents had higher energy intake, and the mean energy intake was above the RNI compared to normal weight and underweight counterparts (Fara Wahida, Chin, & Barakatun Nisak, 2012; Soo et al., 2011). Typical eating patterns such as meal skipping and snacking are known to be associated with overweight and obesity (Dixit et al., 2014). Meanwhile, physical activity is very much related to energy expenditure whereby an increase in moderate-vigorous physical activities predicts reduced in BMI-z-score (Trinh, Campbell, Ukoumunne, Gerner, & Wake, 2013). With the different setting in day-school hostels, do they have the similar dietary practice and physical activity?

Meanwhile, knowledge, attitude, and practice on healthy lifestyle have been reported as factors associated with overweight and obesity (DeBar et al., 2012; Noor Aini et al., 2006; Wilson, 2007). However, previous study revealed that knowledge had a positive association with attitude but no significant association with practice (Jha, Bajracharya, & Shankar, 2013), suggesting that practice on healthy lifestyle might not in accordance with their knowledge on healthy lifestyle. It is recommended that study to be undertaken among adolescents living in day-school hostels as little is known on the level of knowledge, attitude and practice on healthy lifestyle among this group adolescents.

Concern was also given to the body image disturbances whereby higher body image discrepancy scores were more likely to have greater BMI (Farah Wahida, Mohd Nasir, & Hazizi, 2011). This is particularly obvious during adolescence as they experience significant physical changes in their body and more concern on their body image (Khor et al., 2009). Moreover, adolescents living in day-school hostels might have more peers influence as they spend greater time with peers as compared to parents. Hutchinson and Rapee (2007) had highlighted that the importance of the peer environment in body image during adolescence, specifically, girls' perceptions of their peers' body image predicted their own level of body image concern. Therefore, there is a need to understand the influences of body image concern on their body weight status among adolescents living in day-school hostels.

In addition, psychosocial behaviors such as stages of change, self-efficacy, perceived benefits and perceived barriers to healthy diet (Vereecken et al., 2009) and exercise (Lee, Nigg, Diclemente, & Courneya, 2001) were significantly associated with overweight and obesity. A local cross-sectional study reported that respondents who are in action or/maintenance stage had higher self-efficacy and perceived benefits on fruit and vegetable intake (Wong, Zalilah, Mirilanili, & Mohd Nasir, 2014). Moreover, self-efficacy was found to be positively and directly associated with physical activity level in adolescents (de Farias Júnior, Florindo, Santos, Mota, & Barros, 2014). However, little is known about these psychological factors in relation to body weight status among adolescents living in day-school hostels.

In order to fulfill the current knowledge gap, this study assessed the prevalence of body weight status among a representative sample of Malaysian adolescents living in day-school hostels. It also aimed to investigate the possible associations between knowledge, attitude and practice on healthy lifestyle, health behavioral change on nutrition and exercise, dietary practice, physical activity as well as body image with BMI-for-age among these adolescents. Furthermore, the contribution of the above factors to BMI-for-age is determined.

1.3 Significance of the Study

The findings of this study can contribute and update the data on the prevalence of body weight status among adolescents living in day-school hostels in Malaysia. For example, the findings of the study can serve as one of the reference sources for the Ministry of

Health on policy and program planning regarding the issue of malnutrition among adolescents living in day-school hostels in Malaysia. In addition, dietary practice data could be used by Ministry of Health in understanding the dietary practice of the adolescents, and further help in improving menu planning in school hostels.

Since the problem of under- and over-nutrition exists among adolescents in Malaysia, a study on factors associated with body weight status among adolescents is both timely and relevant. This study also determines the potential factors that contribute towards the BMI-for-age of the adolescents. Generally, this study provides an overall view of the factors relating to body weight status, as dietary practice, physical activity, behavioral change, body image, knowledge, attitude and practice on healthy lifestyle are taken into consideration. Moreover, this study provides information concerning factors which are more contributive to the variation in the body weight status of adolescents.

The result of the study is beneficial to health practitioners, such as nutritionists in providing advice to the community on concerning healthy lifestyle practice through the improvement in knowledge, attitude and practice on healthy lifestyle, modification on dietary practice and physical activity, understanding on behavioral change as well as the correct perception of body image. Additionally, policy makers, health agencies, program planners and community leaders can use this information as a reference for future research. For example, the results can serve as baseline data to develop an appropriate health intervention program to educate and promote healthy living and active lifestyle among adolescents living in day-school hostels. At the same time, findings from this study can also be used to develop a module focused on healthy living and active lifestyle as cores for any nutritional intervention program.

1.4 Objectives of the Study

1.4.1 General Objective

To determine the factors associated with BMI-for-age among adolescents living in day-school hostels in Malaysia.

1.4.2 Specific Objectives

1. To determine the body weight status among adolescents living in day-school hostels in Malaysia.
2. To determine the association between socio-demographic characteristics, dietary practice, physical activity, knowledge, attitude and practice (KAP) on healthy lifestyle, transtheoretical model of behavioral change on healthy diet and exercise, body image factors and BMI-for-age among adolescents living in day-school hostels in Malaysia.

3. To determine the contribution of socio-demographic characteristics, dietary practice, physical activity, knowledge, attitude and practice (KAP) on healthy lifestyle, transtheoretical model of behavioral change on healthy diet and exercise, body image factors towards BMI-for-age among adolescents living in day-school hostels in Malaysia.

1.5 Null Hypothesis

1. There is no significant association between socio-demographic characteristics, dietary practice, physical activity, knowledge, attitude and practice (KAP) on healthy lifestyle, transtheoretical model of behavioral change on healthy diet and exercise, body image factors and BMI-for-age among adolescents living in day-school hostels in Malaysia.
2. There is no significant contribution of socio-demographic characteristics, dietary practice, physical activity, knowledge, attitude and practice (KAP) on healthy lifestyle, transtheoretical model of behavioral change on healthy diet and exercise, body image factors and BMI-for-age among adolescents living in day-school hostels in Malaysia.

1.6 Conceptual Framework

Overweight and obesity prevention is necessary because obesity in adolescence is a strong predictor of obesity in adulthood. Moreover, overweight and obesity is associated with a range of serious complication such as type 2 diabetes, cardiovascular disease, and psychosocial problem. Hence, it is imperative to determine the prevalence and factors associated to BMI-for-age among adolescents living in day-school hostels in Malaysia as depicted in Figure 1.1. In the present study, knowledge, attitude and practice on healthy lifestyle, behavioral change, dietary practice, physical activity, body image, are proposed as influencing factors that may predict overweight and obesity among adolescents living in day-school hostels in Malaysia.

Knowledge is considered a prerequisite to intentional performance of health-related behaviors (Worsley, 2002). As knowledge in the health behaviors domain accumulates, changes in attitude are initiated. When changes in attitude accumulate over a period, it results in behavioral changes. For example, knowledge in nutrition knowledge was positively and significantly related to attitude (Webb & Beckford, 2014) and affect the dietary quality (Lin, Yang, Hang, & Pan, 2007). Therefore, this further predicts the occurrences of overweight and obesity.

The Transtheoretical Model (TTM) uses several constructs from other health behavior theories which offer a view of when, how and why people change their behavior (Prochaska & Velicer, 1997). This model includes two levels: i) the stages of change

(SOC), which reflect the temporal dimension of the behavior, and ii) a set of constructs that explain how people evolve along the SOC. These are named decisional balance, representing the perceived benefits and perceived barriers of engaging in behavior, and self-efficacy, reflecting the person's confidence in performing health behavioral change (Prochaska & Velicer, 1997).

As for the dietary practice, the present study includes eating behavior as a risk factor for overweight and obesity. Many studies have confirmed the association between meal skipping behavior and overweight (de Gouw et al., 2010; Stea, Vik, Bere, Svendsen, & Oellingrath, 2014). Adolescents who ate five meals a day have a lower risk for overweight/ obesity (Jääskeläinen et al., 2013) whereas breakfast skippers have higher body mass index-for-age z scores (Deshmukh-Taskar et al., 2010b). Indeed, meal skipping contributes to frequent snacking behavior which in turn contributes to higher energy intake (Teo, Sedek, & Zalifah, 2012).

On the other hand, physical activity, which encompasses physical activity level and energy balance (Al-Haifi et al., 2013; de Gouw et al., 2010; Pahkala et al., 2013) are known to be associated with overweight and obesity. Sustained increase in moderate-vigorous physical activities predicts reduced BMI scores (z-score) (Trinh et al., 2013). Watching more television and using the computer more were associated with overweight and obesity (de Gouw et al., 2010).

The body image perception is also included in the present study. Previous findings have identified the perception of body weight, and perception of body size are directly and indirectly contributing to overweight and obesity (Alwan, Viswanathan, Paccaud, & Bovet, 2011; Perkins, Perkins, & Craig, 2010; Perkins, Perkins, & Craig, 2014). For example, a study by Perkin, Perkin and Craig (2010) found that overestimating peer weight norms was associated with greater risk of being overweight and underestimating peer weight norms was associated with greater risk for being underweight while accurate perception were more likely to have appropriate weight-control behavior (Alwan et al., 2011).

In summary, as shown in Figure 1.1, factors explaining the development of BMI-for-age are supported by previous studies and empirical literature. Hence, the present study would focus on knowledge, attitude and practice on healthy lifestyle, behavioral change, dietary practice, physical activity and body image as independent variables which contribute to body weight status among adolescents living in day-school hostels in Malaysia.




Figure 1.1: Conceptual framework

REFERENCES

- Aboshkair, K. A., Amri, S. B., Yee, K. L., & Samah, B. B. A. (2011). Assessment of weight status of secondary school children in Selangor, Malaysia. *Australian Journal of Basic & Applied Sciences*, 5(9), 1675-1682.
- Afrifa-Anane, E., Agyemang, C., Codjoe, S. N., Ogedegbe, G., & Aikins, A. d.-G. (2015). The association of physical activity, body mass index and the blood pressure levels among urban poor youth in Accra, Ghana. *BMC Public Health*, 15(1), 269.
- Ainsworth, B. E., Haskell, W. L., Herrmann, S. D., Meckes, N., Bassett, D. R., Tudor-Locke, C., Greer, J. L., Vezina, J., Whitt-Glover, M. C., & Leon, A. S. (2011). 2011 compendium of physical activities: a second update of codes and MET values. *Medicine and Science in Sports and Exercise*, 43(8), 1575-1581.
- Al-Haifi, A. R., Al-Fayez, M. A., Al-Athari, B. I., Al-Ajmi, F. A., Allafi, A. R., Al-Hazzaa, H. M., & Musaiger, A. O. (2013). Relative contribution of physical activity, sedentary behaviors, and dietary habits to the prevalence of obesity among Kuwaiti adolescents. *Food and Nutrition Bulletin*, 34(1).
- Al - Sendi, A., Shetty, P., & Musaiger, A. (2004). Body weight perception among Bahraini adolescents. *Child: Care, Health and Development*, 30(4), 369-376.
- Al Junaibi, A., Abdulle, A., Sabri, S., Hag-Ali, M., & Nagelkerke, N. (2013). The prevalence and potential determinants of obesity among school children and adolescents in Abu Dhabi, United Arab Emirates. *International Journal of Obesity*, 37(1), 68-74.
- Alp, H., Eklioglu, B. S., Atabek, M. E., Karaarslan, S., Baysal, T., Altin, H., Karataş, Z., & Şap, F. (2014). Evaluation of epicardial adipose tissue, carotid intima-media thickness and ventricular functions in obese children and adolescents Alp, Hayrullah. *Journal of Pediatric Endocrinology and Metabolism*, 27(9-10), 827-835.
- Alwan, H., Viswanathan, B., Paccaud, F., & Bovet, P. (2011). Is accurate perception of body image associated with appropriate weight-control behavior among adolescents of the Seychelles. *Journal of Obesity*, 2011.
- Andegiorgish, A. K., Wang, J., Zhang, X., Liu, X., & Zhu, H. (2012). Prevalence of overweight, obesity, and associated risk factors among school children and adolescents in Tianjin, China. *European Journal of Pediatrics*, 171(4), 697-703.
- Anuradha, R., Sathyavathi, R., Reddy, T. M., Hemalatha, R., Sudhakar, G., Geetha, P., & Reddy, K. K. (2011). Effect of social and environmental determinants on overweight and obesity prevalence among adolescent school children. *Indian Journal of Endocrinology and Metabolism*, 19(2), 283.
- Aounallah-Skhiri, H., Traissac, P., El Ati, J., Eymard-Duvernay, S., Landais, E., Achour, N., Delpeuch, F., Ben Romdhane, H., & Maire, B. (2011). Nutrition transition among adolescents of a south-Mediterranean country: dietary patterns,

association with socio-economic factors, overweight and blood pressure. A cross-sectional study in Tunisia. *Nutrition Journal*, 10(38), 1-17.

- Baharudin, A., Zainuddin, A. A., Manickam, M., Ambak, R., Ahmad, M. H., Naidu, B. M., Man, C. S., Ying, C. Y., Saad, H. A., & Ahmad, N. A. (2014). Factors associated with physical inactivity among school-going adolescents data from the Malaysian School-Based Nutrition Survey 2012. *Asia-Pacific Journal of Public Health*.
- Baharudin, A., Zainuddin, A. A., Selamat, R., Ghaffar, S. A., Khor, G. L., Poh, B. K., Karim, N. A., Kee, C. C., Ng, C. K., & Ahmad, N. A. (2013). Malnutrition among Malaysian Adolescents: Findings from National Health and Morbidity Survey (NHMS) 2011. *International Journal of Public Health Research*, 3(2), 282-289.
- Bandura, A. (1977). Self-efficacy: toward a unifying theory of behavioral change. *Psychological Review*, 84(2), 191.
- Bandura, A. (1978). Self-efficacy: Toward a unifying theory of behavioral change. *Advances in Behaviour Research and Therapy*, 1(4), 139-161.
- Bauman, A. E., Reis, R. S., Sallis, J. F., Wells, J. C., Loos, R. J., & Martin, B. W. (2012). Correlates of physical activity: why are some people physically active and others not? *The Lancet*, 380(9838), 258-271.
- Berry, T., Naylor, P., & Wharf-Higgins, J. (2005). Stages of change in adolescents: an examination of self-efficacy, decisional balance, and reasons for relapse. *Journal of Adolescent Health*, 37(6), 452-459.
- Bo, S., De Carli, L., Venco, E., Fanzola, I., Maiandi, M., De Michieli, F., Durazzo, M., Beccuti, G., Cavallo-Perin, P., & Ghigo, E. (2014). Impact of Snacking Pattern on Overweight and Obesity Risk in A Cohort of 11-13-y Adolescents. *Journal of Pediatric Gastroenterology and Nutrition*.
- Boutelle, K., Neumark-Sztainer, D., Story, M., & Resnick, M. (2002). Weight control behaviors among obese, overweight, and nonoverweight adolescents. *Journal of Pediatric Psychology*, 27(6), 531-540.
- Boyce, W. T., & Boyce, J. C. (1983). Acculturation and changes in health among Navajo boarding school students. *Social Science and Medicine*, 17(4), 219-226.
- Brambilla, P., Bedogni, G., Moreno, L., Goran, M., Gutin, B., Fox, K., Peters, D., Barbeau, P., De Simone, M., & Pietrobelli, A. (2006). Crossvalidation of anthropometry against magnetic resonance imaging for the assessment of visceral and subcutaneous adipose tissue in children. *International Journal of Obesity*, 30(1), 23-30.
- Brixval, C. S., Rayce, S. L., Rasmussen, M., Holstein, B. E., & Due, P. (2012). Overweight, body image and bullying—an epidemiological study of 11-to 15-years olds. *European Journal of Public Health*, 22(1), 126-130.

- Bühren, K., von Ribbeck, L., Schwarte, R., Egberts, K., Pfeiffer, E., Fleischhaker, C., Wewetzer, C., Kennes, L. N., Dempfle, A., & Herpertz-Dahlmann, B. (2013). Body mass index in adolescent anorexia nervosa patients in relation to age, time point and site of admission. *European Child and Adolescent Psychiatry*, 22(7), 395-400.
- Chambliss, C. A., & Murray, E. J. (1979). Efficacy attribution, locus of control, and weight loss. *Cognitive Therapy and Research*, 3(4), 349-353.
- Chen, L.-J., Fox, K. R., Haase, A., & Wang, J.-M. (2006). Obesity, fitness and health in Taiwanese children and adolescents. *European Journal of Clinical Nutrition*, 60(12), 1367-1375.
- Chen, S., Soo, K., Azriani, A., Van Rostenberghe, H., & Sakinah, H. (2012). Nutrition knowledge, attitude and practice of teachers in rehabilitation centres in northern Malaysia. *Malaysian Journal of Nutrition*, 18(2), 185-205.
- Chin, Y. S., & Mohd Nasir, M. T. (2009). Eating behaviors among female adolescents in Kuantan district, Pahang, Malaysia. *Pakistan Journal of Nutrition*, 8(4), 425-432.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences 3rd edition*. New York, NY: Academic.
- Cole, T. J., Bellizzi, M. C., Flegal, K. M., & Dietz, W. H. (2000). Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ Clinical Research*, 320(7244), 1240.
- Cole, T. J., Flegal, K. M., Nicholls, D., & Jackson, A. A. (2007). Body mass index cut offs to define thinness in children and adolescents: international survey. *BMJ (Clinical Research Ed.)*, 335(7612), 194.
- Collins, A. E., Pakiz, B., & Rock, C. L. (2008). Factors associated with obesity in Indonesian adolescents. *International Journal of Pediatric Obesity*, 3(1), 58-64.
- Cook, C., Swinburn, B., & Stewart, J. (2001). Changing risk behaviours for non-communicable disease in New Zealand working men-is workplace intervention effective? *New Zealand Medical Journal*, 114(1130), 175-178.
- Crocker, P., Bailey, D. A., Faulkner, R. A., Kowalski, K. C., & McGRATH, R. (1997). Measuring general levels of physical activity: preliminary evidence for the Physical Activity Questionnaire for Older Children. *Medicine and Science in Sports and Exercise*, 29(10), 1344-1349.
- Cutler, G. J., Flood, A., Hannan, P., & Neumark-Sztainer, D. (2011). Multiple sociodemographic and socioenvironmental characteristics are correlated with major patterns of dietary intake in adolescents. *Journal of the American Dietetic Association*, 111(2), 230-240.
- Cuypers, K., Kvaløy, K., Bratberg, G., Midthjell, K., Holmen, J., & Holmen, T. L. (2012). Being normal weight but feeling overweight in adolescence may affect weight

development into young adulthood—an 11-year followup: The Hunt Study, Norway. *Journal of Obesity*, 2012.

- Cynthia, J., Zalilah, M., & Lim, M. (2013). Relationship between family meals away from home and nutritional status of adolescents. *Malaysian Journal of Nutrition*, 19(1), 25-35.
- Daniel, W. W. (1999). Determination of sample size for estimating proportions *Biostatistics: A Foundation for Analysis in Health* (pp. 183). New York: John Wiley & Sons Inc.
- Davis, C., Levitan, R. D., Muglia, P., Bewell, C., & Kennedy, J. L. (2004). Decision - making deficits and overeating: A risk model for obesity. *Obesity Research*, 12(6), 929-935.
- De-Jong, E., Visscher, T., HiraSing, R., Heymans, M., Seidell, J., & Renders, C. (2013). Association between TV viewing, computer use and overweight, determinants and competing activities of screen time in 4-to 13-year-old children. *International Journal of Obesity*, 37(1), 47-53.
- de Farias Júnior, J. C., Florindo, A. A., Santos, M. P., Mota, J., & Barros, M. V. (2014). Perceived environmental characteristics and psychosocial factors associated with physical activity levels in adolescents from Northeast Brazil: structural equation modelling analysis. *Journal of Sports Sciences*, 32(10), 963-973.
- de Gouw, L., Klepp, K. I., Vignerova, J., Lien, N., Steenhuis, I. H., & Wind, M. (2010). Associations between diet and (in) activity behaviours with overweight and obesity among 10-18-year-old Czech Republic adolescents. *Public Health Nutrition*, 13(10A), 1701-1707.
- de Onis, M., Onyango, A. W., Borghi, E., Siyam, A., Nishida, C., & Sieckmann, J. (2007). Development of a WHO growth reference for school-aged children and adolescents. *Bulletin of the World Health Organization*, 85:660-667.
- DeBar, L. L., Stevens, V. J., Perrin, N., Wu, P., Pearson, J., Yarborough, B. J., Dickerson, J., & Lynch, F. (2012). A primary care-based, multicomponent lifestyle intervention for overweight adolescent females. *Pediatrics*, 129(3), 611-620.
- Deshmukh-Taskar, P. R., Nicklas, T. A., O'Neil, C. E., Keast, D. R., Radcliffe, J. D., & Cho, S. (2010a). The relationship of breakfast skipping and type of breakfast consumption with nutrient intake and weight status in children and adolescents: the National Health and Nutrition Examination Survey 1999-2006. *Journal of the American Dietetic Association*, 110(6), 869-878.
- Deshmukh-Taskar, P. R., Nicklas, T. A., O'Neil, C. E., Keast, D. R., Radcliffe, J. D., & Cho, S. (2010b). The relationship of breakfast skipping and type of breakfast consumption with nutrient intake and weight status in children and adolescents: the National Health and Nutrition Examination Survey 1999-2006. *Journal of the American Dietetic Association*, 110(6), 869-878.

- Dietz, W. H. (1998). Childhood weight affects adult morbidity and mortality. *The Journal of Nutrition*, 128(2), 411S-414S.
- Dishman, R. K., Dunn, A. L., Sallis, J. F., Vandenberg, R. J., & Pratt, C. A. (2010). Social-cognitive correlates of physical activity in a multi-ethnic cohort of middle-school girls: two-year prospective study. *Journal of Pediatric Psychology*, 35(2), 188-198.
- Dixit, S., Singh, J., Kant, S., Agarwal, G., Dubey, A., & Kumari, N. (2014). A cross-sectional study on predictors and significance of eating behavior of adolescent girls. *Vulnerable Children and Youth Studies*, 9(1), 10-16.
- Doustmohammadian Jr, A., Dorostymotlagh, A., Keshavarz, A., Sadrzadehyeganeh, H., & Mohammadpour-Ahrangani, B. (2009). Socio-demographic factors associated with Body Mass Index of female adolescent students in Semnan City, Iran. *Malaysian Journal of Nutrition*, 15(1), 27-35.
- Drake, K. M., Beach, M. L., Longacre, M. R., MacKenzie, T., Titus, L. J., Rundle, A. G., & Dalton, M. A. (2012). Influence of sports, physical education, and active commuting to school on adolescent weight status. *Pediatrics*, 130(2), e296-e304.
- Duchin, O., Marin, C., Mora-Plazas, M., Mendes de Leon, C., Lee, J. M., Baylin, A., & Villamor, E. (2015). A prospective study of body image dissatisfaction and BMI change in school-age children. *Public Health Nutrition*, 18(02), 322-328.
- Dumith, S. C., Gigante, D. P., Domingues, M. R., & Kohl, H. W. (2011). Physical activity change during adolescence: a systematic review and a pooled analysis. *International Journal of Epidemiology*, 40(3), 685-698.
- Dunn, R. A., Tan, A. K., & Nayga, R. M. (2012). Obesity inequality in Malaysia: decomposing differences by gender and ethnicity using quantile regression. *Ethnicity and Health*, 17(5), 493-511.
- Duong, H. T., & Roberts, R. E. (2014). Perceived weight in youths and risk of overweight or obesity six years later. *Journal of Psychosomatic Research*, 76(1), 23-27.
- Eisenberg, M. E., Neumark-Sztainer, D., & Paxton, S. J. (2006). Five-year change in body satisfaction among adolescents. *Journal of Psychosomatic Research*, 61(4), 521-527.
- Fara Wahida, R., Chin, Y. S., & Barakatun Nisak, M. Y. (2012). Obesity-related behaviors of Malaysian adolescents: a sample from Kajang district of Selangor state. *Nutrition Research and Practice*, 6(5), 458-465.
- Farah Wahida, Z., Mohd Nasir, M., & Hazizi, A. (2011). Physical activity, eating behaviour and body image perception among young adolescents in Kuantan, Pahang, Malaysia. *Malaysian Journal of Nutrition*, 17(3), 325-336.

- Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using G* Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods*, 41(4), 1149-1160.
- Ferrar, K., & Olds, T. (2010). Thin adolescents: Who are they? What do they do? Socio-demographic and use-of-time characteristics. *Preventive Medicine*, 51(3), 253-258.
- Finucane, M. M., Stevens, G. A., Cowan, M. J., Danaei, G., Lin, J. K., Paciorek, C. J., Singh, G. M., Gutierrez, H. R., Lu, Y., & Bahalim, A. N. (2011). National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9· 1 million participants. *The Lancet*, 377(9765), 557-567.
- Fitzgibbon, M. L., & Beech, B. M. (2009). The role of culture in the context of school-based BMI screening. *Pediatrics*, 124(1), 50-62.
- Food and Agriculture Organization, World Health Organization, & United Nations University [FAO/WHO/UNU]. (2004). *Human Energy Requirements: Report of a Joint FAO/WHO/UNU Expert Consultation: Rome, 17-24 October 2001*: Food & Agriculture Org.
- Food and Agriculture Organization/World Health Organization/United Nations University (FAO/WHO/UNU). (2005). Human energy requirements: Report of a joint FAO/WHO/UNU Expert Consultation. *Food and Nutrition Bulletin*, 26, 166.
- Franko, D., Striegel-Moore, R., Thompson, D., Affenito, S., Schreiber, G., Daniels, S., & Crawford, P. (2008). The relationship between meal frequency and body mass index in black and white adolescent girls: more is less. *International Journal of Obesity*, 32(1), 23-29.
- Fredrickson, J., Kremer, P., Swinburn, B., de Silva-Sanigorski, A., & McCabe, M. (2013). Biopsychosocial correlates of weight status perception in Australian adolescents. *Body Image*, 10(4), 552-557.
- Fulkerson, J. A., Neumark - Sztainer, D., Hannan, P. J., & Story, M. (2008). Family meal frequency and weight status among adolescents: cross - sectional and 5 - year longitudinal associations. *Obesity*, 16(11), 2529-2534.
- Garcia , A. W., Antonakos, C. L., & Ronis, D. L. (1998). Changes in physical activity beliefs and behaviors of boys and girls across the transition to junior high school. *Journal of Adolescent Health*, 22(5), 394-402.
- Ghoncheh, R., Gould, M. S., Twisk, J. W., Kerkhof, A. J., & Koot, H. M. (2016). Efficacy of Adolescent Suicide Prevention E-Learning Modules for Gatekeepers: A Randomized Controlled Trial. *JMIR Mental Health*, 3(1), e8.

- Ghosh, A. (2011). Rural–urban comparison in prevalence of overweight and obesity among children and adolescents of Asian Indian origin. *Asia-Pacific Journal of Public Health*, 23(6), 928-935.
- Glanz, K., Rimer, B. K., & Viswanath, K. (2008). *Health behavior and health education: theory, research, and practice*: John Wiley & Sons.
- Glaser, D. R., Haaga, D. A., Hannallah, L., Field, S. E., Kozlosky, M., Reynolds, J., Yanovski, J. A., & Tanofsky - Kraff, M. (2013). Self - efficacy beliefs and eating behavior in adolescent girls at - risk for excess weight gain and binge eating disorder. *International Journal of Eating Disorders*, 46(7), 663-668.
- Gonzalez-Casanova, I., Sarmiento, O. L., Gazmararian, J. A., Cunningham, S. A., Martorell, R., Pratt, M., & Stein, A. D. (2013). Comparing three body mass index classification systems to assess overweight and obesity in children and adolescents. *Revista Panamericana de Salud Publica*, 33(5), 349-355.
- Goyal, R., Mehta, P., & Kaur, G. (2012). Nutritional status and menarche in adolescents of Punjab. *Journal of Life Science*, 4(1), 63-66.
- Gupta, A. K., & Wang, H. (2009). *Getting China and India right: Strategies for leveraging the world's fastest growing economies for global advantage*. New York: John Wiley & Sons.
- Gupta, D. K., Shah, P., Misra, A., Bharadwaj, S., Gulati, S., Gupta, N., Sharma, R., Pandey, R. M., & Goel, K. (2011). Secular trends in prevalence of overweight and obesity from 2006 to 2009 in urban asian Indian adolescents aged 14-17 years. *PLoS ONE*, 6(2), e17221.
- Hagler, A. S., Norman, G. J., Zabinski, M. F., Sallis, J. F., Calfas, K. J., & Patrick, K. (2007). Psychosocial correlates of dietary intake among overweight and obese men. *American Journal of Health Behavior*, 31(1), 3-12.
- Hampel, J., Heaton, C., & Taylor, C. (2003). Snacking patterns influence energy and nutrient intakes but not body mass index. *Journal of Human Nutrition and Dietetics*, 16(1), 3-11.
- Hargreaves, M. K., Schlundt, D. G., Buchowski, M. S., Hardy, R. E., Rossi, S. R., & Rossi, J. S. (1999). Stages of change and the intake of dietary fat in African-American women: improving stage assignment using the Eating Styles Questionnaire. *Journal of the American Dietetic Association*, 99(11), 1392-1399.
- Hassapidou, M., Fotiadou, E., Maglara, E., & Papadopoulou, S. K. (2006). Energy intake, diet composition, energy expenditure, and body fatness of adolescents in northern Greece. *Obesity*, 14(5), 855-862.
- Hiew, C. C., Chin, Y. S., Chan, Y. M., & Mohd Nasir, M. T. (2015). Development and Validation of Knowledge, Attitude and Practice on Healthy Lifestyle Questionnaire (KAP-HLQ) for Malaysian Adolescents. *J Nutr Health Sci*, 2(4), 407.

- Hills, A. P., Okely, A. D., & Baur, L. A. (2010). Addressing childhood obesity through increased physical activity. *Nature Reviews Endocrinology*, 6(10), 543-549.
- Horikawa, C., Kodama, S., Yachi, Y., Heianza, Y., Hirasawa, R., Ibe, Y., Saito, K., Shimano, H., Yamada, N., & Sone, H. (2011). Skipping breakfast and prevalence of overweight and obesity in Asian and Pacific regions: a meta-analysis. *Preventive Medicine*, 53(4), 260-267.
- Hutchinson, D. M., & Rapee, R. M. (2007). Do friends share similar body image and eating problems? The role of social networks and peer influences in early adolescence. *Behaviour Research and Therapy*, 45(7), 1557-1577.
- Huybrechts, I., Grammatikaki, E., Kersting, M., Vyncke, K., Cuenca, M., Gonzalez-Gross, M., Gottrand, F., Moreno, L., Molnar, D., Hallstrom, L., Dallongeville, J., Plada, M., Leclercq, C., Widhalm, K., Manios, Y., & Henauw, S. (2012). Dietary under-reporting by overweight and obese adolescents: Results from the HELENA study. *Public Health Nutrition*, 15(8), 1555.
- Institute for Public Health [IPH]. (2008). *Nutritional Status. In: The Third National Health & Morbidity Survey 2006 (NHMS III)*. National Institutes of Health, Ministry of Health Malaysia.
- Institute for Public Health [IPH]. (2011). *National Health and Morbidity Survey 2011 (NHMS) (Vol. II: Non-Communicable Disease)*. National Institutes of Health, Ministry of Health Malaysia.
- Institute for Public Health [IPH]. (2013). *National Health and Morbidity Survey: Malaysia School Based Nutrition Survey 2012*. Kuala Lumpur: Ministry of Health Malaysia.
- Institute for Public Health [IPH]. (2015). *National Health and Morbidity Survey 2015 (NHMS 2015). Vol. II: Non-Communicable Diseases, Risk Factors & Other Health Problems*. Kuala Lumpur: Ministry of Health Malaysia.
- Intiful, F. D., Ogyiri, L., Asante, M., Mensah, A., Steele-Dadzie, R., & Boateng, L. (2013). Nutritional status of boarding and non-boarding children in selected schools in the Accra metropolis. *Journal of Biology, Agriculture and Healthcare*, 3(7), 156-162.
- Jääskeläinen, A., Schwab, U., Kolehmainen, M., Pirkola, J., Järvelin, M.-R., & Laitinen, J. (2013). Associations of meal frequency and breakfast with obesity and metabolic syndrome traits in adolescents of Northern Finland Birth Cohort 1986. *Nutrition, Metabolism and Cardiovascular Diseases*, 23(10), 1002-1009.
- Jha, N., Bajracharya, O., & Shankar, P. R. (2013). Knowledge, attitude and practice towards medicines among school teachers in Lalitpur district, Nepal before and after an educational intervention. *BMC Public Health*, 13(1), 652.
- Kearney, J. M., de Graaf, C., Damkjaer, S., & Engstrom, L. M. (1999). Stages of change towards physical activity in a nationally representative sample in the European Union. *Public Health Nutrition*, 2(1a), 115-124.

- Kelishadi, R., Alikhani, S., Delavari, A., Alaedini, F., Safaie, A., & Hojatzadeh, E. (2008). Obesity and associated lifestyle behaviours in Iran: findings from the first national non-communicable disease risk factor surveillance survey. *Public Health Nutrition*, 11(03), 246-251.
- Khambalia, A. Z., Lim, S. S., Gill, T., & Bulgiba, A. M. (2012). Prevalence and sociodemographic factors of malnutrition among children in Malaysia. *Food and Nutrition Bulletin*, 33(1), 31-42.
- Khor, G. L., Chee, W. S., Shariff, Z. M., Poh, B. K., Arumugam, M., Rahman, J. A., & Theobald, H. E. (2011). High prevalence of vitamin D insufficiency and its association with BMI-for-age among primary school children in Kuala Lumpur, Malaysia. *BMC Public Health*, 11, 95.
- Khor, G. L., Zalilah, M. S., Phan, Y. Y., Ang, M., Maznah, B., & Norimah, A. K. (2009). Perceptions of body image among Malaysian male and female adolescents. *Singapore Medical Journal*, 50(3), 303-311.
- Kim-Spoon, J., & Farley, J. P. (2014). *Adolescence and Early Adulthood Defining Prevention Science* (pp. 107).
- Kim, Y. h. (2004). Korean adolescents' exercise behavior and its relationship with psychological variables based on stages of change model. *Journal of Adolescent Health*, 34(6), 523-530.
- Kim, Y. H., & Cardinal, B. J. (2009). Effects of a transtheoretical model-based stage-matched intervention to promote physical activity among Korean adults. *International journal of clinical and health psychology*, 9(2), 259-273.
- Kim, Y. H., & Cardinal, B. J. (2010). Psychosocial correlates of Korean adolescents' physical activity behavior. *Journal of Exercise Science & Fitness*, 8(2), 97-104.
- Kimm, S., Glynn, N. W., Kriska, A. M., Fitzgerald, S. L., Aaron, D. J., Similo, S. L., Mc Mahon, R. P., & Barton, B. A. (2000). Longitudinal changes in physical activity in a biracial cohort during adolescence. *Medicine and Science in Sports and Exercise*, 32(8), 1445-1454.
- Koletzko, B., & Toschke, A. M. (2010). Meal patterns and frequencies: do they affect body weight in children and adolescents? *Critical Reviews in Food Science and Nutrition*, 50(2), 100-105.
- Kostanjevec, S., Jerman, J., & Koch, V. (2012). The influence of nutrition education on the food consumption and nutrition attitude of schoolchildren in Slovenia. *US-China Education Review*, 11, 953-964.
- Kromhout, D., Saris, W. H., & Horst, C. H. (1988). Energy intake, energy expenditure, and smoking in relation to body fatness: the Zutphen Study. *The American Journal of Clinical Nutrition*, 47(4), 668-674.

- Kuczmariski, R. J., Ogden, C. L., Grummer-Strawn, L. M., Flegal, K. M., Guo, S. S., Wei, R., Mei, Z., Curtin, L. R., Roche, A. F., & Johnson, C. L. (2000). CDC growth charts: United States. *Advance Data*, 314, 1-27.
- Laforge, R. G., Greene, G. W., & Prochaska, J. O. (1994). Psychosocial factors influencing low fruit and vegetable consumption. *Journal of Behavioral Medicine*, 17(4), 361-374.
- Law, L. S., Mohd Nasir, M. T., & Hazizi, A. S. (2013). Factors associated with Breakfast Skipping among School-going Adolescents in Sarawak, Malaysia. *Malaysian Journal of Nutrition*, 19(3), 401-407.
- Laxmaiah, A., Nagalla, B., Vijayaraghavan, K., & Nair, M. (2007). Factors affecting prevalence of overweight among 12 - to 17 - year - old urban adolescents in Hyderabad, India. *Obesity*, 15(6), 1384-1390.
- Lee, H. G., & Kim, Y. H. (2014). Effects of an obesity intervention integrating physical activity and psychological strategy on BMI, physical activity, and psychological variables in male obese adolescents. *Behavioral Medicine*, 1-8.
- Lee, R. E., Nigg, C. R., Diclemente, C. C., & Courneya, K. S. (2001). Validating motivational readiness for exercise behavior with adolescents. *Research Quarterly for Exercise and Sport*, 72(4), 401-410.
- Lew, K., & Barlow, P. (2005). Dietary practices of adolescents in Singapore and Malaysia. *Singapore Medical Journal*, 46(6), 282-288.
- Li, Y.-P., Hu, X.-Q., Yang, X.-G., & Ma, G.-S. (2009). Application of the WHO growth reference (2007) to assess the nutritional status of children in China. *Biomedical and Environmental Sciences*, 22(2), 130-135.
- Lin, W., Yang, H.-C., Hang, C.-M., & Pan, W.-H. (2007). Nutrition knowledge, attitude, and behavior of Taiwanese elementary school children. *Asia Pacific Journal of Clinical Nutrition*, 16(S2), 534-546.
- Lindelof, A., Nielsen, C. V., & Pedersen, B. D. (2010). Obesity treatment—more than food and exercise: a qualitative study exploring obese adolescents' and their parents' views on the former's obesity. *International Journal of Qualitative Studies on Health and Well-being*, 5(2).
- Liu, J., Bennett, K. J., Harun, N., & Probst, J. C. (2008). Urban - rural differences in overweight status and physical inactivity among US children aged 10 - 17 Years. *The Journal of Rural Health*, 24(4), 407-415.
- López-Azpiazu, I., Martínez-González, M., Leon-Mateos, A., Kearney, J., Gibney, M., & Martínez, J. (2000). Stages of dietary change and nutrition attitudes in the Spanish population. *Public Health*, 114(3), 183-189.
- Lu Ann, A., & Llewellyn, J. C. (1996). *Designing and conducting health surveys: a comprehensive guide* (3rd ed.). United State of America: Jossey-Bass.

- Macfarlane, S. B. (1997). Conducting a descriptive survey: 2. Choosing a sampling strategy. *Tropical Doctor*, 27(1), 14-21.
- Magbuhat, R., Borazon, E. Q., & Villarino, B. J. (2011). Food preferences and dietary intakes of Filipino adolescents in metro Manila, the Philippines. *Malaysian Journal of Nutrition*, 17(1), 31-41.
- Maher, C., Olds, T. S., Eisenmann, J. C., & Dollman, J. (2012). Screen time is more strongly associated than physical activity with overweight and obesity in 9 - to 16 - year - old Australians. *Acta Paediatrica*, 101(11), 1170-1174.
- Maiti, S., Chatterjee, K., De, D., Ali, K. M., Bera, T. K., Jana, K., & Ghosh, D. (2011). The impact of Nutritional Awareness Package (NAP) on secondary school students for the improvement of Knowledge, Attitudes and Practices (KAP) at rural areas of Paschim Medinipur, West Bengal. *Asian Journal of Medical Sciences*, 2(2), 87-92.
- Malabarey, O. T., Balayla, J., Klam, S. L., Shrim, A., & Abenheim, H. A. (2012). Pregnancies in young adolescent mothers: a population-based study on 37 million births. *Journal of Pediatric and Adolescent Gynecology*, 25(2), 98-102.
- Mamun, A. A., O'Callaghan, M. J., Williams, G., & Najman, J. M. (2013). Television watching from adolescence to adulthood and its association with BMI, waist circumference, waist-to-hip ratio and obesity: a longitudinal study. *Public Health Nutrition*, 16(01), 54-64.
- Marcus, B. H., Rakowski, W., & Rossi, J. S. (1992). Assessing motivational readiness and decision making for exercise. *Health Psychology*, 11(4), 257.
- Matsushita, Y., Yoshiike, N., Kaneda, F., Yoshita, K., & Takimoto, H. (2004). Trends in childhood obesity in Japan over the last 25 years from the national nutrition survey. *Obesity Research*, 12(2), 205-214.
- Maximova, K., McGrath, J. J., Barnett, T., O'Loughlin, J., Paradis, G., & Lambert, M. (2008). Do you see what I see? Weight status misperception and exposure to obesity among children and adolescents. *International Journal of Obesity*, 32(6), 1008-1015.
- Mikki, N., Abdul-Rahim, H. F., Awartani, F., & Holmboe-Ottesen, G. (2009). Prevalence and sociodemographic correlates of stunting, underweight, and overweight among Palestinian school adolescents (13-15 years) in two major governorates in the West Bank. *BMC Public Health*, 9(1), 485.
- Ministry of Education Malaysia [MoE]. (2012). Regular. *Day hostel management*. Retrieved from <http://www.moe.gov.my/en/harian>
- Mirmiran, P., Azadbakht, L., & Azizi, F. (2007). Dietary behaviour of Tehranian adolescents does not accord with their nutritional knowledge. *Public Health Nutrition*, 10(09), 897-901.

- Mohd Shariff, Z., Abu Samah, B., Paim, L., Ismail, M., Kasim, M. S., Othman, N., Hashim, N., Buhari, S. S., Jamil Osman, Z., & Hussein, M. (2008). Nutrition education intervention improves nutrition knowledge, attitude and practices of primary school children: a pilot study. *International Electronic Journal of Health Education, 11*(1), 119-132.
- Molnár, D., Jeges, S., Erhardt, E., & Schutz, Y. (1995). Measured and predicted resting metabolic rate in obese and nonobese adolescents. *The Journal of pediatrics, 127*(4), 571-577.
- Mond, J., Van den Berg, P., Boutelle, K., Hannan, P., & Neumark-Sztainer, D. (2011). Obesity, body dissatisfaction, and emotional well-being in early and late adolescence: findings from the project EAT study. *Journal of Adolescent Health, 48*(4), 373-378.
- Moreno, L., De Henauw, S., Gonzalez-Gross, M., Kersting, M., Molnar, D., Gottrand, F., Barrios, L., Sjöström, M., Manios, Y., & Gilbert, C. (2008). Design and implementation of the healthy lifestyle in Europe by nutrition in adolescence cross-sectional study. *International Journal of Obesity, 32*, S4-S11.
- Moy, F. M., Gan, Y. C., & Siti Zaleha, M. K. (2004). Body mass status of school children and adolescents in Kuala Lumpur, Malaysia. *Asia Pacific Journal of Clinical Nutrition, 13*(4).
- Nagl, M., Jacobi, C., Paul, M., Beesdo-Baum, K., Höfler, M., Lieb, R., & Wittchen, H.-U. (2016). Prevalence, incidence, and natural course of anorexia and bulimia nervosa among adolescents and young adults. *European Child and Adolescent Psychiatry, 1-16*.
- Naidu, B. M., Mahmud, S. Z., Ambak, R., Sallehuddin, S. M., Mutalip, H. A., Saari, R., Sahril, N., & Hamid, H. A. (2013). Overweight among primary school-age children in Malaysia. *Asia Pacific Journal of Clinical Nutrition, 22*(3), 408-415.
- National Coordinating Committee on Food and Nutrition [NCCFN]. (2005). *Recommended Nutrient Intake's for Malaysia 2005. A Report of The Technical Working Group on Nutritional Guidelines*. Putrajaya: Ministry of Health Malaysia.
- National Coordinating Committee on Food and Nutrition [NCCFN]. (2010). *Malaysia Dietary Guidelines*. Putrajaya: Ministry of Health Malaysia.
- Nelson, M. C., Gordon-Larsen, P., Song, Y., & Popkin, B. M. (2006). Built and social environments: associations with adolescent overweight and activity. *American Journal of Preventive Medicine, 31*(2), 109-117.
- Neumark-Sztainer, D. (2007). Addressing the spectrum of adolescent weight-related problems: engaging parents and communities. *Prevention Researcher, 14*(3), 11-14.
- Neumark-Sztainer, D., Wall, M., Guo, J., Story, M., Haines, J., & Eisenberg, M. (2006). Obesity, disordered eating, and eating disorders in a longitudinal study of

- adolescents: how do dieters fare 5 years later? *Journal of the American Dietetic Association*, 106(4), 559-568.
- Neumark-Sztainer, D., Wall, M. M., Haines, J. I., Story, M. T., Sherwood, N. E., & van den Berg, P. A. (2007). Shared risk and protective factors for overweight and disordered eating in adolescents. *American Journal of Preventive Medicine*, 33(5), 359-369.
- Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., Mullany, E. C., Biryukov, S., Abbafati, C., & Abera, S. F. (2014). Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet*, 384(9945), 766-781.
- Ng, S. W., Zaghoul, S., Ali, H., Harrison, G., & Popkin, B. M. (2011). The prevalence and trends of overweight, obesity and nutrition - related non - communicable diseases in the Arabian Gulf States. *Obesity Reviews*, 12(1), 1-13.
- Nguyen, P. V. N., Hong, T. K., Hoang, T., & Robert, A. R. (2013). High prevalence of overweight among adolescents in Ho Chi Minh City, Vietnam. *BMC Public Health*, 13(1), 141.
- Nicklas, T. A., Morales, M., Linares, A., Yang, S.-J., Baranowski, T., De Moor, C., & Berenson, G. (2004). Children's meal patterns have changed over a 21-year period: the Bogalusa Heart Study. *Journal of the American Dietetic Association*, 104(5), 753-761.
- Nigg, C. R., & Courneya, K. S. (1998). Transtheoretical model: Examining adolescent exercise behavior. *Journal of Adolescent Health*, 22(3), 214-224.
- Nigg, C. R., Geller, K. S., Motl, R. W., Horwath, C. C., Wertin, K. K., & Dishman, R. K. (2011). A research agenda to examine the efficacy and relevance of the transtheoretical model for physical activity behavior. *Psychology of Sport and Exercise*, 12(1), 7-12.
- Njelekela, M., Muhihi, A., Mpembeni, R., Anaeli, A., Chillo, O., Kubhoja, S., Lujani, B., Ngarashi, D., & Maghembe, M. (2015). *Knowledge and attitudes towards obesity among primary school children in Dar es Salaam, Tanzania* (Vol. 56).
- Noor Aini, M., Pon, L., Ong, F., Adeeb, N., Seri, S., Shamsuddin, K., Mohamad, A., Hapizah, N., Mokhtar, A., & Wan, H. (2006). Lifestyle intervention improved nutritional knowledge, dietary composition and health status of midlife Malaysian women. *Malaysian Journal of Nutrition*, 12(2), 125-146.
- Norcross, J. C., Krebs, P. M., & Prochaska, J. O. (2011). Stages of change. *Journal of Clinical Psychology*, 67(2), 143-154.
- Nothwehr, F., Snetselaar, L., Yang, J., & Wu, H. (2006). Stage of change for healthful eating and use of behavioral strategies. *Journal of the American Dietetic Association*, 106(7), 1035-1041.

- Nurliyana, A. R., Mohd Nasir, M. T., Zalilah, M. S., & Rohani, A. (2014). Dietary patterns and cognitive ability among 12-to 13 year-old adolescents in Selangor, Malaysia. *Public Health Nutrition*, 1-10.
- O'Dea, J. A., & Wilson, R. (2006). Socio-cognitive and nutritional factors associated with body mass index in children and adolescents: possibilities for childhood obesity prevention. *Health Education Research*, 21(6), 796-805.
- Ogden, C. L., Carroll, M. D., Kit, B. K., & Flegal, K. M. (2012a). Prevalence of obesity and trends in body mass index among US children and adolescents, 1999-2010. *JAMA*, 307(5), 483-490.
- Ogden, C. L., Carroll, M. D., Kit, B. K., & Flegal, K. M. (2012b). Prevalence of obesity in the United States, 2009-2010. *NCHS Data Brief*(82), 1-8.
- Ogden, C. L., Carroll, M. D., Kit, B. K., & Flegal, K. M. (2014). Prevalence of childhood and adult obesity in the United States, 2011-2012. *JAMA*, 311(8), 806-814.
- Oh, S.-Y. (2002). Psychosocial characteristics of stages of dietary fat reduction among Korean adults. *Ecology of Food and Nutrition*, 41(6), 515-532.
- Oosthuizen, D., Oldewage-Theron, W. H., & Napier, C. (2011). The impact of a nutrition programme on the dietary intake patterns of primary school children. *South African Journal of Clinical Nutrition*, 24(2).
- Ounpuu, S., Woolcott, D. M., & Greene, G. W. (2000). Defining stage-of-change for lower-fat eating. *Journal of the American Dietetic Association*, 100(6), 674-679.
- Owusu, A., Murdock, P. O. H., & Weatherby, N. L. (2007). Measuring Nutritional Intake of Adolescents in Ghana, West Africa. *International Electronic Journal of Health Education*, 10, 104-113.
- Pahkala, K., Hernelahti, M., Heinonen, O. J., Raitinen, P., Hakanen, M., Lagström, H., Viikari, J. S. A., Rönnemaa, T., Raitakari, O. T., & Simell, O. (2013). Body mass index, fitness and physical activity from childhood through adolescence. *British Journal of Sports Medicine*, 47(2), 71-77.
- Park, E. (2011). Overestimation and underestimation: adolescents' weight perception in comparison to BMI - based weight status and how it varies across socio - demographic factors. *Journal of School Health*, 81(2), 57-64.
- Pasic, M., Milanovic, I., Janic, S. R., Jurak, G., Soric, M., & Mirkov, D. (2014). Physical activity levels and energy expenditure in urban Serbian adolescents-preliminary study. *Nutricion Hospitalaria*, 30(5), 1044-1053.
- Paxton, S. J., Norris, M., Wertheim, E. H., Durkin, S. J., & Anderson, J. (2005). Body dissatisfaction, dating, and importance of thinness to attractiveness in adolescent girls. *Sex roles*, 53(9-10), 663-675.

- Peat, J., & Barton, B. (2008). *Medical statistics: A guide to data analysis and critical appraisal*: John Wiley & Sons.
- Perkins, J., Perkins, H., & Craig, D. (2010). Peer weight norm misperception as a risk factor for being over and underweight among UK secondary school students. *European Journal of Clinical Nutrition*, 64(9), 965-971.
- Perkins, J., Perkins, H. W., & Craig, D. W. (2014). Misperception of peer weight norms and its association with overweight and underweight status among adolescents. *Prevention Science*(1), 70-79.
- Phillips, S. M., Bandini, L. G., Naumova, E. N., Cyr, H., Colclough, S., Dietz, W. H., & Must, A. (2004). Energy - dense snack food intake in adolescence: longitudinal relationship to weight and fatness. *Obesity Research*, 12(3), 461-472.
- Pignatti, R., Bertella, L., Albani, G., Mauro, A., Molinari, E., & Semenza, C. (2006). Decision-making in obesity: a study using the Gambling Task. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*, 11(3), 126-132.
- Plotnikoff, R. C., Lippke, S., Johnson, S. T., Hotz, S. B., Birkett, N. J., & Rossi, S. R. (2009). Applying the stages of change to multiple low-fat dietary behavioral contexts. An examination of stage occupation and discontinuity. *Appetite*, 53(3), 345-353.
- Poh, B. K., Ismail, M. N., Ong, H. F., Norimah, A. K., & Safiah, M. Y. (2004). BMR Predictive Equations for Malaysian Adolescents Aged 12-18 Years. *Kuala Lumpur: Universiti Kebangsaan Malaysia*.
- Poh, B. K., Ng, B. K., Siti Haslinda, M. D., Nik Shanita, S., Wong, J. E., Budin, S. B., Ruzita, A. T., Ng, L. O., Khouw, I., & Norimah, A. K. (2013). Nutritional status and dietary intakes of children aged 6 months to 12 years: findings of the Nutrition Survey of Malaysian Children (SEANUTS Malaysia). *British Journal of Nutrition*, 110 (3), 21- 35.
- Pope, H., Phillips, K. A., & Olivardia, R. (2000). *The Adonis complex: The secret crisis of male body obsession*: Simon and Schuster.
- Popkin, B. M. (2001). Nutrition in transition: the changing global nutrition challenge. *Asia Pacific Journal of Clinical Nutrition*, 10(s1), S13-S18.
- Prochaska, J. O. (2006). Moving beyond the transtheoretical model. *Addiction*, 101(6), 768-774.
- Prochaska, J. O., & DeClemente, C. C. (1983). Stages and processes of self-change of smoking: Toward an integrative model of change. *Journal of Consulting and Clinical Psychology*, 51(3), 390-395.
- Prochaska, J. O., & Marcus, B. (1994). *The transtheoretical model: Applications to exercise* (I. R. Dishman Ed.). Human Kinetics.: Champaign, IL.

- Prochaska, J. O., & Velicer, W. F. (1997). The transtheoretical model of health behavior change. *American Journal of Health Promotion, 12*(1), 38-48.
- Prochaska, J. O., Velicer, W. F., Rossi, J. S., Goldstein, M. G., Marcus, B. H., Rakowski, W., Fiore, C., Harlow, L. L., Redding, C. A., & Rosenbloom, D. (1994). Stages of change and decisional balance for 12 problem behaviors. *Health Psychology, 13*(1), 39.
- Pullen, C., & Noble Walker, S. (2002). Midlife and older rural women's adherence to U.S. dietary guidelines across stages of change in healthy eating. *Public Health Nursing, 19*(3), 170-178.
- Puyau, M. R., Adolph, A. L., Vohra, F. A., Zakeri, I., & Butte, N. F. (2004). Prediction of activity energy expenditure using accelerometers in children. *Medicine and Science in Sports and Exercise, 36*(9), 1625-1631.
- Rampal, L., Sherina, M. S., Rampal, S., Wong, Y. D., Chow, P. L., Liew, J. S., & Shum, Y. S. (2007). Prevalence of overweight among secondary school students in Klang district, Selangor. *Malaysian Journal of Nutrition, 13*(1), 1-8.
- Rampersaud, G. C. (2009). Benefits of breakfast for children and adolescents: update and recommendations for practitioners. *American Journal of Lifestyle Medicine, 3*(2), 86-103.
- Ridley, K., Ainsworth, B. E., & Olds, T. S. (2008). Development of a compendium of energy expenditures for youth. *International Journal of Behavioral Nutrition and Physical Activity, 5*, 45.
- Ridley, K., & Olds, T. (2008). Assigning energy costs to activities in children: a review and synthesis. *Medicine+ Science in Sports+ Exercise, 40*(8), 1439.
- Rieder, J., Khan, U. I., Heo, M., Mossavar-Rahmani, Y., Blank, A. E., Strauss, T., Viswanathan, N., & Wylie-Rosett, J. (2013). Evaluation of a community-based weight management program for predominantly severely obese, difficult-to-reach, inner-city minority adolescents. *Childhood Obesity, 9*(4), 292-304.
- Rohde, P., Stice, E., & Marti, C. N. (2015). Development and predictive effects of eating disorder risk factors during adolescence: Implications for prevention efforts. *International Journal of Eating Disorders, 48*(2), 187-198.
- Rouhani, M. H., Mirseifinezhad, M., Omrani, N., Esmailzadeh, A., & Azadbakht, L. (2012). Fast food consumption, quality of diet, and obesity among Isfahanian adolescent girls. *Journal of Obesity, 2012*.
- Sanaeinasab, H., Saffari, M., Nazeri, M., Karimi Zarchi, A., & Cardinal, B. J. (2013). Descriptive analysis of Iranian adolescents' stages of change for physical activity behavior. *Nursing and Health Sciences, 15*(3), 280-285.
- Sanchez, A., Norman, G. J., Sallis, J. F., Calfas, K. J., Rock, C., & Patrick, K. (2008). Patterns and correlates of multiple risk behaviors in overweight women. *Preventive Medicine, 46*(3), 196-202.

- Sardinha, L. B., Santos, R., Vale, S., Silva, A. M., Ferreira, J. P., Raimundo, A. M., Moreira, H., Baptista, F., & Mota, J. (2011). Prevalence of overweight and obesity among Portuguese youth: A study in a representative sample of 10 - 18 - year - old children and adolescents. *International Journal of Pediatric Obesity*, 6(2), 124-128.
- Savva, S. C., Kourides, Y. A., Hadjigeorgiou, C., & Tornaritis, M. J. (2013). Overweight and obesity prevalence and trends in children and adolescents in Cyprus 2000–2010. *Obesity Research & Cinical Practice*.
- Schönbeck, Y., Talma, H., van Dommelen, P., Bakker, B., Buitendijk, S. E., HiraSing, R. A., & van Buuren, S. (2011). Increase in prevalence of overweight in Dutch children and adolescents: a comparison of nationwide growth studies in 1980, 1997 and 2009. *PLoS ONE*, 6(11), e27608.
- Schwarzer, R., & Renner, B. (2000). Social-cognitive predictors of health behavior: action self-efficacy and coping self-efficacy. *Health Psychology*, 19(5), 487.
- Sedgh, G., Finer, L. B., Bankole, A., Eilers, M. A., & Singh, S. (2015). Adolescent pregnancy, birth, and abortion rates across countries: levels and recent trends. *Journal of Adolescent Health*, 56(2), 223-230.
- Si, Q., Yu, K., J. Cardinal, B., Lee, H., Yan, Z., Loprinzi, P. D., Li, F., & Liu, H. (2011). Promoting exercise behavior among Chinese youth with hearing loss: a randomized controlled trial based on the Transtheoretical Model. *Psychological Reports*, 109(6), 896-906.
- Simko, M. D., Cowell, C., & Hreha, M. S. (1989). *Practical Nutrition: a quick reference for the health care practitioner*.
- Song, Y., Park, M. J., Paik, H. Y., & Joung, H. (2010). Secular trends in dietary patterns and obesity-related risk factors in Korean adolescents aged 10-19 years. *International Journal of Obesity*, 34(1), 48-56.
- Soo, K., Wan, A., Abdul, M., & Lee, Y. (2011). Dietary practices among overweight and obese Chinese children in Kota Bharu, Kelantan. *Malaysian Journal of Nutrition*, 17(1), 87-95.
- Srdić, B., Obradović, B., Dimitrić, G., Stokić, E., & Babović, S. S. (2012). Relationship between body mass index and body fat in children—Age and gender differences. *Obesity Research & Cinical Practice*, 6(2), e167-e173.
- Staiano, A., & Katzmarzyk, P. (2012). Ethnic and sex differences in body fat and visceral and subcutaneous adiposity in children and adolescents. *International Journal of Obesity*, 36(10), 1261-1269.
- Stang, J., & Story, M. (2005). *Guidelines for adolescent nutrition services*. Minneapolis: School of Public Health, University of Minnesota.

- Starc, G., & Strel, J. (2011). Tracking excess weight and obesity from childhood to young adulthood: a 12-year prospective cohort study in Slovenia. *Public Health Nutrition*, 14(01), 49-55.
- Stea, T. H., Vik, F. N., Bere, E., Svendsen, M. V., & Oellingrath, I. M. (2014). Meal pattern among Norwegian primary-school children and longitudinal associations between meal skipping and weight status. *Public Health Nutrition*.
- Steele, M. M., Daratha, K. B., Bindler, R. C., & Power, T. G. (2011). The relationship between self-efficacy for behaviors that promote healthy weight and clinical indicators of adiposity in a sample of early adolescents. *Health Education and Behavior*, 38(6), 596-602.
- Steele, M. M., Steele, R. G., & Cushing, C. C. (2012). Weighing the pros and cons in family-based pediatric obesity intervention: parent and child decisional balance as a predictor of child outcomes. *Children's Health Care*, 41(1), 43-55.
- Steinberger, J., & Kelly, A. S. (2014). Obesity, metabolic syndrome and type 2 diabetes. In E. M. Da Cruz, D. Ivy, & J. Jagers (Eds.), *Pediatric and Congenital Cardiology, Cardiac Surgery and Intensive Care* (pp. 499-507). Springer: London
- Su, T. T., Sim, P. Y., Nahar, A. M., Majid, H. A., Murray, L. J., Cantwell, M. M., Al-Sadat, N., & Jalaludin, M. Y. (2014). Association between self-reported physical activity and indicators of body composition in Malaysian adolescents. *Preventive Medicine*, 67, 100-105.
- Tang, K. H., Trang, N. H., Dibley, M. J., Sibbritt, D. W., Binh, P. N., & Hanh, T. (2010). Factors associated with adolescent overweight/obesity in Ho Chi Minh city. *International Journal of Pediatric Obesity*, 5(5), 396-403.
- Tee, E. S., Mohd Ismail, N., Mohd Nasir, A., & Khatijah, I. (1997). Nutrient composition of Malaysian foods (4th ed.). Kuala Lumpur: Institute for Medical Research.
- Teo, P. S., Nurul-Fadhilah, A., Aziz, M. E., Hills, A. P., & Foo, L. H. (2014). Lifestyle practices and obesity in Malaysian adolescents. *International Journal of Environmental Research and Public Health*, 11(6), 5828-5838.
- Teo, Y. B., Sedek, R., & Zalifah, M. K. (2012). Association between snacking patterns, energy and nutrient intakes, and body mass index among school adolescents in Kuala Lumpur. *American Journal of Food and Nutrition*, 2(3), 69-77.
- Thibault, H., Contrand, B., Saubusse, E., Baine, M., & Maurice-Tison, S. (2010). Risk factors for overweight and obesity in French adolescents: physical activity, sedentary behavior and parental characteristics. *Nutrition*, 26(2), 192-200.
- Thompson, M. A., & Gray, J. J. (1995). Development and validation of a new body-image assessment scale. *Journal of Personality Assessment*, 64(2), 258-269.
- Torun, B., Davies, P. S., Livingstone, M. B., Paolisso, M., Sackett, R., & Spurr, G. B. (1996). Energy requirements and dietary energy recommendations for children

and adolescents 1 to 18 years old. *European Journal of Clinical Nutrition*, 50(1), 37-81.

Toschke, A. M., Küchenhoff, H., Koletzko, B., & Kries, R. (2005). Meal frequency and childhood obesity. *Obesity Research*, 13(11), 1932-1938.

Toschke, A. M., Thorsteinsdottir, K. H., Von Kries, R., & Group, G. S. (2009). Meal frequency, breakfast consumption and childhood obesity. *International Journal of Pediatric Obesity*, 4(4), 242-248.

Triches, R. M., & Giugliani, E. R. J. (2005). Obesity, eating habits and nutritional knowledge among school children. *Revista de Saude Publica*, 39(4), 541-547.

Trinh, A., Campbell, M., Ukoumunne, O. C., Gerner, B., & Wake, M. (2013). Physical activity and 3-year BMI change in overweight and obese children. *Pediatrics*, 131(2), 470-477.

Tung, E., Shamarina, S., & Mohd Nasir, M. (2011). Familial and socio-environmental predictors of overweight and obesity among primary school children in Selangor and Kuala Lumpur. *Malaysian Journal of Nutrition*, 17(2), 151-162.

Tzotzas, T., Kapantais, E., Tziomalos, K., Ioannidis, I., Mortoglou, A., Bakatselos, S., Kaklamanou, M., Lanaras, L., & Kaklamanou, D. (2011). Prevalence of overweight and abdominal obesity in Greek children 6-12 years old: Results from the National Epidemiological Survey. *Hippokratia*, 15(1), 48.

Verdejo - García, A., Pérez - Expósito, M., Schmidt - Río - Valle, J., Fernández - Serrano, M. J., Cruz, F., Pérez - García, M., López - Belmonte, G., Martín - Matillas, M., Martín - Lagos, J. A., & Marcos, A. (2010). Selective alterations within executive functions in adolescents with excess weight. *Obesity*, 18(8), 1572-1578.

Vereecken, C., De Henauw, S., Maes, L., Moreno, L., Manios, Y., Phillipp, K., Plada, M., De Bourdeaudhuij, I., & Group, H. S. (2009). Reliability and validity of a healthy diet determinants questionnaire for adolescents. *Public Health Nutrition*, 12(10), 1830-1838.

Wade, T. D., Hansell, N. K., Crosby, R. D., Bryant-Waugh, R., Treasure, J., Nixon, R., Byrne, S., & Martin, N. G. (2013). A study of changes in genetic and environmental influences on weight and shape concern across adolescence. *Journal of Abnormal Psychology*, 122(1), 119.

Wafa, S. W., Talib, R. A., Hamzaid, N. H., McColl, J. H., Rajikan, R., Ng, L. O., Ramli, A. H., & Reilly, J. J. (2011). Randomized controlled trial of a good practice approach to treatment of childhood obesity in Malaysia: Malaysian Childhood Obesity Treatment Trial (MASCOT). *International Journal of Pediatric Obesity*, 6(2), 62-69.

- Wagner, I. V., Sabin, M. A., Pfaffle, R. W., Hiemisch, A., Sergeev, E., Korner, A., & Kiess, W. (2012). Effects of obesity on human sexual development. *Nature Reviews. Endocrinology*, 8(4), 246-254.
- Wallace, L. S., Buckworth, J., Kirby, T. E., & Sherman, W. M. (2000). Characteristics of exercise behavior among college students: application of social cognitive theory to predicting stage of change. *Preventive Medicine*, 31(5), 494-505.
- Webb, M. C., & Beckford, S. E. (2014). Nutritional knowledge and attitudes of adolescent swimmers in Trinidad and Tobago. *Journal of Nutrition and Metabolism*, 2014.
- Weinsier, R. L., Hunter, G. R., Heini, A. F., Goran, M. I., & Sell, S. M. (1998). The etiology of obesity: relative contribution of metabolic factors, diet, and physical activity. *The American Journal of Medicine*, 105(2), 145-150.
- Wells, J. C. (2007). Sexual dimorphism of body composition. *Best Practice & Research Clinical Endocrinology & Metabolism*, 21(3), 415-430.
- Wilson, L. F. (2007). Adolescents' attitudes about obesity and what they want in obesity prevention programs. *The Journal of School Nursing*, 23(4), 229-238.
- Wilson, M. N., Rahman, S. M. M., Lopez, I. A., Kiros, G. E., James, D. C. S., & Dignan, M. B. (2014). Perceived parental support and self-efficacy for avoiding sedentary behaviors and overweight status among school children. *Universal Journal of Public Health*, 2(1), 1-9.
- Wong, C. Y., Zalilah, M. S., Mirlanili, K., & Mohd Nasir, M. T. (2014). Stages of change to increase fruit and vegetable intake and its relationships with fruit and vegetable intake and related psychosocial factors. *Nutrition Research and Practice*, 8(3), 297-303.
- Wong, C. Y., Zalilah, M. S., & Yap, L. D. (2009). Socio-demographic characteristics and nutritional status of individuals by stages of change for dietary fat reduction. *Pakistan Journal of Nutrition*, 8(12), 1821-1829.
- Woon, F. C., Chin, Y. S., & Mohd Nasir, M. T. (2014). Association between behavioural factors and BMI-for-age among early adolescents in Hulu Langat district, Selangor, Malaysia. *Obesity Research & Clinical Practice*.
- World Health Organization [WHO]. (2005). Nutrition in Adolescence - Issues and Challenges for The Health Sector: Issues in Adolescent Health and Development. Geneva: World Health Organization.
- World Health Organization [WHO]. (2000). Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation. WHO Technical Report Series 894. Geneva: World Health Organization.
- World Health Organization [WHO]. (2007). *WHO Child Growth Standards: length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age* Geneva: WHO. Worsley, A. (2002). Nutrition

knowledge and food consumption: can nutrition knowledge change food behaviour? *Asia Pacific Journal of Clinical Nutrition*, 11(s3), S579-S585.

Yansaneh, I. S. (2003). *Construction and use of sample weights (draft)* (Publication No. ESA/STAT/AC.93/5): New York: United Nations Secretariat, Statistics Division.

Yen, C. F., Hsiao, R. C., Yen, J. Y., Ko, C. H., Liu, S. C., Huang, C. F., & Wang, S. Y. (2010). Body weight statuses and their sociodemographic correlates among adolescents in southern Taiwan: results using two sets of cutoff references. *Journal of Adolescent Health*, 46(1), 62-69.

Yusoff, F., Saari, R., Naidu, B. M., Ahmad, N. A., Omar, A., & Aris, T. (2014). Methodology of the national school-based health survey in Malaysia, 2012. *Asia-Pacific Journal of Public Health*, 26(5 suppl), 9S-17S.

Zabinski, M. F., Daly, T., Norman, G. J., Rupp, J. W., Calfas, K. J., Sallis, J. F., & Patrick, K. (2006). Psychosocial correlates of fruit, vegetable, and dietary fat intake among adolescent boys and girls. *Journal of the American Dietetic Association*, 106(6), 814-821.

Zaccagni, L., Masotti, S., Donati, R., Mazzoni, G., & Gualdi-Russo, E. (2014). Body image and weight perceptions in relation to actual measurements by means of a new index and level of physical activity in Italian university students. *Journal of Translational Medicine*, 12, 42.

Zalilah, M., Khor, G., Mirnalini, K., Norimah, A., & Ang, M. (2006). Dietary intake, physical activity and energy expenditure of Malaysian adolescents. *Singapore Medical Journal*, 47(6), 491-498.

Zhang, C. X., Chen, Y. M., Chen, W. Q., Deng, X. Q., & Jiang, Z. Q. (2007). Energy expenditure and energy intake in 10-12 years obese and non-obese Chinese children in a Guangzhou boarding school. *Asia Pacific Journal of Clinical Nutrition*, 17(2), 235-242.

Zhang, C. X., Chen, Y. M., Chen, W. Q., Su, Y. X., Wang, C. L., & Wu, J. N. (2012). Food group intake among adolescents in Guangzhou city compared with the Chinese dietary guidelines. *Asia Pacific Journal of Clinical Nutrition*, 21(3).

Zhang, Y. X., & Wang, S. R. (2012). Secular trends in body mass index and the prevalence of overweight and obesity among children and adolescents in Shandong, China, from 1985 to 2010. *Journal of Public Health*, 34(1), 131-137.

Zhang, Y. X., Zhou, J. Y., Zhao, J. S., & Chu, Z. H. (2013). The role of 1-h physical activity every day in preventing obesity in adolescents in Shandong, China. *European Journal of Pediatrics*, 172(3), 325-330.