

# **UNIVERSITI PUTRA MALAYSIA**

FACTORS CONTRIBUTING TO BONE MINERAL DENSITY AMONG CHINESE WOMEN IN THE KLANG VALLEY, MALAYSIA

LAU LEE TING

FPSK(M) 2016 8



# FACTORS CONTRIBUTING TO BONE MINERAL DENSITY AMONG CHINESE WOMEN IN THE KLANG VALLEY, MALAYSIA



LAU LEE TING

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science

January 2016

# COPYRIGHT

All materials contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purpose from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Degree of Master of Science

### FACTORS CONTRIBUTING TO BONE MINERAL DENSITY AMONG CHINESE WOMEN IN THE KLANG VALLEY, MALAYSIA

#### By

### LAU LEE TING

#### January 2016

Chairperson Faculty

### : Chan Yoke Mun, PhD : Medicine and Health Sciences

Osteoporosis and its related fractures have become epidemic health problem over the years. However, the studies on bone mineral density (BMD) and its related factors were limited in Malaysia. The purpose of this study was to determine the bone health status and examine the contribution of sociodemographic background, family history of fracture, reproductive history, biochemical measures, dietary intake, lifestyle practices as well as anthropometric measurements towards BMDs among healthy Chinese women residing in the Klang Valley. Subjects were participants from a 12-month milk supplementation study conducted from 2012 to 2014. There were 263 women, of which 137 were premenopausal and 126 were postmenopausal women.

A pre-tested questionnaire and three validated questionnaires, namely International Physical Activity Questionnaire (IPAQ), Weekly Sun Exposure Questionnaire and Food Frequency Questionnaire (FFQ), were administered through interview to collect details of all variables, except biochemical measures which were collected through fasting blood and urine samples. Dual-energy X-ray absorptiometry (DXA) scan examined BMD at total body, L1-L4 spine and total hip. All variables that were significant at bivariate analysis were selected into respective models of BMDs in stepwise multiple linear regression (MLR).

The mean age of the subjects was  $50.0 \pm 10.3$  years. Prevalence of osteoporosis was 1% and 4% at total body and lumbar spine BMD in postmenopausal subjects. There were 1% of premenopausal subjects who had L1-L4 spine and total hip BMD below the expected range for age. Postmenopausal subjects presented significant lower mean BMD at all sites (p<0.05). Older age, lower education level and lower monthly household income as well as higher parity, bone turnover rate [Procollagen Type 1 N-Terminal Propeptide (P1NP), osteocalcin (OC) and C-telopeptide of type I collagen crosslinks (CTX)], energy and nutrients intake, physical activity measure, body mass index (BMI), fat mass, measures of central obesity were observed among postmenopausal women (p<0.05). Vitamin D deficiency (<50 nmol/L) was presented in 42.5%, of which 2/3 was contributed by premenopausal subjects. Mean serum 25-

hydroxyvitamin D, physical activity measure, habitual and current calcium intake were reported as  $56 \pm 20$  nmol/L, 1040 MET-minutes/week,  $499 \pm 202$  mg/day and 330 mg/day, respectively.

Among the variables, menopausal status had shown to develop the strongest correlation with BMDs, especially at L1-L4 spine BMD (r= -0.570, p<0.01). Age and biochemical measures, particularly urinary CTX also demonstrated moderate and negative association with BMDs (p<0.01). Besides, weight and lean mass presented higher magnitude than BMI and fat mass, respectively, in positive relation with BMDs. Despite weaker associations, lower BMDs were associated with lower education level, presence of family history of fractures, multiparous, higher physical activity measure, smoking and higher central obesity measures.

Stepwise multiple linear regression analyses on three skeletal sites showed 40.1%, 42.1% and 17.2% variances in the regulation of BMD at total body, L1-L4 spine and total hip, respectively. Significant contributors of BMDs were as following: urinary CTX towards all BMD sites; menopausal status and lean mass towards total body and L1-L4 spine BMD; weight towards total hip BMD.

In conclusion, this study presented the bone heath status and identified risk of low bone mass significantly contributed by high CTX, being postmenopausal, low anthropometric measurements of lean mass and weight among Chinese women throughout middle-aged and senior adulthood. Where low physical activity level, low dietary calcium intake and vitamin D deficiency were prevalent, appropriate health promotion programs should be carefully planned to optimize bone health status among Chinese women in Malaysia.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

### FAKTOR-FAKTOR MENYUMBANG KEPADA KEPADATAN MINERAL TULANG DALAM KALANGAN WANITA CINA DI LEMBAH KLANG, MALAYSIA

Oleh

#### LAU LEE TING

Januari 2016

### Penge<mark>rusi : Chan Yoke Mun</mark>, PhD Fakulti : Perubatan dan Sains Kesihatan

Osteoporosis dan keretakan tulang telah membawa masalah kesihatan secara epidemik selama ini. Walaubagaimanapun, kajian tentang faktor berkaitan dengan kepadatan mineral tulang (KMT) masih terhad di Malaysia. Kajian ini bertujuan untuk menentukan status kesihatan tulang dan menguji sumbangan latar belakang sosiodemografik, sejarah retak tulang keluarga, sejarah reproduktif, ukuran biokimia, pengambilan makanan, amalan gaya hidup dan ukuran anthropometrik terhadap KMT dalam kalangan wanita cina yang sihat di Lembah Klang. Responden kajian ini terdiri daripada 263 wanita yang menyertai kajian suplementasi susu selama 12 bulan bermula tahun 2012 hingga 2014. Antaranya, terdapat 137 wanita pra-menopaus dan 126 wanita pos-menopaus.

Sesi temubual dijalankan berpandukan satu borang soal selidik yang dipra-uji dan tiga borang soal selidik yang disahkan, iaitu borang soal selidik aktiviti fizikal antarabangsa (IPAQ), borang soal selidik pendedahan cahaya matahari mingguan dan kekerapan pengambilan makanan (FFQ) untuk mengumpul semua butiran pembolehubah kajian kecuali ukuran biokimia yang dikumpul melalui sampel darah dan kencing puasa. Imbasan dual-energy X-ray absorptiometry (DXA) dijalankan untuk mengukur KMT pada bahagian keseluruhan tubuh, lumbar dan pangkal pinggul. Semua pembolehubah yang signifikan pada ujian pekali dipilih untuk model KMT dalam analisis regresi linear berganda.

Min umur responden ialah 50.0  $\pm$  10.3 tahun. Seramai 1% dan 4% responden posmenopaus dikategorikan sebagai osteoporosis pada bahagian keseluruhan tubuh dan lumbar manakala 1% responden pra-menopaus mempunyai KMT di bawah jangkaan umur. Kepadatan mineral tulang dalam kalangan responden pos-menopaus adalah lebih rendah (p<0.05). Secara keseluruhannya, responden pos-menopaus adalah lebih berumur, berpendidikan lebih rendah, pendapatan isi rumah lebih rendah, lebih pariti, mempunyai kadar pertukaran tulang [Procollagen Type 1 N-Terminal Propeptide (P1NP), osteocalcin (OC) dan C-telopeptide of type I collagen crosslinks (CTX)] yang lebih tinggi, mempunyai pengambilan tenaga dan nutrien makanan yang lebih tinggi, lebih aktif secara fizikal dan mempunyai ukuran antropometrik [indek jisim badan (IJT), jisim lemak badan, obesiti abdomen] yang lebih tinggi berbanding dengan responden pra-menopaus (p<0.05). Seramai 42.5% responden menghidapi kekurangan vitamin D (<50 nmol/mL) dan kira-kira 2/3 disumbangkan daripada responden pra-menopaus. Min serum 25-hydroxyvitamin D, ukuran aktiviti fizikal, pengambilan kalsium lazim dan semasa ialah 56  $\pm$  20 nmol/L, 1040 MET-minit/minggu, 499  $\pm$  202 mg/hari and 330 mg/hari masing-masing.

Antara pembolehubah-pembolehubah, ujian pekali menunjukkan kolelasi yang paling kuat di antara status menopaus dengan KMT pada bahagian lumbar (r= -0.570, p<0.01). Umur dan ukuran biokimia, terutamanya CTX mempunyai perkaitan negatif yang serderhana dengan KMT pada semua bahagian (p<0.01). Selain itu, berat badan dan jisim otot menunjukkan magnitud korelasi yang lebih tinggi daripada IJT dan jisim lemak dalam perkaitan positif dengan KMT pada semua bahagian. Walaupun berkorelasi lemah, KMT menurun secara signifikan dengan tahap pendidikan, mempunyai sejarah retak tulang keluarga, lebih pariti, lebih aktif secara fizikal, merokok dan lebih tinggi ukuran obesiti abdomen.

Analisis regresi linear berganda menjelaskan 40.1%, 42.1% dan 17.2% varians KMT pada bahagian keseluruhan tubuh, lumbar dan pangkal pinggul masing-masing. Penyumbang KMT yang signifikan adalah seperti berikut: CTX terhadap semua bahagian KMT; status menopaus dan jisim otot terhadap KMT pada bahagian keseluruhan tubuh dan lumbar; berat badan terhadap KMT pada bahagian pangkal pinggul.

Kesimpulannya, kajian ini menunjukkan status kesihatan tulang dan mengenalpasti risiko KMT yang rendah disumbang secara signifikan oleh CTX yang tinggi, posmenopaus, ukuran anthropometrik yang rendah (jisim otot dan berat badan) dalam kalangan dewasa dan warga emas. Program kesihatan yang mengutamakan pengamalan gaya hidup yang aktif, pengambilan cukup kalsium dari makanan dan peningkatan status vitamin D hendaklah dirancangkan untuk mengoptimumkan status kesihatan tulang dalam kalangan wanita cina di Malaysia.

#### ACKNOWLEDGEMENTS

First and foremost, I would like to dedicate my gratitude to my supervisor, Assoc. Prof. Dr. Chan Yoke Mun for her guidance and continuous support throughout the research execution period and thesis writing. Her enthusiasm and patience moved me in many times throughout the period pursuing studies. Besides, my gratitude also offers to co-supervisors, Dr. Chin Yit Siew and Prof Lye Munn Sann for their advice and constructive comments.

My sincere thanks go to the management and staffs of ANOC Neuroscience and Orthopaedic Centre for their cooperation in ensuring proper execution of procedures and daily research works during data collection period: Dr Chin Chee Howe for offering his medical advice, imaging technicians for ensuring DXA procedure, nurses for administrative issues. I would also like to thank experienced phlembotomists from Quantum Diagnostics Sdn Bhd for their assistance in blood sampling and sample delivery. Moreover, thanks to the lab assistants of nutrition lab for her assistance in obtaining necessary equipment for anthropometric measurements.

I am thankful to meet the respondents from various places in Klang Valley who offered much cooperation during assessment days. I thank my fellow assistants who helped out during the data collection period. In daily work, I have been blessed with a caring colleague, Lau Chin Chin, whom I shared ups and downs with. Last but not least, I thank my family members for their unfailing love and support. This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as following:

# Chan Yoke Mun, PhD

Associate Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Chairman)

# Lye Munn Sann, MBBS (Mal), MPH (Tulane), DrPH (Tulane)

Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Member)

# Chin Yit Siew, PhD

Senior Lecturer Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Member)

# **BUJANG BIN KIM HUAT, PhD** Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date:

### Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property form the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature:

Date:

Name and Matric No: Lau Lee Ting, GS35462

## **Declaration by Members of Supervisory Committee**

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in Rule 41 in Rules 2003 (Revision 2012-2013) were adhered to

Signature: Name of Chairman of Supervisory Committee:

Assoc. Prof. Dr. Chan Yoke Mun

Signature: Name of Member of Supervisory Committee:

Professor Dato' Dr. Lye Munn Sann

Signature: Name of Member of Supervisory Committee:

Dr. Chin Yit Siew

# TABLE OF CONTENTS

	Page
ABSTRACT <i>ABSTRAK</i> ACKNOWLEDGEMENTS APPROVAL DECLARATION LIST OF TABLES LIST OF FIGURES LIST OF APPENDICES LIST OF ABBREVIATIONS GLOSSARY OF TERMS	i iii v vi viii xiii xivi xv xv xvi xviii
CHAPTER	
1       INTRODUCTION         1.1       Introduction         1.2       Problem statement         1.3       Significance of study         1.4       Objective of study         1.4.1       General objective         1.4.2       Specific objective         1.5       Hypotheses         1.6       Conceptual framework	1 2 4 5 5 5 5 6
<ul> <li>2 LITERATURE REVIEW</li> <li>2.1 Bone structure and function</li> <li>2.2 Bone remodeling</li> <li>2.3 Peak bone mass</li> <li>2.4 Loss of bone mass</li> <li>2.5 Osteoporosis</li> <li>2.6 Quantitative assessment of bone mass</li> <li>2.7 Determinants of bone health</li> <li>2.7.1 Genetic factors</li> <li>2.7.2 Race and ethnicity</li> <li>2.7.3 Age</li> <li>2.7.4 Socioeconomic status</li> <li>2.7.5 Reproductive factors</li> <li>2.7.6 Bone turnover markers</li> <li>2.7.7 Dietary intake</li> <li>2.7.8 Physical activity</li> <li>2.7.9 Smoking</li> <li>2.7.10 Alcohol intake</li> </ul>	$ \begin{array}{c} 10\\ 10\\ 10\\ 12\\ 13\\ 14\\ 15\\ 17\\ 17\\ 19\\ 19\\ 20\\ 20\\ 20\\ 23\\ 24\\ 29\\ 30\\ 31\\ \end{array} $
<ul><li>2.7.11 Sun exposure</li><li>2.7.12 Body weight, body mass index and body composition</li></ul>	31 32

3	MET	HODOLOGY	36
	3.1	Study design and data source	36
	3.2	Study population	36
		3.2.1 Inclusion criteria	36
		3.2.2 Exclusion criteria	36
	3.3	Sample size estimation	37
	3.4	Sampling method	38
	3.5	Study location	39
	3.6	Subject recruitment	40
	3.7	Ethical clearance	41
	3.8	Data collection	41
	510	3.8.1 Questionnaire	41
		3.8.2 Anthropometric measurements	42
		3 8 3 Biochemical data	45
		3.8.4 Bone mineral density measurements	45
		3.8.5 Body composition measurements	47
		3.8.6 Dietary intake assessment	47
	39	Data analysis and interpretation	51
	5.7	Dud didiysis and interpretation	01
4	RESI	ILTS	53
-	4.1	Background characteristics	53
		4.1.1 Sociodemographic background	53
		4.1.2 Family history of fractures	54
		4.1.3 Reproductive history	54
		4.1.4 Biochemical measures	55
		4.1.5 Dietary intake	56
		4.1.6 Lifestyle practices	60
		4.1.7 Anthropometric measurements	61
		4.1.8 Bone mineral density measurements	62
	4.2	Comparison between premenopausal and postmenopausal subjects	64
	4.3	Correlation between bone health related variables and BMD	67
		121 Completion between here health related variables with	67
		BMD in premenopausal subjects	07
		4.3.2 Correlation between bone health related variables with BMD in postmenopausal subjects	68
		4.3.3 Correlation between bone health related variables with	70
	4.3	Multiple linear regression	71
5	DISC	CUSSION	74
	5.1	Background characteristics	74
		5.1.1 Sociodemographic background	74
		5.1.2 Family history of fractures	74
		5.1.3 Reproductive history	75
		5.1.4 Biochemical measures	75
		5.1.5 Dietary intake	76
		5.1.6 Lifestyle practices	77
		5.1.7 Anthropometric measurements	77
	5.2	Bone health status of subjects	78

5.3 5.4	<ul> <li>Associations between variables and BMD</li> <li>Regression model</li> </ul>	79 82
6 C(	DNCLUSION AND RECOMMENDATIONS	84
6.1	Conclusion	84
6.2	2 Recommendations	85
6.3	3 Limitations	85
BIBLIC	OGRAPHY	87
APPEN	DICES	115
BIODA	TA OF STUDENT	163



 $\bigcirc$ 

# LIST OF TABLES

Table		Page
3.1	MET score categorization	42
3.2	Classification of BMI	43
3.3	Intrameasurer TEM and acceptable TEM	45
3.4	Intermeasurer TEM and acceptable TEM	45
3.5	T-score and interpretation	46
3.6	Z-score and interpretation	47
4.1	Demographic and socioeconomic background of subjects	53
4.2	Distribution of subjects according to family history of fractures	54
4.3	Reproductive history of subjects	55
4.4	Distribution of subjects according to biochemical measures	56
4.5	Mean or median energy and nutrient intakes of subjects	57
4.6	Distribution of subjects according to dietary adequacy	58
4.7	Distribution of subjects according to acceptable reporters and misreporters	
	on energy intake	60
4.8	Distribution of subjects according to lifestyle practices	60
4.9	Distribution of subjects according to anthropometric measurements	61
4.10	Bone mineral density measurements, T-score and Z-score among subjects	63
4.11	Comparison between bone health related variables and bone health status among premenopausal and postmenopausal subjects	65
4.12	Correlation between bone health related variables with total body BMD, lumbar spine BMD and total hip BMD in premenopausal subjects	68
4.13	Correlation between bone health related variables with total body BMD, lumbar spine BMD and total hip BMD in postmenopausal subjects	69
4.14	Correlation between bone health related variables with total body BMD, lumbar spine BMD and total hip BMD in all subjects	71
4.15	Stepwise multiple linear regression of variables and BMD measurements	73

# LIST OF FIGURES

Figure		Page
1.1	Conceptual framework of the study	9
2.1	Bone remodeling process	12
2.2	Distribution of BMD in women at different ages and the prevalence o	f
	osteoporosis	16
3.1	Klang Valley map and population size	39
3.2	Process of subject recruitment	40
4.1	Distribution of premenopausal subjects according to Z-score classification	63
4.2	Distribution of premenopausal subjects according to T-score classification	64

C

# LIST OF APPENDICES

# Appendix

6

А	Medical conditions for patient exclusion and prohibited	
	concomitant medications during the study	115
В	Results from sample size calculations	116
С	Advertising flyers (English & Mandarin)	117
D	Screening form (premenopausal and postmenopausal)	119
Е	Location and total screened number list	125
F	Study protocol (premenopausal and postmenopausal)	126
G	Informed consent form (premenopausal and postmenopausal)	136
Н	Ethical approval letter	138
Ι	Questionnaire	139
J	International Physical Activity Questionnaire (IPAQ)	147
Κ	Weekly Sun Exposure Questionnaire	149
L	Three days food record form	150
Μ	Food Frequency Questionnaire	156
Ν	Recommended Nutrient Intakes for Malaysia	161

# LIST OF ABBREVIATIONS

AFM	Appendicular fat mass
AOI	Android-to-gynoid ratio
ASEAN	Association of Southeast Asian Nations
BMD	Bone mineral density
BMI	Body mass index
BTM	Bone turnover markers
CSDH	Commission on Social Determinants of Health
СТ	Computed tomography
CTX	C-telopeptide of type I collagen crosslinks
DBP	Vitamin D binding protein
DOSM	Department of Statistics Malaysia
DXA	Dual-energy X-ray Absorptiometry
eGFR	Estimated glomerular filtration rate
FFQ	Food Frequency Questionnaire
GDP	Gross Domestic Product
HRQoL	Health-related quality of life
IGF	Insulin-like growth factor
IGF-1	Insulin like growth factor I
IOF	International Osteoporosis Foundation
IOM	Institute of Medicine
IPAQ	International Physical Activity Questionnaire
IPH	Institute for Public Health
ISAK	International Society for the Advancement of Kinanthropometry
ISCD	International Society for Clinical Densitometry
KNHANES	Korean National Health and Nutrition Examination Survey
MANS	National Malaysian Adult Nutrition Survey
MET	Multiples of the resting metabolic rate
МОН	Ministry of Health Malaysia
MRI	Magnetic resonance imaging
NCCFN	National Coordinating Committee for Food and Nutrition
	Malaysia
NHANES III	The Third National Health and Nutrition Examination Survey
NHMS III	The Third National Health and Morbidity Survey
NHWS	National Health and Wellness Surveys
NIH	National Institute of Health
NOF	National Osteoporosis Foundation
PBM	Peak bone mass
PLI	Poverty Line Income
PPAR	Peroxisome proliferator activated receptor
PTH	Parathyroid hormone
P1NP	Procollagen Type 1 N-Terminal Propeptide
QALYs	Quality-adjusted life years
RNI	Recommended Nutrient Intakes
SD	Standard deviation
SES	Socioeconomic status
SPSS	Statistical Package for Social Sciences
SWAN	Study of Women's Health Across the Nation

TEM	Technical Error of Measurement
UKMMC	Universiti Kebangsaan Malaysia Medical Centre
WHO	World Health Organization
WHO/IOTF/IASO	World Health Organization, the International Association for the
	Study of Obesity and the International Obesity Task Force
WHR	Waist-to-hip ratio



#### **GLOSSARY OF TERMS**

Bone mineral densityBone mineral density is a globally recognized surrogate<br/>measurement of bone strength that could be most accurately<br/>measured by Dual-energy Absorptiometry. It is expressed in<br/>absolute terms of grams of mineral per square centimetre<br/>scanned (g/cm²) and its relationship to two norms, namely<br/>Z-score and T-score (National Osteoporosis Foudation,<br/>2010).Dual-energyX-rayDual-energyX-rayDual-energyX-ray

Absorptiometry (DXA) developed technique that is thoroughly validated biologically to assess bone mass, which is regarded as "gold standard" for the assessment of osteoporosis and for the evaluation of therapies (Blake & Fogelman, 2009; World Health Organization, 2003b).

Osteoporosis is a bone disease characterized by porous and brittle bone, which increase the susceptibility of the individual towards fractures (NIH Consensus Development Panel on Osteoporosis Prevention and Therapy, 2001).

Fracture in this thesis are referring to osteoporotic fracture that is contributed from low energy traumas which may occur during a fall from a standing height, or less, or traumas that would not cause fracture in healthy individual (Melton et al., 1997).

Osteoporosis

Fracture

### **CHAPTER 1**

### INTRODUCTION

# 1.1 Introduction

Bone is essential for growth, mobility, protection, storage of mineral and support to human body (Boskey & Coleman, 2010). Since last century, bone health had received much concern from the healthcare professions and public. National bone health related bodies and organizations such as International Osteoporosis Foundation (IOF), National Osteoporosis Foundation (NOF), Malaysian Osteoporosis Society and Malaysian Orthopaedic Association were established with mission to serve as connection platform to resources and information on bone health. Of which, many were dedicated to create awareness, spread the knowledge and increase understanding among health professionals, patients, populations at risk of bone related diseases and general public. These bodies were passionate about communicating disease prevention, early detection and appropriate treatment of bone related diseases including osteoporosis.

Osteoporosis is a bone disease characterized by porous and brittle bone, which increase the susceptibility of fractures. In the past, osteoporosis was perceived as an inevitable condition along with ageing for an individual lost bone mass as they aged (Holroyd, Cooper, & Dennison, 2008). Susceptibility to fracture was described as the most devastating outcome of osteoporosis. Thus, the incidence of osteoporosis is best measured by incidence of fractures (Handa, Kalla, & Maalouf, 2008). Osteoporosis causes more than 8.9 million fractures every year, of which majority of these incidences were reported in Europe, Western Pacific and Southern Asia. These regions collectively contributed to 96% of all fractures worldwide. Astonishingly, Southeast Asia accounted for 15.3% of the statistic (Johnell & Kanis, 2006).

Osteoporosis is a relevant socioeconomic burden at individual, family, health service and societal level. It is a major public health concern that may bring major implications into future (Guillemin et al., 2013; Harvey, Matthews, Collins, & Cooper, 2013; Qu et al., 2014). In Europe, osteoporotic fractures were estimated to cost  $\in$ 37.4 billion in year 2010. The first year post fracture, subsequent year of long term fracture care, pharmacological costs accounted for 66%, 29% and 6% of the total costs, respectively (Hernlund et al., 2013). In Malaysia, the direct hospitalization cost for hip fractures was estimated at USD6 million in year 1997 (IOF, 2009). Such estimation however did not take into account the costs incurred in rehabilitation and long term nursing care. Quality-adjusted life years (QALYs) lost associated with osteoporosis incorporate both the health-related quality of life (HRQoL) and life years lost due to fracture (Hernlund et al., 2013). In European Union, there were approximately  $\in$ 1.2 million QALYs lost due to osteoporosis in 2010 with women experienced most of the QALYs lost as compare to men. Fractures, especially hip and vertebral fractures that associated with increased mortality, were estimated to cause 2/3 life years lost. When the cost and QALYs lost were combined and valued at 2 times of Gross Domestic Product (GDP), the cost of osteoporosis in European Union was amounted to  $\notin$ 98 billion in 2010 (Hernlund et al., 2013).

Over the past three decades, the incidence of hip fracture incidence has raised two- to three-fold in most Asian countries (IOF, 2009). Recent finding also suggested that 30% of the hip fractures occurred worldwide aroused from Asian population (Cooper et al., 2011). With rapid urbanization happening parallel to the rising fracture rate, it was suggested that urbanization could be the main contributors through decreased physical activity, increased in prevalence of vitamin D deficiency, calcium intake inadequacy and other lifestyle factors (Alves, Castiglione, Oliveira, Sousa, & Pina, 2014; Ballane et al., 2014; Lau et al., 2001). In the Malaysian context, Chinese presented the highest rates of hip fractures compared to Malays and Indians, which constituted 63% of all cases. Race and gender-specific incidence data illustrated that Chinese females reported the highest incidences of hip fractures (220 per 100,000). Female were twice more vulnerable to fractures compared to males (Lee & Khir, 2007), which was in concordance with the findings worldwide (Kanis et al., 2012). The plausible explanation might be due to smaller bone size, weaker bone structure and strength with accelerated loss during advancing age in female comparing to males (Havill, Mahaney, Binkley, & Specker, 2007; Yates, Karasik, Beck, Cupples, & Kiel, 2007).

### 1.2 Problem statement

Despite the magnitude of fracture were widely reported, the epidemiology of osteoporosis was still underrated, underdiagnosed and undertreated (IOF, 2009) as it is a "silent disease" with symptomless nature during disease progression (Barling, 2013; Iacono, 2007). Advance technologies are invented nowadays to assist in the assessment of bone health, but many individuals who have osteoporosis are still unaware of their condition until a fracture occurs [Iacono, 2007; Ministry of Health Malaysia (MOH), 2012; NOF, 2010].

Low bone mineral density is multifactorial. Given that osteoporotic fracture can be predicted from low bone mineral density (BMD) (Kanis, 2002), some available evidence revealed systematically lower BMD being observed among Asian comparing to the White counterparts (Mehta et al., 2004; Roy et al., 2005). Several physical, nutritional and lifestyle aspects were proposed to be responsible for such variations. With smaller body size than the European, Asian was found to have smaller bone size and different rate of bone accrual therefore resulted in lower BMD (Roy et al., 2005). Low calcium intake and greater prevalence of vitamin D deficiency in this region was also been proposed to contribute to the phenomena (Handa et al., 2008; Harinarayan, 2005; Malhotra & Mithal, 2008). Surprisingly, tropical countries also experienced deficiency in vitamin D and the prevalence rate could be as high to 85% as that was observed in Western populations (Chee et al., 2010; Hawkins, 2013; Nguyen et al., 2012; Wat, Leung, Tam, & Kung, 2007).

On the other hand, the pandemic trend in the rising of overweight and obesity over the past three decades predicted a continuous rise in the developing world in near future (Ng et al., 2014). Available evidence showed there were large gap existed between actual and ideal participation in the lifestyle behaviours, including eating a healthy diet and undertaking physical activity. Indeed, the prevalence of healthy lifestyle behaviours was low globally, with lower levels in the lower-income countries as compare to high-income countries. Less than half of the world population has good diet quality while high levels of physical activity related to recreational or leisure activity were more prevalent in high-income countries. The observed lifestyle practices deposed the population especially whom resided in developing countries of middle-income status to overweight and obesity (Teo et al., 2013).

The pandemic issue on overweight and obesity cultivated the interest of many researchers on the relationship between anthropometric measures and BMD in recent years (Fu et al., 2011; Gonnelli, Caffarelli, & Nuti, 2014; Montazerifar et al., 2014; Shao et al., 2014). Till date, there are many investigations from epidemiological, clinical and basic research point of view performed actively, demonstrating that osteoporosis and obesity is no longer opposing in their standings but may present connection (Gonnelli et al., 2014; Hsu et al., 2006; Wang et al., 2013). Malaysia, being one of the nations that has been reported with rapid increase in overweight and obesity rate and rated as having the highest obesity rate among Association of Southeast Asian Nations (ASEAN), has brought to greater concern in healthcare aspect [Dans et al., 2011; World Health Organization (WHO), 2011]. The Third National Health and Morbidity Survey (NHMS III) reported a prevalence of 29.1% overweight and 14.0% obesity among Malaysian which was further increase to 33.3% preobese and 27.2% obese in 2011 [Institute for Public Health (IPH), 2011b]. A similar increasing trend of abdominal obesity was also documented among Malaysian (MOH, 2006). Such unhealthy norm among Malaysian could bring deleterious yet undesired impact in the future.

Acknowledging the magnitude of the bone health related risk factors, related studies in examining the associations between many of the life aspects and BMD were however limited in the Malaysian context with majority of such studies were confined to bivariate analysis where possible interaction between multiple factors are not able to be ascertained (Chan et al., 2014; Chee et al., 2010; Lim et al., 2005; Yee, Zaitun, Chan, & Norhaizan, 2013). The limited studies and national reports called up the needs to initiate study in exploring the contribution of bone health related risk factors towards BMDs using multifactorial analysis approach.

Realizing the epidemic and pandemic trend that would most likely to affect bone health deleteriously, this study proposed the following research questions:

i. What are the sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices, anthropometric measurements and bone health status among premenopausal and postmenopausal Chinese women residing in the Klang Valley?

- ii. Is there any significant difference on sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices, anthropometric measurements and BMD measurements between premenopausal and postmenopausal Chinese women residing in the Klang Valley?
- iii. What is the relationship between sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices, anthropometric measurements and BMDs among premenopausal and postmenopausal Chinese women residing in the Klang Valley?
- iv. What is the contribution of sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices, and anthropometric measurements towards BMDs?

## 1.3 Significance of study

Chinese female, being the most vulnerable group towards low bone mass and hip fracture, has been identified as the study population of this study. Since osteoporosis and its related fractures mostly affected postmenopausal individuals, many studies have been focused on them. Similarly, this study also intends to investigate the bone status and the relevant determinants of BMD among postmenopausal Chinese women from wider perspectives than the previous local studies. Additionally, this study commits into determination of bone health status and the mentioned measures among premenopausal Chinese women, of which data is very limited in local sources. Nonetheless, the determination of bone health status in this study is through the use of Dual-energy X-ray Absorptiometry (DXA), which is the gold standard for BMD measurements.

The present knowledge on bone health and its contributing factors was mostly learnt from abroad research. While research study examining bone health from the above mentioned aspects is scarce in Malaysia, this study may contribute to the pool of local data resources. Conversely, it is important in the study of a disease to take into considerations of cultural, religious, dietary and geographical differences. A broad, racial summarization data would not be more valuable than an ethnic-specific data in bone health studies since variations were reported among Asian ethnic groups (Handa et al., 2008; Lee & Khir, 2007).

With more local data available, the foundation of strategy development in osteoporosis prevention and treatment strategy can be more solid. The findings shall assist healthcare authorities (MOH and non-governmental organizations) in forming strategies targeted to the population to maximize bone formation within the modifiable period, to maintain and to reduce bone loss rate after attainment of peak bone mass. This may result in reducing burden of osteoporosis and fractures while maintaining and improving the quality of life of vulnerable populations and their family members. Lastly, this study will contribute to the existing body of knowledge academically and can be used as future research reference. It can be served as baseline references for future intervention study as well.

### 1.4 Objective of study

## 1.4.1 General objective

To determine factors contributing to bone mineral density among Chinese women in the Klang Valley, Malaysia.

### 1.4.2 Specific objective

- i. To determine sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices, anthropometric measurements and bone health status among premenopausal and postmenopausal Chinese women in the Klang Valley.
- ii. To compare sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices, anthropometric measurements and BMD measurements between premenopausal and postmenopausal Chinese women.
- iii. To determine the relationship between sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices, anthropometric measurements and BMDs among premenopausal and postmenopausal Chinese women.
- iv. To examine the contribution of sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices and anthropometric measurements towards BMDs.

#### 1.5 Hypotheses

- i. H<sub>0</sub>: There is no significant difference in sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices, anthropometric measurements and BMD measurements between premenopausal and postmenopausal Chinese women.
- ii. H<sub>0</sub>: There is no significant relationship between sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices, anthropometric measurements and BMDs among premenopausal and postmenopausal Chinese women.
- iii. H<sub>0</sub>: There is no significant contribution of sociodemographic background, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices and anthropometric measurements towards BMDs.

#### 1.6 Conceptual framework

Bone mineral density is an important determinant of bone strength (NOF, 2010). Bone mineral density measurement defines bone health status as normal bone density, low bone mass (named osteopenia) and osteoporosis according to T-score classification (WHO, 1994). Accumulated BMD before peak bone mass (PBM) achieved and loss of bone mass during advancing age determine the BMD in later life (Cashman, 2007b). Evidence has demonstrated relationships between many factors to play as the determinants of BMD. Of which, the factors are categorized into several categories: sociodemographic factors, family history, reproductive history, biochemical measures, dietary intake, lifestyle practices and anthropometric measurements.

Ageing is associated with impaired bone formation and increased bone resorption. Bone mineral density (BMD) reduction in elderly is secondary to physiological and lifestyle changes that causes impaired production of bone stimulatory agents such as insulin-like growth factor (IGF), increased parathyroid hormone secretion and increased muscle loss (Demontiero, Vidal, & Duque, 2012; Lips, 2001). Socioeconomic status is also found to be one of the risk factors associated with poor bone health along with the emerging evidence on socioeconomic gradient relating to all causes of morbidity and mortality (Begg, Vos, Barker, Stanley, & Lopez, 2008; Turrell, Stanley, Looper, & Oldenburg, 2006). Most of the findings suggested a protective role of greater educational attainment and higher income against low BMD and osteoporotic fracture (Peel, McClure, & Hendrikz, 2007; Wang & Dixon, 2006). While genetic factors contributed more than 60% of the total variance in PBM attainment (Rizzoli, Bianchi, Garabédian, McKay, & Moreno, 2010), assessment on family history, especially parental history of fracture provided insight to the relationship between genetic and BMD for the studied population (Kanis, Johansson, et al., 2004).

Changes in reproductive system over lifespan also alter bone metabolism. Estrogen exposure at menarche positively influenced BMD during puberty years (Yilmaz et al., 2005) and across premenopausal years (Sioka et al., 2010). Estrogen prevents substantial bone loss through inhibiting formation of bone-resorbing cell (namely osteoclast) and stimulating cell death on osteoclast (Clarke & Khosla, 2010). Thus, estrogen enables bone mass accumulation during young age and estrogen deficiency leads to accelerated bone loss after menopause (Khosla et al., 2005; Yilmaz et al., 2005). Age of menarche and number of years since menopause as measures of estrogen exposure are therefore played determining roles towards BMD. Besides, pregnancy induced changes in bone metabolism (Oliveri, Parisi, Zeni, & Mautalen, 2004). Parity was shown to influence BMD both positively (Allali et al., 2007; Streeten et al., 2005) and negatively (Ozdemir, Demirbag, & Rodoplu, 2005; Shin et al., 2010) or produced no significant effect (Lenora, Lekamwasam, & Karlsson, 2009; Sioka, Bougias, Papadopoulos, & Fotopoulos, 2007). Lactation induced approximately 5% maternal bone resorption in six months. However, the bone mass was expected to re-establish and restore after weaning (Oliveri et al., 2004). Duration of lactation also showed either negative association or no association with BMD in studies (Dursun, Akin, Dursun, Sade, & Korkusuz, 2006; Kojima, Douchi, Kosha, & Nagata, 2002).

Parathyroid hormone (PTH) and serum vitamin D are important in the maintenance of normal homeostatic regulation of bone mineral content, particularly through the regulation of serum mineral levels (Moe, 2008). Impairment in PTH regulatory mechanism caused impairment in bone metabolism while serum vitamin D positively influenced BMD (Bischoff-Ferrari, Dietrich, Orav, & Dawson-Hughes, 2004; Moe, 2008). Biochemical measures of bone turnover serve as surrogate markers in the assessment of bone metabolism and management of bone diseases (Niimi et al., 2014). The commonly used bone turnover markers are Procollagen Type 1 N-Terminal Propeptide (P1NP), serum osteocalcin and bone resorption markers, namely C-telopeptide of type I collagen crosslinks (CTX) (Penny, Godber, & Lawson, 2007). These markers demonstrated negative associations with BMD at lumbar spine, femoral neck and total hip (Hu et al., 2013).

Dietary factors such as energy, protein, calcium, phosphorus and vitamin D consumed as part of diet potentially influence BMD. Energy and the mentioned nutrients have been widely studied and have shown to play remarkable roles in preserving bone health (Caporaso, Frisch, & Sumida, 2011; Chevalley, Bonjour, Ferrari, Hans, & Rizzoli, 2005; Heaney, 2007; Holm, Dan, Wilbur, Li, & Walker, 2010; Ihle & Loucks, 2004; Lamberg-Allardt, Karp, & Kemi, 2010; Takeda, Yamamoto, Yamanaka-Okumura, & Taketani, 2012; Thorpe & Evans, 2011). While dietary intake of vitamin D through exogenous food sources ingestion played least role as compare to endogenous synthesis of vitamin D, serum vitamin D is well accepted as a biomarker for vitamin D status which accounted for both endogenous and exogenous sources (Adams & Hewison, 2010; Goff, Cavalier, Souberbielle, González-Antuña, & Delvin, 2015).

Lifestyle practices play significant role in influencing bone health. Regular engagement in physical activity exerted mechanical loading to bone and promoted PBM attainment in early life (DiVasta & Gordon, 2013), maintained bone mass in adulthood (Babatunde, Forsyth, & Gidlow, 2012) and reduced bone loss rate in elderly (Pines & Berry, 2007). Besides, literature suggested a positive relationship between sun exposure and BMD through synthesis of serum 25(OH)D and thereby reduced bone loss (Macdonald et al., 2008; Zhen, Liu, Guan, Zhao, & Tang, 2015). On the other hand, long term cigarette smoking substantially increased risk of osteoporotic fracture to nearly two-fold in hip fracture compare with non-smoker (Kanis et al., 2005). Contrary to harmful effect of smoking, moderate alcohol consumption of less than 150 g or 12 drinks weekly was not associated significant harmful effect on BMD (Waugh et al., 2009). However, excessive and prolonged alcohol intake caused compromised BMD and contributed to increased fracture risk (Berg et al., 2009).

Majority of the literature suggested a positive relationship between body weight, BMI and BMD. Low BMI was associated with increased risk of fracture whereas higher BMI was protective against osteoporotic fracture (Laet et al., 2005). It was generally accepted that larger BMI imposed greater mechanical loading that bone mass increased to accommodate the load. Recent studies, however, challenge the existing findings and demonstrated negative association between body fat and BMD when the mechanical loading effect was controlled for (Zhao et al., 2007). The relative effect of the two major components of body weight, lean mass and fat mass, is still under dispute. Some

studies proposed stronger effect from fat mass towards BMD while others suggested lean mass being more closely related to BMD. Besides, studies also indicated central obesity imposed detrimental effect towards BMD (Fu et al., 2011; Shao et al., 2014) due to increased production of inflammatory cytokines in obesity (Goodpaster et al., 2003). The conceptual framework of this study is illustrated in Figure 1.1.







### BIBLIOGRAPHY

- Adami, S., Bertoldo, F., Braga, V., Fracassi, E., Gatti, D., Gandolini, G., ... Battista Rini, G. (2009). 25-hydroxy vitamin D levels in healthy premenopausal women: association with bone turnover markers and bone mineral density. *Bone*, 45(3), 423–426.
- Adams, J. S., & Hewison, M. (2010). Update in vitamin D. Journal of Clinical Endocrinology and Metabolism, 95(2), 471–478.
- Aday, L.A & Cornelius, L. J. (2006). Designing and conducting health surveys: A Comprehensive Guide.
- Ahlborg, H. G., Johnell, O., Nilsson, B. E., Jeppsson, S., Rannevik, G., & Karlsson, M. K. (2001). Bone loss in relation to menopause: a prospective study during 16 years. *Bone*, 28(3), 327–331.
- Ahn, J. H., Lim, S. W., Song, B. S., Seo, J., Lee, J. A., Kim, D. H., & Lim, J. S. (2013). Age at menarche in the Korean female: secular trends and relationship to adulthood body mass index. *Annals of Pediatric Endocrinology & Metabolism*, 18(2), 60–64. doi:10.6065/apem.2013.18.2.60
- Ainsworth, B. E., Haskell, W. L., Whitt, M. C., Irwin, M. L., Swartz, A. M., Strath, S. J., ... Lean, A. S. (2000). Compendium of physical activities: an update of activity codes and MET intensities. *Medicine and Science in Sports and Exercise*, 32(9), 498–504.
- Akkus, Z., Camdeviren, H., Celik, F., Gur, A., & Nas, K. (2005). Determination of osteoporosis risk factors using a multiple logistic regression model in postmenopausal Turkish women. Saudi Medical Journal, 26(9), 1351–1359.
- Akune, T., Ohba, S., Kamekura, S., Yamaguchi, M., Chung, U., Kubota, N., ... Kawaguchi, H. (2004). PPAR  $\gamma$  insufficiency enhances osteogenesis through osteoblast formation from bone marrow progenitors. *Journal of Clinical Investigation*, 113(6), 846–55.
- Alam, I., Padgett, L. R., Ichikawa, S., Alkhouli, M., Koller, D. L., Lai, D., ... Econs, M. J. (2014). SIBLING family genes and bone mineral density: association and allele-specific expression in humans. *Bone*, 64, 166–172.
- Al-Azzawi, F., & Palacios, S. (2009). Hormonal changes during menopause. *Maturitas*, 63(2), 135–137.
- Albanese, C. V, Terlizzi, F. D., & Passariello, R. (2011). Quantitative ultrasound of the phalanges and DXA of the lumbar spine and proximal femur in evaluating the risk of osteoporotic vertebral fracture in postmenopausal women. *La Radiologia Medica*, *116*(1), 92–101.
- Allali, F., Aichaoui, S. E., Khazani, H., Benyahia, B., Saoud, B., Kabbaj, S. E., ... Hajjaj-Hassouni, N. (2009). High prevalence of hypovitaminosis D in Morocco: relationship to lifestyle, physical performance, bone markers, and bone mineral density. *Seminars in Arthritis and Rheumatism*, 38(6), 444–451.

- Allali, F., Maaroufi, H., Aichaoui, S. E., Khazani, H., Saoud, B., Benyahya, B., ... Hajjaj-Hassouni, N. (2007). Influence of parity on bone mineral density and peripheral fracture risk in Moroccan postmenopausal women. *Maturitas*, 57(4), 392–398.
- Alves, S. M., Castiglione, D., Oliveira, C. M., Sousa, B. D., & Pina, M. F. (2014). Age-period-cohort effects in the incidence of hip fractures: political and economic events are coincident with changes in risk. *Osteoporosis International*, 25(2), 711–720.
- Amin, S., Riggs, B. L., Melton, L. J., Achenbach, S. J., Atkinson, E. J., & Khosla, S. (2007). High serum IGFBP-2 is predictive of increased bone turnover in aging men and women. *Journal of Bone and Mineral Research*, 22(6), 799–807.
- Ardawi, M.-S. M., Qari, M. H., Rouzi, A. A., Maimani, A. A., & Raddadi, R. M. (2011). Vitamin D status in relation to obesity, bone mineral density, bone turnover markers and vitamin D receptor genotypes in healthy Saudi pre- and postmenopausal women. Osteoporosis International, 22(2), 463–475.
- Azmi, M. Y., Junidah, R., Mariam, S. A., Safiah, M. Y., Fatimah, S., Norimah, A. K., ... Tahir, A. (2009). Body mass index (BMI) of adults: findings of the Malaysian Adult Nutrition Survey (MANS). *Malaysian Journal of Nutrition*, 15(2), 97–119.
- Babatunde, O. O., Forsyth, J. J., & Gidlow, C. J. (2012). A meta-analysis of brief highimpact exercises for enhancing bone health in premenopausal women. *Osteoporosis International*, 23(1), 109–119.
- Bacon, L., Stern, J. S., Keim, N. L., & Loan, M. D. V. (2004). Low bone mass in premenopausal chronic dieting obese women. *European Journal of Clinical Nutrition*, 58, 966–971.
- Bainbridge, K. E., Sower, M. F., Crutchfield, M., Lin, X., Jannausch, M., & Harlow, S. D. (2002). Natural history of bone loss over 6 years among premenopausal and early postmenopausal women. *American Journal of Epidemiology*, 156(5), 410–417.
- Bala, Y., Farlay, D., & Boivin, G. (2013). Bone mineralization: from tissue to crystal in normal and pathological contexts. *Osteoporosis International*, 24(8), 2153– 2166.
- Ballane, G., Cauley, J. A., Luckey, M. M., & Fuleihan, G. E.-H. (2014). Secular trends in hip fractures worldwide: Opposing trends east versus west. *Journal of Bone* and Mineral Research, 29(8), 1745–1755.
- Barling, P. M. (2013). Osteoporosis an increasingly important issue for both young and aging citizens of Malaysia. *International E-Journal of Science, Medicine & Education*, 7, 1–3.
- Baron, J. A., Farahmand, B. Y., Weiderpass, E., Michaelsson, K., Alberts, A., Persson, I., & Ljunghall, S. (2001). Cigarette smoking, alcohol consumption, and risk of hip fracture in women. *Achieves of Internal Medicine*, 161, 983–988.
- Barrett-Connor, E., & Goodman-Gruen, D. (1998). Gender differences in insulin-like growth factor and bone mineral density association in old age: the Rancho Bernardo Study. *Journal of Bone and Mineral Research*, *13*(8), 1343–1349.

- Bartl, R., & Frisch, B. (2009). Bone density in osteoporosis. In Osteoporosis (2nd ed., pp. 63–73). Berlin, Germany: Springer-Verlag Berlin Heidelberg.
- Bauer, J. S., Müller, D., Ambekar, A., Dobritz, M., Matsuura, M., Eckstein, F., ... Link, T. M. (2006). Detection of osteoporotic vertebral fractures using multidetector CT. Osteoporosis International, 17(4), 608–615.
- Baxter-Jones, A. D. G., Kontulainen, S. A., Faulkner, R. A., & Bailey, D. A. (2008). A longitudinal study of the relationship of physical activity to bone mineral accrual from adolescence to young adulthood. *Bone*, *43*(6), 1101–1107.
- Beck, T. J., Petit, M. A., Wu, G., LeBoff, M. S., Cauley, J. A., & Chen, Z. (2009). Does obesity really make the femur stronger? BMD, geometry, and fracture incidence in the Women's Health Initiative-Observational Study. *Journal of Bone and Mineral Research*, 24(8), 1369–1379.
- Begg, S. J., Vos, T., Barker, B., Stanley, L., & Lopez, A. D. (2008). Burden of disease and injury in Australia in the new millennium: measuring health loss from diseases, injuries and risk factors. *The Medical Journal of Australia*, 188(1), 36– 40.
- Berg, K. M., Kunins, H. V, Jackson, J. L., Nahvi, S., Chaudhry, A., Harris, K. A. J., ... Arnsten, J. H. (2009). Association between alcohol consumption and both osteoporotic fracture and bone density. *The American Journal of Medicine*, 121(5), 406–418.
- Bernal, P. P. (2010). Osteoporosis in young individuals. *Reumatología Clínica*, 6(4), 217–223.
- Bischoff-Ferrari, H. A., Dietrich, T., Orav, E. J., & Dawson-Hughes, B. (2004). Positive association between 25-hydroxy vitamin D levels and BMD: a population-based study of younger and older adults. *The American Journal of Medicine*, 116(9), 634–639.
- Black, A. E. (2000). Critical evaluation of energy intake using the Goldberg cut-off for energy intake: basal metabolic rate. A practical guide to its calculation, use and limitations. *International Journal of Obesity*, *24*(9), 1119–1130.
- Blake, G. M., & Fogelman, I. (2007). The role of DXA bone density scans in the diagnosis and treatment of osteoporosis. *Postgraduate Medical Journal*, 83(982), 509–517.
- Blake, G. M., & Fogelman, I. (2009). The clinical role of dual energy X-ray absorptiometry. *European Journal of Radiology*, 71(3), 406–414.
- Boivin, G., & Meunier, P. J. (2002). The degree of mineralization of bone tissue measured by computerized quantitative contact microradiography. *Calcified Tissue International*, 70(6), 503–511.
- Bonjour, J.-P., Ammann, P., Chevalley, T., & Rizzoli, R. (2001). Protein intake and bone growth. *Canadian Journal of Applied Physiology*, 26, S153–S166.
- Bonjour, J.-P., Chevalley, T., Ammann, P., Slosman, D., & Rizzoli, R. (2001). Gain in bone mineral mass in prepubertal girls 3.5 years after discontinuation of calcium supplementation: a follow-up study. *Lancet*, 358, 1208–1212.
- Bonjour, J.-P., Schüreh, M.-A., Chevalley, T., Ammann, P., & Rizzoli, R. (1997). Protein intake, IGF -1 and osteoporosis. *Osteoporosis International*, 7(3), 36–42.

- Bonjour, J.-P., Theintz, G., Law, F., Slosman, D., & Rizzoli, R. (1994). Peak bone mass. Osteoporosis International, 1, S7–S13.
- Boskey, A. L., & Coleman, R. (2010). Aging and bone. *Journal of Dental Research*, 89(12), 1333–1348.
- Brennan, S. L., Pasco, J. A., Urquhart, D. M., Oldenburg, B., Wang, Y., & Wluka, A. E. (2011). Association between socioeconomic status and bone mineral density in adults: a systematic review. *Osteoporosis International*, 22(2), 517–527.
- Butler, T. A., & Yingling, V. R. (2013). The effects of delayed puberty on the growth plate. *Journal of Pediatric Orthopedic*, 33(1), 99–105.
- Calvo, M. S., & Tucker, K. L. (2013). Is phosphorus intake that exceeds dietary requirements a risk factor in bone health? *Annals of the New York Academy of Sciences*, 1301, 29–35.
- Calvo, M. S., & Uribarri, J. (2013). Public health impact of dietary phosphorus excess on bone and cardiovascular health in the general population. *The American Journal of Clinical Nutrition*, 98, 6–15.
- Cao, J. J., Johnson, L. K., & Hunt, J. R. (2011). A diet high in meat protein and potential renal acid load increases fractional calcium absorption and urinary calcium excretion without affecting markers of bone resorption or formation in postmenopausal women. *The Journal of Nutrition*, 141(3), 391–397.
- Caporaso, F., Frisch, F., & Sumida, K. D. (2011). Compromised bone health in nonobese, older women with low caloric intake. *Journal of Community Health*, 36, 559–564.
- Carey, J. J., Licata, A. A., & Delaney, M. F. (2006). Biochemical markers of bone turnover. *Clinical Reviews in Bone and Mineral Metabolism*, 4(3), 197–212.
- Cashman, K. D. (2007a). Calcium intake, calcium bioavailability and bone health. *British Journal of Nutrition*, 87(2), S169–S177.
- Cashman, K. D. (2007b). Diet, nutrition, and bone health. *The Journal of Nutrition*, 137, 2507–2512.
- Cech, D. (2012). Prevention of osteoporosis: from infancy through older adulthood. Hong Kong Physiotherapy Journal, 30(1), 6–12.
- Chan, P. J., Nurul, Z. Z., Chuah, J. S., Nabil, M. M. A., Isa, N. M., Sabarul, A. M., & Nazrun, A. S. (2014). Association between risk Factors of osteoporosis and bone mineral density in women of different ethnic groups in a Malaysian hospital. *International Journal of Osteoporosis and Metabolic Disorders*, 7(1), 1–11.
- Charatcharoenwitthaya, N., Khosla, S., Atkinson, E. J., McCready, L. K., & Riggs, B. L. (2007). Effect of blockade of TNF-alpha and interleukin-1 action on bone resorption in early postmenopausal women. *Journal of Bone and Mineral Research*, 22(5), 724–729.
- Chavassieux, P., Seeman, E., & Delmas, P. D. (2007). Insights into material and structural basis of bone fragility from diseases associated with fractures: how determinants of the biomechanical properties of bone are compromised by disease. *Endocrine Reviews*, 28(2), 151–164.

- Chee, S. S., Ismail, M. N., Ng, K. K., & Zawiah, H. (1997). Food intake assessment of adults in rural and urban areas from four selected regions in Malaysia. *Malaysian Journal of Nutrition*, 3, 97–102.
- Chee, W. S. S., Chong, P. N., Chuah, K. A., Karupaiah, T., Mustafa, N., Suniza, S. S., ... Offord-Cavin, E. (2010). Calcium intake, vitamin D and bone health status of post-menopausal Chinese women in Kuala Lumpur. *Malaysian Journal of Nutrition*, 16(2), 233–242.
- Chee, W. S. S., Suriah, A. R., Chan, S. P., Zaitun, Y., & Chan, Y. M. (2003). The effect of milk supplementation on bone mineral density in postmenopausal Chinese women in Malaysia. *Osteoporosis International*, *14*(10), 828–834.
- Chee, W. S. S., Suriah, A. R., Zaitun, Y., Chan, S. P., Yap, S. L., & Chan, Y. M. (2002). Dietary calcium intake in postmenopausal Malaysian women: comparison between the food frequency questionnaire and three-day food records. *Asia Pacific Journal of Clinical Nutrition*, 11(2), 142–146.
- Cheng, S., Lyytikäinen, A., Kroger, H., Lamberg-Allardt, C., Alen, M., Koistinen, A., ... Tylavsky, F. (2005). Effects of calcium, dairy product, and vitamin D supplementation on bone mass accrual and body composition in 10-12-y-old girls: a 2-y randomized trial. *The American Journal of Clinical Nutrition*, 82, 1115–1126.
- Cheung, E., Tsang, S., Bow, C., Soong, C., Yeung, S., Loong, C., ... Kung, A. (2011). Bone loss during menopausal transition among southern Chinese women. *Maturitas*, 69(1), 50–56.
- Chevalley, T., Bonjour, J.-P., Ferrari, S., Hans, D., & Rizzoli, R. (2005). Skeletal site selectivity in the effects of calcium supplementation on areal bone mineral density gain: a randomized, double-blind, placebo-controlled trial in prepubertal boys. *The Journal of Clinical Endocrinology and Metabolism*, 90(6), 3342–3349.
- Cho, G. J., Park, H. T., Shin, J. H., Hur, J. Y., Kim, Y. T., Kim, S. H., ... Kim, T. (2010). Age at menarche in a Korean population: secular trends and influencing factors. *European Journal of Pediatrics*, *169*(1), 89–94. doi:10.1007/s00431-009-0993-1
- Clarke, B. L., & Khosla, S. (2010). Female reproductive system and bone. Archives of Biochemistry and Biophysics, 503(1), 118–128.
- Compston, J. E. (2001). Sex steroids and bone. Physiological Reviews, 81(1), 419-447.
- Cooper, C., Cole, Z. A., Holroyd, C. R., Earl, S. C., Harvey, N. C., Dennison, E. M., ... Kanis, J. A. (2011). Secular trends in the incidence of hip and other osteoporotic fractures. *Osteoporosis International*, 22(5), 1277–1288.
- Cosman, F., Morgan, D. C., Nieves, J. W., Shen, V., Luckey, M. M., Dempster, D. W., ... Parisien, M. (1997). Resistance to bone resorbing effects of PTH in Black women. *Journal of Bone and Mineral Research*, 12(6), 958–966.
- Coulston, A. M., Boushey, C. J., & Ferruzzi, M. G. (2013). *Nutrition in the Prevention and Treatment of Disease* (3rd ed.). San Diego, CA: Elsevier Inc.
- Cure-Cure, C., Cure-Ramírez, P., Teran, E., & Lopez-Jaramillo, P. (2002). Bone-mass peak in multiparity and reduced risk of bone-fractures in menopause. *International Journal of Gynecology & Obstetrics*, *76*, 285–291.

- Damilakis, J., Maris, T. G., & Karantanas, A. H. (2007). An update on the assessment of osteoporosis using radiologic techniques. *European Radiology*, *17*(6), 1591–1602.
- Danielson, M. E., Cauley, J. A., Baker, C. E., Newman, A. B., Dorman, J. S., Towers, J. D., & Kuller, L. H. (1999). Familial resemblance of bone mineral density (BMD) and calcaneal ultrasound attenuation: the BMD in mothers and daughters study. *Journal of Bone and Mineral Research*, 14(1), 102–110.
- Dans, A., Ng, N., Varghese, C., Tai, E. S., Firestone, R., & Bonita, R. (2011). The rise of chronic non-communicable diseases in southeast Asia: time for action. *Lancet*, *377*, 680–689.
- Darling, A. L., Millward, D. J., Torgerson, D. J., Hewitt, C. E., & Lanham-New, S. A. (2009). Dietary protein and bone health: a systematic review and meta-analysis. *The American Journal of Clinical Nutrition*, 90, 1674–1692.
- Das, U. N. (2001). Is obesity an inflammatory condition? Nutrition, 17, 953–966.
- Dawson-Hughes, B., & Harris, S. S. (2002). Calcium intake influences the association of protein intake with rates of bone loss in elderly men and women. *The American Journal of Clinical Nutrition*, 75, 773–779.
- Demir, B., Haberal, A., Geyik, P., Baskan, B., Ozturkoglu, E., Karacay, O., & Deveci, S. (2008). Identification of the risk factors for osteoporosis among postmenopausal women. *Maturitas*, 60(3-4), 253–256.
- Demontiero, O., Vidal, C., & Duque, G. (2012). Aging and bone loss: new insights for the clinician. *Therapeutic Advances in Musculoskeletal Disease*, 4(2), 61–76.
- Department of Statistics Malaysia. (2012a). *Household income and basic amenities survey report*. Retrieved from http://www.statistics.gov.my/portal/index.php?option=com\_content&view=articl e&id=1640&Itemid=111&lang=bm
- Department of Statistics Malaysia. (2012b). *Vital Statistics Malaysia 2012*. Retrieved from https://www.statistics.gov.my/
- Dertina, D., Loro, M. L., Sayre, J., Kaufman, F., & Gilsanz, V. (1998). Childhood bone measurements predict values at young adulthood. *Bone*, 23, S288.
- Desai, M. P., Bhanuprakash, K. V., Khatkhatay, M. I., & Donde, U. M. (2007). Agerelated changes in bone turnover markers and ovarian hormones in premenopausal and postmenopausal Indian women. *Journal of Clinical Laboratory Analysis*, 21(2), 55–60.
- Desai, M. P., Khatkhatay, M. I., Prakash, K. V. B., Savardekar, L. S., Shah, R. S., & Ansari, Z. (2007). Hormonal profiles and biochemical indices of bone turnover in Indian women. *Osteoporosis International*, 18(7), 923–929.
- DiVasta, A. D., & Gordon, C. M. (2013). Exercise and bone: where do we stand? Metabolism Clinical and Experimental, 62(12), 1714–1747.
- Dursun, N., Akin, S., Dursun, E., Sade, I., & Korkusuz, F. (2006). Influence of duration of total breast-feeding on bone mineral density in a Turkish population: does the priority of risk factors differ from society to society? *Osteoporosis International*, 17(5), 651–655.

- Emaus, N., Nguyen, N. D., Almaas, B., Berntsen, G. K., Center, J. R., Christensen, M., ... Fønnebø, V. M. (2013). Serum level of under-carboxylated osteocalcin and bone mineral density in early menopausal Norwegian women. *European Journal of Nutrition*, 52(1), 49–55.
- Engelke, K., Kemmler, W., Lauber, D., Beeskow, C., Pintag, R., & Kalender, W. A. (2006). Exercise maintains bone density at spine and hip EFOPS: a 3-year longitudinal study in early postmenopausal women. *Osteoporosis International*, *17*(1), 133–142.
- Falk, M., & Anderson, C. D. (2012). Measuring sun exposure habits and sun protection behaviour using a comprehensive scoring instrument - an illustration of a possible model based on Likert scale scorings and on estimation of readiness to increase sun protection. *Cancer Epidemiology*, 36(4), e265–e269.
- Farahmand, B. Y., Michaëlsson, K., Baron, J. A., Persson, P. G., & Ljunghall, S. (2000). Body size and hip fracture risk. Swedish Hip Fracture Study Group. *Epidemiology*, 11(2), 214–219.
- Ferrari, S., Rizzoli, R., Slosman, D., & Bonjour, J.-P. (1998). Familial resemblance for bone mineral mass is expressed before puberty. *Journal of Clinical Endocrinology and Metabolism*, 83(2), 358–361.
- Finkelstein, J. S., Brockwell, S. E., Mehta, V., Greendale, G. A., Sowers, M. R., Ettinger, B., ... Neer, R. M. (2008). Bone mineral density changes during the menopause transition in a multiethnic cohort of women. *The Journal of Clinical Endocrinology and Metabolism*, 93(3), 861–868.
- Fox, K. M., Cummings, S. R., Powell-Threets, K., & Stone, K. (1998). Family history and risk of osteoporotic fracture. *Osteoporosis International*, 8(6), 557–562.
- Francucci, C. M., Romagni, P., Camilletti, A., Fiscaletti, P., Amoroso, L., Cenci, G., ... Boscaro, M. (2008). Effect of natural early menopause on bone mineral density. *Maturitas*, 59(4), 323–328.
- Frost, H. M. (1997). On our age-related bone loss: insights from a new paradigm. Journal of Bone and Mineral Research, 12(10), 1539–1546.
- Fu, X., Ma, X., Lu, H., He, W., Wang, Z., & Zhu, S. (2011). Associations of fat mass and fat distribution with bone mineral density in pre- and postmenopausal Chinese women. *Osteoporosis International*, 22(1), 113–119.
- Fujiwara, S., Kasagi, F., Masunari, N., Naito, K., Suzuki, G., & Fukunaga, M. (2003). Fracture prediction from bone mineral density in Japanese men and women. *Journal of Bone and Mineral Research*, 18(8), 1547–1553.
- Garnero, P., Munoz, F., Sornay-Rendu, E., & Delmas, P. D. (2007). Associations of vitamin D status with bone mineral density, bone turnover, bone loss and fracture risk in healthy postmenopausal women. The OFELY study. *Bone*, 40(3), 716–722.
- Gilsanz, V., Chalfant, J., Kalkwarf, H., Zemel, B., Lappe, J., Oberfield, S., ... Winer, K. (2011). Age at onset of puberty predicts bone mass in young adulthood. *The Journal of Pediatrics*, 158(1), 100–105.

- Gilsanz, V., Chalfant, J., Mo, A. O., Lee, D. C., Dorey, F. J., & Mittelman, S. D. (2009). Reciprocal relations of subcutaneous and visceral fat to bone structure and strength. *The Journal of Clinical Endocrinology & Metabolism*, 94(9), 3387–3393.
- Gjesdal, C. G., Halse, J. I., Eide, G. E., Brun, J. G., & Tell, G. S. (2008). Impact of lean mass and fat mass on bone mineral density: The Hordaland Health Study. *Maturitas*, 59(2), 191–200.
- Goff, C. Le, Cavalier, E., Souberbielle, J.-C., González-Antuña, A., & Delvin, E. (2015). Measurement of circulating 25-hydroxyvitamin D: a historical review. *Practical Laboratory Medicine*, *2*, 1–14.
- Goldberg, G. R., Black, A. E., Jebb, S. A., Cole, T. J., Murgatroyd, P. R., Coward, W. A., & Prentice, A. M. (1991). Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *Journal of Clinical Nutrition*, 45, 569–581.
- Gonnelli, S., Caffarelli, C., & Nuti, R. (2014). Obesity and fracture risk. *Clinical Cases in Mineral and Bone Metabolism*, 11(1), 9–14.
- Goodpaster, B. H., Krishnaswami, S., Resnick, H., Kelley, D. E., Haggerty, C., Harris, T. B., ... Newman, A. B. (2003). Association between regional adipose tissue distribution and both type 2 diabetes and impaired glucose tolerance in elderly men and women. *Diabetes Care*, 26(2), 372–379.
- Grainge, M. J., Coupland, C. A. C., Cliffe, S. J., Chilvers, C. E. D., & Hosking, D. J. (1998). Cigarette smoking, alcohol and caffeine consumption, and bone mineral density in postmenopausal women. *Osteoporosis International*, 8(4), 355–363.
- Grainge, M. J., Coupland, C. A. C., Cliffe, S. J., Chilvers, C. E. D., & Hosking, D. J. (1999). Association between a family history of fractures and bone mineral density in early postmenopausal women. *Bone*, 24(5), 507–512.
- Greendale, G. A., Edelstein, S., & Barrett-Connor, E. (1997). Endogenous sex steroids and bone mineral density in older women and men: the Rancho Bernardo Study. *Journal of Bone and Mineral Research*, 12(11), 1833–1843.
- Guillemin, F., Martinez, L., Calvert, M., Cooper, C., Ganiats, T., Gitlin, M., ... Freemantle, N. (2013). Fear of falling, fracture history, and comorbidities are associated with health-related quality of life among European and US women with osteoporosis in a large international study. *Osteoporosis International*, 24(12), 3001–3010.
- Gunter, K., Baxter-Jones, A. D. G., Mirwald, R. L., Almstedt, H., Fuller, A., Durski, S., & Snow, C. (2008). Jump starting skeletal health: a 4-year longitudinal study assessing the effects of jumping on skeletal development in pre and circum pubertal children. *Bone*, 42(4), 710–718.
- Handa, R., Kalla, A. A., & Maalouf, G. (2008). Osteoporosis in developing countries. Best Practice & Research Clinical Rheumatology, 22(4), 693–708.
- Hannan, M. T., Tucker, K. L., Dawson-Hughes, B., Cupples, L. A., Felson, D. T., & Kiel, D. P. (2000). Effect of dietary protein on bone loss in elderly men and women: the Framingham Osteoporosis Study. *Journal of Bone and Mineral Research*, 15(12), 2504–2512.

- Hanwell, H. E. C., Vieth, R., Cole, D. E. C., Scillitani, A., Modoni, S., Frusciante, V., ... Carnevale, V. (2010). Sun exposure questionnaire predicts circulating 25hydroxyvitamin D concentrations in Caucasian hospital workers in southern Italy. *Journal of Steroid Biochemistry and Molecular Biology*, 121, 334–337.
- Harinarayan, C. V. (2005). Prevalence of vitamin D insufficiency in postmenopausal south Indian women. *Osteoporosis International*, *16*(4), 397–402.
- Harvey, N. C., Matthews, P., Collins, R., & Cooper, C. (2013). Osteoporosis epidemiology in UK Biobank: A unique opportunity for international researchers. *Osteoporosis International*, 24(12), 2903–2905.
- Havill, L. M., Mahaney, M. C., Binkley, T. L., & Specker, B. L. (2007). Effects of genes, sex, age, and activity on BMC, bone size, and areal and volumetric BMD. *Journal of Bone and Mineral Research*, 22(5), 737–746.
- Hawkins, R. (2013). Total 25-OH vitamin D concentrations in Chinese, Malays and Indians. *Annals of Laboratory Medicine*, 33(2), 156–158.
- Heaney, R. P. (2007). Effects of protein on the calcium economy. *International Congress Series*, 1297, 191–197.
- Heaney, R. P., Abrams, S., Dawson-Hughes, B., Looker, A., Marcus, R., Matkovic, V., & Weaver, C. (2000). Peak bone mass. *Osteoporosis International*, 11(12), 985– 1009.
- Henderson, P. H., Sowers, M., Kutzko, K. E., & Jannausch, M. L. (2000). Bone mineral density in grand multiparous women with extended lactation. *American Journal of Obstetrics and Gynecology*, 182(6), 1371–1377.
- Hernlund, E., Svedbom, A., Ivergård, M., Compston, J., Cooper, C., Stenmark, J., ... Kanis, J. A. (2013). Osteoporosis in the European Union: medical management, epidemiology and economic burden. *Archives of Osteoporosis*, 8(136), 1–115.
- Hillier, T. A., Rizzo, J. H., Pedula, K. L., Stone, K. L., Cauley, J. A., Bauer, D. C., & Cummings, S. R. (2003). Nulliparity and fracture risk in older women: the Study of Osteoporotic Fractures. *Journal of Bone and Mineral Research*, *18*(5), 893–899.
- Hind, K., & Burrows, M. (2007). Weight-bearing exercise and bone mineral accrual in children and adolescents: a review of controlled trials. *Bone*, 40(1), 14–27.
- Hinriksdóttir, G., Arngrímsson, S. Á., Misic, M. M., & Evans, E. M. (2013). Lean soft tissue contributes more to bone health than fat mass independent of physical activity in women across the lifespan. *Maturitas*, 74(3), 264–269.
- Hinton, P. S., Rector, R. S., Donnelly, J. E., Smith, B. K., & Bailey, B. (2010). Total body bone mineral content and density during weight loss and maintenance on a low- or recommended-dairy weight-maintenance diet in obese men and women. *European Journal of Clinical Nutrition*, 64(4), 392–399.
- Ho, S. C., Chen, Y., & Woo, J. L. F. (2005). Educational level and osteoporosis risk in postmenopausal Chinese women. *American Journal of Epidemiology*, 161(7), 680–690.
- Ho, S. C., Woo, J., Chan, S. G., Chan, C. S. Y., & Leung, P. C. (2007). A study of bone mass and bone loss in peri-menopausal Chinese women. *Hong Kong Medical Journal*, 13, 44–46.

- Hofbauer, L. C., Khosla, S., Dunstan, C. R., Lacey, D. L., Spelsberg, T. C., & Riggs, B. L. (1999). Estrogen stimulates gene expression and protein production of osteoprotegerin in human osteoblastic cells. *Endocrinology*, 140(9), 4367–4370.
- Hofman, M., Landewé-Cleuren, S., Wojciechowski, F., & Kruseman, A. N. (2009). Prevalence and clinical determinants of low bone mineral density in anorexia nervosa. *European Journal of Internal Medicine*, 20(1), 80–84.
- Hoidrup, S., Gronbaek, M., Gottschau, A., Lauritzen, J. B., & Schroll, M. (1999). Alcohol intake, beverage preference, and risk of hip fracture in men and women. *American Journal of Epidemiology*, 149(11), 993–1001.
- Holick, M. E. (2001). A perspective on the beneficial effects of moderate exposure to sunlight: bone health, cancer prevention, mental health and well being. In P. U. Giacomoni (Ed.), *Sun Protection in Man* (pp. 19–24). Amsterdam, The Netherlands: Elsevier Science B. V.
- Holick, M. F. (2007). Vitamin D deficiency. *The New England Journal of Medicine*, 357, 266–281.
- Holick, M. F., Binkley, N. C., Bischoff-Ferrari, H. A., Gordon, C. M., Hanley, D. A., Heaney, R. P., ... Weaver, C. M. (2011). Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society Clinical Practice Guideline. *The Journal of Clinical Endocrinology and Metabolism*, 96(7), 1911–1930.
- Holm, K., Dan, A., Wilbur, J., Li, S., & Walker, J. (2010). A longitudinal study of bone density in midlife women. *Health Care for Women International*, 23(6-7), 678–691.
- Holroyd, C., Cooper, C., & Dennison, E. (2008). Epidemiology of osteoporosis. Best Practice & Research Clinical Endocrinology & Metabolism, 22(5), 671–685.
- Ho-Pham, L. T., & Campbell, L. V. (2011). More on body fat cutoff points. *Mayo Clinic Proceedings*, 86(6), 584–585.
- Ho-Pham, L. T., Nguyen, N. D., Lai, T. Q., & Nguyen, T. V. (2010). Contributions of lean mass and fat mass to bone mineral density: a study in postmenopausal women. *BMC Musculoskeletal Disorders*, 11, 59.
- Hosseinpanah, F., Rambod, M., Hossein-Nejad, A., Larijani, B., & Azizi, F. (2008). Association between vitamin D and bone mineral density in Iranian postmenopausal women. *Journal of Bone and Mineral Metabolism*, 26(1), 86–92.
- Hsu, Y.-H., Venners, S. A., Terwedow, H. A., Feng, Y., Niu, T., Li, Z., ... Xu, X. (2006). Relation of body composition, fat mass, and serum lipids to osteoporotic fractures and bone mineral density in Chinese men and women. *The American Journal of Clinical Nutrition*, 83, 146–154.
- Hu, W.-W., Zhang, Z., He, J.-W., Fu, W.-Z., Wang, C., Zhang, H., ... Zhang, Z.-L. (2013). Establishing reference intervals for bone turnover markers in the healthy Shanghai population and the relationship with bone mineral density in postmenopausal women. *International Journal of Endocrinology*, 2013, 1–7.
- Hui, S. L., Perkins, A. J., Zhou, L., Longcope, C., Econs, M. J., Peacock, M., ... Johnston, C. C. (2002). Bone loss at the femoral neck in premenopausal white women: effects of weight change and sex-hormone levels. *The Journal of Clinical Endocrinology & Metabolism*, 87, 1539–1543.

- Hunt, J. R., Johnson, L. K., & Roughead, Z. K. F. (2009). Dietary protein and calcium interact to influence calcium retention: a controlled feeding study. *American Journal of Clinical Nutrition*, 89, 1357–1365.
- Hunter, D. J., Lange, M. de, Andrew, T., Snieder, H., MacGregor, A. J., & Spector, T. D. (2001). Genetic variation in bone mineral density and calcaneal ultrasound: a study of the influence of menopause using female twins. *Osteoporosis International*, 12(5), 406–411.
- Iacono, M. V. (2007). Osteoporosis: A national public health priority. Journal of Perianesthesia Nursing, 22(3), 175–183.
- Ihle, R., & Loucks, A. B. (2004). Dose-response relationships between energy availability and bone turnover in young exercising women. *Journal of Bone and Mineral Research*, 19(8), 1231–1240.
- Iki, M., Morita, A., Ikeda, Y., Sato, Y., Akiba, T., Matsumoto, T., ... Yoneshima, H. (2006). Biochemical markers of bone turnover predict bone loss in perimenopausal women but not in postmenopausal women-the Japanese Population-based Osteoporosis (JPOS) Cohort Study. Osteoporosis International, 17(7), 1086–1095.
- Institute for Public Health. (2011a). National Health and Morbidity Survey 2011 (NHMS 2011). Vol. II: Non-Communicable Diseases. Kuala Lumpur, Malaysia: Institute for Public Health, Ministry of Health, Malaysia.
- Institute for Public Health. (2011b). National Health and Morbidity Survey 2011 fact sheet.
- Institute of Medicine. (1997). Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride. Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food and Nutrition Board, Institute of Medicine. Washington, DC: National Academy Press.
- Institute of Medicine. (2005). Dietary Reference Intakes for Energy, Carbohydrates, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. Food and Nutrition Board, Institute of Medicine. National Academy Press, Washington DC.
- International Osteoporosis Foundation. (2009). *The Asian Audit Epidemiology, costs* and burden of osteoporosis in Asia 2009. Nyon, Switzerland.
- International Osteoporosis Foundation. (2015). Calcium content of common foods. Retrieved July 20, 2015, from http://www.iofbonehealth.org/osteoporosismusculoskeletal-disorders/osteoporosis/prevention/calcium/calcium-contentcommon-foods
- International Society for Clinical Densitometry. (2007). Official positions of the International Society for Clinical Densitometry. West Hartford, CT.
- Ismail, M. N., Chee, S. S., Nawawi, H., Yusoff, K., Lim, T. O., & James, W. P. T. (2002). Obesity in Malaysia. *Obesity Reviews*, 3, 203–208.
- Jensen, C., Holloway, L., Block, G., Spiller, G., Gildengorin, G., Gunderson, E., ... Marcus, R. (2002). Long term effects of nutrient intervention on markers of bone remodeling and calciotropic hormones in late-postmenopausal women. *American Journal of Clinical Nutrition*, 75(6), 1114–1120.

- Jesudason, D., & Clifton, P. (2011). The interaction between dietary protein and bone health. *Journal of Bone and Mineral Metabolism*, 29, 1–14.
- Johansson, H., Odén, A., Kanis, J. a, McCloskey, E. V, Morris, H. A., Cooper, C., & Vasikaran, S. (2014). A meta-analysis of reference markers of bone turnover for prediction of fracture. *Calcified Tissue International*, 94(5), 560–567.
- Johnell, O., & Kanis, J. A. (2006). An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporosis International*, 17(12), 1726–1733.
- Kalkwarf, H. J., & Specker, B. L. (2002). Bone mineral changes during pregnancy and lactation. *Endocrine*, *17*(1), 49–53.
- Kanis, J. A. (2002). Diagnosis of osteoporosis and assessment of fracture risk. *Lancet*, 359(9321), 1929–1936.
- Kanis, J. A. (2007). Assessment of osteoporosis at the primary health care level. Technical Report. World Health Organization Collaborating Centre for Metabolic Bone Diseases, University of Sheffield, UK. Geneva.
- Kanis, J. A., Johansson, H., Oden, A., Johnell, O., De Laet, C., Eisman, J. A., ... Tenenhouse, A. (2004). A family history of fracture and fracture risk: a metaanalysis. *Bone*, 35(5), 1029–1037.
- Kanis, J. A., Johnell, O., Laet, C. D., Johansson, H., Oden, A., Delmas, P., ... Tenenhouse, A. (2004). A meta-analysis of previous fracture and subsequent fracture risk. *Bone*, 35(2), 375–382.
- Kanis, J. A., Johnell, O., Oden, A., Johansson, H., Laet, C. D., Eisman, J. A., ... Tenenhouse, A. (2005). Smoking and fracture risk: a meta-analysis. *Osteoporosis International*, 16(2), 155–162.
- Kanis, J. A., McCloskey, E. V, Takats, D. De, & Pande, K. (1999). Clinical assessment of bone mass, quality and architecture. *Osteoporosis International*, 9(2), S24–S28.
- Kanis, J. A., Odén, A., McCloskey, E. V, Johansson, H., Wahl, D. A., & Cooper, C. (2012). A systematic review of hip fracture incidence and probability of fracture worldwide. *Osteoporosis International*, 23(9), 2239–2256.
- Kaptoge, S., Reid, D. M., Scheidt-Nave, C., Poor, G., Pols, H. A. P., Khaw, K.-T., ... Reeve, J. (2007). Geographic and other determinants of BMD change in European men and women at the hip and spine. A population-based study from the Network in Europe for Male Osteoporosis (NEMO). *Bone*, 40(3), 662–673.
- Kemi, V. E., Kärkkäinen, M. U. M., & Lamberg-Allardt, C. J. E. (2006). High phosphorus intakes acutely and negatively affect Ca and bone metabolism in a dose-dependent manner in healthy young females. *British Journal of Nutrition*, 96, 545–552.
- Kemmler, W., Lauber, D., Weineck, J., Hensen, J., Kalender, W., & Engelke, K. (2004). Benefits of 2 years of intense exercise on bone density, physical fitness, and blood lipids in early postmenopausal osteopenic women. *Achieves of Internal Medicine*, 164, 1084–1091.

- Kerstetter, J. E., O'Brien, K. O., & Insogna, K. L. (2003). Dietary protein, calcium metabolism, and skeletal homeostasis. *The American Journal of Clinical Nutrition*, 78, 584–592.
- Khaw, K., Sneyd, M., & Compston, J. (1992). Bone density, parathyroid hormone and 25-hydroxyvitaminD concentrations in middle-aged women. *British Medical Journal*, 305, 273–277.
- Khosla, S., Atkinson, E. J., Iii, L. J. M., & Riggs, B. L. (1997). Effects of age and estrogen status on serum parathyroid hormone levels and biochemical markers of bone turnover in women: a population-based study. *Journal of Clinical Endocrinology and Metabolism*, 82(5), 1522–1527.
- Khosla, S., Riggs, B. L., Atkinson, E. J., Oberg, A. L., Mcdaniel, L. J., Holets, M., ... Melton, L. J. (2006). Effects of sex and age on bone microstructure at the ultradistal radius: a population-based noninvasive in vivo assessment. *Journal of Bone and Mineral Research*, 21(1), 124–131.
- Khosla, S., Riggs, B. L., Robb, R. A., Camp, J. J., Achenbach, S. J., Oberg, A. L., ... Melton, L. J. (2005). Relationship of volumetric bone density and structural parameters at different skeletal sites to sex steroid levels in women. *The Journal* of Clinical Endocrinology and Metabolism, 90(9), 5096–5103.
- Kim, K. H., Lee, K., Ko, Y.-J., Kim, S. J., Oh, S. I., Durrance, D. Y., ... Park, S. M. (2012). Prevalence, awareness, and treatment of osteoporosis among Korean women: The Fourth Korea National Health and Nutrition Examination Survey. *Bone*, 50(5), 1039–1047.
- Klotzbuecher, C. M., Ross, P. D., Landsman, P. B., Abbott, T. A., & Berger, M. (2000). Patients with prior fractures have an increased risk of future fractures: a summary of the literature and statistical synthesis. *Journal of Bone and Mineral Research*, 15(4), 721–739.
- Kohrt, W. M., Bloomfield, S. a, Little, K. D., Nelson, M. E., & Yingling, V. R. (2004). American College of Sports Medicine position stand: physical activity and bone health. *Medicine and Science in Sports and Exercise*, 36, 1985–1996.
- Kojima, N., Douchi, T., Kosha, S., & Nagata, Y. (2002). Cross-sectional study of the effects of parturition and lactation on bone mineral density later in life. *Maturitas*, 41(3), 203–209.
- Koller, D. L., Ichikawa, S., Lai, D., Padgett, L. R., Doheny, K. F., Pugh, E., ... Foroud, T. (2010). Genome-wide association study of bone mineral density in premenopausal European-American women and replication in African-American women. *The Journal of Clinical Endocrinology & Metabolism*, 95(4), 1802– 1809.
- Kruger, M. C., Schollum, L. M., Kuhn-Sherlock, B., Hestiantoro, A., Wijanto, P., Li-Yu, J., ... Eastell, R. (2010). The effect of a fortified milk drink on vitamin D status and bone turnover in post-menopausal women from South East Asia. *Bone*, 46(3), 759–767.
- Kučukalić-Selimović, E., Valjevac, A., & Hadžović-džuvo, A. (2013). The utility of procollagen type 1 N-terminal propeptide for the bone status assessment in postmenopausal women. *Bosnian Journal of Basic Medical Sciences*, 13(4), 259–265.

- Kuwahata, A., Kawamura, Y., Yonehara, Y., Matsuo, T., Iwamoto, I., & Douchi, T. (2008). Non-weight-bearing effect of trunk and peripheral fat mass on bone mineral density in pre- and post-menopausal women. *Maturitas*, 60(3-4), 244– 247.
- Kveiborg, M., Flyvbjerg, A., Rattan, S. I. S., & Kassem, M. (2000). Changes in the insulin-like growth factor-system may contribute to in vitro age-related impaired osteoblast functions. *Experimental Gerontology*, 35(8), 1061–1074.
- Laet, C. D., Kanis, J. A., Odén, A., Johanson, H., Johnell, O., Delmas, P., ... Tenenhouse, A. (2005). Body mass index as a predictor of fracture risk: a metaanalysis. *Osteoporosis International*, 16(11), 1330–1338.
- Lamberg-Allardt, C., Karp, H., & Kemi, V. (2010). Phosphorus and bone. In P. Burckhardt, B. Dawnson-Hughes, & C. Weaver (Eds.), *Nutritional influences on bone health* (pp. 87–97). Springer-Verlag London Limited.
- Lau, E. M., Suriwongpaisal, P., Lee, J. K., Das De, S., Festin, M. R., Saw, S. M., ... Sambrook, P. (2001). Risk factors for hip fracture in Asian men and women: the Asian osteoporosis study. *Journal of Bone and Mineral Research : The Official Journal of the American Society for Bone and Mineral Research*, 16(3), 572–80. doi:10.1359/jbmr.2001.16.3.572
- Lau, H. H. L., Ng, M. Y. M., Ho, A. Y. Y., Luk, K. D. K., & Kung, A. W. C. (2005). Genetic and environmental determinants of bone mineral density in Chinese women. *Bone*, 36(4), 700–709.
- Lee, J. K., & Khir, A. S. M. (2007). The incidence of hip fracture in Malaysians above 50 years of age: variation in different ethnic groups. *APLAR Journal of Rheumatology*, 10(4), 300–305.
- Lee, K., Kim, K., Kim, H., Seo, J., & Song, S. (2014). Association between dietary calcium and phosphorus intakes, dietary calcium/ phosphorus ratio and bone mass in the Korean population. *Nutrition Journal*, 13(114), 1–8.
- Lei, S.-F., Deng, F.-Y., Li, M.-X., Dvornyk, V., & Deng, H.-W. (2004). Bone mineral density in elderly Chinese: effects of age, sex, weight, height, and body mass index. *Journal of Bone and Mineral Metabolism*, 22(1), 71–78.
- Lenora, J., Ivaska, K. K., Obrant, K. J., & Gerdhem, P. (2007). Prediction of bone loss using biochemical markers of bone turnover. *Osteoporosis International*, 18(9), 1297–1305.
- Lenora, J., Lekamwasam, S., & Karlsson, M. K. (2009). Effects of multiparity and prolonged breast-feeding on maternal bone mineral density: a community-based cross-sectional study. *BMC Women's Health*, 9(19), 1–6.
- Li, M., Li, Y., Deng, W., Zhang, Z., Deng, Z., Hu, Y., ... Xu, L. (2014). Chinese bone turnover marker study: reference ranges for C-terminal telopeptide of type I collagen and procollagen I N-terminal peptide by age and gender. *PLoS ONE*, 9(8), e103841.
- Li, S., Wagner, R., Holm, K., Lehotsky, J., & Zinaman, M. J. (2004). Relationship between soft tissue body composition and bone mass in perimenopausal women. *Maturitas*, 47(2), 99–105.

- Lim, P. S., Ong, F. B., Adeeb, N., Seri, S. S., Noor-Aini, M. Y., Shamsuddin, K., ... Wan, H. W. H. (2005). Bone health in urban midlife Malaysian women: risk factors and prevention. *Osteoporosis International*, 16(12), 2069–2079.
- Lips, P. (2001). Vitamin D deficiency and secondary hyperparathyroidism in the elderly: consequences for bone loss and fractures and therapeutic implications. *Endocrine Reviews*, *22*, 477–501.
- Liu, J., Zhao, H., Ning, G., Chen, Y., Zhang, L., Sun, L., ... Chen, J. (2008). IGF-1 as an early marker for low bone mass or osteoporosis in premenopausal and postmenopausal women. *Journal of Bone and Mineral Metabolism*, 26(2), 159– 164.
- Livingstone, M. B. E., & Black, A. E. (2003). Markers of the validity of reported energy intake. *The Journal of Nutrition*, 133, 895–920.
- Löfman, O., Magnusson, P., Toss, G., & Larsson, L. (2005). Common biochemical markers of bone turnover predict future bone loss: a 5-year follow-up study. *International Journal of Clinical Chemistry and Diagnostic Laboratory Medicine*, 356(1-2), 67–75.
- Loucks, A. B., & Thuma, J. R. (2003). Luteinizing hormone pulsatility is disrupted at a threshold of energy availability in regularly menstruating women. *Journal of Clinical Endocrinology and Metabolism*, 88(1), 297–311.
- Lu, H., Fu, X., Ma, X., Wu, Z., He, W., Wang, Z., ... Zhu, S. (2011). Relationships of percent body fat and percent trunk fat with bone mineral density among Chinese, black, and white subjects. *Osteoporosis International*, 22(12), 3029–3035.
- Luckey, M., Wallenstein, S., & Lapinski, R. (1996). A prospective study of bone loss in African-American and White women: a clinical research center study. *The Journal of Clinical Endocrinology & Metabolism*, *81*, 2948–2956.
- Lundberg, P., Lundgren, I., Mukohyama, H., Lehenkari, P. P., Horton, M. A., Lerner, U. L. F. H., & Biology, O. C. (2001). Vasoactive Intestinal Peptide (VIP)/pituitary adenylate cyclase-activating peptide receptor subtypes in mouse calvarial osteoblasts : presence of VIP-2 receptors and differentiation-induced expression of VIP-1 receptors. *Endocrinology*, 142(1), 339–347.
- Lynn, H. S., Lau, E. M. C., Au, B., & Leung, P. C. (2005). Bone mineral density reference norms for Hong Kong Chinese. *Osteoporosis International*, 16(12), 1663–1668.
- Macdonald, H. M., Mavroeidi, A., Barr, R. J., Black, A. J., Fraser, W. D., & Reid, D. M. (2008). Vitamin D status in postmenopausal women living at higher latitudes in the UK in relation to bone health, overweight, sunlight exposure and dietary vitamin D. *Bone*, 42(5), 996–1003.
- Maghraoui, A. E., Guerboub, A. A., Mounach, A., Ghozlani, I., Nouijai, A., Ghazi, M., ... Tazi, M. A. (2007). Body mass index and gynecological factors as determinants of bone mass in healthy Moroccan women. *Maturitas*, 56(4), 375– 382.
- Malhotra, N., & Mithal, A. (2008). Osteoporosis in Indians. *Indian Journal of Medical Research*, *127*(3), 263–268.

- Marques, E. A., Moreira, P., Wanderley, F., Pizarro, A. N., Leão-Rosas, J. P., Mota, J., & Carvalho, J. (2012). Appendicular fat mass is positively associated with femoral neck bone mineral density in older women. *Menopause: The Journal of The North American Menopause Society*, 19(3), 311–318.
- Matkovic, V., & Visy, D. (2015). Nutrition and bone health during skeletal modeling and bone consolidation of childhood and adolescence. In M. F. Holick & J. W. Nieves (Eds.), *Nutrition and Bone Health* (2nd ed., pp. 199–216). New York, NY: Springer New York.
- McAuley, K. A., Jones, S., Lewis-Barned, N. J., Manning, P., & Goulding, A. (1997). Low vitamin D status is common among elderly Dunedin women. *New Zealand Medical Journal*, 110, 275–277.
- McDonnell, P., McHugh, P. E., & O'Mahoney, D. (2007). Vertebral osteoporosis and trabecular bone quality. *Annals of Biomedical Engineering*, *35*(2), 170–189.
- McLeod, K. J., Rubin, C. T., Otter, M. W., & Qin, Y. X. (2014). Skeletal cell stresses and bone adaptation. *The American Journal of the Medical Sciences*, 316(3), 176–183.
- McLernon, D. J., Powell, J. J., Jugdaohsingh, R., & Macdonald, H. M. (2012). Do lifestyle choices explain the effect of alcohol on bone mineral density in women around menopause? *The American Journal of Clinical Nutrition*, 95, 1261–1269.
- McTernan, P. G., Anderson, L. A., Anwar, A. J., Eggo, M. C., Crocker, J., Barnett, A. H., ... Kumar, S. (2002). Glucocorticoid regulation of P450 aromatase activity in human adipose tissue: gender and site differences. *The Journal of Clinical Endocrinology & Metabolism*, 87(3), 1327–1336.
- Mehta, G., Taylor, P., Petley, G., Dennison, E., Cooper, C., & Walker-Bone, K. (2004). Bone mineral status in immigrant Indo-Asian women. *Monthly Journal of the Association of Physicians*, 97(2), 95–99.
- Meigs, J. B., Wilson, P. W. F., Fox, C. S., Vasan, R. S., Nathan, D. M., Sullivan, L. M., & D'Agostino, R. B. (2006). Body mass index, metabolic syndrome, and risk of type 2 diabetes or cardiovascular disease. *The Journal of Clinical Endocrinology* and Metabolism, 91, 2906–2912.
- Melton, L. J., Chrischilles, E. A., Cooper, C., Lane, A. W., & Riggs, B. L. (2005). How many women have osteoporosis? *Journal of Bone and Mineral Research*, 20(5), 886–892.
- Melton, L. J., Thamer, M., Ray, N. F., Chan, J. K., Chesnut, C. H., Einhorn, T. A., ... Siris, E. S. (1997). Fractures attributable to osteoporosis: report from the National Osteoporosis Foundation. *Journal of Bone and Mineral Research*, *12*(1), 16–23.
- Michelsen, J., Wallaschofski, H., Friedrich, N., Spielhagen, C., Rettig, R., Ittermann, T., ... Hannemann, A. (2013). Reference intervals for serum concentrations of three bone turnover markers for men and women. *Bone*, 57(2), 399–404.
- Ministry of Education Malaysia. (2012). Preliminary Report: Malaysia Education Blueprint 2013-2025. Retrieved from http://www.moe.gov.my/en/pelanpembangunan-pendidikan-malaysia-2013-2025

- Ministry of Federal Territories. (2014). Official website of Greater Kuala Lumpur/ Klang Valley. Retrieved January 12, 2015, from http://app.kwpkb.gov.my/greaterklkv/home/
- Ministry of Health Malaysia. (2006). *Malaysia NCD surveillance 2006: NCD risk factors in Malaysia.* Putrajaya: Disease Control Division, Ministry of Health, Malaysia.
- Ministry of Health Malaysia. (2012). *Clinical guidance on management of osteoporosis 2012.* (Vol. 12). Selangor, Malaysia: Malaysian Osteoporosis Society.
- Mirnalini, K., Zalilah, M. S., Safiah, M. Y., Tahir, A., Haslinda, M. D. S., Rohana, D. S., ... Normah, H. (2008). Energy and nutrient intakes: findings from the Malaysian Adult Nutrition Survey (MANS). *Malaysian Journal of Nutrition*, 14(1), 1–24.
- Misra, M., & Klibanski, A. (2014). Anorexia nervosa and bone. *Journal of Endocrinology*, 221, 163–176.
- Mitnick, M. A., Grey, A., Masiukiewicz, U., Bartkiewicz, M., Rios-Velez, L., Friedman, S., ... Insogna, K. (2001). Parathyroid hormone induces hepatic production of bioactive interleukin-6 and its soluble receptor. *American Journal* of Physiology Endocrinology and Metabolism, 8020, 405–412.
- Mittal, S., Kumar, A., Gupta, R. K., Kapoor, S., Gulati, P., & Shukla, D. K. (2011). Comparison of bone mineral density and its variables between premenopausal and postmenopausal women. *The Journal of Obstetrics and Gynecology of India*, *61*(2), 200–204.
- Mizuno, K., Suzuki, A., Ino, Y., Asada, Y., Kikkawa, F., & Tomoda, Y. (1995). Postmenopausal bone loss in Japanese women. *International Journal of Gynecology & Obstetrics*, 50(1), 33–39.
- Moayyeri, A., Hammond, C. J., Hart, D. J., & Spector, T. D. (2012). Effects of age on genetic influence on bone loss over 17 years in women: the Healthy Ageing Twin Study (HATS). *Journal of Bone and Mineral Research*, 27(10), 2170–2178.
- Moe, S. M. (2008). Disorders involving calcium, phosphorus, and magnesium. *Primary Care*, 35(2), 215-vi.
- Mol, G. D., Lisdonk, E. H. V. D., Smits, J. P. J. M., Hoogen, J. M. P. V. D., Bor, J. H. J., & Westert, G. P. (2005). A widening health gap in general practice? socioeconomic differences in morbidity between 1975 and 2000 in the Netherlands. *Public Health*, 119(7), 616–625.
- Montazerifar, F., Karajibani, M., Alamian, S., Sandoughi, M., Zakeri, Z., & Dashipour, A. R. (2014). Age, weight and body mass index effect on bone mineral density in postmenopausal women. *Health Scope*, 3(2), e14075.
- Moon, J. H., Koo, B. K., & Moon, M. K. (2015). Optimal high-density lipoprotein cholesterol cutoff for predicting cardiovascular disease: comparison of the Korean and US National Health and Nutrition Examination Surveys. *Journal of Clinical Lipidology*, 9, 334–342.

- Moy, F. M. (2011). Vitamin D status and its associated factors of free living Malay adults in a tropical country, Malaysia. *Journal of Photochemistry and Photobiology B: Biology*, 104(3), 444–448.
- Moy, F.-M., & Bulgiba, A. (2011). High prevalence of vitamin D insufficiency and its association with obesity and metabolic syndrome among Malay adults in Kuala Lumpur, Malaysia. *BMC Public Health*, *11*(1), 735.
- Musa, N., Chee, W. S. S., Rokiah, P., Tan, A. T. B., Chew, Y. Y., Nusaibah, A. R. S., & Chan, S. P. (2013). Effects of sun exposure on 25 (OH) vitamin D concentration in urban and rural women in Malaysia. *Asia Pacific Journal of Clinical Nutrition*, 22(3), 391–399.
- Nakamura, K., Tsugawa, N., Saito, T., Ishikawa, M., Tsuchiya, Y., Hyodo, K., ... Yamamoto, M. (2008). Vitamin D status, bone mass, and bone metabolism in home-dwelling postmenopausal Japanese women: Yokogoshi Study. *Bone*, 42(2), 271–277.
- Namwongprom, S., Rojanasthien, S., Mangklabruks, A., Soontrapa, S., Boonsong, C. W., & Ongphiphadhanakul. (2013). Effect of fat mass and lean mass on bone mineral density in postmenopausal and perimenopausal Thai women. *International Journal of Women's Health*, 5, 87–92.
- National Institute of Health Consensus Conference. (1996). Physical activity and cardiovascular health. *Journal of the American Medical Association*, 276(3), 241–246.
- National Osteoporosis Foudation. (2010). *Clinician's Guide to Prevention and Treatment of Osteoporosis*. Washington, DC: National Osteoporosis Foundation.
- National Osteoporosis Foudation. (2015). A guide to calcium-rich foods. Retrieved July 20, 2015, from http://nof.org/articles/886
- Neer, R. M. (2010). Bone loss across the menopausal transition. Annals of the New York Academy of Sciences, 1192, 66–71.
- New, S. A., Bolton-Smith, C., Grubb, D. A., & Reid, D. M. (1997). Nutritional influences on bone mineral density: sectional study in premenopausal a crosssectional study in premenopausal women. *The American Journal of Clinical Nutrition*, 65, 1831–1839.
- Newton-John, H. F., & Morgan, B. D. (1970). The loss of bone with age: osteoporosis and fractures. *Clinical Orthopaedics and Related Research*, 71, 229–232.
- Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., ... Gakidou, E. (2014). Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*, 384, 766–781.
- Nguyen, H. T. T., Schoultz, B. V., Nguyen, T. V., Dzung, D. N., Duc, P. T. M., Thuy, V. T., & Hirschberg, A. L. (2012). Vitamin D deficiency in northern Vietnam: prevalence, risk factors and associations with bone mineral density. *Bone*, 51(6), 1029–1034.
- Nguyen, T. V, Center, J. R., & Eisman, J. A. (2000). Osteoporosis in elderly men and women: effects of dietary calcium, physical activity, and body mass index. *Journal of Bone and Mineral Research*, *15*(2), 322–331.

- NIH Consensus Development Panel on Osteoporosis Prevention and Therapy. (2001). Osteoporosis prevention, diagnosis, and therapy. *Journal of the American Medical Association*, 285, 785–795.
- Niimi, R., Kono, T., Nishihara, A., Hasegawa, M., Matsumine, A., Kono, T., & Sudo, A. (2014). Determinants associated with bone mineral density increase in response to daily teriparatide treatment in patients with osteoporosis. *Bone*, 66, 26–30.
- Nur, H., Toraman, N. F., Arica, Z., Sarier, N., & Samur, A. (2013). The relationship between body composition and bone mineral density in postmenopausal Turkish women. *Rheumatology International*, 33(3), 607–612.
- Ohtani, K. (2000). Bootstrapping R2 and adjusted R2 in regression analysis. *Economic Modelling*, 17, 473–483.
- Okano, H., Mizunuma, H., Soda, M., Kagami, I., Miyamoto, S., Ohsawa, M., ... Shibata, H. (1998). The long-term effect of menopause on postmenopausal bone loss in Japanese women: results from a prospective study. *Journal of Bone and Mineral Research*, 13(2), 303–309.
- Oliveri, B., Parisi, M. S., Zeni, S., & Mautalen, C. (2004). Mineral and bone mass changes during pregnancy and lactation. *Nutrition*, 20(2), 235–240.
- Olmos, J. M., Valero, C., Barrio, A. G. D., Amado, J. A., Hernández, J. L., Menéndez-Arango, J., & González-Macías, J. (2010). Time course of bone loss in patients with anorexia nervosa. *International Journal of Eating Disorders*, 43, 537–542.
- Ooms, M. E., Lips, P., Roos, J. C., Vijgh, W. J. V. D., Popp-Snijders, C., Bezemer, P. D., & Bouter, L. M. (1995). Vitamin D status and sex hormone binding globulin : determinants of bone turnover and bone mineral density in elderly women. *Journal of Bone and Mineral Research*, 10, 1177–1184.
- Ozdemir, F., Demirbag, D., & Rodoplu, M. (2005). Reproductive factors affecting the bone mineral density in postmenopausal women. *Tohoku Journal of Experimental Medicine*, 205(3), 277–285.
- Pandey, N., Bhola, S., Goldstone, A., Chen, F., Chrzanowski, J., Terranova, C. J., ... Jepsen, K. J. (2009). Interindividual variation in functionally adapted trait sets is established during postnatal growth and predictable based on bone robustness. *Journal of Bone and Mineral Research*, 24(12), 1969–1980.
- Park, J.-H., Song, Y.-M., Sung, J., Lee, K., Kim, Y. S., Kim, T., & Cho, S.-I. (2012). The association between fat and lean mass and bone mineral density: the Healthy Twin Study. *Bone*, 50(4), 1006–1011.
- Peel, N. M., McClure, R. J., & Hendrikz, J. K. (2007). Psychosocial factors associated with fall-related hip fractures. *Age and Ageing*, 36(2), 145–151.
- Pei, L., & Tontonoz, P. (2004). Fat's loss is bone's gain. The Journal of Clinical Investigation, 113(6), 805–806.
- Penny, B., Godber, I. M., & Lawson, N. (2007). Biochemical markers of bone turnov.er. In D. Pearson & C. G. Miller (Eds.), *Clinical Trials in Osteoporosis* (2nd ed., pp. 247–269). London, UK: Springer-Verlag London Limited.

- Perini, T. A., Oliveira, G. L. D., Ornelia, J. D. S., & Oliveira, F. P. D. (2005). Technical error of measurement in anthropometry. *Revista Brasileira de Medicina Do Esporte*, 11, 81–85.
- Perrien, D. S., Achenbach, S. J., Bledsoe, S. E., Walser, B., Suva, L. J., Khosla, S., & Gaddy, D. (2006). Bone turnover across the menopause transition: correlations with inhibins and follicle-stimulating hormone. *The Journal of Clinical Endocrinology and Metabolism*, 91(5), 1848–1854.
- Pines, A., & Berry, E. M. (2007). Exercise in the menopause-an update. *Climacteric*, 10(2), 42–46.
- Pinheiro, M. M., Schuch, N. J., Genaro, P. S., Ciconelli, R. M., Ferraz, M. B., & Martini, L. A. (2009). Nutrient intakes related to osteoporotic fractures in men and women - the Brazilian Osteoporosis Study (BRAZOS). *Nutrition Journal*, 8, 6.
- Pluijm, S. M., Visser, M., Smit, J. H., Popp-Snijders, C., Roos, J. C., & Lips, P. (2001). Determinants of bone mineral density in older men and women: body composition as mediator. *Journal of Bone and Mineral Research*, 16(11), 2142– 2151.
- Poh, B. K., Safiah, M. Y., Tahir, A., Haslinda, M. D. S., Norazlin, N. S., Norimah, A. K., ... Fatimah, S. (2010). Physical activity pattern and energy expenditure of Malaysian adults: findings from the Malaysian Adult Nutrition Survey (MANS). *Malaysian Journal of Nutrition*, 16(1), 13–37.
- Pon, L. W., Noor-Aini, M. Y., Ong, F. B., Adeeb, N., Seri, S. S., Shamsuddin, K., ... Wan, H. W. H. (2006). Diet, nutritional knowledge and health status of urban middle-aged Malaysian women. *Asia Pacific Journal of Clinical Nutrition*, 15(3), 388–399.
- Promislow, J. H. E., Goodman-Gruen, D., Slymen, D. J., & Barrett-Connor, E. (2002). Protein consumption and bone mineral density in the elderly: the Rancho Bernardo Study. *American Journal of Epidemiology*, 155(7), 636–644.
- Puntus, T., Schneider, B., Meran, J., Peterlik, M., & Kudlacek, S. (2011). Influence of age and gender on associations of body mass index with bone mineral density, bone turnover markers and circulating calcium-regulating and bone-active sex hormones. *Bone*, 49(4), 824–849.
- Purohit, V. (2001). Can alcohol promote aromatization of androgens to estrogens? *Alcohol*, 22(2000), 123–127.
- Qu, B., Ma, Y., Yan, M., Wu, H.-H., Fan, L., Liao, D.-F., ... Hong, Z. (2014). The economic burden of fracture patients with osteoporosis in western China. *Osteoporosis International*, 25(7), 1853–1860.
- Radak, T. L. (2004). Caloric restriction and calcium's effect on bone metabolism and body composition in overweight and obese premenopausal women. *Nutrition Reviews*, 62(12), 468–481.
- Recker, R. R., & Deng, H.-W. (2002). Role of genetics in osteoporosis. *Endocrine*, 17(1), 55–66.

- Redman, L. M., Rood, J., Anton, S. D., Champagne, C., Smith, S. R., & Ravussin, E. (2008). Calorie restriction and bone health in young, overweight individuals. *Archives of Internal Medicine*, 168(17), 1859–1866.
- Reid, I. R. (2010). Fat and bone. Archives of Biochemistry and Biophysics, 503(1), 20–27.
- Riancho, J. A. (2010). Genetics of osteoporosis: half-full or half-empty? Clinical Reviews in Bone and Mineral Metabolism, 8(2), 49–50.
- Riedt, C. S., Cifuentes, M., Stahl, T., Chowdhury, H. A., Schlussel, Y., & Shapses, S. A. (2005). Overweight postmenopausal women lose bone with moderate weight reduction and 1 g/day calcium intake. *Journal of Bone and Mineral Research*, 20(3), 455–463.
- Riedt, C. S., Schlussel, Y., Von Thun, N., Ambia-Sobhan, H., Stahl, T., Field, M. P., ... Shapses, S. A. (2007). Premenopausal overweight women do not lose bone during moderate weight loss with adequate or higher calcium intake. *American Journal of Clinical Nutrition*, 85, 972–980.
- Riggs, B. L., Melton, L. J., Robb, R. A., Camp, J. J., Atkinson, E. J., McDaniel, L., ... Khosla, S. (2008). A population-based assessment of rates of bone loss at multiple skeletal sites: evidence for substantial trabecular bone loss in young adult women and men. *Journal of Bone and Mineral Research*, 23(2), 205–214.
- Riggs, B. L., Melton, L. J., Robb, R. A., Camp, J. J., Atkinson, E. J., Peterson, J. M., ... Khosla, S. (2004). Population-based study of age and sex differences in bone volumetric density, size, geometry, and structure at different skeletal sites. *Journal of Bone and Mineral Research*, 19(12), 1945–1954.
- Rizzoli, R., Ammann, P., Chevalley, T., & Bonjour, J.-P. (2001). Protein intake and bone disorders in the elderly. *Joint Bone Spine*, *68*(5), 383–392.
- Rizzoli, R., Bianchi, M. L., Garabédian, M., McKay, H. A., & Moreno, L. A. (2010). Maximizing bone mineral mass gain during growth for the prevention of fractures in the adolescents and the elderly. *Bone*, 46(2), 294–305.
- Robling, A. G., Castillo, A. B., & Turner, C. H. (2006). Biomechanical and molecular regulation of bone remodeling. *Annual Review of Biomedical Engineering*, 8, 455–498.
- Roughead, Z. K. F., Johnson, L. K., Lykken, G. I., & Hunt, J. R. (2003). Controlled high meat diets do not affect calcium retention of indices of bone status in healthy postmenopausal women. *The Journal of Nutrition*, *133*, 1020–1026.
- Roy, D., Swarbrick, C., King, Y., Pye, S., Adams, J., Berry, J., ... O'Neill, T. (2005). Differences in peak bone mass in women of European and South Asian origin can be explained by differences in body size. *Osteoporosis International*, 16(10), 1254–1262.
- Saraví, F. D., & Sayegh, F. (2013). Bone mineral density and body composition of adult premenopausal women with three levels of physical activity. *Journal of Osteoporosis*, 2013, 953271.

- Sato, M., Vietri, J., Flynn, J. A., & Fujiwara, S. (2014). Bone fractures and feeling at risk for osteoporosis among women in Japan: patient characteristics and outcomes in the National Health and Wellness Survey. *Archives of Osteoporosis*, 9, 199.
- Sato, Y., Inose, M., Higuchi, I., Higuchi, F., & Kondo, I. (2002). Changes in the supporting muscles of the fractured hip in elderly women. *Bone*, *30*(1), 325–330.
- Schaffler, M. B., Cheung, W.-Y., Majeska, R., & Kennedy, O. (2014). Osteocytes: master orchestrators of bone. *Calcified Tissue International*, 94, 5–24.
- Schmitt, N. M., Schmitt, J., & Dören, M. (2009). The role of physical activity in the prevention of osteoporosis in postmenopausal women-an update. *Maturitas*, 63(1), 34–38.
- Seeman, E. (2008). Bone quality: the material and structural basis of bone strength. *Journal of Bone and Mineral Metabolism*, 26(1), 1–8.
- Seeman, E. (2008). Structural basis of growth-related gain and age-related loss of bone strength. *Rheumatology*, 47, iv2–8.
- Shao, H. D., Li, G. W., Liu, Y., Qiu, Y. Y., Yao, J. H., & Tang, G. Y. (2014). Contributions of fat mass and fat distribution to hip bone strength in healthy postmenopausal Chinese women. *Journal of Bone and Mineral Research*. doi:10.1007/s00774-014-0613-7
- Shapses, S. A., & Sukumar, D. (2012). Bone metabolism in obesity and weight loss. Annual Review in Nutrition, 32, 287–309.
- Shea, B., Wells, G., Cranney, A., Zytaruk, N., Robinson, V., Griffith, L., ... Guyatt, G. (2002). Meta-analysis of calcium supplementation for the prevention of postmenopausal osteoporosis. *Endocrine Reviews*, 23(4), 552–559.
- Sheng, Z., Xu, K., Ou, Y., Dai, R., Luo, X., Liu, S., ... Liao, E. (2011). Relationship of body composition with prevalence of osteoporosis in central south Chinese postmenopausal women. *Clinical Endocrinology*, 74, 319–324.
- Shin, C. S., Choi, H. J., Kim, M. J., Kim, J. T., Yu, S. H., Koo, B. K., ... Cho, N. H. (2010). Prevalence and risk factors of osteoporosis in Korea: a community-based cohort study with lumbar spine and hip bone mineral density. *Bone*, 47(2), 378– 387.
- Shuster, L. T., Rhodes, D. J., Gostout, B. S., Grossardt, B. R., & Rocca, W. A. (2010). Premature menopause or early menopause: long-term health consequences. *Maturitas*, 65(2), 161–166.
- Sioka, C., Bougias, C., Papadopoulos, A., & Fotopoulos, A. (2007). Is osteoporosis in postmenopausal female patients related to previous pregnancies and/or miscarriages? *Climacteric*, 10, 381–385.
- Sioka, C., Fotopoulos, A., Georgiou, A., Xourgia, X., Papadopoulos, A., & Kalef-Ezra, J. A. (2010). Age at menarche, age at menopause and duration of fertility as risk factors for osteoporosis. *Climacteric*, 13, 63–71.

- Siris, E. S., Brenneman, S. K., Barrett-Connor, E., Miller, P. D., Sajjan, S., Berger, M. L., & Chen, Y.-T. (2006). The effect of age and bone mineral density on the absolute, excess, and relative risk of fracture in postmenopausal women aged 50-99: results from the National Osteoporosis Risk Assessment (NORA). Osteoporosis International, 17(4), 565–574.
- Siris, E. S., Miller, P. D., Barrett-Connor, E., Foulkner, K. G., Wehren, L. E., Abbott, T. A., ... Sherwood, L. M. (2001). Identification and fracture outcomes of undiagnosed low bone mineral density in postmenopausal women: results from the National Osteoporosis Risk Assessment. *Journal of the American Medical Association*, 286(22), 2815–2822.
- Skillington, J., Choy, L., & Derynck, R. (2002). Bone morphogenetic protein and retinoic acid signaling cooperate to induce osteoblast differentiation of preadipocytes. *The Journal of Cell Biology*, 159(1), 135–146.
- Skrzek, A., Kozieł, S., & Ignasiak, Z. (2014). The optimal value of BMI for the lowest risk of osteoporosis in postmenopausal women aged 40-88 years. *HOMO -Journal of Comparative Human Biology*, 65, 232–239.
- Slemenda, C. W., Peacock, M., Hui, S., Zhou, L., & Johnston, C. C. (1997). Reduced rates of skeletal remodeling are associated with increased bone mineral density during the development of peak skeletal mass. *Journal of Bone and Mineral Research*, 12(4), 676–682.
- Sowers, M. R., Jannausch, M., McConnell, D., Little, R., Greendale, G. a, Finkelstein, J. S., ... Ettinger, B. (2006). Hormone predictors of bone mineral density changes during the menopausal transition. *The Journal of Clinical Endocrinology and Metabolism*, 91(4), 1261–1267.
- Streeten, E. A., Ryan, K. A., McBride, D. J., Pollin, T. I., Shuldiner, A. R., & Mitchell, B. D. (2005). The relationship between parity and bone mineral density in women characterized by a homogeneous lifestyle and high parity. *The Journal of Clinical Endocrinology and Metabolism*, 90(8), 4536–4541.
- Sung, J., Song, Y.-M., Stone, J., & Lee, K. (2011). The relationship between bone mineral density and mammographic density in Korean women: the Healthy Twin study. *Breast Cancer Research and Treatment*, 129, 583–591.
- Suzuki, T. (2001). Risk factors for osteoporosis in Asia. Journal of Bone and Mineral Metabolism, 19, 133–141.
- Szulc, P., & Delmas, P. D. (2008). Biochemical markers of bone turnover: potential use in the investigation and management of postmenopausal osteoporosis. *Osteoporosis International*, *19*(12), 1683–1704.
- Takada, H., Washino, K., & Iwata, H. (1997). Risk factors for low bone mineral density among females: the effect of lean body mass. *Preventive Medicine*, *26*, 633–638.
- Takeda, E., Yamamoto, H., Yamanaka-Okumura, H., & Taketani, Y. (2012). Dietary phosphorus in bone health and quality of life. *Nutrition Reviews*, 70(6), 311–321.
- Tanaka, K., Hisada, K., Unno, K., Iijima, M., Unno, A., Tokita, A., ... Yamashiro, Y. (2007). Bone mineral density in children and adolescent girls with anorexia nervosa in Japan. *Pediatrics International*, 49, 637–640.

- Teo, K., Lear, S., Islam, S., Mony, P., Dehghan, M., Li, W., ... Yusuf, S. (2013). Prevalence of a healthy lifestyle among individuals with cardiovascular disease in high-, middle- and low-income countries: the Prospective Urban Rural Epidemiology (PURE) study. *Journal of the American Medical Association*, 309(15), 1613–1621.
- Terracciano, C., Celi, M., Lecce, D., Baldi, J., Rastelli, E., Lena, E., ... Tarantino, U. (2013). Differential features of muscle fiber atrophy in osteoporosis and osteoarthritis. *Osteoporosis International*, 24(3), 1095–1100.
- Thomas, T., & Burguera, B. (2002). Is leptin the link between fat and bone mass? *Journal of Bone and Mineral Research*, 17(9), 1563–1569.
- Thorpe, M. P., & Evans, E. M. (2011). Dietary protein and bone health: harmonizing conflicting theories. *Nutrition Reviews*, 69(4), 215–230.
- Trimpou, P., Bosaeus, I., Bengtsson, B.-A., & Landin-Wilhelmsen, K. (2010). High correlation between quantitative ultrasound and DXA during 7 years of followup. *European Journal of Radiology*, 73(2), 360–364.
- Tsujimoto, M., Chen, P., Miyauchi, A., Sowa, H., & Krege, J. H. (2011). PINP as an aid for monitoring patients treated with teriparatide. *Bone*, 48(4), 798–803.
- Tu, Q., Zhang, J., Dong, L. Q., Saunders, E., Luo, E., Tang, J., & Chen, J. (2011). Adiponectin inhibits osteoclastogenesis and bone resorption via APPL1mediated suppression of Akt1. *The Journal of Biological Chemistry*, 286, 12542– 12553.
- Tucker, K. L., Jugdaohsingh, R., Powell, J. J., Qiao, N., Hannan, M. T., Sripanyakorn, S., ... Kiel, D. P. (2009). Effects of beer, wine, and liquor intakes on bone mineral density in older men and women. *The American Journal of Clinical Nutrition*, 89, 1188–1196.
- Tudor-Locke, C., & McColl, R. S. (2000). Factors related to variation in premenopausal bone mineral status: a health promotion approach. *Osteoporosis International*, *11*(1), 1–24.
- Tuppurainen, M., Kriigerb, H., Saarikoskia, S., Honkanencd, R., & Alhava, E. (1995). The effect of gynecological risk factors on lumbar and femoral bone mineral density in peri- and postmenopausal women. *Maturitas*, 21(2), 137–145.
- Turrell, G., Stanley, L., Looper, M. D., & Oldenburg, B. (2006). Health Inequalities in Australia: Morbidity, health behaviours, risk factors and health service use. Health Inequalities Monitoring Series No. 2. Canberra: Queensland University of Technology and the Australian Institute of Health and Welfare.
- U.S. Department of Health and Human Service. (2004). *Bone Health and Osteoporosis: A Report of the Surgeon General.*
- Uribarri, J., & Calvo, M. S. (2003). Hidden sources of phosphorus in the typical American diet: does it matter in nephrology? *Seminars in Dialysis*, 16(7), 186–188.
- Vasikaran, S., Eastell, R., Bruyère, O., Foldes, A. J., Garnero, P., Griesmacher, A., ... Kanis, J. A. (2011). Markers of bone turnover for the prediction of fracture risk and monitoring of osteoporosis treatment: a need for international reference standards. *Osteoporosis International*, 22(2), 391–420.

- Vestergaard, P., Rejnmark, L., & Mosekilde, L. (2006). Socioeconomic aspects of fractures within universal public healthcare: a nationwide case-control study from Denmark. *Scandinavian Journal of Public Health*, *34*, 371–377.
- Vicente-Rodríguez, G., Ezquerra, J., Mesana, M. I., Fernández-Alvira, J. M., Rey-López, J. P., Casajus, J. A., & Moreno, L. A. (2008). Independent and combined effect of nutrition and exercise on bone mass development. *Journal of Bone and Mineral Metabolism*, 26, 416–424.
- Viljakainen, H. T., Natri, A.-M., Kärkkäinen, M., Huttunen, M. M., Palssa, A., Jakobsen, J., ... Lamberg-Allardt, C. (2006). A positive dose-response effect of vitamin D supplementation on site-specific bone mineral augmentation in adolescent girls: a double-blinded randomized placebo-controlled 1-year intervention. *Journal of Bone and Mineral Research*, 21(6), 836–844.
- Villareal, D. T., Shah, K., Banks, M. R., Sinacore, D. R., & Klein, S. (2008). Effect of weight loss and exercise therapy on bone metabolism and mass in obese older adults: a one-year randomized controlled trial. *Journal of Clinical Endocrinology* and Metabolism, 93, 2181–2187.
- Voort, D. J. M. V. D., Geusens, P. P., & Dinant, G. J. (2001). Risk factors for osteoporosis related to their outcome: fractures. *Osteoporosis International*, 12(12), 630–638.
- Vu, T. T. H., Nguyen, C. K., Nguyen, T. L., Le, B. M., Le, D. N., Bui, T. N., ... Yamamoto, S. (2005). Determining the prevalence of osteoporosis and related factors using quantitative ultrasound in Vietnamese adult women. *American Journal of Epidemiology*, 161(9), 824–830.
- Wagner, H., Melhus, H., Pedersen, N. L., & Michaëlsson, K. (2013). Genetic influence on bone phenotypes and body composition: a Swedish twin Bone and Mineral Metabolism, 31, 681–689.
- Walker, M. D., Babbar, R., Opotowsky, A., McMahon, D. J., Liu, G., & Bilezikian, J. P. (2007). Determinants of bone mineral density in Chinese-American women. *Osteoporosis International*, 18(4), 471–478.
- Wang, J., Thornton, J. C., Ioannidou, E., Soriano, J. M., Gallagher, D., Heymsfield, S. B., ... Allen, L. R. (2005). Four commonly used dual energy X ray absorptiometry scanners do not identically classify subjects for osteopenia or osteoporosis by T-score in four bone regions. *Journal of Clinical Densitometry*, 8(2), 191–198.
- Wang, L., Wang, W., Xu, L., Cheng, X., Ma, Y., Liu, D., ... Wang, Q. (2013). Relation of visceral and subcutaneous adipose tissue to bone mineral density in chinese women. *International Journal of Endocrinology*, 2013, 378632.
- Wang, M. C., Bachrach, L. K., Loan, M. V, Hudes, M., Flegal, K. M., & Crawford, P. B. (2005). The relative contributions of lean tissue mass and fat mass to bone density in young women. *Bone*, 37(4), 474–481.
- Wang, M.-C., & Dixon, L. B. (2006). Socioeconomic influences on bone health in postmenopausal women: findings from NHANES III, 1988-1994. Osteoporosis International, 17(1), 91–98.

- Wang, Q. (2012). Peak bone mass and peak bone strength. In V. R. Preedy (Ed.), Handbook of Growth and Growth Monitoring in Health and Disease (pp. 1317– 1329). New York, NY: Springer New York.
- Wang, Q., Alén, M., Nicholson, P. H. F., Halleen, J. M., Alatalo, S. L., Ohlsson, C., ... Cheng, S. (2006). Differential effects of sex hormones on peri- and endocortical bone surfaces in pubertal girls. *The Journal of Clinical Endocrinology and Metabolism*, 91(1), 277–82.
- Wang, Q., Nicholson, P. H. F., Suuriniemi, M., Lyytikäinen, A., Helkala, E., Alen, M., ... Cheng, S. (2004). Relationship of sex hormones to bone geometric properties and mineral density in early pubertal girls. *The Journal of Clnical Endocrinology and Metabolism*, 89(4), 1698–1703.
- Warensjö, E., Byberg, L., Melhus, H., Gedeborg, R., Mallmin, H., Wolk, A., & Michaëlsson, K. (2011). Dietary calcium intake and risk of fracture and osteoporosis: prospective longitudinal cohort study. *British Medical Journal*, 342, d1473.
- Wat, W. Z. M., Leung, J. Y. Y., Tam, S., & Kung, A. W. C. (2007). Prevalence and impact of vitamin D insufficiency in southern Chinese adults. *Annals of Nutrition* & *Metabolism*, 51, 59–64.
- Waugh, E. J., Lam, M.-A., Hawker, G. A., McGowan, J., Papaioannou, A., Cheung, A. M., ... Jamal, S. A. (2009). Risk factors for low bone mass in healthy 40-60 year old women: a systematic review of the literature. *Osteoporosis International*, 20(1), 1–21.
- Wee, J., Sng, B. Y. J., Shen, L., Lim, C. T., Singh, G., & De, S. D. (2013). The relationship between body mass index and physical activity levels in relation to bone mineral density in premenopausal and postmenopausal women. Archives of Osteoporosis, 8, 162.
- Wehrli, F. W., Song, H. K., Saha, P. K., & Wright, A. C. (2006). Quantitative MRI for the assessment of bone structure and function. *NMR in Biomedicine*, 19, 731–764.
- WHO Commission on Social Determinants of Health. (2008). Closing the gap in a generation: health equity through action on the social determinants of health. Final Report of the Commission on Social Determinants of Health. Geneva, World Health Organization.
- WHO/IASO/IOTF. (2000). The Asia Pacific perspective: redefining obesity and its treatment. Melbourne, Australia: Health Communications Australia.
- Williams, F. M. K., Cherkas, L. F., Spector, T. D., & MacGregor, A. J. (2005). The effect of moderate alcohol consumption on bone mineral density: a study of female twins. *Annals of the Rheumatic Diseases*, 64, 309–310.
- Wilson, R. T., Chase, G. A., Chrischilles, E. A., & Wallace, R. B. (2006). Hip fracture risk among community-dwelling elderly people in the United States: a prospective study of physical, cognitive, and socioeconomic indicators. *American Journal of Public Health*, 96, 1210–1218.
- Winzenberg, T., Shaw, K., Fryer, J., & Jones, G. (2006). Effects of calcium supplementation on bone density in healthy children: meta-analysis of randomised controlled trials. *BMJ (Clinical Research Ed.)*, 333(7572), 775.

- World Health Organization. (2003a). Diet, nutrition and the prevention of chronic diseases. Report of a Joint WHO/FAO Expert Consultation. WHO Technical Report Series No. 916. Geneva, World Health Organization.
- World Health Organization. (2003b). Prevention and management of osteoporosis: report of a WHO scientific group. WHO technical report series 921. Geneva, World Health Organization.
- World Health Organization. (2011). Noncommunicable diseases country profiles 2011.
- Wosje, K. S., & Specker, B. L. (2000). Role of calcium in bone health during childhood. *Nutrition Reviews*, 58(9), 253–268.
- Wu, X. P., Liao, E. Y., Huang, G., Dai, R. C., & Zhang, H. (2003). A comparison study of the reference curves of bone mineral density at different skeletal sites in native Chinese, Japanese, and American Caucasian women. *Calcified Tissue International*, 73, 122–132.
- Wu, X.-Y., Wu, X.-P., Xie, H., Zhang, H., Peng, Y.-Q., Yuan, L.-Q., ... Liao, E.-Y. (2010). Age-related changes in biochemical markers of bone turnover and gonadotropin levels and their relationship among Chinese adult women. *Osteoporosis International*, 21(2), 275–285.
- Yasui, T., Uemura, H., Tomita, J., Miyatani, Y., Yamada, M., Miura, M., & Irahara, M. (2006). Association of serum undercarboxylated osteocalcin with serum estradiol in pre-, peri- and early post-menopausal women. *Journal of Endocrinological Investigation*, 29, 913–918.
- Yates, L. B., Karasik, D., Beck, T. J., Cupples, L. A., & Kiel, D. P. (2007). Hip structural geometry in old and old-old age: similarities and differences between men and women. *Bone*, 41(4), 722–732.
- Yee, Y. S. S., Zaitun, Y., Chan, Y. M., & Norhaizan, M. E. (2013). Association between anthropometric status, dietary intake and physical activity with bone health status among premenopausal Chinese women in Klang Valley, Malaysia. *Malaysian Journal of Nutrition*, 19(3), 293–302.
- Yilmaz, D., Ersoy, B., Bilgin, E., Gümüşer, G., Onur, E., & Pinar, E. D. (2005). Bone mineral density in girls and boys at different pubertal stages: relation with gonadal steroids, bone formation markers, and growth parameters. *Journal of Bone and Mineral Metabolism*, 23, 476–482.
- Yoshimura, N., Muraki, S., Oka, H., Kawaguchi, H., Nakamura, K., & Akune, T. (2011). Biochemical markers of bone turnover as predictors of osteoporosis and osteoporotic fractures in men and women: 10-year follow-up of the Taiji cohort. *Modern Rheumatology*, 21, 608–620.
- Zhao, J., Xia, W., Nie, M., Zheng, X., Wang, Q., Wang, X., ... Xu, L. (2011). The levels of bone turnover markers in Chinese postmenopausal women: Peking Vertebral Fracture Study. *Menopause: The Journal of The North American Menopause Society*, 18(11), 1237–1243.
- Zhao, L.-J., Liu, Y.-J., Liu, P.-Y., Hamilton, J., Recker, R. R., & Deng, H.-W. (2007). Relationship of obesity with osteoporosis. *The Journal of Clinical Endocrinology* & *Metabolism*, 92(5), 1640–1646.

- Zhen, D., Liu, L., Guan, C., Zhao, N., & Tang, X. (2015). High prevalence of vitamin D deficiency among middle-aged and elderly individuals in northwestern China: its relationship to osteoporosis and lifestyle factors. *Bone*, *71*, 1–6.
- Zhu, K., Hunter, M., James, A., Mun, E., & Walsh, J. P. (2015). Associations between body mass index, lean and fat body mass and bone mineral density in middleaged Australians: The Busselton Healthy Ageing Study. *Bone*, 74, 146–152.
- Zillikens, M. C., Uitterlinden, A. G., Leeuwen, J. P. T. M. V, Berends, A. L., Henneman, P., Dijk, K. W. V, ... Rivadeneira, F. (2010). The role of body mass index, insulin, and adiponectin in the relation between fat distribution and bone mineral density. *Calcified Tissue International*, 86, 116–125.

