



UNIVERSITI PUTRA MALAYSIA

***FACTORS ASSOCIATED WITH DISORDERED EATING AMONG
PRIMARY
SCHOOL CHILDREN IN SELANGOR, MALAYSIA***

CHONG LIN SIEW

FPSK(M) 2016 19



**FACTORS ASSOCIATED WITH DISORDERED EATING AMONG PRIMARY
SCHOOL CHILDREN IN SELANGOR, MALAYSIA**

By

CHONG LIN SIEW

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

January 2016

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

FACTORS ASSOCIATED WITH DISORDERED EATING AMONG PRIMARY SCHOOL CHILDREN IN SELANGOR, MALAYSIA

By

CHONG LIN SIEW

January 2016

Chair : Chin Yit Siew, PhD
Faculty : Medicine and Health Sciences

The prevalence of disordered eating is on the rise among children. This study aimed to determine the prevalence of disordered eating and the associations between socio-demographic characteristics, pubertal development, body weight status, psychological factors, socio-cultural factors and behavioral factors with disordered eating among primary school children (aged 10 to 11 years) in Selangor.

A total of 816 children (35.1% males and 64.9% females) from 12 randomly selected primary schools involved in this cross-sectional study. Children completed a set of self-administered questionnaire on disordered eating, socio-demographic characteristics (sex, ethnicity, parental monthly income, parental education level), pubertal development, psychological factors (self-esteem, depression, health specific self-efficacy, body size satisfaction), socio-cultural factors (perceived pressure to lose weight, gain weight and increase muscle tone from parents, peers and media) and behavioral factors (meal skipping behaviors, snacking behaviors, fast food consumption, dietary intake, physical activity level). Their body weight and height were measured and body weight status was categorized based on WHO Growth Reference (2007). Multivariate logistic regression analysis was conducted to determine factors associated with disordered eating among the children.

The prevalence of disordered eating was 30.8% [Males: 32.8% (95% CI: 27.3, 38.8); Females: 29.7% (95% CI: 25.8, 33.9)]. Older age children (10.7; 95% CI: 10.7, 10.8) reported higher prevalence of disordered eating than younger age children (10.5; 95% CI: 10.5, 10.6). Indian (39.9%; 95% CI: 29.3, 51.40) showed the highest prevalence of disordered eating, followed by Malay (33.6%; 95% CI: 29.7, 37.8) and Chinese (17.0%; 95% CI: 11.8, 23.8). Children who were in advanced and post-pubertal stage had the highest prevalence of disordered eating (44.6%; 95% CI: 33.0, 56.7). Those who consumed fast food at least once a week showed the highest prevalence of disordered eating (38.4%; 95% CI: 32.6, 44.5). Socio-cultural pressures to lose weight among children with disordered eating (32.4; 95% CI: 31.0, 33.8) was higher than those without disordered eating (27.6; 95% CI: 26.7, 28.4). No associations were found between sex, parental monthly income, parental education level, body weight status,

self-esteem, depression, health specific self-efficacy, body size satisfaction, meal skipping behaviors, snacking behaviors, energy and macronutrient intakes, energy expenditure and physical activity level with disordered eating of the children.

Factors that associated with disordered eating were being an Indian (OR = 2.048; 95% CI: 1.050, 3.995), perceived pressure to lose weight from parents, peers and media (OR = 1.035; 95% CI: 1.017, 1.053) and fast food consumption at least once a week (OR = 1.540; 95% CI: 1.098, 2.162) after age was controlled.

In conclusion, one third of the children had disordered eating. After controlling for age, factors of disordered eating in both males and females were being an Indian, perceived pressure to lose weight from parents, peers and media and fast food consumption at least once a week. Future ethnic specific intervention programs on the prevention of disordered eating should be taken consideration of age, socio-cultural factors and fast food consumption.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

FAKTOR-FAKTOR BERKAITAN DENGAN GANGGUAN TINGKAH LAKU PEMAKANAN DALAM KALANGAN KANAK-KANAK SEKOLAH RENDAH DI SELANGOR, MALAYSIA

Oleh

CHONG LIN SIEW

Januari 2016

Pengerusi : Chin Yit Siew, PhD
Fakulti : Perubatan dan Sains Kesihatan

Prevalens gangguan tingkah laku pemakanan semakin meningkat di kalangan kanak-kanak. Kajian ini bertujuan untuk menentukan prevalens gangguan tingkah laku pemakanan dan perkaitan antara ciri-ciri sosio-demografi, perkembangan akil baligh, status berat badan, faktor-faktor psikologi, faktor-faktor sosio-budaya dan faktor-faktor tingkah laku dengan gangguan tingkah laku pemakanan dalam kalangan kanak-kanak sekolah rendah berumur 10 - 11 tahun di Selangor.

Sejumlah 816 kanak-kanak (35.1% lelaki dan 64.9% perempuan) dari 12 sekolah rendah yang dipilih secara rawak terlibat dalam kajian ini. Kanak-kanak tersebut menjawab satu set borang soal-selidik tentang gangguan tingkah laku pemakanan, ciri-ciri sosio-demografi (jantina, keturunan, pendapatan bulanan ibu bapa, taraf pendidikan ibu bapa), perkembangan akil baligh, faktor-faktor psikologi (harga diri, kemurungan, efikasi diri khusus untuk kesihatan, kepuasan terhadap saiz tubuh badan), faktor-faktor sosio-budaya (tekanan yang dirasakan untuk menurunkan berat badan, meningkatkan berat badan, meningkatkan tona otot daripada ibu bapa, rakan sebaya dan media) dan faktor-faktor tingkah laku (melangkau waktu makan utama, pengambilan snek, pengambilan makanan segera, pengambilan pemakanan, tahap aktiviti fizikal). Berat badan dan ketinggian mereka telah diukur. Status berat badan telah dikategorikan berdasarkan Rujukan Pertumbuhan WHO (2007). Analisis regresi logistik multivariat telah dijalankan untuk menentukan faktor-faktor yang berkaitan dengan tingkah laku makan yang tidak teratur dalam kalangan kanak-kanak.

Prevalens gangguan tingkah laku pemakanan adalah 30.8% [Lelaki: 32.8% (95% CI: 27.3, 38.8); Perempuan: 29.7% (95% CI: 25.8, 33.9)]. Kanak-kanak yang lebih tua (10.7; 95% CI: 10.7, 10.8) menunjukkan prevalens gangguan tingkah laku pemakanan yang lebih tinggi berbanding dengan kanak-kanak yang lebih muda (10.5; 95% CI: 10.5, 10.6). India (39.9%; 95% CI: 29.3, 51.40) menunjukkan prevalens gangguan tingkah laku pemakanan yang tertinggi, diikuti dengan Melayu (33.6%; 95% CI: 29.7, 37.8) dan Cina (17.0%; 95% CI: 11.8, 23.8). Kanak-kanak yang berada di peringkat akhir dan pasca akil baligh mempunyai prevalens gangguan tingkah laku pemakanan yang

tertinggi (44.6%; 95% CI: 33.0, 56.7). Mereka yang mengamalkan pengambilan makanan segera sekurang-kurangnya seminggu sekali menunjukkan prevalens gangguan tingkah laku pemakanan yang tertinggi (38.4%; 95% CI: 32.6, 44.5). Tekanan sosio-budaya untuk menurunkan berat badan dalam kalangan kanak-kanak yang mempunyai gangguan tingkah laku pemakanan (32.4; 95% CI: 31.0, 33.8) adalah lebih tinggi daripada mereka yang tidak mempunyai gangguan tingkah laku pemakanan (27.6; 95% CI: 26.7, 28.4). Tiada perkaitan didapati antara jantina, pendapatan bulanan ibu bapa, taraf pendidikan ibu bapa, status berat badan, harga diri, kemurungan, keberkesanan diri khusus untuk kesihatan, kepuasan terhadap saiz tubuh badan, tingkah laku melangkaui waktu makan utama, tingkah laku pengambilan snek, pengambilan tenaga dan nutrien, penggunaan tenaga dan tahap aktiviti fizikal dengan gangguan tingkah laku pemakanan dalam kalangan kanak-kanak tersebut.

Faktor-faktor yang berkaitan dengan gangguan tingkah laku pemakanan adalah sebagai seorang India (OR = 2.048; 95% CI: 1.050, 3.995), tekanan yang dirasakan untuk menurunkan berat badan daripada ibubapa, rakan sebaya dan media (OR = 1.035; 95% CI: 1.017, 1.053) dan pengambilan makanan segera sekurang-kurangnya seminggu sekali (OR = 1.540; 95% CI: 1.098, 2.162) selepas umur diselaraskan.

Kesimpulannya, satu pertiga daripada kanak-kanak mempunyai gangguan tingkah laku pemakanan. Faktor-faktor berkaitan dengan gangguan tingkah laku pemakanan dalam kalangan lelaki dan perempuan adalah sebagai seorang India, tekanan yang dirasakan untuk menurunkan berat badan daripada ibubapa, rakan sebaya dan media dan pengambilan makanan segera sekurang-kurangnya seminggu sekali selepas umur diselaraskan. Program intervensi untuk mencegah gangguan tingkah laku pemakanan yang khusus untuk etnik tertentu perlu dijalankan dengan mengambil kira umur, faktor-faktor sosio-budaya dan pengambilan makanan segera.

ACKNOWLEDGEMENTS

My sincerest appreciation must go to my supervisor, Dr. Chin Yit Siew, who took the risk of supervising me even knowing that I was from different background. Many thanks for her brilliance, guidance, advice, patience, and constant care; which I am grateful to have her as the supervisor and it will be a priceless experience that I will never forget. Besides my supervisor, my appreciation and gratitude also goes to the members of supervisory committee, Associate Professor Dr. Mohd Nasir Mohd Taib and Dr. Gan Wan Ying for their guidance on thesis writing, encouragement, and insightful comments.

I am thankful for the scholarship given by the Ministry of Higher Education (MOHE) Malaysia and Universiti Putra Malaysia (UPM). Without this, I will never be able to set my feet on this top university. I would like to thank the financial support of Research University Grant Scheme (RUGS) for this study. I would also like to thank the school authorities, and students for their full cooperation during the study. Sincere thanks also extend to my colleague, Lai Soke Ching for sharing her ideas, knowledge, and invaluable helps throughout the study. My appreciation and gratitude also goes to my friends, Lew Kok Fang, Chai Kong Fei and Yau Chwan Yu for their continuous support and encouragement throughout the study.

My love and thanks to my parents and brothers for their constant support. They have sacrificed a lot to make sure I get the best education possible, which I can never ever repay. Last but not least, to those who have made contributions direct or indirectly and cannot all be named, thank you very much.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Chin Yit Siew, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Mohd Nasir Mohd Taib, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Gan Wan Ying, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

BUJANG KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____

Date: _____

Name and Matric No.: Chong Lin Siew GS35384

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of

Chairman of

Supervisory Dr. Chin Yit Siew

Committee: _____

Signature: _____

Name of

Member of

Supervisory Assoc. Prof. Mohd Nasir Mohd Taib

Committee: _____

Signature: _____

Name of

Member of

Supervisory Dr. Gan Wan Ying

Committee: _____

TABLE OF CONTENTS

	ABSTRACT	Page
	<i>ABSTRAK</i>	i
	ACKNOWLEDGEMENTS	iii
	APPROVAL	v
	DECLARATION	vi
	LIST OF TABLES	viii
	LIST OF FIGURES	xiii
	LIST OF APPENDICES	xv
	LIST OF ABBREVIATIONS	xvi
	GLOSSARY OF TERMS	xvii
CHAPTER		
1	INTRODUCTION	1
	1.1 Background of the Study	1
	1.2 Problem Statement	2
	1.3 Significance of the Study	4
	1.4 Objectives of the Study	4
	1.4.1 General Objective	4
	1.4.2 Specific Objectives	4
	1.5 Null Hypotheses	5
	1.6 Conceptual Framework	15
2	LITERATURE REVIEW	9
	2.1 Growth and Development in Childhood	9
	2.2 Eating Disorders	9
	2.3 Disordered Eating	11
	2.3.1 Definition of Disordered Eating	11
	2.3.2 Assessment of Disordered Eating	11
	2.3.3 Prevalence of Disordered Eating - Worldwide	12
	2.3.4 Prevalence of Disordered Eating - Malaysia	15
	2.3.5 Consequences of Disordered Eating	16
	2.4 Factors Associated with Disordered Eating	17
	2.4.1 Socio-demographic Characteristics and Disordered Eating	17
	2.4.2 Pubertal Development	21
	2.4.3 Body Weight Status	22
	2.4.4 Psychological Factors and Disordered Eating	23
	2.4.5 Socio-Cultural Factors and Disordered Eating	26
	2.4.6 Behavioral Factors and Disordered Eating	28
	2.5 Factors Contributing to Disordered Eating	31

3	METHODOLOGY	33
3.1	Study Design	33
3.2	Study Location and Subjects	33
3.3	Inclusion and exclusion criteria	33
3.4	Sample Size Determination	33
3.5	Sampling	35
3.6	Study Instruments	36
	3.6.1 Anthropometric Measurements	36
	3.6.2 Self-administered Questionnaire	36
	3.6.3 Two-day 24-hour Dietary Recall	41
	3.6.4 Two-day Physical Activity Recall	42
3.7	Data Collection	44
3.8	Data Analysis	44
4	RESULTS	46
4.1	Socio-demographic Characteristics	46
4.2	Disordered Eating	48
4.3	Pubertal Development	51
4.4	Body Weight Status	53
4.5	Psychological Factors	55
	4.5.1 Self-Esteem	55
	4.5.2 Depression	56
	4.5.3 Health Specific Self-Efficacy	57
	4.5.4 Body Size Satisfaction	59
4.6	Socio-Cultural Factors	61
	4.6.1 Perceived Pressure to Lose Weight from Parents, Peers and Media	61
	4.6.2 Perceived Pressure to Gain Weight from Parents, Peers and Media	62
	4.6.3 Perceived Pressure to Increase Muscle Tone from Parents, Peers and Media	63
4.7	Behavioral Factors	65
	4.7.1 Meal Skipping Behaviors	65
	4.7.2 Snacking Behaviors	66
	4.7.3 Frequency of Fast Food Consumption	67
	4.7.4 Energy and Macronutrients Intake	68
	4.7.5 Micronutrients Intake	70
	4.7.6 Energy Expenditure and Physical Activity Level	71
	4.7.7 Energy Balance	72
4.8	Association between Socio-Demographic Characteristics And Disordered Eating	73
4.9	Association between Body Weight Status and Disordered Eating	75
4.10	Association between Psychological Factors and Disordered Eating	75
4.11	Association between Socio-cultural Factors and Disordered Eating	76
4.12	Association between Behavioral Factors and Disordered Eating	77

4.13	Contributions of Socio-demographic Characteristics, Psychological, Socio-Cultural and Behavioral Factors towards Disordered Eating	78
5	DISCUSSION	80
5.1	Prevalence of Disordered Eating	80
5.2	Association between Socio-demographic Characteristics and Disordered Eating	80
5.3	Association between Body Weight Status and Disordered Eating	83
5.4	Association between Psychological Factors and Disordered Eating	84
5.5	Association between Socio-cultural Factors and Disordered Eating	85
5.6	Association between Behavioral Factors and Disordered Eating	86
5.7	Contribution of Socio-demographic Characteristics, Body Weight Status, Psychological, Socio-cultural and Behavioral Factors towards Disordered Eating	88
6	CONCLUSION AND RECOMMENDATIONS	91
6.1	Conclusion	91
6.2	Limitation of the Study	91
6.3	Recommendations	91
	REFERENCES	93
	APPENDICES	109
	BIODATA OF STUDENT	171
	LIST OF PUBLICATIONS	172

LIST OF TABLES

Table	Page
2.1 The prevalence of disordered eating among children using ChEAT or EAT instruments	13
2.2 The prevalence of disordered eating among children in Malaysia	16
3.1 Sample size calculation based on the cross-sectional (one group) estimate: proportion for prevalence study	34
3.2 Sample size calculation based on formula for logistic regression described by Aday and Cornelius (2011).	34
3.3 BMI-for-age (z-score) classification for children aged 5 to 19 years old	36
3.4 Classification of physical activity level	43
4.1 Socio-demographic characteristics of respondents (n=816)	47
4.2 Distribution of respondents by items based on subscale of disordered eating (n=816)	49
4.2 Distribution of respondents by items based on subscale of disordered eating (n=816) (continued)	50
4.3 Distribution of respondents by perceived pubertal timing (n=816)	51
4.4 Distribution of respondents by pubertal development (n=816)	52
4.5 Distribution of respondents by pubertal development (n=816)	53
4.6 Distribution of respondents by body weight, height, BMI, BMI-for-age (z-score) (n = 816)	54
4.7 Distribution of respondents by body weight status (n = 816)	54
4.8 Distribution of respondents by self-esteem (n=816)	56
4.9 Distribution of respondents by depression (n=816)	57
4.10 Distribution of respondents by nutrition self-efficacy (n=816)	58
4.11 Distribution of respondents by physical activity self-efficacy (n=816)	59
4.12 Distribution of respondents by current body size perception and ideal body size perception (n=816)	60
4.13 Distribution of respondents by body size discrepancy score (n =816)	61
4.14 Distribution of respondents by body size satisfaction (n =816)	61
4.15 Distribution of respondents by items based on perceived pressure to lose weight from parents, peers and media (n =816)	62
4.16 Distribution of respondents by items based on perceived pressure to gain weight from parents, peers and media (n =816)	63
4.17 Distribution of respondents by items based on perceived pressure to increase muscle tone from parents, peers and media (n =816)	64
4.18 Distribution of respondents by meal consumption frequency (n =816)	65
4.19 Distribution of respondents by meal skipping behaviors (n=816)	65
4.20 Distribution of respondents by snacking frequency (n =816)	66
4.21 Distribution of respondents by snacking behaviors (n=816)	66
4.22 Distribution of respondents by frequency of fast food consumption (n=816)	67

4.23	Distribution of respondents by fast food consumption category (n=816)	67
4.24	Distribution of respondents by mean energy and macronutrient intakes (n=816)	68
4.25	Distribution of respondents according to energy and macronutrients adequacy (n=816)	69
4.26	Distribution of respondents by under-reporting, acceptable-reporting, and over-reporting of energy intake (n=816)	70
4.27	Mean micronutrient intakes among respondents (n=816)	70
4.28	Distribution of respondents according to micronutrients adequacy (n=816)	71
4.29	Mean energy expenditure and physical activity value of respondents (n=816)	72
4.30	Distribution of respondents by physical activity level (n=816)	72
4.31	Distribution of respondents by energy balance (n=816)	73
4.32	Association between socio-demographic characteristics, pubertal status and disordered eating	74
4.33	Association between BMI-for-age and body weight status with disordered eating	75
4.34	Association between psychological factors and disordered eating	76
4.35	Association between socio-cultural factors and disordered eating	76
4.36	Association between behavioral factors and disordered eating	77
4.37	Multivariate logistic regression of socio-demographic characteristics, psychological, socio-cultural and behavioral factors towards disordered eating (Overall model)	77

LIST OF FIGURES

Figure	Page
1.1 Conceptual framework	6
3.1 Multistage stratified sampling method	35
3.2 Body Figure Rating Scale for male and female	40
4.1 Distribution of respondents by disordered eating (n = 816)	51

LIST OF APPENDICES

Appendix	Page
A Letter of Approval	110
B Information Sheet	114
C Consent forms	121
D Self-Administered Questionnaire	128



LIST OF ABBREVIATIONS

AN	Anorexia Nervosa
BED	Binge Eating Disorder
BMI	Body Mass Index
BMR	Basal Metabolic Rate
BN	Bulimia Nervosa
ChEAT	Children's Eating Attitude Test
DOSM	Department of Statistics Malaysia
DSM-5	Diagnostic and Statistical Manual of Mental Disorders-5
EAT	Eating Attitude Test
EBQ	Eating Behaviors Questionnaire
FAO	Food and Agriculture Organization
IOTF	International Obesity Task Force
IPH	Institute of Public Health
MET	Metabolic Equivalent
MOHR	Ministry of Human Resource Malaysia
NCCFN	National Coordinating Committee on Food and Nutrition of Malaysia
OSFED	Other Specified Feeding Or Eating Disorder
PAL	Physical Activity Level
RMR	Resting Metabolic Rate
RNI	Recommended Nutrient Intakes for Malaysians
SES	Socio-economic Status
SPSS	Statistical Package for Social Sciences
TDEE	Total Daily Energy Expenditure
UNICEF	United Nations Children's Fund
WHO	World Health Organization

GLOSSARY OF TERMS

Disordered eating	Troublesome eating behaviors such as restrictive dieting, bingeing, or purging which occur less frequently or are less severe than those required to meet the full criteria for the diagnosis of an eating disorder (U.S. Department of Health and Human Services, 2005)
Child	A person under the age of 18 years stated in the Children Act 2001 (Aldgate & Stratham, 2001)
Meal skipping	Skipped at least one of the three main meals per day (Fara Wahida, Chin, & Barakatun Nisak, 2012)
Snacking	Snacked between meals at least once per day (Fara Wahida et al., 2012)

CHAPTER 1

INTRODUCTION

1.1 Background of the Study

Disordered eating is defined as troublesome eating behaviors including restrictive dieting, bingeing, or purging, which occur less frequently or are less severe than those required to meet the full criteria for the diagnosis of an eating disorder (U.S. Department of Health and Human Services, 2005). Disordered eating is a recognised predictor of the onset of eating disorders involving serious eating disturbances, such as extreme and restriction of food intake, severe overeating, as well as feelings of distress or extreme concern about body shape or weight (Abebe, Torgersen, Lien, Hafstad, & Soest, 2013; Stice, Davis, Miller, & Marti, 2008). According to The Diagnostic and Statistical Manual of Mental Disorders (5th ed.; DSM-5; American Psychiatric Association [APA], 2013), eating disorders are currently classified into four types; anorexia nervosa, bulimia nervosa, binge eating disorder and other specified feeding or eating disorder (OSFED). In the general population, the prevalence estimates of eating disorders are between 1% to 4% (Lofrano-Prado et al., 2011). Although studies have estimated that eating disorders affect only 0.3 - 1.6% of the children population (Swanson, Crow, Le Grange, Swendsen, & Merikangas, 2011), complications may lead to life-threatening situations.

Disordered eating occurs across all age groups from adults to children. In general, the prevalence of disordered eating among adults ranges from 11.2% to 31.4%, while around 2.7% to 17.1% of university students had disordered eating (Eisenberg, Nicklett, Roeder, & Kirz, 2011; Forrester-Knauss & Zemp Stutz, 2012; Paulson & Rutledge, 2014; Reba-Harrelson et al., 2009; Uzun et al., 2006). Disordered eating is most commonly found in children and increased prevalence of disordered eating has been observed in the children population (Neumark-Sztainer, Wall, Larson, Eisenberg, & Loth, 2011). The prevalence of disordered eating among children in Western countries ranges from 7.6% to 16.7%, namely Ireland (7.6%); United States (12.0%); Kosovo (13.1%) and Greece (16.7%) (Bilali, Galanis, Velonakis, & Katostaras, 2010; McNicholas et al., 2010; Santos, Richards, & Bleckley, 2007). On the contrary, studies in Asian countries have reported that the prevalence of disordered eating varied from 5.1% to 12.7%, including Hong Kong (5.1%); Taiwan (10.5%) and Korea (12.7%) (Lee et al., 2013; Tam, Ng, Yu, & Young, 2007; Wong, Chang, & Tsao, 2014). In general, disordered eating is more prevalent among Western populations than Asian populations, but the prevalence of disordered eating in Asian populations has shown an increasing trend over the last decades (Chisuwa & O'Dea, 2010). Since the industrialization era, the Western countries have emphasized the thin ideal body image as the beauty standard in Western cultures, which may lead to high prevalence of disordered eating (Mallick, Ray, & Mukhopadhyay, 2014). However, in recent decades, the globalization has exposed the Western beauty culture to the Asian countries that increased the prevalence of disordered eating in Asian countries (Yang, Kim, & Yoon, 2010). Based on these findings, although the prevalence estimates of eating disorders is relatively low (0.3% - 1.6%) (Swanson et al., 2011), disordered eating has been reported within a

significant number of children (5.1% - 41.2%) (Musaiger, Al-Mannai, & Al-Lalla, 2014; Tam et al., 2007). This shows that attention is needed in this area.

Disordered eating is associated with various adverse behavioral, psychological, and physical consequences such as increased risk of weight-gain, obesity, and poorer dietary intake (Chang, Lin, & Wong, 2011; Field et al., 2003; Neumark-Sztainer, Wall, Story, & Standish, 2012; Tsai, Chang, Lien, & Wong, 2011). In terms of psychological and mental health consequences, disordered eating is associated with increased fatigue, depression, anxiety and suicidal behavior (Solmi, Hatch, Hotopf, Treasure, & Micali, 2014; Wille, Hölling, Vloet, & Ravens-Sieberer, 2008). In addition, findings has shown that disordered eating reduced the quality of life for children (Wille et al., 2008). Due to the high prevalence of disordered eating during childhood and the negative health implications, there is a need to clearly understand the factors associated with disordered eating among children.

1.2 Problem Statement

There are three broad stages of growth in childhood: early childhood (birth to eight years), middle childhood (eight to twelve years), and adolescence (twelve to eighteen years) (Tomanari, 2011). Previous findings reported that disordered eating started in the age range 10 to 11 years among children (Combs, Pearson, & Smith, 2011; McVey, Tweed, & Blackmore, 2004; Pearson, Combs, & Smith, 2010; Tam et al., 2007; Wong et al., 2014; Yang et al., 2010). Studies from the United States reported that disordered eating behavior was present in males and females at the age of 11 years (Combs et al., 2011; Pearson et al., 2010). A Canadian study published in 2004 reported that females as young as 10 years of age reported disordered eating (McVey et al., 2004). Studies from non-Western countries such as South Korea, Hong Kong and Taiwan also found that disordered eating appeared among children since middle childhood (Tam et al., 2007; Wong et al., 2014; Yang et al., 2010). For instance, study from South Korea indicated that disordered eating is more prevalent among fourth grade students (aged 9-10 years) than seventh grade students (aged 12-13 years). Moreover, in Hong Kong, the youngest case of disordered eating reported was 11 years old (Tam et al., 2007). Therefore, it can be concluded that disordered eating started to emerge during middle childhood, particularly between the age of 10 to 13 years. However, little is known about the disordered eating behavior on children among this age group in the Malaysian context.

There are few studies on disordered eating in Malaysia focusing on young children (8-9 years) and adolescents (13-17 years) (Dan, Mohd Nasir, & Zalilah, 2011; Farah, Mohd Nasir, & Hazizi, 2011; Law, Mohd Nasir, & Abu Saad, 2014; Soo, Zalilah, Mohd Nasir, & Bahaman, 2008; Zalilah & Zaidah, 2005). For instance, a study conducted on 107 Malay female children aged 8 to 9 years in Selangor reported that 38.0% of the Malay females reported disordered eating (Zalilah & Zaidah, 2005). Besides, Soo et al. (2008) found that the prevalence of disordered eating was 35.4% to 36.0% among female adolescents aged between 15 to 17 years in Kelantan. However, the study by Zalilah and Zaidah (2005) was limited to a sample of Malay females at a younger age group (8 to 9 years) while the study by Soo et al. (2008) was focused on a sample of female

adolescents aged 15 to 17 years. There is a lack of published studies on disordered eating in Malaysia which focus on children of various ethnic groups, considering that Malaysia is a multi-ethnic country.

A number of studies have examined the prevalence of disordered eating focused primarily on female children, however, disordered eating is not a problem solely by a single sex (Chang et al., 2011; Evans, Tovee, Boothroyd, & Drewett, 2013; Hamel, Zaitsoff, Taylor, Menna, & Le Grange, 2012; Harrison & Hefner, 2006; Jones, Bennett, Olmsted, Lawson, & Rodin, 2001; McVey et al., 2004; Toro et al., 2006; Tsai et al., 2011). Evidence suggested that disordered eating among males were in increasing trends. In fact, disordered eating is prevalent among both male and female children (Neumark-Sztainer et al., 2011). Evidence from a 10-year longitudinal study on disordered eating behaviors reported that the prevalence of dieting among males had significantly increased from 21.9% to 27.9%; while the prevalence of extreme weight control behaviors increased from 2.1% to 7.3% after 10 years of follow-up from adolescence to young adulthood (Neumark-Sztainer et al., 2011). However, the local studies only focused primarily on female children and adolescents but not on male children and adolescents (Soo et al., 2008; Zalilah & Zaidah, 2005). Limited number of published studies included both sexes (Farah et al., 2011; Law et al., 2014). Farah et al. (2011) reported that both males and females in Pahang reported high prevalence of disordered eating (Males: 28.8%; Females: 26.9%); however there is no significance between sexes. Another local study by Law et al. (2014) reported that 18.3% of males and 20.4% of females in Sarawak had disordered eating. Nevertheless, the difference according to sex in disordered eating was not examined. Thus, there is a need to conduct a study on disordered eating which includes both male and female children and examine the difference in terms of different sexes in Malaysia.

The development of disordered eating is complex and no single factor can explain it. It has been reported that various socio-demographic characteristics, body weight status, psychological, socio-cultural and behavioral factors were found to be associated with disordered eating (Muris, Meesters, van de Blom, & Mayer, 2005; Neumark-Sztainer, Wall, Story, & Sherwood, 2009). Previous findings reported that socio-demographic characteristics (age and pubertal development), body weight status, familial and social factors were associated with disordered eating among female children in Jordan (Mousa, Al-Domi, Mashal, & Jibril, 2010). Furthermore, Evans et al. (2013) reported that psychological factors including depression and body image were correlated with disordered eating among female children in North East England. Besides that, psychological factors and socio-cultural factors were associated with disordered eating among children aged 9-15 years in United States (Field et al., 2008). A recent research by Veses et al. (2014) reported that behavioral factors particularly physical activity level was a significant factor of disordered eating among a sample of Spanish children. However, all these studies focused on single domains and were conducted in Western countries. Factors associated with disordered eating in children from non-Western countries have not been fully investigated. There is still lack of studies that investigate the combination of psychological, body weight status, socio-cultural, and behavioral factors on disordered eating among school children in Malaysia.

The following are the research questions to be addressed in this study: -

1. What is the prevalence of disordered eating among male and female primary school children in Selangor?
2. Do socio-demographic characteristics, body weight status, psychological factors, socio-cultural factors and behavioral factors contribute to disordered eating among primary school children?

1.3 Significance of the Study

At present, there is insufficient information on disordered eating among primary school children in Malaysia. Thus, the findings from this study can provide the baseline information on disordered eating for future research. This study contributes and updates the data on the prevalence of disordered eating among primary school children.

In addition, the present study covers socio-demographic characteristics, body weight status, psychological factors, socio-cultural factors as well as behavioral factors on disordered eating whereas others studies only focused on certain factors on disordered eating. Therefore, this study can provide a more comprehensive picture of the factors associated to disordered eating among primary school children.

Besides that, this study also provides information about factors contributed to disordered eating, particularly among primary school children, which may serve as potential targets for prevention of disordered eating that can be tracked into adulthood. The factors associated to disordered eating among primary school children in this study are useful as reference for developing prevention and intervention programs for successful improvement of children's physical and psychological well-being.

1.4 Objectives of the Study

1.4.1 General Objective

To determine the factors associated with disordered eating among primary school children in Selangor.

1.4.2 Specific Objectives

1. To determine the prevalence of disordered eating among primary school children in Selangor.

2. To determine the socio-demographic characteristics (age, sex, ethnicity, parental education level, parental monthly income, socio-economic status); pubertal development; body weight status; psychological factors (self-esteem, depression, health specific self-efficacy, and body size satisfaction); socio-cultural factors (perceived pressure to lose weight, gain weight and increase muscle tone from parents, peers and media); and behavioral factors (meal skipping and snacking behaviors, frequency of fast food consumption, dietary intake and physical activity) of the primary school children in Selangor.
3. To determine the associations between socio-demographic characteristics (age, sex, ethnicity, parental education level, parental monthly income, socio-economic status); pubertal development; body weight status; psychological factors (self-esteem, depression, health specific self-efficacy, and body size satisfaction); socio-cultural factors (perceived pressure to lose weight, gain weight and increase muscle tone from parents, peers and media); and behavioral factors (meal skipping and snacking behaviors, frequency of fast food consumption, dietary intake and physical activity) with disordered eating among primary school children in Selangor.
4. To determine the contribution of socio-demographic characteristics, pubertal development, body weight status, psychological, socio-cultural and behavioral factors towards disordered eating among primary school children in Selangor.

1.5 Null Hypotheses

1. There are no significant associations between socio-demographic characteristics, pubertal development, body weight status, psychological factors, socio-cultural factors and behavioral factors with disordered eating among primary school children in Selangor.
2. There are no significant contributions of socio-demographic characteristics, pubertal development, body weight status, psychological factors, socio-cultural factors and behavioral factors towards disordered eating among primary school children in Selangor.

1.6 Conceptual Framework

Figure 1.1 shows the dependent and independent variables in this study. The dependent variable in this study is disordered eating. The independent variables in this study are categorized as six sections, including socio-demographic characteristics, pubertal

development, body weight status, psychological factors, socio-cultural factors and behavioral factors.




Figure 1.1. Conceptual framework

Socio-demographic characteristics include age, sex, ethnicity, parental education level, and parental monthly income. Previous studies have shown that socio-demographic characteristics, including sex, age, ethnicity, parental education level, and parental monthly income may serve as potential contributing factors to disordered eating (Downs, DiNallo, Savage, & Davison, 2007; Ho, Tai, Lee, Cheng, & Liow, 2006; Yang et al., 2010). For instance, findings from South Korea reported that younger students (aged 9-10 years) reported higher prevalence of disordered eating than older students (aged 12-13 years) (Yang et al., 2010).

Moreover, numerous studies showed that females were more likely to develop disordered eating as compared to males (Bilali et al., 2010; Ferreiro, Seoane, & Senra, 2012; Hadjigeorgiou, Tornaritis, Savva, Solea, & Kafatos, 2012; Musaiger et al., 2013). In addition, Ho et al. (2006) reported that Malay female children were more likely to develop disordered eating than Chinese female children in Singapore. Previous research found pubertal development increased the risk of disordered eating (Baker, Thornton, Lichtenstein, & Bulik, 2012; McNicholas, Dooley, McNamara, & Lennon, 2012; Mousa et al., 2010). Findings from Ireland reported that female children who are more matured was correlated with disordered eating (McNicholas et al., 2012).

Body weight status is one of the hypothesized risk factor of disordered eating in current study. For instance, findings from Hong Kong conducted by Tam et al. (2007) found that overweight children were at higher risk to have disordered eating as compared to normal weight children. Goldschmidt, Aspen, Sinton, Tanofsky-Kraff, and Wilfley (2008) reported that childhood obesity is associated with an increased risk of disordered eating, weight concern, dieting and bingeing.

Psychological factors comprise of self-esteem, depression, health specific self-efficacy and body size satisfaction. Previous research reported that low self-esteem associated with increased risk of disordered eating (Ata et al., 2007; Fan et al., 2010; Rhea & Thatcher, 2013). Downs et al. (2007) reported that high levels of depression has been recognized as a risk factor for disordered eating among children. In addition, according to Bilali et al. (2010), body dissatisfaction was a risk factor for disordered eating among Greek children.

Socio-cultural factors encompass perceived pressure to lose weight, gain weight and increase muscle tone from parents, peers and media. Ata, Ludden, and Lally (2007) reported that children with disordered eating reported higher levels of pressures from peers and family to lose weight. In EAT Project (Eating Among Teens), Neumark-Sztainer et al. (2009) reported that reading magazines articles regarding weight loss was correlated with increased prevalence of disordered eating among children.

Behavioral factors cover meal skipping behaviors, snacking behaviors, frequency of fast food consumption, dietary intake and physical activity. A study by Chang et al. (2011) on Taiwanese female high school students indicated that caloric intake in students with disordered eating was significantly lower than students without disordered eating. Veses et al. (2014) in their AVENA and AFINOS studies found that

disordered eating was associated with medium and low levels of physical activity among children. Moreover, Stice et al. (2008) reported that meal skipping and obsessive exercise are recognized predictors of disordered eating.

Based on the literature review, the development of disordered eating is multifactorial; socio-demographic characteristics, pubertal development, body weight status, psychological factors, socio-cultural factors and behavioral factors are important contributing factors of disordered eating (Ata et al., 2007; Bilali et al., 2010; Downs et al., 2007; Lai et al., 2013; Lee et al., 2013; Tsai et al., 2011). In short, socio-demographic characteristics, pubertal development, body weight status, psychological factors, socio-cultural factors and behavioral factors were hypothesized as contributors to disordered eating among primary school children.



REFERENCES

- Abebe, D. S., Torgersen, L., Lien, L., Hafstad, G. S., & Soest, T. v. (2013). Predictors of disordered eating in adolescence and young adulthood: A population-based, longitudinal study of females and males in Norway. *International Journal of Behavioral Development*, 38(2), 128-138.
- Aday, L. A., & Cornelius, L. J. (2011). *Designing and Conducting Health Surveys: A Comprehensive Guide*: John Wiley & Sons.
- Ainsworth, B. E., Haskell, W. L., Herrmann, S. D., Meckes, N., Bassett, D. R., Tudor-Locke, C., Greer, J. L., Vezina, J., Whitt-Glover, M. C., & Leon, A. S. (2011). 2011 compendium of physical activities: a second update of codes and MET values. *Medicine and Science in Sports & Exercise*, 43(8), 1575-1581.
- Aldgate, J., & Stratham, J. (2001). *The Children Act Now: Messages from Research*: London, UK: Stationery Office.
- Anderson, B., Lyon-Callos, S., Fussman, C., Imes, G., & Rafferty, A. P. (2011). Peer Reviewed: Fast-food consumption and obesity among Michigan adults. *Preventing Chronic Disease*, 8(4), 1-11.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Ata, R. N., Ludden, A. B., & Lally, M. M. (2007). The effects of gender and family, friend, and media influences on eating behaviors and body image during adolescence. *Journal of Youth and Adolescence*, 36(8), 1024-1037.
- Austin, S. B., Ziyadeh, N. J., Forman, S., Prokop, L. A., Keliher, A., & Jacobs, D. (2008). Peer reviewed: screening high school students for eating disorders: results of a national initiative. *Preventing Chronic Disease*, 5(4), 1-9.
- Babio, N., Canals, J., Pietrobelli, A., Pérez, S., & Arija, V. (2009). A two-phase population study: relationships between overweight, body composition and risk of eating disorders. *Nutrición Hospitalaria*, 24(4), 485-491.
- Baharudin, A., Zainuddin, A. A., Manickam, M. A., Ambak, R., Ahmad, M. H., Naidu, B. M., Cheong, S. M., Ying, C. Y., Saad, H. A., & Ahmad, N. A. (2014). Factors associated with physical inactivity among school-going adolescents: data from the Malaysian school-based nutrition survey 2012. *Asia-Pacific Journal of Public Health*, 26(5 Suppl), 27S-35S.
- Baker, J. H., Thornton, L. M., Lichtenstein, P., & Bulik, C. M. (2012). Pubertal development predicts eating behaviors in adolescence. *International Journal of Eating Disorders*, 45(7), 819-826.
- Berkman, N. D., Lohr, K. N., & Bulik, C. M. (2007). Outcomes of eating disorders: a systematic review of the literature. *International Journal of Eating Disorders*, 40(4), 293-309.

- Berman, E. S. (2006). The relationship between eating self-efficacy and eating disorder symptoms in a non-clinical sample. *Eating Behaviors*, 7(1), 79-90.
- Bilali, A., Galanis, P., Velonakis, E., & Katostaras, T. (2010). Factors associated with abnormal eating attitudes among Greek adolescents. *Journal of Nutrition Education and Behavior* 42(5), 292-298.
- Boon, T., & Sedek, R. (2012). Association between snacking patterns, energy and nutrient intakes, and body mass index among school adolescents in Kuala Lumpur. *American Journal of Food and Nutrition*, 2(3), 69-77.
- Brausch, A. M., & Gutierrez, P. M. (2009). The Role of Body Image and Disordered Eating as Risk Factors for Depression and Suicidal Ideation in Adolescents. *Suicide and Life-Threatening Behavior*, 39(1), 58-71.
- Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Reports*, 100(2), 126-131.
- Chang, Y. J., Lin, W., & Wong, Y. (2011). Survey on eating disorder-related thoughts, behaviors, and their relationship with food intake and nutritional status in female high school students in Taiwan. *Journal of the American College of Nutrition*, 30(1), 39-48.
- Chin, Y. S., & Mohd Nasir, M. T. (2009). Eating behaviors among female adolescents in Kuantan district, Pahang, Malaysia. *Pakistan Journal of Nutrition*, 8(4), 425-432.
- Cohen, M. L., Tanofsky-Kraff, M., Young-Hyman, D., & Yanovski, J. A. (2005). Weight and its relationship to adolescent perceptions of their providers (WRAP): a qualitative and quantitative assessment of teen weight-related preferences and concerns. *Journal of Adolescent Health*, 37(2), 163.e169-163.e116.
- Collins, M. E. (1991). Body figure perceptions and preferences among preadolescent children. *International Journal of Eating Disorders*, 10(2), 199-208.
- Combs, J. L., Pearson, C. M., & Smith, G. T. (2011). A risk model for preadolescent disordered eating. *International Journal of Eating Disorders*, 44(7), 596-604.
- Combs, J. L., Pearson, C. M., Zapolski, T. C., & Smith, G. T. (2013). Preadolescent disordered eating predicts subsequent eating dysfunction. *Journal of Pediatric Psychology*, 38(1), 41-49.
- Courtney, E. A., Gamboz, J., & Johnson, J. G. (2008). Problematic eating behaviors in adolescents with low self-esteem and elevated depressive symptoms. *Eating Behaviors*, 9(4), 408-414.

- Croll, J., Neumark-Sztainer, D., Story, M., & Ireland, M. (2002). Prevalence and risk and protective factors related to disordered eating behaviors among adolescents: relationship to gender and ethnicity. *Journal of Adolescent Health, 31*(2), 166-175.
- Cynthia, J., Zalilah, M., & Lim, M. (2013). Relationship between family meals away from home and nutritional status of adolescents. *Malaysian Journal of Nutrition, 19*(1), 25-35.
- da Costa, N. F., Schtscherbyna, A., Soares, E. A., & Ribeiro, B. G. (2013). Disordered eating among adolescent female swimmers: dietary, biochemical, and body composition factors. *Nutrition (Burbank, Los Angeles County, Calif.), 29*(1), 172-177.
- Dan, S. P., Mohd Nasir, M. T., & Zalilah, M. (2011). Determination of factors associated with physical activity levels among adolescents attending school in Kuantan, Malaysia. *Malaysian Journal of Nutrition, 17*(2), 175-187.
- Department of Statistics Malaysia. (2010). Population distribution and basic demographic characteristic, 2010. Retrieved from http://www.statistics.gov.my/portal/download_Population/files/population/05Jadual_Mukim_negeri/Mukim_Selangor.pdf
- de Souza Ferreira, J. E., & da Veiga, G. V. (2008). Eating disorder risk behavior in Brazilian adolescents from low socio-economic level. *Appetite, 51*(2), 249-255.
- DeLeel, M. L., Hughes, T. L., Miller, J. A., Hipwell, A., & Theodore, L. A. (2009). Prevalence of eating disturbance and body image dissatisfaction in young girls: An examination of the variance across racial and socioeconomic groups. *Psychology in the Schools, 46*(8), 767-775.
- Denney-Wilson, E., Crawford, D., Dobbins, T., Hardy, L., & Okely, A. D. (2009). Influences on consumption of soft drinks and fast foods in adolescents. *Asia Pacific Journal of Clinical Nutrition, 18*(3), 447-452.
- Downs, D. S., DiNallo, J. M., Savage, J. S., & Davison, K. K. (2007). Determinants of eating attitudes among overweight and non-overweight adolescents. *Journal of Adolescent Health, 41*(2), 138-145.
- Eccles, J. S. (1999). The development of children ages 6 to 14. *The future of children, 30*-44.
- Eisenberg, D., Nicklett, E. J., Roeder, K., & Kirz, N. E. (2011). Eating disorder symptoms among college students: Prevalence, persistence, correlates, and treatment-seeking. *Journal of American College Health, 59*(8), 700-707.
- Eisenberg, M. E., & Neumark-Sztainer, D. (2010). Friends' Dieting and Disordered Eating Behaviors Among Adolescents Five Years Later: Findings From Project EAT. *Journal of Adolescent Health, 47*(1), 67-73.

- EPU. (2012). *Brief Household Income & Poverty Statistics. Statistics Newsletter 2009 & 2012*. Putrajaya. Economic Planning Unit, Prime Minister's Department Malaysia.
- Evans, E. H., Tovee, M. J., Boothroyd, L. G., & Drewett, R. F. (2013). Body dissatisfaction and disordered eating attitudes in 7- to 11-year-old girls: testing a sociocultural model. *Body Image, 10*(1), 8-15.
- Fan, Y., Li, Y., Liu, A., Hu, X., Ma, G., & Xu, G. (2010). Associations between body mass index, weight control concerns and behaviors, and eating disorder symptoms among non-clinical Chinese adolescents. *BMC Public Health, 10*(1), 314.
- Fairburn, C. G., & Beglin, S. J. (1994). Assessment of eating disorders: interview or self-report questionnaire? *International Journal of Eating Disorders, 16*(4), 363-370.
- Food and Agriculture Organization/World Health Organization/United Nations University (FAO/WHO/UNU). (1985). *Energy and Protein Requirements Report of a Joint FAO/WHO/UNU Expert Consultation. Technical Report Series No. 724*. Geneva: World Health Organization.
- Food and Agriculture Organization/World Health Organization/United Nations University (FAO/WHO/UNU). (2004). Human energy requirements: Report of a joint FAO/WHO/UNU Expert Consultation.
- Food and Agriculture Organization/World Health Organization/United Nations University (FAO/WHO/UNU). (2005). Human energy requirements: Report of a joint FAO/WHO/UNU Expert Consultation. *Food and Nutrition Bulletin, 26*, 166.
- Fara Wahida, R., Chin, Y. S., & Barakatun Nisak, M. Y. (2012). Obesity-related behaviors of Malaysian adolescents: a sample from Kajang district of Selangor state. *Nutrition Research and Practice, 6*(5), 458-465.
- Farah, W. Z., Mohd Nasir, M. T., & Hazizi, A. (2011). Physical activity, eating behaviour and body image perception among young adolescents in Kuantan, Pahang, Malaysia. *Malaysian Journal of Nutrition, 17*(3), 325-336.
- Ferreiro, F., Seoane, G., & Senra, C. (2012). Gender-related risk and protective factors for depressive symptoms and disordered eating in adolescence: a 4-year longitudinal study. *Journal of Youth and Adolescence, 41*(5), 607-622.
- Field, A. E., Austin, S., Taylor, C., Malspeis, S., Rosner, B., Rockett, H. R., Gillman, M. W., & Colditz, G. A. (2003). Relation between dieting and weight change among preadolescents and adolescents. *Pediatrics, 112*(4), 900-906.
- Field, A. E., Javaras, K. M., Aneja, P., Kitos, N., Camargo, C. A., Taylor, B., & Laird, N. M. (2008). Family, peer, and media predictors of becoming eating disordered. *Archives of Pediatrics & Adolescent Medicine, 162*(6), 574-579.

- Field, A. E., Sonneville, K. R., Micali, N., Crosby, R. D., Swanson, S. A., Laird, N. M., Treasure, J., Solmi, F., & Horton, N. J. (2012). Prospective association of common eating disorders and adverse outcomes. *Pediatrics*, 130(2), e289-e295.
- First Data Bank. (2003). Nutritionist Pro™ Diet Analysis software. United States: Axxya System.
- Fitzgerald, A., Heary, C., Kelly, C., Nixon, E., & Shevlin, M. (2013). Self-efficacy for healthy eating and peer support for unhealthy eating are associated with adolescents' food intake patterns. *Appetite*, 63(0), 48-58.
- Fleischhacker, S. E., Evenson, K. R., Rodriguez, D. A., & Ammerman, A. S. (2011). A systematic review of fast food access studies. *Obesity Reviews*, 12(5), e460-e471.
- Follansbee-Junger, K., Janicke, D. M., & Sallinen, B. J. (2010). The influence of a behavioral weight management program on disordered eating attitudes and behaviors in children with overweight. *Journal of the American Dietetic Association*, 110(11), 1653-1659.
- Fontenot, B. (2011, 9 August 2011). When Eating Becomes Disordered. Retrieved 09 January, 2015, from <http://www.thedoctorwillseeyounow.com/content/dieting/art3393.html>
- Forrester-Knauss, C., & Zemp Stutz, E. (2012). Gender differences in disordered eating and weight dissatisfaction in Swiss adults: Which factors matter? *BMC Public Health*, 12(1), 809-817.
- Garner, D. M. (1991). EDI-2. *Eating disorder inventory-2. Professional manual*.
- Garner, D. M. (2004). *The Eating Disorder Inventory-3: Psychological Assessment Resources*.
- Garner, D. M., & Olmsted, M. P. (1984). *Manual for eating disorder inventory (EDI): Psychological Assessment Resources, Incorporated*.
- Garner, D. M., Olmsted, M. P., Bohr, Y., & Garfinkel, P. E. (1982). The eating attitudes test: psychometric features and clinical correlates. *Psychological Medicine*, 12(04), 871-878.
- Goldschmidt, A. B., Aspen, V. P., Sinton, M. M., Tanofsky-Kraff, M., & Wilfley, D. E. (2008). Disordered eating attitudes and behaviors in overweight youth. *Obesity (Silver Spring)*, 16(2), 257-264.
- Goñi, A., & Rodríguez, A. (2007). Variables associated with the risk of eating disorders in adolescence. *Salud Mental*, 30(4), 16-23.

- Gowey, M. A., Lim, C. S., Clifford, L. M., & Janicke, D. M. (2014). Disordered Eating and Health-Related Quality of Life in Overweight and Obese Children. *Journal of Pediatric Psychology*, 39(5), 552-561.
- Grucza, R. A., Przybeck, T. R., & Cloninger, C. R. (2007). Prevalence and correlates of binge eating disorder in a community sample. *Comprehensive Psychiatry*, 48(2), 124-131.
- Hadjigeorgiou, C., Tornaritis, M., Savva, S., Solea, A., & Kafatos, A. (2012). Secular trends in eating attitudes and behaviours in children and adolescents aged 10-18 years in Cyprus: A 6-year follow-up, school-based study. *Public Health*, 126(8), 690-694.
- Haines, J., Gillman, M. W., Rifas-Shiman, S., Field, A. E., & Austin, S. B. (2009). Family dinner and disordered eating behaviors in a large cohort of adolescents. *Eating Disorders*, 18(1), 10-24.
- Haines, J., Kleinman, K. P., Rifas-Shiman, S. L., Field, A. E., & Austin, S. B. (2010). Examination of shared risk and protective factors for overweight and disordered eating among adolescents. *Archives of Pediatrics & Adolescent Medicine*, 164(4), 336-343.
- Hair, J. F., Black, W. C., Babin, B. J., Anderson, R. E., & Tatham, R. L. (2006). *Multivariate data analysis*. Upper Saddle River, NJ [etc.]: Pearson Prentice Hall.
- Hamel, A. E., Zaitsoff, S. L., Taylor, A., Menna, R., & Le Grange, D. (2012). Body-related social comparison and disordered eating among adolescent females with an eating disorder, depressive disorder, and healthy controls. *Nutrients*, 4(9), 1260-1272.
- Harrison, K., & Hefner, V. (2006). Media Exposure, Current and Future Body Ideals, and Disordered Eating Among Preadolescent Girls: A Longitudinal Panel Study. *Journal of Youth and Adolescence*, 35(2), 146-156.
- Hautala, L. A., Junnila, J., Helenius, H., Väänänen, A.-M., Liuksila, P.-R., Räihä, H., Välimäki, M., & Saarijärvi, S. (2008). Towards understanding gender differences in disordered eating among adolescents. *Journal of Clinical Nursing*, 17(13), 1803-1813.
- Herpertz-Dahlmann, B., Dempfle, A., Konrad, K., Klasen, F., & Ravens-Sieberer, U. (2014). Eating disorder symptoms do not just disappear: the implications of adolescent eating-disordered behaviour for body weight and mental health in young adulthood. *European Child and Adolescent Psychiatry*, 1-10.
- Ho, T. F., Tai, B. C., Lee, E. L., Cheng, S., & Liow, P. H. (2006). Prevalence and profile of females at risk of eating disorders in Singapore. *Singapore Medical Journal*, 47(6), 499-503.

- Hutchinson, D. M., & Rapee, R. M. (2007). Do friends share similar body image and eating problems? The role of social networks and peer influences in early adolescence. *Behaviour Research and Therapy*, 45(7), 1557-1577.
- Ismail, M. N. (2002). The nutrition and health transition in Malaysia. *Public Health Nutrition*, 5(1A), 191-195.
- Jacobi, C., Hayward, C., de Zwaan, M., Kraemer, H. C., & Agras, W. S. (2004). Coming to terms with risk factors for eating disorders: application of risk terminology and suggestions for a general taxonomy. *Psychological Bulletin*, 130(1), 19-65.
- Johnson, J. G., Cohen, P., Kasen, S., & Brook, J. S. (2014). Childhood adversities associated with risk for eating disorders or weight problems during adolescence or early adulthood. *American Journal of Psychiatry*, 39(4), 394-400.
- Jones, J. M., Bennett, S., Olmsted, M. P., Lawson, M. L., & Rodin, G. (2001). Disordered eating attitudes and behaviours in teenaged girls: a school-based study. *Canadian Medical Association Journal*, 165(5), 547-552.
- Kadriu, F., Kelpi, M., & Kalyva, E. (2014). Eating-disordered Behaviours in Kosovo School-based Population: Potential Risk Factors. *Procedia - Social and Behavioral Sciences*, 114(0), 382-387.
- Kaluski, D. N., Natamba, B. K., Goldsmith, R., Shimony, T., & Berry, E. M. (2008). Determinants of Disordered Eating Behaviors among Israeli Adolescent Girls. *Eating Disorders*, 16(2), 146-159.
- Kandel, D. B., & Davies, M. (1982). Epidemiology of depressive mood in adolescents: an empirical study. *Archives of General Psychiatry*, 39(10), 1205-1212.
- Kaushik, J. S., Narang, M., & Parakh, A. (2011). Fast food consumption in children. *Indian Pediatrics*, 48(2), 97-101.
- Kerr, M. A., Rennie, K. L., McCaffrey, T. A., Wallace, J. M., Hannon-Fletcher, M. P., & Livingstone, M. B. (2009). Snacking patterns among adolescents: a comparison of type, frequency and portion size between Britain in 1997 and Northern Ireland in 2005. *British Journal of Nutrition*, 101(1), 122-131.
- Khor, G. L., Cobiac, L., & Skrzypiec, G. (2002). Gender Differences in Eating Behavior and Social Self Concept among Malaysian University Students. *Malaysian Journal of Nutrition*, 8(1), 75-98.
- Khor, G. L., Zalilah, M., Phan, Y. Y., Ang, M., Maznah, B., & Norimah, A. K. (2009). Perceptions of body image among Malaysian male and female adolescents. *Singapore Medical Journal*, 50(3), 303-311.
- Killen, J. D., Taylor, C. B., Hayward, C., Haydel, K. F., Wilson, D. M., Hammer, L., Kraemer, H., Blair-Greiner, A., & Strachowski, D. (1996). Weight concerns

- influence the development of eating disorders: a 4-year prospective study. *Journal of Consulting and Clinical Psychology*, 64(5), 936-940.
- Kim, H.-Y., Lee, N.-R., Lee, J.-S., Choi, Y.-S., Kwak, T.-K., Chung, H. R., Kwon, S., Choi, Y.-J., Lee, S.-K., & Kang, M.-H. (2012). Meal skipping relates to food choice, understanding of nutrition labeling, and prevalence of obesity in Korean fifth grade children. *Nutrition Research and Practice*, 6(4), 328-333.
- Kluck, A. S. (2010). Family influence on disordered eating: The role of body image dissatisfaction. *Body Image*, 7(1), 8-14.
- Klump, K. L. (2013). Puberty as a critical risk period for eating disorders: A review of human and animal studies. *Hormones and Behavior*, 64(2), 399-410.
- Koletzko, B., & Toschke, A. M. (2010). Meal Patterns and Frequencies: Do They Affect Body Weight in Children and Adolescents? *Critical Reviews in Food Science and Nutrition*, 50(2), 100-105.
- Lai, C. M., Mak, K. K., Pang, J. S., Fong, S. S., Ho, R. C., & Guldán, G. S. (2013). The associations of sociocultural attitudes towards appearance with body dissatisfaction and eating behaviors in Hong Kong adolescents. *Eating Behaviors*, 14(3), 320-324.
- Larson, N. I., Neumark-Sztainer, D., & Story, M. (2009). Weight Control Behaviors and Dietary Intake among Adolescents and Young Adults: Longitudinal Findings from Project EAT. *Journal of the American Dietetic Association*, 109(11), 1869-1877.
- Law, L. S., Mohd Nasir, M. T., & Abu Saad, H. (2014). Factors associated with physical activity level among adolescents in Sarawak, Malaysia. *Journal of Physical Activity, Sports and Exercise* 2(1): 07-14.
- Lee, H. J., Park, S., Kim, C. I., Choi, D. W., Lee, J. S., Oh, S. M., Cho, E., Park, H. K., Kwon, K. I., & Oh, S. W. (2013). The association between disturbed eating behavior and socioeconomic status: the Online Korean Adolescent Panel Survey (OnKAPS). *PLoS ONE*, 8(3), e57880.
- Liechty, J. M., & Lee, M. J. (2013). Longitudinal predictors of dieting and disordered eating among young adults in the U.S. *International Journal of Eating Disorders*, 46(8), 790-800.
- Loewenthal, K. M. (2001). *An introduction to psychological tests and scales*: Psychology Press.
- Lofrano-Prado, M. C., Prado, W. L. d., de Piano, A., Tock, L., Caranti, D. A., Nascimento, C. M. O. d., Oyama, L. M., Tufik, S., de Mello, M. T., & Dâmaso, A. R. (2011). Eating disorders in adolescents: Correlations between symptoms and central control of eating behavior. *Eating Behaviors*, 12(1), 78-82.

- López-Guimerà, G., Levine, M. P., Sánchez-carracedo, D., & Fauquet, J. (2010). Influence of Mass Media on Body Image and Eating Disordered Attitudes and Behaviors in Females: A Review of Effects and Processes. *Media Psychology*, 13(4), 387-416.
- Loth, K. A., MacLehose, R., Bucchianeri, M., Crow, S., & Neumark-Sztainer, D. (2014). Predictors of Dieting and Disordered Eating Behaviors From Adolescence to Young Adulthood. *Journal of Adolescent Health*, 55(5), 705-712.
- Luszczynska, A., Scholz, U., & Schwarzer, R. (2005). The general self-efficacy scale: Multicultural validation studies. *Journal of Psychology*, 139(5), 439-457.
- Mallick, N., Ray, S., & Mukhopadhyay, S. (2014). Eating Behaviours and Body Weight Concerns among Adolescent Girls. *Advances in Public Health*, 2014.
- Maloney, M. J., McGuire, J. B., & Daniels, S. R. (1988). Reliability Testing of a Children's Version of the Eating Attitude Test. *Journal of the American Academy of Child and Adolescent Psychiatry*, 27(5), 541-543.
- Mayer, B., Muris, P., Meesters, C., & Zimmermann-van Beuningen, R. (2009). Brief report: Direct and indirect relations of risk factors with eating behavior problems in late adolescent females. *Journal of Adolescence*, 32(3), 741-745.
- McCabe, M., Ricciardelli, L., Yeow, J., Daliza, N., & Hapidzal, N. F. (2009). Sociocultural influences on body dissatisfaction and body change behaviors among Malaysian adolescents. *Body Image*, 6(2), 121-128.
- McCabe, M. P., & Ricciardelli, L. A. (2001). The structure of the perceived sociocultural influences on body image and body change questionnaire. *International Journal of Behavioral Medicine*, 8(1), 19-41.
- McNicholas, F., Dooley, B., Keogh, L., Lydon, A., Lennon, R., Ahern, S., Coyle, C., Whelan, A., & Donoghue, L. O. (2010). Eating problems in Irish children and adolescence – EPICA. *Irish Journal of Psychological Medicine*, 27(04), 172-178.
- McNicholas, F., Dooley, B., McNamara, N., & Lennon, R. (2012). The impact of self-reported pubertal status and pubertal timing on disordered eating in Irish adolescents. *European Eating Disorders Review*, 20(5), 355-362.
- McNicholas, F., Lydon, A., Lennon, R., & Dooley, B. (2009). Eating concerns and media influences in an Irish adolescent context. *European Eating Disorders Review*, 17(3), 208-213.
- McVey, G., Tweed, S., & Blackmore, E. (2004). Dieting among preadolescent and young adolescent females. *Canadian Medical Association Journal*, 170(10), 1559-1561.

- Meyer, T. A., & Gast, J. (2008). The Effects of Peer Influence on Disordered Eating Behavior. *The Journal of School Nursing*, 24(1), 36-42.
- Micali, N., Ploubidis, G., De Stavola, B., Simonoff, E., & Treasure, J. (2014). Frequency and Patterns of Eating Disorder Symptoms in Early Adolescence. *Journal of Adolescent Health*, 54(5), 574-581.
- Ministry of Human Resources Malaysia. (2010). *Malaysia standard classification of occupations 2008* (3rd ed.). Putrajaya: Human Resource Policy Division, Ministry of Human Resources
- Molnár, D., Jeges, S., Erhardt, E., & Schutz, Y. (1995). Measured and predicted resting metabolic rate in obese and nonobese adolescents. *The Journal of Pediatrics*, 127(4), 571-577.
- Monir, Z. M., Khalifa, A. G., Hassaballa, F., Tawfeek, S., Abdelmonem, M., & Abu, M. (2010). Eating behavior and problems in Egyptian adolescents; relation to dietary intake. *Journal of American Science*, 6(12), 1145-1159.
- Mousa, T. Y., Al-Domi, H. A., Mashal, R. H., & Jibril, M. A.-K. (2010). Eating disturbances among adolescent schoolgirls in Jordan. *Appetite*, 54(1), 196-201.
- Moy, F. M., Ying, G. C., & Kassim, S. Z. M. (2006). Eating patterns of school children and adolescents in Kuala Lumpur. *Malaysian Journal of Nutrition*, 12(1), 1-10.
- Muris, P., Meesters, C., van de Blom, W., & Mayer, B. (2005). Biological, psychological, and sociocultural correlates of body change strategies and eating problems in adolescent boys and girls. *Eating Behaviors*, 6(1), 11-22.
- Murrin, C., McNicholas, F., Keogh, L., Shiely, F., Corrigan, H., Gabhainn, S. N., & Kelleher, C. C. (2007). Correlates of increased risk of eating disorders in Irish school children. *Irish Medical Journal*, 100(8), suppl 40-42.
- Musaiger, A. O., Al-Mannai, M., & Al-Lalla, O. (2014). Risk of disordered eating attitudes among male adolescents in five Emirates of the United Arab Emirates. *International Journal of Eating Disorders*, 47(8), 898-900.
- Musaiger, A. O., Al-Mannai, M., Tayyem, R., Al-Lalla, O., Ali, E. Y., Kalam, F., Benhamed, M. M., Saghir, S., Halahleh, I., Djoudi, Z., & Chirane, M. (2013). Risk of disordered eating attitudes among adolescents in seven Arab countries by gender and obesity: a cross-cultural study. *Appetite*, 60(1), 162-167.
- National Coordinating Committee on Food Nutrition [NCCFN] (2005). Recommended Nutrient Intakes for Malaysia (2005): Putrajaya: Ministry of Health, Malaysia.
- Nattiv, A., Loucks, A. B., Manore, M. M., Sanborn, C. F., Sundgot-Borgen, J., & Warren, M. P. (2007). American College of Sports Medicine position stand. The female athlete triad. *Medicine and Science in Sports & Exercise*, 39(10), 1867-1882.

- Neumark-Sztainer, D. (1996). School - based programs for preventing eating disturbances. *Journal of School Health*, 66(2), 64-71.
- Neumark-Sztainer, D., Levine, M. P., Paxton, S. J., Smolak, L., Piran, N., & Wertheim, E. H. (2006a). Prevention of body dissatisfaction and disordered eating: What next? *Eating Disorders: The Journal of Treatment and Prevention*, 14(4), 265-285.
- Neumark-Sztainer, D., Wall, M., Guo, J., Story, M., Haines, J., & Eisenberg, M. (2006b). Obesity, disordered eating, and eating disorders in a longitudinal study of adolescents: how do dieters fare 5 years later? *Journal of the American Dietetic Association*, 106(4), 559-568.
- Neumark-Sztainer, D., Wall, M., Haines, J., Story, M., & Eisenberg, M. (2007a). Why does dieting predict weight gain in adolescents? Findings from project EAT-II: a 5-year longitudinal study. *Journal of the American Dietetic Association*, 107(3), 448-455.
- Neumark-Sztainer, D., Wall, M., Haines, J., Story, M., Sherwood, N. E., & van den Berg, P. (2007b). Shared risk and protective factors for overweight and disordered eating in adolescents. *American Journal of Preventive Medicine*, 33(5), 359-369.
- Neumark-Sztainer, D., Wall, M., Larson, N. I., Eisenberg, M. E., & Loth, K. (2011). Dieting and disordered eating behaviors from adolescence to young adulthood: findings from a 10-year longitudinal study. *Journal of the American Dietetic Association*, 111(7), 1004-1011.
- Neumark-Sztainer, D., Wall, M., Story, M., & Sherwood, N. E. (2009). Five-year longitudinal predictive factors for disordered eating in a population-based sample of overweight adolescents: implications for prevention and treatment. *International Journal of Eating Disorders*, 42(7), 664-672.
- Neumark-Sztainer, D., Wall, M., Story, M., & Standish, A. R. (2012). Dieting and Unhealthy Weight Control Behaviors During Adolescence: Associations With 10-Year Changes in Body Mass Index. *Journal of Adolescent Health*, 50(1), 80-86.
- Neumark-Sztainer, D. R., Bauer, K. W., Friend, S., Hannan, P. J., Story, M., & Berge, J. M. (2010). Family weight talk and dieting: how much do they matter for body dissatisfaction and disordered eating behaviors in adolescent girls? *Journal of Adolescent Health*, 47(3), 270-276.
- Norimah, A. K., Mohd Nasir, M. T., Hazizi, A. S., Suraya, I., Loh, S., & Nurliyana, A. R. (2014). Association of body weight status and socio-demographic factors with food habits among preschool children in Peninsular Malaysia. *Malaysian Journal of Nutrition*, 20(3), 303-315.

- Onis, M. d., Onyango, A. W., Borghi, E., Siyam, A., Nishida, C., & Siekmann, J. (2007). Development of a WHO growth reference for school-aged children and adolescents. *Bulletin of the World Health Organization*, 85(9), 660-667.
- Palma-Coca, O., Hernandez-Serrato, M. I., Villalobos-Hernandez, A., Unikel-Santoncini, C., Olaiz-Fernandez, G., & Bojorquez-Chapela, I. (2011). Association of socioeconomic status, problem behaviors, and disordered eating in Mexican adolescents: results of the Mexican National Health and Nutrition Survey 2006. *Journal of Adolescent Health*, 49(4), 400-406.
- Patton, G. C., Selzer, R., Coffey, C., Carlin, J. B., & Wolfe, R. (1999). Onset of adolescent eating disorders: population based cohort study over 3 years. *BMJ : British Medical Journal*, 318(7186), 765-768.
- Patton, G. C., & Viner, R. (2007). Pubertal transitions in health. *The Lancet*, 369(9567), 1130-1139.
- Paulson, L. R., & Rutledge, P. C. (2014). Effects of perfectionism and exercise on disordered eating in college students. *Eating Behaviors*, 15(1), 116-119.
- Pearson, C. M., Combs, J. L., & Smith, G. T. (2010). A risk model for disordered eating in late elementary school boys. *Psychology of Addictive Behaviors*, 24(4), 696-704.
- Peat, J., & Barton, B. (2008). *Medical statistics: A guide to data analysis and critical appraisal*: John Wiley & Sons.
- Petersen, A. C., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report measure of pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*, 17(2), 117-133.
- Pon, L. W., Jr., Kandiah, M., & Mohd Nasir, M. T. (2004). Body image perception, dietary practices and physical activity of overweight and normal weight Malaysian female adolescents. *Malaysian Journal of Nutrition*, 10(2), 131-147.
- Pourghassem Gargari, B., Kooshavar, D., Seyed Sajadi, N., Safoura, S., Hamed Behzad, M., & Shahrokhi, H. (2011). Disordered Eating Attitudes and Their Correlates among Iranian High School Girls. *Health Promotion Perspectives*, 1(1), 41-49.
- Quiles-Marcos, Y., Balaguer-Sola, I., Pamies-Aubalat, L., Quiles-Sebastian, M. J., Marzo-Campos, J. C., & Rodriguez-Marin, J. (2011). Eating habits, physical activity, consumption of substances and eating disorders in adolescents. *Spanish Journal of Psychology*, 14(2), 712-723.
- Ranjana, S., Mahomoodally, F. M., & Ramasawmy, D. (2013). Is healthy eating behaviour common among school adolescents in Mauritius? *Current Research in Nutrition and Food Science Journal*, 1(1), 11-22.

- Rayner, K. E., Schniering, C. A., Rapee, R. M., & Hutchinson, D. M. (2013). A longitudinal investigation of perceived friend influence on adolescent girls' body dissatisfaction and disordered eating. *Journal of Clinical Child and Adolescent Psychology*, 42(5), 643-656.
- Reba-Harrelson, L., Von Holle, A., Hamer, R. M., Swann, R., Reyes, M. L., & Bulik, C. M. (2009). Patterns and prevalence of disordered eating and weight control behaviors in women ages 25-45. *Eating and Weight Disorders*, 14(4), e190-198.
- Rhea, D. J., & Thatcher, W. G. (2013). Ethnicity, ethnic identity, self-esteem, and at-risk eating disordered behavior differences of urban adolescent females. *Eating Disorders: The Journal of Treatment and Prevention*, 21(3), 223-237.
- Ricciardelli, L. A., & McCabe, M. P. (2001). Children's body image concerns and eating disturbance: a review of the literature. *Clinical Psychology Review*, 21(3), 325-344.
- Ricciardelli, L. A., & McCabe, M. P. (2004). A biopsychosocial model of disordered eating and the pursuit of muscularity in adolescent boys. *Psychological Bulletin*, 130(2), 179-205.
- Ridley, K., Ainsworth, B. E., & Olds, T. S. (2008). Development of a compendium of energy expenditures for youth. *International Society of Behavioral Nutrition and Physical Activity*, 5, 4552.
- Rodgers, R. F., Ganchou, C., Franko, D. L., & Chabrol, H. (2012). Drive for muscularity and disordered eating among French adolescent boys: A sociocultural model. *Body Image*, 9(3), 318-323.
- Rosenberg, M. (1965). The measurement of self-esteem. *Society and the Adolescent Self Image*, 297, V307.
- Sakinah, H., Seong-Ting, C., Rosniza, R., & Jayah, K. (2012). Socio-demographic, dietary and physical activity determinants of adolescents overweight and obesity in Kelantan. *Health and the Environment Journal*, 3(1), 44-53.
- Santos, M., Richards, C. S., & Bleckley, M. K. (2007). Comorbidity between depression and disordered eating in adolescents. *Eating Behaviors*, 8(4), 440-449.
- Schwarzer, R., & Fuchs, R. (1996). Self-efficacy and health behaviours. *Predicting Health Behaviour*, 163-196.
- Schwarzer, R., & Renner, B. (2000). Social-cognitive predictors of health behavior: action self-efficacy and coping self-efficacy. *Health Psychology*, 19(5), 487-495.

- Sekiyama, M., Roosita, K., & Ohtsuka, R. (2012). Snack foods consumption contributes to poor nutrition of rural children in West Java, Indonesia. *Asia Pacific Journal of Clinical Nutrition*, 21(4), 558-567.
- Shomaker, L. B., & Furman, W. (2009). Interpersonal influences on late adolescent girls' and boys' disordered eating. *Eating Behaviors*, 10(2), 97-106.
- Sjoberg, A., Hallberg, L., Hoglund, D., & Hulthen, L. (2003). Meal pattern, food choice, nutrient intake and lifestyle factors in The Goteborg Adolescence Study. *European Journal of Clinical Nutrition*, 57(12), 1569-1578.
- Solmi, F., Hatch, S. L., Hotopf, M., Treasure, J., & Micali, N. (2014). Prevalence and correlates of disordered eating in a general population sample: the South East London Community Health (SELCoH) study. *Social Psychiatry and Psychiatric Epidemiology*, 1-12.
- Soo, K. L., Zalilah, M., Mohd Nasir, M. T., & Bahaman, A. S. (2008). Eating behaviour, body image, and self-esteem of adolescent girls in Malaysia. *Perceptual and Motor Skills*, 106(3), 833-844.
- Stang, J., & Story, M. (2005). *Guidelines for adolescent nutrition services*: Center for Leadership, Education and Training in Maternal and Child Nutrition.
- Stice, E., Davis, K., Miller, N. P., & Marti, C. N. (2008). Fasting increases risk for onset of binge eating and bulimic pathology: a 5-year prospective study. *Journal of Abnormal Psychology*, 117(4), 941-946.
- Strien, Tatjana van & Thames Valley Test Company (2002). *Dutch eating behaviour questionnaire : manual*. Thames Valley Test Company, Bury St. Edmunds
- Strong, W. B., Malina, R. M., Blimkie, C. J. R., Daniels, S. R., Dishman, R. K., Gutin, B., Hergenroeder, A. C., Must, A., Nixon, P. A., Pivarnik, J. M., Rowland, T., Trost, S., & Trudeau, F. (2005). Evidence Based Physical Activity for School-age Youth. *The Journal of Pediatrics*, 146(6), 732-737.
- Swanson, S. A., Crow, S. J., Le Grange, D., Swendsen, J., & Merikangas, K. R. (2011). Prevalence and correlates of eating disorders in adolescents: Results from the national comorbidity survey replication adolescent supplement. *Archives of General Psychiatry*, 68(7), 714-723.
- Tam, C. K., Ng, C. F., Yu, C. M., & Young, B. W. (2007). Disordered eating attitudes and behaviours among adolescents in Hong Kong: prevalence and correlates. *Journal of Paediatric and Child Health*, 43(12), 811-817.
- Tanofsky-Kraff, M., Yanovski, S. Z., Schvey, N. A., Olsen, C. H., Gustafson, J., & Yanovski, J. A. (2009). A prospective study of loss of control eating for body weight gain in children at high risk for adult obesity. *International Journal of Eating Disorders*, 42(1), 26-30.

- Tee, Ismail, Nasir, & Khatijah. (1997). Nutrient composition of Malaysian foods. Malaysian Food Composition Database Programme, c. o Institute for Medical Research, Kuala Lumpur.
- Tomanari, R. (2011). Stages of growth child development-early childhood (birth to eight years), middle childhood (eight to twelve years). Retrieved 13 August, 2015, from <http://education.stateuniversity.com/pages/1826/Child-Development-Stages-Growth.html>
- Toro, J., Gomez-Peresmitré, G., Sentis, J., Vallés, A., Casulà, V., Castro, J., Pineda, G., Leon, R., Platas, S., & Rodriguez, R. (2006). Eating disorders and body image in Spanish and Mexican female adolescents. *Social Psychiatry and Psychiatric Epidemiology*, 41(7), 556-565.
- Torun, B., Davies, P. S., Livingstone, M. B., Paolisso, M., Sackett, R., & Spurr, G. B. (1996). Energy requirements and dietary energy recommendations for children and adolescents 1 to 18 years old. *European Journal of Clinical Nutrition*, 50 Suppl 1, S37-80; discussion S80-31.
- Tremblay, L., & Lariviere, M. (2009). The influence of puberty onset, Body Mass Index, and pressure to be thin on disordered eating behaviors in children and adolescents. *Eating Behaviors*, 10(2), 75-83.
- Tsai, M. R., Chang, Y. J., Lien, P. J., & Wong, Y. (2011). Survey on eating disorders related thoughts, behaviors and dietary intake in female junior high school students in Taiwan. *Asia Pacific Journal of Clinical Nutrition*, 20(2), 196-205.
- United Nations Children's Fund (UNICEF). (2005). *Adolescent Development: Perspectives and Frameworks - A Discussion Paper*. New York, US: United Nations Children's Fund.
- U.S. Department of Health and Human Services. (2005). Body wise handbook. Eating disorders information for middle school personnel (3rd ed.). Washington, DC: US Department of Health and Human Services.
- Uzun, Ö., Güleç, N., Özşahin, A., Doruk, A., Özdemir, B., & Çalışkan, U. (2006). Screening disordered eating attitudes and eating disorders in a sample of Turkish female college students. *Comprehensive Psychiatry*, 47(2), 123-126.
- van den Berg, P. A., Mond, J., Eisenberg, M., Ackard, D., & Neumark-Sztainer, D. (2010). The Link Between Body Dissatisfaction and Self-Esteem in Adolescents: Similarities Across Gender, Age, Weight Status, Race/Ethnicity, and Socioeconomic Status. *Journal of Adolescent Health*, 47(3), 290-296.
- van Strien, T., Frijters, J. E. R., Bergers, G. P. A., & Defares, P. B. (1986). The Dutch Eating Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external eating behavior. *International Journal of Eating Disorders*, 5(2), 295-315.

- Veses, A. M., Martínez-Gómez, D., Gómez-Martínez, S., Vicente-Rodriguez, G., Castillo, R., Ortega, F. B., González-Gross, M., Calle, M. E., Veiga, O. L., & Marcos, A. (2014). Physical fitness, overweight and the risk of eating disorders in adolescents. The AVENA and AFINOS studies. *Pediatric Obesity*, 9(1), 1-9.
- Wang, Z., Zhai, F., Zhang, B., & Popkin, B. M. (2012). Trends in Chinese snacking behaviors and patterns and the social-demographic role between 1991 and 2009. *Asia Pacific Journal of Clinical Nutrition*, 21(2), 253-262.
- Wille, D.-P. N., Hölling, D.-P. H., Vloet, T. D., & Ravens-Sieberer, U. (2008). Disordered eating behaviour and attitudes, associated psychopathology and health-related quality of life: results of the BELLA study. *European Child and Adolescent Psychiatry*, 17(1), 82-91.
- Wong, Y., Chang, Y. J., & Tsao, S. W. (2014). Disturbed eating tendency and related factors in grade four to six elementary school students in Taiwan. *Asia Pacific Journal of Clinical Nutrition*, 23(1), 112-120.
- Yang, S. J., Kim, J. M., & Yoon, J. S. (2010). Disturbed eating attitudes and behaviors in South Korean boys and girls: a school-based cross-sectional study. *Yonsei Medical Journal*, 51(3), 302-309.
- Yannakoulia, M., Matalas, A.-L., Yiannakouris, N., Papoutsakis, C., Passos, M., & Klimis-Zacas, D. (2004). Disordered eating attitudes: an emerging health problem among Mediterranean adolescents. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*, 9(2), 126-133.
- Yusoff, F., Saari, R., Naidu, B. M., Ahmad, N. A., Omar, A., & Aris, T. (2014). Methodology of the national school-based health survey in malaysia, 2012. *Asia-Pacific Journal of Public Health*, 26(5 Suppl), 9S-17S.
- Zalilah, M., & Zaidah, M. (2005). Correlates of Children's Eating Attitude Test scores among primary school children. *Perceptual and Motor Skills*, 100(2), 463-472.