



UNIVERSITI PUTRA MALAYSIA

***TUBERCULOSIS-RELATED KNOWLEDGE, ATTITUDE AND
PREVENTIVE PRACTICES AMONG ABATTOIR WORKERS
IN GUSAU, NIGERIA***

ISMAILA USMAN GUSAU

FPSK(M) 2016 24



**TUBERCULOSIS-RELATED KNOWLEDGE, ATTITUDE AND
PREVENTIVE PRACTICES AMONG ABATTOIR WORKERS
IN GUSAU, NIGERIA**

By

ISMAILA USMAN GUSAU

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfillment of the Requirements for the Degree of
Master of Science**

January 2016

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree of Master of Science

TUBERCULOSIS-RELATED KNOWLEDGE, ATTITUDE AND PREVENTIVE PRACTICES AMONG ABATTOIR WORKERS IN GUSAU, NIGERIA

By

ISMAILA USMAN GUSAU

January 2016

Chair : Assoc. Prof. Hejar Binti Abd Rahman, PhD
Faculty : Medicine and Health Sciences

Introduction: Tuberculosis (TB) is still a major health concern worldwide. The global burden of TB is estimated to be 8.5 to 9.2 million with 1.2 to 1.5 million deaths annually. Asia and Africa are affected with 57% and 26% of the disease burden respectively. Nigeria ranked third among the world's six countries with high burden of TB and has the highest estimated number of new cases with nearly 590,000 new cases annually. Nigeria is one of the African countries where bovine TB is wide spread in both cattle and humans. Although, the current status on the actual prevalence rate of bovine TB at the national level is unknown but from the limited surveys conducted by some researchers reported that, the prevalence of TB due to *M. bovis* is up to 14%. The isolation and identification of *M. bovis* in fresh and sour milk sold at local markets, sputum and tissue samples from humans especially herdsmen, abattoir and slaughter houses has been reported. This indicates that the disease occurs both in cattle and humans in Nigeria. Thus, reason to assess the knowledge, attitude and practices of abattoir workers on TB.

Objectives: To determine the level of knowledge, attitude and preventive practices on tuberculosis among abattoir workers in Gusau, Zamfara State, Nigeria.

Methodology: A cross-sectional study was conducted in Gusau Modern Abattoir, Zamfara State, Nigeria. The abattoir was chosen because it is the only one in the state. A total of 416 eligible abattoir workers were selected using proportionate stratified random sampling. Data was collected on knowledge, attitude and preventive practices on TB among the abattoir workers from May to July 2014, through face to face interview and self-administered questionnaire, which was validated by content and face validities. Reliability was measured by checking the internal consistency using Cronbach's alpha and values above 0.7 were accepted. Descriptive analysis, chi-square and multivariate logistic regression were conducted. Significant level (p) was set at 0.05.

Results: The response rate was 91.3%, the overall mean knowledge score of the respondents was 29.76 (SD) 14.15, with 53.2% having high knowledge and 46.8% having low knowledge on TB. However, 52.9%, mean (SD) is 42.76 (6.23) of the respondents have positive attitude and 58.7%, mean (SD) is 9.13 (4.50) practices preventive measures towards TB. Predictors of knowledge, attitude and preventive practices were determined using logistic regression analysis and results from Chi-square tests with $p < 0.25$ were included. Years of working experience (AOR = 1.98; CI = 1.231 – 3.189), TB screening (AOR = 3.04; CI = 1.446 – 6.419) and family member history of TB infection (AOR = 17.35; CI = 2.274 – 132.421) were found to be the predictors of knowledge on TB. Immunization (AOR = 2.45; CI = 1.279 – 4.688) and training (AOR = 4.60; CI = 2.254 – 9.385) were found to be the predictors of attitude towards TB. For preventive practices, history of TB infection (AOR = 5.21; CI = 1.547 – 17.560) and family member history of TB infection (AOR = 3.54; CI = 1.330 – 9.429) were found to be the predictors of preventive practices regarding TB.

Conclusion: This study has identified that there is low level of knowledge, negative attitude and poor preventive practices on TB among abattoir workers in Zamfara State. However, predictors of knowledge on TB were years of working experience, TB screening and family member infected with TB. Predictors of attitude towards TB were immunization and training. Predictors of TB preventive practices were personal and family member infected with TB. The study findings will help in development of interventional programs for efficient control of TB (human and animal).

Keywords: Knowledge, attitude, preventive practices, tuberculosis, abattoir, Zamfara State

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

**TUBERKULOSIS BERKAITAN PENGETAHUAN, SIKAP DAN AMALAN
PENCEGAHAN DALAM KALANGAN PEKERJA ABATOIR
DI GUSAU, NIGERIA**

Oleh

ISMAILA USMAN GUSAU

Januari 2015

Pengerusi : Prof. Madya Hejar Binti Abd Rahman, PhD
Fakulti : Perubatan dan Sains Kesihatan

Pengenalan: Tuberkulosis (TB) masih merupakan masalah kesihatan utama di seluruh dunia. Beban global TB dianggarkan pada 8.5 hingga 9.2 juta dengan 1.2 hingga 1.5 juta kematian setiap tahun. Asia dan Afrika telah dijangkiti dengan beban penyakit sebanyak 57% dan 26% masing-masing. Nigeria menduduki tempat ketiga di antara enam negara di dunia dengan beban tinggi TB dan mempunyai bilangan anggaran kes baru tertinggi iaitu hampir 590,000 kes baru setiap tahun. Nigeria merupakan salah satu negara Afrika di mana TB bovin luas tersebar pada lembu dan juga manusia. Walaupun, status semasa kadar kekerapan sebenar TB bovin di peringkat kebangsaan tidak diketahui tetapi dari kajian terhad dijalankan oleh beberapa penyelidik melaporkan bahawa, kekerapan TB disebabkan oleh *M. bovis* adalah sehingga 14%. Pengasingan dan pengenalpastian *M. bovis* dalam susu segar dan masam dijual di pasar tempatan, kahak dan sampel tisu daripada manusia terutama sekali pada penggembala, abatoir dan rumah-rumah penyembelihan telah dilaporkan. Ini menunjukkan bahawa penyakit ini berlaku pada lembu dan juga pada manusia di Nigeria. Justeru, alasan untuk menilai pengetahuan, sikap dan amalan pekerja abatoir mengenai TB.

Objektif: Untuk mengenalpasti tahap pengetahuan, sikap dan amalan pencegahan mengenai tuberkulosis dalam kalangan pekerja abatoir di Gusau, Zamfara, Nigeria.

Metodologi: Satu kajian keratan rentas telah dijalankan di Abatoir Moden Gusau, Zamfara, Nigeria. Abatoir ini telah dipilih kerana ia merupakan satu-satunya abatoir di negeri tersebut. Seramai 416 pekerja abatoir yang layak telah dipilih menggunakan persampelan rawak berstrata berkadaran. Data yang dikumpulkan adalah mengenai pengetahuan, sikap dan amalan pencegahan mengenai TB dalam kalangan pekerja abatoir dari Mei hingga Julai 2014, melalui temuduga bersemuka dan soal selidik tadbir sendiri yang telah disahkan melalui kesahan kandungan dan muka. Kebolehpercayaan diukur dengan memeriksa ketekalan dalaman menggunakan nilai alfa Cronbach dan nilai 0.7 keatas boleh diterima. Analisis deskriptif, khi kuasa dua

dan regresi multivariat logistik telah dijalankan. Aras signifikan (p) telah ditetapkan pada 0.05.

Hasil: Kadar sambutan ialah 91.3%, skor min keseluruhan pengetahuan responden adalah 29.76 (SP) 14.15, dengan 53.2% mempunyai pengetahuan yang tinggi dan 46.8% mempunyai pengetahuan yang rendah mengenai TB. Walau bagaimanapun, 52.9% min (SP) adalah 42.76 (6.23) daripada responden mempunyai sikap positif dan 58.7%, min (SP) adalah 9.13, (4.50) mengamalkan langkah-langkah pencegahan terhadap TB. Peramal pengetahuan, sikap dan amalan pencegahan telah ditentukan dengan menggunakan analisis regresi logistik dan hasil daripada ujian khi kuasa dua dengan $p < 0.25$ telah dimasukkan. Tahun pengalaman bekerja (AOR = 1.98; CI = 1.231 – 3.189), saringan TB (AOR = 3.04; CI = 1.446 – 6.419) dan sejarah ahli keluarga yang dijangkiti TB (AOR = 17.35; CI = 2.274 – 132.421) didapati merupakan peramal pengetahuan mengenai TB. Imunisasi (AOR = 2.45; CI = 1.279 – 4.688) dan latihan (AOR = 4.60; CI = 2.254 – 9.385) didapati merupakan peramal sikap terhadap TB. Bagi amalan pencegahan, sejarah jangkitan TB (AOR = 5.21; CI = 1.547 – 17.560) dan sejarah ahli keluarga yang dijangkiti TB (AOR = 3.54; CI = 1.330 – 9.429) didapati merupakan peramal amalan pencegahan mengenai TB.

Kesimpulan: Kajian ini telah mengenalpasti bahawa terdapat tahap pengetahuan yang rendah, sikap negatif dan amalan pencegahan yang rendah terhadap TB dalam kalangan pekerja abatoir di Zamfara. Walau bagaimanapun, peramal pengetahuan terhadap TB adalah tahun pengalaman bekerja, saringan TB dan ahli keluarga dijangkiti TB. Peramal bagi sikap terhadap TB adalah imunisasi dan latihan. Peramal bagi amalan pencegahan TB adalah diri sendiri dan ahli keluarga yang dijangkiti oleh TB. Hasil kajian boleh membantu dalam pembangunan program intervensi bagi pengawalan TB yang berkesan (manusia dan haiwan).

Kata kunci: Pengetahuan, sikap, amalan pencegahan, tuberkulosis, abatoir, Zamfara

ACKNOWLEDGEMENT

I wish to acknowledge the support of my supervisor Assoc. Prof. Dr. Hejar Binti Abd Rahman and my co-supervisor Dr. Suhainizam Muhamad Saliluddin for their guidance, encouragement and commitment during the period of this study.

I am sincerely grateful to Ministry of Animal Health and Livestock Development, Gusau, Zamfara State for granting me the permission to undergo this programme. Special thanks to Department of Public Health and management of Gusau Modern Abattoir for their kind support during my research.

I am very grateful to my parents, wife, kids and siblings who continued to support and encourage me in all aspects throughout my program. I would also like to appreciate the efforts of all my lecturers for the invaluable knowledge they impacted on us, colleagues and friends who helped directly or indirectly in making this study a success.

I certify that a Thesis Examination Committee has met on 22 January 2016 to conduct the final examination of Ismaila Dr Usman Gusau on his thesis entitled "Tuberculosis-Related Knowledge, Attitude and Preventive Practices among Abattoir Workers in Gusau, Nigeria" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Kulanthayan a/I K.C. Mani @ Subramanian, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Malina binti Osman, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Rahmah Mohd Amin, PhD

Associate Professor
Universiti Sultan Zainal Abidin
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 24 March 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Hejar Binti Abd Rahman, M.D., Master of Community Health

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairperson)

Suhainizam Muhamad Saliluddin, MB, BCh, BAO, MPH

Senior Lecturer (Medical)
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

BUJANG BIN KIN HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Ismaila Usman Gusau, GS35365

Declaration by Members of Supervisory Committee

This is to confirm that:

- The research conducted and the writing of this thesis was under our supervision;
- Supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of
Chairperson of
Supervisory

Committee: Hejar Binti Abd Rahman, M.D., Master of Community Health

Signature: _____

Name of
Member of
Supervisory

Committee: Suhainizam Muhamad Saliluddin, MB, BCh, BAO, MPH

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENT	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xvi
CHAPTER	
1 INTRODUCTION	1
1.1 Background	1
1.2 Problem Statement	3
1.3 Significance of the Study	5
1.4 Objectives of the Study	5
1.4.1 General Objective	5
1.4.2 Specific Objectives	5
1.5 Research Hypotheses	6
2 LITERATURE REVIEW	7
2.1 Historical Perspective of TB	7
2.2 Etiology of TB	8
2.3 Global Epidemiology of Tuberculosis	8
2.4 Epidemiology of Tuberculosis in Africa	9
2.5 Epidemiology of Tuberculosis in Nigeria	9
2.6 Human Tuberculosis	10
2.6.1 Mode of Transmission and Pathogenesis	10
2.6.2 Signs and Symptoms	11
2.6.3 Diagnosis of Human TB	12
2.6.4 Treatment of Human TB	13
2.6.5 Prevention and Control of Human TB in Nigeria adopted from WHO	13
2.6.5.1 Primary prevention of Human TB	13
2.6.5.2 Secondary prevention of Human TB	13
2.6.5.3 Tertiary prevention of Human TB	17
2.7 Animal Tuberculosis (Bovine TB)	18
2.7.1 Mode of Transmission and Pathogenesis of Bovine TB	18
2.7.2 Signs and Symptoms of Bovine TB	18
2.7.3 Diagnosis of Bovine TB	18
2.7.4 Prevention and Control of Bovine TB in Nigeria	20
2.8 Zoonotic Tuberculosis (TB)	21
2.9 Knowledge on TB	22
2.10 Attitude towards TB	24

2.11	Preventive practices regarding TB	25
2.12	Conceptual Framework	26
3	MATERIALS AND METHODS	28
3.1	Study Location	28
3.2	Study Design	29
3.3	Study Duration	29
3.4	Study Population	29
3.5	Inclusion Criteria	29
3.6	Sampling Frame	30
3.7	Sample Size Estimation	30
3.8	Sampling Method	31
3.9	Instrument of the Study	33
3.10	Components of the Questionnaire	33
3.10.1	Section A: Socio-Demography	33
3.10.2	Section B: Personal and Medical Histories	33
3.10.3	Section C: Knowledge on Tuberculosis	33
3.10.4	Section D: Attitude regarding Tuberculosis	34
3.10.5	Section E: Preventive Practices regarding TB	34
3.11	Data Collection	34
3.12	Study Variables	35
3.13	Dependent Variables	35
3.14	Independent Variables	35
3.15	Definition of Operational Variables	35
3.16	Quality Control	37
3.16.1	Content Validity	37
3.16.2	Face Validity	37
3.16.3	Internal Consistency of the Questionnaire	37
3.17	Data Analysis	37
3.18	Ethical Consideration	38
4	RESULTS	39
4.1	Response Rate	39
4.2	Test of Normality	39
4.3	Socio-Demographic Characteristics of the Respondents	40
4.4	Personal and Medical Histories of the Respondents	41
4.5	Sources of Information on TB among Abattoir Workers	42
4.6	Knowledge on TB among Abattoir Workers	42
4.7	Level of Knowledge on TB among Abattoir Workers	45
4.8	Attitude towards TB among Abattoir Workers	45
4.9	Level of Attitude towards TB among Abattoir Workers	46
4.10	Preventive Practices regarding TB among Abattoir Workers	47
4.11	Level of Preventive Practices regarding TB among Abattoir Workers	48
4.12	Association between Socio-Demography, Personal and Medical History and Level of Knowledge of the Respondents	48
4.13	Association between Socio-Demography, Personal and Medical History and Level of Attitude of the Respondents	50

4.14	Association between Socio-Demography, Personal and Medical History and Level of Preventive Practices of the Respondents	53
4.15	Predictors of Knowledge	54
4.16	Predictors of Attitude	56
4.17	Predictors of Preventive Practices	57
5	DISCUSSION	59
5.1	Socio-Demographic Characteristics, Medical and Personal History of the Abattoir Workers	59
5.2	Sources of Information about TB	60
5.3	Knowledge of Abattoir Workers on TB	60
5.4	Factors Associated with Knowledge of the Abattoir Workers on TB	61
5.5	Attitude of Abattoir Workers towards TB	63
5.6	Factors Associated with Attitude of Abattoir Workers towards TB	63
5.7	Preventive Practices of Abattoir Workers regarding TB	64
5.8	Factors Associated with Preventive Practices of Abattoir regarding TB	65
5.9	Predictors of Knowledge, Attitude and Preventive Practices of Abattoir Workers on TB	66
6	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	68
6.1	Summary and Conclusion	68
6.2	Strengths and Limitation of the Study	68
6.2.1	Strength of the Study	68
6.2.2	Limitations of the Study	69
6.3	Recommendations and Further Studies	69
	REFERENCES	70
	APPENDICES	78
	BIODATA OF STUDENT	101
	LIST OF PUBLICATIONS	102

LIST OF TABLES

Table	Page
3. 1: Variables used in sample size estimation	30
3. 2: Range of knowledge scores	34
3. 3: Internal consistency of questionnaire	37
4. 1: Test of normality	39
4. 2: Socio-demographic characteristics of the respondents (N = 380)	40
4. 3: Personal and medical histories of the respondents (N = 380)	41
4. 4: Sources of information on TB among abattoir workers	42
4. 5: Knowledge on TB among abattoir workers	43
4. 6: Level of knowledge on TB among abattoir workers	45
4. 7: Attitude towards TB among abattoir workers	46
4. 8: Practice regarding TB among abattoir workers	47
4. 9: Level of practice regarding TB among abattoir workers	48
4. 10: Association between socio-demography, personal and medical history and knowledge scores of the respondents	49
4. 11: Association between socio-demography, personal and medical history and attitude scores of the respondents	51
4. 12: Association between socio-demography, personal and medical history and practice scores of the respondents	53
4. 13: Simple logistic regression showing crude odd ratio (OR) of predictors of knowledge	55
4. 14: Multiple logistic regression showing adjusted odd ratio (OR) of predictors of knowledge	56
4. 15: Simple logistic regression showing crude odd ratio (OR) of predictors of attitude	56
4. 16: Multiple logistic regression showing adjusted odd ratio (OR) of predictors of attitude	57

4. 17:	Simple logistic regression showing crude odd ratio (OR) of predictors of practice	58
4. 18:	Multiple logistic regression showing adjusted odd ratio (OR) of predictors of practice	58



LIST OF FIGURES

Figure	Page
2. 1: Factors associated with knowledge, attitude and practices of abattoir workers on tuberculosis	27
3. 1: Map of Nigeria showing Zamfara State	29
3. 2: Flow chart showing proportionate stratified random sampling	32



LIST OF ABBREVIATIONS

<	Less than
>	Greater than
≤	Less than or equal to
≥	Greater than or equal to
ACSM	Advocacy, communication and social mobilization
AIDS	Acquired immunodeficiency syndrome
AOR	Adjusted odds ratio
BCG	Bacillus Calmette–Guérin
BTB	Bovine tuberculosis
CDR	Case detection rate
CI	Confidence interval
COR	Crude odds ratio
df	Degree of freedom
DOTS	Directly observed therapy – short course
HIV	Human immunodeficiency virus
IQR	Inter quartile range
KAP	Knowledge, attitude and practice
LGA	Local government area
LGTBLS	Local government tuberculosis and leprosy supervisor
LTBI	Latent tuberculosis infection
MDR-TB	Multi drug resistant tuberculosis
n	Number
NASCAP	National AIDS and STDs control program
NPI	National program on immunization
NTBLCP	National tuberculosis and leprosy control program
OR	Odds ratio
p	Level of significance
PHC	Primary health care
PPE	Personal protective equipment
PPM	Public - private mix

PTB	Pulmonary tuberculosis
RM	Malaysian ringgit
SD	Standard deviation
STBLCP	State tuberculosis and leprosy control program
TB	Tuberculosis
TSR	Treatment success rate
TST	Tuberculin skin test
WHO	World health organization



CHAPTER 1

INTRODUCTION

1.1 Background

Tuberculosis (TB) has remained as a major public health concern globally, responsible for poor health condition suffered among millions of people every year. This disease is the second main causes of death from an infectious disease worldwide, after the human immunodeficiency virus (HIV). In 2013, about 9 million new TB cases and 1.5 million TB deaths were estimated. About 85% of the disease burden is found in Asia and Africa (WHO, 2014). The TB situation has worsened over the last three decades, which can be attributed to the HIV/AIDS pandemic (Getahun, Gunneberg, Granich, & Nunn, 2010).

Statistically, the African region captured one quarter of the world's cases, and the highest rates of cases and deaths relative to the population (280 incident cases per 100,000 on average, more than double the global average of 126) (WHO, 2014). The disease burden is high in this region due to the connection between the two diseases; TB and the human immunodeficiency virus as well as the acquired immune deficiency syndrome (HIV/AIDS), which Africa accounts for 78% of the global cases (Getahun et al., 2010). The totals in 2014 global TB report are higher than those reported in 2013, mainly because of upward revisions made to estimate the number of TB cases and deaths in Nigeria following the finalisation of results from Nigeria's first-ever national TB prevalence survey (completed in 2012). Given the size of the population and the high TB burden in Nigeria, these revisions made have affected the estimation globally (WHO, 2014).

Tuberculosis (TB) is caused by *Mycobacterium tuberculosis* complex species. TB mainly affects the lungs, but it can also affect other organs located in the central nervous system, lymphatic system, and circulatory system among others. The disease is airborne transmitted, depending on the degree of infectiousness of the source patient and the current situation of the environment at the time when the TB exposure took place. This initial infection (primary TB) will progress rapidly to a full blown TB disease in some individuals (especially young and immune-compromised individuals), but it will also resolves spontaneously in most individuals. This condition whereby the organism lies dormant is called as latent TB infection (LTBI) (National Tuberculosis Controllers Association & Centers for Disease Control and Prevention (CDC), 2005).

In humans, tuberculosis (TB) is a pulmonary and systemic disease primarily caused by a bacterium called *Mycobacterium tuberculosis*. Small fraction of disease is caused by *M. bovis* (LoBue, Enarson & Thoen, 2010; Thoen, LoBue, Enarson, Kaneene & Kantor, 2009). Pulmonary tuberculosis (PTB) is characterized by a series of symptoms namely fever, cough, night sweats and weight loss; and occasionally infected individual may cough up blood while the extra pulmonary tuberculosis (EPTB) is

characterized by pleural effusion, lymphadenopathy, pericardial effusion, miliary disease, and meningitis. TB patients will usually present with constitutional symptoms and local features related to the site of disease (Jamison, 2006).

On the other hand, animal tuberculosis (TB) also known as bovine TB is mostly caused by *Mycobacterium bovis*. The major reservoir of *M. bovis* is infected cattle. Cattle TB also has the ability of spreading to other domestic and wild animals through direct or indirect contact with the infected animal, inhalation of the infected droplets and ingestion of the contaminated milk in calves (Anaelom, Ikechukwu, Sunday, & Nnaemeka, 2010; Taylor, Murphy, Hopkins, Rutland, & Chistov, 2007). Bovine TB is an important zoonotic disease that is transmissible to humans through various routes such as inhalation of aerosols, which will lead to pulmonary TB. While contaminated milk and meat ingested mostly will cause extra pulmonary TB. The lesions of extra pulmonary TB are mostly present in the lymph nodes, bones and joints, genitourinary system, and the meninges. Bovine TB is also transmitted through handling contaminated meat which is a less common route of traumatic inoculation into the skin. There are rare anecdotal records shown of human to human *M. bovis* transmission. The disease caused by *M. bovis* is clinically indistinguishable from that caused by *M. tuberculosis* (Ayele, Neill, Zinsstag, Weiss, & Pavlik, 2004; Grange, 2001).

In industrialized countries such as the UK, through several initiatives such as milk pasteurization, animal TB control and elimination programmes, have managed to reduce the incidence of disease caused by *M. bovis* drastically in both cattle and humans, where control measures and herd testing have been practiced since the 1930s bovine TB accounts for less than 1% of human TB cases, and tends to be a disease that is associated with individuals being in close proximity to livestock, such as farmers, veterinary surgeons and abattoir workers (Taylor et al., 2007). However, in tropic countries, the risk factors for *M. bovis* in both animals and humans are visible. Although there is lack of data regarding the spread of bovine TB amongst the human population in the developing country, the global estimation reveals around 2.1% of population infected with pulmonary TB and 9.4% of extra-pulmonary TB cases are attributed to *M. bovis* (Anaelom et al., 2010).

The role of human behaviour has been increasingly recognized as part of the contributing factors in health and illness. Health is no longer considered simply as a biomedical problem; rather, it is also influenced by other factors such as social, cultural, physiological, economic and political factors that determine the behaviour of the people concerned (Suleiman, Sahal, Sodemann, Elsony, & Aro, 2014). In social and behavioural sciences, several models have been developed to explain and enhance health behaviours and sustained behavioural changes, also in different cultural contexts (Petrie, Jago & Devcich, 2007; Whitehead, 2001). Some of the models widely used that support program planning processes in health promotion and education are health belief model, social learning theory, social cognitive theory, theory of reasoned action, theory of planned behavior and the transtheoretical model (Glanz, Rimer & Viswanath, 2008).

The task of health behavior and health education is both to understand health behavior and to transform the knowledge about the behavior into effective strategies that will improve health, to close the gap between what is known about ideal health practice and what is actually practiced. Application of effective health behavior interventions can reduce substantial suffering, premature mortality and medical costs. Changes in the lifestyle can be facilitated by a combination of efforts to enhance awareness, change behavior and create environment that support good health practices (Glanz et al., 2008).

Knowledge, Attitude and Practice (KAP) surveys in TB can identify factors that may facilitate understanding and action, as well as pose problems or create barriers for TB control efforts. Among the factors are knowledge gaps, cultural beliefs, or behavioural patterns. They can identify factors influencing behaviour that are not known and, reasons for the attitudes. Communication processes and sources also assessed by the KAP survey as the key to define effective activities and messages in TB prevention and control. These surveys may be used to identify problems, need, as well as barriers in program delivery and solutions for improving quality and accessibility of services. From the data collected, this will allow program managers to determine TB program priorities, to calculate resources needed for various activities, to choose the most effective communication channels and messages, to set up baseline levels and to ascertain change that results from the interventions. For advocacy purposes, KAP data provides national TB program managers and their staff with key information that is required in order to make strategic decisions (WHO, 2008).

Knowledge and perception among the community members in regards to the causes of TB is very important and it may reduce or increase the transmission of the disease. In developing countries like Nigeria, among the factors that may delay the diagnosis of TB are practices and beliefs by the community such as high stigma attached to the disease, poor sanitation, crowded dwellings and failure to recognize symptoms early hence increasing the spread of the disease in the community. Due to that, it increases the spread of the disease in the community.

Other factors would be due to close proximity between animals and humans as large proportion of the populations of this settings are depending on agriculture as their livelihoods, the practice of drinking unpasteurised milk and eating uncooked or partially cooked meat, and social as well as economic challenges that limit use of effective control measures such as poverty and low health literacy. The high HIV infection rates and the associated increase in vulnerability to TB and other disease infections is also a critical factor in these settings (Abubakar et al., 2011; Shitaye, Tsegaye & Pavlik, 2007; Woith & Larson, 2008).

1.2 Problem Statement

Tuberculosis has been portrayed as a complex, long standing problem in Nigeria as it is embedded in deep social, economic and cultural determinants such as high stigma of the society that is attached to the disease, low awareness and low health knowledge,

poverty, poor hygiene, and crowded dwellings. The disease is exacerbated due to poor knowledge, negative attitude and social and economic challenges that constrain use of effective control measures (Ayele et al., 2004; Grange, 2001; Mangtani et al., 2014; Shitaye et al., 2007).

The World Health Organisation reported that Nigeria has a population of over 170 million people and ranked 3rd among the world's six countries with high rate of TB burden. This report also revealed that among all African countries, Nigeria has the highest estimated number of new TB cases, with nearly 590,000 new TB cases annually (WHO, 2014). While there is shortage of data with regards to the contribution of animal TB to the total overall human TB prevalence in Nigeria, molecular analysis of mycobacterial strains isolated from both pulmonary and extra pulmonary TB cases have shown that up to 14% of them belong to *M. bovis* and it is associated with individuals in close proximity to livestock (including abattoir workers) (Abubakar et al., 2011; Taylor et al., 2007).

Prevalence of the disease among abattoir workers has not been documented in Nigeria; however prevalence of the disease among livestock traders who are among the high risk group due to their close proximity to livestock was reported to be 10%. Among which 7.1% was caused by *M. tuberculosis* and 2.9% due to *M. bovis* (Adesokan et al., 2012).

Prevention of spread of TB can only be achieved with high level of knowledge on various modes of transmission, treatment and preventive measures, positive attitude towards the disease and preventive practices especially among the high risk group such as the abattoir workers. Although, high awareness about TB was reported among meat handlers in Nigeria (80%), knowledge on transmission of the disease from animals to humans and consumption of meat infected with TB as a source of BTB in humans was low (31% and 34.1% respectively) (Hambolu et al., 2013).

Similarly, 36.7% and 34.7% among meat handlers do not agree that they have an increased risk of contracting the BTB because of their work and the disease is treatable respectively. About 21.5% among them reported eating the visible infected parts of the lung in-order to convince customers to buy meat (Hambolu et al., 2013). It has also been reported in another study conducted among butchers in Nigeria that, only 15.5% practice hand washing, 14.4% wear personal protective equipment (PPE) and 12.6% were immunized against TB (Rosiji & Adesokan, 2012).

Despite this, currently there is no study that assessed the level of knowledge, attitude and preventive practices on tuberculosis among abattoir workers in Guasu, Zamfara State, Nigeria.

1.3 Significance of the Study

The significance of this study is to determine the level of knowledge, attitude and preventive practices on TB among abattoir workers in Gusau, Zamfara State, Nigeria. The study is necessary as no research has been conducted in this area and other studies revealed low level of knowledge, negative attitude and little or no compliance with TB preventive practices among different study populations.

Therefore, the results from this study if proven positive would provide evidence base information which would help to enhance the development of effective interventional programs between veterinary and medical professions for efficient control of TB in the state and Nigeria as a whole.

1.4 Objectives of the Study

1.4.1 General Objective

The aim of this research is to determine the level of knowledge, attitude and preventive practices on tuberculosis among abattoir workers in Gusau, Zamfara State.

1.4.2 Specific Objectives

- i. To determine the socio-demographic characteristics (such as age, gender, marital status, occupation, ethnicity, level of education and monthly income), personal and medical histories (working experience, training on meat handling, TB screening and immunization) of the abattoir workers.
- ii. To determine the level of knowledge on human and animal TB (in terms of clinical signs, mode of transmission, treatment and preventive measures) among abattoir workers.
- iii. To determine the level of attitude towards human and animal tuberculosis with respect to risk of contraction and preventive measures among abattoir workers.
- iv. To determine the preventive practices of abattoir workers about human and animal TB.
- v. To determine the relationship between socio-demographic characteristic, personal and medical histories and knowledge of abattoir workers on TB.
- vi. To determine the relationship between socio-demographic characteristics, personal and medical histories and attitude of abattoir workers towards TB.
- vii. To determine the relationship between socio-demographic characteristics, personal and medical histories and preventive practices of abattoir workers regarding TB.
- viii. To determine the predictors of knowledge, attitude and preventive practices of abattoir workers in Zamfara State, Nigeria towards TB.

1.5 Research Hypotheses

- i. There is a significant association between socio-demographic characteristics, personal and medical histories and knowledge on TB among abattoir workers in Zamfara State, Nigeria.
- ii. There is a significant association between socio-demographic characteristics, personal and medical histories and attitude towards TB among abattoir workers in Zamfara State, Nigeria.
- iii. There is a significant association between socio-demographic characteristics, personal and medical histories and preventive practices regarding TB among abattoir workers in Zamfara State, Nigeria.



REFERENCES

- Abdelwahab, A. A., Sadek, R. R., Kotb, S. A. M., & Ahmed, M. A. K. (2013). Knowledge and Attitude of Patients and their Families about Tuberculosis in Minia Governorate. *AAMJ*, *11*(3), 101-130.
- Abdulraheem, I., Onajole, A., Jimoh, A., & Oladipo, A. (2011). Reasons for incomplete vaccination and factors for missed opportunities among rural Nigerian children. *J Public Health Epidemiol*, *3*(4), 194-203.
- Abubakar, U., Ameh, J., Abdulkadir, I., Salisu, I., Okaiyeto, S., & Kudi, A. (2011). Bovine tuberculosis in Nigeria: A review. *Veterinary Research*, *4*(1), 24-27.
- Addo, K. K., Mensah, G. I., Nartey, N., Nipah, G. K., Mensah, D., Aning, K. G., & Smits, H. L. (2011). Knowledge, attitudes and practices (KAP) of herdsmen in Ghana with respect to milk-borne zoonotic diseases and the safe handling of milk. *Journal of Basic and Applied Scientific Research*, *1*(10), 1556-1562.
- Adebanjo, O. D. (2011). Knowledge, attitudes and practices of healthcare workers about prevention and control of multidrug-resistant tuberculosis at Botsabelo Hospital Maseru, Lesotho. Unpublished manuscript.
- Adesokan, H. K., & Ocheja, S. E. (2014). Knowledge, attitudes and practices of traceability among livestock traders in South-Western Nigeria: Implications for sustainable livestock industry. *Tropical Animal Health and Production*, *46*(1), 159-165.
- Adesokan, H. K., Jenkins, A. O., van Soolingen, D., & Cadmus, S. I. B. (2012). *Mycobacterium bovis* infection in livestock workers in Ibadan, Nigeria: evidence of occupational exposure. *The International Journal of Tuberculosis and Lung Disease*, *16*(10), 1388-1392.
- Aliyu, A. (2013). Nigeria: The world of Illorin female butchers. *Daily Trust*. Retrieved September 10, 2014, from: <http://allafrica.com/stories/201311100048.html>.
- Amanfu, W. (2006). The situation of tuberculosis and tuberculosis control in animals of economic interest. *Tuberculosis*, *86*(3), 330-335.
- Anaelom, N. J., Ikechukwu, O. J., Sunday, W., & Nnaemeka, U. C. (2010). Zoonotic tuberculosis: A review of epidemiology, clinical presentation, prevention and control. *Journal of Public Health and Epidemiology*, *2*, 118-124.
- Ansari-Lari, M., Soodbakhsh, S., & Lakzadeh, L. (2010). Knowledge, attitudes and practices of workers on food hygienic practices in meat processing plants in Fars, Iran. *Food Control*, *21*(3), 260-263.
- Awofeso, N., Schelokova, I., & Dalhatu, A. (2008). Training of front-line health workers for tuberculosis control: Lessons from Nigeria and Kyrgyzstan. *Human Resources for Health*, *6*(20), 1-9.

- Ayele, W., Neill, S., Zinsstag, J., Weiss, M., & Pavlik, I. (2004). Bovine tuberculosis: An old disease but a new threat to Africa. *The International Journal of Tuberculosis and Lung Disease*, 8(8), 924-937.
- Barnes, D. S. (2000). Historical perspectives on the etiology of tuberculosis. *Microbes and Infection*, 2(4), 431-440.
- Basu, M., Sinha, D., Das, P., Roy, B., Biswas, S., & Chattopadhyay, S. (2013). Knowledge and practice regarding pulmonary tuberculosis among private practitioners. *Indian Journal of Community Health*, 25(4), 403-412.
- Bati, J., Legesse, M., & Medhin, G. (2013). Community's knowledge, attitudes and practices about tuberculosis in Itang special district, Gambella region, South Western Ethiopia. *BMC Public Health*, 13(1), 1-9.
- Beals, F. (2007). The risk of bovine tuberculosis from raw milk consumption with A focus on Michigan in wise traditions in food, farming and the healing arts, the quarterly magazine of the Weston A. Price Foundation.
- Brosch, R., Gordon, S. V., Marmiesse, M., Brodin, P., Buchrieser, C., Eiglmeier, K., Cole, S. T. (2002). A new evolutionary scenario for the *Mycobacterium tuberculosis* complex. *Proceedings of the National Academy of Sciences of the United States of America*, 99(6), 3684-3689.
- Cadmus, S., Palmer, S., Okker, M., Dale, J., Gover, K., Smith, N., Gordon, S. V. (2006). Molecular analysis of human and bovine tubercle bacilli from a local setting in Nigeria. *Journal of Clinical Microbiology*, 44(1), 29-34.
- California Department of Food and Agriculture (2015). Bovine Tuberculosis Information for Livestock Producers. Retrieved February 6, 2016, from: https://www.cdfa.ca.gov/ahfss/animal_health
- Cambridge Dictionaries Online (2016). Retrieved February 2, 2016, from: <http://dictionary.cambridge.org/dictionary/english>
- Centers for Disease Control and Prevention (2006). *Morbidity and mortality weekly report*. (No. 55 (No.RR-9)). U.S. Department of Health and Human Services, Atlanta: Center for Health Information and Service, CDC.
- Christopher, O., & Bosedede, I. (2010). Health seeking behaviour of tuberculosis patients in Ekiti State, Nigeria. *Studies on Ethno-Medicine*, 4(3), 191-197.
- Cosivi, O., Grange, J. M., Daborn, C., Raviglione, M. C., Fujikura, T., Cousins, D., Meslin, F. (1998). Zoonotic tuberculosis due to *Mycobacterium bovis* in developing countries. *Emerging Infectious Diseases*, 4(1), 59.
- Courtenay, W. H. (2000). Constructions of masculinity and their influence on men's well-being: A theory of gender and health. *Social Science & Medicine*, 50(10), 1385-1401.

- Cousins D. V., Francis B. R. and Gow B. L. (1989). Advantages of a new agar medium in the primary isolation of *Mycobacterium bovis*. *Veterinary microbiology*. (pp. 89-95)
- Daniel, T. M. (2006). The history of tuberculosis. *Respiratory Medicine*, 100(11), 1862-1870.
- Depra, . (2013). Department for environment, food and rural affairs, Bovine tuberculosis: The tuberculin test. www.depra.gov.uk.
- Erah, P., & Ojieabu, W. (2009). Success of the control of tuberculosis in Nigeria: A review. *International Journal of Health Research*, 2(1), 3-14.
- Fatiregun, A., & Ejeckam, C. (2010). Determinants of patient delay in seeking treatment among pulmonary tuberculosis cases in a government specialist hospital in Ibadan, Nigeria. *Tanzania Journal of Health Research*, 12(2), 113-120.
- FMOH. (2010). *National tuberculosis and leprosy control programme (NTBLCP) worker's manual*. (No. Revised 5th Edition). Abuja, Nigeria: Department of Public Health, Federal Ministry of Health, Nigeria.
- Getahun, H., Gunneberg, C., Granich, R., & Nunn, P. (2010). HIV Infection—Associated tuberculosis: The epidemiology and the response. *Clinical Infectious Diseases*, 50(Supplement 3), S201-S207.
- Grace, D., Dipeolu, M., Olawoye, J., Ojo, E., Odebode, S., Agbaje, M., Randolph, T. (2012). Evaluating a group-based intervention to improve the safety of meat in Bodija market, Ibadan, Nigeria. *Tropical Animal Health and Production*, 44(1), 61-66.
- Grange, J. (2001). *Mycobacterium bovis* infection in human beings. *Tuberculosis*, 81(1), 71-77.
- Gusau Modern Abattoir. (2014). *Daily slaughter record*, Gusau Modern Abattoir, Ministry of Animal Health and Livestock Development. Unpublished manuscript.
- Gutierrez, M. C., Brisse, S., Brosch, R., Fabre, M., Omaïs, B., Marmiesse, M., Vincent, V. (2005). Ancient origin and gene mosaicism of the progenitor of *Mycobacterium tuberculosis*. *PLoS Pathogens*, 1(1), 55-61.
- Hambolu, D., Freeman, J., & Taddese, H. B. (2013). Predictors of bovine TB risk behaviour amongst meat handlers in Nigeria: A cross-sectional study guided by the health belief model. *PloS One*, 8(2), e56091, 1-9.
- Hosmer, D. W., & Lemeshow, S. (Eds.). (1989). *Applied logistic regression*. N Y: John Wiley & Sons Inc, New York.

- Ibrahim, L. M., Hadjia, I. S., Nguku, P., Waziri, N. E., Akhimien, M. O., Patrobas, P., & Nsubuga, P. (2014). Health care workers' knowledge and attitude towards TB patients under direct observation of treatment in Plateau State Nigeria, 2011. *The Pan African Medical Journal*, 18(Suppl 1):8, 1-4.
- Ibrahim, S., Cadmus, S., Umoh, J., Ajogi, I., Farouk, U., Abubakar, U., & Kudi, A. (2012). Tuberculosis in humans and cattle in Jigawa State, Nigeria: Risk factors analysis. *Veterinary Medicine International*, 2012. 1-5. doi:10.1155/2012/865924
- Ismail, A., & Josephat, P. (2014). Knowledge and perception on tuberculosis transmission in Tanzania: Multinomial logistic regression analysis of secondary data. *Tanzania Journal of Health Research*, 16(1). doi: <http://dx.doi.org/10.4314/thrb.v16i1.5>
- Jamison, D. T. (2006). *Disease and mortality in Sub-Saharan Africa (2nd edition)*. World Bank Publications.
- Jekel, J. F., Katz, D. L., Elmore, J. G., & Wild, D. (2007). *Epidemiology, biostatistics and preventive medicine*. Elsevier Health Sciences.
- Kaufmann, S. H., & Gengenbacher, M. (2012). Recombinant live vaccine candidates against tuberculosis. *Current Opinion in Biotechnology*, 23(6), 900-907.
- Kaufmann, S. H., Hussey, G., & Lambert, P. (2010). New vaccines for tuberculosis. *The Lancet*, 375(9731), 2110-2119.
- Legesse, M., Ameni, G., Mamo, G., Medhin, G., Shawel, D., Bjune, G., & Abebe, F. (2010). Knowledge and perception of pulmonary tuberculosis in pastoral communities in the middle and lower Awash valley of Afar region, Ethiopia. *BMC Public Health*, 10, 187-2458-10-187.
- Lemeshow, S., Hosmer, D. W., Klar, J., & Lwanga, S. K. (1990). Adequacy of sample size in health studies.
- Longo, D., Fauci, A., Kasper, D., Hauser, S., Jameson, J., & Loscalzo, J. (2012). Infectious Diseases: Mycobacterial Diseases (Tuberculosis). *Harrison's Principles of Internal Medicine, 18th Edition. Vol. 1: 8:8(165)*. McGraw Hill Professional.
- LoBue, P., Enarson, D., & Thoen, C. (2010). Tuberculosis in humans and animals: An overview [serialised article. tuberculosis: A re-emerging disease in animals and humans. number 1 in the series]. *The International Journal of Tuberculosis and Lung Disease*, 14(9), 1075-1078.
- Macmillan Dictionary. (2015). *Macmillan publishers limited*. Retrieved August 10, 2015, from <http://www.macmillandictionary.com/dictionary/british>

- Maher, D., Chaulet, P., Spinaci, S., & Harries, A. (1997). Treatment of tuberculosis: Guidelines for national programmes. *Treatment of Tuberculosis: Guidelines for National Programmes (2nd Edition)*, 1-77.
- Mandell, G. E., Gerald L., Bennett, J. E. (2010). *Principles and practices of infectious diseases (7th edition)*, 2(250), 3129-3163. Churchill Livingstone Elsevier.
- Mangtani, P., Abubakar, I., Ariti, C., Beynon, R., Pimpin, L., Fine, P. E., Sterne, J. A. (2014). Protection by BCG vaccine against tuberculosis: A systematic review of randomized controlled trials. *Clinical Infectious Diseases : An Official Publication of the Infectious Diseases Society of America*, 58(4), 470-480.
- Mushtaq, M., Majrooh, M., Ahmad, W., Rizwan, M., Luqman, M., Aslam, M., Shad, M. (2010). Knowledge, attitudes and practices regarding tuberculosis in two districts of Punjab, Pakistan. *The International Journal of Tuberculosis and Lung Disease*, 14(3), 303-310.
- National Tuberculosis Controllers Association & Centers for Disease Control and Prevention (CDC). (2005). Guidelines for the investigation of contacts of persons with infectious tuberculosis. Recommendations from the national tuberculosis controllers association and CDC. *MMWR.Recommendations and Reports : Morbidity and Mortality Weekly Report.Recommendations and Reports / Centers for Disease Control*, 54(RR-15), 1-47.
- OIE. (2009). Bovine tuberculosis. Retrieved from: www.oie.int/fileadmin/home/eng/health
- Okoh, A. C. (2009). Prevention and Control of Tuberculosis. *Benin Journal of Postgraduate Medicine*, 11(1), 97-99.
- Okuonghae, D., & Omosigho, S. (2010). Determinants of TB case detection in Nigeria: A survey. *Global Journal of Health Science*, 2(2), p123.
- Onlinenigeria. (2013). CommunityPortalof Nigeria, Zamfara State natural resources and potential for development. Retrieved from <http://www.onlinenigeria.com/links/zamfarastateadv.asp?blurb=397>
- Orcau, À., Caylà, J. A., & Martínez, J. A. (2011). Present epidemiology of tuberculosis. prevention and control programs. *Enfermedades Infecciosas y Microbiologia Clinica*, 29, 2-7.
- O'Reilly, L. M., & Daborn, C. (1995). The epidemiology of *Mycobacterium bovis* infections in animals and man: A review. *Tubercle and Lung Disease*, 76, 1-46.
- Orogade, A., Ahmed, P., Onazi, S., Abubakar, U., & Isa, H. (2013). BCG status in children with tuberculosis: A multicenter study in Northern Nigeria. *Journal of Medicine in the Tropics*, 15(2), 126-130.

- Otu, A. A. (2013). A review of the national tuberculosis and leprosy control programme (NTBLCP) of Nigeria: Challenges and prospects. *Annals of Tropical Medicine and Public Health*, 6(5), 491-500.
- Otupiri, E., Adam, M., Laing, E., & Akanmori, B. (2000). Detection and management of zoonotic diseases at the Kumasi slaughterhouse in Ghana. *Acta Tropica*, 76(1), 15-19.
- Peat, J., & Barton, B. (2005). Continuous variables: Descriptive statistics. *Medical statistics: A guide to data analysis & critical appraisal* (pp. 31-32).
- Petrie, K. J., Jago, L. A., & Devcich, D. A. (2007). The role of illness perceptions in patients with medical conditions. *Current Opinion in Psychiatry*, 20(2), 163-167.
- Radostits, O. M., Gay, C. C., Blood, D. C., & Henscheliff, K. W. (2000). *Veterinary medicine: A text book of disease of cattle, sheep, pigs, goats and horses (9th edition)*, Harcourt publisher ltd., London.
- Rodríguez, E., Sánchez, L., Pérez, S., Herrera, L., Jiménez, M., Samper, S., & Iglesias, M. (2009). Human tuberculosis due to *Mycobacterium bovis* and *M. caprae* in Spain, 2004-2007. *The International Journal of Tuberculosis and Lung Disease*, 13(12), 1536-1541.
- Rosiji, C., & Adesokan, H. (2012). Measures taken for bovine tuberculosis prevention among butchers in two major abattoirs in Ibadan, South-Western Nigeria. *Global Veterinaria*, 9(1), 107-112.
- Shitaye, J., Tsegaye, W., & Pavlik, I. (2007). Bovine tuberculosis infection in animal and human populations in Ethiopia: A review. *Veterinarni Medicina-Praha*, 52(8), 317-332.
- Stead, W., Aufderheide, A., Buikstra, J., & Holcomb, T. (1995). When did mycobacterium tuberculosis infection first occur in the new world. *American Journal of Respiratory and Critical Care Medicine*, 151, 1267-1268.
- Suleiman, M., Sahal, N., Sodemann, M., Elsony, A., & Aro, A. R. (2014). Tuberculosis awareness on Gezira, Sudan: Knowledge, attitudes and practice case-control survey. *Eastern Mediterranean Health Journal*, 20(2), 120-129.
- Swai, E. S., Schoonman, L., & Daborn, C. (2010). Knowledge and attitude towards zoonoses among animal health workers and livestock keepers in Arusha and Tanga, Tanzania. *Tanzania Journal of Health Research*, 12(4), 272-277.
- Tameris, M. D., Hatherill, M., Landry, B. S., Scriba, T. J., Snowden, M. A., Lockhart, S., Hanekom, W. A. (2013). Safety and efficacy of MVA85A, a new tuberculosis vaccine, in infants previously vaccinated with BCG: A randomised, placebo-controlled phase 2b trial. *The Lancet*, 381(9871), 1021-1028.

- Taylor, G. M., Murphy, E., Hopkins, R., Rutland, P., & Chistov, Y. (2007). First report of *Mycobacterium bovis* DNA in human remains from the Iron age. *Microbiology*, 153(4), 1243-1249.
- Tohen, C. O., LoBue, P. A., Enarson, D. A., Kaneene, J. B., & de Kantor, I. N. (2009). Tuberculosis: A re-emerging disease in animals and humans. *Veternaria Italiana*, 45(1), 135-181.
- Tohen, C., LoBue, P., & De Kantor, I. (2006). The importance of *Mycobacterium bovis* as a zoonosis. *Veterinary Microbiology*, 112(2), 339-345.
- Tobin, E. A., Okojie, P. W., & Isah, E. C. (2013). Community knowledge and attitude to pulmonary tuberculosis in rural Edo State, Nigeria. *Annals of African Medicine*, 12(3), 148-154.
- United States Department of Agriculture (2014). Animal and Plant Health Inspection Service (APHIS) Factsheet: *Questions and Answers: Preventing Bovine Tuberculosis*. Retrieved February 6, 2016, from: https://www.aphis.usda.gov/publications/animal_health
- Wammanda, R., Gambo, M., & Abdulkadir, I. (2004). Short communication: Age at BCG administration during routine immunization. *Journal of Community Medicine and Primary Health Care*, 16(1), 33-35.
- Wedlock, D. N., Skinner, M. A., de Lisle, G. W., & Buddle, B. M. (2002). Control of *Mycobacterium bovis* infections and the risk to human populations. *Microbes and Infection*, 4(4), 471-480.
- Whelan, A. O., Clifford, D., Upadhyay, B., Breadon, E. L., McNair, J., Hewinson, G. R., & Vordermeier, M. H. (2010). Development of a skin test for bovine tuberculosis for differentiating infected from vaccinated animals. *Journal of clinical microbiology*, 48(9), 3176-3181.
- White, Z. N. (2011). Survey on the knowledge, attitudes and practices on tuberculosis (TB) among health care workers in Kingston & St. Andrew, Jamaica. Unpublished manuscript.
- Whitehead, D. (2001). Health education, behavioural change and social psychology: Nursing's contribution to health promotion? *Journal of Advanced Nursing*, 34(6), 822-832.
- WHO. (2008). *Advocacy, communication and social mobilization for TB control: A guide to developing knowledge, attitude and practice surveys*. (No. WHO/HTM/STB/2008.46). Switzerland: World Health Organisation.
- WHO. (2014). *Global tuberculosis report 2014* World Health Organization.
- Woith, W. M., & Larson, J. L. (2008). Delay in seeking treatment and adherence to tuberculosis medications in Russia: A survey of patients from two clinics. *International Journal of Nursing Studies*, 45(8), 1163-1174.

World Health Organization. (2008). *Implementing the WHO stop TB strategy: A handbook for national TB control programmes* World Health Organization.

World Health Organization. (2009). *Global tuberculosis control: Epidemiology, strategy, financing: WHO report 2009* World Health Organization.

World Health Organization. (2010). *Treatment of tuberculosis: Guidelines* World Health Organization.

Zamfara State Government. (2010). Zamfara state, Nigeria. Retrieved from www.nigeriagallery.com/Nigeria/States_Nigeria/Zamfara_State.html

