

UNIVERSITI PUTRA MALAYSIA

CYTOTOXICITY EFFECT OF COCOA (Theobroma cacao L.) POLYPHENOL EXTRACT ON MCF-7 CELLS, AND MODE OF CELL DEATH

HAZIRAH BINTI ABDUL RADZAK

FPSK(M) 2014 3



CYTOTOXICITY EFFECT OF COCOA (*Theobroma cacao* L.) POLYPHENOL EXTRACT ON MCF-7 CELLS, AND MODE OF CELL DEATH



HAZIRAH BINTI ABDUL RADZAK

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfillment of the Requirements for the Degree of Master of Science.

May 2014

COPYRIGHT

All materials contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright matrials of Universiti Putra Malaysia unless otherwise stated. Use may be made of any materials contained within thesis for non-commercial purposes from the copyright holder. Commercial use of materials may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree of Master of Science.

CYTOTOXICITY EFFECT OF COCOA (Theobroma cacao L.) POLYPHENOL EXTRACT ON MCF-7 CELLS, AND MODE OF CELL DEATH

By

HAZIRAH BINTI ABDUL RADZAK

May 2014

Chairman : Abdah binti Md Akim, PhD

Faculty : Medicine and Health Sciences

The Incidence of breast cancer in Malaysia is alarming and it increased every year. It is one of the common causes of deaths among cancer patients in women. Despite of several drugs have been formulated for breast cancer treatment, these drugs can cause undesired side effects to the breast cancer patients. Therefore, scientists are searching for potential cancer chemopreventive and chemotherapeutic agents through dietary approaches. Cocoa (Theobroma cacao L.) is rich in specific antioxidants such as catechin, epicatechin and procyanidins. Previous research exhibited that cocoa possessed antioxidant and cytotoxic properties. The objective of the present study was to investigate the cytotoxicity effect of cocoa polyphenol extract (CPE) towards MCF-7 cells and its effect on mode of cell death. The phenolic constituents of CPE were evaluated by phytochemical screening, HPLC profiling and total phenolic content (TPC) assay. The antioxidant activity of CPE was determined using DPPH radical scavenging and ferric reducing antioxidant power (FRAP) assay. Cell viability was measured using MTT assay. The morphological alteration was observed using inverted light and fluorescence microscope (acridine orange/ propidium iodide dual staining). The mode of cell death was investigated using annexin V FITC-PI and DNA fragmentation assay. The apoptotic marker of cell death was carried out using p53 and caspase-9 ELISA kits. The phytochemical screening and HPLC profiling exhibited CPE contained phenolic compound particularly saponins, flavonoids and condensed tannins. TPC, DPPH IC₅₀ and FRAP value of CPE were 13558.99±420.10 mg GAE/100g dry weight of sample, 14.73±1.47 µg/ml and 2130.33±2.33 µM FE/1 mg dry weight of sample respectively. CPE exhibited highest cytotoxicity towards MCF-7 cells with the lowest IC₅₀ value (3 mg/ml) and exhibited significant difference (p<0.05) compared to other cancer cell lines. The difference of IC_{50} value was significant (p<0.05) between 24 h (4.50±0.50 mg/ml), 48 h (2.85±0.20 mg/ml) and 72 h (1.60±0.10 mg/ml). The morphological alteration of MCF-7 cells upon 48 h CPE treatment showed apoptosis and necrosis characteristics including cell membrane blebbing, cell

ii

shrinkage, nuclear condensation, apoptotic bodies and cell membrane rupture. The cell cycle analysis revealed that CPE was able to cause mild cell cycle arrest at G0/G1 phase and also induced sub-G1 peak, indicating apoptosis. Annexin V-PI assay proved that CPE induced early and late apoptosis in treated MCF-7 cells. The DNA fragmentation assay confirmed that DNA fragmentation had occurred during apoptosis in treated MCF-7 cells. The expression level of p53 and caspase-9 were increased upon CPE treatment in MCF-7 cells indicating that apoptosis was executed via mitochondria pathway. In conclusion, these findings suggested that CPE demonstrated cytotoxicity effect towards MCF-7 cells through inhibition of cell proliferation by arresting G0/G1 phase and apoptosis execution via p53 and caspase-9 activation. Based on the current findings, further research is required to develop CPE as chemopreventive agents for breast cancer.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Sarjana Sains.

KESAN SITOTOKSISITI EKSTRAK (Theobroma cacao L.) POLIFENOL KOKO KE ATAS SEL MCF-7, DAN MOD KEMATIAN SEL

Oleh

HAZIRAH BINTI ABDUL RADZAK

Mei 2014

Pengerusi : Abdah binti Md Akim, PhD

Fakulti : Fakulti Perubatan dan Sains Kesihatan

Insiden kanser payudara di Malaysia semakin membimbangkan dan meningkat setiap tahun. Ia merupakan satu daripada penyebab umum kematian di kalangan pesakit kanser wanita. Walaupun pelbagai ubat telah diformulasi untuk rawatan kanser payudara, ubat-ubat ini akan menyebabkan kesan sampingan yang tidak diingini kepada para pesakit kanser payudara. Oleh itu, para saintis sedang mencari agen yang berpotensi sebagai pencegah kanser dan perawatan kanser melalui pendekatan pemakanan. Koko (*Theobroma cacao* L.) kaya dengan kandungan antioksidan yang spesifik seperti katekin, epikatekin dan prosianidin. Penyelidikan sebelum ini menunjukkan koko mempunyai ciri-ciri antioksidan dan sitotoksik. Objektif kajian ini untuk menyiasat kesan sitotoksisiti ekstrak polifenol koko (CPE) terhadap sel MCF-7 dan kesannya ke atas mod kematian sel. Kandungan fenolik CPE telah diuji dengan saringan fitokimia, pemprofilan HPLC dan asai jumlah kandungan fenolik (TPC). Aktiviti antioksidan yang terdapat pada CPE ditentukan menggunakan asai penyahbebas radikal DPPH dan kuasa penurunan ferik (FRAP). Viabiliti sel diukur menggunakan asai MTT. Perubahan morfologi diperhatikan menggunakan mikroskop inversi cahaya dan fluoressen (dwi perwarnaan akridina oren/propidium iodida). Mod kematian sel disiasat menggunakan asai annexin V FITC-PI dan fragmentasi DNA. Mekanisme kematian sel dijalankan menggunakan kit ELISA p53 dan caspase-9. Saringan fitokimia dan pemprofilan HPLC menunjukkan CPE mengandungi bahan fenolik terutamanya saponins, flavonoid dan tannin tersejat. TPC, nilai IC₅₀ untuk asai DPPH dan FRAP masing-masing ialah 13558.99±420.10 mg GAE/100g berat kering sampel, 14.73±1.47 µg/ml dan 2130.33±2.33 µM FE/1 mg berat kering sampel. CPE menunjukkan sitotoksik yang tertinggi terhadap sel MCF-7 dengan nilai IC₅₀ yang terendah (3 mg/ml) dan menunjukkan perbezaan yang signifikan (p<0.05) berbanding dengan kanser sel yang lain. Perbezaan nilai IC₅₀ adalah signifikan (p<0.05) diantara 24 jam (4.50±0.50 mg/ml), 48 jam (2.85±0.20 mg/ml) dan 72 jam (1.60±0.10 mg/ml). Perubahan morfologi sel MCF-7 sebaik sahaja rawatan CPE menunjukkan ciri-ciri apoptosis dan nekrosis termasuklah pembengkakkan membran sel, pengecutan sel, kondensasi nukleus dan sel membran ruptur. Analisis kitar sel mendedahkan CPE boleh menyebabkan sedikit penahanan kitar sel pada fasa G0/G1 dan juga merangsang fasa sub-G1 yang menandakan apoptosis. Asai annexin V-PI membuktikan bahawa CPE telah merangsang apoptosis peringkat awal dan apoptosis peringkat lewat pada sel MCF-7 yang dirawat. Asai fragmentasi DNA mengesahkan fragmentasi DNA telah berlaku semasa apoptosis pada sel MCF-7 yang dirawat. Peringkat ekspresi p53 dan kaspase-9 meningkat sebaik sahaja rawatan CPE pada sel MCF-7 menandakan bahawa apoptosis dilaksanakan melalui aliran mitokondria. Secara kesimpulannya, penemuan-penemuan diatas ini membuktikan CPE menunjukkan kesan sitotoksisiti terhadap sel MCF-7 melalui perencatan pembahagian sel dengan penahanan pada fasa G0/G1 dan pelaksanaan apoptosis melalui pengaktifan p53 dan kaspase-9. Berdasarkan pada penemuan-penemuan ini, penyelidikan lanjutan diperlukan untuk menjadikan CPE sebagai agen pencegah kanser untuk kanser payudara.

AKNOWLEDGEMENTS

In the name of Allah, the Most Gracious, the Most Merciful. Praise to Allah for the blessing, courage and guidance that have been given to me in completing my Master research project.

First of all, I wish to show my deepest gratitude and appreciation to my upmost beloved supervisor, Dr Abdah Md Akim. She gave me a lot of knowledge about cell culture and some techniques to suceed the works. She also gave me emotional support and some useful suggestions when I met with some difficulties during the project. Honestly, Dr Abdah is a very nice lecturer and supervisor that I ever met before. Also, not forget to my respectful co-supervisor, Dr Huzwah Khaza'ai and Associate Professor Dr Cheah Yoke Kqueen for their great guidance, encouragement and support throughout this whole research project. I maybe cannot finish my work without their great and brilliant suggestions.

Apart from that, I would like to express my appreciation to my beloved parents, Mrs Norasiah and Mr Abdul Radzak for being there for me and giving me support, prayers and love advice. Their love have given me strength to accomplish my work.

My sincere gratitude and appreciation also forwarded to my best friends, Nurhayatie Sajali, Thilaga Rati, Nurul Akmaryanti, Siti Sarah, Siti Aisyah, Mr Zainal, Miss Armania, Mr Foo Jhi Biao, Siti Syariah, Hamizah Sulaiman and Siti Rukhaiyah who always helping me in trouble, giving me motivations when I feel give up, sharing hardness, happiness and sadness in research and life together. Thanks make my research life meaningful and may Allah give excellent in your Master and PhD studies.

I would like to thank lab personals in Biochemistry, Cell Signaling and Immunology Lab, Mrs Fatimah, Miss Norellina, Mrs Intan, Mr Zul, Mrs Ayien and Mrs Marsyitah for giving permission to use any equipment and helping me to use some laboratory equipment in the lab for my research project.

At last, I would like to express my thankful again to all the people who directly and indirectly related to my research project for helping me to suceed in this chapter of my life.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfiillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Abdah binti Md Akim, PhD

Senior Lecturer Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Chairman)

Huzwah binti Khaza'ai, PhD

Senior Lecturer Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Member)

Cheah Yoke Kqueen, PhD

Associate Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Member)

BUJANG BIN KIM HUAT, PhD

Professor Deputy Dean School of Graduate Studies Universiti Putra Malaysia

Date:

DECLARATION

Declaration by Graduate Student:

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not submitted previously or concurrently for other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, preceedings, popular writtings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature:

Date:

Name and Matric No: Hazirah Binti Abdul Radzak (GS28709)

Declaration by Members Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature :	Signature :
Name of	Name of
Chairman of	Member of
Supervisory	Supervisory
Committee : Dr Abdah Binti Md Akim	Committee: Associate Professor
	Dr Cheah Yoke

Kqueen

Signature :_____ Name of Members of Supervisory

Committee : Dr Huzwah Binti Khaza'ai

TABLE OF CONTENTS

ABSTR ABSTR ACKNO APPRO DECLA LIST OI LIST OI LIST OI	ACT AK OWLEDGEMENTS VAL RATION F TABLES F FIGURES F ABBREVIATIONS	Page ii iv vi vii ix xv xviii xx
CHAP1	LK	1
1	1.1 Background of study 1.2 Problem statement 1.3 Research objectives 1.3.1 General objective 1.3.2 Specific objectives 1.4 Research hypothesis	1 2 3 3 3 3 3
2	LITEDATUDE DEVIEW	Λ
4	2 1 Cancor	4
	2.1 Callet 2.2 Epidemiology of concer	4
	2.2 Epidemiology of cancer	4
	2.5 Breast cancer	5
	2.4 Incidence of breast cancer	2
	2.5 Risk factors	5
	2.5.1 Family history and genetic	6
	2.5.2 Hormone Replacement Therapy (HRT)	6
	2.5.3 Lifestyle	6
	2.5.4 Enviromental	7
	2.6 Antioxidants	7
	2.7 Polyphenol	8
	2.8 Cocoa polyphenol	10
	2.9 Cancer cell lines	13
	2.9.1 HepG2	13
	2.9.2 HT-29	14
	2.9.3 HeLa	14
	2.9.4 MCF-7	14
	2.9.5 MDA-MB-231	15
	2.10 High Performance Liquid Chromatography (HPLC)	15
	2.11 Cell cycle	15
	2.12 Apoptosis	17
	2.13 Necrosis	19
	2.14 Tumor protein suppressor gene (p53)	20
	2.15 Cysteine-aspartic protease-9 (Caspase-9)	22
3	MATERIALS AND METHODOLOGY	23
	3.1 Materials	23
	3.2 Compound	23

3.3 Phytochemical screening	23
3.3.1 Test for Alkaloids	24
3.3.2 Test for Saponins	24
3.3.3 Test for Flavonoids	24
3.3.4 Test for Tannins and Polyphenolic compounds	24
3.3.5 Test for Triterpenes/Steroid	24
3.4 High Performance Liquid Chromatography (HPLC)	24
profiling of CPE	
3.5 Total Phenolic Content	25
3.6 Antioxidant assay	25
3.6.1 DPPH radical scavenging activity	25
3.6.2 Ferric Reducing Antioxidant Power (FRAP) assay	26
3.7 Cell culture	26
3.7.1 Media preparation	26
3.7.2 Cells recovery and thawing	27
3.7.3 Cell culturing	27
3.7.4 Changing media	27
3.7.5 Subculture of cells	27
3.7.6 Cells cryopreservation	28
3.7.7 Cell counting	28
3.8 Cytotoxicity assay of CPE on various cancer cell lines	29
3.8.1 Cell seeding	29
3.8.2 Treatment with CPE	29
3.8.3 MTT assay	29
3.8.4 Cytotoxicity assay of CPE on MCE-7 at different	29
time points	27
3 9 Morphological studies	30
3.9.1 Cell seeding	30
3.9.2 Treatment with CPE	30
393 Inverted microscope	30
3.9.4 Acridine Orange and Propidium Iodide	30
(AO/PI)dual staining	50
3 10 Flow cytometry	31
3 10 1 Cell seeding	31
3 10 2 Treatment with CPE	31
3 10 3 Cell cycle assay	31
3.10.4 Annexin V FITC/PI assay	31
3.11 DNA fragmentation assay	32
3.11.1 Cell seeding	32
3 11.2 Treatment with CPE	32
3 11 3 DNA extraction	32
3 11 4 Gel electrophoresis	33
3 12 Determination of apoptotic marker	34
(Human p53 and Caspase-9)	51
3.12.1 Cell seeding	34
3.12.2 Treatment with CPE	34
3.12.2 Fleatment with CFE	34 34
3.12.5 Con rysis 3.12.4 Quantification of total protein using Bradford	34
method	57
3.12.5 FLISA of human n53	35
5.12.3 LLISA OF HUIHAII p55	55

	3.12.6 ELISA of human caspase-9	35
	3.13 Statistical Analysis	36
4	RESULTS	38
	4.1 Phytochemical screening of CPE	38
	4.2 Qualitative analysis of High Performance Liquid	39
	Chromatography (HPLC) profiling of CPE	
	4.3 Total phenolic content	42
	4.4 Antioxidant assay	43
	4.4.1 DPPH assay	43
	4.4.2 FRAP assay	44
	4.5 MTT assay	45
	4.5.1 Screening of different cancer cell lines	45
	4.5.2 Cytotoxicity assay of MCF-7 cells	47
	4.6 Morphological studies of MCF-7 cells	49
	4.6.1 Inverted microscope	49
	4.6.2 AO/PI dual staining assay	52
	4.7 Flow cytometry analysis	54
	4.7.1 Cell cycle assay	54
	4.7.2 Annexin V FITC/PI assay	57
	4.8 DNA fragmentation	00 61
	4.9 Expression of human asspace 0 level	01 62
	4.10 Expression of numan caspase-9 lever	02
5	DISCUSSION	63
	5.1 Phytochemical screening and HPLC profiling of CPE	63
	5.2 Total phenolic content of CPE	63
	5.3 Antioxidant activity of CPE	64
	5.4 Cytotoxicity study of CPE	65
	5.5 Morphological study of MCF-7 cells upon CPE	66
	5.6 Cell cycle analysis of MCE-7 cells against CPE	67
	treatment	07
	5.7 Annexin V FITC/PI analysis of MCF-7 cells against	68
	CPE treatment	00
	5.8 DNA fragmentation study of MCF-7 cells against CPE	68
	treatment	
	5.9 The expression of human p53 level in MCF-7 cells	69
	upon CPE treatment	
	5.10 The expression of human caspase-9 level in MCF-7	69
	cells upon CPE treatment	
6	SUMMARY, CONCLUSION AND	71
	RECOMMENDATIONS FOR FUTURE RESEARCH	
	6.1 Summary and conclusion	71
	6.2 Recommendations for future research	72

REFERENCES/BIBLIOGRAPHY	74
APPENDICES	91
BIODATA OF STUDENT	107
LIST OF PUBLICATIONS	108



LIST OF TABLES

Table		Page
3.1	Materials for preparing complete media.	26
4.1	Phytochemical constitutuent of CPE	38
4.2	TPC value of trolox, ascorbic acid and CPE	42
4.3	FRAP value of trolox, ascorbic acid and CPE.	44
4.4	IC ₅₀ value of CPE at 24, 48 and 72 h	47
4.5	Percentage of viable, apoptosis and necrosis of MCF-7 cells at different concentration of CPE incubated for 48 h.	53
4.6	Percentage of MCF-7 cells in Sub-G1, G0/G1, S and G2/M phase at different concentration of CPE incubated for 48 h.	56
4.7	Percentage of viable, early apoptosis, late apoptosis and necrosis of MCF-7 cells at different concentration of CPE incubated for 48 h.	59
4.8	Percentage of human p53 in MCF-7 cells treated at different concentration of CPE incubated for 48 h.	61
4.9	Percentage of human caspase-9 in MCF-7 cells treated at different concentration of CPE incubated for 48 h.	62
D.1	Analysis of variance (ANOVA) table for TPC values of trolox, ascorbic acid and CPE.	100
D.2	Analysis of variance (ANOVA) table for DPPH IC_{50} values of trolox, ascorbic acid and CPE.	100
D.3	Analysis of variance (ANOVA) table for FRAP values of trolox, ascorbic acid and CPE.	100
D.4	Analysis of variance (ANOVA) table for IC_{50} value of CPE in cells screening for cytotoxicity assay at 48 h.	101
D.5	Analysis of variance (ANOVA) table for IC_{50} value of CPE in MCF-7 cells treated with 1-10 mg/ml of CPE for 24, 48 and 72 h.	101

	D.6	Analysis of variance (ANOVA) table for viable cells percentages upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in AOPI assay.	102
	D.7	Analysis of variance (ANOVA) table for apoptotic cells percentages upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in AOPI assay.	102
	D.8	Analysis of variance (ANOVA) table for necrotic cells percentages upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in AOPI assay.	102
	D.9	Analysis of variance (ANOVA) table for cells percentages in sub-G1 phase upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in cell cycle assay.	103
	D.10	Analysis of variance (ANOVA) table for cells percentages in G0/G1 phase upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in cell cycle assay.	103
	D.11	Analysis of variance (ANOVA) table for cells percentages in S phase upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in cell cycle assay.	103
	D.12	Analysis of variance (ANOVA) table for cells percentages in G2/M phase upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in cell cycle assay.	104
	D.13	Analysis of variance (ANOVA) table for viable cells percentages upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in Annexin V FITC/PI assay.	104
	D.14	Analysis of variance (ANOVA) table for early apoptotic cells percentages upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in Annexin V FITC/PI assay.	104
	D.15	Analysis of variance (ANOVA) table for late apoptotic cells percentages upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in Annexin V FITC/PI assay.	105
	D.16	Analysis of variance (ANOVA) table for necrotic cells percentages upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE in Annexin V FITC/PI assay.	105
	D.17	Analysis of variance (ANOVA) table for concentration of human p53 upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE	105

xvi

D.18 Analysis of variance (ANOVA) table for concentration of human caspase-9 upon cells treatment with 0, 1, 2 and 3 mg/ml of CPE.

106



LIST OF FIGURES

Figure		Page
2.1	Chemicals structure of flavonoids, phenolic acids and polyphenolic amide.	9
2.2	Chemical structure of catechin, epicatechin and procyanidin B2.	10
2.3	Cocoa tree	11
2.4	Extrinsic and intrinsic pathway of apoptosis.	18
2.5	Morphological alteration of normal or cancer cell undergoes necrosis and apoptosis process.	19
2.6	Activation of p53-dependent growth arrest or apoptosis	21
3.1	Flow chart of overall study	37
4.1	HPLC spectrum of CPE at a) 254 nm, b) 280 nm and c) 300 nm using formic acid and acetonitrile as solvent.	40
4.2	HPLC spectrum of CPE and standard procyanidin B2 at 280 nm using formic acid and acetonitrile as solvent.	41
4.3	Scavenging activity percentage in trolox, ascorbic acid and CPE at different concentration (0-1000 μ g/ml).	43
4.4	Cell viability percentage in HepG2, HT- 29, HeLa, MDA-MB-231, MCF-7 and WRL-68 cell lines against CPE concentration (0.001-10 mg/ml) after 48 h treatment.	46
4.5	Percentage of cell viability in untreated and treated MCF-7 cells with different concentration of CPE (1-10 mg/ml) for 24, 48 and 72 h.	48
4.6	Morphological of MCF-7 cells upon treatment with CPE at the concentration of a) 0 mg/ml, b) 1 mg/ml, c) 2 mg/ml and d) 3 mg/ml for 48 h viewed under inverted light microscope (200x magnifications).	50

4.7	Morphological of MCF-7 cells upon treatment with CPE at the concentration of a) 0 mg/ml, b) 1 mg/ml, c) 2 mg/ml and d) 3 mg/ml for 48 h viewed under inverted light microscope (800x magnifications).	51
4.8	Morphological alteration of MCF-7 cells treated with CPE for 48 h at the concentration of a) 0 mg/ml, b) 1 mg/ml, c) 2 mg/ml and d) 3 mg/ml under fluorescence microscope (400x magnifications).	53
4.9	Cell cycle analysis of untreated and treated MCF-7 cells with different concentration of CPE for 48 h.	55
4.10	Flow cytometric analysis of PI-FITC to quantify CPE-induced apoptosis and necrosis in MCF-7 cells. Dot plots of MCF-7 treated with CPE at concentration of a) 0 mg/ml, b) 1 mg/ml, c) 2 mg/ml and d) 3 mg/ml for 48 h.	58
4.11	DNA fragmentation of MCF-7 cells incubated with different concentration of CPE for 48 h.	60
B.1	Gallic acid graph standard.	95
B.2	Ferric sulphate (FeSO ₄) graph standard.	95
B.3	Human p53 graph standard.	96
B.4	Human caspase-9 graph standard.	96
B.5	Bovine serum albumin (BSA) graph standard.	97
C.1	Cocoa fruits.	98
C.2	Cocoa beans.	98
C.3	Cocoa Polyphenol Extract.	99

LIST OF ABBREVIATIONS

AIP1	Actin-interacting protein 1
Akt	Protein kinase
ANOVA	Analysis of variance
AO	Acridine orange
Apaf-1	Apoptotic protease activation factor-1
ARF	Alternative reading frame
ASPP1/2	Apoptosis-stimulating protein of p53 1 and 2
ATCC	American Type Culture Collection
ATM	Ataxiatelangiectasia mutated
АТР	Adenosine triphosphate
Bak	Bcl-2 antagonist killer
Bax	Bcl-2-associated X protein
Bcl-2	B-cell-lymphoma 2
BRCA 1	Breast cancer type 1 susceptibility protein
BRCA 2	Breast cancer type 2 susceptibility protein
BSA	Bovine serum albumin
CADP	Collagen-ADP
cAMP	cyclic adenosine monophosphate
CARD	Caspase recruitment domain
Caspase	Cysteine-aspartic protease
CAT	Catalase
CDC	Cell division cycle
Cdc2	Cell division cycle 2
Cdc25	Cell division cycle 25

CDK	Cyclin-dependent kinase
CDKI	Cyclin-dependent kinase inhibitor
CEPI	Collagen epinephrine
CF	Cocoa rich fiber
chk2	Checkpoint kinase 2
cIAP	cellular Inhibitor of Apoptosis
CO ₂	Carbon dioxide
СРЕ	Cocoa Polyphenol Extract
CPF	Cocoa procyanidin fraction
Cu ⁺	Copper(I) ion
DAPI	4',6-diamidino-2-phenylindole
DBP	Diastolic blood pressure
DD	Death domain
DED	Death effector domain
DISC	Death-inducing signaling complex
DMEM	Dulbecco's Minimum Eagle Medium
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
DPPH	2,2-diphenyl-1-picrylhydrazyl
dATP	Deoxyadenosine triphosphate
DU145	Androgen nonresponsive prostate cancer cell line
EDTA	Ethylenediamine Tetraacetic Acid
ELISA	Enzyme-Linked Immunoabsorbent Assay
ER	Estrogen receptor

ERK/Cx43	Extracellular signal- regulated kinase/connexin 43
FADD	Fas-Associated protein with Death Domain
FBS	Fetal Bovine Serum
FE	Ferric equivalents
Fas/CD95	Cell death signaling receptor
Fe ²⁺	Ferric (II) ion
Fe ²⁺ -TPTZ	Ferrous-tripyridyltriazine
Fe ³⁺ -TPTZ	Ferric tripyridyltriazine
FFAs -	Free fatty acids
FITC	Fluorescein isothiocyanate
FMD	Flow mediated dilation
FRAP	Ferric reducing antioxidant power
FRIM	Forest Research Institute of Malaysia
GAE	Gallic acid equivalent
GJIC	Gap junctional intracellular communication
GPx	Glutathione peroxidase
GPIIb/IIa-act	Glycoprotein lib/IIIa receptor
GRx	Glutathione reductase
GSH	Glutathione
g	Gram
H_2O_2	Hydrogen peroxide
HCl	Hydrochloric acid
HDL	High-density lipoprotein
HeLa	Helacyton gartleri
HepG2	Hepatocellular carcinoma

xxii

HIPK2	Homedomain-interacting protein kinase 2	
HNE	4-hydroxynenal	
HPLC	High Performance Liquid Chromatography	
HRP	Horseradish peroxide	
HRT	Hormone Replacement Therapy	
HT-29	Human colon adenocarcinoma grade II cel line	
h	Hours	
IC ₅₀	Inhibition concentration 50% of cell viability	
IgG	Immunoglobulin G	
IU	International Unit	
i.d	Internal diameter	
ЈМҮ	Junction mediating and regulatory protein	
JNK	c-Jun N-terminal protein kinase	
kb	Kilobase	
kDa	Kilodaltons	
LDL	Low-density lipoprotein	
MCF-7	Michigan Cancer Foundation -7	
MCP-1	Monocyte chemoattractant protein-1	
MDA	Malondialdehyde	
MDA-MB-231	Mammary gland adenocarcinoma	

xxiii

Mammary gland adenocarcinoma

Mammary gland adenocarcinoma

Mouse doubling minute 2 homolog

Mitogen activated protein kinase kinase 4-c-Jun N-terminal protein kinase

MDA-MB-436

MDA-MB-468

MDM2

MKK4-JNK

MOMP	Mitochondrial outer membrane permeabilization		
MPTPs	Mitochondrion permeability transition pores		
MTS	3-(4,5-dimethylthiazol-2-yl)-5-(3- carboxymethoxyphenyl)-2-(4-sulfophenyl)- 2H-tetrazolium		
MTT	3-(4,5-dimethylthiazol-2-yl)-2,5 diphenyltetrazolium bromide		
mg	Milligram		
min	Minutes		
ml	Mililiter		
mM	Milimolar		
mm	Milimeter		
NCR	National Cancer Registry		
NF-κB	Nuclear factor-kB		
NO	Nitric oxide		
NOX-1	NADPH oxidase-1		
ng	Nanogram		
nm	Nanometer		
ОН∙	Hydroxyl radical		
РА	α-amylase		
РАН	Polycyclic aromatic hydrocarbon		
PARP	Poly (ADP-ribose) polymerase		
PBS	Phosphate buffer saline		
PC12	Pheocromocytoma		
PI	Propidium iodide		
PIG3	p53-inducible gene 3		

 \bigcirc

РКА	Protein Kinase A
PL	Pancreatic lipase
PLA ₂	Phospholipase A ₂
PolII	Polymerase II
pRb	Retinoblastoma protein
PS	Phosphatidylserine
Puma	p53-upregulated modulator of apoptosis
p38-MAPK-	p38 mitogen-activated-protein-kinase-
p53	Tumor suppressor gene
p63	Tumor protein 63
p73	Tumor protein 73
p107	Retinoblastoma-like protein 1
p130	Retinoblastoma-like protein 2
p53AIP1	p53-regulated apoptosis-inducing protein 1
P53DINP1	p53-dependent damage-inducible nuclear protein 1
RIP	Receptor-interacting protein
RLE	Rat liver epithelial
RNA	Ribonucleic acid
ROS	Reactive species oxygen
rpm	Revolutions per minute
RWEP-1	Normal prostate cancer cell line
SBP	Systolic blood pressure
SEM	Standard error mean
Ser46	Serine 46
SHR	Spotaneously hypertensive rats

XXV

	SKRB-3	Retinoid receptor- positive breast cancer cell line
	SOD	Superoxide dismutase
	SPSS	Statistical package for social science
	S	Seconds
	TAB	TAK-1 binding protein
	ТАК	Transforming growth factor- β activated kinase
	TBE	Tris/Borate/Ethylenediaminetetraacetic acid
	t-BOOH	tert-butylhydroperoxide
	TGs	Triacylglycerols
	TNF	Tumor necrosis factor
	TNF- α	Tumor necrosis factor-α
	TNFR	TNF receptor
	ТРС	Total phenolic content
	TPTZ	2,4,6-tri(2-pyridyl)-1,3,5-triazine
	TRADD	TNFR-associated death domain
	TRAF	TNFR-associated factor
	TUNEL	TdT-mediated dUTP nick-end-labeling
	t	Time
	U	Units
	UK	United Kingdom
	USA	United States of America
	UV	Ultraviolet
	UV-VIS	Ultraviolet-visible
	V	Volt

vWF	von Willebrand Factor
WHO	World Health Organization
WISP-1	WNT1-inducible-signaling protein 1
WRL-68	Hepatic human cell line
WST	Water soluble Tetrazolium salts
XIAP	X-linked inhibitor of apoptosis
XTT	Sodium 2,3,-bis(2-methoxy-4-nitro-5- sulfophenyl)-5-[(phenylamino)-carbonyl]- 2H-tetrazolium inner salt
22Rv1	Androgen responsive prostate cancer cell line
cm ²	Square centimetre
μg	Microgram
μ	Microliter
μm	Micrometer
μΜ	Micromolar
±	Approximately or about
°C	Celcius
%	Percentage
v/v	Volume per volume
w/v	Weight per volume

CHAPTER 1

INTRODUCTION

1.1 Background of study

Cancer is an abnormal mass of tissue as a result of uncontrollable proliferation which continued in excessive manner once the stimuli that induce the alteration terminated (Stricker and Kumar, 2007). Development of cancer frequently occurs in any region of organ or tissues such as skin, breast, lung, colon, nerve tissue and bones. According to the GLOBOCAN in the year of 2008, it was estimated approximately 7.6 million of people died due to cancer and about 12.7 million of people were diagnosed with cancer. From this statistic, about 56% of the incidence and 64% of the mortality arise from the economically developed countries (Jemal et al., 2011).

In Malaysia, cancer is one of the major public health problems. A sum of 18 219 new cancer cases was discovered in 2007 as recorded in the National Cancer Registry (NCR). The cancer incidence rate for males and females were 8123 (44.6%) and 10 096 (55.4%) peoples respectively. Five most frequent cancer cases among Malaysian males in 2007 were lung, colorectal, nasopharynx, prostate and lymphoma, while the five most common cancer cases in females were breast, colorectal, cervix, ovary and lung (Zainal Ariffin and Nor Saleha, 2011).

Breast cancer is the most common cancer diagnosed in women and the principal cause of death among women in most parts of the world. According to statistic, breast cancer contributed approximately 410 000 death cases per year (Coughlin and Ekwueme, 2009). In Malaysia, breast cancer is prevalent in women amongst all races as early age of 20. It is most common cancer disease in Malays, followed by Chinese and Indian. Based on the NCR 2007, breast cancer incidence was approximately 32.1% of overall cancer occurring among women population in Malaysia (Zainal Ariffin and Nor Saleha, 2011).

Polyphenol is vital part of the human diet and mainly present in berries, grapes/wine, chocolate/cocoa, coffee, soybeans and other fruits and vegetables. Several studies reported that polyphenol possessed antimutagenic properties and powerful free radical scavengers (Stoner and Mukhtar, 1995; Rice-Evans et al., 1995). These properties exhibited that polyphenol is one of the dietary compounds which obviously emerged as potential cytotoxicity agents against cancer.

Currently, cocoa is one of the famous dietary polyphenol that contain rich source of polyphenol which possessed beneficial effect for human health. Cocoa tree or scientifically known as *Theobroma cacao* L., belongs to family of *Sterculiaceae* which originated from the area of central, southern and southeastern Mexico (Rusconi and Conti, 2010). Catechin, epicatechin, flavanol glycosides, anthocyanins

procyanidins are among polyphenol identified in cocoa beans and cocoa products (Rimbach et al., 2009). Furthermore, cocoa demonstrated higher antioxidant capacity and contains more phenolic phytochemicals than teas and red wine (Arteel and Sies, 1999). Research on pharmacological potential of cocoa has been done over the past few years. Numerous investigators reported that cocoa phenolic contained bioactive phenolic compounds that possessed antioxidant, anticarcinogenic, and antiradical properties (Sanbongi et al., 1998; Wollgast and Anklam, 2000; Ren et al., 2003). However, the cytotoxicity study against cancer cell lines that had been done in previous research is insufficient to answer the present research objective due to different origin of cocoa polyphenol extract (CPE) (United States of America and France) and different cancer cell lines used such as Caco-2, 22Rv1 and DU145, MDA MB-436, MDA MB-468 and SKRB-3. Furthermore, the data on mode of cell death including AOPI, annexin V FITC-PI, DNA fragmentation, p53 and caspase-9 assay were scanty. Thus, the aforementioned parameters were conducted to elucidate the mode of cell death and prove the pharmacological potential of cocoa for cytotoxicity study in vitro.

1.2 Problem statement

The prevalence of breast cancer and mortality rate among women suffered with this disease is increasing every year. In fact, there are a number of factors which contribute to the breast cancer risk such as age, genetic hereditary, radiation, lifestyle, environmental and hormonal (Washbrook, 2006). Moreover, breast cancer was confirmed to be a challenging disease to cure and only several effective drugs are available. To date, surgery, hormone therapy, chemotherapy and radiotherapy are several conventional strategies for breast cancer treatment (El Saghir et al., 2011). However, these treatments were insufficient in order to prevent breast cancer patients from intermittence and metastasis of the tumor.

In addition, slowing action of chemotherapeutic drugs in breast cancer treatment will cause body developing resistance towards the drugs and trigger tumor recurrence (Yaacob et al., 2010). On top of that, the drugs cause critical side-effects to the patients such as cardiac and other toxicities (Beer and Bubalo, 2001; Leonard et al., 2009; Wonders and Reigle, 2009). Due to adverse effects of drug, a great deal of research has been conducted to explore natural products from plants and to search for potential cytotoxic activity towards breast cancer. Instead of medicinal herbs, dietary approaches have attracted tremendous attention among nutritionist, scientist and consumers due to its role to prevent, slow and delay the development of breast cancer without causing excessive damage to normal cells in human body. Therefore, this study is designed to elucidate the cytotoxicity effect of CPE originated from cocoa bean clone KKM4, KKM22, PBC123 and PBC159 towards breast cancer cell line (MCF-7).



1.3 Research objectives

1.3.1 General objective:

To investigate the cytotoxicity effect of CPE towards MCF-7 cells and elucidates mode of cell death.

1.3.2 Specific objectives:

- 1. To determine CPE constituents by qualitative phytochemical screening and HPLC profiling analysis.
- 2. To determine total phenolic and antioxidant activity of CPE using TPC, DPPH and FRAP assay.
- 3. To determine the IC_{50} value of CPE on various cancers cell lines using MTT assay.
- 4. To determine the IC₅₀ value of CPE against MCF-7 at different time points (24, 48 and 72 h) using MTT assay.
- 5. To determine the mode of cell death of CPE towards MCF-7 using inverted light microscope, acridine orange/propidium iodide, cell cycle, annexin V FITC and DNA fragmentation assay.
- 6. To determine the apoptotic pathway of CPE towards MCF-7 using p53 and caspase-9 ELISA kits.

1.4 Research hypothesis

CPE possess high total phenolic content, antioxidant activity and exhibit cytotoxicity towards MCF-7 cells through morphological alteration of cells, cell cycle phase arrest, phosphatidylserine (PS) externalization, DNA fragmentation and inducing apoptosis via increasing level of p53 and caspase-9 expression.

REFERENCES

- Abd Ghafar, S.A., Ismail, M., Saiful Yazan, L., Fakurazi, S., Ismail, N., Chan, K.
 W. and Md Tahir, P. (2013). Cytotoxic Activity of Kenaf Seed Oils from Supercritical Carbon Dioxide Fluid Extraction towards Human Colorectal Cancer (HT-29) Cell Lines. *Evidence-Based Complementary and Alternative Medicine*. 2013: 8.
- Abdul, A.B., Abdel-Wahab, S.I., Fong, H.K., Mohan, S.M., Al-Zubairi, A.S. and Elhassan, M.M. (2009). *In vitro* Response of Cancer Cells to the Growth-Inhibitory Effects of Dichloromethane Extract of *Goniothalamus umbrosus*. *Research Journal of Pharmacology*. 3(1): 1-6.
- Adami, H.O., Persson, I., Ekbom, A., Wolk, A., Pontén, J. and Trichopoulos, D. (1995). The aetiology and pathogenesis of human breast cancer. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*. 333(1–2): 29-35.
- Agarwal, M.L., Agarwal, A., Taylor, W.R. and Stark, G.R. (1995). p53 controls both the G2/M and the G1 cell cycle checkpoints and mediates reversible growth arrest in human fibroblasts. *Proceedings of the National Academy of Sciences*. 92(18): 8493-8497.
- Al-Sader, H., Abdul-Jabar, H., Allawi, Z. and Haba, Y. (2009). Alcohol and Breast Cancer : The Mechanism Explained. *Journal of Clinical Medicine Research*. 1(3): 125-131.
- Amundson, S.A., Myers, T.G., Albert, J. and Fornace, Jr. (1998). Roles for p53 in growth arrest and apoptosis : putting on the brakes after genotoxic stress. Oncogene. 17(25): 3287-3299.
- Armania, N., Yazan, L. S., Musa, S. N., Ismail, I. S., Foo, J. B., Chan, K. W., Noreen, H., Hisham, A.H., Zulfahmi, S. and Ismail, M. (2013). *Dillenia suffruticosa* exhibited antioxidant and cytotoxic activity through induction of apoptosis and G2/M cell cycle arrest. *Journal of Ethnopharmacology*. 146(2): 525-535.
- Arnoult, D., Gaume, B., Karbowski, M., Sharpe, J.C., Cecconi, F. and Youle, R.J. (2003). Mitochondrial release AIF and EndoG requires caspase activation downstream of Bax/Bak-mediated permeabilization. *The EMBO Journal*. 22: 4385-4399.
- Arteel, G.E. and Sies, H. (1999). Protection against peroxynitrite by cocoa polyphenol oligomers. *FEBS Letters*. 462(1–2): 167-170.
- Arts, I.C. and Hollman, P.C. (2005). Polyphenols and disease risk in epidemiologic studies. *The American Journal of Clinical Nutrition*. 81(1): 317S-325S.

- Bahorun, T., Luximon-Ramma, A., Crozier, A. and Aruoma, O.I. (2004). Total phenol, flavonoid, proanthocyanidin and vitamin C levels and antioxidant activities of Mauritian vegetables. *Journal of the Science of Food and Agriculture*. 84(12): 1553-1561.
- Bai, L. and Zhu, W.G. (2006). p53: Structure, Function and Therapeutic Applications. *Journal of Cancer Molecules*. 2(4): 141-153.
- Balasubramanian, K. and Ragunathan. (2012). Study of antioxidant and anticancer activity of natural sources. *Journal of Natural Product and Plant Resources*. 2(1): 192-197.
- Banin, S., Moyal, L., Shieh, S.Y., Taya, Y., Anderson, C.W., Chessa, L., Smorodinsky, C., Prives, Y., Reiss, Y., Shiloh, Y. and Ziv, Y. (1998).
 Enhanced Phosphorylation of p53 by ATM in Response to DNA Damage. *Science*. 281(5383): 1674-1677.
- Bates, S. E., Valverius, E. M., Ennis, B. W., Bronzert, D. A., Sheridan, J. P., Stampfer, M. R., Mendelsohn, J.,Lippman, M.E. and Dickson, R. B. (1990). Expression of the Transforming Growth Factor-α/Epidermal Growth Factor Receptor Pathway in Normal Human Breast Epithelial Cells. *Endocrinology*. 126(1): 596-607.
- Beer, T.M. and Bubalo, J.S. (2001). Complications of chemotherapy for prostate cancer. *Seminar Urologic Oncolology*. 19(3): 222-230.
- Benzie, I.F., Chung, W. and Strain, J.J. (1999). "Antioxidant" (reducing) efficiency of ascorbate in plasma is not affected by concentration. *The Journal of Nutritional Biochemistry*. 10(3): 146-150.
- Benzie, I.F.F. and Strain, J.J. (1996). The Ferric Reducing Ability of Plasma (FRAP) as a Measure of "Antioxidant Power": The FRAP Assay. *Analytical Biochemistry*. 239(1): 70-76.
- Beyer, J., Drummer, O.H. and Maurer, H.H. (2009). Analysis of toxic alkaloids in body samples. *Forensic Science International*. 185(1–3): 1-9.
- Bradford, M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*. 72(1–2): 248-254.
- Brady, C.A. and Attardi, L.D. (2010). p53 at a glance. *Journal of Cell Science*. 123(15): 2527-2532.
- Broeks, A., Braaf, L. M., Wessels, L.F.A., van de Vijver, M., De Bruin, M.L., Stovall, M., Russell, N.S., E.van Leewen, F. and Van 't Veer, L.J. (2010). Radiation-Associated Breast Tumors Display a Distinct Gene Expression Profile. *International Journal of Radiation Oncology*Biology*Physics*. 76(2): 540-547.

- Busch, S. J., Barnhart, R. L., Martin, G. A., Flanagan, M. A. and Jackson, R. L. (1990). Differential regulation of hepatic triglyceride lipase and 3-hydroxy-3methylglutaryl-CoA reductase gene expression in a human hepatoma cell line, HepG2. *Journal of Biological Chemistry*. 265(36): 22474-22479.
- Cailleau, R., Olivé, M. and Cruciger, Q. J. (1978). Long-term human breast carcinoma cell lines of metastatic origin: Preliminary characterization. *In Vitro*. 14(11): 911-915.
- Canman, C.E., Lim, D.S., Cimprich, K.A., Taya, Y., Tamai, K., Sakaguchi, K., Apella, E., Kastan, M.B. and Siliciano, J.D. (1998). Activation of the ATM Kinase by Ionizing Radiation and Phosphorylation of p53. *Science*. 281(5383): 1677-1679.
- Cardone, M.H., Roy, N., Stennicke, H.R., Salvesen, G.S., Franke, T.F., Stanbridge, E., Frisch, S. and Reed, J.C. (1998). Regulation of Cell Death Protease Caspase-9 by Phosphorylation. *Science*. 282(5392): 1318-1321.
- Carnésecchi, S., Schneider, Y., Lazarus, S.A., Coehlo, D., Gossé, F. and Raul, F. (2002). Flavanols and procyanidins of cocoa and chocolate inhibit growth and polyamine biosynthesis of human colonic cancer cells. *Cancer Letters*. 175(2): 147-155.
- Cecconi, F., Alvarez-Bolado, G., Meyer, B.I., Roth, K.A. and Gruss, P. (1998). Apaf1 (CED-4 Homolog) Regulates Programmed Cell Death in Mammalian Development. *Cell*. 94(6): 727-737.
- Chan, K.M., Rajab, N.F., Ishak, M.H.A., Ali, A.M., Yusoff, K., Din, L.B. and Inayat-Hussain, S.H. (2006). Goniothalamin induces apoptosis in vascular smooth muscle cells. *Chemico-Biological Interactions*. 159(2): 129-140.
- Ching, B., Chen, X.L., Yong, J.H.A., Wilson, J.M., Hiong, K.C., Sim, E.W.L., Wong, W.P., Lam, S.H., Chew, S.F. and Ip, Y.K. (2013). Increases in apoptosis, caspase activity and expression of p53 and bax, and the transition between two types of mitochondrion-rich cells, in the gills of the climbing perch, Anabas testudineus, during a progressive acclimation from freshwater to seawater. *Frontiers in Physiology*. 4.
- Cho, E.S., Jang, Y.J., Kang, N.J., Hwang, M.K., Kim, Y.T., Lee, K.W. and Lee, H.J. (2009). Cocoa procyanidins attenuate 4-hydroxynonenal-induced apoptosis of PC12 cells by directly inhibiting mitogen-activated protein kinase kinase 4 activity. *Free Radical Biology and Medicine*. 46(10): 1319-1327.
- Coughlin, S.S. and Ekwueme, D.U. (2009). Breast cancer as a global health concern. *Cancer Epidemiology*. 33(5): 315–318.
- Cui, H., Schroering, A. and Ding, H.F. (2002). p53 Mediates DNA Damaging Druginduced Apoptosis through a Caspase-9-dependent Pathway in SH-SY5 Neuroblastoma Cells. *Molecular Cancer Therapeutics*. 1(9): 679-686.

- Dai, J. and Mumper, R.J. (2010). Plant Phenolics: Extraction, Analysis and Their Antioxidant and Anticancer Properties. *Molecules*. 15(10): 7313-7352.
- Darlington, G. J., Kelly, J. H. and Buffone, G. J. (1987). Growth and hepatospecific gene expression of human hepatoma cells in a defined medium. *In Vitro Cellular & Developmental Biology*. 23(5): 349-354.
- Devedjian, J. C., Fargues, M., Denis-Pouxviel, C., Daviaud, D., Prats, H. and Paris, H. (1991). Regulation of the alpha 2A-adrenergic receptor in the HT29 cell line. Effects of insulin and growth factors. *Journal of Biological Chemistry*. 266(22): 14359-14366.
- Deveraux, Q.L., Roy, N., Stennicke, H.R., Arsdale, T.V., Zhou, Q., Srinivasula, S.M., Alnemri, E.S., Salvesen, G.Y. and Reed, J.C. (1998). IAPs block apoptotic events induced by caspase-8 and cytochrme c by direct inhibition of distinct caspases. *The EMBO Journal*. 17(8): 2215-2223.
- Dobashi, Y., Shoji, M., Jiang, S.X., Kobayashi, M., Kawakubo, Y. and Kameya, T. (1998). Active Cyclin A-CDK2 Complex, a Possible Critical Factor for Cell Proliferation in Human Primary Lung Carcinomas. *The American Journal of Pathology*. 153(3): 963-972.
- Eckelman, B.P., Salvesen, G.S. and Scott, F.L (2006). Human inhibitor of apoptosis proteins: why XIAP is the black sheep of the family. *EMBO Reports*. 7(10): 988-994.
- Elmore, S. (2007). Apoptosis: A Review of Programmed Cell Death. *Toxicologic Pathology*. 35(4): 495-516.
- El Saghir, N.S., Tfayli, A., Hatoum, H.A., Nachef, Z., Dinh, P. and Awada, A. (2011). Treatment of metastatic breast cancer: State-of-the-art, subtypes and perspectives. *Critical Reviews in Oncology/Hematology*. 80(3): 433-449.
- Fan, T.J., Han, L.H., Cong, R.S. and Liang, J. (2005). Caspase Family Proteases and Apoptosis. *Acta Biochimica et Biophysica Sinica*. 37(11): 719-727.
- Fantini, J., Cook, D. G., Nathanson, N., Spitalnik, S. L. and Gonzalez-Scarano, F. (1993). Infection of colonic epithelial cell lines by type 1 human immunodeficiency virus is associated with cell surface expression of galactosylceramide, a potential alternative gp120 receptor. *Proceedings of the National Academy of Sciences*. 90(7): 2700-2704.
- Fatland, B.L., Ke, J., Anderson, M.D., Mentzen, W.I., Cui, L.W., Allred, C.C., Johnston, J.L., Nikolau, B.J. and Wurtele, E.S. (2002). Molecular Characterization of a Heteromeric ATP-Citrate Lyase That Generates Cytosolic Acetyl-Coenzyme A in Arabidopsis. *Plant Physiology*. 130(2): 740-756.

- Flatt, P.M., Polyak, K., Tang, L.J., Scatena, C.D., Westfall, M.D., Rubinstein, L.A., Yu, J., Kinzler, K.W., Vogelstein, B., Hill, D.E. and Pietenpol, J.A. (2000). p53-dependent expression of PIG3 during proliferation, genotoxic stress, and reversible growth arrest. *Cancer Letters*. 156(1): 63-72.
- Fogh, J., Fogh, J. M. and Orfeo, T. (1977). One Hundred and Twenty-Seven Cultured Human Tumor Cell Lines Producing Tumors in Nude Mice. *Journal* of the National Cancer Institute. 59(1): 221-226.
- Franks, L.M. and Knowles, M.A. (2005). What is cancer? In Knowles, M. and Selby,
 P. *Introduction to the Cellular and Molecular Biology of Cancer* (pp. 1-24).
 4th edition: Oxford University Press.
- Freshney, R.I. (2005). Culture of Animal Cells: A manual of Basic Technique (pp.1-645). 5th edition. A John Willey and Sons, Inc Publication.
- Gammon, M.D., Santella, R.M., Neugut, A.I., Eng, S.M., Teitelbaum, S.L., Paykin, A., Levin, B., Terry, M.B., Young, T.L., Wang, L.W., Wang, Q., Britton, Wolff, M.S., Stellman, S.D., Hatch, M., Kabat, G.C., Senie, R., Garbowski, G., Maffeo, C., Montalvan, P., Berkowitz, G., Kemeny, M., Citron, M., Schnabel, F., Schuss, A., Hadju, S. and Vinceguerra, V. (2002). Environmental Toxins and Breast Cancer on Long Island. I. Polycyclic Aromatic Hydrocarbon DNA Adducts. *Cancer Epidemiology Biomarkers and Prevention.* 11(8): 677-685.
- Gasser, P., Lati, E., Peno-Mazzarino, L., Bouzoud, D., Allegaert, L. and Bernaert, H. (2008). Cocoa polyphenols and their influence on parameters involved in ex vivo skin restructuring. *International Journal of Cosmetic Science*. 30(5): 339-345.
- Gillett, C., Fantl, V., Smith, R., Fisher, C., Bartek, J., Dickson, C., Barnes, D. and Peters, G. (1994). Amplification and Overexpression of Cyclin D1 in Breast Cancer Detected by Immunohistochemical Staining. *Cancer Research*. 54(7): 1812-1817.
- Graf, B.A., Milbury, P.E. and Blumberg, J.B. (2005). Flavanols, flavones, flavanones, and human health: epidemiological evidence. *Journal of Medicinal Food.* 8(3): 281-290.
- Green, D.R. and Kroemer, G. (2009). Cytoplasmic function of the tumor suppressor p53. *Nature*. 458: 1127-1130.
- Green, D.R. and Reed, J.C. (1998). Mitochondria and Apoptosis. *Science*. 28(5381): 1309-1312.
- Greene, M.H. (1997). Genetics of Breast Cancer. Mayo Clinic proceedings. Mayo Clinic, 72(1): 54-65.
- Gu, Y., Hurst, W.J., Stuart, D.A. and Lambert, J.D. (2011). Inhibition of Key Digestive Enzymes by Cocoa Extracts and Procyanidins. *Journal of Agricultural and Food Chemistry*. 59(10): 5305-5311.

- Hague, A., Díaz, G.D., Hicks, D.J., Krajewski, S., Reed, J.C. and Paraskeva, C. (1997). Bcl-2 and bak may play a pivotal role in sodium butyrate-induced apoptosis in colonic epithelial cells; however overexpression of bcl-2 does not protect against bak-mediated apoptosis. *International Journal of Cancer*. 72(5): 898-905.
- Hall, P., Ploner, A., Bjohle, J., Huang, F., Lin, C.Y., Liu, E.T., Miller, L.D., Nordgren, H., Pawitan, Y., Shaw, P., Skoog, L., Smeds, J., Wedren, S., Öhd, J. and Bergh, J. (2006). Hormone-replacement therapy influences gene expression profiles and is associated with breast-cancer prognosis: a cohort study. *BMC Medicine*. 4(1): 16.
- Halliwell, B., Aeschbach, R., Löliger, J. and Aruoma, O.I. (1995). The characterization of antioxidants. *Food and Chemical Toxicology*. 33(7): 601-617.
- Hanski, C., Stolze, B. and Riecken, E. G. (1992). Tumorigenicity, mucin production and AM-3 epitope expression in clones selected from the HT-29 colon carcinoma cell line. *International Journal of Cancer.* 50(6): 924-929.
- Harbour, J.W., Luo, R.X., Santi, A.D., Postigo, A.A. and Dean, D.C. (1999). Cdk Phosphorylation Triggers Sequential Intramolecular Interactions that Progressively Block Rb Functions as Cells Move through G1. *Cell*. 98(6): 859-869.
- Harper, J.W., Adami, G. R., Wei, N., Keyomarsi, K. and Elledge, S.J. (1993). The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell*. 75(4): 805-816.
- Hii, C.L., Law, C.L., Suzannah, S., Misnawi. and Cloke, M. (2009). Polyphenols in cocoa (*Theobroma cacao* L.). Asian Journal of Food and Agro-Industry. 2(04): 702-722.
- Himes, R. H., Kersey, R. N., Heller-Bettinger, I. and Samson, F. E. (1976). Action of the Vinca Alkaloids Vincristine, Vinblastine, and Desacetyl Vinblastine Amide on Microtubules in Vitro. *Cancer Research*, 36(10), 3798-3802.
- Hong, S.J., Tran, Q.A. and Keiler, K.C. (2005). Cell cycle-regulated degradation of tmRNA is controlled by RNase R and SmpB. *Molecular Microbiology*. 57(2): 565-575.
- Horsthemke, B. (1992). Genetics and cytogenetics of retinoblastoma. *Cancer Genetics and Cytogenetics*. 63(1): 1-7.
- Huang, X., Halicka, H.D., Traganos, F., Tanaka, T., Kurose, A. and Darzynkiewicz, Z. (2005). Cytometric assessment of DNA damage in relation to cell cycle phase and apoptosis. *Cell Proliferation*. 38(4): 223-243.
- Huda-Faujan, N., Noriham, A., Norrakiah, A.S. and Babji, A.S. (2007). Antioxidative Activities of Water Extracts of Some Malaysian Herbs. *ASEAN Food Journal*. 14(1): 61-68.

- Huguet, E. L., McMahon, J. A., McMahon, A. P., Bicknell, R. and Harris, A. L. (1994). Differential Expression of Human Wnt Genes 2, 3, 4, and 7B in Human Breast Cell Lines and Normal and Disease States of Human Breast Tissue. *Cancer Research*. 54(10): 2615-2621.
- Iida, H., Towatari, M., Tanimoto, M., Morishita, Y., Kodera, Y. and Saito, H. (1997). Overexpression of Cyclin E in Acute Myelogenous Leukemia. *Blood*. 90(9): 3707-3713.
- Inayat-Hussain, S.H., Osman, A.B., Din, L.B. and Taniguchi, N. (2002). Altholactone, a novel styryl-lactone induces apoptosis via oxidative stress in human HL-60 leukemia cells. *Toxicology Letters*. 131(3): 153-159.
- Ishikawa, T., Furihata, M., Ohtsuki, Y., Murukami, H., Inoue, A. and Ogoshi, S. (1998). Cyclin D1 overexpression related to retinoblastoma protein expression as prognostic marker in human oesophgeal squamos cell carcinoma, *British Journal of Cancer*. 77(1): 92-97.
- Isobe, M., Emanuel, B.S., Givol, D., Oren, M. and Croce, C.M. (1986). Localization of gene for human p53 tumor antigen to band 17p13. *Nature*. 320: 84-85.
- Jemal, A., Bray, F., Center, M.M., Ferlay, J., Ward, E. and Forman, D. (2011). Global cancer statistics: GLOBOCAN 2008. *CA: A Cancer Journal for Clinicians*. 61(2): 69-90.
- Jones, H. W. J., Mc Kusick, V. A., Harper, P. S. and Wuu, K.-D. (1971). George Otto Gey (1899-1970): The HeLa Cell and a Reappraisal of its Origin. *Obstetrics and Gynecology*. 38(6): 945-949.
- Jourdain, C., Tenca, G., Deguercy, A., Troplin, P. and Poelman, D. (2006). In-vitro effects of polyphenols from cocoa and β-sitosterol on the growth of human prostate cancer and normal cells. *European Journal of Cancer Prevention*. 15(4): 353-361.
- Kelvin, J.F. and Tyson, L.B. (2010). Cancer symptoms and cancer treatment side effects. *Cancer and cancer treatment* (pp. 1-30). 2nd edition: Jones and Bartlett Publishers.
- Kenny, T. (2012). Breast cancer, <u>http://www.patient.co.uk/health/breast-cancer-leaflet</u>. Egton Medical Information System (Accessed by 30 October 2013).
- Kerr, J.F.R., Wylie, A.H. and Currie, A.R. (1972). Apoptosis: A Basic Biological Phenomenon with Wide-ranging implications in Tissue Kinetics. *British Journal of Cancer*. 26(4): 239-257.
- Keyomarsi, K., O'Leary, N., Molnar, G., Lees, E., Fingert, H.J. and Pardee, A.B. (1994). Cyclin E, a Potential Prognostic Marker for Breast Cancer. *Cancer Research*. 54(2): 380-385.

- Keyomarsi, K. and Pardee, A.B. (1993). Redundant cyclin overexpression and gene amplification in breast cancer cells. *Proceedings of the National Academy of Sciences*. 90(3): 1112-1116.
- Khalid, M., Siddiqui, H.H. and Freed, S. (2011). Free Radical Scavenging and Total Phenolic Content of *Saccharum spontaneum* L. Root Extract. *International Journal of Research In Pharmacy and Chemistry*. 1(14): 2231-2781.
- Kim, Y.S., Morgan, M.J., Choksi, S. and Liu, Z.G. (2007). TNF-Induced Activation of the Nox1 NADPH Oxidase and Its Role in the Induction of Necrotic Cell Death. *Molecular Cell*. 26(5): 675-687.
- Kischkel, F.C., Hellbardt, S., Behrmann, I., Germer, M., Pawlita, M., Krammer, P.H. and Peter, M.E. (1995). Cytotoxicity-dependent APO-1 (Fas/CD95)associated proteins form a death-inducing signaling complex (DISC) with the receptor. *The EMBO Journal*. 14(22): 5579-5588.
- Klener Jr., Anděra, P., Klener, L., Klener, P., Nečas, P. and Živný. E. (2006). Cell Death Signaling Pathways in the Pathogenesis and Therapy of Haematologic Malignancies: Overview of Apoptotic Pathways. *Folia Biologica*. 52: 34-44.
- Konfink, M., Papagiannopoulos, M. and Galensa, R. (2007). (-)-Catechin in Cocoa and Chocolate: Occurence and Analysis of an Atypical Flavan-3-ol Enantiomer. *Molecules*. 12(7): 1274-1288.
- Kuida, K. (2000). Caspase-9. The International Journal of Biochemistry and Cell Biology. 32(2): 121-124.
- Kumarappan, C.T., Thilagam, E. and Mandal, C.M. (2012). Antioxidant activity of polyphenolic extracts of *Ichnocarpus frutescens*. Saudi Journal of Biological Sciences. 19(3): 349-355.
- Knowles, B. B. and Aden, D. P. (1983). Human hepatoma derived cell line, process for preparation thereof, and uses therefor: Google Patents.
- Laín, S. and Lane, D.P. (2005). Tumor suppressor genes. In Knowles, M. and Selby,
 P. *Introduction to the Cellular and Molecular Biology of Cancer* (pp. 135-155). 4th edition: Oxford University Press.
- Lamb, P. and Crawford, L. (1986). Characterization of the human p53 gene. *Molecular and Cellular Biology*. 6(5): 1379-1385.
- Lapidot, T., Walker, M.D. and Kanner, J. (2002). Antioxidant and Prooxidant Effects of Phenolics on Pancreatic β-Cells *in vitro*. *Journal of Agricultural and Food Chemistry*. 50(25): 7220-7225.
- Laptenko, O. and Prives, C. (2006). Transcriptional regulation by p53: one protein, many possibilities. *Cell Death and Differentiation*. 13: 951-961.

- Lattanzio, V., Lattanzio, V.M.T. and Cardinali, A. (2006). Role of phenolics in the resistance mechanism of plants against fungal pathogens and insects. In F. Imperato. *Phytochemistry: Advances in Research* (pp. 23-67). Research Signpost Publishers, Kerala, India.
- Leach, F.S., Elledge, S.J., Sherr, C.J., Willson, J.K.V., Markowitz, S., Kinzler, K.W. and Vogelstein, B. (1993). Amplification of Cyclin Genes in Colorectal Carcinomas. *Cancer Research*. 53(9): 1986-1989.
- Lecumberri, E., Goya, L., Mateos, R., Alía, M., Ramos, S., Izquierdo-Pulido, M. and Bravo, L. (2007). A diet rich in dietary fiber from cocoa improves lipid profile and reduces malondialdehyde in hypercholesterolemic rats. *Nutrition*. 23(4): 332-341.
- Lee, Y.J., Kim, D.B., Lee, J., Cho, J.H., Kim, B., Choi, H.S., Lee, B.Y. and Lee, O.H. (2013). Antioxidant Activity and Anti-Adipogenic Effects of Wild Herbs Mainly Cultivated in Korea. *Molecules*. 18(10): 12937-12950.
- Lee, D.E., Kang, N.J., Lee, K.M., Lee, B.K., Kim, J.H., Lee, K.W. and Lee, H.J. (2010). Cocoa polyphenols attenuate hydrogen peroxide-induced inhibition of gap-junction intercellular communication by blocking phosphorylation of connexin 43 via the MEK/ERK signaling pathway. *The Journal of Nutritional Biochemistry*. 21(8): 680-686.
- Lee, K.W., Kundu, J.K., Kim, S.O., Chun, K.S., Lee, H.J. and Surh, Y.J. (2006). Cocoa Polyphenols Inhibit Phorbol Ester-Induced Superoxide Anion Formation in Cultured HL-60 Cells and Expression of Cyclooxygenase-2 and Activation of NF-κB and MAPKs in Mouse Skin In Vivo. *The Journal of Nutrition.* 136(5): 1150-1155.
- Leonard, R.C., Williams, S., Tulpule, A., Levine, A.M. and Oliveros, S. (2009). Improving the therapeutic index of anthracycline chemotherapy: focus on liposomal doxorubicin (Myocet). *Breast.* 18(4): 218-224.
- Leong, S.L., Cady, B., Jablons, D., Garcia-Aguilar, J., Reintgen, D., Jakub, J., Pendas, S., Duhaime, L., Gardner, M., Giuliano, R., Archie, V., Calvin, D., Mensha, L., Shivers, S., Cox, C., Werner, J.A., Kitaqawa, Y. and Kitajima, M. (2006). Clinical patterns of metastasis. *Cancer and Metastasis Reviews*. 25(2): 221-232.
- Lesuffleur, T., Porchet, N., Aubert, J. P., Swallow, D., Gum, J. R., Kim, Y. S., Real, F.X. and Zweibaum, A. (1993). Differential expression of the human mucin genes MUC1 to MUC5 in relation to growth and differentiation of different mucus-secreting HT-29 cell subpopulations. *Journal of Cell Science*. 106(3): 771-783.
- Levenson, A. S. and Jordan, V. C. (1997). MCF-7: The First Hormone-responsive Breast Cancer Cell Line. *Cancer Research*. 57(15): 3071-3078.
- Li, J. and Yuan, J. (2008). Caspases in apoptosis and beyond. Oncogene. 27: 6194–6206.

- Li, P., Nijhawan, D., Budihardjo, I., Srinivasula, S.M., Ahmad, M., Alnemri, E.S. and Wang, X. (1997). Cytochrome c and dATP-Dependent Formation of Apaf 1/Caspase-9 Complex Initiates an Apoptotic Protease Cascade. *Cell*. 91(4): 479-489.
- Lucey, B. P., Nelson-Rees, W. A. and Hutchins, G. M. (2009). Henrietta Lacks, HeLa Cells, and Cell Culture Contamination. *Archives of Pathology and Laboratory Medicine*. 133(9): 1463-1467.
- Luximon-Ramma, A., Bahorun, T., Crozier, A., Zbarsky, V., Datla, K.P., Dexter, D.T. and Aruoma, O.I. (2005). Characterization of the antioxidant functions of flavonoids and proanthocyanidins in Mauritian black teas. *Food Research International*. 38(4): 357-367.
- Luximon-Ramma, A., Bahorun, T. and Crozier, A. (2003). Antioxidant actions and phenolic and vitamin C contents of common Mauritian exotic fruits. *Journal of the Science of Food and Agriculture*. 83(5): 496-502.
- Malumbres, M. and Barbacid, M. (2005). Mammalian cyclin-dependent kinases. *Trends in Biochemical Sciences*. 30(11): 630-641.
- Mandal, S., Yadav, S. and Nema, R.K. (2009). Antioxidant: A Review. Journal of Chemical and Pharmaceutical Research. 1(1): 102-104.
- Martín, M.A., Ramos, S., Mateos, R., Granado Serrano, A.B., Izquierdo-Pulido, M., Bravo, L. and Goya, L. (2008). Protection of Human HepG2 Cells against Oxidative Stress by Cocoa Phenolic Extract. *Journal of Agricultural and Food Chemistry*. 56(17): 7765-7772.
- Martin, M.C., Allan, L.A., Lickrish, M., Sampson, C., Morrice, N. and Clarke, P.R. (2005). Protein Kinase A Regulates Caspase-9 Activation by Apaf-1 Downstream of Cytochrome c. *Journal of Biological Chemistry*. 280(15): 15449-15455.
- Martin, A.M. and Weber, B.L. (2000). Genetic and Hormonal Risk Factors in Breast Cancer. *Journal of the National Cancer Institute*. 92(14): 1126-1135.
- Matsuda, K., Yoshida, K., Taya, Y., Nakamura, K., Nakamura, Y. and Arakawa, H. (2002). p53AIP1 Regulates the Mitochondrial Apoptotic Pathway. *Cancer Research*. 62(10): 2883-2889.
- Matsuoka, S., Huang, M. and Elledge, S.J. (1998). Linkage of ATM to Cell Cycle Regulation by the Chk2 Protein Kinase. *Science*. 282(5395): 1893-1897.
- Mbzima, V.G., Mokgotho, M.P., February, F., Rees, D.J.G. and Mampuru, L.J. (2008). Alteration of Bax-to-Bcl-2 ratio modulates the anticancer activity of methanolic extract of *Commelina benghalensis* (Commelinaceae) in Jurkat T cells. *African Journal of Biotechnology*. 7(20): 3569-3576.
- Meek, D.W. (2004). The p53 response to DNA damage. *DNA Repair*. 3(8–9): 1049-1056.

- Mihara, M., Erster, S., Zaika, A., Petrenko, O., Chittenden, T., Pancoska, P. and Moll, U.M. (2003). p53 Has a Direct Apoptogenic Role at the Mitochondria. *Molecular Cell*. 11(3): 577-590.
- Miller, J.P., Yeh, N., Vidal, A. and Koff, A. (2007). Interweaving the Cell Cycle Machinery with Cell Differentiation. *Cell Cycle*. 6(23): 2932-2938.
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*. 65(1–2): 55-63.
- Murray, B.K., Brown, B., Scherer, P.M., Tomer, D.P., Garvin, K.R., Hughes, B.G. and O'Neill, K.L. (2006). Induction of apoptosis in HT-29 human colon adenocarcinoma cells by 13-cis-retinoic acid and vitamin E succinate. *Nutrition Research*. 26(4): 186-192.
- Nakano, K. and Vousden, K.H. (2001). PUMA, a Novel Proapoptotic Gene, Is Induced by p53. *Molecular Cell*. 7(3): 683-694.
- Nanji, A.A. and Hiller-Sturmhöfel, S. (1997). Apoptosis and Necrosis: Two Type of Cell Death in Alkoholic Liver Disease. *Alcohol Health and Research World*. 21(4): 325-330.
- Nilsson, I. and Hoffmann, I. (2000). Cell cycle regulation by the Cdc25 phosphatase family. In Meijer, L., Jézéquel, A. and Ducommun, B. (Eds.). *Progress in Cell Cycle Research* (pp. 107-114). Springer US.
- Norbury, C. and Nurse, P. (1992). Animal Cell Cycles and Their Control. Annual Review of Biochemistry. 61(1): 441-468.
- O'Connor, L., Harris, A.W. and Strasser, A. (2000). CD95 (Fas/APO-1) and p53 Signal Apoptosis Independently in Diverse Cell Types. *Cancer Research*. 60: 1217-1220.
- Oda, E., Ohki, R., Murasawa, H., Nemoto, J., Shibue, T., Yamashita, T., Tokino, T., Taniguchi, T. and Tanaka, N. (2000). Noxa, a BH3-Only Member of the Bcl-2 Family and Candidate Mediator of p53-Induced Apoptosis. *Science*. 288(5468): 1053-1058.
- Ormerod, M.G. (2004). Cell-Cycle Analysis of Asynchronous Populations. In Hawley, T.S. and Hawley, R.G. (Ed). *Methods in Molecular Biology: Flowcytometry protocols* (pp. 345-353). 2nd edition: Humana Press Inc., Totowa, New Jersey.
- Osawa, T. (1994). Novel natural antioxidants for utilization in food and biological systems. In Uritani, I., Garcia, V.V. and Mendoza, E.M. (Eds). *Postharvest Biochemistry of plant Food-Materials in the Tropics* (pp. 241-251). Japan Scientific Soceities Press: Tokyo, Japan.

- Pandey, K.B. and Rizvi, S.I. (2009). Plant polyphenols as dietary antioxidants in human health and disease. Oxidative Medicine and Cellular Longetivity. 2(5): 270-278.
- Park, C., Moon, D.O., Rhu, C.H., Choi, B.T., Lee, W.H., Kim, G.Y. and Choi, Y.H. (2007). β-Sitosterol Induces Anti-proliferation and Apoptosis in Human Leukemic U937 Cells through Activation of Caspase-3 and Induction of Bax/Bcl-2 Ratio. *Biological and Pharmaceutical Bulletin*. 30(7): 1317-1323.
- Pearson, D.A., Paglieroni, T.G., Rein, D., Wun, T., Schramm, D.D., Wang, J.F., Holt, R.R., Gosselin, R., Schmitz, H.H. and Keen, C.L. (2002). The effects of flavanol-rich cocoa and aspirin on ex vivo platelet function. *Thrombosis Research*. 106(4–5): 191-197.
- Pecorino, L. (2005). What is cancer? *Molecular Biology of Cancer*. *Mechanisme*, *Targets and Therapeutics* (pp. 1-20). 1st edition: Oxford University Press.
- Perron, N.R., Wang, H.C., DeGuire, S.N., Jenkins, M., Lawson, M. and Brumaghim, J.L. (2010). Kinetics of iron oxidation upon polyphenol binding. *Dalto Transactions*. 39: 9982-9987.
- Podolak, I., Galanty, A. and Sobolewska. (2010). Saponins as cytotoxic agents: a review. *Phytochemistry Reviews*. 9(3): 425-474.
- Pollack, M. S., Heagney, S. D., Livingston, P. O. and Fogh, J. (1981). HLA-A,B,C and DR Alloantigen Expression on Forty-Six Cultured Human Tumor Cell Lines. *Journal of the National Cancer Institute*. 66(6): 1003-1012.
- Qui[~]nones, M., Miguel, M., Muguerza, B. and Aleixandre, A. (2011). Effect of a cocoa polyphenol extract in spontaneously hypertensive rats. *Food and Function*. 2(11): 649-653.
- Ramiro, E., Franch, A., Castellote, C., Perez-Cano, F., Permanyer, J., Izquierdo-Pulido, M. and Castell, M. (2005). Flavonoids from *Theobroma cacao* Down-Regulate Inflammatory Mediators. *Journal of Agricultural and Food Chemistry*. 53(22): 8506-8511.
- Ramljak, D., Romanczyk, L.J., Metheny-Barlow, L.J., Thompson, N., Knezevic, V., Galperin, M., Ramesh, A. and Dickson, R.B. (2005). Pentameric procyanidin from *Theobroma cacao* selectively inhibits growth of human breast cancer cells. *Molecular Cancer Therapeutics*. 4(4): 537-546.
- Reiter, L. S., Kruithof, E. K. O., Cajot, J.-F. and Sordat, B. (1993). The role of the urokinase receptor in extracellular matrix degradation by ht29 human colon carcinoma cells. *International Journal of Cancer*. *53*(3): 444-450.
- Ren, W., Qiao, Z., Wang, H., Zhu, L. and Zhang, L. (2003). Flavonoids: promising anticancer agents. *Medicinal Research Reviews*. 23(4): 519–534.

- Rice-Evans, C., Miller, N. and Paganga, G. (1997). Antioxidant properties of phenolic compounds. *Trends in Plant Science*. 2(4): 152-159.
- Rice-Evans, C.A., Miller, N.J., Bolwell, P.G., Bramley, P.M. and Pridham, J.B. (1995). The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radical Research*. 22(4): 375-383.
- Richie, R.C. and Swanson, J.O. (2003). Breast Cancer: A Review of the Literature. *Journal of Insurance Medicine*. 35(2): 85-101.
- Riley, T., Sontag, E., Chen, P. and Levine, A. (2008). Transcriptional control human p53-regulated genes. *Nature Reviews Molecular Cell Biology*. 9: 402-412.
- Rimbach, G., Melchin, M., Moehring, J. and Wagner. (2009). Polyphenols from Cocoa and Vascular Health-A Critical Review. *International Journal of Molecular Science*. 10(10): 4290-4309.
- Rogakou, E.P., Nieves-Neira, W., Boon, C., Pommier, Y. and Bonner, W.M. (2000). Initiation of DNA Fragmentation during Apoptosis Induces Phosphorylation of H2AX Histone at Serine 139. *Journal of Biological Chemistry*. 275(13): 9390-9395.
- Rusconi, M. and Conti, A. (2010). *Theobroma cacao* L., the Food of the Gods: A scientific approach beyond myths and claims. *Pharmacological Research*. 61(1): 5-13.
- Sanbongi, C., Osakabe, N., Natsume, M., Takizawa, T., Gomi, S. and Osawa, T. (1998). Antioxidative polyphenols isolated from *Theobroma cocoa*. Journal of Agricultural and Food Chemistry. 46(2): 454-457.
- Sánchez-Rabaneda, F., Jáuregui, O., Casals, I., Andrés-Lacueva, C., Izquierdo-Pulido, M. and Lamuela-Raventós, R.M. (2003). Liquid chromatographic/ electrospray ionization tandem mass spectrometric study of the phenolic composition of cocoa (*Theobroma cacao*). Journal of Mass Spectrometry. 38(1): 35-42.
- Santos-Buelga, C. and Scalbert, A. (2000). Proanthocyanidins and tannin-like compounds–nature, occurrence, dietary intake and effects on nutrition and health. *Journal of the Science of Food and Agriculture*. 80(7): 1094-1117.
- Schuler, M., Bossy-Wetzel, E., Goldstein, J.C., Fitzgerald, P. and Green, D.R. (2000). p53 Induces Apoptosis by Caspase Activation through Mitochondrial Cytochrome c Release. *Journal of Biological Chemistry*. 275(10): 7337-7342.
- Scuderi, R., Palucka, K., Pokrovskaja, K., Bjorkholm, M., Wiman, K. and Pisa, P. (1996). Cyclin E overexpression in relapsed adult acute lymphoblastic leukemias of B-cell lineage. *Blood.* 87(8): 3360-3367.

- Shi, M., Lin, T.H., Appell, K.C. and Berg, L.J. (2009). Cell Cycle Progression following Naive T Cell Activation Is Independent of Jak3/Common γ-Chain Cytokine Signals. *The Journal of Immunology*. 183(7): 4493-4501.
- Shi, J., Yu, J., Pohorly, J. and Young, C. (2003). Optimization of the extraction of polyphenols from grape seed meal by aqueous ethanol solution. *Food, Agriculture and Environment*. 1(2): 42-47.
- Shinde, A., Ganu, J. and Naik, P. (2012). Effect of Free Radicals and Antioxidants on Oxidative Stress: A Review. *Journal of Dental and Allied Sciences*. 1(2): 63-66.
- Shiozaki, E.N., Chai, J. and Shi, Y. (2002). Oligomerization and activation of caspase-9 induced by Apaf-1 CARD. Proceedings of the National Academy of Sciences. 99(7): 4197-4202.
- Siegel, R., Naishadham, D. and Jemal, A. (2013). Cancer statistics, 2013. CA: A Cancer Journal for Clinicians. 63(1): 11-30.
- Sies, H., Schewe, T., Heiss, C., and Kelm, M. (2005). Cocoa polyphenols and inflammatory mediators. *The American Journal of Clinical Nutrition*. 81(1): 304S-312S.
- Silverstein, A., Silverstein, V. and Nunn, L.S. (2006). Cells gone wild. *In Cancer* (pp.7-12). United state of America: Lerner Publishing Group.
- Sisa, M., Bonnet, S.L., Ferreira, D. and Van der Westhuizen, J.H. (2010). Photochemistry of Flavonoids. *Molecules*. 15(8): 5196-5245.
- Soule, H. D., Vazquez, J., Long, A., Albert, S. and Brennan, M. (1973). A Human Cell Line From a Pleural Effusion Derived From a Breast Carcinoma. *Journal of the National Cancer Institute*. 51(5): 1409-1416.
- Stapleton, A.E. and Walbot, V. (1994). Flavonoids Can Protect Maize DNA from the Induction of Ultraviolet Radiation Damage. *Plant Physiology*. 105(3): 881-889.
- Stoner, G.D. and Mukhtar, H. (1995). Polyphenols as cancer chemoprevention agents. *Journal of Cellular Biochemistry Supplement*. 22: 169–180.
- Stork, C.J. and Li, Y.V. (2006). Measuring cell viability with membrane impermeable zinc fluorescent indicator. *Journal of Neuroscience Methods*. 155(2): 180-186.
- Stricker, T.P. and Kumar, V. (2007). Neoplasia. In Kumar, V., Abbas, A.K., Fausto, N. and Mitchell, R.N. *Robbins Basic Pathology* (pp. 173-223). 8th edition. Saunders Elsevier. Philadelphia.
- Subhashini, R., Mahadeva Rao, U.S., Sumathi, P. and Gunalan, G. (2010). A Comparative Phytochemical Analysis of Cocoa and Green Tea, *Indian Journal Science and Technology*. 3(2): 188-192.

- Sykes, J.A. (1975). Separation of Tumor Cells from Fibroblast. In Fogh, J (Ed). *Human tumor cells in vitro* (pp. 1-22). New York: Plenum Press.
- Suzuki, K. and Matsubara, H. (2011). Recent Advances in p53 Research and Cancer Treatment. *Journal of Biomedicine and Biotechnology*. 2011(978312): 1-7.
- Tajudin, T.J.S.A., Mat, N., Siti Aishah, A.B., Yusran, A.A.M., Alwi, A. and Ali, A.M. (2012). Cytotoxicity, Antiproliferative Effects and Apoptosis Induction of Methanolic Extract of *Cynometra cauliflora* Linn. Whole Fruit on Human Promyelocytic Leukemia HL-60 Cells. *Evidence-Based Complementary and Alternative Medicine*. 2012: 1-6.
- Takimoto, R. and El-Deiry, W.S. (2000). Wild-type p53 transactivates the KILLER/DR5 gene through an intronic sequence-specific DNA-binding site. *Oncogene*. 19(14): 1735-1743.
- Taylor, R.B., Cullen, S.P. and Martin, S.J. (2008). Apoptosis: controlled demolition at the cellular level. *Nature Reviews Molecular Cell Biology*. 9: 231-241.
- Taylor, W.R. and Stark, G.R. (2001). Regulation of the G2/M transition by p53. Oncogene. 20(15): 1803-1815.
- Tomas-Barberán, F.A., Cienfuegos-Jovellanos, E., Marín, A., Muguerza, B., Gil-Izquierdo, A., Cerdá, B., Zafrilla, P., Morillas, J., Mulero, J., Ibarra, A., Pasamar, M.A., Ramo, D. and Espín, J.C. (2007). A New Process To Develop a Cocoa Powder with Higher Flavonoid Monomer Content and Enhanced Bioavailability in Healthy Humans. *Journal of Agricultural and Food Chemistry*. 55(10): 3926-3935.
- Tsao, R. (2010). Chemistry and Biochemistry of Dietary Polyphenols. *Nutrients*. 2(12): 1231-1246.
- Tsao, R. and McCallum, J. (2009). Chemistry of Flavonoids. In: de la Rosa, L.A., Alvarez-Parrilla, E. and Gonzalez-Aguilar, G. Fruit and Vegetable Phytochemicals: Chemistry, Nutritional Value and Stability (pp. 131-153). Blackwell Publishing; Ames, Iowa (IA), United States of America (USA).
- Uygun, B. E., Uygun, K. and Yarmush, M. L. (2011). 5.541 Liver Tissue Engineering. In P. Ducheyne (Ed.), *Comprehensive Biomaterials* (pp. 575-585). Oxford: Elsevier.
- Van Cruchten, S. and Van den Broeck, W. (2002). Morphological and Biochemical Aspects of Apoptosis, Oncosis and Necrosis. *Anatomia, Histologia, Embryologia.* 31(4): 214-223.
- Velioglu, Y.S., Mazza, G., Gao, L. and Oomah, B.D. (1998). Antioxidant Activity and Total Phenolics in Selected Fruits, Vegetables, and Grain Products. *Journal of Agricultural and Food Chemistry*. 46(10): 4113-4117.

- Vermeulen, K., Van Bockstaele, D.R. and Berneman, Z.N. (2003). The cell cycle: a review of regulation, deregulation and therapeutic targets in cancer. *Cell Proliferation*. 36(3): 131-149.
- Washbrook, E. (2006). Risk factors and epidemiology of breast cancer. *Women's Health Medicine*. 3(1): 8-14.
- Waters. (2014). High Performance Liquid Chromatography. <u>http://www.waters.com/waters/en_US/HPLC---High-Performance-Liquid</u> Chromatography/nav.htm?cid=10048919. Copyright ©1995 Waters Corporation, 34 Maple Street, Milford, Massachusetts, 01757, USA. All rights reserved. (Accessed by 13 May 2014).
- Wink, M. (1993). Allelochemical properties or the Raison d'Être of Alkaloids. In Cordell,G.A. (Ed). The Alkaloids. (pp.1-104). Volume 48. Academic Press Inc.California, United State of America.
- Wollgast, J. and Anklam, E. (2000). Review on polyphenols in *Theobroma cacao*: changes in composition during the manufacture of chocolate and methodology for identification and quantification. *Food Research International*. 33(6): 423-447.
- Wonders, K.Y. and Reigle, B.S. (2009). Trastuzumab and doxorubicin-related cardiotoxicity and the cardioprotective role of exercise. *Integrative Cancer Therapies*. 8(1): 17-21.
- Wyllie, A.H., Kerr, J.F.R. and Currie, A.R. (1980). Cell death: the significance of apoptosis. *International Review of Cytology*. 68: 251–306.
- Xiong, Y., Hannon, G.J., Zhang, H., Casso, D., Kobayashi, R. and Beach, D. (1993). p21 is a universal inhibitor of cyclin kinases. *Nature*. 366(6456): 701-704.
- Yaacob, N.S., Hamzah, N., Nik, M. K., Zainal Abidin, S.A., Lai, C., Navaratnam, V. and Norazmi, M.N. (2010). Anticancer activity of a sub-fraction of dichloromethane extract of *Strobilanthes crispus* on human breast and prostate cancer cells *in vitro*. *BMC Complementary and Alternative Medicine*. 10(42): 1-14.
- Yazan, L.S., Foo, J.B., Chan, K.W., Tahir, P.M. and Ismail, M. (2011). Kenaf seed oil from supercritical carbon dioxide fluid extraction shows cytotoxicity effects towards various cancer cell lines. *African Journal of Biotechnology*. 10(27): 5381-5388.
- Yazan, L.S., Ng, W.K., Al-Naqeeb, G. and Ismail, M. (2009). Cytotoxicity of Thymoquinone (TQ) from *Nigella sativa* Towards Human Cervical Carcinoma Cells (HeLa). *Journal of Pharmacy Research*. 2(4): 585-589.
- Yu, J., Zhang, L., Hwang, P.M., Kinzler, K.W. and Vogelstein, B. (2001). PUMA Induces the Rapid Apoptosis of Colorectal Cancer Cells. *Molecular Cell*. 7(3): 673-682.

- Zahin, M., Aqil, F. and Ahmad, I. (2009). The *in vitro* antioxidant and total phenolic content of four Indian medicinal plants. *International Journal of Pharmacy and Pharmaceutical Sciences*. 1: 88-95.
- Zainal Ariffin, O. and Nor Saleha, I.T. (2011). National Cancer Registry (NCR) Report 2007, Ministry of Health, Malaysia.
- Zhou, J., Ahn, J., Wilson, S.H. and Prives, C. (2001). A role for p53 in base excision. *The EMBO Journal*. 20(4): 914-923.
- Zilfou, J.T. and Lowe, S.W. (2009). Tumor Supressive Functions of p53. *Cold Spring Habor Perspectives in Biology*. 1(5): 1-40.

