



UNIVERSITI PUTRA MALAYSIA

***HEDGEHOG PATHWAY PROTEINS AND THEIR ASSOCIATION WITH
CLINICOPATHOLOGICAL PARAMETERS OF BLADDER CANCER***

KHAIRUNNISA BINTI MOHD ARIFFIN

FPSK(M) 2016 42



**HEDGEHOG PATHWAY PROTEINS AND THEIR ASSOCIATION WITH
CLINICOPATHOLOGICAL PARAMETERS OF BLADDER CANCER**

By

KHAIRUNNISA BINTI MOHD ARIFFIN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfillment of the Requirements for the Degree of Master of Science**

August 2016

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

HEDGEHOG PATHWAY PROTEINS AND THEIR ASSOCIATION WITH CLINICOPATHOLOGICAL PARAMETERS OF BLADDER CANCER

By

KHAIRUNNISA BINTI MOHD ARIFFIN

August 2016

Chairman : Maizatun Atmadini Abdullah, PhD
Faculty : Medicine and Health Sciences

Hedgehog pathway is important for growth and patterning during embryonic development. Previous studies have shown that constitutive activation of Hedgehog pathway can lead to various types of malignancies including medulloblastoma, basal cell carcinoma, gastrointestinal, breast and prostate cancer. The purpose of this study was to investigate the expression of Hedgehog pathway proteins in bladder cancer and determine their association with clinicopathological parameters. The expression of Sonic hedgehog (SHH), its receptor Patched (PTCH1), downstream transcription factor GLI1 and its signal transducer SMO in 112 bladder cancer tissues from Hospital Kuala Lumpur were determined by immunohistochemistry. SHH was expressed in 108 (96.4%) cases, GLI1 in 104 (92.9%) cases, PTCH1 in 111 (99.1%) cases and SMO in all cases. The relationship between the expression of these four proteins and its clinicopathological parameters were analysed statistically. Immunohistochemical staining results showed SHH, GLI1 and SMO proteins were mainly located in the cytoplasm of tumour cells, whereas PTCH1 was mainly located in the nucleus of tumour cells. Positive expression of SHH, PTCH1, GLI1 and SMO proteins were correlated with a few variables which include grade and stage of bladder cancer, lymph node metastasis and distant metastasis (Mx). Among all of the clinicopathological parameters, SMO was found to be the only protein that showed statistically significant association with higher grade ($p=0.001$) and higher stage ($p=0.042$) of bladder cancer. SMO expression also showed borderline association with lymph node metastasis ($p=0.056$). These findings indicate that SMO expression may be a poor prognostic marker in bladder cancer.

Keywords: Hedgehog signalling pathway, Sonic hedgehog, Gli1, Patched, Smo, bladder cancer.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

LALUAN HEDGEHOG PROTEIN DAN HUBUNGKAITNYA DENGAN PARAMETER PATOLOGI KLINIKAL KANSER PUNDI KENCING

Oleh

KHAIRUNNISA BINTI MOHD ARIFFIN

Ogos 2016

Pengerusi : Maizatun Atmadini Abdullah, PhD
Fakulti : Perubatan dan Sains Kesihatan

Laluan hedgehog penting untuk pertumbuhan dan corak semasa perkembangan embrio. Kajian sebelum ini menunjukkan bahawa pengaktifan berterusan laluan hedgehog boleh membawa kepada pelbagai jenis tumor termasuk medulloblastoma, karsinoma sel basal, kanser gastrousus, kanser payudara dan kanser prostat. Kajian ini bertujuan untuk menyiasat ekspresi protein laluan Hedgehog dalam kanser pundi kencing dan menentukan hubungkaitnya dengan parameter patologi klinikal. Ekspresi Sonic hedgehog (SHH), penerima (PTCH), faktor transkripsi, (GLI1) dan isyarat transduser, (SMO) bagi 112 tisu kanser pundi kencing daripada Hospital Kuala Lumpur telah ditentukan dengan teknik immunohistokimia. SHH menunjukkan ekspresi dalam 108 (96.4%) kes, GLI1 dalam 104 (92.9%) kes, PTCH1 dalam 111 (99.1%) kes dan SMO dalam semua kes. Hubungan ekspresi untuk keempat-empat protein dan parameter patologi klinikal telah dianalisa secara statistik. Keputusan pewarnaan immunohistokimia menunjukkan protein SHH, GLI1 dan SMO kebanyakannya terletak di sitoplasma sel-sel tumor, manakala PTCH1 kebanyakannya terletak di dalam nukleus sel-sel tumor. Ekspresi positif daripada SHH, GLI1, PTCH1 dan SMO protein telah dikorelasi dengan beberapa pemboleh ubah yang merangkumi gred dan peringkat kanser pundi kencing, metastasis nodus limfa dan metastasis lebih jauh. Antara semua parameter patologi klinikal, SMO protein sahaja yang menunjukkan hubungkait yang signifikan secara statistik dengan gred tertinggi ($p=0.001$) dan peringkat tertinggi ($p=0.042$) kanser pundi kencing. Ekspresi SMO juga menunjukkan garis batas berkait dengan nodus limfa ($p=0.056$). Keputusan ini menunjukkan bahawa ekspresi SMO mungkin merupakan petanda prognostik yang lemah untuk kanser pundi kencing.

Kata kunci: Laluan hedgehog protein, Sonic hedgehog, GLI1, PTCH1, SMO, kanser pundi kencing.

ACKNOWLEDGEMENTS



Firstly, I would like to thank to Allah who gave me such a great strength and courage throughout this journey and without His blessing I do not think I could complete this thesis successfully.

I would like to acknowledge my special gratitude and appreciation to my supervisor, Dr. Maizatun Atmadini Abdullah for her endless help, support, advice and guidance in completing this research. In addition, I would like to express my thankfulness to my co-supervisor, Dr. Fauzah Abd Ghani and Dr. Salmiah Md Said for their advice and comments throughout the entire project.

My sincere gratitude is also accorded to all staffs of the Department of Pathology, Hospital Kuala Lumpur for their kind assistance during collection of data and samples at the department. Not forgetting all staffs of Histopathology Laboratory, Faculty of Medicine and Health Science for their help in assisting me to use the equipment and facilities required to complete this research.

I would like to express my special gratitude to my beloved mother and my family who gave me such a great support in every possible way and always prayed for me whenever they could. With their support, I was able to complete my research with great success. They are always my supporters ever.

Last but not least, thanks to all my friends who were involved in my research directly or indirectly, with all their helps, this thesis will be able to be completed with flying colours.

I certify that a Thesis Examination Committee has met on 3 August 2016 to conduct the final examination of Khairunnisa binti Mohd Ariffin on her thesis entitled "Hedgehog Pathway Proteins and their Association with Clinicopathological Parameters of Bladder Cancer" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Sabariah binti Md. Noor, PhD

Senior Lecturer
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Norhafizah binti Mohtarrudin, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Siti Aishah Md. Ali, PhD

Professor
Pusat Perubatan Universiti Kebangsaan Malaysia
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 September 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Maizatun Atmadini Abdullah, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Fauzah Abd Ghani, MBBS, M.Path

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Salmiah Md. Said, MD, MPH

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

BUJANG KIM HUAT , PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date :

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No: Khairunnisa Binti Mohd Ariffin (GS37160)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to

Signature: _____

Name of
Chairman of
Supervisory
Committee : Maizatun Atmadini Abdullah, PhD

Signature: _____

Name of
Member of
Supervisory
Committee : Fauzah Abd.Ghani, MBBS, MPath

Signature: _____

Name of
Member of
Supervisory
Committee : Salmiah Md. Said, MD, MPH

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	ii
ACKNOWLEDGEMENTS	iii
APPROVAL	iv
DECLARATION	vi
LIST OF TABLES	xi
LIST OF FIGURES	xii
LIST OF ABBREVIATIONS	xiii
CHAPTER	
1 INTRODUCTION	
1.1 General introduction	1
1.2 Problem statement	2
1.3 Significance of the study	2
1.4 Research objectives	
1.4.1 General objective	3
1.4.2 Specific objectives	3
1.5 Research hypothesis	3
1.6 Conceptual framework	4
2 LITERATURE REVIEWS	
2.1 Anatomy of the urinary bladder	
2.1.1 Gross (macroscopic) appearance of the urinary bladder	5
2.1.2 Histology (microscopic) of the urinary bladder	5
2.2 Background of bladder cancer	
2.2.1 Epidemiology of bladder cancer	6
2.2.2 Risk factors of bladder cancer	7
2.2.3 Histological type of bladder cancer	7
2.2.4 Staging and grading of bladder cancer	8
2.3 Hedgehog (HH) signalling pathway	
2.3.1 An overview of Hedgehog (HH) signalling pathway	11
2.3.2 Function of each of Hedgehog (HH) pathway protein	
2.3.2.1 SHH	11
2.3.2.2 GLI1	11
2.3.2.3 PTCH1	12
2.3.2.4 SMO	12
2.3.3 Mechanism of the HH signalling pathway	12
2.3.4 Aberrant of HH signalling pathway in human cancer	16
2.3.5 Role of HH signalling in bladder development	17
2.3.6 HH signalling in bladder tumourigenesis	18

3	MATERIALS AND METHODS	
3.1	Location of the study	19
3.2	Study design	19
3.3	Sampling	
3.3.1	Study population	19
3.3.2	Sampling frame	19
3.3.3	Sampling unit	19
3.3.4	Sample size	20
3.3.5	Sampling method	
3.3.5.1	Tissue sectioning	20
3.3.5.2	Hematoxylin and Eosin (H&E) staining	20
3.3.5.3	Principles of immunohistochemistry	21
3.3.5.4	Control	21
3.3.5.5	Preparation of tissue biopsy sections for immunohistochemistry	22
3.3.5.6	Antigen retrieval and blocking of endogenous peroxidase activity	22
3.3.5.7	Detection of antigen	22
3.3.5.8	Visualization and counterstaining	23
3.4	Variables	
3.4.1	Dependent variables	24
3.4.2	Independent variables	25
3.5	Data collection	25
3.6	Operational definition of terms	26
3.7	Data analysis	27
3.8	Ethical validation	28
4	RESULTS	
4.1	Demographic distribution of bladder cancer	29
4.2	Immunohistochemical expression of hedgehog signalling proteins in bladder cancer	
4.2.1	Morphology analysis of the samples	31
4.2.2	Analyses of SHH protein expression in bladder cancer	33
4.2.3	Analyses of GLI1 protein expression in bladder Cancer	36
4.2.4	Analyses of PTCH1 protein expression in bladder cancer	39
4.2.5	Analyses of SMO protein expression in bladder cancer	42
4.3	Correlation between HH pathway proteins in bladder cancer	45
4.4	Association of Hedgehog pathway proteins in bladder cancer	
4.4.1	Association of SHH expression with demographic and clinicopathological parameters	46

4.4.2	Association of GLI1 expression with demographic and clinicopathological parameters	48
4.4.3	Association of PTCH1 expression with demographic and clinicopathological parameters	50
4.4.4	Association of SMO expression with demographic and clinicopathological parameters	52
5	DISCUSSION	
5.1	Demographic distribution of bladder cancer	54
5.2	Analyses of HH pathway proteins expression in bladder cancer	54
5.3	Correlation of immunohistochemical expression between HH pathway proteins	55
5.4	Association of demographic factors and clinicopathological parameters with the expression of HH pathway proteins	55
6	CONCLUSION AND FUTURE RECOMMENDATIONS	
6.1	Conclusions	56
6.2	Future recommendations	56
	REFERENCES	57
	APPENDICES	64
	BIODATA OF STUDENT	70

LIST OF TABLES

Table		Page
2.1	Old and new classification of bladder cancer	9
2.2	TNM staging system of bladder cancer	10
2.3	Cancers linked to aberrant of HH signalling pathway	17
3.1	Positive control used in this study	22
3.2	Primary antibodies for immunohistochemical staining	23
3.3	Site of immune-positive staining of each protein	23
3.4	The contextual definition of all dependent study variables	24
3.5	The contextual definitions of all independent study variables	25
3.6	The operational definitions of all dependent study variables	26
3.7	The operational definitions of all independent study variables	27
3.8	Immunoreactivity score system used in this study	28
4.1	Demographic distribution and clinicopathological data of bladder cancer patients in Hospital Kuala Lumpur	30
4.2	Correlation among HH pathway proteins expression in bladder cancer	45
4.3	Association of SHH protein expression with demographic and clinicopathological parameters	47
4.4	Association of GLI1 protein expression with demographic and clinicopathological parameters	49
4.5	Association of PTCH1 protein expression with demographic and clinicopathological parameters	51
4.6	Association of SMO protein expression with demographic and clinicopathological parameters	53

LIST OF FIGURES

Figure		Page
2.1	The anatomy and histology of urinary bladder	6
2.2	Simplified diagram of HH signalling pathway	14
2.3	Diagram of Ligand-dependent signalling and Ligand-independent signalling pathway	15
4.1	Representative photos of the bladder cancer stained with H&E staining	32
4.2	Score distribution of SHH immunoreactivity	33
4.3	Representative photo of tonsil as a positive control of SHH	34
4.4	SHH expression in bladder cancer	35
4.5	Score distribution of GLI1 immunoreactivity	36
4.6	Representative photo of gastric carcinoma as a positive control of GLI1	37
4.7	GLI1 expression in bladder cancer	38
4.8	Score distribution of PTCH1 immunoreactivity	39
4.9	Representative photo of gastric carcinoma as a positive control of PTCH1	40
4.10	PTCH1 expression in bladder cancer	41
4.11	Score distribution of SMO immunoreactivity	42
4.12	Representative photo of gastric carcinoma as a positive control of SMO	43
4.13	SMO expression bladder cancer	44

LIST OF ABBREVIATIONS

WHO	World Health Organization
TCC	Transitional cell carcinoma
HH	Hedgehog
Shh	Sonic hedgehog
PTCH1	Patched homologue 1
SMO	Smoothened homologue
GLI1	GLI family zinc finger 1
mL	mililitre
G1	well differentiated
G2	moderately differentiated
G3	poorly differentiated
WHO/ISUP	World Health Organization/International Society of Urological Pathology
PUNLMP	papillary urothelial neoplasms of low malignant
TNM	Tumour nodes metastasis
AJCC	American Joint Committee on Cancer
Ihh	Indian Hedgehog
Dhh	Desert Hedgehog
SUFU	Suppressor of fused
DNA	Deoxyribonucleic acid
Gli ^R	GLI family zinc finger repressor
Gli ^A	GLI family zinc finger activator
BCC	Basal cell carcinomas
MB	Medulloblastoma
CSCs	Cancer stem cells
FFPE	Formalin-fixed paraffin embedded
H&E	Hematoxylin & Eosin
DPX	Di-N-Butyle Phthalate in Xylene
Ag-Ab	Antigen antibody
ABC	Avidin-biotin complex
LSAB	Labelled streptavidin-biotin
PAP	Peroxidase-antiperoxidase
mM	mili molar
PBS	Phosphate buffered saline
TBST	Tris Buffered Saline with Tween 20
HRP	Horse radish peroxidase
µl	microlitre
DAB	Diaminobenzidine
SPSS	Statistical Package for the Social Sciences
Mx	Distant metastasis

CHAPTER 1

INTRODUCTION

1.1 General Introduction

Cancer is one of the common health problems worldwide. In the United States, one in four deaths is due to cancer (Siegel, Naishadham, & Jemal, 2013). In year 2007, based on Ministry of Health Malaysia Hospitals after Heart Diseases & Diseases of Pulmonary Circulation and Septicaemia, cancer was the third common cause of death in Malaysia. A total of 18,219 new cancer cases were diagnosed in 2007 and registered at the National Cancer Registry. Among the cases reported, there were 8,123 (44.6%) males and 10,096 (55.4%) females (Zainal & Nor Saleha, 2011). Worldwide, bladder cancer ranks eleventh most common malignancy. It is the 7th most common cancer in men and the 19th most common cancer in women (Jacques Ferlay, Isabelle Soerjomataram, Rajesh Dikshit, 2012). The disease is more common in whites than blacks, and the average age at diagnosis is 65 years. In addition, it has been recommended that the stage survival of bladder cancer in female is worse than in male (Kirkali et al., 2005).

Approximately 90% of bladder tumours arise from the urothelial lining, and transitional cell carcinoma (TCC) is the most common histological type in cases arising in the United States. TCC or as known as urothelial carcinoma starts in the cells lining of the bladder and later will spread to the nearby organ or other body parts if it is not treated at early stage. Smoking habit, chemical exposure, age, sex and chronic bladder inflammation are risk factors that can contribute to the development of bladder cancer (Jemal et al., 2011). However, the pathogenesis of bladder cancer still largely remains unknown. Bladder cancer is treated based on the tumour stage and grade. Besides the treatment protocols, the incidence of therapeutic resistance and failure are still present.

Malignant cells aberrantly reactivate oncogenic developmental pathways such as Hedgehog, Wnt and Notch to provide growth and survival advantage to tumour cells (Ingham, McMahon, Ingham, & McMahon, 2001). Emerging evidence clearly suggesting the activation of Hedgehog pathway has been implicated in the tumour development of a large number of human cancers (Beachy, Karhadkar, & Berman, 2004).

The exact genetic events leading to urothelial transformation involve the activation and inactivation of oncogenes, inactivation or loss of tumour suppressor genes and alterations in the apoptotic gene products (Sandberg and Berger, 1994). Study shows that there are a lot of genetic abnormalities but the specific genes involved are still unidentified. This is because there are a lot of gene mutations detectable in bladder cancer (Cantile et al., 2003).

In 2010, Chen et al. have genotyped 177 single-nucleotide polymorphisms (SNP) in 11 SHH pathway genes in a study including 803 bladder cancer cases and 803 controls. They also analysed SNP associations with cancer risk and clinical outcomes in 318

cases of muscle-invasive and 419 cases of non-muscle-invasive bladder cancer and metastatic bladder cancer. As their findings, it has been confirmed that germ-line genetic variations in the SHH pathway could predict the clinical outcomes of non-muscle-invasive bladder cancer patients received transurethral resection and Bacillus Calmette-Guérin.

Hedgehog signalling pathway operates in organogenesis, control of proliferation, and the differentiation of embryonic adult stem cells, thus, not surprisingly it has been linked with several different tumours (Sverrisson et al., 2014) including gastrointestinal cancers and is frequently activated in esophageal (Ingham et al., 2001), pancreatic (Morton et al., 2007) and gastric cancer (Merchant, Saqui-Salces, & El-Zaatari, 2010). On the other hand, the role of hedgehog signalling pathway has been less investigated in the development of bladder cancer and its involvement to the progression of transitional cell carcinoma of bladder is not clear (Haraguchi et al., 2007). However, He et al. successfully proved that the positive expression of SHH, GLI1 and PTCH1 proteins in bladder cancer were significantly higher than in adjacent normal bladder. Ha, Y. S. et al. (2007) has shown that SMO is closely related with the differentiation and progression of bladder cancer. Therefore, SHH, GLI1, PTCH1 and SMO may play important roles during development of bladder cancer (He et al., 2012). This study is necessary to contribute the knowledge in understanding the role of HH pathway proteins in bladder cancer.

1.2 Problem statement

Bladder cancer is generally diagnosed by cystoscopy and biopsy. It has a very high frequency of recurrence and therefore requires follow up cystoscopy as well as urine cytology as periodic surveillance to identify early recurrence. Failure to treat this cancer at an early stage will lead to advance stage which may further complicate the management of the patient. Currently, radical cystectomy is the treatment for urothelial carcinoma. As this cancer needs long-term follow up and surveillance procedure to monitor for tumour recurrence, many of the patients tend to default their follow up. The role of hedgehog signalling pathway in bladder development is well recognized but its role in human bladder cancer progression is uncertain and has been quite controversial. So, this study was done to determine the association of hedgehog pathway protein expression with the progression of bladder cancer so that it can be treated at an early stage with more effective treatments.

1.3 Significance of the study

The findings from this study will contribute knowledge in understanding the role of SHH, GLI1, PTCH1 and SMO in bladder cancer. This study will also offer overview on the expression of these proteins in bladder cancer to better evaluate the function of these proteins in tumour progression. It also provides fundamental background for further investigations on clinical use of related antibodies for personalised cancer management.

1.4 Research objectives

1.4.1 General objective

- To study the HH signalling pathway in bladder cancer.


1.4.2 Specific objectives

- 1) To determine the demographic distribution of bladder cancer in Hospital Kuala Lumpur from 2008 to 2013.
- 2) To analyse the immunohistochemical expression of SHH, GLI1, PTCH1 and SMO in bladder cancer.
- 3) To study the correlation in the immunohistochemical expression between Hedgehog pathway proteins (SHH, GLI1, PTCH1 and SMO) in bladder cancer.
- 4) To study the association of demographic factors (age, gender, ethnicity) and clinicopathological parameters (grade, stage, lymph node metastasis, distant metastasis) of bladder cancer with the expression of hedgehog pathway protein.

1.5 Research hypothesis

- 1) There is high expression of hedgehog pathway proteins (SHH, GLI1, PTCH1 and SMO) in bladder cancer.
- 2) There is significant correlation among the expression of hedgehog signalling proteins (SHH, GLI1, PTCH1 and SMO).
- 3) There is significant association between demographic factor and clinicopathological parameters of bladder cancer with the expression of hedgehog pathway proteins (SHH, GLI1, PTCH1 and SMO).

1.6 Conceptual Framework



REFERENCES

- Al-Hajj, M., & Clarke, M. F. (2004). Self-renewal and solid tumor stem cells. *Oncogene*, 23(43), 7274–7282.
- Ara, J. A. R. A. (2005). Technical Aspects of Immunohistochemistry. *Veterinary Pathology*, 42(4), 405–426.
- Ayers, K. L., & Thérond, P. P. (2010). Evaluating Smoothed as a G-protein-coupled receptor for Hedgehog signalling. *Trends in Cell Biology*, 20(5), 287–298.
- Baris, D., Karagas, M. R., & Verrill, C. et al. (2009). A case-control study of smoking and bladder cancer risk: emergent patterns over time. *Journal of the National Cancer Institute*, 101, 1553–1561.
- Beachy, P. A., Karhadkar, S. S., & Berman, D. M. (2004). Tissue repair and stem cell renewal in carcinogenesis. *Nature*, 432(7015), 324–331.
- Berman, D. M., Karhadkar, S. S., Hallahan, A. R., Pritchard, J. I., Eberhart, C. G., Watkins, D. N., Beachy, P. A. (2002). Medulloblastoma Growth Inhibition by Hedgehog Pathway Blockade. *Science*, 297(5586), 1559–1561.
- Berman, D. M., Karhadkar, S. S., Maitra, A., Montes De Oca, R., Gerstenblith, M. R., Briggs, K., Beachy, P. A. (2003). Widespread requirement for Hedgehog ligand stimulation in growth of digestive tract tumours. *Nature*, 425(6960), 846–851.
- Bitgood, M. J., Shen, L., & McMahon, A. P. (1996). Sertoli cell signaling by Desert hedgehog regulates the male germline. *Current Biology* 6(3), 298–304.
- Braud, F. De, Maffezzini, M., Vitale, V., Bruzzi, P., Gatta, G., Hendry, W. F., Tabor, M. (2002). Bladder cancer. *Crit Rev Oncol Hematol*, 41, 89–106.
- Cantile, M., Cindolo, L., Napadono, G., Altieri, V., Cillo, C. (2003). Hyperexpression of *ocus C* gene in the HOX network is strongly associated in vivo with human bladder transitional cell carcinomas. *Oncogene*, 22, 6562–6468.
- Chalasanani, V., Chin, J. L., & J. I. Izawa. (2009). Histologic variants of urothelial bladder cancer and nonurothelial histology in bladder cancer. *Canadian Urological Association Journal*, 3, S193–198.
- Chan, K. S., Espinosa, I., Chao, M., Wong, D., Ailes, L., & Diehn, M. et al. (2009). Identification, molecular characterization, clinical prognosis, and therapeutic targeting of human bladder tumor-initiating cells. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 14016–14021.
- Chen, C. J., Chuang, Y. C., Lin, T. M., & Wu, H. Y. (1985). Malignant neoplasms among residents of a blackfoot disease-endemic area in Taiwan: high-arsenic artesian well water and cancers. *Cancer Research*, 45, 5895–5899.
- Chen, M., Hildebrandt, M. A., Clague, J. et al. (2010). Genetic variation in the sonic hedgehog pathway affect clinical outcomes in non-muscle-invasive bladder cancer. *Cancer Prev Res (Phila)*, 3, 1235–1245.
- Cheng, W., Yeung, C. K., Ng, Y. K., Zhang, J. R., Hui, C. C., & Kim, P. C. W. (2008). Sonic Hedgehog Mediator *Gli2* Regulates Bladder Mesenchymal Patterning.

Journal of Urology, 180(4), 1543–1550.

- Clark, P. E., Agarwal, N., Biagioli, M. C., Eisenberger, M. A., Greenberg, R. E., Herr, H. W., Wilson, T. G. (2013). NCCN.org Bladder Cancer.
- Coons, A. H., & Kaplan, M. H. (1950). Localization of antigen in tissue cells; improvements in a method for the detection of antigen by means of fluorescent antibody. *The Journal of Experimental Medicine*, 91(1), 1–13.
- Dahmane, N., Lee, J., Robins, P., Heller, P., & Ruiz i Altaba, A. (1997). Activation of the transcription factor Gli1 and the Sonic hedgehog signalling pathway in skin tumours. *Nature*, 389, 876–881.
- DeSouza, K. R., Saha, M., Carpenter, A. R., Scott, M., & McHugh, K. M. (2013). Analysis of the Sonic Hedgehog Signaling Pathway in Normal and Abnormal Bladder Development. *PLoS ONE*, 8(1), e53675.
- Droller, M. J. (2001). *Bladder cancer: current diagnosis and treatment*. Retrieved from <https://books.google.com.my/books?id=Pn31BwAAQBAJ>
- Echelard, Y., Epstein, D. J., St-Jacques, B., Shen, L., Mohler, J., McMahon, J. A., & McMahon, A. P. (1993). Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in the regulation of CNS polarity. *Cell*, 75(7), 1417–1430.
- Edge, S., Byrd, D., & Compton, C. (2010). *AJCC cancer staging manual* (7th ed.). New York: Springer.
- Edward, B. K., Ward, E., & Kohler, B. A. et al. (2010). Annual report to the nation on the status of cancer, 1975– 2006, featuring colorectal cancer trends and impact of interventions (risk factors, screening, and treatment) to reduce future rates. *Cancer*, 116, 544–573.
- Elisabeth, H.-V., Davis, O.-W., & Philip, M.-I. (2000). The Sonic hedgehog-patched-Gli pathway in human development and disease. *American Journal of Human Genetics*, 67, 1047–1054.
- Fajkovic, H., Halpern, J. A., & Cha, E. K. et al. (2011). Impact on gender on bladder cancer incidence, staging and prognosis. *World Journal of Urology*, 29, 457–463.
- Fei, D. L., Sanchez-Mejias, A., Wang, Z., Flaveny, C., Long, J., Singh, S., Robbins, D. J. (2012). Hedgehog signaling regulates bladder cancer growth and tumorigenicity. *Cancer Research*, 72(17), 4449–4458.
- Freedman, N. D., Silverman, D. T., Hollenbeck, A. R., Schatzkin, A., & Abnet, C. C. (2011). Association between smoking and risk of bladder cancer among men and women. *Journal of the American Medical Association*, 306, 737–745.
- Gonnissen, A., Isebaert, S., & Haustermans, K. (2013). Hedgehog signaling in prostate cancer and its therapeutic implication. *International Journal of Molecular Sciences*, 14(7), 13979–14007.
- Goodrich, L. V., Milenković, L., Higgins, K. M., & Scott, M. P. (1997). Altered neural cell fates and medulloblastoma in mouse patched mutants. *Science (New York, N.Y.)*, 277(5329), 1109–1113.

- Gupta, P., Jain, M., Kapoor, R., et al (2009). Impact of age and gender on the clinicopathological characteristics of bladder cancer. *Indian J Urol*, 25, 207-10.
- Gupta, S., Takebe, N., & Lorusso, P. (2010). Targeting the Hedgehog pathway in cancer. *Therapeutic Advances in Medical Oncology*, 2(4), 237–250.
- Ha, Y., Yun, S., Kim, Y., Lee, S., & Kim, W. (2007). 방광암의 예후 인자로서 Smo의 가치 Utility of Smo as a Prognostic Marker for Human Bladder Tumors, 997–1003.
- Hahn, H., Wicking, C., Zaphiropoulos, P. G., Gailani, M. R., Shanley, S., Chidambaram, A., Bale, A. E. (1996). Mutations of the human homolog of Drosophila patched in the nevoid basal cell carcinoma syndrome. *Cell*, 85(6), 841-851.
- Hamed, S., Larue, H., & Hovington, H. (2004). Accelerated Induction of Bladder Cancer in Patched Heterozygous Mutant Mice Advances in Brief Accelerated Induction of Bladder Cancer in Patched Heterozygous Mutant Mice, 1938–1942.
- Haraguchi, R., Motoyama, J., Sasaki, H., Satoh, Y., Miyagawa, S., Nakagata, N., Yamada, G. (2007). Molecular analysis of coordinated bladder and urogenital organ formation by Hedgehog signaling. *Development*, 134, 525–533.
- Hartge, P., Harvey, E. B., Linehan, W.M., et al (1990). Unexplained excess risk of bladder cancer in men. *J Natl Cancer Inst*, 82, 1636-40.
- He, H. C., Chen, J. H., Chen, X. Bin, Qin, G. Q., Cai, C., Liang, Y. X., Zhong, W. De. (2012). Expression of hedgehog pathway components is associated with bladder cancer progression and clinical outcome. *Pathology and Oncology Research*, 18(2), 349–355.
- Horn, E. P., Tucker, Ma., Lambert, G., et al (1995). A study of genderbased cytochrome P450 1A2 variability: A possible mechanism for the male excess of bladder cancer. *Cancer Epidemiol Biomarkers Prev*, 4, 529-33
- Ingham, P. W., & McMahon, A. P. (2001). Hedgehog signaling in animal development. *Genes Dev*, 15(10), 1330-1337.
- Islam, S. S., Mokhtari, R. B., Noman, A. S., Uddin, M., Rahman, M. Z., & Azadi, M. A. et al. (2015). Sonic hedgehog (Shh) signaling promotes tumorigenicity and stemness via activation of epithelial-to-mesenchymal transition (EMT) in bladder cancer. *Molecular Carcinogenesis*, 55(5), 537-551.
- Jacques Ferlay, Isabelle Soerjomataram, Rajesh Dikshit, S. E. (2012). Cancer incidence and mortality worldwide: sources, method and major patterns in Globocan 2012. *International Journal of Cancer*, Vol. 5.
- Jayendran, D., & Manoj, S. (2012). Bladder cancer. Retrieved from <http://www.malaysiaoncology.org/article.php?aid=9>
- Jemal, A., Bray, F., & Ferlay, J. (2011). Global Cancer Statistics: 2011. *CA Cancer J Clin*, 61, 69–90.
- Johnson, R. L., Rothman, a L., Xie, J., Goodrich, L. V, Bare, J. W., Bonifas, J. M., Scott, M. P. (1996). Human homolog of patched, a candidate gene for the basal cell nevus syndrome. *Science (New York, N.Y.)*, 272(5268), 1668–1671.

- Kappler, R., Bauer, R., Calzada-Wack, J., Rosemann, M., Hemmerlein, B., & Hahn, H. (2004). Profiling the molecular difference between Patched- and p53-dependent rhabdomyosarcoma. *Oncogene*, 23, 8785–8795.
- Karhadkar, S. S., Bova, G. S., Abdallah, N., Dhara, S., Gardner, D., Maitra, A., Beachy, P. A. (2004). Hedgehog signalling in prostate regeneration, neoplasia and metastasis. *Nature*, 431(7009), 707–712.
- Kim, J. E., Singh, R. R., Cho-Vega, J. H., Drakos, E., Davuluri, Y., Khokhar, F. A., Vega, F. (2009). Sonic hedgehog signaling proteins and ATP-binding cassette G2 are aberrantly expressed in diffuse large B-cell lymphoma. *Mod Pathol*, 22(10), 1312–1320.
- Kinzler, K. W., Bigner, S. H., Bigner, D. D., Trent, J. M., Law, M. L., O'Brien, S. J., Vogelstein, B. (1987). Identification of an amplified, highly expressed gene in a human glioma. *Science*, 236(70), 70–73.
- Kirkali, Z., Chan, T., Manoharan, M., Algaba, F., Busch, C., Cheng, L., Weider, J. (2005). Bladder cancer: Epidemiology, staging and grading, and diagnosis. *Urology*, 66, 4–34.
- Kogevinas, M., Mannetje, A., & Cordier, S. et al. (2003). Occupation and bladder cancer among men in Western Europe. *Cancer Cause Control*, 14, 907–914.
- Kong, C. H. C., Singam, P., Hong, G. E., Cheok, L. B., Azrif, M., Tamil, A. M., et al. (2010). Clinicopathological features of bladder tumours in a single institution in Malaysia. *Asian Pacific J Cancer Prev.*, 11(1), 149–52.
- Kubo, M., Nakamura, M., Tasaki, A., Yamanaka, N., Nakashima, H., Nomura, M., Katano, M. (2004). Hedgehog signaling pathway is a new therapeutic target for patients with breast cancer. *Cancer Research*, 64(17), 6071–6074.
- Lindemann, R. K. (2008). Stroma-initiated Hedgehog signaling takes center stage in B-cell lymphoma. *Cancer Research*, 68(4), 961–964.
- Liu, C. Z., Yang, J. T., Yoon, J. W., Walterhouse, D., & Iannaccone, P. (1998). Characterization of the promoter region and genomic organization of GLI, a member of the Sonic hedgehog-Patched signaling pathway. *Gene*, 209(1-2), 1–11.
- Luo, W., Li, S., Peng, B., Ye, Y., Deng, X., & Yao, K. (2013). Embryonic Stem Cells Markers SOX2 , OCT4 and Nanog Expression and Their Correlations with Epithelial- Mesenchymal Transition in Nasopharyngeal Carcinoma, 8(2), e56324.
- Malkowicz, S. B., van Poppel, H., Mickisch, G., Pansadoro, V., Thüroff, J., Soloway, M. S., Fukui, I. (2007). Muscle-Invasive Urothelial Carcinoma of the Bladder. *Urology*, 69, 3–16.
- Marigo, V., Roberts, D. J., Lee, S. M., Tsukurov, O., Levi, T., Gastier, J. M., et al. (1995). Cloning, expression and chromosomal location of SHH and IHH: two human homologues of the Drosophila segment polarity gene hedgehog. *Genomics*, 28(1), 44–51.
- Merchant, J. L., Saqui-Salces, M., & El-Zaatari, M. (2010). *Hedgehog signaling in gastric physiology and cancer. Progress in Molecular Biology and Translational*

Science (Vol. 96). Elsevier Inc.

- Montironi, R., & Lopez-Beltran, A. (2005). The 2004 WHO classification of bladder tumors: a summary and commentary. *International Journal of Surgical Pathology*, 13, 143–153.
- Morton, J. P., Mongeau, M. E., Klimstra, D. S., Morris, J. P., Lee, Y. C., Kawaguchi, Y., Lewis, B. C. (2007). Sonic hedgehog acts at multiple stages during pancreatic tumorigenesis. *Proceedings of the National Academy of Sciences of the United States of America*, 104(12), 5103–5108.
- Ning, Z. F., Huang, Y. Z., Lin, T. X., Zhou, Y. X., Jiang, C., & Xu, K. W. et al. (2009). Subpopulation of stem-like cells in side population cells from the human bladder transitional cell cancer cell line T24. *Journal of International Medical Research*, 37, 621–630.
- Nishimaki, H., Kasai, K., Kozaki, K., Takeo, T., Ikeda, H., Saga, S., Itoh, G. (2004). A role of activated Sonic hedgehog signaling for the cellular proliferation of oral squamous cell carcinoma cell line. *Biochemical and Biophysical Research Communications*, 314(2), 313–20.
- Nüsslein-Volhard, C., & Wieschaus, E. (1980). Mutations affecting segment number and polarity in *Drosophila*. *Nature*, 287(5785), 795–801.
- Pasca di Magliano, M., & Hebrok, M. (2003). Hedgehog signalling in cancer formation and maintenance. *Nature Reviews. Cancer*, 3(12), 903–911.
- Reuter, V. (1997). *Urinary bladder, ureter, and renal pelvis*. In: Sternberg SS, ed. *Histology for Pathologists* (2nd ed.). New York: Lippincott-Raven.
- Ruiz I Altaba, A. (1999). Gli proteins encode context-dependent positive and negative functions: implications for development and disease. *Development (Cambridge, England)*, 126(14), 3205–3216.
- Ruiz I Altaba, A., Mas, C., & Stecca, B. (2007). The Gli code: an information nexus regulating cell fate, stemness and cancer. *Trends in Cell Biology*, 17(9), 438-447.
- Sanchez, P., Hernández, A. M., Stecca, B., Kahler, A. J., DeGueme, A. M., Barrett, A., Ruiz i Altaba, A. (2004). Inhibition of prostate cancer proliferation by interference with SONIC HEDGEHOG-GLI1 signaling. *Proceedings of the National Academy of Sciences of the United States of America*, 101(34), 12561–12566.
- Sandberg, A. A., and Berger, C. S. (1994). Review chromosomes studies in urologic tumors cytogenetics and molecular genetics of bladder. *Cancer Journals of Urology*, 151(3), 545–560.
- Scales, S. J., & de Sauvage, F. J. (2009). Mechanisms of Hedgehog pathway activation in cancer and implications for therapy. *Trends in Pharmacological Sciences*, 30(6), 303–312.
- She, J. J., Zhang, P. G., Wang, Z. M., Gan, W. M., & Che, X. M. (2008). Identification of side population cells from bladder cancer cells by dyecycle violet staining. *Cancer Biology and Therapy*, 7(10), 1663–1668.
- Shiroyanagi, Y., Liu, B., Cao, M., Agras, K., Li, J., Hsieh, M. H., ... Baskin, L. S.

- (2007). Urothelial sonic hedgehog signaling plays an important role in bladder smooth muscle formation. *Differentiation; Research in Biological Diversity*, 75(10), 968–77.
- Siegel, R., Naishadham, D., & Jemal, A. (2013). Cancer statistics, 2013. *A Cancer Journal for Clinicians*, 63, 11–30.
- Stanton, B. Z., Peng, L. F., Maloof, N., Nakai, K., Wang, X. et al. (2009). A small molecule that binds Hedgehog and blocks its signaling in human cells. *Nat. Chem. Biol.*, 5(3), 154–156.
- Stenzl, A., Cowan, N. C., & Santis, M. D. et al. (2009). The updated EAU guidelines on muscle-invasive and metastatic bladder cancer. *European Urology*, 55(4), 815–825.
- Sverrisson, E. F., Zens, M. S., Liang Fei, D., Andrews, A., Schned, A., Robbins, D., Seigne, J. D. (2014). Clinicopathological correlates of Gli1 expression in a population-based cohort of patients with newly diagnosed bladder cancer. *Urologic Oncology: Seminars and Original Investigations*, 32(5), 539–545.
- Syed, I. S., Pedram, A., & Farhat, W. A. (2016). Role of Sonic Hedgehog (Shh) Signaling in Bladder Cancer Stemness and Tumorigenesis. *Current Urology Reports*, 17(2), 11.
- Szkandera, J., Kiesslich, T., Haybaeck, J., Gerger, A., & Pichler, M. (2013). Hedgehog signaling pathway in ovarian cancer. *International Journal of Molecular Sciences*, 14(1), 1179–1196.
- Taipale, J., Cooper, M. K., Maiti, T., & Beachy, P. A. (2002). Patched acts catalytically to suppress the activity of Smoothed. *Nature*, 418(6900), 892–7.
- Taylor, M. D., Northcott, P. A., Korshunov, A., Remke, M., et al. (2011). Molecular subgroups of medulloblastoma: the current consensus. *Acta Neuropathologica*, 123(4), 465–472.
- Teglund, S., & Toftgård, R. (2010). Hedgehog beyond medulloblastoma and basal cell carcinoma. *Biochimica et Biophysica Acta*, 1805(2), 181–208.
- Thayer, S. P., di Magliano, M. P., Heiser, P. W., Nielsen, C. M., Roberts, D. J., Lauwers, G. Y., Hebros, M. (2003). Hedgehog is an early and late mediator of pancreatic cancer tumorigenesis. *Nature*, 425(6960), 851–6.
- Theunissen, J. W., & de Sauvage, F. J. (2009). Paracrine Hedgehog signaling in cancer. *Cancer Research*, 69(15), 6007–6010.
- Tian, H., Callahan, C. A., DuPree, K. J., Darbonne, W. C., Ahn, C. P., Scales, S. J., & de Sauvage, F. J. (2009). Hedgehog signaling is restricted to the stromal compartment during pancreatic carcinogenesis. *Proceedings of the National Academy of Sciences of the United States of America*, 106(11), 4254–9.
- Tortora, G. J., & Derrickson, B. H. (2009). *Principle of anatomy and physiology* (12th ed.). Chichester: Wiley.
- Tripathi, K., Mani, C., Barnett, R., Nalluri, S., Bachaboina, L., Rocconi, R. P., Palle, K. (2014). Gli1 protein regulates the S-phase checkpoint in tumor cells via bid protein, and its inhibition sensitizes to DNA Topoisomerase 1 inhibitors. *Journal*

of Biological Chemistry, 289(45), 31513–31525.

- VanPutte, C. L., & Seeley, R. R. (2014). *Seeley's anatomy & physiology*. New York: McGraw-Hill.
- Watkins, D. N., Berman, D. M., Burkholder, S. G., Wang, B., Beachy, P. A., & Baylin, S. B. (2003). Hedgehog signalling within airway epithelial progenitors and in small-cell lung cancer. *Nature*, 422, 313–317.
- Watkins, D. N., & Peacock, C. D. (2004). Hedgehog signalling in foregut malignancy. *Biochemical Pharmacology*, 68, 1055–1060.
- Wayne W. Daniel. (2009). *Biostatistics: A foundation of analysis in the health sciences* (9th ed.). USA: Wiley.
- Wicking, C., Smyth, I., & Bale, A. (1999). The hedgehog signalling pathway in tumorigenesis and development. *Oncogene*, 18(55), 7844–7851.
- Xie, J., Murone, M., Luoh, S. M., Ryan, A., Gu, Q., Zhang, C., de Sauvage, F. J. (1998). Activating Smoothed mutations in sporadic basal-cell carcinoma. *Nature*, 391, 90–92.
- Yauch, R. L., Gould, S. E., Scales, S. J., Tang, T., Tian, H., Ahn, C. P., de Sauvage, F. J. (2008). A paracrine requirement for hedgehog signalling in cancer. *Nature*, 455(7211), 406–410.
- Yoon, J. W., Kita, Y., Frank, D. J., Majewski, R. R., Konicek, B. A., Nobrega, M. A., Iannaccone, P. (2002). Gene expression profiling leads to identification of GLI1-binding elements in target genes and a role for multiple downstream pathways in GLI1-induced cell transformation. *Journal of Biological Chemistry*, 277(7), 5548–5555.
- Zainal, A. O., & Nor Saleha, I. T. (2011). *National Cancer Registry Report. Ministry of Health Malaysia*.
- Zhu, G., Zhau, H. E., Wu, D., Zhang, L., Li, L., He, D., & Chung, L. W. K. (2013). Sonic hedgehog signaling in normal human bladder development. *Journal of Urology*, 189(4), e222.