



UNIVERSITI PUTRA MALAYSIA

***APPLICATION OF FUNCTIONAL METAGENOMICS FOR ISOLATION
AND SCREENING OF ANTIMICROBIAL ACTIVITY OF ANTARCTIC SOIL
MICROORGANISMS***

PREMMALA A/P RANGASAMY

FPSK(M) 2016 68



**APPLICATION OF FUNCTIONAL METAGENOMICS FOR ISOLATION
AND SCREENING OF ANTIMICROBIAL ACTIVITY OF ANTARCTIC
SOIL MICROORGANISMS**

By

PREMMALA A/P RANGASAMY

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

August 2016

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright© Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

**APPLICATION OF FUNCTIONAL METAGENOMICS FOR ISOLATION
AND SCREENING OF ANTIMICROBIAL ACTIVITY OF ANTARCTIC
SOIL MICROORGANISMS**

By

PREMMALA A/P RANGASAMY

August 2016

Chairman : Leslie Than Thian Lung, PhD
Faculty : Medicine and Health Sciences

Thousands of secondary metabolites have been identified from the culture of Gram positive, Gram negative and filamentous fungi that have been isolated from the Antarctic environment and since the year 2002. About 130 to 140 of these natural products have been used in human medicine, veterinary medicine and agriculture as pesticides. Bacteriocin production is also known as common ability among soil bacteria to outcompete competitor for resources including nutrient and water. Such ability is expected to be essential to the survival in the harsh environmental condition in Antarctica. In order to understand if Antarctic soils harbour novel bacteriocins with potential medical application, a functional metagenomic screening need to be conducted using the environmental DNA of the soil. Soil DNA extraction were done by using the Epicenter Meta-g-nome DNA extraction kit and the construction of metagenomic library were done by using the CopyControl Fosmid Library Construction kit from Epicenter. Two libraries were constructed and screened with different types of pathogens using the double agar layer method and incubated overnight at 37°C. Inhibition around the clones were observed the next day. Positive clones were picked for further screening. Positive clones were sent for Illumina next generation sequencing (NGS). The responsible gene that were identified is further cloned into an expression vector to confirm the production of antibiotic activity. Further screening were done by purifying the selected clone with C18 cartridge using the solid phase extraction method (SPE) with different percentage of solvent fractions. SPE products were evaporated using the microcentrifuge vapour evaporator at 30°C and were tested on the bacteria using the well diffusion method at the concentration of 350mg/ml. Inhibition zone were observed. By constructing two separate fosmid libraries which in total encompassed >15, 000 clones, 4 fosmid clones showing positive inhibition against *Klebsiella pneumoniae* were detected. BLAST results based on the DNA contigs generated using illumina NGS revealed that the clones carried periplasmic thiol-disulfide interchange protein DsbA gene and pathways related to isoprenoids for quinones and tyrosine, and phenylalanine

branches from chorismate. Virtual screening using Antibiotics and Secondary Metabolites Shell (Antismash) server corroborated our observation that the selected clones harboured bacteriocin production gene. We have attempted to purify the bacterial product using C-18 solid phase extraction and had obtained fraction (80% water, 20% methanol) which conserved the antibacterial effect against *Klebsiella pneumoniae*. The fraction was sent for a liquid chromatography – mass spectrometry (LCMS) analysis and it showed Nalidixic acid which could be the responsible metabolite. Further screening need to be done.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

APLIKASI METAGENOMIK BERFUNGSI UNTUK PENGASINGAN DAN PEMERIKSAAN AKTIVITI ANTIMIKROBIOLOGI DARIPADA MIKROORGANISMA YANG DIDAPATI DALAM TANAH

Oleh

PREMMALA A/P RANGASAMY

Ogos 2016

Pengerusi : Leslie Than Thian Lung, PhD
Fakulti : Medicine and Health Sciences

Beribu-ribu metabolit sekunder telah dikenal pasti daripada kultur Gram positif, kulat dan Gram negatif yang telah diasingkan daripada persekitaran Antartika dan sejak tahun 2002. Sehingga kini, 130-140 produk-produk semulajadi yang telah digunakan dalam perubatan manusia, perubatan veterinar dan pertanian sebagai racun perosak. Pengeluaran bakteriosin juga dikenali sebagai keupayaan biasa di kalangan bakteria tanah untuk bersaing dengan pesaing untuk keperluan sumber termasuk nutrien dan air. Keupayaan tersebut dijangka menjadi penting untuk terus hidup dalam keadaan alam sekitar yang keras di Antartika. Untuk memahami jika tanah Antartika ialah pelabuhan untuk bacteriocins novel yang mempunyai potensi aplikasi dalam bidang perubatan, pemeriksaan metagenomic yang berfungsi perlu dijalankan dengan menggunakan DNA alam sekitar tanah. Pengekstrakan DNA dari tanah telah dilakukan dengan menggunakan Epicenter Meta-g-nome kit pengekstrakan DNA dan pembinaan perpustakaan metagenomic telah dilakukan dengan menggunakan kit CopyControl Fosmid Perpustakaan Pembinaan daripada Epicenter. Dua perpustakaan telah dibina dan ditapis dengan pelbagai jenis patogen menggunakan kaedah lapisan agar berganda dan dieram semalaman pada 37 ° C. Inhibisi sekitar klon dapat diperhatikan pada hari berikutnya. Klon positif dikumpul untuk pemeriksaan lanjut. Klon positif telah dihantar untuk Illumina Next generation sequencing (NGS). Gen bertanggungjawab yang telah dikenal pasti seterusnya diklon ke dalam vektor ungkapan untuk mengesahkan pengeluaran aktiviti antibiotik. Pemeriksaan selanjutnya dilakukan dengan memurnikan klon yang dipilih dengan kartrij C18 menggunakan kaedah pengekstrakan fasa pepejal (SPE) dengan peratusan yang berbeza pecahan pelarut. Produk SPE telah sejat menggunakan microcentrifuge wap penyejat pada 30 ° C dan telah diuji pada bakteria menggunakan kaedah penyebaran baik pada kepekatan 350mg / ml. Zon perencatan diperhatikan. Dengan membina dua perpustakaan fosmid berasingan yang secara keseluruhan meliputi > 15, 000 klon, 4 fosmid klon menunjukkan perencatan positif terhadap *Klebsiella pneumoniae* dikesan. Keputusan BLAST berdasarkan contigs DNA dijana menggunakan Illumina

NGS mendedahkan bahawa klon dijalankan periplasmic thiol-disulfida pertukaran protein DsbA gen dan laluan yang berkaitan dengan isoprenoids untuk quinone dan tyrosine, dan cawangan phenylalaline dari chorismate. Antismash menyokong pemerhatian kami bahawa klon terpilih memendam bakteriosin gen pengeluaran. Kami telah cuba untuk membersihkan produk bakteria menggunakan C-18 pengekstrakan fasa pepejal dan telah mendapat sebahagian kecil (80% air, 20% metanol) yang mempunyai kesan antibakteria terhadap *Klebsiella pneumoniae*. Pecahan telah dihantar untuk liquid chromatography – mass spectrometry (LCMS) analisis dan ia menunjukkan asid Nalidixic yang boleh menjadi metabolit yang bertanggungjawab. Pemeriksaan lanjut perlu dilakukan.



ACKNOWLEDGEMENTS

I would like to express my sincere gratitude and deepest appreciation to my respectful supervisor Dr. Leslie Than Thian Lung for his continuous support, invaluable guidance, advice and unfailing help throughout my entire research project. His immeasurable kindness and patience are commendable.

I also would like to thank my co-supervisors namely Dr. Chong Chun Wie, Prof. Dr. Peter Convey and Assoc. Prof. Dr. Vasanthakumari Neela for all the humble thoughts, guidance, suggestions and supports throughout my study. In addition, I would like to thank Dr. Ivan Yap Kok Seng and Dr. Cheong Kok Whye for their kind help and assistance in performing solid phase extraction (SPE) and high-performance liquid chromatography (HPLC) experiments.

Not forgetting also to my dedicated lab mates and friends namely Chew Shu Yih, Ng Tzu Shan, Siti Aisya bt Saud Gany, Tan Swee Ching, Pn Suzana, Jocelyn Toh, Wong Li Zhe and many others whom I did not mention here. I would like to thank them for their endless supports, helps and guidance. I am forever grateful to all of them for sharing their experience and knowledge with me throughout the highs and lows of my research project.

Last but not least, I would like to extend my acknowledgements to all the members in International Medical University research office staffs, Media Lab, and Mycology Lab for their kind assistances. Also to all others who have attributed and involved one way another to the successful completion of my study, they are conferred with my sincere appreciation. Finally, a special thanks to my beloved parents for their love, understanding and utmost moral support throughout my study. Thank you.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for degree of Master of Science. The members of the Supervisory Committee were as follows:

Leslie Than Thian Lung, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Vasanthakumari Neela, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Chong Chun Wie, PhD

Lecturer
School of Pharmacy
International Medical University
(Member)

Peter Convey, PhD

Professor
Natural Environment Research Council
British Antarctic Survey
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before the thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other material as stated in the Universiti Putra Malaysia (Research) rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ **Date:** _____

Name and Matric No: Premmala A/P Rangasamy GS35733

Declaration by Members of Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to

Signature: _____
Name of Chairman
of Supervisory
Committee: Leslie Than Thian Lung, PhD

Signature: _____
Name of Member
of Supervisory
Committee: Vasanthakumari Neela, PhD

Signature: _____
Name of Member
of Supervisory
Committee: Chong Chun Wie, PhD

Signature: _____
Name of Member
of Supervisory
Committee: Peter Convey, PhD

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF APPENDICES	xviii
LIST OF ABBREVIATIONS	xix
 CHAPTER	
 1. INTRODUCTION	 1
1.1 Introduction	2
1.2 Objectives	2
 2. LITERATURE REVIEW	 4
2.1 Antibiotics and the limitation of current antimicrobial therapy	4
2.2 Classification and nomenclature of antibiotics	5
2.3 Antibiotic-producing microorganisms	5
2.3.1 Actinomycetes	6
2.3.2 True bacteria and fungi	6
2.3.3 Bacteriocin	7
2.3.3.1 Classification of Bacteriocin	7
2.3.3.2 Bacteriocin vs Antibiotics	9
2.3.4 Thioredoxin Superfamily	9
2.3.4.1 Thiol disulfide protein (DsbA)	10
2.3.4.2 Quinone and quinolone	11
2.4 Biochemistry of Antibiotics	12
2.5 Antarctica	13
2.5.1 Antarctica soils	14
2.6 Psychrophiles	14
2.6.1 Growth at low temperatures	15
2.6.2 Lipid composition and membrane fluidity at low temperatures	15
2.6.3 Protein structure	16
2.6.4 Cold shock and proteins	16
2.7 Metagenomic libraries	16
2.7.1 Function-based screening	18
2.7.2 Sequence-based screening	18
2.7.3 Extraction of environmental genome	19
2.7.4 Vectors	19
2.7.5 Host	19
2.7.6 Metagenomic library screening	20
2.7.6.1 Enzyme discovery	20

2.7.6.2	Antibiotic discovery	20
2.8	Next generation sequencing (NGS)	21
2.9	Solid phase extraction (SPE)	23
2.9.1	Introduction	23
2.9.2	Principle of SPE	23
2.9.3	Methods	23
2.9.3.1	Conditioning	23
2.9.3.2	Loading of sample	24
2.9.3.3	Rising and washing	24
3.	MATERIALS AND METHODS	
3.1	Study design	25
3.1.1	Growth media and culture condition	27
3.1.2	Agarose gel electrophoresis	27
3.1.3	Sample collection	28
3.2	Construction of metagenomic library	28
3.2.1	DNA extraction	28
3.2.2	Filtration, lysis and protein precipitation	28
3.2.3	End-repair of the insert DNA	30
3.2.4	Ligation reaction	30
3.2.5	Packaging the CopyControl Fosmid clones	30
3.2.6	Titering the packaged CopyControl Fosmid clones	31
3.2.7	Colony fast screening for fosmid clones (40kb)	32
3.3	Screening and identification of metagenomic clones	32
3.3.1	Fosmid library induction	32
3.3.2	Double agar overlay assay (antimicrobial screening)	32
3.3.3	Microplate titer assay (96 well plate)	33
3.4	Identifying genes that are responsible for antimicrobial activity	34
3.5	Subcloning and assessing the potential putative genes with antimicrobial activity	34
3.5.1	Restriction digest of PCR product and acceptor Flexi vector	35
3.5.2	Ligation	36
3.5.3	Transformation	36
3.6	Solid phase extraction	37
3.7	High performance - liquid chromatography	38
3.8	Liquid chromatography – mass spectrometry	38
3.9	Statistical analysis	38
4.	RESULTS	39
4.1	Construction of metagenomic library from Sub-Antarctica soil	39
4.2	Screening and identification of the metagenomic library clones with antimicrobial activity	41
4.2.1	Molecular identification and confirmation of clones with antimicrobial properties	45
4.3	Identifying genes that are responsible for antimicrobial activities	45
4.4	Subcloning and assessing the potential putative genes with Antimicrobial activities	53

5. DISCUSSION	
5.1 Construction of metagenomic library from Sub-Antarctica soil	60
5.2 Screening and identification of the metagenomic library clones resulted in inhibition observed in <i>K.pneumoniae</i>	60 61
5.3 Potential gene candidate for antimicrobial activities	63
5.4 Subcloning and assessing the potential putative genes with Antimicrobial activities	63
6. CONCLUSION AND RECOMMENDATIONS FOR FUTURE WORK	
6.1 Conclusions	65
6.2 Recommendation for future research	65
REFERENCES	67
APPENDICES	79
BIODATA OF STUDENT	83

LIST OF TABLES

Table	Page
2.1 Summary of classification of bacteriocins	8
2.2 Comparison between bacteriocins and antibiotics	9
2.3 Example of virulence processes modified by DsbA activity	11
2.4 Classification of quinolone antimicrobial	12
3.1 List of tested microorganisms	27
3.2 The design of DsbA primer	35
3.3 Different percentages of methanol-water fractions for SPE	37
4.1 Concentration of DNA isolated from five different stations from Bird Island, South Georgia, Sub-Antarctic	39
4.2 NCBI BLAST strain identification	46
4.3 Identity of putative Bacteriocin gene CL54 and CL58 inferred from Anti-SMASH	48
4.4 Comparison of compounds identified between CL39, CL54 and CL58 (dark shading indicates presence)	49

LIST OF FIGURES

Figure	Page
2.1 Map of the Antarctic continent and surrounding Southern Ocean, indicating the three commonly recognized terrestrial Biogeographical zones within the region	13
2.2 Construction and screening of metagenomic libraries	17
2.3 Strategy to access and exploit the soil metagenome through the construction screening of DNA libraries derived from soil samples	18
2.4 Schematic illustration of the NGS Illumina flow	22
2.5 Solid phase extraction operation steps	24
3.1 Schematic illustration of study design	26
3.2 Schematic overview of the procedure for isolating DNA from soil using the Meta-G-Nome DNA isolation kit	30
3.3 Schematic overview of the process for preparing a fosmid library using the CopyControl™ Fosmid Library production kit	32
3.4 Schematic illustration of the NGS Illumina flow	32
4.1 Agarose gel electrophoresis of extracted soil DNA with the size of 40kb	39
4.2 Culture plates showing examples of metagenomic libraries constructed from the soil DNA from (A) station 1 (B) station 5	40
4.3 Plate showing inhibition zone around the active clone that was overlaid with <i>K. pneumoniae</i> (A) inhibition zone around a group of clones that have clumped together (B) inhibition zone around two different clones	41
4.4 Representative double agar overlay assay showing the inhibitory growth effect of the selected clone against <i>K.pneumonie</i> on LB agar plate. The clones were with streaked and spot pattern (A) Clone 39 (no inhibition) (B) Clone 40 (no inhibition)	42
4.5 Representative double agar overlay assay showing the inhibitory growth effect of the selected clone against <i>K.pneumonie</i> on LB agar plate. The clones were with streaked and spot pattern (C) Clone 52 and (D) Clone 53	43

4.6	Representative double agar overlay assay showing the inhibitory growth effect of the selected clone against <i>K.pneumonie</i> on LB agar plate. The clones were with streaked and spot pattern (E)Clone 54 and (F) clone 58	44
4.7	Agarose gel electrophoresis of 11 clones been amplified using rep-PCR with msp1 restriction enzyme	45
4.8	Anti-SMASH result of (A) CL54 and (B) CL58	47
4.9	(A) Subsystem features in CL39 generated by RAST	50
	(B) Subsystem features in CL54 generated by RAST	51
	(C) Subsystem features in CL58 generated by RAST	52
4.10	Agarose gel electrophoresis of DsbA present in Sub-Antarctic Soil. These PCR product were amplicons from TS-F and TS-R primer. The PCR product of 424bp indicates the present of DsbA in the soil	53
4.11	Two colonies of DsbA transformed on a LB agar plate supplemented with Ampicillin 100µg/mL	54
4.12	Growth inhibitory effect of the filtered broth on bacteria and yeast	55
4.13	Well diffusion assay of CL54 (SPE semi crude extract, 350mg/mL)	56
4.14	(A) Inhibitory effect of F254+TS and F254 on <i>K.pneumoniae</i> clinical strain evaluated using resazurin assay	57
	(A) Inhibitory effect of F254+TS and F254 on <i>K.pneumoniae</i> BAA 1705 evaluated using resazurin assay	58
4.15	Liquid chromatography – mass spectrometry (LCMS) output of the SPE crude extract	59
5.1	Illustration of the result that has been detected by SSDB	62

LIST OF APPENDICES

Appendix	Page
A Preparation of solutions and reagents	79
B Chamber locations	80



LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
Anti-SMASH	Antibiotics and secondary metabolite analysis Shell
ATCC	American Type Culture Collection
BAC	Bacterial Artificial Chromosomes
BLAST	Basic local alignment search tool
bp	Base pair
CaCl ₂	Calcium chloride
CAMP	Cationic antimicrobial peptide
DAHPh	3-deoxy-D-arabino-heptulosonate-7-phosphate
Dha	Didehydroalanine
Dhb	Didehydrobutyrine
DNA	Deoxyribonucleic acid
dNTP	Deoxyribonucleis triphosphate
Dsb	Disulfide bond
EDTA	Ethylenediaminetetraacetic acid
GCMS	Gas chromatography – mass spectrometry
GPS	Global positioning system
GST	Gluthathione transferase
HPLC	High performance liquid chromatography
LCMS	Liquid chromatography – mass spectrometry
LLE	Liquid – liquid extraction
LB	Luria – Bertani
Lan	Lanthionine
MgCl ₂	Magnesium chloride
MRSA	Methicillin-resistant <i>Staphylococcus aureus</i>
NaOH	Sodium hydroxide
NCBI	National Center of Biotechnology Information
NGS	Next generation sequencing
NMR	Nuclear magnetic resonance
OD	Optical density
ORF	Open reading frame

PBS	Phosphate buffered saline
PCR	Polymerase chain reaction
PDB	Phage dilution buffer
PEP	Phosphenolpyruvate
RAST	Rapid annotation using subsystem technology
RE	Restriction enzyme
RNA	Ribonucleic acid
SDA	Sabouraud dextrose agar
SOC	Super optimal broth with catabolite repression
SPE	Solid phase extraction
SSDB	Sequence similarity data base
TAE	Tris-Acetate-EDTA
TE	Tris-EDTA
TLC	Thin-layer chromatography
TRX	Thioredoxin
UTI	Urinary tract infection
UV	Ultraviolet

CHAPTER 1

INTRODUCTION

Antarctica is a continent almost completely covered by thick ice sheets. The functional roles of bacteria are particularly significant in Antarctica, where terrestrial trophic interactions are generally dominated by microorganisms (Chong et al., 2012; Hogg et al., 2006). It is almost twice the size of Australia, with regions of cold desert soils that have little free water and temperatures that rarely if ever rise above freezing. Antarctica is covered by ice that is on average at least two kilometers in thickness. It is the coldest, driest continent on Earth, and its habitats and organisms have little access to free water. Only during summer are its simple soil and rock habitats free from snow and ice (O'Brien et al., 2004).

The Southern Ocean is a broad physical barrier to the colonization of Antarctica by microorganisms and other biota, compounded by the lack of habitats suitable for their establishment. The islands and ice-free exposures of the sub-Antarctic and maritime Antarctic have somewhat more benign environmental conditions due to their warmer and moister climate, buffered by the surrounding ocean. Temperature, length of growing season and moisture availability play important roles in the colonization of these microorganisms (Smith, 1992).

Antarctic soils have fascinated researchers throughout the last century, being the focus of microbiological and other environmental studies (Pearce et al., 2012). As early as 1903, Elkelof found a significant numbers of bacteria in Antarctic soils, with greater numbers during summer (Boyd & Boyd, 1963).

Prokaryotes dominate many Antarctic ecosystems, where they play a major role in food chains, biogeochemical cycling of nutrients, and the mineralization of pollutants. Antarctica is an important example of an environment dominated by prokaryotes, playing important roles in trophic networks and biogeochemical cycles. Bacteria are very important members of trophic webs, and they are extremely adaptable enabling survival under a wide range of environmental conditions (Russell et al., 1990). There has been a focus on research on cold-adapted microorganisms, generally known as psychrophilic or psychrotolerant, which are capable of growth and reproduction at low temperature. They have specific modifications to their enzymes, membranes and other cellular components which allow them to function at low temperature (Russell et al., 2006).

Thousands of secondary metabolites have been identified from cultures of Gram-positive and Gram-negative bacteria, and filamentous fungi, that have been isolated from the Antarctic environment (Lewis et al., 2010). Since 2002, 130-140 of these secondary metabolites have seen application in human medicine, veterinary medicine and agriculture (Moncheva et al., 2002). Although the first commercialized

antibiotic, penicillin, was discovered by Alexander Fleming by chance, most present-day antibiotics are discovered and developed through systematic searching – ‘bioprospecting’. The systemic screening path introduced by Paul Ehrlich has come to form the core of the search for new drugs in clinical practice and trials, and in the pharmaceutical industry generally (Aminov, 2010).

According to Crag et al. (1997), roughly 60% of the world’s population depends almost completely on plants and natural products from various environmental sources such as water and soils that have been identified as an important source of therapeutically effective medicines. These natural products show biological activity in assays and are generally small molecules with drug-like properties which are capable of being absorbed and metabolized by the body. In order to counter the rapidly increasing challenge of multidrug resistant pathogens such as methicillin resistant *Staphylococcus aureus* (MRSA) (Enright et al., 2002), new antibiotics need to be developed and one of the richest sources is from the uncultured microorganisms of soil. Although many antibiotics are available today, the need remains for the discovery of new compounds to help solve urgent therapeutic challenges such as the continuing problem of the development of drug resistance and new infective properties amongst pathogens, both of which are encouraged by the widespread and often uncontrolled use of antibiotics (Gillespie et al., 2002).

The inability to culture most microorganisms obtained from the environment has frustrated microbiologists for decades (Daniel, 2004). One means of overcoming this challenge is provided by rapid advances in molecular biological approaches. One such advance is the development of the field of metagenomics, which involves the cloning and analysis of genomes without the need for culturing of the source microorganisms (Sabree et al., 2009). This new field offers a path to studying and investigating microbial communities as entire units without cultivating their individual members (Handelsman, 2004).

1.1 Hypothesis

Microorganisms cultured from sub-Antarctic soils using metagenomic screening will produce novel active antimicrobial compounds.

1.2 Objective

General Objective:

To screen and identify prospective compounds with antimicrobial activities from metagenomic library clones constructed from DNA found in sub-Antarctic soil.

Specific objectives

1. To construct a metagenomic library of DNA isolated found in sub-Antarctic soil.
2. To screen for clones which have antimicrobial activities.

3. To identify genes that are responsible for antimicrobial activities.
4. To sub-clone and assess the potential putative genes with antimicrobial activities.



REFERENCES

- Aakvik, T., Lale, R., Liles, M., & Valla, S. (2011). Metagenomic Libraries for Functional Screening. In F. J. d. Bruijn (Ed.), *Handbook of Molecular Microbial Ecology I Metagenomics and Complementary Approaches* (1st ed., pp. 171–173). Hoboken, New Jersey: John Wiley & Sons.
- Abraham, E. P., & Chain, E. (1940). An Enzyme from Bacteria able to Destroy Penicillin. *Nature*, 146(3713), 837–837.
- Adams, B. J., Bardgett, R. D., Ayres, E., Wall, D. H., Aislabie, J., Bamforth, S. Stevens, M. I. (2006). Diversity and distribution of Victoria Land biota. *Soil Biology and Biochemistry*, 38(10), 3003–3018.
- Aminov, R. I. (2010). A brief history of the antibiotic era: lessons learned and challenges for the future. *Frontiers in Microbiology*, 1(December), 134.
- Andriole, V. T. (2005). The Quinolones: Past, Present, and Future. *Clinical Infectious Diseases*, 41(Suppl 2), 113–119.
- Ansorge, W. J. (2009). Next-generation DNA sequencing techniques. *New Biotechnology*, 25(4), 195–203.
- Ashton, P. (1984). A botanical explorer: Frank N. Meyer. *Science (New York, N.Y.)*, 225(4667), 1140–1141.
- B M Duggar, and, & Singleton, V. L. (1953). Biochemistry of Antibiotics. *Annual Review of Biochemistry*, Vol. 22, 459–496.
- Bader, M. W., Xie, T., Yu, C. A., & Bardwell, J. C. a. (2000). Disulfide bonds are generated by quinone reduction. *Journal of Biological Chemistry*, 275(34), 26082–26088.
- Ball, P. (2000). Quinolone generations: natural history or natural selection? *The Journal of Antimicrobial Chemotherapy*, 46, 17–24.
- Barber, M. (1947). Staphylococcal infection due to penicillin-resistant strains. *British Medical Journal*, 2(4534), 863–5.
- Bastos, M. D. C. D. F., Coelho, M. L. V., & Santos, O. C. D. S. (2014). Resistance to bacteriocins produced by Gram-positive bacteria. *Microbiology (Reading, England)*, 161, 683–700.
- Beales, N. (2004). Adaptation of Microorganisms to Cold Temperatures, Weak Acid Preservatives, Low pH, and Osmotic Stress: A Review. *Comprehensive Reviews in Food Science and Food Safety*, 3, 1–20.
- Beasley, S. S., & Saris, P. E. J. (2004). Nisin-Producing *Lactococcus lactis* Strains Isolated from Human Milk. *Applied and Environmental Microbiology*, 70(8), 5051–5053.

- Béjà, O. (2004). To BAC or not to BAC: marine ecogenomics. *Current Opinion in Biotechnology*, 15(3), 187–190.
- Beloqui, A., de María, P. D., Golyshin, P. N., & Ferrer, M. (2008). Recent trends in industrial microbiology. *Current Opinion in Microbiology*, 11(3), 240–248.
- Berdy, J. (2005). Bioactive Microbial Metabolites. *The Journal of Antibiotics*, 58(1), 1–26.
- Berry, E. D., & Foegeding, P. M. (1997). Cold Temperature Adaptation and Growth of Microorganisms. *Journal of Food Protection*, 60(12), 1583–1594.
- Biziuk, M. (2006). Solid Phase Extraction Technique – Trends , Opportunities and Applications. *Polish Journal of Environmental Studies*, 15(5), 677–690.
- Bockheim, J. G. (1997). Properties and Classification of Cold Desert Soils from Antarctica. *Soil Science Society of America Journal*, 61(1), 224.
- Bogomolovas, J., Simon, B., Sattler, M., & Stier, G. (2009). Screening of fusion partners for high yield expression and purification of bioactive viscotoxins. *Protein Expression and Purification*, 64(1), 16–23.
- Bornemann, S., Lawson, D. M., & Thorneley, R. N. F. (2003). A Branch Point in Chorismate Synthase Research. *Structure*, 11(12), 1463–1465.
- Boyd, W. L., & Boyd, J. W. (1963). Soil microorganisms of the McMurdo Sound area, Antarctica. *Applied Microbiology*, 11(2), 116–121.
- Braun, P., Ockhuijsen, C., Eppens, E., Koster, M., Bitter, W., & Tommassen, J. (2001). Maturation of *Pseudomonas aeruginosa* elastase: Formation of the disulfide bonds. *Journal of Biological Chemistry*, 276(28), 26030–26035.
- Butcher, B. G., & Helmann, J. D. (2006). Identification of *Bacillus subtilis* sigma-dependent genes that provide intrinsic resistance to antimicrobial compounds produced by Bacilli. *Molecular Microbiology*, 60(3), 765–82.
- Carvalho, A. P., Fernandes, P. a., & Ramos, M. J. (2006). Similarities and differences in the thioredoxin superfamily. *Progress in Biophysics and Molecular Biology*, 91, 229–248.
- Chong, C. W., Pearce, D. a., Convey, P., Yew, W. C., & Tan, I. K. P. (2012). Patterns in the distribution of soil bacterial 16S rRNA gene sequences from different regions of Antarctica. *Geoderma*, 181-182, 45–55.
- Cleveland, J., Montville, T. J., Nes, I. F., & Chikindas, M. L. (2001). Bacteriocins: Safe, natural antimicrobials for food preservation. *International Journal of Food Microbiology*, 71, 1–20.
- Convey, P. (2011). Antarctic terrestrial biodiversity in a changing world. *Polar Biology*, 34(11), 1629–1641.
- Convey, P., & Stevens, M. I. (2007). Ecology. Antarctic biodiversity. *Science (New*

York, N.Y.), 317(5846), 1877–1878.

- Cottrell, M. T., Moore, J. A., & Kirchman, D. L. (1999). Chitinases from uncultured marine microorganisms. *Applied and Environmental Microbiology*, 65(6), 2553–7.
- Courtois, S., Cappellano, C. M., Ball, M., Francou, F., Normand, P., Martinez, A., Hopke, J. (2003). Recombinant Environmental Libraries Provide Access to Microbial Diversity for Drug Discovery from Natural Products Recombinant Environmental Libraries Provide Access to Microbial Diversity for Drug Discovery from Natural Products. *Applied and Environmental Microbiology*, 69(1), 49–55.
- Cowan, D. A., Russell, N. J., Mamais, A., & Sheppard, D. M. (2002). Antarctic Dry Valley mineral soils contain unexpectedly high levels of microbial biomass. *Extremophiles : Life under Extreme Conditions*, 6(5), 431–6
- Cowan, D., Meyer, Q., Stafford, W., Muyanga, S., Cameron, R., & Wittwer, P. (2005). Metagenomic gene discovery: past, present and future. *Trends in Biotechnology*, 23(6), 321–329.
- Cui, X. L., Mao, P. H., Zeng, M., Li, W. J., Zhang, L. P., Xu, L. H., & Jiang, C. L. (2001). *Streptimonospora salina* gen. nov., sp. nov., a new member of the family Nocardiopsaceae. *International Journal of Systematic and Evolutionary Microbiology*, 51(2), 357–363.
- D’Costa, V. M., McGrann, K. M., Hughes, D. W., & Wright, G. D. (2006). Sampling the antibiotic resistome. *Science (New York, N.Y.)*, 311(5759), 374–377.
- Dairi, T. (2005). Studies on biosynthetic genes and enzymes of isoprenoids produced by actinomycetes. *Japanese Journal of Antibiotics*, 58(1), 87–98.
- Dairi, T., Hamano, Y., Kuzuyama, T., Itoh, N., Furihata, K., & Seto, H. (2001). Eubacterial diterpene cyclase genes essential for production of the isoprenoid antibiotic terpentecin. *Journal of Bacteriology*, 183(20), 6085–94.
- Daniel, R. (2004). The soil metagenome--a rich resource for the discovery of novel natural products. *Current Opinion in Biotechnology*, 15(3), 199–204.
- Daniel, R. (2005). The metagenomics of soil. *Nature Reviews. Microbiology*, 3(6), 470–478.
- Davies, J., & Davies, D. (2010). Origins and evolution of antibiotic resistance. *Microbiology and Molecular Biology Reviews : MMBR*, 74(3), 417–433.
- De Voss, J. J., Rutter, K., Schroeder, B. G., Su, H., Zhu, Y., & Barry, C. E. (2000). The salicylate-derived mycobactin siderophores of *Mycobacterium tuberculosis* are essential for growth in macrophages. *Proceedings of the National Academy of Sciences*, 97(3), 1252–1257.
- Deegan, L. H., Cotter, P. D., Hill, C., & Ross, P. (2006). Bacteriocins: Biological tools for bio-preservation and shelf-life extension. *International Dairy Journal*,

16, 1058–1071.

- Demain, A. L. (2009). Antibiotics: natural products essential to human health. *Medicinal Research Reviews*, 29(6), 821–842.
- Demain, A. L., & Sanchez, S. (2009). Microbial drug discovery: 80 years of progress. *The Journal of Antibiotics*, 62(1), 5–16.
- Dorenbos, R., Stein, T., Kabel, J., Bruand, C., Bolhuis, A., Bron, S., ... Van Dijl, J. M. (2002a). Thiol-disulfide oxidoreductases are essential for the production of the lantibiotic sublancin 168. *The Journal of Biological Chemistry*, 277(19), 16682–8.
- Dorenbos, R., Stein, T., Kabel, J., Bruand, C., Bolhuis, A., Bron, S., ... Van Dijl, J. M. (2002b). Thiol-disulfide oxidoreductases are essential for the production of the lantibiotic sublancin 168. *The Journal of Biological Chemistry*, 277(19), 16682–8.
- Drider, D., Fimland, G., Héchard, Y., McMullen, L. M., & Prévost, H. (2006). The continuing story of class IIa bacteriocins. *Microbiology and Molecular Biology Reviews : MMBR*, 70(2), 564–582.
- Ennahar, S., Sashihara, T., Sonomoto, K., & Ishizaki, A. (2000). Class IIa bacteriocins: Biosynthesis, structure and activity. *FEMS Microbiology Reviews*, 24, 85–106.
- Enright, M. C., Robinson, D. A., Randle, G., Feil, E. J., Grundmann, H., & Spratt, B. G. (2002). The evolutionary history of methicillin-resistant *Staphylococcus aureus* (MRSA). *Proceedings of the National Academy of Sciences of the United States of America*, 99(11), 7687–7692.
- Fairbairn, D. J., & Law, B. A. (1986). Proteinases of psychrotrophic bacteria: their production, properties, effects and control. *Journal of Dairy Research*, 53(01), 139–177.
- Ferrer, M., Beloqui, A., Timmis, K. N., & Golyshin, P. N. (2009). Metagenomics for mining new genetic resources of microbial communities. *Journal of Molecular Microbiology and Biotechnology*, 16(1-2), 109–123.
- Field, D., Cotter, P., Hill, C., & Ross, R. P. (2007). Bacteriocin Biosynthesis, Structure and Function. In M. A. Riley & O. Gillor (Eds.), *Bacteriocins Current Research and Applications* (p. 218). Norfolk: Horizon Scientific Press.
- Fontaine, L., & Hols, P. (2008a). The inhibitory spectrum of thermophilin 9 from *Streptococcus thermophilus* LMD-9 depends on the production of multiple peptides and the activity of BlpG(St), a thiol-disulfide oxidase. *Applied and Environmental Microbiology*, 74(4), 1102–10.
- Fontaine, L., & Hols, P. (2008b). The inhibitory spectrum of thermophilin 9 from *Streptococcus thermophilus* LMD-9 depends on the production of multiple peptides and the activity of BlpG(St), a thiol-disulfide oxidase. *Applied and Environmental Microbiology*, 74(4), 1102–10.

- Fraga, M. F., & Esteller, M. (2007). Epigenetics and aging: the targets and the marks. *Trends in Genetics : TIG*, 23(8), 413–8.
- Gamenara, D., Seoane, G., Mendez, P. S., & Maria, P. D. de. (2012). Enzymes Involved in Redox Reactions: Natural Sources and Mechanistic Overview. In *Redox Biocatalyst: Fundamentals and Applications* (pp. 1–86). John Wiley & Sons.
- Godlewska-zylkiewicz, B. (2003). Biosorption of platinum and palladium for their separation y preconcentration prior to graphite furnace atomic absorption. *Spectrochimica Acta*, 58, 1531–1540.
- Gounot, A. M. (1991). Bacterial life at low temperature: physiological aspects and biotechnological implications. *The Journal of Applied Bacteriology*, 71(5), 386–397.
- Handelsman, J. (2004). Metagenomics : Application of Genomics to Uncultured Microorganisms Metagenomics : Application of Genomics to Uncultured Microorganisms. *Microbiology and Molecular Review*, 68(4), 669–685.
- Handelsmanl, J., Rondon, M. R., Goodman, R. M., Brady, S. F., & Clardy, J. (1998). Molecular biological access to the chemistry of unknown soil microbes: a new frontier for natural products. *Chemistry and Biology*, 5(10), 245–249.
- Hansen, J. N. (1993). Antibiotics synthesized by posttranslational modification. *Annual Review of Microbiology*, 47, 535–564.
- Heeb, S., Fletcher, M. P., Chhabra, S. R., Diggle, S. P., Williams, P., & Cámara, M. (2011). Quinolones: From antibiotics to autoinducers. *FEMS Microbiology Reviews*, 35(2), 247–274.
- Hennion, M. C. (1999). Solid-phase extraction: method development, sorbents, and coupling with liquid chromatography. *Journal of Chromatography. A*, 856, 3–54.
- Heras, B., Shouldice, S. R., Totsika, M., Scanlon, M. J., Schembri, M. a, & Martin, J. L. (2009). DSB proteins and bacterial pathogenicity. *Nature Reviews. Microbiology*, 7(3), 215–225.
- Hogg, I. D., Craig Cary, S., Convey, P., Newsham, K. K., O'Donnell, A. G., Adams, B. J., Wall, D. H. (2006). Biotic interactions in Antarctic terrestrial ecosystems: Are they a factor? *Soil Biology and Biochemistry*, 38(10), 3035–3040.
- Holmgren, A. (1985). Thioredoxin. *Annual Review of Biochemistry*, 54, 237–271.
- Hurst, A. (1981). *Advances in Applied Microbiology Volume 27. Advances in Applied Microbiology* (Vol. 27). Elsevier.
- Inglis, R. F., Bayramoglu, B., Gillor, O., & Ackermann, M. (2013). The role of bacteriocins as selfish genetic elements. *Biology Letters*, 9(3), 20121173–20121173.

- Ito, K., & Inaba, K. (2008). The disulfide bond formation (Dsb) system. *Current Opinion in Structural Biology*, 18(4), 450–458.
- Iyalomhe, G. B. S., Iyalomhe, S. I., & Eholor, R. E. (2011). Antibiotic prescription and resistance: A contemporary literature review. *International Journal of Medicine and Medical Sciences*, 3(14), 376–380.
- Jacob-Dubuisson, F., Pinkner, J., Xu, Z., Striker, R., Padmanabhan, A., & Hultgren, S. J. (1994). PapD chaperone function in pilus biogenesis depends on oxidant and chaperone-like activities of DsbA. *Proceedings of the National Academy of Sciences of the United States of America*, 91(24), 11552–11556.
- Jaenicke, R., Heber, U., Franks, F., Chapman, D., Griffin, M. C. A., Hvidt, A., & Cowan, D. A. (1990). Protein Structure and Function at Low Temperatures [and Discussion]. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 326(1237), 535–553.
- Jones, P. G., Cashel, M., Glaser, G., & Neidhardt, F. C. (1992). temperature downshifts in Escherichia coli . Function of a Relaxed-Like State following Temperature Downshifts in Escherichia coli. *Journal of Bacteriology*, 174(12), 3903–3914.
- Kawasaki, T., Hayashi, Y., Kuzuyama, T., Furihata, K., Itoh, N., Seto, H., & Dairi, T. (2006). Biosynthesis of a natural polyketide-isoprenoid hybrid compound, furaquinocin A: identification and heterologous expression of the gene cluster. *Journal of Bacteriology*, 188(4), 1236–44.
- Kimura, N. (2006). Minireview Metagenomics : Access to Unculturable Microbes in the Environment. *Microbes and Environments*, 21(4), 201–215.
- Klaenhammer, T. R. (1993). Genetics of bacteriocins produced by lactic acid bacteria. *FEMS Microbiology Reviews*, 12, 39–85.
- Koehn, F. E., & Carter, G. T. (2005). The evolving role of natural products in drug discovery. *Nature Reviews. Drug Discovery*, 4(3), 206–220.
- Koma, D., Yamanaka, H., Moriyoshi, K., Ohmoto, T., & Sakai, K. (2012). Production of aromatic compounds by metabolically engineered Escherichia coli with an expanded shikimate pathway. *Applied and Environmental Microbiology*, 78(17), 6203–6216.
- LaVallie, E. R., DiBlasio, E. A., Kovacic, S., Grant, K. L., Schendel, P. F., & McCoy, J. M. (1993). A thioredoxin gene fusion expression system that circumvents inclusion body formation in the E. coli cytoplasm. *Bio/technology (Nature Publishing Company)*, 11(2), 187–193.
- Lee, H. S., Kwon, K. K., Kang, S. G., Cha, S.-S., Kim, S.-J., & Lee, J.-H. (2010). Approaches for novel enzyme discovery from marine environments. *Current Opinion in Biotechnology*, 21(3), 353–357.
- Leveau, J. H. J. (2007). The Magic and Menace of Metagenomics : prospects for the study of plant growth-promoting rhizobacteria. *Eur J.Plant Pathol*, 119, 279–

- Lewis, K., Epstein, S., D'Onofrio, A., & Ling, L. L. (2010). Uncultured microorganisms as a source of secondary metabolites. *The Journal of Antibiotics*, 63(8), 468–476.
- Li, H. R., Yu, Y., Luo, W., & Zeng, Y. X. (2010). *Marisediminicola antarctica* gen. nov., sp. nov., an actinobacterium isolated from the Antarctic. *International Journal of Systematic and Evolutionary Microbiology*, 60(11), 2535–2539.
- Li, Y. (2009). Carrier proteins for fusion expression of antimicrobial peptides in *Escherichia coli*. *Biotechnology and Applied Biochemistry*, 54(1), 1–9.
- Lim, H. K., Chung, E. J., Kim, J., Ja, G., Jang, K. S., Chung, Y. R., Cho, K. Y. (2005). Characterization of a Forest Soil Metagenome Clone That Confers Indirubin and Indigo Production on *Escherichia coli*. *Applied and Environmental Microbiology*, 71(12), 7768–7777.
- Lorenz, P., & Schleper, C. (2002). Metagenome—a challenging source of enzyme discovery. *Journal of Molecular Catalysis B: Enzymatic*, 19-20, 13–19.
- Mardis, E. R. (2008). Next-generation DNA sequencing methods. *Annual Review of Genomics and Human Genetics*, 9, 387–402.
- Martinez, A., Kolvek, S. J., Lai, C., Yip, T., Hopke, J., Brown, K. A., ... Osburne, M. S. (2004). Genetically Modified Bacterial Strains and Novel Bacterial Artificial Chromosome Shuttle Vectors for Constructing Environmental Libraries and Detecting Heterologous Natural Products in Multiple Expression Hosts. *APPLIED AND ENVIRONMENTAL MICROBIOLOGY*, 70(4), 2452–2463.
- Martinez, A., & S.Osburne, M. (2013). Preparation of Fosmid Libraries and Functional Metagenomic Analysis. In E. DeLong (Ed.), *Microbial Metagenomics, Metatranscriptomics, and Metaproteomics* (Volume 531., pp. 124–139). Academic Press, 2013.
- Messens, J., & Collet, J.-F. (2006). Pathways of disulfide bond formation in *Escherichia coli*. *The International Journal of Biochemistry & Cell Biology*, 38(7), 1050–1062.
- Mevs, U., Stackebrandt, E., Schumann, P., Gallikowski, C. a., & Hirsch, P. (2000). *Modestobacter multiseptatus* gen. nov., sp. nov., a budding actinomycete from soils of the Asgard Range (Transantarctic Mountains). *International Journal of Systematic and Evolutionary Microbiology*, 50(1), 337–346.
- Meyerdierks, A., & Glockner, F. O. (2010). Metagenome Analysis. In J. M. Cock (Ed.), *Introduction to Marine Genomics* (Volume 1 o., pp. 33–46). Springer Science & Business Media.
- Missiakas, D., & Raina, S. (1997). Protein folding in the bacterial periplasm. *Journal of Bacteriology*, 179(8), 2465–2471.

Moldoveanu, S. C., & David, V. (2002). *Sample Preparation in Chromatography*. Elsevier.

Moncheva, P., Tishkov, S., Dimitrova, N., Chipeva, V., Antonova-nikolova, S., & Bogatzevska, N. (2002). CHARACTERISTICS OF SOIL ACTINOMYCETES FROM ANTARCTICA. *Journal of Culture Collections*, 3, 3–14.

Morozova, O., & Marra, M. a. (2008). Applications of next-generation sequencing technologies in functional genomics. *Genomics*, 92, 255–264.

Myrcha, A., & Tatur, A. (1991). Ecological role of the current and abandoned penguin rookeries in the land environment of the maritime Antarctic. *Polish Polar Research*, 12(1), 3–24.

Nowicka, B., & Kruk, J. (2010). Occurrence, biosynthesis and function of isoprenoid quinones. *Biochimica et Biophysica Acta - Bioenergetics*, 1797(9), 1587–1605.

O'Brien, A., Sharp, R., Russell, N. J., & Roller, S. (2004). Antarctic bacteria inhibit growth of food-borne microorganisms at low temperatures. *FEMS Microbiology Ecology*, 48(2), 157–167.

Overbeek, R., Olson, R., Pusch, G. D., Olsen, G. J., Davis, J. J., Disz, T., ... Stevens, R. (2014). The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Research*, 42(D1), 206–214.

Paik, S. H., Chakicherla, A., & Hansen, J. N. (1998). Identification and Characterization of the Structural and Transporter Genes for, and the Chemical and Biological Properties of, Sublancin 168, a Novel Lantibiotic Produced by *Bacillus subtilis* 168. *Journal of Biological Chemistry*, 273(36), 23134–23142.

Pan, S., Tan, G., Convey, P., Pearce, D., & Tan, I. (2013). Diversity and bioactivity of actinomycetes from Signy Island terrestrial soils, maritime Antarctic. *Advances in Polar Sciences*, 24(4), 208–212.

Pawliszyn, J. (Ed.). (2002). *Sampling and Sample Preparation for Field and Laboratory: Fundamentals and New Directions in Sample Preparation, Volume 37* (Volume 37.). Amsterdam: Elsevier.

Pearce, D. a, Newsham, K. K., Thorne, M. a S., Calvo-Bado, L., Krsek, M., Laskaris, P., Wellington, E. M. (2012). Metagenomic analysis of a southern maritime antarctic soil. *Frontiers in Microbiology*, 3(December), 403.

Peláez, J., Alarcón, J., ... Bouza, E. (2008). Metronidazole Resistance in *Clostridium difficile* Is Heterogeneous. *JOURNAL OF CLINICAL MICROBIOLOGY*, 46(9), 3028–3032.

Pettit, R. K. (2004). Soil DNA libraries for anticancer drug discovery. *Cancer Chemotherapy and Pharmacology*, 54(1), 1–6.

Pitchandi, P., Hopper, W., & Rao, R. (2013). Comprehensive database of Chorismate synthase enzyme from shikimate pathway in pathogenic bacteria. *BMC*

- Poffé, R., & Mertens, W. (1988). Heat-stable proteases of psychrotrophic bacteria isolated from cooled raw milk. *Applied Microbiology and Biotechnology*, 27(5-6), 437–442.
- R.N.J Comans, & Roskam, G. D. (2002). *Methodologies in Soil and Sediment Fractionation Studies: Single and Sequential Extraction Procedures*. (P. Quevauviller, Ed.). Cambridge: Royal Society of Chemistry.
- Riley, M. A., & Wertz, J. E. (2002). Bacteriocins: Evolution, Ecology, and Application. *Annual Review of Microbiology*, 56(1), 117–137.
- Rohmer, M., Grosdemange-Billiard, C., Seemann, M., & Tritsch, D. (2004). Isoprenoid biosynthesis as a novel target for antibacterial and antiparasitic drugs. *Current Opinion in Investigational Drugs (London, England : 2000)*, 5(2), 154–162.
- Rupp, M. E., & Fey, P. D. (2003). Extended spectrum beta-lactamase (ESBL)-producing Enterobacteriaceae: considerations for diagnosis, prevention and drug treatment. *Drugs*, 63(4), 353–65.
- Russell, N. J. (1997). Psychrophilic bacteria--molecular adaptations of membrane lipids. *Comparative Biochemistry and Physiology. Part A, Physiology*, 118(3), 489–493.
- Russell, N. J., Harrison, P., Johnston, I. a., Jaenicke, R., Zuber, M., Franks, F., & Wynn-Williams, D. (1990). Cold Adaptation of Microorganisms [and Discussion]. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 326(1237), 595–611.
- Russell, N. J., London, I. C., Campus, W., & Tn, K. (2006). Antarctic micro-organisms : coming in from the cold. *Oxoid*, 27(2).
- Sabree, Z. L., Rondon, M. R., & Handelsman, J. (2009). Metagenomics. *Gentic, Genomics*, 41(7), 622–632.
- Sar, P., & Islam, E. (2012). Metagenomic Approaches in Microbial Bioremediation of Metals and Radionuclides. In T.Satyanarayana, B. N. Johri, & A. Prakash (Eds.), *Microorganisms in Environmental Management: Microbes and Environment* (p. 525). Springer Science & Business Media.
- Sauvonnet, N., & Pugsley, A. P. (1998). The requirement for DsbA in pullulanase secretion is independent of disulphide bond formation in the enzyme. *Molecular Microbiology*, 27(3), 661–667.
- Schaechter, M. (2010). Sequenced Based Metagenomic Analysis. In *Desk Encyclopedia of Microbiology* (2nd ed., p. 754). Academic Press, 2010.
- Schmeisser, C., Steele, H., & Streit, W. R. (2007). Metagenomics, biotechnology with non-culturable microbes. *Applied Microbiology and Biotechnology*, 75(5), 955–962.

- Schnier, J. (1987). The Synthesis of Heat-shock Proteins after a Decrease in Translational Capacity in *Escherichia coli*. *Microbiology*, 133(11), 3151–3158.
- Scholz, M. B., Lo, C.-C., & Chain, P. S. G. (2012). Next generation sequencing and bioinformatic bottlenecks: the current state of metagenomic data analysis. *Current Opinion in Biotechnology*, 23(1), 9–15.
- Schumann, P., Prauser, H., Rainey, F. a, Stackebrandt, E., & Hirsch, P. (1997). *Friedmanniella antarctica* gen. nov., sp. nov., an LL-diaminopimelic acid-containing actinomycete from Antarctic sandstone. *International Journal of Systematic Bacteriology*, 47(2), 278–283.
- Schuster, S. C. (2008). Next-generation sequencing transforms today's biology. *Nature Methods*, 5(1), 16–8.
- Shaikh, S., Fatima, J., Shakil, S., Rizvi, S. M. D., & Kamal, M. A. (2015). Antibiotic resistance and extended spectrum beta-lactamases: Types, epidemiology and treatment. *Saudi Journal of Biological Sciences*, 22(1), 90–101.
- Sheibani, N. (1999). Prokaryotic gene fusion expression systems and their use in structural and functional studies of proteins. *Preparative Biochemistry & Biotechnology*, 29(1), 77–90.
- Shin-Young Park, Yun-Jeong Heo, Young-Seok Choi, Eric Déziel, Y.-H. C. (2005). *Pseudomonas aeruginosa* *Bacillus subtilis*. *The Journal of Microbiology*, 43(5), 443–450.
- Shouldice, S. R., Heras, B., Walden, P. M., Totsika, M., Schembri, M. a, & Martin, J. L. (2011). Structure and function of DsbA, a key bacterial oxidative folding catalyst. *Antioxidants & Redox Signaling*, 14(9), 1729–1760.
- Simon, C., & Daniel, R. (2011). Metagenomic analyses: past and future trends. *Applied and Environmental Microbiology*, 77(4), 1153–1161.
- Simpson, N. J. K. (2010). *Solid-Phase Extraction: Principles, Techniques, and Applications*. CRC Press.
- Singh, B. K., & Macdonald, C. a. (2010). Drug discovery from uncultivable microorganisms. *Drug Discovery Today*, 15(17-18), 792–799.
- Smith, H. (1992). Distribution and ecology of the testate rhizopod fauna of the continental Antarctic zone. *Polar Biology*, 12(6), 629–634.
- Smith, V. R. (2002). Climate change in the sub-Antarctic: An illustration from Marion Island. *Climatic Change*, 52(3), 345–357.
- Stein, T. (2005). *Bacillus subtilis* antibiotics: structures, syntheses and specific functions. *Molecular Microbiology*, 56(4), 845–857.
- Stenson, T. H. (2002). DsbA and DsbC Are Required for Secretion of Pertussis Toxin by *Bordetella pertussis*. *Infection and Immunity*, 70(5), 2297–2303.

- Stenson, T. H., & Weiss, A. A. (2002). DsbA and DsbC Are Required for Secretion of Pertussis Toxin by *Bordetella pertussis*. *INFECTION AND IMMUNITY*, 70(5), 2297–2303.
- Streit, W. R., & Schmitz, R. A. (2004). Metagenomics--the key to the uncultured microbes. *Current Opinion in Microbiology*, 7(5), 492–498.
- Torok, E., Moran, E., & Cooke, F. (2009). *Oxford Handbook of Infectious Diseases and Microbiology*. OUP Oxford.
- Vaklu, J., Ambardar, S., & Johri, B. N. (2012). Analysis of Isolated Metagenomic DNA for Taxonomic and Metabolic Diversity. In T. Satyanarayana, B. N. Johri, & Anil Prakash (Eds.), *Microorganisms in Sustainable Agriculture and Biotechnology Life sciences* (illustrate., p. 280). Springer Science & Business Media, 2012.
- Vandamme, E. J. (2010). Industrial microbiology. *Current Opinion in Microbiology*, 13(3), 253–254.
- Varma, A., & Oelmüller, R. (2007). Metagenomics. In *Advanced Techniques in Soil Microbiology* (volume 11., p. 371). Springer Science & Business Media.
- Vlamiş-Gardikas, A., & Holmgren, A. (2002). Thioredoxin and glutaredoxin isoforms. *Methods in Enzymology*, 347, 286–296.
- Waksman, S. a, Schatz, A., & Reynolds, D. M. (2010). Production of antibiotic substances by actinomycetes. *Annals of the New York Academy of Sciences*, 1213, 112–124.
- Wang, G.-Y.-S., Graziani, E., Waters, B., Pan, W., Li, X., McDermott, J., ... Davies, J. (2000). Novel Natural Products from Soil DNA Libraries in a Streptomyces Host. *Organic Letters*, 2(16), 2401–2404.
- Ward, N. (2006). New directions and interactions in metagenomics research. *FEMS Microbiology Ecology*, 55(3), 331–338.
- Workinger, M. S., Huber, C. J., Crais, E. R., Campbell, S. K., Dalton, J. M., Murray, E. a, & Features Submission, H. C. (1989). Antarctica: soils, weathering processes and environment. *Antarctic Science*, 1(1), 77–82
- Wright, G. D. (2005). Bacterial resistance to antibiotics: enzymatic degradation and modification. *Advanced Drug Delivery Reviews*, 57(10), 1451–1470.
- Wülfing, C., & Rappuoli, R. (1997). Efficient production of heat-labile enterotoxin mutant proteins by overexpression of dsbA in a degP-deficient *Escherichia coli* strain. *Archives of Microbiology*, 167(5), 280–293.
- Xu, J. (2006). Microbial ecology in the age of genomics and metagenomics: concepts, tools, and recent advances. *Molecular Ecology*, 15(7), 1713–1731.
- Yu, J. (1998). Inactivation of DsbA, but not DsbC and DsbD, affects the intracellular survival and virulence of *Shigella flexneri*. *Infection and Immunity*, 66(8),

3909–3917.

Yu, J., & Kroll, J. S. (1999). DsbA: A protein-folding catalyst contributing to bacterial virulence. *Microbes and Infection*, 1(14), 1221–1228.

Zeller, T., & Klug, G. (2006). Thioredoxins in bacteria: Functions in oxidative stress response and regulation of thioredoxin genes. *Naturwissenschaften*, 93(6), 259–266.

Zheng, G., Yan, L. Z., Vederas, J. C., & Zuber, P. (1999). Genes of the sbo-alb locus of *Bacillus subtilis* are required for production of the antilisterial bacteriocin subtilisin. *Journal of Bacteriology*, 181(23), 7346–55.

