



**UNIVERSITI PUTRA MALAYSIA**

**EFFECTS OF IRRADIANCE AND CUTTING INTERVAL ON GROWTH,  
YIELD, PERSISTENCE AND NUTRITIVE VALUE OF *ASYSTASIA*  
*INTRUSA***

**SUPARJO NOORDIN MOKHTAR**

**FP 2008 35**



**EFFECTS OF IRRADIANCE AND CUTTING  
INTERVAL ON GROWTH, YIELD,  
PERSISTENCE AND NUTRITIVE VALUE OF  
*ASYSTASIA INTRUSA***

**SUPARJO NOORDIN MOKHTAR**

**DOCTOR OF PHILOSOPHY  
UNIVERSITY PUTRA MALAYSIA**

**2008**



**EFFECTS OF IRRADIANCE AND CUTTING INTERVAL ON GROWTH,  
YIELD, PERSISTENCE AND NUTRITIVE VALUE OF *ASYSTASIA INTRUSA***

By

**SUPARJO NOORDIN MOKHTAR**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

**JULY 2008**



## DEDICATION

*Years have passed with memories like yesterday  
It seems like you're still around the same in every way  
Though the tears have barely dried from what we tried to say  
With daily thoughts of you in our hearts and in our prayer.*

*We remember the charm the light the spirit you left behind  
The love of life the love that binds  
The love that only you Noor Hailan Sulaiman can find  
And teach us to love and to love divine.*

*Atiqah and Hanum need you, and Arief and Azrin miss you  
Haikal asks for you and Suparjo has to live without you  
They say life for us must go on when life left you  
But how can we begin to find another you?*

*The loss in you and the loss we feel  
But will time heal what our hearts reveal?  
The Lord's way of life is meant to be  
Is what life we are meant to see.*

*None had been a better mother and wife  
If only you had some more years of life  
We cherish the memory we hold dear  
Darling mother and wife you had no peer.*

*Wassallam.*

Source: A modification of poem by James, Melissa, Julian and Jonathan in the Sun  
30<sup>th</sup> August 1996, page 43.

Suparjo Noordin Mokhtar  
July 2008



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Doctor of Philosophy

**EFFECTS OF IRRADIANCE AND CUTTING INTERVAL ON GROWTH, YIELD, PERSISTENCE AND NUTRITIVE VALUE OF *ASYSTASIA INTRUSA***

**By**

**SUPARJO NOORDIN MOKHTAR**

**July 2008**

**Chairman: Associate Professor Mohd Ridzwan Abd Halim, PhD.**

**Faculty: Agriculture**

*Asystasia intrusa*, a species introduced 90 years ago, is now a naturalized weed in Malaysia that is prevalent under oil palm plantations. However, this species is known to be very palatable to cattle and goats. Despite being recognized as a shade plant, its response to shade has not been studied. Similarly, the appropriate cutting interval for optimum yield and quality has not been determined. A study was, therefore, conducted to determine the performance of animals consuming *A. intrusa* and the effects of shading and cutting interval on its yield, chemical composition, growth and persistence.

Experiments were conducted encompassing the; i) intake and nutritive value of *A. intrusa* in herbivores when harvested from the field; ii) effects of cutting interval and shade levels on yield, growth, persistence of survival, chemical composition, in vitro dry matter digestibility (IVDMD), essential amino acid (EAA) and mineral



concentrations of *A. intrusa* grown in a glasshouse; and iii) optimum light requirements for photosynthesis.

*Asystasia intrusa* is very palatable and nutritious for ruminant and non-ruminant herbivores. Adult rams consumed *A. intrusa* dry matter (DM) at a rate of 33 g kg<sup>-1</sup> body weight. Both acid detergent fiber (ADF) and neutral detergent fiber (NDF) were digested at a rate of 590 g kg<sup>-1</sup> DM. The digestible energy (DE) and dry matter digestibility (DMD) were 2.1 mcal kg<sup>-1</sup> DM and 620 g kg<sup>-1</sup> DM, respectively. In rabbits, the DE and DMD value in this experiment was 2.2 mcal kg<sup>-1</sup> DM and 509 g kg<sup>-1</sup> DM, respectively. The mean IVDMD was 612 g kg<sup>-1</sup> DM for leaf and 451 g kg<sup>-1</sup> DM for stem which was within the normal digestibility of most forages. The digestibility of CP was quite low with 496 g kg<sup>-1</sup> CP in rabbits and 500 g kg<sup>-1</sup> CP for rams. As a result, *A. intrusa* had DCP of 68 g kg<sup>-1</sup> DM in rams and 101 g kg<sup>-1</sup> DM in rabbits.

The CP concentration in leaf, stem and aerial part was 326, 167, and 253 g kg<sup>-1</sup> DM, respectively. Both cutting interval and shade had an effect on the crude protein (CP) concentration. Crude protein in leaf decreased from 354 g kg<sup>-1</sup> DM at 4 week cutting interval to 303 g kg<sup>-1</sup> DM at 12 weeks. In stem, CP concentration of young *A. intrusa* was 213 g kg<sup>-1</sup> DM which declined to 144 g kg<sup>-1</sup> DM at longer cutting interval. They showed no difference between 8 and 12 week-cutting interval. On the other hand, with increased shading, the leaf CP concentration increased from 278 g kg<sup>-1</sup> DM in the open to 361 g kg<sup>-1</sup> DM at 93% shade. Shade has no effect on the CP concentration of stem.

An essential amino acid profile of leaves showed that cystine concentration decreased with higher shade levels, while histidine increased with longer cutting interval. The most limiting EAA were histidine, lysine and methionine in leaf, and histidine and methionine in stem. Lysine concentration of stem was higher than leaf, especially when cut early. Essential amino acid constituted 35.2% of the leaf CP, and 37.7% of stem CP.

Mineral concentrations increased with more shade and longer cutting interval in leaf and stem, except for Mg in leaf. Similarly, the P concentration in leaf increased with age. The ratio of Ca:P was 4.3 in leaf and 5.9 in stem, thus formulation of ration for calcium would be made easier.

The concentration of fiber in *A. intrusa* was low relative to other tropical forages. The concentration of ADF in leaf, stem and aerial parts of *A. intrusa* were 158, 393 and 258 g kg<sup>-1</sup> DM, respectively. Neutral detergent fiber in leaf, stem and aerial parts were 463, 536 and 494 g kg<sup>-1</sup> DM, respectively. The ADF, as well as NDF, was highest with long cutting interval in both leaf and stem. They showed no difference in concentrations between 8 and 12 week cutting interval. Shading had contrasting effects on ADF and NDF and on leaf and stem. In the leaf, the ADF decreased while the NDF increased with increasing shade. In contrast, stem ADF increased as shade increased.

The photosynthetic rate of *A. intrusa* increased when incident light was increased from 59  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  to a peak at 600  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  after which it declined. This shows that it is a shade plant with sun tolerant property. The light

compensation point for *A. intrusa* occurred at  $81.2 \mu\text{Mol photons m}^{-2} \text{ s}^{-1}$ . The water use efficiency of *A. intrusa* was 1.46. The efficiency of *A. intrusa* to convert solar energy to chemical energy was 0.018. Shade and cutting intervals affected the morphology and growth of *A. intrusa*. Increasing shade reduced the number of branches and nodes. Plant was shorter (15.5 cm) in the open than in heavy shade (27.0 cm). They flower at all height, with shorter plants tending to have more flowers. In contrast to flower, seed production of *A. intrusa* was higher in taller plants. Canopy size and leafiness was influenced more by cutting interval than by shade. The cumulative annual yield of leaf was only affected by shade, while in stem, both cutting interval and shade affected the annual yield.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah.

**KESAN CAHAYA DAN KERAPAN POTONGAN KEATAS TUMBESARAN,  
PENGELUARAN, KETAHANAN DAN NILAI MAKANAN *ASYSTASIA*  
*INTRUSA***

**Oleh**

**SUPARJO NOORDIN MOKHTAR**

**Julai 2008**

**Pengerusi : Profesor Madya Mohd Ridzwan Abd Halim, PhD.**

**Fakulti : Pertanian**

*Asystasia intrusa* yang diperkenal 90 tahun dahulu dari luar negara kini menjadi rumpai alamiah di dalam kebun kelapa sawit. Rumpai ini sangat disukai lembu, dan kambing. Walaupun dikenali sebagai pokok yang suka naungan, tindak balas naungan keatas tumbuhan tidak diketahui. Juga, umur potongan yang sesuai untuk pengeluaran yang optima dan mutu yang baik belum ditentukan. Satu kajian telah dijalankan untuk mengkaji nilai prestasi ternakan yang memakannya dan respon *A. intrusa* kepada naungan dan umur potongan ke atas pengeluaran bahan kering, kandungan kimia, pertumbuhan dan ketahanan.

Uji kaji merangkumi i) kadar pengambilan oleh ternakan, dan nilai pemakanan *A. intrusa* yang dituai dari lading, ii) kesan umur pemotongan dan naungan terhadap pengeluaran, pertumbuhan, ketahanan dari mati, kandungan kimia, pencernaan in



vitro, dan kandungan asid amino perlu dan galian pada *A. intrusa* yang ditanam di dalam rumah kaca, dan, iii) keperluan cahaya untuk proses fotosintesis.

Ternakan ruminan herbivor dan bukan herbivor menggemari *A. intrusa* kerana zatnya. Biri-biri jantan dewasa memakan bahan kering *A. Intrusa* pada kadar 33 g kg<sup>-1</sup> berat badan. Kadar cerna kedua-dua ADF dan NDF ialah 590 g kg<sup>-1</sup> BK. Ia mengandungi tenaga dan BKT masing-masing sebanyak 2.1 mcal kg<sup>-1</sup> BK dan 620 g kg<sup>-1</sup> BK. Nilai tenaga dan BKT pada arnab ialah masing-masing 2.2 mcal kg<sup>-1</sup> BK dan 509 g kg<sup>-1</sup> BK. Berat kering tercerna daun dan batang masing-masing pada kadar 612 g kg<sup>-1</sup> BK dan 451 g kg<sup>-1</sup> BK menggunakan cara invitro meletakkan ia setanding dengan rumput lain. Protin kasar dicerna arnab adalah pada kadar 496 g kg<sup>-1</sup> PK dan 500 g kg<sup>-1</sup> PK pada biri-biri. Dengan itu, terdapat PKT sebanyak 68 g kg<sup>-1</sup> BK dari *A. intrusa* bagi biri-biri jantan dan 101 g kg<sup>-1</sup> BK bagi arnab.

Daun, batang dan bahagian atas tumbuhan ini masing-masing mengandungi PK pada kadar 326 g kg<sup>-1</sup> BK, 167 g kg<sup>-1</sup> BK dan 253 g kg<sup>-1</sup> BK. Kedua-dua umur potongan dan naungan memberi kesan kepada kandungan protein kasar. Protin kasar dalam daun menurun dari 354 g kg<sup>-1</sup> BK pada 4 minggu umur potongan ke 303 g kg<sup>-1</sup> BK pada umur 12 minggu. Batang muda *A. intrusa* mengandungi PK yang tinggi sebanyak 213 g kg<sup>-1</sup> BK dan menurun hingga 144 g kg<sup>-1</sup> apabila lebih tua. Tidak terdapat perbezaan kandungan PK antara umur potongan 8 dan 12 minggu. Sebaliknya, kandungan sebanyak 278 g kg<sup>-1</sup> DM protin kasar pada daun di tempat terang meningkat ke 361 g kg<sup>-1</sup> DM pada 93% naungan. Naungan tidak memberi kesan kepada batang.

Profil asid amino perlu dalam daun menunjukkan penurunan kandungan cistena dengan peningkatan naungan. Histidina pula bertambah dengan kenaikan umur potongan. Asid amino perlu yang paling terhad ialah histidina, lisina dan metionina pada daun, dan histidina dan metionina pada batang. Batang mengandungi lisina yang lebih tinggi daripada daun apabila dipotong awal. Kandungan asid amino perlu dalam daun dan batang masing-masing ialah 35.2% dan 37.7% daripada kandungan PK.

Kandungan galian dalam daun dan batang pada amnya bertambah dengan peningkatan naungan dan umur potongan, kecuali Mg pada daun. Juga, kandungan P dalam daun menurun dengan peningkatan umur potongan. Kadar Ca:P ialah 4.3 pada daun dan 5.9 pada batang memudahkan proses membuat ransum makanan.

*Asystasia intrusa* mempunyai serabut yang rendah berbanding dengan forage lain dari iklim tropika. Bahagian daun, batang dan atas tumbuhan masing-masing mengandungi ADF sebanyak 158, 393 dan 258 g kg<sup>-1</sup> BK. Manakala kandungan NDF pada bahagian daun, batang dan atas tumbuhan masing-masing mengandungi 463, 536 dan 494 g kg<sup>-1</sup> BK. Kandungan ADF dan juga NDF sentiasa meningkat dengan pertambahan umur potongan pada kedua daun berlaku juga pada batang. Peningkatan itu tidak perbezaan di antara potongan 8 dan 12 minggu. Kesan naungan keatas serabut pula bergantung kepada samada ia ADF atau NDF dan samada daun atau batang. Kandungan ADF menurun dan NDF meningkat jika ia daun dengan penambahan kegelapan. Sebaliknya, ADF pada batang meningkat dalam gelap.

Kadar fotosintesis *A. intrusa* mulai cahaya sebanyak  $59 \mu\text{mol photons m}^{-1} \text{ s}^{-1}$  meningkat ke kemuncak pada  $600 \mu\text{mol photons m}^{-1} \text{ s}^{-1}$  dan kembali turun selepas itu. Ini menunjukkan ia pokok jenis hidup bawah naungan tetapi boleh tahan matahari. Titik pampasan cahaya untuk *A. intrusa* ialah  $81.2 \mu\text{Mol photons m}^{-2} \text{ s}^{-1}$ . Kecekapan kegunaan cahaya *A. intrusa* ialah 1.46. Kecekapan *A. intrusa* menukar tenaga suria ke tenaga kimia ialah 0.018. Naungan dan umur potongan memberi kesan kepada morfologi dan pertumbuhan *A. intrusa*. Bilangan buku-ruas dan cabang menurun dengan naungan. *Asystasia intrusa* tumbuh rendah tanpa naungan (15.5 cm) berbanding naungan gelap (27.0 cm). Tumbuhan akan sentiasa berbunga pada semua ketinggian, terutamanya apabila rendah. Bertentangan dengan bunga, biji banyak kelihatan pada *A. intrusa* yang tinggi. Saiz kanopi ditentukan oleh umur potongan dan bukan oleh naungan. Kerimbunannya ditentukan oleh umur potongan dan bukan oleh naungan. Naungan memberi kesan kepada bahan kering terkumpul dalam setahun pada daun, manakala kedua-dua umur potongan dan naungan memberi kesan pada batang.

## ACKNOWLEDGEMENTS

In the name of Allah, the most compassionate, and the most merciful. All praise upon Allah (*swt*). Selawat and salam to the Prophet Muhammad (*saw*), his family and close companions. Bless Allah (*swt*) for his innayah that the thesis finally completed after a long ten years. Though slow, I was happy as it ran without blunders, as would when doing things in a rush. And, I would like to thank many people who made this thesis possible.

Before that, I felt satisfied about the choice of *A. intrusa* as a theme for the thesis. Historically, I stumbled on the species by coincidence. All the while, I have been watching for a plant in Malaysia which is nutritively comparable with alfalfa. In 1986 when touring MARDI with FELDA trainees, Dr Wong Choi Chee showed an *A. intrusa* weed that could reach 240 g kg<sup>-1</sup> DM protein. I immediately jumped on it. My experience with 18 dairy farmers in USA studying alfalfa hay quality committed me to *A. intrusa*. A high crude protein value and low ADF content was the quality sought when buying hay lots.

Besides, in late 1970s I observed something was lacking in the way animals were fed. For every two wagons of fodder fed to goats, one wagon of left-over remained. It was painful to see time and effort being wasted on securing feed yet the animals was skinny. That was where I began to develop interest on quality of roughage.

Later on, it developed to the effect of the climate on the quality of roughage, as well as on the animal itself.



I thanked Allah swt who bestowed many good people who aided me whenever needed. Many individuals have to be thanked for their assistance. All of them contributed the best they could. Among them were:

Abd Rahim Hamid MARDI : Former Staff, Livestock Department Office.  
Abdul Aziz Mahat, MARDI : Dry matter disappearance.  
Abdul Halim Radzi, MARDI : LCA-2 infrared gas analyzer for photosynthesis.  
Abdul Rahim Parman, MARDI : Computer scanning.  
Abdul Rashid bin Dimin, MARDI : Computer problem.  
Ahmad Affandi Kamaruddin, MARDI : Ketua Pusat Komputer.  
Ahmad Shokri Othman, MARDI : Statistical Analysis.  
Ahmad Yusof Hamali, MARDI : Deputy Director, Livestock Research Center.  
Ahmad Tajuddin Zainuddin, MARDI : Former Director of Livestock Research.  
Ahmad Zam Zam Mohamad, Dr., MARDI : Former Director of Central Analytical Service.  
Aidah Sahim, MARDI : Computer Department  
Chan Kam Seng, MARDI : LCA-2 infrared gas analyzer for photosynthesis.  
Cek Tah Md Yasin, MARDI : Former Staff, Livestock Department Office  
Chen Chin Peng, MARDI : Discussion  
Daud bin Endam, MARDI : Computer problem.  
Engku Azahan bin Engku Ahmed, Dr : Former Deputy Director, Livestock Nutrition Program.  
Faizah Mohamed, MARDI : Preparation & Analysis of samples and NIR.  
Hamidon Bahari, MARDI (late) : Data management in IBM 4282 mainframe.  
Hilaliah : Computer Department  
Jamaluddin Salim : Pengarah, MARDI Negeri Johor.  
Kamaruzaman Saman, MARDI : Former Staff, Livestock Department Office.  
Mahyuddin Dahan, Dr., UPM : Inspiration and advice.  
Malek Yaacob, MARDI : GC of amino acids.  
Maslan bin Sabtu, MARDI : Retrieval of problematic files.  
Md Sharif Ahmad, Dr., MARDI : Former Director General.  
Md Yunus Jaafar, MARDI : Non-linear.  
Moh Esa Saiman, MARDI : GC of amino acids.  
Mohamed Ariff Omar, Dr., MARDI : Former Director of Livestock Research.  
Mohamed Aziz Dollah, Dr., MARDI : Inspiration and advice.  
Mohamed Khusahry bin Mohamed Yusuf, Dr., MARDI : Former Director, Strategic Livestock Research Department.  
Mohd Hafiz Ibrahim, Plant Physiology Graduate Student, UPM : Photosynthesis Tutor  
Mohd Nor Abdullah, MARDI : Chemical analysis.  
Mohd Othman Rahmat, MARDI : Chemical analysis.  
Mohd Ramli Kadir, MARDI : Chemical analysis – Bomb calorimeter.



Mohd Rum Anuar : Computer  
 Mohd Salleh Subi : Former, Livestock Department Office  
 Mohd Senawi Tamin Dr., MARDI : Encouragement and comments.  
 Mohd Yusoff bin Abdullah, Dr., Director, Strategic Resource Research Departement,  
 MARDI : LCA-2 infrared gas analyzer for photosynthesis.  
 Mohd. Yunus Ismail, MARDI : Sample drying and preparation.  
 Mohd Zahar bin Abu Samah, MARDI : Computer problem.  
 Morshidi Mostapa, MARDI : PC problems.  
 Nantha Kumaran, MARDI : Analysis & preparation.  
 Noorlaila Zainal : Former, Livestock Department Office  
 Normah Ab Majid, MARDI :  
 Norzaina Abbas, MARDI (late) : Livestock Department Office  
 Omar Yahyer, MARDI : Collection & measurement.  
 Osman Omar, MARDI : Supply of Asystasia  
 Ramadesan, K., Dr., MARDI : Encouragement and comments.  
 Ramli Mohd Noor, MARDI : Supply of Asystasia.  
 Rasidah Seman, MARDI : Former, Livestock Department Office  
 Robiah Sarit, MARDI : Livestock Department Office  
 Rohaya Ismail, MARDI : Former Staff, Livestock Department Office  
 Rosiah bt Hamzah, MARDI : Publication, Technical Services Center.  
 Rosminah Sodar, MARDI : Chemical analysis.  
 Rosnizah Ismail, MARDI : Chemical analysis.  
 Rozinita Jalaluddin, MARDI : Livestock Department Office  
 Roziyah Majid, MARDI : Computer scanning.  
 Saari Awang, MARDI : Former Staff, Livestock Department Office  
 Sabariah Samsuddin, MARDI : Chemical analysis.  
 Saridah Pilus, MARDI : Chemical analysis.  
 Samiyah Mohamad Nasir : Literature.  
 Sarimah Kahar, MARDI : Computer problem.  
 Selamat Basar, MARDI : Former Staff, Livestock Department Office.  
 Siti Rubinah bt Senin, MARDI : Computer service.  
 Subahir Samijan, MARDI : LCA-2 infrared gas analyzer for photosynthesis.  
 Syed Jalaluddin bin Syed Salim, Dr : Former Vice Chancellor, UPM as former  
 Chairman of Examination Committee.  
 Thani Hussien, MARDI (late) : Computer graphics.  
 Wan Zahari Mohamed, Dr., MARDI : Former Head of Nutrition Program, Director  
 of Livestock Research Department, MARDI.  
 Wong Choi Chee, Dr., MARDI : Supply of Literature.  
 Yusof Hashim, Dr., MARDI : Former Director General.  
 Zainal Abidin A Rahman, MARDI : Collection & measurement.  
 Zainol Abd Aziz, MARDI : LCA-2 infrared gas analyzer for photosynthesis.

Also, sincere thanks to Associate Professor Dr Ridzwan Halim, my advisor, for the patience in guiding me throughout the course fulfilling the requirements of the university.

Similar thank is also indebted to Professor Dr Razak Alimon and Senior Research



Officer Dr Abu Bakar Chik, who are members of the supervisory committee, for continually encouraging me to complete the study. Also, thanks to Dr Mohamad Aziz Dollah who initially pursued me to join the course. The thesis would not have been possible if Professor Dr Syed Jalaluddin bin Syed Salim, Vice Chancellor, UPM, have not envisioned the search for new fodder crops as feed for ruminants in this young country of mine.

Lastly, my family contributed many ways in laboring the thesis. I honor my wife Noor Hailan Sulaiman whose character is analogous to the deeds of Siti Khadijah, wife of prophet Muhammad (*saw*). Among others, she abides and conforms to her husband while others would protests. She shares her wealth while others would dissents. She is honest, gentle and loving. I was fortunate to have a wife that is “*menyenangkan hati suami*“. I was just a passenger in the family’s high reputation built by her. However, she departed early and could not see the completed thesis. Al-fatiha to her. Doa for her soul to be among the solihah, sins be forgiven, and may she rest in peace. In a time like this, my children Joanna Azrin, Muhammad Arief, Farah Hanum, Nur Atiqah and Nuzul Haikal, and I could only doa her.





I certify that a Thesis Examination Committee has met on 18<sup>th</sup> July 2008 to conduct the final examination of Suparjo Noordin Mokhtar on his thesis entitled “Effects of Irradiance and Cutting Interval on Growth, Yield, Persistence and Nutritive Value of *Asystasia intrusa*” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

**Mohd. Razi Ismail, PhD.**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Dr Dahlan Ismail, PhD.**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Dr Hawa Jaafar, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Kevin D. Kephart, PhD**

Professor  
Ag Experiment Station  
South Dakota University  
United States of America  
(External Examiner)

---

**BUJANG BIN KIM HUAT, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 28 April 2009

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirements for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Mohd Ridzwan Abd Halim, PhD.**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Abdul Razak Alimon, PhD.**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Abu Bakar bin Chik, PhD.**

Senior Research Officer  
MARDI  
(Member)

---

**Hasanah Mohd. Ghazali, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 14 May 2009

## **DECLARATION**

I hereby declare that the thesis is based on my original work except for quotations and citations that have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

---

**SUPARJO NOORDIN MOKHTAR**

Date: 30 June 2009

## TABLE OF CONTENTS

	<b>Page</b>
<b>DEDICATION</b>	i
<b>ABSTRACT</b>	ii
<b>ABSTRAK</b>	vi
<b>ACKNOWLEDGEMENTS</b>	x
<b>APPROVAL</b>	xiv
<b>DECLARATION</b>	xvi
<b>LIST OF TABLES</b>	xvii
<b>LIST OF FIGURES</b>	xx
<b>LIST OF PLATES</b>	xxiv
<b>LIST OF ABBREVIATIONS / NOTATIONS / GLOSSARY OF TERMS</b>	xxv
<b>LIST OF APPENDICES</b>	xxviii
<b>TABLE OF CONTENTS</b>	xxxix
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	<b>5</b>
2.1 Introduction	5
2.2 <i>Asystasia intrusa</i>	5
2.2.1 Taxonomy / Botany	9
2.2.2 Plant Introduction	12
2.2.3 Forage Quality	14
2.2.4 Uses of <i>A. intrusa</i>	16
2.3 Plant Composition and Forage Quality	18
2.3.1 Chemical Composition	18
2.3.1.1 Cell Wall	19
2.3.1.2 Detergent Fiber Analysis	21
2.3.1.3 Lignin	23
2.3.1.4 Available Carbohydrates	24
2.3.1.5 Crude Protein	25
2.3.1.6 Minerals	27
2.3.2 Leaf to Stem Ratio	29
2.3.3 Digestibility	31
2.3.4 Intake	34
2.3.5 Species	35
2.4 Effects of Light on Plant Growth and Quality	36
2.4.1 Day Length	38
2.4.2 Sun and Shade Plants	39
2.4.3 C <sub>3</sub> and C <sub>4</sub> Plants	43
2.4.4 Sun Flecks	45
2.4.5 Artificial Shade Cloth	46
2.4.6 The Shade Avoidance Effect	46
2.4.7 Photosynthetically Active Radiation (PAR)	47



	<b>Page</b>	
2.4.8	Height	49
2.4.9	Node	50
2.4.10	Dry Matter Yield	51
2.4.11	Light Effects on Crude Protein Concentration	53
2.4.12	ADF, NDF and Water Soluble Carbohydrates	54
2.4.13	Lignin	56
2.5	Effect of Temperature on Forage Growth, Yield and Quality	57
2.5.1	Growth	59
2.5.2	ADF, NDF and Water Soluble Carbohydrates	60
2.5.3	Lignin	60
2.5.4	Crude Protein	61
2.5.5	Minerals	61
2.6	Effect of Age of Cutting on Plant Growth, Yield and Quality	61
2.6.1	Root	64
2.6.2	Mineral	65
2.6.3	Crude Protein	65
2.6.4	ADF, NDF and Water Soluble Carbohydrates	67
2.6.5	Yield	68
2.7	Effect of Soil Moisture on Forage Growth, Yield and Quality	72
2.8	Persistence	72
<b>3</b>	<b>GENERAL MATERIALS AND METHODS</b>	<b>76</b>
3.1	Experimental Outline	76
3.1.1	Experiment 1 (Chapter 4)	76
3.1.2	Experiment 2 (Chapter 5)	76
3.1.3	Experiment 3 (Chapter 6)	77
3.1.4	Experiment 4 (Chapter 7)	77
3.1.5	Experiment 5 (Chapter 8)	77
<b>4</b>	<b>QUALITY OF <i>A. INTRUSA</i> AS ANIMAL FEED</b>	<b>78</b>
4.1	Experiment 1 <b>Nutritive Value and Digestibility of <i>A. intrusa</i> in Rams</b>	78
4.1.1	Introduction	78
4.1.2	Material and Method	79
4.1.2.1	Hay Lots	79
4.1.2.2	Animals	80
4.1.2.3	General Procedure	80
4.1.2.4	Intake Measurement and Fecal Collection	81
4.1.2.5	Sample Preparation	83
4.1.2.6	Calculation	84
4.1.2.7	Chemical Analysis	85
4.1.3	Results and Discussion	86
4.1.4	Conclusion	92

	<b>Page</b>	
4.2	<b>Experiment II : Nutritive Value and Digestibility of A. intrusa in Rabbits.</b>	93
4.2.1	Introduction	93
4.2.2	Materials and Methods	93
	4.2.2.1 Digestibility Trial	93
	4.2.2.2 Performance Trial	95
4.2.3	Results and Discussion	97
	4.2.3.1 Digestibility Trial in Rabbit	97
	4.2.3.2 Performance Trial in Rabbit	99
4.2.4	Conclusion	101
5	<b>THE EFFECT OF SHADE AND CUTTING INTERVAL ON THE GROWTH AND YIELD OF A. INTRUSA</b>	102
5.1	Introduction	102
5.2	Materials and Methods	104
	5.2.1 Harvest Schedule	108
	5.2.2 Analysis of Data	109
	5.2.3 Climatic Variables	110
5.3	Results and Discussion	111
	5.3.1 Plant Height	114
	5.3.2 Canopy	116
	5.3.3 Length of Branch	117
	5.3.4 Number of Branch	119
	5.3.5 Number of Nodes	120
	5.3.6 Flower	123
	5.3.7 Seed	125
	5.3.8 Cumulative Seed at Different Time of the Year	129
	5.3.9 Total Leaf DM Yield per Year	131
	5.3.10 Total Stem DM Yield per Year	133
	5.3.11 Total Aerial DM Yield per Year	136
	5.3.12 Leaf to Stem Ratio	139
	5.3.13 Effect of Climate	142
5.4	Conclusion	146
6	<b>THE EFFECT OF SHADE AND CUTTING INTERVAL ON THE CHEMICAL COMPOSITIONS AND NUTRITIVE VALUE OF A. INTRUSA.</b>	150
6.1	Introduction	150
6.2	Materials and Method	154
6.3	Results and Discussion	156
	6.3.1 Leaf Crude Protein	156
	6.3.2 Leaf Essential Amino Acid	160
	6.3.3 Leaf Acid Detergent Fiber	162
	6.3.4 Leaf Neutral Detergent Fiber	164
	6.3.5 Leaf Hemicellulose	166
	6.3.6 Leaf Calcium	167
	6.3.7 Leaf Phosphorus	168

	<b>Page</b>
6.3.8 Leaf Magnesium	168
6.3.9 Leaf Gross Energy	170
6.3.10 Stem Crude Protein	171
6.3.11 Stem Essential Amino Acid	173
6.3.12 Stem Acid Detergent Fiber	177
6.3.13 Stem Neutral Detergent Fiber	178
6.3.14 Stem Hemicellulose	179
6.3.15 Stem Calcium	182
6.3.16 Stem Phosphorus	183
6.3.17 Stem Magnesium	183
6.3.18 Stem Gross Energy	185
6.3.19 Crude Protein of Aerial Part (Calculated)	186
6.3.20 Acid Detergent Fiber of Aerial Part (Calculated)	188
6.3.21 Neutral Detergent of Aerial Part (Calculated)	189
6.3.22 Hemicellulose of Aerial Part (Calculated)	191
6.3.23 Dry Matter Content of Aerial Part	192
6.3.24 In Vitro Dry Matter Digestibility	194
6.4 Conclusion	197
<b>7 PERSISTENCE OF A. INTRUSA AT DIFFERENT SHADE AND CUTTING INTERVAL</b>	<b>198</b>
7.1 Introduction	198
7.2 Materials And Method	200
7.3 Results And Discussion	200
7.4 Conclusion	206
<b>8 OPTIMUM LIGHT REQUIREMENTS OF A. INTRUSA</b>	<b>207</b>
8.1 Introduction	207
8.2 Materials and Method	211
8.3 Results and Discussion	213
8.4 Conclusion	217
<b>9 GENERAL DISCUSSION AND CONCLUSION</b>	<b>218</b>
<b>BIBLIOGRAPHY</b>	<b>227</b>
<b>APPENDICES</b>	<b>259</b>
<b>BIODATA OF THE STUDENT</b>	<b>304</b>

## LIST OF ABBREVIATIONS / NOTATIONS / GLOSSARY OF TERMS

AA	Amino acid
ACIAR	Australian Center for International Agricultural Research
AIA	Acid insoluble ash.
ADF	Acid detergent fiber.
AOAC	American Organization of Analytical Chemistry
Aug	August
BK	<i>Bahan kering</i>
BKT	<i>Bahan kering tercerna</i>
BW	Body weight
CIAT	Centro Internacional de Agricultura Tropical
CF	Crude fiber
CP	Crude protein.
CV	Coefficient of variation.
DBP	Dewan Bahasa dan Pustaka
DCP	Digestible crude protein
DDM	Digestible dry matter.
DE	Digestible energy
DM	Dry matter.
DMD	Dry matter digestibility
DMI	Dry matter intake
EAA	Essential Amino Acid
Feb	February
g	Gram





GE	Gross energy
IVDMD	In vitro dry matter digestibility
Jan	January
kg	Kilogram
kcal	Kilocalory
LSR	Leaf-to-stem ratio
mcg kg <sup>-1</sup>	Megacalorie per kilogram
MARDI	Malaysian Agricultural Research and Development Institute.
μmol	micromole
μM	micromole (used in graphs)
NAHQC	National Alfalfa Hay Quality Committee
NDF	Nitrogen detergent fiber.
NIR	Near Infra Red
OM	Organic matter
PAR	Photosynthetic active radiation
PD	Partially dried
PFD	Photon Flux Density
PK	<i>Protin kasar</i>
PKT	<i>Protin kasar tercerna</i>
PPFD	Photosynthetic photon flux density
RHI	Relative humidity (going inside)
RHO	Relative humidity (going outside)
RRIM	Rubber Research Institute of Malaysia
s	Second
s.d.	Standard deviation