



UNIVERSITI PUTRA MALAYSIA

***ANTIDEPRESSANT-LIKE EFFECTS OF MENHADEN FISH OIL IN
POSTPARTUM-INDUCED MODEL OF DEPRESSION IN RATS***

LEILA ARBABI

FPSK(M) 2014 15



**ANTIDEPRESSANT-LIKE EFFECTS OF MENHADEN FISH OIL IN
POSTPARTUM-INDUCED MODEL OF DEPRESSION IN RATS**

By

LEILA ARBABI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science**

March 2014

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

I dedicate this thesis to my beloved parents for their unconditional love and support



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Master of Science

**ANTIDEPRESSANT-LIKE EFFECTS OF MENHADEN FISH OIL
IN POSTPARTUM-INDUCED MODEL OF DEPRESSION IN RATS**

By

LEILA ARBABI

March 2014

Chair: Mohamad. Taufik Hidayat Baharuldin, PhD

Faculty: Faculty of Medicine and Health Sciences

Postpartum depression (PPD) is a psychiatric disorder that defined as a subtype of major depressive disorder (MDD) which may result from reproductive hormone fluctuations subsequent to childbirth. Approximately 10-15% of women experience postpartum depression (PPD) after baby delivery. It is hypothesized that the depression symptoms may be attenuated with omega-3 fatty acids. In order to examine this hypothesis, ovariectomized female rats underwent hormone-simulated pregnancy (HSP) regimen and received progesterone and estradiol benzoate or vehicle for 23 days, mimicking the actual rat's pregnancy. Forced feeding of menhaden fish oil (rich in omega-3) with three doses of 1, 3 and 9 g/kg/d, fluoxetine (an antidepressant drug) 15mg/kg/d and distilled water 2ml/d per rat started in six experimental groups on postpartum day 1 and continued for 15 consecutive days. On postpartum day 2, 8 and 15 all groups were tested in the forced swimming test (FST) and open field test (OFT). The seventh experimental group (normal control), which had not experienced neither ovariectomy nor injection, received only distilled water 2ml/d through oral gavage for 15 consecutive days and underwent the same behavioral tests as well. Animals were sacrificed by decapitation on postpartum day 15 following exposure to behavioral tests and blood and brain samples were collected. Plasma samples were utilized to assay levels of corticosterone and pro-inflammatory cytokines using ELISA and Procarta immunossay technique respectively. The corticosterone levels of hippocampus were measured by ELISA as well. Significant differences between groups were evaluated using one-way analysis of variance (ANOVA), followed by the post hoc Tukey's multiple comparison test when appropriate $P < 0.05$ was considered significant. The results showed that following 15 consecutive days' treatment with three different doses of menhaden fish oil, the immobility time of animals seen in FST was reduced compared to negative control group. The effect was found to be dose dependent where menhaden fish oil 3g and 9g/kg/d have shown significant reduction in immobility time. Furthermore, menhaden fish oil did not change locomotor activity; therefore, the decrease of immobility time observed in FST

following menhaden fish oil supplementation is due to its ability to attenuate depression. The results of biochemical analysis showed that the plasma levels of corticosterone, interleukin 1- β and interferon- γ were decreased significantly following menhaden fish oil treatment at doses of 3 and 9 g/kg/d. In addition, supplementation with 9 g/kg menhaden fish oil significantly decreased the plasma levels of tumor necrosis factor- α and hippocampal levels of corticosterone. However, menhaden fish oil at 1 g/kg produced a slight reduction ($p>0.05$) in immobility time in FST and in the levels of corticosterone and pro-inflammatory cytokines. Taken together, these results suggest that menhaden fish oil, rich in omega-3, exerts beneficial effect on postpartum depression and decreases the biomarkers related to depression.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Sarjana Sains

KESAN ANTI-KEMURUNGAN MINYAK IKAN MENHADEN KEATAS TIKUS MODEL KEMURUNGAN POSTPARTUM

Oleh

LEILA ARBABI

Mac 2014

Pengerusi: Mohamad. Taufik Hidayat Baharuldin, PhD
Fakulti: Perubatan dan Sains Kesihatan

Kemurungan postpartum (PPD) ialah gangguan psikiatri yang berlaku selepas kelahiran anak. PPD didefinisikan sebagai satu bentuk gangguan kemurungan (MDD) disebabkan oleh ketidakstabilan hormon reproduktif selepas kelahiran anak. Hampir 10-15% wanita mengalami PPD setelah kelahiran bayi. Adalah dihipotesiskan bahawa simptom-simptom kemurungan boleh dikurangkan dengan pengambilan asid omega 3 lemak. Untuk mengkaji hipotesis ini, tikus betina yang diaruh ovariektomi disimulasi kehamilan oleh hormon (HSP) melalui pemberian benzoat progesteron dan estradiol atau bahan kawalan selama 23 hari, bagi menyerupai tempoh kehamilan tikus sebenar. Pemberian secara oral minyak ikan menhaden (kaya dengan omega-3) dengan tiga dos 1, 3 dan 9 g/kg/d, 15mg/kg/d fluoxetine (dadah antidepresan) dan 2mL/d air suling setiap tikus dimulakan keatas 6 kumpulan eksperimen pada hari pertama pospartum dan berterusan untuk 15 hari berikutnya. Pada hari ke-2, 8 dan 15 pospartum, semua kumpulan diuji dengan ujian paksa-renang (FST) dan lapangan terbuka (OFT). Kumpulan uji kaji ketujuh, yang tidak diaruh ovariektomi mahupun suntikan hormon, menerima 2mL/d air suling secara oral untuk 15 hari berturut-turut serta menjalani ujian tingkahlaku yang sama. Haiwan dikorbankan dengan dipenggal lehernya pada hari ke-15 pospartum selepas ujian tingkahlaku serta sampel darah dan hipokampus diambil. Sampel plasma digunakan untuk ujian kortikosteron dan sitokin pro-inflamatori menggunakan ELISA dan teknik ujian imun Procarta. Tahap kortikosteron hipokampus diukur menggunakan ELISA. Perbezaan yang signifikan antara kumpulan dinilai menggunakan ANOVA, diikuti dengan ujian perbandingan pelbagai Tukey post hoc. Setelah 15 hari rawatan dengan tiga dos minyak ikan menhaden, masa immobiliti haiwan ketika FST lebih pendek berbanding kumpulan kawalan negatif. Dos 3g dan 9g/kg/d minyak ikan menhaden menunjukkan pengurangan yang signifikan terhadap masa immobiliti haiwan. Tambahan pula, pengambilan minyak ikan menhaden tidak mengubah aktiviti lokomotor; maka, pengurangan masa immobiliti haiwan ketika FST mencadangkan bahawa pengambilan minyak ikan menhaden berupaya mengurangkan kemurungan.

Keputusan analisis biokimia menunjukkan aras kortikosteron plasma, interleukin1- β dan interferon- γ menurun dengan signifikan pada kumpulan rawatan 3 dan 9 g/kg/d minyak ikan menhaden. Pengambilan 9 g/kg minyak ikan menhaden juga mengurangkan aras plasma nekrosis tumor faktor α dan kortikosteron hipokampus dengan signifikan. Walaubagaimanapun, minyak ikan menhaden 1g/kg menunjukkan sedikit penurunan ($p>0.05$) terhadap tempoh imobiliti FST dan paras kortikosteron serta sitokin. Oleh itu minyak ikan menhaden, yang kaya dengan omega-3, dicadangkan mempunyai kesan antidepresan serta berupaya mengurangkan paras penunjuk biologi keatas tikus-tikus yang diaruh PPD.



ACKNOWLEDGEMENTS

First of all, I would like to thank God for giving me this opportunity. I am honored and grateful to have the full support and commitment from a number of people during my study. I would like to express my sincere appreciation and gratitude to my supervisor Assoc. Prof. Dr. Mohamad. Taufik Hidayat Baharudin, my co-supervisor Assoc. Prof. Dr. Mohamad. Aris Mohamad. Moklas and member of the Supervisory Committee Assoc. Prof. Dr. Sharida Fakurazi for their kindness, support and guidance all throughout my study.

I am also indebted to the staffs of Anatomy laboratory, Department of Human Anatomy of FMHS for helping me to set up the behavioral room and assisted during the laboratory work. I would like to express special thanks to Dr. Sani Ismaila Muhammad for his technical support regarding surgery on rats and Mr. Ramli Suhaimi and staffs of Animal Unit for their kind support. I am also thankful to Mr. Ariyo Movahedi and Mr. Sobhan Ghafourian for their technical support.

Last but not least, I would like to express my deepest thanks to my beloved family and parents for their patience, continuous support and encouragement.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Mohamad. Taufik Hidayat Baharuldin, PhD

Associate. Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Mohamad. Aris Mohamad. Moklas, PhD

Associate. Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Sharida Fakurazi, PhD

Associate. Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, Ph.D

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 16 June 2014

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the University Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: _____

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of
Chairman of
Supervisory
Committee: _____

Signature: _____

Name of
Member of
Supervisory
Committee: _____

Signature: _____

Name of
Member of
Supervisory
Committee: _____

TABLE OF CONTENTS

	Page
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xv
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xviii
 CHAPTER	
1 INTRODUCTION	1
1.1 Introduction	1
1.2 Problem statement	3
1.3 Hypothesis	3
1.4 Objectives	3
1.4.1 General objectives	3
1.4.2 Specific objectives	3
 2 LITERATURE REVIEW	4
2.1 Postpartum depression	4
2.1.1 Pathophysiology of postpartum depression	5
2.1.2 Animal Model of Postpartum Depression (PPD)	5
2.1.3 Reproductive system in female rats	8
2.1.4 The effect of postpartum depression on family	11
2.1.5 Medical treatment options for PPD	11
2.2 Omega-3 fatty acids	12
2.2.1 The effects of omega-3 fatty acids on the body	13
2.2.2 Omega-3 and depression	14
2.2.4 Proposed mechanisms of omega-3 in depression	14
2.3 Hypothalamic-pituitary-adrenal (HPA) axis	15
2.3.1 HPA axis dysregulation in depression	17
2.3.2 Hippocampus and depression	18
2.4 Immune system and depression	18
2.4.1 Relation between pro-inflammatory cytokines and depression	19
2.4.2 Pathways linking the immune system and depression	19
2.5 Behavioral tests to study depression	20
2.5.1 Forced swim test (FST)	20
2.5.2 Open field test (OFT)	22
 3 MATERIALS AND METHODS	23
3.1 Animals	23
3.2 Postpartum model induction	25
3.2.1 Ovariectomy	25

3.2.2 Hormone Simulating Pregnancy (HSP)	25
3.3 Maternal behavior	26
3.4 Treatment regimen	26
3.5 Behavioral test	28
3.5.1 Open Field Test (OFT)	28
3.5.2 Forced Swim Test (FST)	30
3.6 Biochemical analysis	30
3.6.1 Blood and brain samples collection	32
3.6.2 Preparation of blood plasma sample	32
3.6.3 Preparing and homogenization of brain hippocampus	32
3.6.4 Hippocampus and plasma corticosterone analysis	32
3.6.5 Measurement of plasma level of pro-inflammatory cytokines including IL1 β , TNF- α , and INF- γ	32
3.7 Statistics	33
4 RESULTS AND DISCUSSION	34
4.1 Maternal behavior assessment	34
4.2 Body weight profile	35
4.3 The effects of menhaden fish oil on forced swim test (FST) according to postpartum days	36
4.3.1 The effects of menhaden fish oil on FST on postpartum day 2	36
4.3.2 The effects of menhaden fish oil on FST on Postpartum day 8	37
4.3.3 The effects of menhaden fish oil on FST on Postpartum day 15	38
4.3.4 Discussion	39
4.4 The effects of menhaden fish oil on open field test (OFT)	42
4.4.1 Results	42
4.4.2 Discussion	45
4.5 The effects of 15 days supplementation of menhaden fish oil on plasma and hippocampal levels of corticosterone	46
4.5.1 The effects of 15 days supplementation of menhaden fish oil on plasma level of corticosterone	46
4.5.2 The effects of 15 days forced feeding of menhaden fish oil on hippocampus corticosterone level	47
4.5.3 Discussion	48
4.6 The effects of 15 days supplementation of menhaden fish oil on pro-inflammatory cytokines	51
4.6.1 The effects of menhaden fish oil on interleukin 1 β (IL-1 β)	51
4.6.2 The effects of menhaden fish oil on Tumor necrosis factor-alpha (TNF- α)	52
4.6.3 The effects of menhaden fish oil on interferon-gamma (INF- γ)	53
4.6.4 Discussion	54
5 GENERAL DISCUSSION, CONCLUSION AND RECOMMENDATION FOR FUTURE STUDY	56
5.1 General discussion and conclusion	56
5.2 Recommendation for future study	57

REFERENCES	58
APPENDICES	75
BIODATA OF STUDENT	99
LIST OF PUBLICATIONS	100



LIST OF TABLES

Table	Page
2.1 Studies on postpartum-induced model of rat	7
3.1 Hormone-simulated pregnancy (HSP) regimen	26



LIST OF FIGURES

Figure	Page
2.1 A Schematic Picture of Reproductive System of Female Rat	8
2.2 Human and rat Reproductive Menstrual Cycle	9
2.3 Figure 2.3 Pattern of Estrogen and Progesterone in Rat's Normal Gestation	10
2.4 Plasma Levels of Estradiol and Progesterone in Postpartum Rats (Following a Normal Gestation) Compared to Virgin Control	10
2.5 The Hypothalamic-Pituitary-Adrenal (HPA) Axis and the Glucocorticoids (GC) Feedback	16
3.1 The Summary of Experimental Design	24
3.2 Timeline of Hormone and Treatment Regimen	27
3.3 Open Field Test for Locomotor Activity Evaluation	29
3.4 Forced Swim Test for Depressive-like Behavior Evaluation	31
4.1 Positive Nesting Behavior at the End of HSP Regimen	34
4.2 Percentages of Body Weight Gain of Experimental Rats from postpartum day 1 to postpartum day 15	35
4.3 Effects of Menhaden Fish Oil (1 g/kg, 3 g/kg and 9 g/kg) on Immobility Time in the Forced Swim Test on Postpartum Day 2	36
4.4 Effects of Menhaden Fish Oil (1 g/kg, 3 g/kg and 9 g/kg) on Immobility Time in the Forced Swim Test on Postpartum Day 8	37
4.5 Effects of Menhaden Fish Oil (1 g/kg, 3 g/kg and 9 g/kg) on Immobility Time in the Forced Swim Test on Postpartum Day 15	38
4.6 Effects of Menhaden Fish Oil (1 g/kg, 3 g/kg and 9 g/kg) and Fluoxetine (15 mg/kg) on Open Field Test (OFT) on Postpartum Day 2	42
4.7 Effects of Menhaden Fish Oil (1 g/kg, 3 g/kg and 9 g/kg) and Fluoxetine (15 mg/kg) on Open Field Test (OFT) on Postpartum Day 8	43

4.8	Effects of Menhaden Fish Oil (1 g/kg, 3 g/kg and 9 g/kg) and Fluoxetine (15 mg/kg) on Open Field Test (OFT) on Postpartum Day 15	44
4.9	Effects of Menhaden Fish Oil (1 g/kg/d, 3 g/kg/d and 9 g/kg/d) on Plasma Levels of Corticosterone in the Postpartum Depression-induced Rats	46
4.10	Effects of Menhaden Fish Oil (1 g/kg/d, 3 g/kg/d and 9 g/kg/d) on Hippocampus Level of Corticosterone in the Postpartum Depression-induced Rats	47
4.11	Effects of Menhaden Fish Oil (1 g/kg/d, 3 g/kg/d and 9 g/kg/d) on Plasma Levels of IL-1 β in the Postpartum Depression-induced Rats	51
4.12	Effects of Menhaden Fish Oil (1 g/kg/d, 3 g/kg/d and 9 g/kg/d) on Plasma Levels of TN- α in the Postpartum Depression-induced Rats	52
4.13	Effects of Different Doses of Menhaden Fish Oil on Plasma Level of INF-gamma in the Postpartum Depression-induced Rats	53

LIST OF ABBREVIATIONS

AA	Arachidonic Acid
ACTH	Adrenocorticotrophic Hormone
ACUC	Animal Care and Use Committee
ALA	Alpha-Linolenic Acid
ANOVA	Analysis Of Variance
C	Centigrade
CORT	Corticosterone
CRF	Corticotropin-Releasing Factor
d	day
DHA	Docosahexaenoic Acid
EB	Estradiol
EDTA	Ethylenediaminetetraacetic Acid
ELISA	Enzyme-Linked Immunosorbent Assay
EPA	Eicosapentaenoic Acid
FLX	Fluoxetine
FSH	Follicle-Stimulating Hormone
FST	Forced Swim Test
g	gravitational acceleration
GnRH	Gonadotropin-Releasing Hormone
GR	Glucocorticoid Receptor
HPA	Hypothalamic-Pituitary-Adrenal
HSP	Hormone-Simulated Pregnancy
i.m.	Intramuscular
IDO	Indoleamine 2,3-dioxygenase
IL	Interleukin
INF- γ	Interferon gamma
IRS	Inflammatory Response System
JHU	John Hopkins University
LA	Linoleic Acid
LCPUFA	Long Chain Polyunsaturated Fatty Acids
LH	Luteinizing Hormone
MAP	Multi-Analyte Profiling beads
MD	Major Depression
MDD	Major Depression Disorder
MFO	Menhaden Fish Oil
mL	milliliter
MR	Mineralocorticoid Receptor
MSc	Master of Science
N	Number
ng	nanogram
NK	Natural killer
OFT	Open Field Test

OVX	Ovariectomy
p	probability
PBS	Phosphate Buffered Saline
PG	Prostaglandin
Pg	picogram
PLA2	Phospholipases A2
PP	Postpartum
PPD	Postpartum Depression
PUFA	Polyunsaturated Fatty Acids
PVN	Paraventricular Nucleus
s.c.	subcutaneous
SEM	Standard Error of the Mean
SIDS	Sudden Infant Death Syndrome
SSRI	Selective Serotonin Reuptake Inhibitor
TCA	Tricyclic Antidepressants
TNF- α	Tumor Necrosis Factor alpha
UPM	Universiti Putra Malaysia
USA	United States of America
W	Watt

CHAPTER 1

INTRODUCTION

1.1 Introduction

Postpartum depression (PPD) is a psychiatric disorder, which occurs following childbirth. PPD is defined as a subtype of major depressive disorder (MDD) with several symptoms such as depressed mood, loss of interest in activities, loss of pleasure, sleep and appetite disturbance, guilty feeling, lack of concentration, and thought of suicide (Pearlstein *et al.*, 2009).

The onset of PPD occurs within the first 2-3 month after baby delivery and may last one year (Cuijpers *et al.*, 2008). Mood disorder is the most frequent morbidity for mothers during postpartum period (Stocky & Lynch, 2000).

Dramatically changes in reproductive hormones in all women at parturition results in “baby blues” with an estimated rate of 75 % of mothers (Moses-Kolko *et al.*, 2009). Symptoms of baby blues, which are similar to PPD, occur in the first week following baby delivery, peaked on the 5th day, and usually diminish by the 12th day. If the symptoms of baby blues last more than two weeks, it will be considered as postpartum depression (Leitch, 2002). It has been reported that 10-15% women suffer from PPD following childbirth (Halbreich, 2005).

Although the underlying etiology of postpartum depression remained unknown, scientists suggest a number of possible theories in this respect. Biological, psychological and social factors or combination of them may contribute to cause and progress of PPD (Berggren-Clive, 1998). One of the most potent hypotheses of its etiology is the hormonal hypothesis. Abrupt changes in reproductive hormones that women undergo in post-delivery period may cause postpartum depression (Moses-Kolko *et al.*, 2009; O'Hara, 2009; Parry *et al.*, 2003). In addition to the role of estrogen and progesterone, some other biological factors such as hypothalamic-pituitary-adrenal (HPA) axis hormones, altered immune system and cytokines, and altered fatty acids have been proposed to play a role in causing postpartum depression (Corwin & Pajer, 2008; Zonana & Gorman, 2005).

Not diagnosing and treating postpartum depression has significant adverse effects on depressed individuals and their families (Dennis, 2004). Increased risk for marital disruption and divorce as well as child abuse, infanticide and maternal suicide are some of the problems of untreated PPD (Sit *et al.*, 2006). Besides, children from depressed mothers have a high rate of cognitive and behavioral problems and have lower vocabulary skills (Jones & Venis, 2001).

Generally, two main classes of antidepressant drugs, prescribed to decrease PPD symptoms, are serotonin-specific reuptake inhibitor (SSRIs) and tricyclic antidepressants (TCAs). The best example for the former is fluoxetine (Prozac) and for the latter are amitriptyline (Elavil) and imipramine (Tofranil). Due to side effects of medical treatment on breastfed infants and negative effects of untreated depression, mothers face dilemma over how to deal with depression symptoms. Therefore, another alternative treatment should be considered to lessen depressive symptoms with lower side effects for both mother and baby.

The benefits of omega-3 to attenuate depression symptoms have been reported in previous studies. There are many reasons indicating an inverse relationship between omega-3 fatty acids and depression. This link is seen in both observational and experimental research (Hallahan & Garland, 2005; Frasere-Smith *et al.*, 2004; Tiemeier *et al.*, 2003; Maes *et al.*, 1999).

Omega-3 fatty acids are long chain polyunsaturated fatty acids (LCPUFA). Alpha-linolenic acid (ALA) is the precursor for omega-3 fatty acids and it is converted to eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) through elongation (Nettleton, 1995). EPA (20:5 n-3) and DHA (22:6 n-3) which are both vital compounds should be supplied via individuals' diet since the human body cannot convert ALA to EPA and DHA efficiently (Pawlosky *et al.*, 2001). Fish and fish oil, which are rich in omega-3 fatty acids, are the best dietary sources of EPA and DHA (McGregor *et al.*, 2001).

In the nervous system, omega-3 has vital functions such as controlling membrane function (Tinoco, 1982), acting as antioxidant, reducing cerebral lipid peroxides (Choi-Kwon *et al.*, 2004; Hossain *et al.*, 1999) and protecting cells from toxicants (Baker, 2007). Immune system, vision and motor skills are the other parts of the body which are affected by omega-3 (Baker, 2007).

Although it is believed that omega-3 plays a vital role in the body and particularly in the nervous system and mood disorder, the mechanism involved is poorly understood. While numerous studies have been done to evaluate the effects of omega-3 on depression and other mood disorders, researches carried out to determine the effects of omega-3 on postpartum depression are few and the results are inconsistent.

Due to the contradictions among the results of previous researches in this regard, the present study was performed to clarify these discrepancies by investigating the effects of menhaden fish oil (rich in omega-3) on postpartum-induced rats. In this study, the locomotor activity and antidepressant-like effect of omega-3 was evaluated using standard behavioral tests, OFT and FST respectively. In addition, corticosterone levels in plasma and hippocampus were assayed and to determine the relation between omega-3 and immune system responses, plasma levels of pro-inflammatory cytokines were also measured.

1.2 Problem statement

Due to disadvantages of using antidepressant drugs such as costs and side effects on mother and breastfed infant, another alternative treatment should be considered to lessen depressive symptoms with lower drawbacks.

1.3 Hypothesis

Menhaden fish oil (rich in omega-3 fatty acids) have antidepressant-like effects on postpartum model of depression in rats.

1.4 Objectives

1.4.1 General objectives

To investigate the beneficial effects of menhaden fish oil (rich in omega-3 fatty acids) on behavioral activity, corticosterone levels and immunological indicators in rat model of postpartum depression.

1.4.2 Specific objectives

1. To determine the effects of menhaden fish oil on behavioral activity in forced swimming test (FST) and locomotor activity in PPD-induced rats
2. To measure levels of corticosterone in the hippocampus and plasma of PPD-induced rats following menhaden fish oil treatment
3. To measure plasma levels of IL1- β , TNF- α and INF- γ in PPD-induced rats following menhaden fish oil treatment.

REFERENCES

- Abdel-Malek, Z. (2001). Melanocortin receptors: their functions and regulation by physiological agonists and antagonists. *Cellular and Molecular Life Sciences CMLS*, 58(3), 434-441 .
- Abel, E. L., & Bilitzke, P. (1990). A possible alarm substance in the forced swimming test. *Physiology & behavior*, 48(2), 233-239 .
- Abreu, A. C., & Stuart, S. (2005). Pharmacologic and Hormonal Treatments for Postpartum Depression-This article addresses management of postpartum depression using medications, including hormone therapy. Medication use during. *Psychiatric Annals*, 35(7), 568-576 .
- Ahmad, A., Moriguchi, T., & Salem Jr, N. (2002). Decrease in neuron size in docosahexaenoic acid-deficient brain. *Pediatric neurology*, 26(3), 210-218 .
- Akana, S., Chu, A., Soriano, L & ,Dallman, M. (2001). Corticosterone Exerts Site-Specific and State-Dependent Effects in Prefrontal Cortex and Amygdala on Regulation of Adrenocorticotrophic Hormone, Insulin and Fat Depots. *Journal of neuroendocrinology*, 13(7), 625-637 .
- Anisman, H., Ravindran, A., Griffiths, J., & Merali, Z. (1999). Endocrine and cytokine correlates of major depression and dysthymia with typical or atypical features. *Molecular psychiatry*, 4(2), 182-188 .
- Arab, L. (2003). Biomarkers of fat and fatty acid intake. *The Journal of nutrition*, 133(3), 925S-932S .
- Austin, M., Janosky, J., & Murphy, H. (2003). Increased corticotropin-releasing hormone immunoreactivity in monoamine-containing pontine nuclei of depressed suicide men. *Molecular psychiatry*, 8(3), 324-332 .
- Bagga, D., Wang, L., Farias-Eisner, R., Glaspy, J. A., & Reddy, S. T. (2003). Differential effects of prostaglandin derived from ω -6 and ω -3 polyunsaturated fatty acids on COX-2 expression and IL-6 secretion. *Proceedings of the National Academy of Sciences*, 100(4), 1 .1756-751
- Baker, J. (2007). Interactive Effects of Fish Oil and Methylmercury on the Fatty Acid Profile of Adult Rat Forebrain Phospholipids .
- Bardi, M., Shimizu, K., Barrett, G. M., Borgognini-Tarli, S. M., & Huffman, M. A. (2003). Peripartum cortisol levels and mother-infant interactions in Japanese macaques. *American journal of physical anthropology*, 120(3), 298-304 .
- Bate, L. A. (2013). REPRODUCTIVE ENDOCRINOLOGY. from <http://people.upei.ca/bate/html/pregnancyandparturition1.html#Top13>. Retrieved April 19, 2014
- Battle, C. L., Zlotnick, C., Pearlstein, T., Miller, I. W., Howard, M., Salisbury, A., & Stroud, L. (2008). Depression and breastfeeding: which postpartum patients take antidepressant medications? *Depression and anxiety*, 25(10), 888-891 .
- Beck, C. T. (1998). The effects of postpartum depression on child development: a meta-analysis. *Archives of psychiatric nursing*, 12(1), 12-20 .
- Becker, J. B., Arnold, A. P., Berkley, K. J., Blaustein, J. D., Eckel, L. A., Hampson, E., . . . Steiner, M. (2005). (Strategies and methods for research on sex differences in brain and behavior. *Endocrinology*, 146(4), 1650-1673 .

- Berggren-Clive, K. (1998). Out of the darkness and into the light: women's experiences with depression after childbirth. *Canadian Journal of Community Mental Health (Revue canadienne de santé mentale communautaire)*, 17(1), 103-120 .
- Bernardi, M., Vergoni, A., Sandrini, M., Tagliavini, S., & Bertolini, A. (1989). Influence of ovariectomy, estradiol and progesterone on the behavior of mice in an experimental model of depression. *Physiology & behavior*, 45(5), 1067-1068 .
- Bielajew, C., Konkle, A., Kentner, A., Baker, S., Stewart, A., Hutchins, A., . . . Fouriezos, G. (2003). Strain and gender specific effects in the forced swim test: effects of previous stress exposure. *Stress: The International Journal on the Biology of Stress*, 6(4), 269-280 .
- Binder, E., Künzel, H., Nickel, T., Kern, N., Pfennig, A., Majer, M., . . . Holsboer, F. (2009). HPA-axis regulation at in-patient admission is associated with antidepressant therapy outcome in male but not in female depressed patients. *Psychoneuroendocrinology*, 34(1), 99-109 .
- Bledsoe, S. E., & Grote, N. K. (2006). Treating depression during pregnancy and the postpartum: a preliminary meta-analysis. *Research on Social Work Practice*, 16(2), 109-120 .
- Bloch, M., Daly, R. C., & Rubinow, D. R. (2003). Endocrine factors in the etiology of postpartum depression. *Comprehensive Psychiatry*, 44(3), 234-246 .
- Bloch, R. G., Dooneief, A. S., Buchberg, A. S., & Spellman, S. (1954). The clinical effect of isoniazid and iproniazid in the treatment of pulmonary tuberculosis. *Annals of internal medicine*, 40(5), 881 .
- Boldrini, M., Underwood, M. D., Hen, R., Rosoklija, G. B., Dwork, A. J., Mann, J. J., & Arango, V. (2009). (Antidepressants increase neural progenitor cells in the human hippocampus. *Neuropsychopharmacology*, 34(11), 2376-2389 .
- Bonaccorso, S., Puzella, A., Marino, V., Pasquini, M., Biondi, M., Artini, M., . . . Bosmans, E. (2001). Immunotherapy with interferon-alpha in patients affected by chronic hepatitis C induces an intercorrelated stimulation of the cytokine network and an increase in depressive and anxiety symptoms. *Psychiatry research*, 105(1), 45-55 .
- Borsini, F., & Meli, A. (1988). Is the forced swimming test a suitable model for revealing antidepressant activity? *Psychopharmacology*, 94(2), 147-160 .
- Bourin, M., Fiocco, A., & Clenet, F. (2001). How valuable are animal models in defining antidepressant activity? *Human Psychopharmacology: Clinical and Experimental*, 16(1), 9-21 .
- Bourre, J.-M. (2007). Dietary omega-3 fatty acids for women. *Biomedicine & pharmacotherapy*, 61(2), 105-112 .
- Brennan, P. A., Hammen, C., Andersen, M. J., Bor, W., Najman, J. M., & Williams, G. M. (2000). Chronicity, severity, and timing of maternal depressive symptoms: relationships with child outcomes at age 5. *Developmental psychology*, 36(6), 759 .
- Bridges, N., Slais, K., & Syková, E. (2008). The effects of chronic corticosterone on hippocampal astrocyte numbers: a comparison of male and female Wistar rats. *Acta neurobiologiae experimentalis*, 68(2), 131 .
- Bridges, R. S. (1984). A quantitative analysis of the roles of dosage, sequence, and duration of estradiol and progesterone exposure in the regulation of maternal behavior in the rat. *Endocrinology*, 114(3), 930-940 .

- Brummelte, S., & Galea, L. A. (2010a). Chronic corticosterone during pregnancy and postpartum affects maternal care, cell proliferation and depressive-like behavior in the dam. *Hormones and behavior*, 58(5), 769-779.
- Brummelte, S., & Galea, L. A. (2010b). Chronic high corticosterone reduces neurogenesis in the dentate gyrus of adult male and female rats. *Neuroscience*, 168(3), 680-690.
- Brummelte, S., Pawluski, J. L., & Galea, L. A. (2006). High post-partum levels of corticosterone given to dams influence postnatal hippocampal cell proliferation and behavior of offspring: a model of post-partum stress and possible depression. *Hormones and behavior*, 50(3), 370-382.
- Brunton, P., Russell, J., & Douglas, A. (2008). Adaptive Responses of the Maternal Hypothalamic-Pituitary-Adrenal Axis during Pregnancy and Lactation. *Journal of neuroendocrinology*, 20(6), 764-776.
- Burr, G. O., & Burr, M. M. (1929). A new deficiency disease produced by the rigid exclusion of fat from the diet. *Journal of Biological Chemistry*, 82(2), 345-367.
- Butterweck, V., Christoffel, V., Nahrstedt, A., Peterleit, F., Spengler, B., & Winterhoff, H. (2003). Step by step removal of hyperforin and hypericin: activity profile of different *Hypericum* preparations in behavioral models. *Life sciences*, 73(5), 627-639.
- Calder, P. (2001). The scientific basis for fish oil supplementation in rheumatoid arthritis *Food and Nutritional Supplements* (pp. 175-197): Springer.
- Calder, P., & Grimble, R. (2002). Polyunsaturated fatty acids, inflammation and immunity. *European Journal of Clinical Nutrition*, 56, S14-19.
- Cao, D.-H., Xu, J.-F., Xue, R.-H., Zheng, W.-F., & Liu, Z.-L. (2004). Protective effect of chronic ethyl docosahexaenoate administration on brain injury in ischemic gerbils. *Pharmacology Biochemistry and Behavior*, 79(4), 651-659.
- Capuron, L., Gumnick, J. F., Musselman, D. L., Lawson, D. H., Reemsnyder, A., Nemeroff, C. B., & Miller, A. H. (2001). Neurobehavioral effects of interferon- α in cancer patients: phenomenology and paroxetine responsiveness of symptom dimensions.
- Capuron, L., Raison, C. L., Musselman, D. L., Lawson, D. H., Nemeroff, C. B., & Miller, A. H. (2003). Association of exaggerated HPA axis response to the initial injection of interferon-alpha with development of depression during interferon-alpha therapy. *American Journal of Psychiatry*, 160(7), 1342-1345.
- Capuron, L., Ravaut, A., Miller, A. H., & Dantzer, R. (2004). Baseline mood and psychosocial characteristics of patients developing depressive symptoms during interleukin-2 and/or interferon-alpha cancer therapy. *Brain, behavior, and immunity*, 18(3), 205-213.
- Carlezon Jr, W. A., Mague, S. D., Parow, A. M., Stoll, A. L., Cohen, B. M., & Renshaw, P. F. (2005). Antidepressant-like effects of uridine and omega-3 fatty acids are potentiated by combined treatment in rats. *Biological psychiatry*, 57(4), 343-350.
- Carrié, I., Clément, M., de Javel, D., Francès, H., & Bourre, J.-M. (2000). Phospholipid supplementation reverses behavioral and biochemical alterations induced by n-3 polyunsaturated fatty acid deficiency in mice. *Journal of Lipid Research*, 41(3), 473-480.
- Cerqueira, J. J., Taipa, R., Uylings, H. B., Almeida, O. F., & Sousa, N. (2007). Specific configuration of dendritic degeneration in pyramidal neurons of the medial

- prefrontal cortex induced by differing corticosteroid regimens. *Cerebral Cortex*, 17(9), 1998-2006 .
- Charan, J., & Kantharia, N. (2013). How to calculate sample size in animal studies? *Journal of pharmacology & pharmacotherapeutics*, 4(4), 303 .
- Charmandari, E., Tsigos, C., & Chrousos, G. (2005). Endocrinology of the stress response 1. *Annu. Rev. Physiol.*, 67, 259-284 .
- Checkley, S. (1996). The neuroendocrinology of depression and chronic stress. *British medical bulletin*, 52(3), 597-617 .
- Chen, H.-F., & Su, H.-M. (2012). Exposure to a maternal n-3 fatty acid-deficient diet during brain development provokes excessive hypothalamic–pituitary–adrenal axis responses to stress and behavioral indices of depression and anxiety in male rat offspring later in life. *The Journal of Nutritional Biochemistry* .
- Chitme, H., Muchandi, I., & Burli, S. (2009). Effect of Asparagus Racemosus Willd Root Extract on Ovariectomized Rats. *The Open Natural Products Journal*, 2, 16-23 .
- Choi-Kwon, S., Park, K., Lee, H.-J., Park, M.-S., Lee, J.-H., Jeon, S.-E., . . . Park, K.-C. (2004). Temporal changes in cerebral antioxidant enzyme activities after ischemia and reperfusion in a rat focal brain ischemia model: effect of dietary fish oil. *Developmental brain research*, 152(1), 11-18 .
- Choleris, E., Thomas, A., Kavaliers, M., & Prato, F. (2001). A detailed ethological analysis of the mouse open field test: effects of diazepam, chlordiazepoxide and an extremely low frequency pulsed magnetic field. *Neuroscience & Biobehavioral Reviews*, 25(3), 235-260 .
- Chung, M., Raman, G., Chew, P., Magula, N., Trikalinos, T., & Lau, J. (2007). Breastfeeding and maternal and infant health outcomes in developed countries. *Evid Technol Asses (Full Rep)*, 153, 1-186 .
- Colangelo, L. A., He, K., Whooley, M. A., Daviglus, M. L., & Liu, K. (2009). Higher dietary intake of long-chain ω -3 polyunsaturated fatty acids is inversely associated with depressive symptoms in women. *Nutrition*, 25(10), 1011-1019 .
- Corwin, E. J & ,Pajer, K. (2008). The psychoneuroimmunology of postpartum depression. *Journal of women's health*, 17(9), 1529-1534 .
- Cott, J., & Hibbeln, J. R. (2001). Lack of seasonal mood change in Icelanders. *American Journal of Psychiatry*, 158(2), 328-328 .
- Craft, R ,Kostick, M., Rogers, J., White, C., & Tsutsui, K. (2010). Forced swim test behavior in postpartum rats. *Pharmacology Biochemistry and Behavior*, 96(4), 402-412 .
- Cryan, J., & Mombereau, C. (2004). In search of a depressed mouse: utility of models for studying depression-related behavior in genetically modified mice. *Molecular psychiatry*, 9(4), 326-357 .
- Cryan, J., Valentino, R. J., & Lucki, I. (2005). Assessing substrates underlying the behavioral effects of antidepressants using the modified rat forced swimming test. *Neuroscience & Biobehavioral Reviews*, 29(4), 547-569 .
- Cuijpers, P., Brännmark, J. G., & van Straten, A. (2008). Psychological treatment of postpartum depression: a meta-analysis. *Journal of Clinical Psychology*, 64(1), 103-118 .
- D'Amato, F ,Rizzi, R., & Moles, A. (2001). A model of social stress in dominant mice: effects on sociosexual behaviour. *Physiology & behavior*, 73(3), 421-426 .

- Dalla, C., Antoniou, K., Drossopoulou, G., Xagoraris, M., Kokras, N., Sfikakis, A., & Papadopoulou-Daifoti, Z. (2005). Chronic mild stress impact: are females more vulnerable? *Neuroscience*, 135(3), 703-714 .
- Dalvi, A., & Lucki, I. (1999). Murine models of depression. *Psychopharmacology*, 147(1), 14-16 .
- De Chaves, G., Moretti, M., Castro, A., Dagostin, W., Da Silva, G., Boeck, C., . . . Gavioli, E. (2009). Effects of long-term ovariectomy on anxiety and behavioral despair in rats. *Physiology & behavior*, 97(3), 420-425 .
- De Kloet, E. R., Vreugdenhil, E., Oitzl, M. S., & Joëls, M. (1998). Brain corticosteroid receptor balance in health and disease. *Endocrine reviews*, 19(3), 269-301 .
- De Vriese, S., Christophe, A., & Maes, M. (2003). Lowered serum n-3 polyunsaturated fatty acid (PUFA) levels predict the occurrence of postpartum depression: further evidence that lowered n-PUFAs are related to major depression. *Life sciences*, 73(25), 3181-3187 .
- Delgado, P. L. (2000). Depression: the case for a monoamine deficiency. *Journal of Clinical Psychiatry* .
- Delion, S., Chalon, S., Hérault, J., Guilloteau, D., Besnard, J.-C & ,Durand, G. (1994). Chronic dietary alpha-linolenic acid deficiency alters dopaminergic and serotonergic neurotransmission in rats. *The Journal of nutrition*, 124(12), 2466-2476 .
- DeMar, J., Ma, K., Bell, J. M., Igarishi, M., Greenstein, D., & Rapoport, S. (2005). Dietary n-3 polyunsaturated fatty acid deprivation in rats following weaning increases their behavioral depression and aggression test scores. *J. Lipid Res*, 47, 172-180 .
- Dennis, C. (2004). Treatment of postpartum depression, part 2: a critical review of nonbiological interventions. *The Journal of clinical psychiatry*, 65(9), 1252-1265 .
- Dennis, C., & Kavanagh, J. (2002). Psychosocial interventions for preventing postpartum depression. *The Cochrane Library* .
- Dennis, C., & McQueen, K. (2007). (Does maternal postpartum depressive symptomatology influence infant feeding outcomes? *Acta paediatrica*, 96(4), 590-594 .
- Detke, M. J., & Lucki, I. (1995). Detection of serotonergic and noradrenergic antidepressants in the rat forced swimming test: the effects of water depth. *Behavioural brain research*, 73(1), 43-46 .
- Detke, M. J., Rickels, M., & Lucki, I. (1995). Active behaviors in the rat forced swimming test differentially produced by serotonergic and noradrenergic antidepressants. *Psychopharmacology*, 72-66, (1)121 ,
- Donahue Jennings, K., Ross, S., Popper, S., & Elmore, M. (1999). Thoughts of harming infants in depressed and nondepressed mothers. *Journal of affective disorders*, 54(1), 21-28 .
- Droste, S., De Groote, L., Lightman, S., Reul, J., & Linthorst, A. (2009). The ultradian and circadian rhythms of free corticosterone in the brain are not affected by gender: an in vivo microdialysis study in Wistar rats. *Journal of neuroendocrinology*, 21(2), 132-140 .
- Duman, R. (2002). Pathophysiology of depression: the concept of synaptic plasticity. *European psychiatry*, 17, 306-310 .

- Dunn, A. J. (2000). Cytokine activation of the HPA axis. *Annals of the New York Academy of Sciences*, 917(1), 608-617 .
- Dunn, A. J., Swiergiel, A. H., & Beaupaire, R. d. (2005). Cytokines as mediators of depression: what can we learn from animal studies? *Neuroscience & Biobehavioral Reviews*, 29(4), 891-909 .
- Emanuele, M. A., Wezeman, F., & Emanuele, N. V. (2002). Alcohol's effects on female reproductive function. *Alcohol Research and Health*, 26(4), 274-281 .
- Endres, S., Ghorbani, R., Kelley, V. E., Georgilis, K., Lonnemann, G., van der Meer, J. W., . . . Weber, P. C. (1989). The effect of dietary supplementation with n—3 polyunsaturated fatty acids on the synthesis of interleukin 1-and tumor necrosis factor by mononuclear cells. *New England Journal of Medicine*, 320(5), 265-271 .
- Engblom, D., Ek, M., Saha, S., Ericsson-Dahlstrand, A., Jakobsson, P.-J., & Blomqvist, A. (2002). Prostaglandins as inflammatory messengers across the blood-brain barrier. *Journal of molecular medicine*, 80(1), 5-15 .
- Epperson, N., Czarkowski, K. A., Ward-Oâ, D., Weiss, E., Gueorguieva, R., Jatlow, P., & Anderson, G. M. (2001). Maternal sertraline treatment and serotonin transport in breast-feeding mother-infant pairs. *American Journal of Psychiatry*, 158(10), 1631-1637 .
- Estrada-Camarena, E., Contreras, C. M., Saavedra, M., Luna-Baltazar, I., & López-Rubalcava, C. (2002). Participation of the lateral septal nuclei (LSN) in the antidepressant-like actions of progesterone in the forced swimming test (FST). *Behavioural brain research*, 134(1), 175-183 .
- Estrada-Camarena, E., Fernandez-Guasti, A., & Lopez-Rubalcava, C. (2003). Antidepressant-like effect of different estrogenic compounds in the forced swimming test. *Neuropsychopharmacology: official publication of the American College of Neuropsychopharmacology*, 28(5), 830-838 .
- Evans, J., Sun, Y., McGregor, A., & Connor, B. (2012). Allopregnanolone regulates neurogenesis and depressive/anxiety-like behaviour in a social isolation rodent model of chronic stress. *Neuropharmacology* .
- Fedorova, I., & Salem Jr, N. (2006). Omega-3 fatty acids and rodent behavior. *Prostaglandins, leukotrienes and essential fatty acids*, 75(4), 271-289 .
- Ferraz, A. C., Delattre, A. M., Almendra, R. G., Sonagli, M., Borges, C., Araujo, P., . . . Lima, M. (2011). Chronic ω-3 fatty acids supplementation promotes beneficial effects on anxiety, cognitive and depressive-like behaviors in rats subjected to a restraint stress protocol. *Behavioural brain research*, 219(1), 116-122 .
- Festing, M. F. (2006). Design and statistical methods in studies using animal models of development. *ILAR journal*, 47(1), 5-14 .
- Festing, M. F., & Altman, D. G. (2002). Guidelines for the design and statistical analysis of experiments using laboratory animals. *ILAR journal*, 43(4), 244-258 .
- Frasure-Smith, N., Lespérance, F., & Julien, P. (2004). Major depression is associated with lower omega-3 fatty acid levels in patients with recent acute coronary syndromes. *Biological psychiatry*, 55(9), 891-896 .
- Frye, C. A., & Wawrzycki, J. (2003). Effect of prenatal stress and gonadal hormone condition on depressive behaviors of female and male rats. *Hormones and behavior*, 44(4), 319-326 .

- Galani, R., Duconseille, E., Bildstein, O., & Cassel, J.-C. (2001). Effects of room and cage familiarity on locomotor activity measures in rats. *Physiology & behavior*, 74(1), 1-4 .
- Galea, L. A., Wide, J. K., & Barr, A. M. (2001). Estradiol alleviates depressive-like symptoms in a novel animal model of post-partum depression. *Behavioural brain research*, 122(1), 1-9 .
- Garrido, P., de Blas, M., Del Arco, A., Segovia, G., & Mora, F. (2012). Aging increases basal but not stress-induced levels of corticosterone in the brain of the awake rat. *Neurobiology of aging*, 33(2), 375-382 .
- Garza, J. C., Guo, M., Zhang, W., & Lu, X.-Y. (2011). Leptin restores adult hippocampal neurogenesis in a chronic unpredictable stress model of depression and reverses glucocorticoid-induced inhibition of GSK-3 β / β -catenin signaling. *Molecular psychiatry*, 17(8), 790-808 .
- Gil, A. (2002). Polyunsaturated fatty acids and inflammatory diseases. *Biomedicine & pharmacotherapy*, 56(8), 388-396 .
- Gould, E. (2007). How widespread is adult neurogenesis in mammals? *Nature Reviews Neuroscience*, 8(6), 481-488 .
- Gould, T. D., Dao, D. T., & Kovacsics, C. E. (2009). The open field test *Mood and Anxiety Related Phenotypes in Mice* (pp. 1-20): Springer.
- Grace, S. L., Evindar, A., & Stewart, D. (2003). The effect of postpartum depression on child cognitive development and behavior: a review and critical analysis of the literature. *Archives of Women's Mental Health*, 6(4), 263-274 .
- Green, A. D., Barr, A. M., & Galea, L. A. (2009). Role of estradiol withdrawal in 'anhedonic' sucrose consumption: a model of postpartum depression. *Physiology & behavior*, 97(2), 259-265 .
- Green, A. D., & Galea, L. A. (2008). Adult hippocampal cell proliferation is suppressed with estrogen withdrawal after a hormone-simulated pregnancy. *Hormones and behavior*, 54(1), 20 .211-3
- Green, P., Gispan-Herman, I., & Yadid, G. (2005). Increased arachidonic acid concentration in the brain of Flinders Sensitive Line rats, an animal model of depression. *Journal of Lipid Research*, 46(6), 1093-1096 .
- Gregus, A., Wintink, A. J., Davis, A. C., & Kalynchuk, L. E. (2005). Effect of repeated corticosterone injections and restraint stress on anxiety and depression-like behavior in male rats. *Behavioural brain research*, 156(1), 105-114 .
- Halbreich, U. (2005). Postpartum disorders: multiple interacting underlying mechanisms and risk factors. *Journal of affective disorders*, 88(1), 1-7 .
- Hall, C. S. (1934). Emotional behavior in the rat. I. Defecation and urination as measures of individual differences in emotionality. *Journal of Comparative Psychology*, 18(3), 385 .
- Hallahan, B., & Garland, M. R. (2005). Essential fatty acids and mental health. *The British Journal of Psychiatry*, 186(4), 275-277 .
- Hallberg, P., & Sjöblom, V. (2005). The use of selective serotonin reuptake inhibitors during pregnancy and breast-feeding: a review and clinical aspects. *Journal of clinical psychopharmacology*, 25(1), 59-73 .
- Hamilton, J. A., & Harberger, P. N. (1992). *Postpartum psychiatric illness: a picture puzzle*: University of Pennsylvania Press.

- Handa, R. J & ,Weiser, M. J. (2013). Gonadal steroid hormones and the hypothalamo–pituitary–adrenal axis. *Frontiers in neuroendocrinology* .
- Hansen, H., Sanchez, C., & Meier, E. (1997). Neonatal administration of selective serotonin reuptake inhibitors Lu 10-17-C increased forced swimming immobility in adult rats: a putative model of depression. *J Pharmacol Exp Ther*, 3, 1333-1341 .
- Hasliza, A., Noordin, M., & Goh, Y. (2011). Effects of exercise and dietary polyunsaturated fatty acid on blood lipid profiles of streptozotocin-induced diabetes in rats. *Pertanika Journal of Tropical Agricultural Science*, 34(1), 151-155 .
- Heinzmann, J.-M. (2012). *Central and peripheral aspects of hypothalamic-pituitary-adrenal (HPA) axis dysfunction: insights from mice selectively bred for extremes in stress reactivity*. München, Ludwig-Maximilians-Universität, Diss., 2012 .
- Hennebelle, M., Balasse, L., Latour, A., Champeil-Potokar, G., Denis, S., Lavielle, M., . . . Vancassel, S. (2012). Influence of omega-3 fatty acid status on the way rats adapt to chronic restraint stress. *PloS one*, 7(7), e42142 .
- Herbert, T. B., & Cohen, S. (1993). Depression and immunity: a meta-analytic review. *Psychological bulletin*, 113(3), 472 .
- Herman, J. P., & Cullinan, W. E. (1997). Neurocircuitry of stress :central control of the hypothalamo–pituitary–adrenocortical axis. *Trends in neurosciences*, 20(2), 78-84 .
- Herman, J. P., Figueiredo, H., Mueller, N. K., Ulrich-Lai, Y., Ostrander, M. M., Choi, D. C., & Cullinan, W. E. (2003). Central mechanisms of stress integration: hierarchical circuitry controlling hypothalamo–pituitary–adrenocortical responsiveness. *Frontiers in neuroendocrinology*, 24(3), 151-180 .
- Hibbeln, J. R. (1998). Fish consumption and major depression. *Lancet*, 351(9110), 1213 .
- Hibbeln, J. R .(2002) .Seafood consumption, the DHA content of mothers' milk and prevalence rates of postpartum depression: a cross-national, ecological analysis. *Journal of affective disorders*, 69(1), 15-29 .
- Hibbeln, J. R., Linnoila, M., Umhau, J. C., Rawlings, R., George, D. T., & Salem Jr, N. (1998a). Essential fatty acids predict metabolites of serotonin and dopamine in cerebrospinal fluid among healthy control subjects, and early-and late-onset alcoholics. *Biological psychiatry*, 44(4), 235-242 .
- Hibbeln, J. R., Umhau, J. C., Linnoila, M., George, D. T., Ragan, P. W., Shoaf, S. E., . . . Salem Jr, N. (1998b). A replication study of violent and nonviolent subjects: cerebrospinal fluid metabolites of serotonin and dopamine are predicted by plasma essential fatty acids .*Biological psychiatry*, 44(4), 243-249 .
- Hill, M. N., Brotto, L. A., Lee, T. T.-Y., & Gorzalka, B. B. (2003). Corticosterone attenuates the antidepressant-like effects elicited by melatonin in the forced swim test in both male and female rats. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 27(6), 905-911 .
- Hinojosa, F. R., Spricigo Jr, L., Izidio, G. S., Brüske, G. R., Lopes, D. M., & Ramos, A. (2006). Evaluation of two genetic animal models in behavioral tests of anxiety and depression .*Behavioural brain research*, 168(1), 127-136 .
- Holmes, P. V. (2003). Rodent models of depression: reexamining validity without anthropomorphic inference. *Critical Reviews™ in Neurobiology*, 15(2) .(

- Holsboer, F. (2001). Stress, hypercortisolism and corticosteroid receptors in depression: implications for therapy. *Journal of affective disorders*, 62(1), 77-91 .
- Hossain, M., Hashimoto, M., Gamoh, S., & Masumura, S. (1999). Antioxidative effects of docosahexaenoic acid in the cerebrum versus cerebellum and brainstem of aged hypercholesterolemic rats. *Journal of neurochemistry*, 72(3), 1133-1138 .
- Huang, S.-Y., Yang, H.-T., Chiu, C.-C., Pariante, C. M., & Su, K.-P. (2008). Omega-3 fatty acids on the forced-swimming test. *Journal of psychiatric research*, 42(1), 58 .63-
- Hvid, H., Thorup, I., Sjögren, I., Oleksiewicz, M. B., & Jensen, H. E. (2012). Mammary gland proliferation in female rats: Effects of the estrous cycle, pseudo-pregnancy and age. *Experimental and Toxicologic Pathology*, 64(4), 321-332 .
- Hyman, S. E .(2009) .How adversity gets under the skin. *Nature neuroscience*, 12(3), 241-243 .
- Irwin, M. R., & Miller, A. H. (2007). Depressive disorders and immunity: 20 years of progress and discovery. *Brain, behavior, and immunity*, 21(4), 374-383 .
- Janeway, C., Travers, P., & Hunt, S. (1996). *Immunobiology: the immune system in health and disease* (Vol. 7): Current Biology.
- Jayatissa, M. N., Bisgaard, C. F., West, M. J., & Wiborg, O. (2008). The number of granule cells in rat hippocampus is reduced after chronic mild stress and re-established after chronic escitalopram treatment. *Neuropharmacology*, 54(3), 530-541 .
- JHU. (2009). Rodent Drug Formulary: Quick Reference Guide. from <http://web.jhu.edu/animalcare/rdf/index.html>. Retrieved December 20, 2013
- Johnson, S. A .,Fournier, N. M., & Kalynchuk, L. E. (2006). Effect of different doses of corticosterone on depression-like behavior and HPA axis responses to a novel stressor. *Behavioural brain research*, 168(2), 280-288 .
- Jolley, S. N., Elmore, S., Barnard, K. E., & Carr ,D. B. (2007). Dysregulation of the hypothalamic-pituitary-adrenal axis in postpartum depression. *Biological Research for Nursing*, 8(3), 210-222 .
- Jones, H. W., & Venis, J. A. (2001). Identification and classification of postpartum psychiatric disorders .*Journal of psychosocial nursing and mental health services*, 39(12), 23-30 .
- Kalynchuk, L. E., Gregus, A., Boudreau, D., & Perrot-Sinal, T. S. (2004). Corticosterone increases depression-like behavior, with some effects on predator odor-induced defensive behavior, in male and female rats. *Behavioral neuroscience*, 118(6), 1365 .
- Kent, S., Bluthé, R.-M., Kelley, K. W., & Dantzer, R. (1992). Sickness behavior as a new target for drug development. *Trends in Pharmacological Sciences*, 13, 24-28 .
- Kessing, L. V .,Agerbo, E., & Mortensen, P. (2003). Does the impact of major stressful life events on the risk of developing depression change throughout life? *Psychological medicine*, 33(07), 1177-1184 .
- Kiecolt-Glaser, J. K., Belury, M. A., Porter, K., Beversdorf, D .Q., Lemeshow, S., & Glaser, R. (2007). Depressive symptoms, omega-6: omega-3 fatty acids, and inflammation in older adults. *Psychosomatic medicine*, 69(3), 217-224 .

- Kodas, E., Galineau, L., Bodard, S., Vancassel, S., Guilloteau, D., Besnard, J. C., & Chalon, S. (2004). Serotonergic neurotransmission is affected by n-3 polyunsaturated fatty acids in the rat. *Journal of neurochemistry*, 89(3), 695-702 .
- Kronfol, Z. (2002). Immune dysregulation in major depression: a critical review of existing evidence. *The International Journal of Neuropsychopharmacology*, 5(04), 333-343 .
- Kronfol, Z., & Remick, D. G. (2000). Cytokines and the brain: implications for clinical psychiatry. *American Journal of Psychiatry*, 157(5), 683-694 .
- Kurata, A., Morinobu, S., Fuchikami, M., Yamamoto, S., & Yamawaki, S. (2009). Maternal postpartum learned helplessness (LH) affects maternal care by dams and responses to the LH test in adolescent offspring. *Hormones and behavior*, 56(1), 112-120 .
- Lacroix, S., Vallières, L., & Rivest, S. (1996). C-fos mRNA pattern and corticotropin-releasing factor neuronal activity throughout the brain of rats injected centrally with a prostaglandin of E2 type. *Journal of Neuroimmunology*, 70(2), 163-179 .
- Lands, W. E., Libelt, B., Morris, A., Kramer, N. C., Prewitt, T. E., Bowen, P., . . . Burns, J. H. (1992). Maintenance of lower proportions of (< i> n</i>- 6) eicosanoid precursors in phospholipids of human plasma in response to added dietary (< i> n</i>- 3) fatty acids. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, 1180(2), 147-162 .
- Leitch, S. (2002). Postpartum depression: A review of the literature. *St. Thomas, Ontario: Elgin-St. Thomas Health Unit* .
- Lightman, S. L., Windle, R. J., Wood, S. A., Kershaw, Y. M., Shanks, N & ,Ingram, C. D. (2001). Peripartum plasticity within the hypothalamo-pituitary-adrenal axis. *Progress in brain research*, 133, 111-129 .
- Lindgren, K. (2001). Relationships among maternal-fetal attachment, prenatal depression, and health practices in pregnancy. *Research in Nursing & Health*, 24(3), 203-217 .
- Little, H., Croft, A., O'callaghan, M., Brooks, S., Wang, G., & Shaw, S. (2008). Selective increases in regional brain glucocorticoid: a novel effect of chronic alcohol. *Neuroscience*, 156(4), 1017-1027 .
- Logan, A. C. (2003). Neurobehavioral aspects of omega-3 fatty acids: possible mechanisms and therapeutic value in major depression. *Alternative medicine review*, 8(4), 410-425 .
- Loomer, H. P., Saunders, J. C., & Kline, N. S. (1957). A clinical and pharmacodynamic evaluation of iproniazid as a psychic energizer. *Psychiatric research reports* .
- Lotrich, F. E., Sears, B., & McNamara, R. K. (2013). Elevated ratio of arachidonic acid to long-chain omega-3 fatty acids predicts depression development following interferon-alpha treatment: relationship with interleukin-6. *Brain, behavior, and immunity*, 31, 48-53 .
- Lucki, I. (1997). The forced swimming test as a model for core and component behavioral effects of antidepressant drugs. *Behavioural pharmacology*, 8(6-7),(532-523
- Maes, M. (1995). Evidence for an immune response in major depression: a review and hypothesis. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 19(1), 11-38 .

- Maes, M. (1999). Major depression and activation of the inflammatory response system *Cytokines, stress, and depression* (pp. 25-46): Springer.
- Maes, M. (2011). Depression is an inflammatory disease, but cell-mediated immune activation is the key component of depression. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 35(3), 664-675 .
- Maes, M., Bosmans, E., Suy, E., Vandervorst, C., De Jonckheere, C., & Raus, J. (1990). Immune disturbances during major depression: upregulated expression of interleukin-2 receptors. *Neuropsychobiology*, 24(3), 115-120 .
- Maes, M., Christophe, A., Delanghe, J., Altamura, C., Neels, H., & Meltzer, H. Y. (1999). Lowered ω 3 polyunsaturated fatty acids in serum phospholipids and cholesteryl esters of depressed patients. *Psychiatry research*, 85(3), 275-291 .
- Maes, M., & Smith, R. S. (1998). Fatty acids, cytokines, and major depression. *Biological psychiatry*, 43(5), 313 .
- Magiakou, M., Mastorakos, G., Rabin, D., Dubbert, B., Gold, P., & Chrousos, G. (1996). Hypothalamic corticotropin-releasing hormone suppression during the postpartum period: implications for the increase in psychiatric manifestations at this time. *Journal of Clinical Endocrinology & Metabolism*, 81(5), 1912-1917 .
- Makrides, M., Neumann, M. A., Byard, R. W., Simmer, K., & Gibson, R. A. (1994). Fatty acid composition of brain, retina, and erythrocytes in breast-and formula-fed infants. *The American journal of clinical nutrition*, 60(2), 189-194 .
- Mandl, A. M. (1951). The phases of the oestrous cycle in the adult white rat. *Journal of Experimental Biology*, 28(4), 576-584 .
- Martínez-Mota, L. a., Contreras, C. M., & Saavedra, M. (1999). Progesterone reduces immobility in rats forced to swim. *Archives of medical research*, 30(4), 286-289 .
- Mayer, A. D., Monroy, M. A., & Rosenblatt, J. S. (1990). Prolonged estrogen-progesterone treatment of nonpregnant ovariectomized rats: factors stimulating home-cage and maternal aggression and short-latency maternal behavior. *Hormones and behavior*, 24(3), 342-364 .
- McEwen, B. S. (2007). Physiology and neurobiology of stress and adaptation: central role of the brain. *Physiological reviews*, 87(3), 873-904 .
- McGregor, J. A., Allen, K. G., Harris, M. A., Reece, M., Wheeler, M., French, J. I., & Morrison, J. (2001). The Omega-3 Story:: Nutritional Prevention of Preterm Birth and Other Adverse Pregnancy Outcomes. *Obstetrical & gynecological survey*, 56(5), S1-S13 .
- McIsaac, S. A., & Young, A. H. (2009). The role of hypothalamic pituitary-adrenal axis dysfunction in the etiology of depressive disorders. *Drugs of today (Barcelona, Spain: 1998)*, 45(2), 127 .
- McKinnon, M. C., Yucel, K., Nazarov, A., & MacQueen, G. M. (2009). A meta-analysis examining clinical predictors of hippocampal volume in patients with major depressive disorder. *Journal of psychiatry & neuroscience: JPN*, 34(1), 41 .
- Melmed, S., Polonsky, K. S., Larsen, P. R., & Kronenberg, H. M. (2011). *Williams Textbook of Endocrinology: Expert Consult*: Elsevier Health Sciences.
- Merali, Z., Du, L., Hrdina, P., Palkovits, M., Faludi, G., Poulter, M. O., & Anisman, H. (2004). Dysregulation in the suicide brain: mRNA expression of corticotropin-releasing hormone receptors and GABAA receptor subunits in frontal cortical brain region. *The Journal of neuroscience*, 24(6), 1478-1485 .

- Merali, Z., Kent, P., Du, L., Hrdina, P., Palkovits, M., Faludi, G . . . ,Anisman, H. (2006). Corticotropin-releasing hormone, arginine vasopressin, gastrin-releasing peptide, and neuromedin B alterations in stress-relevant brain regions of suicides and control subjects. *Biological psychiatry*, 59(7), 594-602 .
- Mezadri, T .,Batista, G., Portes, A., Marino-Neto, J., & Lino-de-Oliveira, C. (2011). Repeated rat-forced swim test: reducing the number of animals to evaluate gradual effects of antidepressants. *Journal of neuroscience methods*, 195(2), 200-205 .
- Miguelé, M., Anisman ,H., Weber, J.-M., & Merali, Z. (2006). Effects of acute or chronic omega-3 and omega-6 polyunsaturated fatty acid treatment on behavioral, neuroendocrine and cytokine changes elicited by exogenous interleukin-1 β challenge. *Journal of Neuroimmunology*, 18 .28-19 ,(1)1
- Mirescu, C., & Gould, E. (2006). Stress and adult neurogenesis. *Hippocampus*, 16(3), 233-238 .
- Mitra, R., & Sapolsky, R. M. (2008). Acute corticosterone treatment is sufficient to induce anxiety and amygdaloid dendritic hypertrophy. *Proceedings of the National Academy of Sciences*, 105(14), 5573-5578 .
- Molina-Hernández, M., & Téllez-Alcántara, N. P. (2001). Antidepressant-like actions of pregnancy, and progesterone in Wistar rats forced to swim. *Psychoneuroendocrinology*, 26(5), 479-491 .
- Moses-Kolko, E. L., Berga, S. L., Kalro, B., Sit, D. K., & Wisner, K. L. (2009). Transdermal estradiol for postpartum depression: a promising treatment option. *Clinical obstetrics and gynecology*, 52(3), 516 .
- Musselman, D. L., Miller, A. H., Porter, M. R., Manatunga, A., Gao, F., Penna, S., . . . McDaniel, J. S. (2001). Higher than normal plasma interleukin-6 concentrations in cancer patients with depression: preliminary findings. *American Journal of Psychiatry*, 158(8), 1252-1257 .
- Naliwaiko, K., Araújo, R., Da Fonseca, R., Castilho, J., Andreatini, R., Bellissimo, M., . . . Fernandes, L. (2004). Effects of fish oil on the central nervous system: a new potential antidepressant? *Nutritional neuroscience*, 7(2), 91-99 .
- Navarre, B. M. (2008). Sucrose preference testing as a measure of anhedonia in "postpartum" rats. *Washington State University Honors College thesis, Fall* .
- Navarre, B. M., Laggart, J. D., & Craft, R. M. (2010). Anhedonia in postpartum rats. *Physiology & behavior*, 99(1), 59-66 .
- Nettleton, J. A. (1 .(995*Omega-3 fatty acids and health*: Springer.
- Newport, D. J., Hostetter, A., Arnold, A., & Stowe, Z. N. (2002). The treatment of postpartum depression: minimizing infant exposures. *The Journal of clinical psychiatry*, 63, 31 .
- Nielsen, A., Jørgensen, L .,Nielsen, J. N., Eivindson, M., Grønbæk, H., Vind, I., . . . Munkholm, P. (2005). Omega-3 fatty acids inhibit an increase of proinflammatory cytokines in patients with active Crohn's disease compared with omega-6 fatty acids. *Alimentary pharmacology & therapeutics*, 22(11-12), 1121-1128 .
- Nunes, S., Reiche, E., Morimoto, H., Matsuo, T., Itano, E., Xavier, E., . . . Silva, S. (2002). Immune and hormonal activity in adults suffering from depression. *Brazilian Journal of Medical and Biological Research*, 35(5 .587-581 ,(

- O'Hara, M. W. (2009). Postpartum depression: what we know. *Journal of Clinical Psychology*, 65(12), 1258-1269 .
- Okada, M., Hayashi, N., Kometani, M., Nakao, K., & Inukai, T. (1997). Influences of ovariectomy and continuous replacement of 17beta-estradiol on the tail skin temperature and behavior in the forced swimming test in rats. *Japanese journal of pharmacology*, 73(1), 93-96 .
- Otto, S., De Groot, R., & Hornstra, G. (2003). Increased risk of postpartum depressive symptoms is associated with slower normalization after pregnancy of the functional docosahexaenoic acid status. *Prostaglandins, leukotrienes and essential fatty acids*, 69(4), 237-243 .
- Pace, T. W., Hu, F., & Miller, A. H. (2007). Cytokine-effects on glucocorticoid receptor function :relevance to glucocorticoid resistance and the pathophysiology and treatment of major depression. *Brain, behavior, and immunity*, 21(1), 9-19 .
- Pariante, C. M., & Miller, A. H. (2001). Glucocorticoid receptors in major depression: relevance to pathophysiology and treatment. *Biological psychiatry*, 49(5), 391-404 .
- Parry, B. L., Sorenson, D. L., Meliska, C. J., Basavaraj, N., Zirpoli, G. G., Gamst, A., & Hauger, R. (2003). Hormonal basis of mood and postpartum disorders. *Current women's health reports*, 3(3) .235-230 ,(
- Pawlosky, R. J., Hibbeln, J. R., Novotny, J. A., & Salem, N. (2001). Physiological compartmental analysis of α -linolenic acid metabolism in adult humans. *Journal of Lipid Research*, 42(8), 1257-1265 .
- Pawluski, J. L., Charlier, T. D., Lieblich ,S. E., Hammond, G. L., & Galea, L. A. (2009). Reproductive experience alters corticosterone and CBG levels in the rat dam. *Physiology & behavior*, 96(1), 108-114 .
- Pearlstein, T., Howard, M., Salisbury, A., & Zlotnick, C. (2009). Postpartum depression. *American journal of obstetrics and gynecology*, 200(4), 357-364 .
- Pearlstein, T., Zlotnick, C., Battle, C., Stuart, S., O'Hara, M., Price, A., . . . Howard, M. (2006). Patient choice of treatment for postpartum depression: a pilot study. *Archives of women's mental health*, 9(6), 303-308 .
- Peng, W.-H., Lo, K.-L., Lee, Y.-H., Hung, T.-H., & Lin, Y.-C. (2007). Berberine produces antidepressant-like effects in the forced swim test and in the tail suspension test in mice. *Life sciences*, 81(11), 933-938 .
- Pérez, M .Á., Terreros, G., & Dagnino-Subiabre, A. (2013). Long-term omega-3 fatty acid supplementation induces anti-stress effects and improves learning in rats. *Behavioral and Brain Functions*, 9(1), 25 .
- Pham, K., Nacher, J., Hof, P. R., & McEwen, B. S. (2003). Repeated restraint stress suppresses neurogenesis and induces biphasic PSA-NCAM expression in the adult rat dentate gyrus. *European Journal of Neuroscience*, 17(4), 879-886 .
- Phillis, J. W., & O'Regan, M. H. (2004). A potentially critical role of phospholipases in central nervous system ischemic, traumatic, and neurodegenerative disorders. *Brain research reviews*, 44(1), 13-47 .
- Pike, J. L., & Irwin, M. R. (2006). Dissociation of inflammatory markers and natural killer cell activity in major depressive disorder. *Brain, behavior, and immunity*, 20(2), 169-174 .

- Pischon, T., Hankinson, S. E., Hotamisligil, G. S., Rifai, N., Willett, W. C., & Rimm, E. B. (2003). Habitual dietary intake of n-3 and n-6 fatty acids in relation to inflammatory markers among US men and women. *Circulation*, 108(2), 155-160 .
- Plamondon, H., & Roberge, M.-C. (2008). Dietary PUFA supplements reduce memory deficits but not CA1 ischemic injury in rats. *Physiology & behavior*, 95(3), 492-500 .
- Porsolt, R. D., Anton, G., Blavet, N., & Jalfre, M. (1978). Behavioural despair in rats: a new model sensitive to antidepressant treatments. *European journal of pharmacology*, 47(4), 379-391 .
- Raison, C. L., & Miller, A. H. (2003). When not enough is too much: the role of insufficient glucocorticoid signaling in the pathophysiology of stress-related disorders. *American Journal of Psychiatry*, 160(9), 1554-1565 .
- Riediger, N. D., Othman, R. A., Suh, M., & Moghadasian, M. H. (2009). A systemic review of the roles of n-3 fatty acids in health and disease. *Journal of the American Dietetic Association*, 109(4), 668-679 .
- Riemer, S., Maes, M., Christophe, A., & Rief, W. (2010). Lowered ω -3 PUFAs are related to major depression, but not to somatization syndrome. *Journal of affective disorders*, 123(1), 173-180 .
- Rodrigues, A. L. S., Rocha, J. B. T., Mello, C. F., & Souza, D. O. (1996). Effect of Perinatal Lead Exposure on Rat Behaviour in Open-Field and Two-Wky Avoidance Tasks. *Pharmacology & toxicology*, 79(3), 150-156 .
- Rosenberger, T. A., Villacreses, N. E., Hovda, J. T., Bosetti, F., Weerasinghe, G., Wine, R. N., . . . Rapoport, S. I. (2004). Rat brain arachidonic acid metabolism is increased by a 6-day intracerebral ventricular infusion of bacterial lipopolysaccharide. *Journal of neurochemistry*, 88(5), 1168-1178 .
- Rosenblatt, J. S., Olufowobi, A., & Siegel, H. I. (1998). Effects of pregnancy hormones on maternal responsiveness, responsiveness to estrogen stimulation of maternal behavior, and the lordosis response to estrogen stimulation. *Hormones and behavior*, 33(2), 114-124 .
- Sachar, E. J., & Baron, M. (1979). The biology of affective disorders. *Annual review of neuroscience*, 2(1), 505-518 .
- Sastry, P. (1985). Lipids of nervous tissue: composition and metabolism. *Progress in lipid research*, 24(2), 69-176 .
- Schiepers, O. J., Wichers, M. C., & Maes, M. (2005). Cytokines and major depression. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 29(2), 201-217 .
- Schüle, C., Baghai, T. C., Eser, D., Häfner, S., Born, C., Herrmann, S., & Rupprecht, R. (2009). The combined dexamethasone/CRH Test (DEX/CRH test) and prediction of acute treatment response in major depression. *PloS one*, 4(1), e4324 .
- Sebastian, L. (1998). *Understanding Postpartum Depression and Anxiety*: Addicus Books.
- Sebastian, L. (1999). *Overcoming Postpartum Depression and Anxiety*: LWW.
- Selikoff, I. J., & Robitzek, E. H. (1952). Tuberculosis Chemotherapy with Hydrazine Derivatives of Isonicotinic Acid. *CHEST Journal*, 21(4), 385-438 .
- Siegel, H. I., & Rosenblatt, J. S. (1975). Estrogen-induced maternal behavior in hysterectomized-ovariectomized virgin rats. *Physiology & behavior*, 14(4), 465-471 .

- Silvers, K. M., & Scott, K. M. (2002). Fish consumption and self-reported physical and mental health status. *Public health nutrition*, 5(3), 427-43 .2
- Simopoulos, A. P. (2002). The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomedicine & pharmacotherapy*, 56(8), 365-379 .
- Simopoulos, A. P., Leaf, A., & Salem Jr, N. (1999). Workshop on the essentiality of and recommended dietary intakes for omega-6 and omega-3 fatty acids. *Journal of the American College of Nutrition*, 18(5), 487-489 .
- Sit, D. K., Rothschild, A. J., & Wisner, K. L. (2006). A review of postpartum psychosis. *Journal of women's health*, 15(4), 352-368 .
- Sit, D. K & ,Wisner, K. L. (2005). Decision Making for Postpartum Depression Treatment-Effective treatment decisions in postpartum depression must be a collaborative process between patient and provider. *Psychiatric Annals*, 35(7), 577-585 .
- Slattery, D. A., & Neumann ,I. D. (2008). No stress please! Mechanisms of stress hyporesponsiveness of the maternal brain. *The Journal of physiology*, 586(2), 377-385 .
- Smith, R. (1991). The macrophage theory of depression. *Medical hypotheses*, 35(4), 298-306 .
- Song, C., Li, X., Kang, Z., & Kadotomi, Y. (2007). Omega-3 Fatty Acid Ethyl-Eicosapentaenoate Attenuates IL-1 β -Induced Changes in Dopamine and Metabolites in the Shell of the Nucleus Accumbens: Involved with PLA2 Activity and Corticosterone Secretion. *Neuropsychopharmacology*-1207 ,(5)32 , .1207
- Sonino, N., & Fava, G. A. (2002). Residual symptoms in depression: an emerging therapeutic concept. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 26(4), 763-770 .
- Sontrop, J., & Campbell, M. K. (2006). ω -3 polyunsaturated fatty acids and depression: a review of the evidence and a methodological critique. *Preventive medicine*, 42(1), 4-13 .
- Sowash, J. R. (2009). Dissection of the Rat. from <http://www.freebookez.com/ebook.php?u=aHR0cDovL2pyc293YXNoLndpa2lzcGFjZXMuY29tL2ZpbGUvdmlldy9yYXQuc3R1ZGVudC5wZGYKRGlzc2VjdGlvbiBvZiB0aGUgUmF0>. Retrieved April 20, 2014
- Spedding, M., Neau, I., & Harsing, L. (2003). Brain plasticity and pathology in psychiatric disease: sites of action for potential therapy. *Current opinion in pharmacology*, 3(1), 33-40 .
- Starkman, M. N., Giordani, B., Gebarski, S. S., & Schteingart, D. E. (2007). Improvement in mood and ideation associated with increase in right caudate volume. *Journal of affective disorders*, 101(1), 139-147 .
- Steiner, M. (1998). Perinatal mood disorders: position paper. *Psychopharmacology Bulletin* .
- Stern, E. Y., & Kalynchuk, L. E. (2010). Behavioral and neurobiological consequences of prolonged glucocorticoid exposure in rats: relevance to depression. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 34(5), 777-790 .
- Stocky, A., & Lynch, J. (2000). Acute psychiatric disturbance in pregnancy and the puerperium. *Best Practice & Research Clinical Obstetrics & Gynaecology*, 14(1), 73-87 .

- Stoffel, E. C., & Craft, R. M. (2004). Ovarian hormone withdrawal-induced "depression" in female rats. *Physiology & behavior*, 83(3), 505-513 .
- Stoll, A. L., Severus, W. E., Freeman, M. P., Rueter, S., Zboyan, H. A., Diamond, E., . . . Marangell, L. B. (1999). Omega 3 fatty acids in bipolar disorder: a preliminary double-blind, placebo-controlled trial. *Archives of General Psychiatry*, 56(5), 407 .
- Su, K.-P., Huang, S.-Y., Chiu, C.-C., & Shen, W. W. (2003). Omega-3 fatty acids in major depressive disorder: a preliminary double-blind, placebo-controlled trial. *European Neuropsychopharmacology*, 13(4), 267-271 .
- Su, K.-P., Huang, S.-Y., Chiu, T.-H., Huang, K.-C., Huang, C.-L., Chang, H.-C., & Pariante, C. M. (2008). Omega-3 fatty acids for major depressive disorder during pregnancy: results from a randomized, double-blind, placebo-controlled trial. *Journal of Clinical Psychiatry*, 69(4), 644 .
- Suda, S., Segi-Nishida, E., Newton, S. S., & Duman, R. S. (2008). A postpartum model in rat: behavioral and gene expression changes induced by ovarian steroid deprivation. *Biological psychiatry*, 64(4), 311-319 .
- Sun, G. Y., Xu, J., Jensen, M. D., & Simonyi, A. (2004). Phospholipase A2 in the central nervous system implications for neurodegenerative diseases. *Journal of Lipid Research*, 45(2), 205-213 .
- T Abou-Saleh, M., Ghubash, R., Karim, L., Krymski, M., & Bhai, I. (1998). Hormonal aspects of postpartum depression. *Psychoneuroendocrinology*, 23(5), 465-475 .
- Tanskanen, A., Hibbeln, J. R., Hintikka, J., Haatainen, K., Honkalampi, K., Viinamäki, H., & Stoll, A. L. (2001a). Fish consumption, depression, and suicidality in a general population. *Archives of General Psychiatry*, 58(5), 512 .
- Tanskanen, A., Hibbeln, J. R., Tuomilehto, J., Uutela, A., Haukkala, A., Viinamäki, H., . . . Vartiainen, E. (2001b). Fish consumption and depressive symptoms in the general population in Finland. *Psychiatric Services*, 52(4), 529-531 .
- Tiemeier, H., van Tuijl, H. R., Hofman, A., Kiliaan, A. J., & Breteler, M. M. (2003). Plasma fatty acid composition and depression are associated in the elderly: the Rotterdam Study. *The American journal of clinical nutrition*, 78(1), 40-46 .
- Tinoco, J. (1982). Dietary requirements and functions of α -linolenic acid in animals. *Progress in lipid research*, 21(1), 1-45 .
- Tsigos, C., & Chrousos, G. P. (2002). Hypothalamic-pituitary-adrenal axis, neuroendocrine factors and stress. *Journal of psychosomatic research*, 53(4), 865-871 .
- Vancassel, S., Leman, S., Hanonick, L., Denis, S., Roger, J., Nollet, M., . . . Chalon, S. (2008). n-3 polyunsaturated fatty acid supplementation reverses stress-induced modifications on brain monoamine levels in mice. *Journal of Lipid Research*, 49(2), 340-348 .
- Venna, V. R., Deplanque, D., Allet, C., Belarbi, K., Hamdane, M., & Bordet, R. (2009). PUFA induce antidepressant-like effects in parallel to structural and molecular changes in the hippocampus. *Psychoneuroendocrinology*, 34(2), 199-211 .
- Walker, C. D., Trottier, G., Rochford, J., & Lavallée, D. (1995). Dissociation between behavioral and hormonal responses to the forced swim stress in lactating rats. *Journal of neuroendocrinology*, 7(8), 615-622 .
- Weiss, J. M., Goodman, P. A., Losito, B. G., Corrigan, S., Charry, J. M., & Bailey, W. H. (1981). Behavioral depression produced by an uncontrollable stressor:

- relationship to norepinephrine, dopamine, and serotonin levels in various regions of rat brain. *Brain research reviews*, 3(2), 167-205 .
- Willner, P. (1984). The validity of animal models of depression. *Psychopharmacology*, 83(1), 1-16 .
- Wong, E., & Herbert, J. (2006).(Raised circulating corticosterone inhibits neuronal differentiation of progenitor cells in the adult hippocampus. *Neuroscience*, 137(1), 83-92 .
- Xiang, H., Liu, Y., Zhang, B., Huang, J., Li, Y., Yang, B., . . . Zhang, H. (2011). The antidepressant effects and mechanism of action of total saponins from the caudexes and leaves of *Panax notoginseng* in animal models of depression. *Phytomedicine*, 18(8), 731-738 .
- Yi, L.-T., Li, J., Li, H.-C., Su, D.-X., Quan, X.-B., He, X.-C., & Wang, X.-H. (2012). Antidepressant-like behavioral, neurochemical and neuroendocrine effects of naringenin in the mouse repeated tail suspension test. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 39(1), 175-181 .
- Yi, L.-T., Li, J., Liu, Q., Geng, D., Zhou ,Y.-F., Ke, X.-Q., . . . Weng, L.-J. (2013). Antidepressant-like effect of oleanolic acid in mice exposed to the repeated forced swimming test. *Journal of Psychopharmacology*, 27(5), 459-468 .
- Yi, L.-T., Xu, Q., Li, Y.-C., Yang, L., & Kong, L.-D. (2009). Antidepressant-like synergism of extracts from magnolia bark and ginger rhizome alone and in combination in mice. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 33(4), 616-624 .
- Zarrow, M., Gandelman, R., & Denenberg, V. (1971). Prolactin :Is it an essential hormone for maternal behavior in the mammal? *Hormones and behavior*, 2(4), 343-354 .
- Zeller, E., & Barsky, J. (1952). *In vivo inhibition of liver and brain monoamine oxidase by 1-Isonicotinyl-2-isopropyl hydrazine*. Paper presented at the Proceedings of the Society for Experimental Biology and Medicine. Society for Experimental Biology and Medicine (New York, NY.)
- Zhu, C.-B., Blakely, R. D., & Hewlett, W. A. (2006). The proinflammatory cytokines interleukin-1beta and tumor necrosis factor-alpha activate serotonin transporters. *Neuropsychopharmacology*, 31(10), 2121-2131 .
- Zonana, J., & Gorman, J. M. (2005). The neurobiology of postpartum depression. *CNS spectrums*, 10(10), 792 .
- Zorrilla, E. P., Luborsky, L., McKay, J. R., Rosenthal, R., Houldin, A., Tax, A., . . . Schmidt, K. (2001). The relationship of depression and stressors to immunological assays: a meta-analytic review. *Brain, behavior, and immunity*, 15(3), 199-226 .