



UNIVERSITI PUTRA MALAYSIA

***ANTITUMOUR EFFECTS OF *Morinda citrifolia* L. FRUIT AND
Clinacanthus nutans (Burm. F.) LINDAU LEAF MIXTURES IN
LEUKAEMIA-LYMPHOMA BEARING SPRAGUE DAWLEY RATS***

**SAJJARATTUL NURUL NADIA ASYURA MOHAMAD
IFFENDI**

FPV 2016 37



ANTITUMOUR EFFECTS OF *Morinda citrifolia* L. FRUIT AND *Clinacanthus nutans* (Burm. F.) LINDAU LEAF MIXTURES IN LEUKAEMIA-LYMPHOMA BEARING SPRAGUE DAWLEY RATS

By

SAJJARATTUL NURUL NADIA ASYURA MOHAMAD IFFENDI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Doctor Philosophy**

December 2016

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION


This Thesis is Dedicated to

My Beloved Parents, Sisters,

Lecturers and Friends

Without Whom, None of My Success Would Be Possible

With Love, Nadia Asyura



Abstract of thesis prepared to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

ANTITUMOUR EFFECTS OF MIXTURES OF *Morinda citrifolia* L. FRUIT AND *Clinacanthus nutans* (Burm. F.) LINDAU LEAF EXTRACTS IN LEUKAEMIA-LYMPHOMA BEARING SPRAGUE DAWLEY RATS

By

SAJJARATTUL NURUL NADIA ASYURA MOHAMAD IFFENDI

December 2016

Chairman: Hazilawati Hj. Hamzah, PhD
Faculty: Veterinary Medicine

Morinda citrifolia L. (*M. citrifolia*) fruit known as ‘mengkudu’ and *Clinacanthus nutans* (Burm. F.) Lindau (*C. nutans*) leaf known as ‘rumpit belalai gajah’, are widely known herbs that have anti-cancer properties. The antitumour effects of the combination of these herbs in leukaemia-lymphoma have never been reported. The aim of this study were to evaluate toxicity of ethanol extracts of *M. citrifolia* fruits and *C. nutans* leaves and the antitumour effects of these ethanol extracts in MNU-induced leukaemia-lymphoma. The antitumour of MNU-induced leukaemia-lymphoma rats was evaluated via haematological and serum biochemical parameters, histopathology, levels of an oxidative stress biomarker (malondialdehyde [MDA]) and antioxidative stress enzymes (glutathione peroxidase [GPx] and superoxide dismutase [SOD], transcription levels of VEGF and COX-2 mRNA in the blood, spleen and lymph nodes and expression levels of VEGF and COX-2 protein in the spleen and lymph nodes. A 14-day acute toxicity was conducted for determination of lethal dose 50 (LD₅₀), followed by a 28-day subacute toxicity study and a 13-week subchronic toxicity study of both extracts. In acute toxicity, rats were divided into four groups, namely, control, 10% DMSO (vehicle) and 2000 mg/kg of body weight of each extract. In the subacute and subchronic toxicities, rats were divided into eight groups including control, 10% DMSO (vehicle), 75 mg/kg body weight, 125 mg/kg body weight and 250 mg/kg body weight of each extract. At the end of the experimental period, blood samples were collected for the hematological and serum biochemical parameters. The liver and kidneys were collected for histopathological analysis. The results showed a single dose (2000 mg/kg of body weight) of ethanol extracts of *M. citrifolia* fruits and *C. nutans* leaf induced no acute toxicity to the male Sprague Dawley rats. The oral administration of 75 mg/kg of *M. citrifolia* fruit and *C. nutans* leaf for 28 and 90 days induced no subacute and subchronic toxicity in male Sprague Dawley rats. Administration of 125 mg/kg of *M. citrifolia* fruit and *C. nutans* leaf for 28 and 90 days induced hepatotoxicity in male Sprague Dawley rats. Administration of 250

mg/kg of *M. citrifolia* fruit and *C. nutans* leaf for 28 and 90 days induced hepatotoxicity and nephrotoxicity in male Sprague Dawley rats.

In the MNU-induced leukaemia-lymphoma study, a total of 56 rats were divided into seven groups namely control (group A), MNU-induced leukaemia-lymphoma rats (group B), MNU-induced leukaemia-lymphoma rats treated with four different combinations of *M. citrifolia* and *C. nutans* extracts (mixture C [group C], mixture D [group D], mixture E [group E] and mixture F [group F]) and MNU-induced leukaemia-lymphoma rats treated with chemotherapy regimen (CHOP) (group G). Leukaemia-lymphoma was induced using *N*-methyl-*N*-nitrosourea (MNU) which was administered intraperitoneally (i.p) at a dose of 60 mg/kg of body weight for four times in the 2-week period. Mixtures of herbs for treatment of the leukaemia-lymphoma rats were formulated not to exceed 250 mg/kg of body weight per administration. The herbal mixed extracts were given via oral gavage, once daily starting from week 12 post MNU administration until week 24 of the experiment period. Blood samples were collected and analysed on the 18th and 24th weeks of the experiment. Heart, lungs, liver kidneys, lymph nodes and spleen were collected for histopathological evaluation. The results showed the presence of blast cells in the blood smear in both untreated and treated leukaemia-lymphoma rats at week 18th and 24th. Groups B and G had a significant elevation in the number of blast cells from week 18 to week 24. The oral administration of herbal mixtures (groups C, D, E and F) was effective in limiting the degree of leukaemic cells progression. The serum biochemical parameter results showed CK level was significantly increased in groups C, D and F from week 18 to week 24. The LDH level had also significantly increased in groups C and F from week 18 to week 24. Histopathology results showed neoplastic lymphocytes were present in all spleen and lymph nodes of the untreated and treated MNU-induced leukaemia-lymphoma rats. The oral administration of mixtures E and F in MNU-induced leukaemia-lymphoma rats inhibited metastasise of neoplastic lymphocytes to the selected vital organs (liver, lungs, heart and kidneys) as compared to other herbal mixtures and CHOP regimen. Treatment of CHOP regimen in MNU-induced leukaemia-lymphoma rats induced significant liver and spleen fibrosis and necrosis in the testis. However, treatment of herbal mixtures in MNU-induced leukaemia-lymphoma rats induced no fibrosis and necrosis (testis).

The levels of serum MDA in the MNU-induced leukaemia-lymphoma rats treated with all herbal mixtures and CHOP regimen were decreased from week 18 to week 24. The level of serum SOD in MNU-induced leukaemia-lymphoma rats treated with herbal mixtures and CHOP regimen increased from week 18 to week 24. A decrease of serum GPx was observed from week 18 to week 24 in all groups. Administration of MNU-induced leukaemia-lymphoma rats with herbal mixtures succeeded in inhibiting angiogenesis and inflammation in blood, lymph node and spleen compared to untreated MNU-induced leukaemia-lymphoma rats. A down-regulated of VEGF and COX-2 mRNA transcription was observed especially in MNU-induced leukaemia-lymphoma rats treated with mixtures E and F compared to untreated MNU-induced leukaemia-lymphoma. On the other hand, treatment of mixture F succeeded to inhibit COX-2 expression in the lymph nodes of MNU-induced leukaemia-lymphoma rats via immunohistochemistry. A more sensitive result was obtained from Western blot

whereby mixtures E and F were able to decrease VEGF and COX-2 expression in MNU-induced leukaemia-lymphoma rats.

In conclusion, oral administration of 2000 mg/kg of body weight of both extracts in male Sprague Dawley rats induced no acute toxicity. Oral administration of 125 mg/kg of body weight for 29 and 90 days of both extracts induced hepatotoxicity. Oral administration of 250 mg/kg of body weight for 29 and 90 days of both extracts induced hepatotoxicity and nephrotoxicity. Oral administration of herbal extracts (group C, D, E and F) limited leukaemic cell progression in MNU-induced leukaemia-lymphoma rats. Oral administration of mixtures E and F inhibited the metastasise of neoplastic lymphocytes in heart, lungs, liver and kidneys. Oral administration of all herbal mixtures down-regulated the serum MDA level in the MNU-induced leukaemia-lymphoma rats. It also enhanced the production of serum SOD level in the MNU-induced leukaemia-lymphoma rats treated with all herbal mixtures. Oral administration of mixtures E and F was able to down-regulate VEGF and COX-2 transcription and expression in MNU-leukaemia-lymphoma rats.

Abstrak thesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai mematuhi keperluan untuk Ijazah Doktor Falsafah

**KESAN ANTITUMOR CAMPURAN EKSTRAK BUAH *Morinda citrifolia* L.
DAN DAUN *Clinacanthus nutans* (Burm. F.) LINDAU TERHADAP MODEL
TIKUS SPRAGUE DAWLEY LEUKEMIA-LIMFOMA**

Oleh

SAJJARATTUL NURUL NADIA ASYURA MOHAMAD IFFENDI

Disember 2016

Pengerusi: Hazilawati Hj. Hamzah, PhD
Fakulti: Perubatan Veterinar

Buah *Morinda citrifolia* L. (*M. citrifolia*) atau dikenali sebagai ‘mengkudu’ dan daun *Clinacanthus nutans* (Burm. F.) Lindau (*C. nutans*) atau dikenali sebagai ‘rumpun belalai gajah’ adalah tumbuhan herba yang dikenali umum dan mempunyai ciri-ciri anti-kanser. Kesan anti-tumor daripada kombinasi ekstrak-ekstrak ini ke atas leukemia dan limfoma belum lagi dilaporkan. Tujuan bahagian pertama kajian ini dijalankan adalah untuk menilai ketoksikan oral ekstrak etanol buah *M. citrifolia* dan daun *C. nutans* dan kesan anti-tumor kombinasi ekstrak etanol ini ke atas MNU-induksi leukemia-limfoma di dalam tikus Sprague Dawley melalui penilaian hematologi dan biokimia serum, histopatologi, tahap penanda-bio tekanan oksidatif [malondialdehyde] dan enzim tekanan anti-oksidatif (glutathione peroksidase) [GSH-Px] dan superoksida dismutase [SOD], tahap transkripsi mRNA VEGF dan COX-2 di dalam darah, limfa dan nodus limfa dan tahap ekspresi protein VEGF dan COX-2 di dalam darah, limfa dan nodus limfa. Kajian ketoksikan akut selama 14 hari telah dijalankan untuk menilai dos kematian purata (LD₅₀), diikuti dengan kajian 28 hari ketoksikan subakut dan 13 minggu kajian ketoksikan subkronik untuk kedua-dua ekstrak. Di dalam kajian ketoksikan akut, tikus-tikus dibahagikan kepada empat kumpulan iaitu termasuk kawalan, 10% DMSO (pengangkutan) dan 2000 mg/kg daripada berat badan untuk kedua-dua ekstrak. Manakala di dalam kajian ketoksikan subakut, tikus-tikus dibahagikan kepada lapan kumpulan iaitu kawalan, 10% DMSO (pengangkutan), 75 mg/kg daripada berat badan, 125 mg/kg daripada berat badan dan 250 mg/kg daripada berat badan untuk kedua-dua ekstrak. Penilaian piawai toksikologi termasuk kadar kematian, patologi kasar, hematologi, biokimia serum, dan pemeriksaan histopatologi ke atas hati dan buah pinggang telah dilakukan di akhir kajian. Keputusan kajian menunjukkan dos tunggal ekstrak etanol (2000 mg/kg) untuk buah *M. citrifolia* dan daun *C. nutans* tidak mengakibatkan ketoksikan akut ke atas tikus jantan Sprague Dawley. Pemberian oral buah *M. citrifolia* dan daun

C. nutans pada dos 75 mg/kg daripada berat badan untuk 28 dan 90 hari tidak mengakibatkan ketoksikan subakut dan subkronik ke atas tikus jantan Sprague Dawley. Pemberian oral buah *M. citrifolia* dan daun *C. nutans* pada dos 125 mg/kg daripada berat badan untuk 28 dan 90 hari mengakibatkan hepatotoksikiti ke atas tikus jantan Sprague Dawley. Pemberian oral buah *M. citrifolia* dan daun *C. nutans* pada dos 250 mg/kg daripada berat badan untuk 28 dan 90 hari mengakibatkan hepatotoksikiti dan nefrotoksikiti ke atas tikus jantan Sprague Dawley.

Di dalam kajian MNU-induksi leukemia-limfoma, tikus dibahagi kepada tujuh kumpulan termasuk tikus kawalan (kumpulan A), tikus MNU-induksi leukemia-limfoma (kumpulan B), tikus leukemia-limfoma dirawat dengan empat kombinasi buah *M. citrifolia* dan daun *C. nutans* yang berlainan (kombinasi C [kumpulan C], kombinasi D [kumpulan D], kombinasi E [kumpulan E] dan kombinasi F [kumpulan F]) dan tikus leukemia-limfoma dirawat dengan regimen kimoterapi (CHOP) (kumpulan G). Tikus leukemia-limfoma telah diinduksi dengan menggunakan *N*-metil-*N*-nitrosourea (MNU) secara intraperitoneum (i.p) pada dos 60 mg/kg daripada berat badan, diberi sebanyak empat kali dalam tempoh dua minggu. Ekstrak etanol buah *M. citrifolia* dan daun *C. nutans* tersebut telah diberi secara gajaj oral sekali sehari bermula pada minggu ke-12 sehingga minggu ke-24 setelah MNU-diinduksikan. Sampel darah telah diambil dan dianalisa pada minggu ke-12 dan minggu ke-24. Jantung, paru-paru, hati, buah pinggang, modul limfa dan limfa telah diambil untuk dianalisa dan dinilai dengan lebih lanjut. Keputusan darah menunjukkan kehadiran sel-sel blastik pada minggu ke-12 dan minggu ke-24 di dalam tikus tidak dirawat dan dirawat. Kumpulan B and G mempunyai penakikan yang ketara dalam nombor sel blastik dari minggu 18 ke 24. Pemberian oral kombinasi herba (kumpulan C, D, E dan F) telah menunjukkan keberkesanan dalam mengawal perkembangan sel-sel leukemia. Keputusan biokimia serum menunjukkan aras CK mempunyai peningkatan yang ketara dari minggu 18 ke 24 di dalam kumpulan C, D dan F. Aras LDH juga menunjukkan peningkatan yang ketara dari minggu 18 ke 24 di dalam kumpulan C dan F. Keputusan histopatologi menunjukkan kehadiran limfosit neoplastik di dalam semua limfa dan nodul limfa pada tikus yang tidak dirawat dan dirawat (kombinasi herba dan CHOP). Pemberian oral kombinasi E dan F ke atas MNU-induksi leukaemia-lymphoma merencat metastasis limfosit neoplastik ke jantung, paru-paru, hati dan buah pinggang. Rawatan regimen CHOP ke atas tikus MNU-induksi leukaemia-lymphoma menyebabkan fibrosis yang ketara pada hati dan limfa serta nekrosis pada testis. Walau bagaimanapun, rawatan kombinasi herba ke atas tikus MNU-induksi leukaemia-lymphoma tidak mengakibatkan fibrosis dan nekrosis (testis). Tahap serum MDA telah menurun daripada minggu 18 ke minggu 24 di dalam tikus MNU-induksi leukemia-limfoma yang dirawat (kombinasi herba dan regimen CHOP). Tahap serum SOD di dalam tikus MNU-induksi leukemia-limfoma telah menaik daripada minggu 18 ke minggu 24 setelah dirawat dengan kombinasi herba dan regimen CHOP. Penurunan serum GPx telah diperhatikan pada minggu 18 ke minggu 24 dalam kesemua kumpulan tikus. Pemberian oral kombinasi herba kepada tikus MNU-induksi leukemia-limfoma telah berjaya merencat angiogenesis dan keradangan di dalam darah, limfa dan nodus limfa. Penurunan transkripsi mRNA VEGF dan COX-2 telah diperhatikan di dalam kumpulan tikus MNU-induksi leukemia-limfoma yang dirawat dengan kombinasi E dan F apabila dibandingkan dengan tikus MNU-induksi leukemia-limfoma yang tidak dirawat. Selain itu juga,

kombinasi F berjaya merencat ekspresi COX-2 ke atas nodul limfa tikus. MNU-induksi leukemia-limfoma melalui penilaian immunohistokimia. Keputusan yang lebih tepat dan sensitif diperolehi daripada ujian Western blot di mana kombinasi E and F berjaya merencat ekspresi VEGF dan COX-2 di dalam tikus MNU-induksi leukemia-limfoma.

Secara konklusinya, pemberian dos tunggal (2000 mg/kg daripada berat badan) secara oral ekstrak etanol buah *M. citrifolia* dan daun *C. nutans* pada tidak mengakibatkan ketosikan akut ke atas tikus Sprague Dawley jantan. Pemberian oral ekstrak etanol buah *M. citrifolia* dan daun *C. nutans* pada dos 125 mg/kg daripada berat badan mengakibatkan hepatotoksiti ke atas tikus Sprague Dawley jantan. Pemberian oral ekstrak etanol buah *M. citrifolia* dan daun *C. nutans* pada dose 250 mg/kg daripada berat badan mengakibatkan hepatotoksiti dan kenefrotoksikan ke atas tikus Sprague Dawley jantan. Pemberian oral kombinasi herba (kombinasi C, D, E dan F) merencat perkembangan sel-sel leukaemia di dalam tikus MNU-induksi leukemia-limfoma. Pemberian oral kombinasi E dan F merencat metastasis limfosit neoplastic ke dalam jantung, paru-paru, hati dan buah pinggang. Pemberian oral kesemua kombinasi herba menurunkan tahap serum MDA di dalam tikus MNU-induksi leukemia-limfoma. Ia juga meningkatkan penghasilan serum SOD di dalam tikus MNU-induksi leukemia-limfoma. Pemberian oral kombinasi E dan F menurunkan transkripsi dan ekspresi VEGF dan COX-2 di dalam tikus MNU-induksi leukemia-limfoma.

ACKNOWLEDGEMENTS

Alhamdulillah and praise to Allah S.W.T for HIS blessings and guidance in my life, particularly during the completion of my degree of Doctor of Philosophy in Universiti Putra Malaysia (UPM).

First of all, I would like to thank my beloved mother, Haslina binti Hj. Hamdan and my beloved father, Mohamad Iffendi bin Hj. Ibrahim. Without their love, blessings and supports over the years, none of this would have been possible. I also would like to show my greatest gratitude to my lovely sisters, Nadrah Nabilah binti Mohamad Iffendi and Nadirah Atikah binti Mohamad Iffendi for their supports, understanding and help during the difficult times I faced.

I would like to express my greatest gratitude and appreciation to my supervisor, Associate Prof. Dr. Hazilawati Hj. Hamzah, the chairman of the supervisory committee, for her continuous supervision, help and guidance. Without her help, I would not be where I am today. I also wish to thank her for the knowledge she has shared, and I will always be grateful for having the opportunity to be under her supervision throughout my research and during my study period at the Faculty of Veterinary Medicine, UPM.

My special thanks also go to the supervisory committee members, Prof. Dr. Noordin Mohamed Mustapha, Dr. Shanmugavelu a/l Sithambaram and Associate Prof. Dr. Norhaizan Md. Esa for their guidance, constructive comments and for providing me with all the facilities required for my research. Last but not least, my special thanks go to Dr. Mohd. Rosly Shaari for his guidance in toxicity study and also his assistance in helping me conduct the animal study.

I would like to extend my greatest appreciation to all laboratory technicians in the Department of Microbiology and Pathology Laboratories, Faculty Veterinary Medicine, UPM, especially Mr. Halmi, Mr. Abdullah, Miss Darul, Mr. Arman and Miss Yana for their never-ending assistance during my research. Special appreciation also goes to my labmates, Syahirah, Farhan, Hutheyfa, Shayma, Hakimi and Yusnaini for their friendship and assistance. To my best friends, Fareeza Farihah Basri and Noor Farhana Bachek, I am grateful to have them in my life, and I am thankful for their friendship.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as a fulfillment of the requirement for the degree of Doctor of Philosophy. The members of Supervisory Committee were as follows:

Hazilawati Hamzah, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Noordin Mohamed Mustapha, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Norhaizan Mohamad Esa, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Shanmugavelu Sithabaram, PhD

Director
Animal Research Centre
Malaysian Agriculture Research and Development
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by Graduate Student

I hereby confirm that

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules, 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is publish (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____

Date: _____

Name and matric no: Sajjarattul Nurul Nadia Asyura; GS35670

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of Chairman of Supervisory Committee: Associate Professor
Hazilawati Hj. Hamzah

Signature: _____

Name of Member of Supervisory Committee: Professor
Noordin Mohamed Mustapha

Signature: _____

Name of Member of Supervisory Committee: Associate Professor
Norhaizan Mohamad Esa

Signature: _____

Name of Member of Supervisory Committee: Director
Dr. Shanmugavelu Sithabaram

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iv
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xvi
LIST OF FIGURES	xxii
LIST OF ABBREVIATIONS	xxv
 CHAPTER	
1 INTRODUCTION	1
 2 LITERATURE REVIEW	5
2.1. <i>Morinda citrifolia</i>	5
2.1.1. Plant	5
2.1.2. Traditional use	6
2.1.3. Toxicity of <i>M. citrifolia</i>	6
2.1.4. Bioactive compounds	7
2.1.5. Anti-cancer properties	8
2.2. <i>Clinacanthus nutans</i>	8
2.2.1. Plant	8
2.2.1. Traditional use	9
2.2.3. Toxicity of <i>C. nutans</i>	10
2.1.4. Bioactive compounds	10
2.3. Toxicity lesions of liver and kidneys induced chemicals	11
2.4. Cancer	12
2.4.1. Leukaemia	12
2.4.2. Lymphoma	14
2.4.3. <i>N</i> -methyl- <i>N</i> -nitrosurea induced leukaemia-lymphoma	15
2.4.4. Treatment of leukaemia-lymphoma	15
2.5. Angiogenesis and cancer	16
2.6. Inflammation and cancer	18
2.7. Oxidative stress and cancer	19
2.8. Animal model for leukaemia-lymphoma (Clinico-Pathological changes of MNU-induced leukaemia-lymphoma)	22
2.8.1. Mortality, clinical signs, necroscopy findings, body weight and organ relative weight of MNU-induced leukaemia-lymphoma.	22
2.8.2. Haematological parameters	23
2.8.3. Serum biochemical parameters	23
2.8.4. Histopathological evaluation	24

3	TOXICITY EVALUATION OF <i>Morinda citrifolia</i> L. fruit AND <i>Clinacanthus nutans</i> (Burm. F) LINDAU LEAF EXTRACTS IN MALE SPRAGUE DAWLEY RATS	25
3.1.	Introduction	25
3.2.	Materials and methods	26
3.2.1.	Plant preparation	26
3.2.2.	Plant extraction	26
3.2.3.	Animal management and routine	27
3.2.4.	Experimental design	27
3.2.5.	Blood samplings	28
3.2.6.	Haematology analysis	29
3.2.6.1.	Complete blood count	29
3.2.6.2.	Packed cell volume	29
3.2.6.3.	Icterus index	29
3.2.6.4.	Plasma protein determination	29
3.2.7.	Serum biochemical analysis	30
3.2.8.	Histopathology analysis	30
3.2.8.1.	Toxicity lesions characteristics	30
3.3.	Results	31
3.3.1.	Acute toxicity of the ethanol extract of <i>M. citrifolia</i> fruit	33
3.3.1.1.	Body weight	33
3.3.1.2.	Haematological and serum biochemical parameters and organ to body weight ratio	33
3.3.1.3.	Lesion score	37
3.3.2.	Subacute toxicity of the ethanol extract of <i>M. citrifolia</i> fruit	41
3.3.2.1.	Body weight	41
3.3.2.2.	Haematological and serum biochemical parameters and organ to body weight ratio	41
3.3.2.3.	Lesion score	45
3.3.3.	Subchronic toxicity of the ethanol extract of <i>M. citrifolia</i> fruit	49
3.3.3.1.	Body weight	49
3.3.3.2.	Haematological and serum biochemical parameters and organ to body weight ratio	50
3.3.3.3.	Lesion score	54
3.3.4.	Acute toxicity of the ethanol extract of <i>C. nutans</i> leaf	58
3.3.4.1.	Body weight	58
3.3.4.2.	Haematological and serum biochemical parameters and organ to body weight ratio	58
3.3.4.3.	Lesion score	62
3.3.5.	Subacute toxicity of the ethanol extract of <i>C. nutans</i> leaf	66
3.3.5.1.	Body weight	66
3.3.5.2.	Haematological and serum biochemical parameters and organ to body weight ratio	66
3.3.5.3.	Lesion score	70
3.3.6.	Subchronic toxicity of the ethanol extract of	

	<i>C. nutans</i> leaf	74
	3.3.6.1. Body weight	74
	3.3.6.2. Haematological and serum biochemical parameters and organ to body weight ratio	74
	3.3.6.3. Lesion score	78
3.4.	Discussion	84
3.5.	Conclusion	88
4	CLINICAL AND HISTOPATHOLOGICAL EVALUATION OF MIXTURES OF <i>Morinda citrifolia</i> FRUIT L. AND <i>Clinacanthus nutans</i> LEAF (Burm. F.) LINDAU EXTRACTS IN LEUKAEMIA-LYMPHOMA BEARING SPRAGUE DAWLEY RATS	89
4.1.	Introduction	89
4.2.	Materials and methods	90
4.2.1.	Plant preparation	90
4.2.2.	Plant extraction	90
4.2.3.	Animal management and routine	90
4.2.4.	Experimental design	90
4.2.5.	<i>N</i> -methyl- <i>N</i> -nitrosourea preparation and administration	91
4.2.6.	Preparation and administration of <i>M. citrifolia</i> fruit and <i>C. nutans</i> leaf extracts	91
4.2.7.	Preparation of chemotherapeutic drugs	93
4.2.8.	Haematology analysis	95
4.2.9.	Serum biochemistry analysis	95
4.2.10.	Histopathological analysis	95
4.2.10.1.	Lesions characteristics	95
4.2.11.	Statistical analysis	98
4.3.	Results	98
4.3.1.	Body weight	98
4.3.2.	Haematological and serum biochemical parameters	99
4.3.3.	Organ to body weight ratio	112
4.3.4.	Gross lesion	114
4.3.5.	Lymphoma lesion score	118
4.3.5.1.	Lymphoma lesion characteristics	120
4.4.	Discussion	150
4.5.	Conclusion	157
5	OXIDATIVE STRESS BIOMARKER, ANTIOXIDATIVE STRESS ENZYMES, VEGF AND COX-2 EXPRESSION IN LEUKAEMIA-LYMPHOMA RATS TREATED WITH MIXTURES OF <i>Morinda citrifolia</i> L. FRUIT AND <i>Clinacanthus nutans</i> (Burm. F) LINDAU leaf extracts	158
5.1.	Introduction	159
5.2.	Materials and methods	160
5.2.1.	Blood samplings	160
5.2.2.	Serum malondialdehyde (MDA) assay	160

5.2.3.	Serum glutathione peroxidase (GSH-Px)	160
5.2.4.	Serum superoxide dismutase (SOD)	160
5.2.5.	RNA extraction	161
5.2.4.1.	RNA extraction from blood	161
5.2.4.2.	RNA extraction from spleen and lymph nodes	162
5.2.4.3.	RNA sample quantification	162
5.2.4.4.	Conversion of RNA to DNA via reverse transcriptase	163
5.2.4.5.	Primer sequence	163
5.2.4.6.	Quantitative real time RT-PCR assay	166
5.2.4.7.	Gene expression analysis	166
5.2.5.	Immunohistochemistry examination	167
5.2.6.	Analysis of protein expression	168
5.2.6.1	Protein extraction from spleen and lymph nodes	168
5.2.6.2.	Bicinchonic (BCA) assay	168
5.2.6.3.	Protein lysis	169
5.2.6.4.	Western blot	169
5.2.7.	Statistical analysis	170
5.3.	Results	170
5.3.1.	Malondialdehyde (MDA) assay	170
5.3.2.	Superoxide dismutase (SOD) assay	171
5.3.3.	Glutathione peroxidase (GSG-Px) assay	172
5.3.4.	Quantitative real-time PCR assay of VEGF and COX-2 RNA transcription	173
5.3.5.	Immunohistochemistry evaluation	176
5.3.5.1.	VEGF and COX-2 immunostaining in lymph nodes	176
5.3.5.2.	VEGF and COX-2 immunostaining in spleen	180
5.3.6.	Western blot	188
5.4.	Discussion	189
5.5.	Conclusion	199
6	GENERAL DISCUSSION, CONCLUSION AND FUTURE STUDIES	200
6.1.	General Discussion	200
6.2.	Conclusion	201
6.3.	Recommendation and Future Studies	202
	REFERENCES	203
	APPENDICES	235
	BIODATA OF STUDENT	251
	LIST OF PUBLICATIONS	252

LIST OF TABLES

Table	Page
2.1 Parts of <i>M. citrifolia</i> used in Polynesian medicinal practice	6
2.2 Bioactive compounds in <i>M. citrifolia</i>	7
2.3 Benefits of <i>C. nutans</i> leaves	9
2.4 Types and description of VEGF family	17
2.5 Subspecies and description of glutathione peroxidase family	20
2.6 Forms and description of superoxide dismutase family	21
3.1 Treatment group in the acute oral toxicity study of <i>M. citrifolia</i> fruit extract in male Sprague Dawley rats	28
3.2 Treatment group in the subacute and subchronic oral toxicity studies of <i>M. citrifolia</i> fruit and <i>C. nutans</i> leaf extract in male Sprague Dawley rats	28
3.3 Scoring system used for histopathological evaluation of liver	31
3.4 Scoring system used for histopathological evaluation of kidney	32
3.5 Haematological parameters, icterus index and plasma protein values of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	34
3.6 Total white blood cell (WBC) and WBC differential counts of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	34
3.7 Serum biochemical parameters of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	35
3.8 Ratio of organ to relative body weight (%) of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	36
3.9 Number of male Sprague Dawley (SD) rats with hepatic and renal lesion in the 14-day (acute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	38

3.10	Scores of liver and kidney lesions of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	39
3.11	Results of Mann-Whitney U test for comparison between groups for lesions in the liver and kidneys of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	40
3.12	Haematological parameters, icterus index and plasma protein of male Sprague Dawley (SD) rats in a repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	42
3.13	Total white blood cell (WBC) and WBC differential counts of male Sprague Dawley (SD) rats in a repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> Fruit	42
3.14	Serum biochemical parameters of male Sprague Dawley (SD) rats in a repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	43
3.15	The organ relative weights of male Sprague Dawley (SD) rats in a repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	44
3.16	Number of male Sprague Dawley (SD) rats with hepatic and renal lesions in the repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	46
3.17	Scores of liver and kidneys lesions of male Sprague Dawley (SD) rats in the repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	47
3.18	Results of Mann-Whitney U test for comparison between groups of the toxicity lesions in liver and kidneys of male Sprague Dawley (SD) rats in the repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	48
3.19	Haematological parameters, icterus index and plasma protein values of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	51
3.20	Total white blood cell (WBC) and WBC differential count of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	51

3.21	Serum biochemical parameters of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	52
3.22	The organ relative weights of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	53
3.23	Number of male Sprague Dawley (SD) rats with hepatic and renal lesions in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	55
3.24	Scores of liver and kidneys lesions of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	56
3.25	Results of Mann-Whitney U test for comparison between groups of the toxicity lesions in liver and kidneys of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>M. citrifolia</i> fruit	57
3.26	Haematological parameters, icterus index and plasma protein values of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	59
3.27	Total white blood cell (WBC) and WBC differential count of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	59
3.28	Serum biochemical parameter of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	60
3.29	Ratio organ to relative body weight (%) of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	61
3.30	Number of male Sprague Dawley (SD) rats with hepatic and renal lesions in the 14-day (acute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	63
3.31	Scores of liver and kidneys lesions of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf.	64
3.32	Results of Mann-Whitney U test for comparison between groups of the toxicity lesions in liver and kidneys of male Sprague Dawley (SD) rats in the 14-day (acute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	65

3.33	Haematological parameters, icterus index and plasma protein values of male Sprague Dawley (SD) rats in repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	67
3.34	Total white blood cell (WBC) and WBC differential count of male Sprague Dawley (SD) rats in a repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	67
3.35	Serum biochemical parameter of male Sprague Dawley (SD) rats in the repeated dose 28-day (subacute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	68
3.36	Ratio of organ to relative body weight (%) of male Sprague Dawley (SD) rats in a repeated day (subacute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	69
3.37	Number of male Sprague Dawley (SD) rats with hepatic and renal lesions in therepeated 28-day (subacute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	71
3.38	Scores of the liver and kidneys lesions of male Sprague Dawley (SD) rats in the repeated dose 28-day (acute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	72
3.39	Results of Mann-Whitney U test for comparison between groups of the toxicity lesions in liver and kidney of male Sprague Dawley (SD) rats in the repeated day 28-day (subacute) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	73
3.40	Haematological parameters, icterus index and plasma protein values of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	75
3.41	Total white blood cell (WBC) and WBC differential count of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	75
3.42	Serum biochemical parameter of male Sprague Dawley (SD) rats in the 90-day (subchronic) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	75
3.43	Ratio of organ relative body weight (%) of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	77
3.44	Number of male Sprague Dawley (SD) rats with hepatic and renal lesions in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	79

3.45	Scores of liver and kidneys of male Sprague Dawley (SD) rats in the repeated 90-day (subchronic) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	80
3.46	Results of Mann-Whitney U test for comparison between groups of the toxicity lesions in liver and kidney of male Sprague Dawley (SD) rats in the repeated dose 90-day (subchronic) oral toxicity study of ethanol extract of <i>C. nutans</i> leaf	81
4.1	Experimental design of the antitumour effects of mixtures of <i>M. citrifolia</i> fruit and <i>C. nutans</i> leaf extracts in leukaemia-lymphoma Sprague Dawley rats	91
4.2	Scoring system for lymphoma lesion in the liver, kidneys, heart and Lungs	96
4.3	Scoring system for lymphoma lesion in the spleen	97
4.4	Scoring system for lymphoma lesion in the lymph nodes	97
4.5	Haematological parameters, icterus index and plasma protein values of male Sprague Dawley rats at week 18 in the leukaemia-lymphoma study	100
4.6	Haematological parameters, icterus index and plasma protein values of male Sprague Dawley rats at week 24 (end of study period) in leukaemia-lymphoma study	101
4.7	Leukogram of male Sprague Dawley rats at week 18 in the leukaemia-lymphoma study	102
4.8	Leukogram of male Sprague Dawley rats at week 24 in the leukaemia-lymphoma study	103
4.9	Blats cells differential count in blood smear at week 18 and week 24 in the leukaemia-lymphoma study	104
4.10	Percentage of leukaemic rats incidence in MNU-induced group	108
4.11	Serum biochemical parameters of male Sprague Dawley rats at week 18 in the leukaemia-lymphoma study	110
4.12	Serum biochemical parameters of male Sprague Dawley rats at week 24 in the leukaemia-lymphoma study	111
4.13	Organ to body weight ratio of male Sprague Dawley rats in the leukaemia-lymphoma study	113
4.14	Number of Sprague Dawley rats gross lesions in the leukaemia-lymphoma study (n =8 in each group)	117

4.15	Percentage of nodal and organ lymphoma in the male Sprague Dawley rat in the leukaemia-lymphoma study	119
4.16	Overall severity of lymphoma lesion in male Sprague Dawley of the leukaemia-lymphoma study (n=8 in each group)	138
4.17	Lesion scoring results of lymphoma lesions in the liver, lung, heart and kidneys of all groups	140
4.18	Lesion scoring results of lymphoma lesions in the spleen and lymph nodes of all groups	141
4.19	Comparison of lymphoma lesion scores in the liver, heart, kidneys and lungs of all groups using Mann-Whitney test	142
4.20	Comparison of lymphoma lesion scores in the spleen and lymph nodes Of all groups using Mann-Whitney test	144
4.21	Lesion scoring results of fibrous tissue formation in the liver and spleen of all groups	146
4.22	Comparison of the scores of fibrous tissue formation in the liver and spleen of all groups using Mann-Whitney test	147
4.23	Lesion scoring results of testicular necrosis in all groups	148
4.24	Comparison on the scores of the testicular necrosis in all groups using Mann-Whitney test	149
5.1	Forward and reverse sequences of target genes and house-keeping genes	164
5.2	Volumes of components for standard curve setup	165
5.3	Amplification protocol for each primer	165
5.4	Summary of VEGF and COX-2 expression in the lymph nodes of all groups.	184
5.5	VEGF and COX-2 expression in the lymph nodes between the groups	185
5.6	Summary of VEGF and COX-2 expression in the spleen of all groups	186
5.7	VEGF and COX-2 expression in the spleen between the groups	187

LIST OF FIGURES

Figure	Page
2.1 <i>Morinda citrifolia</i> tree (A) and fruit (B)	5
2.2 <i>Clinacanthus nutans</i> plant	10
2.3 Age specific cancer incidences per 100,000 population by sex, Malaysia 2007	13
3.1 The body weights (Mean±SEM) of male Sprague Dawley (SD) in 2 weeks experimental period	33
3.2 The body weights (Mean±SEM) of male Sprague Dawley (SD) in 4 weeks experimental period	41
3.3 The body weights (Mean±SEM) of male Sprague Dawley (SD) in 12 weeks experimental period	49
3.4 The body weights (Mean±SEM) of male Sprague Dawley (SD) in 2 weeks experimental period	58
3.5 The body weights (Mean±SEM) of male Sprague Dawley (SD) in 4 weeks experimental period	66
3.6 The body weights (Mean±SEM) of male Sprague Dawley (SD) in 12 weeks experimental period	74
3.7 Rat's liver sections at the end of experimental period (day 90)	82
3.8 Rat's kidney sections at the end of experimental period (day 90)	83
4.1 Timeline of herbal mixture administration	92
4.2 Schedule of the CHOP regimen administrations	94
4.3 The body weights (Mean±SEM) of male Sprague Dawley (SD) in 24 weeks experimental period	98
4.4 Photomicrograph from the blood smear of rat in the control group (Wright stain, x200)	105
4.5 Photomicrograph from the blood smear of rat in group C showing leukocytosis with the presence of large leukaemic cells (arrows) (Wright stain, x200)	106

4.6	Gross lesions of treated and untreated leukaemia lymphoma rats at the end of experimental period	115
4.7	Normal heart histology and mild cardiac lymphoma at week 24 of the study period	120
4.8	Normal kidney histology and moderate renal lymphoma at week 24 of the study period	121
4.9	Severe renal lymphoma at week 24 of the study period	122
4.10	Normal liver histology and mild hepatic lymphoma at week 24 of the study period	123
4.11	Moderate and severe hepatic lymphoma at week 24 of the study period	124
4.12	Normal lung histology and moderate pulmonary lymphoma at week 24 of the study period	125
4.13	Severe pulmonary lymphoma at week 24 of the study period	126
4.14	Normal histology of lymph node in one of the group A rats at week 24 of study the period	127
4.15	Mild and moderate nodal lymphoma lesions in rats at week 24 of the study period	128
4.16	Moderate nodal lymphoma lesion in a rat at week 24 of the study period	129
4.17	Moderate and severe nodal lymphoma lesions in rats at week 24 of the study period	130
4.18	Normal histology of spleen in one of the group A rats at week 24 of study period	131
4.19	Mild lymphoma lesion in a rat at week 24 of the study period	132
4.20	Severe lymphoma lesion in a rat at week 24 of the study period	133
4.21	Mild and moderate splenic lymphoma lesion in a rat at week 24 of study period	134
4.22	Normal histology and mild splenic lymphoma lesion of subcapsular sinuses of lymph node in rats at week 24 of study period	135
4.23	Moderate and severe fibrous tissue in the liver and spleen of rats in group G at week 24 of study period	136

4.24	Normal and necrosis of testis tissues in rats of group A and G at week 24 of study period	137
5.1	The serum MDA levels of rats at weeks 18 and 24 in the leukaemia-lymphoma study	171
5.2	The serum SOD levels of rats at weeks 18 and 24 in the leukaemia-lymphoma study	172
5.3	The serum GSH-Px levels of rats at week 18 and 24 in the leukaemia-lymphoma study	173
5.4	The VEGF RNA transcription levels of rats in the leukaemia-lymphoma study	174
5.5	The COX-2 RNA transcription levels of leukaemia-lymphoma study	175
5.6	Immunohistochemical staining of lymph node of control rat at week 24 of the experiment	159
5.7	Immunohistochemical staining of lymph node of treated rat at week 24 of the experiment	176
5.8	Immunohistochemical staining of lymph node VEGF of treated rat at week 24 of the experiment	178
5.9	Immunohistochemical staining of lymph node COX-2 of treated rat at week 24 of the experiment	179
5.10	Immunohistochemical staining of spleen of control rat at week 24 of the experiment	180
5.11	Immunohistochemical staining of spleen of treated rat at week 24 of the experiment	181
5.12	Immunohistochemical staining of spleen of treated rat at week 24 of the experiment	182
5.13	Western blot analysis of VEGF protein expression in lymph nodes	188
5.14	Western blot analysis of COX-2 protein expression in lymph nodes	189

LIST OF ABBRREVIATIONS

AMTREC	Animal Metabolism, Toxicology and Reproductive Centre
COX-2	Cylooxygenase-2
CHOP	Cyclophosphamide, hydroxydaunorubicin, oncovin, prednisone
CBC	Complete blood count
GPx	Glutathione Peroxidase
HL	Hodgkin's lymphoma
MDA	Malondialdehyde
MNU	<i>N</i> -methyl- <i>N</i> -nitrosourea
NHL	Non Hodgkin's lymphoma
SD	Sprague Dawley
SOD	Superoxide dismutase
VEGF	Vascular endothelial growth factor

CHAPTER 1

INTRODUCTION

Cancer or malignant neoplasm is a group of diseases involving abnormal cells growth with the potential to invade or spread to other parts of the body. It is one of the major health problems in many parts of the world including Malaysia; it is also the second leading cause of death in the United States after heart disease and will become the leading cause of death in the next few years (Siegel *et al.*, 2015). According to the National Cancer for Health (2015), 1,658,370 new cancer cases and 589,430 cancer deaths were estimated to occur in the United States in 2015. In Malaysia, cancer cases increased from 32,000 to 38,000 cases between 2008 and 2012 (International Agency of Research on Cancer) [IARC], 2012), and the top leading cancers are breast cancer (18.1%), head and neck cancer (13.2%), colorectal cancer (12.3%), tracheal, bronchial and lung cancers (10.2%), and cervix cancer (4.6%) (National Cancer Registry [NCR], 2012). Leukaemia and lymphoma are also among the top leading cancers in the United States (US). Approximately in every 3 minutes, one person in the US is diagnosed with blood cancer (Leukemia & Lymphoma Society, 2016).

Leukaemia is divided into four groups known as chronic lymphocytic leukaemia (CLL), acute lymphocytic leukaemia (ALL), chronic myeloid leukaemia (CML) and acute myeloid leukaemia (AML) (Bain, 2005) while lymphoma is divided into two main categories known as non-Hodgkin's lymphoma (NHL) and Hodgkin's lymphoma (HL). Leukaemia and lymphoma are haematopoietic cancers derived from haematopoietic and lymphatic systems (Mughal *et al.*, 2006). Leukaemia is a type of blood cancer caused by the overproduction of abnormal white blood cells (WBCs) in the bone marrow. Abnormal WBCs are unable to carry their normal functions to fight infections. Lymphoma, on the other hand, is cancer that starts in the cells of the lymphatic systems that include lymph nodes, thymus and spleen (Mughal *et al.*, 2006). Both cancers are diagnosed by the presence of abnormal cells in the blood circulation and organs, (Sathiya & Muthuchelian, 2009). In 2013, 7.35% people with leukaemia in the United States died because of the disease (Leukaemia-lymphoma Society, 2013). As the seventh most common cancer in Malaysia, 741 to 1162 leukaemia cases were reported between the year 2007 and 2009. According to the National Cancer Registry Report (2007), 322 were females while the remaining 419 were males. As for lymphoma, it was estimated that there were between 761,659 and 841,649 people living with lymphoma worldwide from 2013 to 2014 (Leukemia-Lymphoma Society of United States, 2014).

Cyclooxygenases (COXs) or commonly known as prostaglandin H synthase are members of the myeloperoxidase family. These enzymes are mostly expressed and located in the endoplasmic reticulum and nuclear membrane of cells (Sobolewski *et al.*, 2010). Among COXs, cyclooxygenase-2 (COX-2) is a rate limiting enzyme involved in inflammatory reactions and synthesis of angiogenic prostaglandins (PG), for example prostaglandin-endoperoxidase-2 (PGE2) (Fosslien *et al.*, 2001). Activation of PGE2 induces angiogenesis factors which include vascular endothelial

growth factor (VEGF) and inflammatory cytokines, particularly tumour necrosis factor (TNF). Overexpression of COX-2 is found in a variety of human malignancies including leukaemia (Egyudova *et al.*, 2011) and lymphoma (Wun *et al.*, 2004). The first evidence of a strong COX-2 expression was immunohistologically detected in the colorectal carcinoma (Wadell & Loughry, 1983). The immunohistological staining showed marked expression of COX-2 in tumour cells that were surrounded by numerous small vessels (Wadell & Loughry, 1983). Furthermore, expression of VEGF in the same area as of the tumour strongly suggests that overexpression of COX-2 induces angiogenesis (Fosslien *et al.*, 2001). It is shown that the expression of COX-2 is reduced by non-steroidal anti-inflammatory drugs (NSAIDs); however, prolonged use of the drugs could cause adverse gastrointestinal and cardiovascular side effects (Cha *et al.*, 2007). Previous *in vitro* study has shown that *M. citrifolia* fruit extract inhibited COX-2 expression in the *Helicobacter pylori*-infected glandular cells (AG) cells. However, there is no report on the effects on the expression of this enzyme on cancer cell lines or patients or animals with cancer. Similar to *M. citrifolia*, the inhibition effects of *C. nutans* on the COX-2 expression have never been studied for both *in vitro* and *in vivo* studies of inflammatory or neoplastic diseases.

Angiogenesis is the growth of new blood vessels from the pre-existing blood vessels. It plays important roles in the pathogenesis of malignant tumour (Prager *et al.*, 2012). Tumours require sustenance in the form of nutrients and oxygen as well as the ability to evacuate metabolic wastes and carbon dioxide as normal tissues. Tumour-associated neovasculature plays an important role in sustaining nutrients and oxygen in the microenvironment of the tumour. The development of vasculogenesis during embryogenesis involves the development of new endothelial cells and the sprouting of new vessels from the existing ones. During tumour progression, angiogenesis process is always activated, and it remains on. This process leads to the continuity of new blood vessels production that helps to sustain and expand the growth of the tumour (Hanahan & Folkman, 1996). Vascular endothelial growth factor (VEGF) is involved in the development and growth of new blood vessels during embryonic development, in homeostatic survival of endothelial cells, as well as in physiological and pathological process of cells (Hanahan & Weinberg, 2011). The expression of VEGF can be upregulated via hypoxia and signaling pathway (Ferrara, 2004). Ferrara (2004) reported that overexpression of VEGF could lead to carcinogenesis and tumorigenesis. Upregulation of VEGF was discovered in various types of cancer including leukaemia, lymphoma, ovarian cancer, gastrointestinal tract cancer and prostate cancer (Brown *et al.*, 1993; Bellamy *et al.*, 2001; Duncan *et al.*, 2008; Botelho *et al.*, 2010; Roorda *et al.*, 2012). Blocking of angiogenesis is one of the therapeutic ways in blocking cancer growth. No research has been done on *Morinda citrifolia* fruit and *Clinacanthus nutans* leaf in inhibiting angiogenesis in cancer. However, a study on stress-induced impairment cognitive function, showed low level of VEGF protein in the hypothalamus of restrained rats given *M. citrifolia* fruit juice (Muto *et al.*, 2010).

Most herbal plants are rich in beneficial compounds such as fiber, vitamins, flavonoids, alkaloids, sulphur compounds, phytosterols, carotenoids and organic acids, which contribute positively to human health (Mohd Shukri *et al.*, 2010). Herbal plants are also rich in antioxidant properties that play important roles in protecting

cells against oxidative stress-induced by diseases (Hertoq *et al.*, 1992; WHO, 2003). Herbal plants such as *Morinda citrifolia* and *Clinacanthus nutans* are among the two most popular local plants in Malaysia that are being studied for market potential in the medicinal and pharmaceutical industries. Traditionally, *M. citrifolia* was used to treat bowel disorder by Polynesians (Wang *et al.*, 2008) while *C. nutans* was traditionally used to treat Herpes simplex virus infection (Sakdarat *et al.*, 2006). *M. citrifolia* is already known for its *in vitro* anti-cancer activity. In the non-small human lung cancer (NCI-H23), human cervical cancer (HeLa and SiHa) and human colorectal cancer cell lines, it exhibited anti-proliferative effects (Kharis *et al.*, 2012; Nualsanit *et al.*, 2012; Gupta *et al.*, 2013). Meanwhile the *in vivo* anti-cancer activity of *M. citrifolia* was discovered in rats with Lewis lung carcinoma (Hirazumi *et al.*, 1994) and mammary tumour (Chew, *et al.*, 1999; Clafshenkel *et al.*, 2012). In contrast to *M. citrifolia*, the anti-tumour potential of *C. nutans* has not been extensively explored. Previous study by Yong *et al.* (2013) reported that *C. nutans* extract possessed anti-proliferative effects on selected cancer cell lines. The treatment of *C. nutans* chloroform leaf extract showed the highest anti-proliferative activity on human erythroleukaemia cell lines (K-562) and human Burkitt's lymphoma cell lines (Raji). Meanwhile, aqueous extract of *C. nutans* leaves showed the highest anti-proliferative effects on human cervical cancer (Hela) and K-562 cell lines. Previous mentioned studies investigated the effects of non-mixtures of *M. citrifolia* fruit and *C. nutans* leaf on cancer. Therefore, this study was conducted to determine the toxicity of *M. citrifolia* fruit and *C. nutans* leaf extracts and the antitumour effects of mixtures of *M. citrifolia* fruit and *C. nutans* leaf extracts in MNU-induced leukaemia-lymphoma Sprague Dawley rats.

Research problem:

New cases of leukaemia and lymphoma had increased 10.2 % from 2014 to 2016 (Leukaemia and Lymphoma Society, 2016). The use of chemotherapy drugs had increased the five-year survival rates of leukaemia and lymphoma patients by only 2 % (2004-2010 to 2005-2011) (National Cancer Institute, 2015). Side effects of CHOP regimen in treating leukaemia and lymphoma such as hepatic fibrosis (Lim *et al.*, 2010) and low sperm production in men (azoospermic) (Chan *et al.*, 2009) have been reported. Thus, this study was conducted to evaluate the effects of mixture of *M. citrifolia* fruit and *C. nutans* leaf in alleviating the side effects of chemotherapy drugs.

Justification of the study:

The study is conducted to seek alternative treatment for leukaemia-lymphoma.

The present study hypothesised that:

- 1) A single dose of 2000 mg/kg of ethanol extracts of *M. citrifolia* fruit and *C. nutans* leaf will induce no toxicity in male Sprague Dawley rats.
- 2) Once daily oral administration of ethanol extracts of *Morinda citrifolia* fruit and *Clinacanthus nutans* leaf at a dose of 75 mg/kg of body weight will induce no toxicity in male Sprague Dawley rats.
- 3) Once daily oral administration of ethanol extracts of *Morinda citrifolia* fruit and *Clinacanthus nutans* leaf at a dose of 125 mg/kg and 250 mg/kg of body weights will induce toxicity in male Sprague Dawley rats.
- 4) Mixtures of *M. citrifolia* fruit and *C. nutans* leaf extracts will enhance the anticancer activities and reduce leukaemia-lymphoma metastasised in male MNU-induced leukaemia-lymphoma rats.

The objectives of the present study are:

- 1) to determine the acute (14 days), subacute (28 days) and subchronic (90 days) toxicity studies of ethanol extracts of *M. citrifolia* fruit and *C. nutans* leaf in male Sprague Dawley rats;
- 2) to investigate and evaluate the antitumour effects of mixtures of *M. citrifolia* and *C. nutans* in MNU-induced leukaemia-lymphoma Sprague Dawley rats; and
- 3) to evaluate the oxidative stress biomarker, antioxidative stress enzymes, VEGF and COX-2 in MNU-induced leukaemia-lymphoma rats treated with mixture of *M. citrifolia* and *C. nutans* extracts.

REFERENCES

- Abalaka, S.E., Fatihu, M.Y., Ibrahim, N.D.G. and Ambali, S.F (2014). Hepatotoxicity of ethanol extract of *Adenium Obesum* stem bark in Wistar rats. *British Journal of Pharmaceutical Research*, 4(9), 1041-1052.
- Abd. Aziz, R. (2003). 'Turning Malaysia into Global Herbal Producer: A Personal Perspective', *Siri Syarahan Perdana Professor*, Universiti Teknologi Malaysia.
- Abdalla, S. I., Sanderson, I. R. and Fitzgerald, R. C. (2005). Effect of inflammation on cyclooxygenase (COX)-2 expression in benign and malignant esophageal cells. *Carcinogenesis*, 26(9), 1627-1633.
- Abdou, A.G., Aiad, M., Asaad, N., Abd El-Waheed, M. and Serag El-Dien, M. (2006). Immunohistochemical evaluation of vascular endothelial growth factor (VEGF) on colorectal carcinoma. *Journal of the Egyptian National Cancer Institute*, 18(4), 311-322.
- Abdul-Raheem, I.T. and Abdel-Ghany, A.A. (2009). Hesperidin alleviates doxorubicin-induced cardiotoxicity in rats. *Journal of Egypt National Cancer Institute*, 21(2), 175-184.
- Adedapo, A. A., Abatan, M. O., and Olorunsogo, O. O. (2004). Toxic effects of some plants in the genus *Euphorbia* on haematological and biochemical parameters of rats. *Veterinarski Arhiv*, 74(1), 53-62.
- Adekeye, A. O., Akintayo, C. O., Adewale, O., Sabiu, S. and Ogundele, O. M. (2014). Carbon tetrachloride (CCl₄)-induced hepatic damage in experimental Sprague Dawley rats: Antioxidant potential of *Xylopi aethiopica*. *Journal of Phytopharmacology*, 3(2), 118-123.
- Agnani, D., Camacho-Vanegas, O., Camacho, C., Lele, S., Odunsi, K., Cohen, S., Dottina, P. and Martignetti, J.A. (2011). Decreased levels of serum glutathione peroxidase 3 are associated with papillary serous ovarian cancer and disease progression. *Journal of Ovarian Research*, 4(18), doi:10.1186/1757-2215-4-18.
- Ahmad, M., Lone, A. R., Kuchai, S. U., Zargar, A. H., Aziz, S. A., Bhat, G. M., Banday, M.A., Jelani, S. and Bhat, J. R. (2008). Post-treatment testicular activity in lymphoma patients. *Indian Journal of Medical and Paediatric Oncology*, 29(1), 8-15.
- Al-Dissi, A. N., Haines, D. M., Singh, B. and Kidney, B. A. (2010). Immunohistochemical expression of vascular endothelial growth factor and vascular endothelial growth factor receptor-2 in canine simple mammary gland adenocarcinomas. *Canadian Veterinary Journal*, 51(10), 1109-1114.

- Amin, H. M., Yang, Y., Shen, Y., Estey, E. H., Giles, F. J., Pierce, S. A., Kantarjian, M., O'Brien, S.M., Jilani, I and Albitar, M. (2005). Having a higher blast percentage in circulation than bone marrow: clinical implications in myelodysplastic syndrome and acute lymphoid and myeloid leukemias. *Leukemia*, 19(9), 1567-1572.
- Amresh, G.R., Singh, P.N. and Rao, C.V. (2008). Toxicological screening of traditional medicine Laghupatha (*Cissampelos pareira*) in experimental animals. *Journal of Ethnopharmacology*, 116 (3), 454–460.
- Anitha, T. and Mohandass, S. (2006). Anti-oxidant activity of *Morinda citrifolia* on lymphoma-bearing mice. *Ancient Science of Life*, 26(1-2), 85-88.
- Arruda, V.R., Salles, T.C., Costa, F.F. and Saad, S.T. (1996). Glutathione peroxidase, reduced glutathione, superoxide dismutase and catalase in red cells of patients with hairy cell leukemia. *Neoplasma*, 43(2), 99-102.
- Arsad, S.S., Esa, N.M. and Hazilawati, H. (2014) Histopathologic changes in liver and kidneys tissue from male Sprague Dawley rats treated with *Rhaphidophora decursiva* (Roxb.) shoot extract. *Journal of Cytology and Histology*, 4(1), doi:10.4172/2157-7099.S4-001.
- Arthur, J.R. (2000).The glutathione peroxidases. *Cellular and Molecular Life Sciences*. 57(13-14), 1825-1835.
- Ashafa, A.O.T., Orekoya, L.O. and Yakubu, M.T. (2012). Toxicity profile ethanolic extract of *Azadirachta indica* stem bark in male Wistar rats. *Asian Pacific Journal of Tropical Biomedic*, 2(10), 811-817.
- Asproni, P., Vignoli, M., Cancedda, S., Millanta, F., Terragni, R. and Poli, A. (2014). Immunohistochemical expression of cyclooxygenase-2 in normal, hyperplastic and neoplastic canine lymphoid tissues. *Journal of Comparative Pathology*, 151(1), 35-41.
- Asyura, S.N.N., Hamzah, H., Shaari, R.M., Sithambaram, S., Mustapha, N.M. (2016). Blood profiles and histopathological changes of liver and kidney tissues from male Sprague Dawley rats treated with extracts of *Clinacanthus nutans*. *Journal Clinical of Toxicology*, 6(6), DOI: 10.4172/2161-0495.1000329
- Ayala, A., Muñoz, M. F. and Argüelles, S. (2014). Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxidative Medicine and Cellular Longevity*, dx.doi.org/10.1155/2014/360438.
- Baccarani, M., Dreyling, M. and ESMO Guidelines Working Group. (2010). Chronic myeloid leukaemia: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology*, 21(5), 165-167.

- Badcock, N.R., Zoanetti, G.D. and Martin, E.S. (1997). Nonchromatographic assay for malondialdehyde-thiobarbituric acid adduct with HPLC equivalence. *Clinical Chemistry*, 43(9), 1655-1657.
- Bae, S. H., Jung, E. S., Park, Y. M., Kim, B. S., Kim, B. K., Kim, D. G. and Ryu, W. S. (2001). Expression of cyclooxygenase-2 (COX-2) in hepatocellular carcinoma and growth inhibition of hepatoma cell lines by a COX-2 inhibitor. *Clinical Cancer Research*, 7(5), 1410-1418.
- Bahadur, G., Ozturk, O., Muneer, A., Wafa, R., Ashraf, A., Jaman, N. and Ralph, D. J. (2005). Semen quality before and after gonadotoxic treatment. *Human Reproduction*, 20(3), 774-781.
- Baiju, N., Mathews, P.P., Skariah, K.S. and Ambika, K. (2012). *Morinda citrifolia*-Noni: A brief review of its anti-cancer activity. *Dentistry Reviews*, 1(3), 104-107.
- Bain, B.J. (2005). Diagnosis from the blood smear. *The New England Journal of Medicine*, 353, 498 – 507.
- Baker, R.D., Baker, S.S., LaRosa, K., Whitney, C. and Newburger, P.E. (1993). Selenium regulation of glutathione peroxidase in human hepatoma cell line Hep3B. *Archives of Biochemistry and Biophysics*. 304(1), 53-57.
- Balkwill, F. and Mantovano, A. (2001). Inflammation and cancer. Back to Virchow? *The Lancet*, 357, 539-545.
- Baskol, G., Demir, H., Baskol, M., Kilic, E., Ates, F., Karakukcu, C. and Ustdal, M. (2006). Investigation of protein oxidation and lipid peroxidation in patients with rheumatoid arthritis. *Cell Biochemistry and Function*, 24(4), 307-312.
- Bataller, R. and Brenner, D. A. (2005). Liver fibrosis. *Journal of Clinical Investigation*, 115(2), 209-218.
- Bautz, F., Rafii, S., Kanz, L. and Mohle, R. (2000). Expression and secretion of vascular endothelial growth factor-A by cytokine-stimulated hematopoietic progenitor cells. Possible role in the hematopoietic microenvironment. *Experimental Hematology*, 28(6), 700-706.
- Bellamy, W.T., Richter, L., Sirjani, D., Roxas, C., Glinsmann-Gibson, B., Frutiger, Y., Grogan, T.M. and List, A.F. (2001). Vascular endothelial cell growth factor is an autocrine promoter of abnormal localized immature myeloid precursors and leukemia progenitor formation in myelodysplastic syndromes. *Blood*, 97(5), 1427-1434.
- Belson, M., Kingsly, B. and Holmes, A. (2006). Risk factors for acute leukaemia in children: A review. *Environmental Health Perspectives*, 115 (1), 138-145.

- Bennett, J. M., Catovsky, D., Daniel, M. T., Flandrin, G., Galton, D. A. G., Gralnick, H. R., & Sultan, C. (1976). Proposals for the Classification of the Acute Leukaemias French-American-British (FAB) Co-operative Group. *British Journal of Haematology*, 33(4), 451-458.
- Bennett, J. M., Catovsky, D., Daniel, M. T., Flandrin, G., Galton, D. A., Gralnick, H. R., and Sultan, C. (1985). Proposed revised criteria for the classification of acute myeloid leukemia: a report of the French-American-British Cooperative Group. *Annals of Internal Medicine*, 103(4), 620-625.
- Bergendahl, V., Glaser, B.T. and Burgess, R.R. (2003). A fast western blot procedure improved for quantitative analysis by direct fluorescence labelling primary antibodies. *Journal of Immunological Methods*, 277(1-2), 117-125.
- Birgegård, G. (2008). Managing anemia in lymphoma and multiple myeloma. *Therapeutics and Clinical Risk Management*, 4(2), 527-539.
- Bocca, C., Ievollela, M., Autelli, R., Motta, M., Mosso, L., Torchio, B., Bozzo, F., Canitto S., Paternostro, C., Collombato, S., Parola, M. and Miglietta, A. (2014). Expression of COX-2 in human breast cancer cells as a critical determine of epithelial-to-mesenchymal transition and invasiveness. *Expert Opinions on Therapeutic Targets*, 18(2), 121-135.
- Borre, M., Nerstrøm, B. and Overgaard, J. (2000). Association between immunohistochemical expression of vascular endothelial growth factor (VEGF), VEGF-expressing neuroendocrine-differentiated tumor cells, and outcome in prostate cancer patients subjected to watchful waiting. *Clinical Cancer Research*, 6(5), 1882-1890.
- Botelho, F., Pina, F. and Lunet, N. (2010). VEGF and prostate cancer: a systematic review. *European Journal of Cancer Prevention*, 19(5), 385-392.
- Brancaccio, P., Lippi, G. and Maffulli, N. (2010). Biochemical markers of muscular damage. *Clinical Chemistry and Laboratory Medicine*, 48(6), 757-767.
- Brown, A. C. (2012). Anticancer activity of *Morinda citrifolia* (Noni) fruit: a review. *Phytotherapy Research*, 26(10), 1427-1440.
- Brown, F.L., Berse, B., Jackman, R.W., Tognazzi, K., Manseau, E.J., Senger, D.R. and Dvorak, H.F. (1993). Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in adenocarcinomas of the gastrointestinal tract. *Cancer Research*, 53(19), 4727-4735.
- Burkill, I. H. (1966). A Dictionary of Economic Products of the Malay Peninsular, volume 1, Ministry of Agriculture and Cooperative Malaysia, Kuala Lumpur, Malaysia. Vol. I, A-H: 1-1220; vol. II, I-Z: 1221-2402.
- Burrell, K. and Zadeh, G. (2012). Molecular Mechanisms of Tumor Angiogenesis, Tumor Angiogenesis, Dr. Sophia Ran (Ed.), ISBN: 978-953-51-0009-6.

- Byrd, J. C. (2015). Introduction to a series of reviews on chronic lymphocytic leukemia. *Blood*, 126(4), 427-427.
- Cairns, D., Michalitsi, E., Jenkins, T.C. and Mackay, S.P. (2002). Molecular modelling and cytotoxicity of substituted anthraquinones as inhibitors of human telomerase. *Bioorganic and Medicinal Chemistry*, 10(3), 803-807.
- Ceccarelli, C., Piazzzi, G., Paterini, P., Pantaleo, M. A., Taffurelli, M., Santini, D., Martinelli, G.N. and Biasco, G. (2005). Concurrent EGFr and Cox-2 expression in colorectal cancer: proliferation impact and tumour spreading. *Annals of Oncology*, 16(4), 74-79.
- Cha, Y. I., and DuBois, R. N. (2007). NSAIDs and cancer prevention: targets downstream of COX-2. *Annual Review of Medicine*, 58, 239-252.
- Chan, P. T. (2009). Fertility after cancer in men. *Canadian Urological Association Journal*, 3(3), 223-224.
- Chan-Blanco, Y., Fabrice, V., Ana, M.P., Max, R., Jean-Marc, B. and Pierre, B. (2006). The noni fruit (*Morinda citrifolia* L.): A review of agricultural research, nutritional and therapeutic properties. *Journal of Food Composition and Analysis*, 19(6), 645-654.
- Chang, Y.C., Hsu, J.D., Lin, W.L., Lee, Y.J. and Wang, C.J. High incidence of acute promyelocytic leukemia specifically induced by *N*-nitroso-*N*-methylurea (NMU) in Sprague Dawley rats. (2012). *Achieves of Toxicology*. 86(2), 315-327.
- Chen, J., Rider, D.A. and Ruan, R. (2006). Identification of valid housekeeping genes and antioxidant enzyme gene expression change in the aging rat liver. *Journal of Gerontology: Biological Sciences*, 21A, 20 – 27.
- Cheng, J. and Haas, M. (1990). Frequent mutations in the p53 tumor suppressor gene in human leukaemia T-cell lines. *Molecular and Cell Biology*, 10 (10), 5502-5509.
- Cheson, B. D. (2002). CHOP plus rituximab—balancing facts and opinion. *New England Journal of Medicine*, 346(4), 280-282.
- Chew, B.P., Park, J.S., Wong, M.W. and Wong, T.S. (1999). A comparison of the anticancer activities of dietary beta-carotene, canthaxanthin, and astaxanthin in mice in vivo. *Anticancer Research*, 19(3A), 1849-1853.
- Chin, J.H., Akowuah, G.A., Sabu, M.C. and Khalivulla, S.I. (2014). Sub-acute (28 days) toxicity study of methanol leaves extract of *Cinacanthus nutans* in rats. *International Journal of Pharmacy*, 4(2): 61-69.

- Chole, R. H., Patil, R. N., Basak, A., Palandurkar, K. and Bhowate, R. (2010). Estimation of serum malondialdehyde in oral cancer and precancer and its association with healthy individuals, gender, alcohol, and tobacco abuse. *Journal of Cancer Research and Therapeutics*, 6(4), 487.
- Chung, J.C., Oh, M.J., Choi, S.H. and Bae, C.D. (2008). Proteomic analysis to identify biomarker proteins in pancreatic ductal adenocarcinoma. *ANZ Journal of Surgery*, 78(4), 245-251.
- Clafshenkel, W.P., King, T.L., Kotlarczyk, M.P., Cline, J.M., Foster, W.G., Davis, V.L. and Witt-Enderby, P.A. (2012). *Morinda citrifolia* (noni) juice augments mammary gland differentiation and reduces mammary tumour growth in mice expressing the unactivated c-erbB2 transgene. *Evidence-Based Complementary and Alternative Medicine*, doi:10.1155/2012/487423.
- Close, B., Banister, K., Baumans, V., Bernoth, E-M., Bromage, N., Bunyan, J., Erhardt, W., Flecknell, P., Gregory, N., Hackbarth, H., Morton, D. and Warwick, C. (1997). Recommendations for euthanasia of experimental animals: Part 1, working party report. *Laboratory Animals*, 30 (2), 293 – 316.
- Coiffier, B., Lepage, E., Briere, J., Herbrecht, R., Tilly, H., Bouabdallah, R., Morel, P., Van De Neste, E., Salles, G., Gaulard, P. and Reyes, F. (2002). CHOP chemotherapy plus rituximab compared with CHOP alone in elderly patients with diffuse large-B-cell lymphoma. *New England Journal of Medicine*, 346(4), 235-242.
- Coombs, C. C., Tavakkoli, M., and Tallman, M. S. (2015). Acute promyelocytic leukemia: where did we start, where are we now, and the future. *Blood Cancer Journal*, 5(4), e304.
- Coons, A.H., Creech, H.J. and Jones, R.N. (1941). Immunology properties of an antibody containing a fluorescent group. *Proceedings of the Society for Experimental Biology and Medicine*, 47 (3), 200-202.
- Corless, J. K. and Middleton, H. M. (1983). Normal liver function: A basis for understanding hepatic disease. *Archives of Internal Medicine*, 143(12), 2291-2294.
- Coussens, L.M. and Werb, Z., (2002). Inflammation and cancer. *Nature*, 420, 860-867.
- Da, Silva, Franchi, C.A., Bacchi, M.M., Padovani, C.R. and de, Camargo, J.L. (2003). Thymic lymphomas in Wistar rats exposed to *N*-methyl-*N*-nitrosourea (MNU). *Cancer Science*, 94 (3), 240-243.
- Dias, S., Hattori, K., Heissiq, B., Zhu, Z., Wu, Y., Witte, L., Hicklin, D.J., Tateno, M., Bohlen, M., Moore, M.A. and Rafii, S. (2001). Inhibition of both paracrine and autocrine VEGF/VEGFR-2 pathway is essential to induce long-term remission

of xenotransplanted human leukemias. *Proceedings of the International Academy of Sciences*. 98(19), 10857-10862.

David, M.M. (2000). Hematology of the rat (*Rattus Norvegicus*). In B.E. Feldman, J.G. Zinkl and N.C. Jain (Eds.), *Schalm's veterinary hematology* (5th ed.). (pp. 1210-1218). Baltimore, Maryland: Lippincott Williams and Wilkins.

Dennis-Skyes, C.A., Miller, W.J., McAleer, W.J (1985). A quantitative western blot method for protein measurement. *Journal of Biological Standardization*, 13 (2), 309-314.

Deschpande, S.S., Lalitha, V.S., Ingle, A.D., Raste, A.S., Gadre, S.G. and Maru, G.B. (1998). Subchronic oral toxicity of turmeric and ethanolic turmeric extract in female mice and rats. *Toxicology Letter*, 95(3): 183-195.

Dhar, S.K. and St. Clair, D.K. (2012). Manganese superoxide dismutase regulation and cancer. *Free Radical and Biology and Medicine*, 52, 2209-2222.

Domrongkitchaiporn, S., Sritara, P., Kitiyakara, C., Stitchantrakul, W., Krittaphol, V., Lolekha, P., Cheepudomwit, S. and Yipintsoi, T. (2005). Risk factors for development of decreased kidney function in a Southeast Asian population: a 12-year cohort study. *Journal American Society of Nephrology*, 16, 791-799.

Drumond, A. L., Weng, C. C., Wang, G., Chiarini-Garcia, H., Eras-Garcia, L. and Meistrich, M. L. (2011). Effects of multiple doses of cyclophosphamide on mouse testes: Accessing the germ cells lost, and the functional damage of stem cells. *Reproductive Toxicology*, 32(4), 395-406.

Dubois, R. N., Abramson, S. B., Crofford, L., Gupta, R. A., Simon, L. S., Van De Putte, L. B. and Lipsky, P. E. (1998). Cyclooxygenase in biology and disease. *The FASEB Journal*, 12(12), 1063-1073.

Duffy, A.M., Boucheir-Hayes, D.J. and Harmey, J.H. (2000). Vascular endothelial growth factor (VEGF) and its role in non-endothelial cells: Autocrine signaling by VEGF. *Madame Curie Bioscience Database [Internet]*. Austin (TX). Landes Bioscience, 94(3), 240-243.

Duncan, T.J., Al-Attar, A., Rolland, P., Scott, I.V., Deen, S., Liu, D.T.Y., Spendlove, I. and Durrant, L.G. (2008). Vascular endothelial growth factor expression in ovarian cancer: A model for targeted of novel therapies?, *Clinical Cancer Research*, 14, 3030.

Duse, A.O., Ceausu, R.A., Mezei, T., Cimpean, A.M., Gaje, P., Ionita, H., Jung, I. and Raica, M. (2013). Expression and possible significance of vascular endothelial growth factor in non-Hodgkin lymphoma. *Archives of Biological Science Belgrade*, 65(2), 487-491.

Egyudova, K., Siltanen, A., Kankuri, E. and Bizik J. (2011). Leukemic cells modulate induction of COX-2 in human stromal fibroblast. *Neoplasma*, 58(6), 525-531.

- Eissa, L. A. and Esmaeel, M. I. (2008). Relevance of some serum biomarkers (E Cadherin, GAGs & MDA in patients with diffuse large B-cell lymphoma. *Pakistan Journal of Pharmaceutical Sciences*, 21(1), 29-35.
- Ellis, L. M. and Hicklin, D. J. (2008). VEGF-targeted therapy: mechanisms of anti-tumour activity. *Nature Reviews Cancer*, 8(8), 579-591.
- Espinoza, S.E., Guo, H., Fedarko, N., DeZern, A., Fried, L.P., Xue, Q-L., Leng, S., Beamer, B. and Walston, J.D. (2008). Glutathione peroxidase enzyme activity in aging. *The Journals of Gerontology. Series A: Biological Sciences and Medical Sciences*, 63(5), 505-509.
- Esterbauer, H., Cheeseman, K. H., Dianzani, M. U., Poli, G. and Slater, T. F. (1982). Separation and characterization of the aldehydic products of lipid peroxidation stimulated by ADP-Fe²⁺ in rat liver microsomes. *Biochemical Journal*, 208(5), 129-140.
- Faezah, A., Mohd, Azlan, S.Z., Noorasiah, S., Fadzilah, A.A.M. (2015). *Prosiding Persidangan Kebangsaan Ekonomi Malaysia*, 10(2), 227-238.
- Farell, D.J., Bower, L (2003). Fatal water intoxication. *Journal of Clinical Pathology*, 56(10), 803-804.
- Fasola, G., Fanin, R., Gherlinzoni F., Galeini, P., Taruscio, D., Frezza, G., Mazza, P., Pileri, S. and Baccarani, M. (1984). Serum LDH concentration in non-Hodgkin's lymphomas. Relationship to histologic type, tumour mass and presentation features. *Acta Haematology*, 74(2), 231-238.
- Fattman, C.L., Schaefer, L.M. and Oury, T.D. (2003). Extracellular superoxide dismutase in biology and medicine. *Free Radical Biology and Medicine*, 35(3), 236-256.
- Ferrara, N. (2004). Vascular endothelial growth factor: basics science and clinical progress. *Endocrine reviews*, 25(4), 581-611.
- Ferrara, N., Huock, K. and Jakeman, L. (1992). Molecular and biological properties of the vascular endothelial growth factor family of proteins. *Endocrine Reviews*, 13 (1), 18-32.
- Feugier, P., Van Hoof, A., Sebban, C., Solal-Celigny, P., Bouabdallah, R., Ferme, C., Christian, B., Lepage, G., Tilly, H., Morschhauser, P., Gaulard, P., Salles, G., Bosly, A., Gisselbrecht, C., Reyes, F., Coiffier, B. (2005). Long-term results of the R-CHOP study in the treatment of elderly patients with diffuse large B-cell lymphoma: a study by the Groupe d'Etude des Lymphomes de l'Adulte. *Journal of Clinical Oncology*, 23(18), 4117-4126.
- Fiedler, W., Graeven, U., Ergun, S., Verago, S., Kilic, N., Stockschrader, M. and Hossfeld, D.K. (1997). Vascular endothelial growth factor, a possible

- paracrine growth factor in human acute myeloid leukemia. *Blood*, 89(6), 1870-1875.
- Flora, G., Gupta, D., and Tiwari, A. (2012). Toxicity of lead: a review with recent updates. *Interdisciplinary Toxicology*, 5(2), 47-58.
- Folkman, J. (1971). Tumor angiogenesis; therapeutics implications. *The New England Journal of Medicine*, 285, 1182-1186.
- Franceschino, A., Tornaghi, L., Benemacher, V., Assouline, S. and Gambacorti-Passerini, C. (2008). Alterations in creatine kinase, phosphate and lipid values in patients with chronic myeloid leukemia during treatment with imatinib. *Haematologica*, 93(2), 317-318.
- Freedman, M.D., Stewart, P., Kleinerman, R.A., Wacholder, S., Hatcher, E.E. and Tarone, R.E. (2001). Household solvent exposures and childhood acute lymphoblastic leukaemia. *American Journal of Public Health*, 91, 564-567.
- Fridovich, I. (1974). Superoxide dismutases. *Advances in Enzymology and Related areas of Molecular Biology*, 41, 35-97.
- Furusawa, E. (2002). Anti-cancer activity of noni fruit juice against tumors in mice. In *Proceedings of the 2002 Hawai'i Noni Conference. University of Hawaii at Manoa, College of Tropical Agriculture and Human Resources*, 9(8), 23-24.
- Galarraga, B., Sinclair, D., Fahie-Wilson, M. N., McCrae, F. C., Hull, R. G. and Ledingham, J. M. (2003). A rare but important cause for a raised serum creatinine kinase concentration: two case reports and a literature review. *Rheumatology*, 42(1), 186-188.
- Garibyan, L. and Avashia, N. (2013). Polymerase chain reaction. *Journal of Investigative Dermatology*, 133 (3), doi:10.1038/jid.2013.1.
- Ghadyalpatil, N. S., Chandrasekar, R., Snehalatha, D. and Reddy, B. M. (2011). A case of primary ovarian lymphoma with autoimmune hemolytic anemia achieving complete response with Rituximab-based combination chemotherapy. *Indian Journal of Medical and Paediatric Oncology: Official Journal of Indian Society of Medical and Paediatric Oncology*, 32(4), 207-210.
- Ghalaut, V. S., Ghalaut, P. S. and Singh, S. (1999). Lipid peroxidation in leukaemia. *The Journal of the Association of Physicians of India*, 47(4), 403-405.
- Glass, D.C. (2003). Leukaemia risk associated with low-level benzene exposure. *Epidemiology*, 14, 569-577.
- Goldmann, E. (1908). The growth of malignant disease in man and the lower animals, with special reference to the vascular system. *Journal of the Royal Society of Medicine*, 1, 1-13.

- Gomez-Esquer, F., Agudo, D., Martinez-Arribas, F., Nunez-Villar, M. J. and Schneider, J. (2004). mRNA expression of the angiogenesis markers VEGF and CD105 (endoglin) in human breast cancer. *Anticancer Research*, 24(3A), 1581-1586.
- Gönenc, A., Ozkan, Y., Torun, M. and Simsek, B. (2001). Plasma malondialdehyde (MDA) levels in breast and lungs cancer patients. *Journal of Clinical Pharmacy and Therapeutics*, 26(2), 141-144.
- Goppelt-Strube, M. (1995). Regulation of prostaglandin endoperoxide synthase (cyclooxygenase) isozyme expression. *Prostaglandins Leukotrienes Essential Fatty Acids*, 52, 213-222.
- Gowda, S., Desai, P. B., Kulkarni, S. S., Hull, V. V., Math, A. A. and Vernekar, S. N. (2010). Markers of renal function tests. *North American Journal of Medical Sciences*, 2(4), 170.
- Greenhough, A., Smartt, H.J.M., Moore, A.E., Roberts, H.R., Williams, A.C., Paraskeva, C. and Kaidi, A. (2009). The COX-2/ PGE2 pathway: key roles in the hallmarks of cancer and adaptation to the tumor microenvironment. *Carcinogenesis*, 30(3), 377-386.
- Gregory, S.A. and Trumper, L. (2005). Chemotherapy dose intensity in non-Hodgkin's lymphoma: is dose intensity an emerging paradigm for better outcomes? *Annals of Oncology*, 16, 1413- 1424.
- Guenin, S., Mauriat, M., Pelloux, J., Wuytswinkel, O.V., Bellini, C. and Gutierrez, L. (2009). Normalization of qRT-PCR data: the necessity of adopting a systematic, experimental conditions-specific, validation of references. *Journal of Experimental Botany*. 60(2), 487-493.
- Güney, Y., Bilgihan, A., Ciftci, T.U., Cimen, F. and Coskun, O. (2004). Serum malondialdehyde levels and superoxide dismutase activities in pulmonary tuberculosis and lung cancers. *Meslek Yüksekokulu Dergisi*, 6(2), 33-38.
- Gupta, R.K. and Patel, A.K. (2013). Review- Do the health claims made for *Morinda citrifolia* (Noni) harmonize with current scientific knowledge and evaluation of its biological effects. *Asian Pacific Journal of Cancer Prevention*, 14 (8): 4495-4499.
- Hadija, H., Ayub, M.Y., Zaridah, H. and Normah, A. Acute and subchronic toxicity study of aqueous extract of *Morinda citrifolia* fruit in rat. (2003). *Journal of Tropical Agriculture and Food Science*, 31(1), 67-73.
- Hadija, H., Normah, A. and Ahmad Tarmizi, S. (2008). Thirteen-week toxicity study of mengkudu juice (*Morinda citrifolia*): Effect on blood analysis. *Journal of Tropical Agriculture and Food Science*, 36(2), 000-000.

- Hafez, E., and Sahar Issa, Y. (2015). Parenchymatous Toxicity of Tramadol: Histopathological and Biochemical Study. *Journal of Alcoholism and Drug Dependence*, 4(1), 3-5.
- Hanahan, D. and Folkman, J. (1996). Patterns and emerging mechanism of the angiogenic switch during tumorogenesis. *Cell*, 86, 352-364.
- Hanahan, D. and Weinberg, R.A. (2011). Hallmarks of cancer: the next generation. *Cell*, 144(5), 646-674.
- Harada, S., Hamabe, W., Kamiya, K., Satake, T., Yamamoto, J. and Tokuyama, S. (2009). Preventive effect of Morinda citrifolia fruit juice on neuronal damage induced by focal ischemia. *Biological and Pharmaceutical Bulletin*, 32(3), 405-409.
- Hasegawa, R., Futakuchi, M., Mizoquchi, Y., Yamauchi, T., Shirai, T., Ito, N. and Lijinsky, W. (1998). Studies of initiation and promotion of carcinogenesis by N-nitroso compounds. *Cancer Letters*. 123(2), 185-191.
- Hashino, Y., Ichijo, S., Osame, S and Takahashi, E. (1989). Studies on serum tocopherol, selenium levels and blood glutathione peroxidase activities in calves with white muscle disease. *Nihon Juigaku Zasshi*, 51(4), 741-748.
- Hazar, B., Ergin, M., Seyrek, E., Erdogan, S., Tuncer, I. and Hakverdi, S. (2004). Cyclooxygenase-2 (COX-2) expression in lymphomas. *Leukemia and Lymphoma*, 45(7), 1395-1399.
- Hazilawati, H., Abdullah, M., Hutheyfa, A.H., Rosly, S.M., Jasni, S., Noordin, M.M. and Shanmugavelu, S. (2009). High concentration of N-Methyl-Nnitrourea (MNU) induced intravascular hemolytic anaemia rather than leukaemia-lymphoma in Sprague Dawley rats. In Proceedings of the 1st Malaysian Association of Veterinary Pathology Conference Harbour View Hotel, Kuching, Sarawak: *Malaysian Association of Veterinary Pathology*, 95-99.
- Hazilawati, H., Farah Dina, P., Rosly, S.M., Shanmugavelu, S. and Noordin, M.M. (2010). Phyllanthus niruri reduces renal azotaemia in rats induced with chronic renal failure. *The Medical Journal of Malaysia*, 65, 132 – 134.
- He, R., Liu, B., Yang, R.C., Tobelem, G. and Han, Z.C. (2003). Inhibition of K562 leukaemia angiogenesis and growth by expression of antisense vascular endothelial growth factor (VEGF) sequence. *The Journal of Cancer Gene and Cellular Therapies*, 10, 879-886.
- Hermans, N., Cos, P., Maes, L., De Bruyne, T., Vanden Berghe, D., Vlietinck, A.J. and Pieters, L. (2007). Challenges and pitfalls in antioxidant research. *Current Medicinal Chemistry*, 14(4), 417- 430.
- Hertoq, M.G.I., Hollman, P.C.H. and Katan, M.B. (1992). Content of potentially anticarcinogenic flavanoids of 28 vegetables and 9 fruits commonly consumed in the Netherlands. *Journal of Agricultural and Food Chemistry*, 40, 3954-3962.

- Hirazumi, A., Furusawa, E., Chow, S.C., and Hokama, Y. (1994). Anticancer activity of *Morinda citrifolia* (noni) in intraperitoneally implanted Lewis lung carcinoma in syngeneic mice. *Proceedings of the Western Pharmacology Society*, 37, 145-146.
- Hoeben, A., Landuyt, B., Highley, M.S., Wildiers, H., Van, Osteroom, A.T. and De, Bruijn, E.A. (2004). Vascular endothelial factor and angiogenesis. *Pharmacological Reviews*. 56, 549-580.
- Hou, Z., Falcone, D. J., Subbaramaiah, K. and Dannenberg, A. J. (2011). Macrophages induce COX-2 expression in breast cancer cells: role of IL-1 β autoamplification. *Carcinogenesis*, 32(5), 695-702.
- Hu, Y., Rosen, D. G., Zhou, Y., Feng, L., Yang, G., Liu, J. and Huang, P. (2005). Mitochondrial manganese-superoxide dismutase expression in ovarian cancer role in cell proliferation and response to oxidative stress. *Journal of Biological Chemistry*, 280(47), 39485-39492.
- Huang, D., Guo, W., Gao, J., Chen, J., and Olatunji, J. O. (2015). *Clinacanthus nutans* (Burm. f.) Lindau Ethanol Extract Inhibits Hepatoma in Mice through Upregulation of the Immune Response. *Molecules*, 20(9), 17405-17428.
- Huang, H. L., Ko, C. H., Yan, Y. Y. and Wang, C. K. (2014). Antiadhesion and anti-inflammation effects of noni (*Morinda citrifolia*) fruit extracts on AGS cells during *Helicobacter pylori* infection. *Journal of Agricultural and Food Chemistry*, 62(11), 2374-2383.
- Huang, P., Feng, L., Oldham, E.A., Keating, M.J. and Plunkett, W. (2000). Superoxide dismutase as a target for the selective killing of cancer cells. *Nature*, 407(6802), 390-395.
- Huggins, C.B., Grand, L. and Ueda, N. Specific induction of erythroleukemia and myelogenous leukemia in Sprague-Dawley rats. (1982). *Proceedings of National Academy of Sciences USA* 79, 5144-5414.
- Hussain, T., Gupta, R. K., Sweet, K., Eswaran, B., Vijayakumar, M. and Rao, C. V. (2012). Nephroprotective activity of *Solanum xanthocarpum* fruit extract against gentamicin-induced nephrotoxicity and renal dysfunction in experimental rodents. *Asian Pacific Journal of Tropical Medicine*, 5(9), 686-691.
- Hutheyfa, A.H. (2010). Effects of *Morinda citrifolia* on N-methyl N-nitrosourea induced peripheral T cell non-Hodgkin's lymphoma in Sprague Dawley rats. Master's Thesis, Universiti Putra Malaysia, Malaysia.
- Hutheyfa, A.H., Hazilawati, H., Rosly, S.M., Jasni, S., Noordin, M.M. and Shanmugavelu, S. Histopathological features of peripheral T-cell lymphoma in Sprague-Dawley rats induced with N-methyl-N-nitrosourea. (2011). *Pertanika Journal of Tropical Agricultural Science* 34, 357-368.

- Ilmie, U.M., Jaafar, H., Mansor, S. M. and Abdullah, J. M. (2015). Subchronic Toxicity Study of Standardized Methanolic Extract of *Mitragyna Speciosa* Korth in Sprague-Dawley Rats. *Name: Frontiers in Neuroscience*, 9, 189-193.
- Ismail, Z., Halim, S.Z., Abdullah, N.R., Afzan, A., Abdul Rashid, B.A. and Jantan, I. (2014). Safety evaluation of oral toxicity of *Carica papaya* Linn. Leaves: A subchronic toxicity study in Sprague Dawley rats. *Evidence Based Complementary and Alternative Medicine*, 741470. doi: 10.1155/2014/741470.
- Itzkowitz, S.H. and Yio, X. (2004). Inflammation and cancer. IV. Colorectal cancer in bowel disease: the role of inflammation. *American Journal Physiology-Gastrointestinal and Liver Physiology*, 287, 7-17.
- Jaffe, E. S. (Ed.). (2001). Pathology and genetics of tumours of haematopoietic and lymphoid tissues. World Health Organisation Classification of Tumours. *Annals Oncology*, 13(3): 490-491.
- Jardim, B. V., Moschetta, M. G., Leonel, C., Gelaleti, G. B., Regiani, V. R., Ferreira, L. C., Lopes, J.R. and de Campos Zuccari, D. A. (2013). Glutathione and glutathione peroxidase expression in breast cancer: An immunohistochemical and molecular study. *Oncology reports*, 30(3), 1119-1128.
- Jebbink, J., Bai, X., Rogers, B.B., Dawson, D..B, Scheuermann, R.H. and Diomati-Saad R. (2003). Development of real-time PCR assay for quantitative detection of Epstein Barr Virus and Cytomegalovirus, comparison of TaqMan probes, and molecular beacons. *Journal Molecular Diagnosis*. 5(1), 15-20.
- Jr-Shiuan, J. M. P. Y., Wang, H. L. J., Lin, C. C. and Liang, J. Y. (2012). Effects of *Clinacanthus nutans* (Burm. f) Lindau leaf extracts on protection of plasmid DNA from riboflavin photoreaction. *MC-Transaction on Biotechnology*, 4(1), 5-8.
- Kaefer, C. M., and Milner, J. A. (2008). The role of herbs and spices in cancer prevention. *The Journal of Nutritional Biochemistry*, 19(6), 347-361.
- Kamal, M.S.A., Ghazali, A.R., Yahya, N.A., Wasiman, M.I. and Ismail, Z. (2012). Acute toxicity study of standardized *Mitragyna speciose* Korth aqueous extract in Sprague Dawley rats. *Journal of Plant Studies*, 1(2), 120-129.
- Kazem, A., El Sayed, K. and El Kerm, Y. (2014). Prognostic significance of COX-2 and β -catenin in colorectal carcinoma. *Alexandria Journal of Medicine*, 50(3), 211-220.
- Keeley, D. and Rees, J. (1997). New guidelines in asthma management. *British Medical Journal*, 314, 315-316.
- Kelaidi, C., Adès, L., and Fenaux, P. (2011). Treatment of Acute Promyelocytic Leukemia with High White Cell Blood Counts. *Mediterranean Journal of Hematology and Infectious Diseases*, 3(1), doi.org/10.4084/MJHID.2011.038.

- Keller, G-A., Warner, T.G., Steimer, K.S. and Hallawell, R.A. (1991). Cu, Zn superoxide dismutase is a perixomal enzyme in humans and hepatoma cells. *Cell Biology*, 88 (16), 7381-7385.
- Kerbel, R.S. (2008). Tumor angiogenesis. *The New England Journal of Medicine*, 385 (19), 2039-2049.
- Khanna, S., Pande, D., Negi, R., Karki, K., Khanna, R.S. and Khanna, H.D. (2012). Oxidative stress induced damage in benign and malignant breast diseases: histopathological and biochemical aspects. *Journal of Stress Physiology and Biochemistry*, 8 (1), 210-214.
- Kharis, Z., Azimathol Hawariah, L.P., Normah, A. And Aishah, R. (2010). *Morinda citrifolia* extract inhibits proliferation of non-small human lung cancer cell line NCI-H23 via apoptosis by modulating the Bax: Bcl-2 ratio. *Journal of Tropical Agriculture and Food Science*, 38(2), 203-209.
- Kim, W. R., Brown, R. S., Terrault, N. A. and El-Serag, H. (2002). Burden of liver disease in the United States: summary of a workshop. *Hepatology*, 36(1), 227-242.
- Kinoshita, S., Kase, S., Ando, R., Dong, Z., Fukuhara, J., Dong, Y., Inafuku, S., Noda, K., Noda, M., Kanda, A. and Ishida, S. (2014). Expression of vascular endothelial growth factor in human ocular adnexal lymphoma. *Investigative Ophthalmology and Visual Science*, 55(6), 3461-3467.
- Kipps, T. J. (2001). Chronic lymphocytic leukemia and related diseases. *Williams Hematology*, 6, 1163-1194.
- Kobayashi, H. and Lin, P. C. (2009). Angiogenesis links chronic inflammation with cancer. *Inflammation and Cancer: Methods and Protocols: Volume 1: Experimental Models and Practical Approaches*, 9(7), 185-191.
- Kohji, U., Masumi, Y., Shuzo, S., Akiyoshi, F., Kohji, K., and Toshimi, U. (2001). Sequential histopathological study on Tg-rasH2 mice after administration of MNU. *Toxicologic Pathology*, 29(3), 263–265.
- Kohn, H.I. and Liversedge, M. (1944). On a new aerobic material metabolite whose production by brain is inhibited by apomorphine emetine, ergotamine, epinephrine and menadione. *Journal of Pharmacology and Experimental Therapeutics*, 82(9), 292-300.
- Kotoh, K., Enjoji, M., Kato, M., Kohjima, M., Nakamuta, M. and Takayanagi, R. (2008). A new parameter using serum lactate dehydrogenase and alanine aminotransferase level is useful for predicting the prognosis of patients at an early stage of acute liver injury: A retrospective study. *Comparative Hepatology*, 7(6), 1-8.

- Kreja, L. and Seidel, H. J. (1986). T-cell markers in the thymus during methylnitrosourea-induced leukemogenesis. *Immunology Letters*, 12(2), 115-119.
- Krug, T., Ganser, A. and Koeffler, H.P. (2002). Tumor suppressor gene in normal and malignant hematopoiesis. *Oncogene*, 21(12), 34755-3495.
- Kshirsagar, A., Vetat, Y., Ashak, P., Bhosle, P. and Ingawale D. (2008). Drug induced hepatotoxicity: A comprehensive review, *The Internet Journal of Pharmacology*, 7(1), 1-17.
- Kumar, S., Witzig, T.E., Timm, M., Haug, J., Wellik, K., Fonseca, R., Greipp, P.R. and Rajkumar, S.V. (2003). Epression of VEGF and its receptors by myeloma cells. *Leukemia*, 17(3), 2025-2031.
- Kune, G. A., Kune, S. and Watson, L. F. (1988). Colorectal cancer risk, chronic illnesses, operations, and medications: case control results from the Melbourne Colorectal Cancer Study. *Cancer Research*, 48(15), 4399-4404.
- Kunsorn, P., Ruangrunsi, N., Lipipun, V., Khanboon, A. and Rungsihirunrat, K. (2013). The identities and anti-herpes simplex virus activity of *Clinacanthus nutans* and *Clinacanthus siamense*. *Asian Pacific Journal of tropical Biomedicine*, 3(4), 284-290.
- Lazar, D., Taban, S., Ardeleanu, C., Simionescu, C., Sporea, I., Cornianu, M., & Vernic, C. (2008). Immunohistochemical expression of the cyclooxygenase-2 (COX-2) in gastric cancer. The correlations with the tumor angiogenesis and patients' survival. *Romania Journal Morphology Embryology*, 49(3), 371-9.
- Lee, R., Margaritis, M., Channon, K.M. and Antoniadis, C. (2012). Evaluating oxidative stress in human cardiovascular disease: methodological aspects and considerations. *Current in Medical Chemistry*, 19(16), 2504-2520.
- Lee, R., Woo, W., Wu, B., Kummer, A., Duminy, H. and Xu, Z. (2003). Zinc Accumulation in N-methy-N-Nitrosourea-Induced rat mammary tumors is accompanied by an altered expression of ZnT-1 and metallothionein. *Experimental Biology and Medicine*, 228, 689-696.
- Leonel, C., Gelaleti, G. B., Jardim, B. V., Moschetta, M. G., Regiani, V. R., Oliveira, J. G. and Zuccari, D. A. (2014). Expression of glutathione, glutathione peroxidase and glutathione S-transferase pi in canine mammary tumors. *BMC Veterinary Research*, 10(1), 1-9.
- Lewis, P., Stefanovic, N., Pete, J., Calkin, A. C., Giunti, S., Thallas-Bonke, V., Jandeleit-Dahm, K.A., Allen, T.J., Kola, I., Copper, M.E and de Haan, J. B. (2007). Lack of the antioxidant enzyme glutathione peroxidase-1 accelerates atherosclerosis in diabetic apolipoprotein e-deficient mice. *Circulation*, 115(16), 2178-2187.

- Li, S., Yan, T., Yang, J. Q., Oberley, T. D. and Oberley, L. W. (2000). The role of cellular glutathione peroxidase redox regulation in the suppression of tumor cell growth by manganese superoxide dismutase. *Cancer Research*, 60(14), 3927-3939.
- Lijinsky, W. (1999). N-nitroso compounds in the diet. *Mutation Research-Genetic Toxicology and Environmental Mutagenesis*. 443, 129-138.
- Lim, K. H., Yoon, H. I., Kang, Y., Lee, K. W., Kim, J. H., Bang, S. M., and Lee, J. S. (2010). Severe pulmonary adverse effects in lymphoma patients treated with cyclophosphamide, doxorubicin, vincristine, and prednisone (CHOP) regimen plus rituximab. *The Korean Journal of Internal Medicine*, 25(1), 86-92.
- Lim, S. L., Goh, Y. M., Noordin, M. M., Rahman, H. S., Othman, H. H., Bakar, N. A. A., and Mohamed, S. (2016). *Morinda citrifolia* edible leaf extract enhanced immune response against lung cancer. *Food and Function*, 10.1039/C5FO01475A.
- Limat, S., Demesmay, K., Voillat, L., Bernard, Y., Deconinck, E., Brion, A., Sabah, A., Woronoff-Lemsi, M.C. and Cahn, J. Y. (2003). Early cardiotoxicity of the CHOP regimen in aggressive non-Hodgkin's lymphoma. *Annals of Oncology*, 14(2), 277-281.
- Lionel, C., Gelaleti, Gabriela, B., Jardin, Bruna, V., Moschetta, Marina, G., Regiani, Victor, R., Oliviera, Juliana, G. and Zuccari, Deborah, A.P.C. (2014). Expression of glutathione, glutathione peroxidase and glutathione S-transferase pi in canine mammary tumours. *BMC Veterinary research*, 10(49), doi:10.1186/1746-6148-10-49.
- Liu, J., Hinkhouse, M. M., Sun, W., Weydert, C. J., Ritchie, J. M., Oberley, L. W., and Cullen, J. J. (2004). Redox regulation of pancreatic cancer cell growth: role of glutathione peroxidase in the suppression of the malignant phenotype. *Human Gene Therapy*, 15(3), 239-250.
- Liu, J.M., Haroun-Bouhedja, F. and Boisson-Vidal, C. (2000). Analysis of the in vitro inhibition of mammary adenocarcinoma cell adhesion by sulphated polysaccharides. *Anticancer Research*, 20(5A), 3265-3271.
- Liu, Y., Cox, Shanna. R., Morita, T. and Kourembanas, S. (1995). Hypoxia regulates vascular endothelial growth factor gene expression in endothelial cells. *Circulation Research*, 77, 638-643.
- Lo, C. Y., Lam, K. Y., Leung, P. P. and Luk, J. M. (2005). High prevalence of cyclooxygenase 2 expression in papillary thyroid carcinoma. *European Journal of Endocrinology*, 152(4), 545-550.
- Lohr, J. W., Willsky, G. R. and Acara, M. A. (1998). Renal drug metabolism. *Pharmacological Reviews*, 50(1), 107-142.

- Lu, H., Ouyang, W. and Huang, C. (2006). Inflammation, a key event in cancer development. *Molecular Research Center*, 7, 4230-4238.
- Lubet, R. A., Gordon, G. B., Prough, R. A., Lei, X. D., You, M., Wang, Y and Moon, R. D. (1998). Modulation of methylnitrosourea-induced breast cancer in Sprague Dawley rats by dehydroepiandrosterone: dose-dependent inhibition, effects of limited exposure, effects on peroxisomal enzymes, and lack of effects on levels of Ha-Ras mutations. *Cancer Research*, 58(5), 921-926.
- Ma, D. L., Chen, M., Su, C. X. and West, B. J. (2013). In vivo antioxidant activity of deacetylasperulosidic acid in Noni. *Journal of Analytical Methods in Chemistry*, doi:10.1155/2013/804504.
- Maae, E., Nielsen, M., Steffensen, K. D., Jakobsen, E. H., Jakobsen, A. and Sørensen, F. B. (2011). Estimation of immunohistochemical expression of VEGF in ductal carcinomas of the breast. *Journal of Histochemistry and Cytochemistry*, 59(8), 750-760.
- Machado, D. E., Berardo, P. T., Palmero, C. Y. and Nasciutti, L. E. (2010). Higher expression of vascular endothelial growth factor (VEGF) and its receptor VEGFR-2 (Flk-1) and metalloproteinase-9 (MMP-9) in a rat model of peritoneal endometriosis is similar to cancer diseases. *Journal of Experimental and Clinical Cancer Research*, 7, 8-12.
- Macotpet, A., Suksawat, F., Sukon, P., Pimpakdee, K., Pattarapanwichien, E., Tangrassameeprasert, R. and Boonsiri, P. (2013). Oxidative stress in cancer-bearing dogs assessed by measuring serum malondialdehyde. *BMC Veterinary Research*, 9(101), doi:10.1186/1746-6148-9-101.
- Madinah, N., Nozmo, M. and Ezekiel, I. (2015). The protective effects of aqueous extract of Carica papaya seeds in paracetamol induced nephrotoxicity in male Wistar rats. *African Health Sciences*, 15(2), 598-605.
- Mahmood, T. and Yang, P.C. (2012). Western blot: technique, theory and trouble shooting. *North American Journal of Medical Sciences*, 4(9): 429-434.
- Mai, C. W., Yap, I. K. S., Kho, M. T., Ismail, N. H., Yusoff, K. M., Shaarie, K. and Lim, S. H. E. (2016). Mechanisms underlying the anti-inflammatory effects of Clinacanthus nutans Lindau extracts: inhibition of cytokine production and Toll-like receptor-4 activation. *Frontiers in Pharmacology*, 7, 7.
- Manaf, A., Hazilah, N., Johari, R. and Marikar, K. (2010). Positioning Malaysia in medical tourism, 1-21.
- Manaresi, E., Pasini, P., Gallinella, G., Gentilomi, S., Venturoli, A., Roda, M., Zerbini, M. and Musiani, M. (1999). Chemiluminescence western blot assay for the detection of immunity against parvovirus B19 VP1 and VP2 linear epitopes using a video camera based luminograph. *Journal of Virological Methods*, 81, 91-99.

- Marino, D. J. (2012). Age-specific absolute and relative organ weight distributions for B6C3F1 mice. *Journal of Toxicology and Environmental Health, Part A*, 75(2), 76-99.
- Markovic, A., MacKenzie, K.L. and Lock, R.B. (2011). Induction of vascular endothelial growth factor secretion by childhood acute lymphoblastic leukemia cells via the FLT-3signaling pathways. *Molecular Cancer Therapeutics*, 11(1), 183-193.
- Marum, L., Miguel, A., Ricardo, C. P., & Miguel, C. (2012). Reference gene selection for quantitative real-time PCR normalization in *Quercus suber*. *PloS One*, 7(4), 114-119.
- Masood, A., Kairouz, S., Hudhud, K. H., Hegazi, A. Z., Banu, A. and Gupta, N. C. (2009). Primary non-Hodgkin lymphoma of liver. *Current Oncology*, 16(4), 74-78.
- Massaad, C. A., Washington, T. M., Pautler, R. G. and Klann, E. (2009). Overexpression of SOD-2 reduces hippocampal superoxide and prevents memory deficits in a mouse model of Alzheimer's disease. *Proceedings of the National Academy of Sciences*, 106(32), 13576-13581.
- MatÉs, J. M., Pérez-Gómez, C. and De Castro, I. N. (1999). Antioxidant enzymes and human diseases. *Clinical biochemistry*, 32(8), 595-603.
- McClatchey, W. (2002). From Polynesians healers to health food stores: changing perspectives of *Morinda citrifolia* (Rubiaceae). *Integrative Cancer Therapies*, 1(2), 110-120.
- McKay, L.I and Cidlowski, J.A. (2003). Corticosteroids in the treatment of neoplasms. 6th. Hamilton. Holland-Frei Cancer Medicine.
- McKelvey, E.M., Gottlieb, J.A. and Wilson, H.E. (1976). Hydroxyldaunomycin (Adrimycin) combination chemotherapy in malignant lymphoma. *Cancer*, 38, 1484- 1493.
- McLeay, Y., Barnes, M.J., Mundel, T., Hurst, S.M., Hurst, R.D. and Stannard, S.R. (2012). Effect of New Zealand blueberry consumption on recovery from eccentric exercise-induced muscle damage. *Journal of the International Society of Sports Nutrition*, 9(1), 9-19.
- McMicheal, M.A. (2007). Oxidative stress, antioxidants, and assessment of oxidative stress in dogs and cats. *Journal of the American Veterinary Medical Association*, 231(5), 714-720.
- Medinger, M., Fischer, N. and Tzankov, A. (2010). Vascular endothelial growth factor-related pathways in hemato-lymphoid malignancies. *Journal of Oncology*, 28, 1935-1944.

- Meng, H., Guo, J., Sun, J. Y., Pei, J. M., Wang, Y. M., Zhu, M. Z., & Huang, C. (2008). Angiogenic effects of the extracts from Chinese herbs: Angelica and Chuanxiong. *The American Journal of Chinese Medicine*, 36(03), 541-554.
- Milionis, J., Bourantas, C.L., Siamopoulus, K.C. and Elisaf, M. S. (1999). Acid-base and electrolyte abnormalities in patients with acute leukaemia. *American Journal of Haematology*, 62(4), 201-207.
- Millanta, F., Asproni, P., Canale, A., Citi, S. and Poli, A. (2014). COX-2, mPGES-1 and EP2 receptor immunohistochemical expression in canine and feline malignant mammary tumours. *Veterinary and Comparative Oncology*, doi: 10.1111/vco.12096.
- Miller, J.A., Lang, J.E., Ley, M., Nagle, R., Hsu, C-H., Thompson, P.A., Cordova, C., Waer, A. and Sherry, Chow, H-H. (2013). Human breast tissue deposition and bioactive of limonene in women with early stage breast cancer. *Cancer Prevention Research*, 6(6), 577-584.
- Millonig, G., Stadlmann, S. and Vogel, W. (2005). Herbal hepatotoxicity: acute hepatitis caused by a Noni preparation (*Morinda citrifolia*). *European Journal of Gastroenterology and Hepatology*, 17(4), 445-447.
- Mohamed, E.A.S., Lim, C.P., Ebrika, O.S., Asmawi, M.Z., Sadikun, A. and Yam, M.F. (2011). Toxicity evaluation of a standardised 50% ethanol extract of *Orthosiphon stamineus*. *Journal of Ethnopharmacology*, 133, 358-363.
- Mohammed, S. I., Knapp, D. W., Bostwick, D. G., Foster, R. S., Khan, K. N. M., Masferrer, J. L., Woerner, B.M., Snyder, P.W and Koki, A. T. (1999). Expression of cyclooxygenase-2 (COX-2) in human invasive transitional cell carcinoma (TCC) of the urinary bladder. *Cancer Research*, 59(22), 5647-5650.
- Mohd. Shukri, M.A. and Alan, C. (2010). Analysis of phenolics in *Anacardium occidentale* shoots extracts using a reversed-phase high performance liquid chromatography tandem mass spectrometry (RP-HPLC-MS). *Journal of Tropical Agriculture and Food Science*, 38(2), 221-230.
- Mokuno, K., Riku, S., Sugimura, K., Takahashi, A., Kato, K. and Osugi, S. (1987). Serum creatine kinase isoenzymes in Duchenne muscular dystrophy determined by sensitive enzyme immunoassay methods. *Muscle and Nerve*, 10(5), 459-463.
- Moldovan, L. and Moldovan, N. I. (2004). Oxygen free radicals and redox biology of organelles. *Histochemistry and Cell Biology*, 122(4), 395-412.
- Monacci, W.T., Merrill, M.J. and Oldfield, E.H. (1993). Expression of vascular permeability factor vascular endothelial growth factor in normal rat tissues. *American Journal of Physiology*, 4, 995-1002.

- Moore, K. P. and Aithal, G. P. (2006). Guidelines on the management of ascites in cirrhosis. *Gut*, 55(6), 1-12.
- Mori, N, Inoue, H., Yoshida, T., Tanabe, T. and Yamamoto, N. (2001). Constitutive expression of the cyclooxygenase-2 gene in the T-cell lines infected with human T cell leukaemia virus type 1. *International Journal of Cancer*, 94, 813-819.
- Morrone, F.B., Spiller, F., Edelwises, M.I.A., Meurer, L., Engroff, P. and Barrios, C.H. (2009). Effect of temozolomide treatment on the adenine nucleotide hydrolysis in blood serum of rats with implanted gliomas. *Applied Cancer Research*, 29,118–124.
- Morton, D., Bailey, K. L., Stout, C. L., Weaver, R. J., White, K. A., Lorenzen, M. J. and Ball, D. J. (2008). N-Methyl-N-Nitrosourea (MNU): A positive control chemical for p53+/- mouse carcinogenicity studies. *Toxicologic Pathology*, 36(7), 926-931.
- Mozos, I. (2015). Mechanisms Linking Red Blood Cell Disorders and Cardiovascular Diseases. *BioMed research international*, 682054-682054.
- Mudie, N. Y., Swerdlow, A. J., Higgins, C. D., Smith, P., Qiao, Z., Hancock, B. W., Hoskin, P.J. and Linch, D. C. (2006). Risk of second malignancy after non-Hodgkin's lymphoma: a British Cohort Study. *Journal of Clinical Oncology*, 24(10), 1568-1574.
- Mughal, T.I., Goldman, J.M. and Mughal, S.T. (2006). Understanding Leukaemias, Lymphoma and Myelomas. 1st edition, United Kingdom: Taylor and Francis, 117–122.
- Muhamad Hakimi Bin Mohd. Kassim. (2014). Carcinogenic effects of N-methyl-N-nitrosourea administration via oral and intraperitoneal routes in female Sprague Dawley rats. Master Thesis. Universiti Putra Malaysia.
- Murnah, M. (2014). Effect of Ethanol Noni (*Morinda citrifolia* L) Extract on Diabetic Nephropathy Sprague Dawley Rats Induced Streptozotocin (STZ). *Journal of Nutrition and Health*, 2(1), 1-15.
- Muthukkaruppan, V.R., Kubai, L. and Auerbach, R. (1982). Tumor-induced neovascularization in the mouse eyes. *Journal of the National Cancer Institute*, 69(3), 699-708.
- Muto, J., Hosung, L., Uwaya, A., Isami, F., Ohno, M. and Mikami, T. (2010). *Morinda citrifolia* fruit reduces stress-induced impairment of cognitive function accompanied by vasculature improvement in mice. *Physiology and Behaviour*, 101(2), 211-217.
- Na-Bangchang, K., Ketsa-ard, K., Wiwatwittaya, S., Thongprasert, S., Vongsakul, M., Picha, P. and Karbwang, J. (2012). Anticancer activity and immunostimulating

effect on NK-cell activity of a well-known Thai folkloric remedy. *International Research Journal of Pharmacology*, 2, 1-11.

Nagakawa, T., Mazzali, M., Kang, D.H., Kanellis, J., Watanabe, S., Sanchez-Lozada, L.G., Rodriguez-Iturbe, B., Herrera-Costa, J. and Johnson, R.J. (2003). Hyperuricemia causes glomerular hypertrophy in the rat. *American Journal of Nephrology*, 23, 2-7.

Nakanishi, Y., Kamijo, R., Takizawa, K., Hatori, M. and Nagumo, M. (2001). Inhibitors of cyclooxygenase-2 (COX-2) suppressed the proliferation and differentiation of human leukaemia cell lines. *European Journal of Cancer*, 37(12), 1570-1578.

Nelson, S.C. (2001). Noni cultivation in Hawaii. *Fruit and Nuts*, 4, 1-14.

Netto, G.J., Saad, R.D., Dysert, P.A. (2003). Diagnostic molecular pathology: current techniques and clinical applications. *Baylor University Medical Center Proceedings*, 16(4), 379-383.

Nishida, N., Yano, H., Nishida, T., Kamura, T. and Kojiro, M. (2006). Angiogenesis in cancer. *Journal of Vascular Health Risk and Management*, 2(3), 213-219.

Noronha, V., Shafi, N. Q., Obando, J. A. and Kummar, S. (2005). Primary non-Hodgkin's lymphoma of the liver. *Critical Reviews in Oncology/Hematology*, 53(3), 199-207.

Nowakowski, G. S., Hoyer, J. D., Shanafelt, T. D., Zent, C. S., Call, T. G., Bone, N. D., LaPlant, B., Dewald, G.W., Tschumper, R.C., Jelinek, D.F., Witzig, T.E. and Kay, N.E. (2009). Percentage of smudge cells on routine blood smear predicts survival in chronic lymphocytic leukemia. *Journal of Clinical Oncology*, 27(11), 1844-1849.

Nualsanit, T., Rojanapanthu, P., Gritsanapan, W., Lee, S-H., Lawson, D. and Baek, S-J. (2012). Damnacanthol, a Noni component, exhibits anti-tumorigenic activity in human colorectal cancer cells. *The Journal of Nutritional Biochemistry*, 23(8), 914-923.

Nursyuhada Haron. (2012). Preventive effects of consuming nutritional supplement of *Morinda citrifolia* on the transcription of Bax and Bcl-2 in early leukaemic rats. Master Thesis. Universiti Putra Malaysia.

Nursyuhada, H., Hazilawati, H., Hutheyfa, A.H., Rosly, S.M., Shanmugavellu, S., Noordin, M.M. and Jasni, S. (2011). Detection of Bcl-2 in leukaemic rats using an EvaGreen real-time RT-PCR assay. *Pertanika Journal of Tropical Agriculture Science*. 34(2), 372-380.

- Nurul, Husna, R., Noriham, A., Noorain, A., Aziziah, H. and Farah Amna, O. (2013). Acute oral toxicity effects of *Momordica Charantia* in Sprague Dawley rats. *International Journal of Bioscience, Biochemistry and Bioinformatics*, 3(4), 408-410.
- Ogbonnia, S.O., Mbaka, G.O., Anyika, E.N., Osegbo, O.M. and Igbokwe, N.H. Evaluation of acute toxicity of hydro-ethanolic extract of *chromolaena odorata* (L.) king and robinson (Fam. Asteracea) in rats (2010). *Agriculture and Biology Journal of North America*, 1, 869-865.
- Ohkawa, H., Ohishi, N. and Yagi, K. (1979). Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry*, 95(2), 351-358.
- Okokon, J. E., Nwafor, P. A., and Ekpo, M. D. (2010). Subchronic toxicity studies of the ethanolic root extract of *Croton zambesicus*. *Pakistan Journal of Pharmaceutical Science*, 23(2), 160-169.
- Olusola, L., Matthew, O. and Oluwatosin, A. (2015). Comparative study on the effects of aqueous extracts of *Viscum album* (mistletoe) from three host plants on hematological parameters in albino rats. *African Health Sciences*, 15(2), 606-612.
- Ono, M. (2008). Molecular links between tumor angiogenesis and inflammation: inflammatory stimuli of macrophages and cancer cells as targets for therapeutic strategy. *Cancer Science*, 99(8), 1501-1506.
- Ostapowics, G., Fontana, R.J., Schiodt, F.V. (2002). Results of a prospective study of acute liver failure at 17 tertiary care centers in the United States. *Annals of Intern Medicine*, 137(12), 947-954.
- Osterberg, E. C., Ramasamy, R., Masson, P. and Brannigan, R. E. (2014). Current practices in fertility preservation in male cancer patients. *Urology Annals*, 6(1), 13.
- Othman, A., Ismail, A., Ghani, A.N. and Adenan, I. (2007). Antioxidant capacity and phenolic content of cocoa beans. *Food Chemistry*, 100, 1523-1530.
- P'ng, X.W., Akowuah, G.A. and Chin, J.H (2012). Acute oral toxicity study of *Clinacanthus nutans* in mice. *International Journal of Pharmaceutical Sciences and Research*, 3(11), 4202-4205.
- Palu, A. K., Deng, S., West, B. J., Jensen, J. and Sabin, R. A. (2012). Sunburn (fohia) Healing Effects of Noni: Is it a Mechanism Involving Its Inhibitory Effects on MMP, COX-2 and Cat-G Enzymes?. *Journal of Applied Pharmaceutical Science*, 2(8), 40-50.
- Peng, T.W., P'ng, X.W., Chin, J.H and Akowuah, G.A. (2014). Effect of methanol extract of *Clinacanthus nutans* on serum biochemical parameters in rats. *Journal of Applied Pharmaceutical Science*, 6(1), 77-86.

- Perse, M., Cerar, A., Injac, R. and Strukelj, B. (2008). *N*-methylnitrosoourea induced breast cancer in rat, the histopathology of the resulting tumours and its drawbacks as a model. *Pathology and Oncology Research*, 15(1), 115-121.
- Petrola, Maria, J., Montenegro, de, Castro, A.J. and Goncalves, R.P. (2012). Serum concentrations of nitrite and malondialdehyde as markers of oxidative stress of chronic myeloid leukemia patients treated with tyrosine kinase inhibitors. *Revista Brasileira de Hematologica e Hemoterapia*, 34(4), 352-255.
- Petterino, C. and Argentino-Storino, A. (2006). Clinical chemistry and hematology historical data in control Spague-Dawley rats from pre-clinical toxicity studies. *Experimental Toxicology Pathology*, 57(3), 213-219.
- Pettersson, J., Karlsson, P. C., Göransson, U., Rafter, J. J. and Bohlin, L. (2008). The flavouring phytochemical 2-pentanone reduces prostaglandin production and COX-2 expression in colon cancer cells. *Biological and Pharmaceutical Bulletin*, 31(3), 534-537.
- Phillip, M., Rowley, D.A. and Schreiber, H. (2004). Inflammation as a tumor promoter in cancer induction. *Seminars in Cancer Biology*, 14, 433-439.
- Poli, G., Dianzani, M. U., Cheeseman, K. H., Slater, T., Lang, J., and Esterbauer, H. (1985). Separation and characterization of the aldehydic products of lipid peroxidation stimulated by carbon tetrachloride or ADP-iron in isolated rat hepatocytes and rat liver microsomal suspensions. *Biochemical Journal*, 227, 629-638.
- Potti, A., Ganti, A. K., Kargas, S. and Koch, M. (2001). Immunohistochemical detection of C-kit (CD117) and vascular endothelial growth factor (VEGF) overexpression in mantle cell lymphoma. *Anticancer Research*, 22(5), 2899-2901.
- Prager, G. W., Poettler, M., Unseld, M. and Zielinski, C. C. (2012). Angiogenesis in cancer: anti-VEGF escape mechanisms. *Translational lung cancer research*, 1(1), 14.
- Psathakis, D., Wedemeyer, N., Oevermann, E., Krug, F., Siegers, C. P. and Bruch, H. P. (1998). Blood selenium and glutathione peroxidase status in patients with colorectal cancer. *Diseases of the Colon and Rectum*, 41(3), 328-335.
- Pujari, K.N. and Jadkar, S.P. (2011). Superoxide dismutase levels in leukemias. *International Journal of Basic Medical Science*, 2(2), 96-100.
- Rafati, A., Yasini, S. M., Dashti-R, M. H., Pakdel, S. and Norani, F. (2006). Tramadol dependence rate as compared with morphine in rats. *World Journal of Medical Sciences*, 1(1), 40-43.
- Ramos-Vara, J.A. (2011). Principles and methods of immunohistochemistry. *Methods of Molecular Biology*, 691, 83-96.

- Reuter, S., Gupta, S. C., Chaturvedi, M. M. and Aggarwal, B. B. (2010). Oxidative stress, inflammation, and cancer: how are they linked?. *Free Radical Biology and Medicine*, 49(11), 1603-1616.
- Rhiouani, H.R., Nazari, P., Kamli-Nejad, M. and Lyoussi, B. (2008). Acute and subchronic oral toxicity of an aqueous extract of leaves of *Herniaria glabra* in rodents. *Journal of Ethnopharmacology*, 118, 378-386.
- Roelofs, H. M., te Morsche, R. H., van Heumen, B. W., Nagengast, F. M. and Peters, W. H. (2014). Over-expression of COX-2 mRNA in colorectal cancer. *BMC Gastroenterology*, 14(1), 1-9
- Roomi, M. W., Roomi, N. W., Ivanov, V., Kalinovsky, T., Niedzwiecki, A. and Rath, M. (2005). Modulation of N-methyl-N-nitrosourea induced mammary tumors in Sprague–Dawley rats by combination of lysine, proline, arginine, ascorbic acid and green tea extract. *Breast Cancer Research*, 7(3), 1-9.
- Roorda, B.D., Ter Elst, A., Scherpen, F.J., Meeuwssen-de Boer, T.G., Kamps, W.A. and Bont, E.S. (2010). VEGF-A promotes lymphoma tumor growth by activation of STAT proteins and inhibition of p27 (K1P1) via paracrine mechanisms. *European Journal of Cancer*, 46(5), 974-982.
- Rosidah, Yam, M.F., Sadikun, A., Ahmad, A., Akowuah, G.A. and Asmawi, M.Z. (2009). Toxicology evaluation of standardized methanol extract of *Gynura procumbens*. *Journal of Ethnopharmacology*, 123, 244–249.
- Rosly, S.M., Shanmugavelu, S., Murugaiyah, M., Hadija, H., Ahmad Tarmizi, S., Noridayusni, Y. and Subramaniam, K. (2011). Subchronic oral toxicity of *Morinda citrifolia* (mengkudu) in Sprague Dawley rats. *Pertanika Journal of Tropical Agriculture Science*, 34(2), 341-349.
- Rozenberg, G. (2011). Microscopic haematology: A practical guide for the laboratory. *Elsevier Australia*, 106–107.
- Saedi, T.A., Md, Noor, S., Ismail, P. and Othman, F. (2014). The effects of herbs and fruits on leukaemia. Evidence-Based Complementary and Alternative Medicine, 2014 (2014), dx.doi.org/10.1155/2014/494136.
- Safarzadeh, E., Sandoghchian Shotorbani, S. and Baradaran, B. (2014). Herbal medicine as inducers of apoptosis in cancer treatment. *Advanced Pharmaceutical Bulletin*, 4(1), 421-427.
- Saganuwan, S.A., Aondoaver, A.D., Roman I.T. (2014). Reassessment of acute and chronic toxicity effect of aqueous leaf extract of *Morinda Lucida* in *Rattus norvegicus*. *Journal of Haematology Research*, 1, 36-46.
- Said, N. M. and Abiola, O. (2014). Haematological profile shows that Inbred Sprague Dawley rats have exceptional promise for use in biomedical and pharmacological studies. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 4(37), 33-37.

- Saito, T., Kurasaki, M., Kaji, H. and Saito, K. (1984). Deficiency of erythrocytes superoxide dismutase and catalase activities in patients with malignant lymphoma and acute myeloid leukemia. *Cancer letters*, 24(2), 141-146.
- Sajjarattul, N.N.A., Hazilawati, H., Rosly, S.M., Hakimi, M.K., Nursyuhada, H., Yusnani, M.Y., Hutjeyfa, A.H., Nurul, Huda, M.Z., Shanmugavelu, S., Noordin, M.M., Jasni, S., and Tan, W. (2014). Detection of VEGF mRNA transcript in rat spleen exposed to *N*-methyl-*N*-nitrosourea via oral and intraperitoneal routes. *Online Journal of Veterinary Medicine*. 18(9), 756-766.
- Sakdarat, S., Shuyprom, A., Dechatiwongse Na Ayudhya T., Waterman P.G. and Karagianis, G. (2006). Chemical composition investigation of the *Clinacanthus nutans* Lindau leaves. *Thai Journal of Phytopharmacy*, 13(2), 2459-2551.
- Sakdarat, S., Shuyprom, A., Pientong, C., Ekalaksananan, T. and Thongchai, S. (2009). Bioactive constituents from the leaves of *Clinacanthus nutans* Lindau. *Bioorganic and Medicinal Chemistry*, 17, 1857-1860.
- Sakran, M., Selim, Y. and Zidan, N. (2014). A new isoflavonoid from seeds of *Lepidium sativum* L. and its protective effect on hepatotoxicity induced by paracetamol in male rats. *Molecules*, 19(10), 15440-15451.
- Saladin, K.S. (2004). Anatomy and physiology-the unity of form and function. 3rd. New York: McGraw-Hill Publishers; 2004: 94–128.
- Saminathan, M., Rai, R. B., Dhama, K., Jangir, B. L., Suresh, S., Ranganath, G. J., Sophia, I., Karupanasammy, K., Barathiraja, S. and Gopalakrishnan, A. (2014). Effect of *Morinda citrifolia* (Noni) Fruit Juice on Antioxidant, Hematological and Biochemical Parameters in N-Methyl-N-Nitrosourea (NMU) Induced Mammary Carcinogenesis in Sprague-Dawley Rats. *International Journal of Pharmacology*, 10(2), 109-119.
- Sandalio, L. M., López-Huertas, E., Bueno, P., & Del Río, L. A. (1997). Immunocytochemical Localization of Copper, Zinc Superoxide Disrnutase in Peroxisomes from Wihermelon (*Citrullus vulgaris* Schrad.) Cotyledons. *Free Radical Raeseach*, 26(3), 187-194.
- Sano, H., Kawahito, Y., Wilder, R. L., Hashiramoto, A., Mukai, S., Asai, K., Kimura, S., Kato, H., Koto, M. and Hla, T. (1995). Expression of cyclooxygenase-1 and-2 in human colorectal cancer. *Cancer Research*, 55(17), 3785-3789.
- Sanyal, J., Bandyopadhyay, S. K., Banerjee, T. K., Mukherjee, S. C., Chakraborty, D. P., Ray, B. C. and Rao, V. R. (2009). Plasma levels of lipid peroxides in patients with Parkinson's disease. *European. Review for Medical Pharmacology Sciences*, 13(2), 129-132.
- Sarega, N., Imam, M. U., Ooi, D. J., Chan, K. W., Md Esa, N., Zawawi, N. and Ismail, M. (2016). Phenolic Rich Extract from *Clinacanthus nutans* Attenuates

Hyperlipidemia-Associated Oxidative Stress in Rats. *Oxidative Medicine and Cellular Longevity*, 4137908-4137908.

- Sathiya, M. and Muthuchelian, K (2009). Significance of immunologic markers in the diagnosis of lymphoma. *Academic Journal of Cancer Research*. 2, 40 -50.
- Satomi, A., Murakami, S., Hashimoto, T., Ishida, K., Matsuki, M. and Sonoda, M. (1995). Significance of superoxide dismutase (SOD) in human colorectal cancer tissue: correlation with malignant intensity. *Journal of gastroenterology*, 30(2), 177-182.
- Schacht, V., and Kern, J. S. (2015). Basics of immunohistochemistry. *Journal of Investigative Dermatology*, 135(3), 1-4.
- Schmidt GW and Delaney SK. Stable internal reference genes for normalization of real-time RT-PCR in tobacco (*Nicotiana tabacum*) during development and abiotic stress. *Molecule Genetics and Genomics*. 283(3), 233-241. 2010.
- Schwane, J.A., Johnson, S.R., Vandenakker, C.B. and Armstrong R.B (1983). Delayed-onset muscular soreness and plasma CPK and LDH activities after downhill running. *Medicine and Science in Sports and Exercise*, 15(1), 51-56.
- Secchiero, P., Barbarotto, E., Gonelli, A., Tiribelli, M., Zerbinati, C., Celeghini, C. and Zauli, G. (2005). Potential pathogenetic implications of cyclooxygenase-2 overexpression in B chronic lymphoid leukemia cells. *The American Journal of Pathology*, 167(6), 1599-1607.
- Serafini, M.R., Moreira de Lima, C., Santos, R.C., Doria, G.A.A., Yohanna de Jesus, J., Doria de Melo, M., Paulo, J., dos Santos, A., Rabello, T.K., Fortes, V.S., Fonceca, M.J.V., Calvancanti de Albuquerque, R.L., Quintans-Junior, L.J. and Antunes de Saoza Araujo, A. (2011). Pre-clinical toxicity of *Morinda citrifolia* Linn leaf extract. *African Journal of Biotechnology*, 10(65), 14566-14572.
- Shahid, M., Subhan, F., Ullah, I., Ali, G., Alam, J. and Shah, R. (2016). Beneficial effects of *Bacopa monnieri* extract on opioid induced toxicity. *Heliyon*, 2(2), e00068.
- Sheng, H., Shao, J., Morrow, J. D., Beauchamp, R. D. and DuBois, R. N. (1998). Modulation of apoptosis and Bcl-2 expression by prostaglandin E₂ in human colon cancer cells. *Cancer Research*, 58(2), 362-366.
- Shetty, S.R., Babu, S.G., Kumari, S., Karikal, A., Shetty, P. and Hedge, S. (2013). Salivary superoxide dismutase levels in oral leukoplakia and oral squamous cell carcinoma; a clinicopathology study. *Oxidants and Antioxidants in Medical Sciences*, 2(1), 69-71.
- Shimoyama, M. (1991). Diagnostic criteria and classification of clinical subtypes of adult T-cell leukaemia-lymphoma. *British Journal of Haematology*, 79(3), 428-437.

- Shweiki, D., Itin, A., Soffer, D. and Keshet, E. (1992). Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis. *Nature*, 359(6398), 843-845.
- Siegel, R.L., Miller, K.D. and Jemal, A. (2015). Cancer statistics, 2015. *A cancer Journal for Clinicians*, 6, 5-29.
- Singh, B., Berry, J. A., Shoher, A., Ramakrishnan, V. and Lucci, A. (2005). COX-2 overexpression increases motility and invasion of breast cancer cells. *International Journal of Oncology*, 26(5), 1393-1399.
- Sinnhuber, R. O., Yu, T. C. and Yu, T. C. (1958). Characterization of the red pigment formed in the 2-thiobarbituric acid determination of oxidative rancidity a, b. *Journal of Food Science*, 23(6), 626-634.
- Slongo, M. L., Molena, B., Brunati, A. M., Frasson, M., Gardiman, M., Carli, M., Perilongo G., Roselon, A. and Onisto, M. (2007). Functional VEGF and VEGF receptors are expressed in human medulloblastomas. *Neuro-Oncology*, 9(4), 384-392.
- Sloss, A. and Kubler, P. (2009). Prescribing in liver, *Australian Prescriber*, 32, 32-35.
- Smolková, K., and Ježek, P. (2012). The role of mitochondrial NADPH-dependent isocitrate dehydrogenase in cancer cells. *International Journal of Cell Biology*, 2012, 273947-273947.
- Sobolewski, C., Cerella, C., Dicato, M., Ghibelli, L., & Diederich, M. (2010). The role of cyclooxygenase-2 in cell proliferation and cell death in human malignancies. *International Journal of Cell Biology*, dx.doi.org/10.1155/2010/215158
- Song, G., Li, Y. and Jiang, G. (2012). Role of VEGF/VEGFR in the pathogenesis of leukemias and as treatment targets (Review). *Oncology Reports*, 28(6), 1935-1944.
- Sosa, V., Moliné, T., Somoza, R., Paciucci, R., Kondoh, H. and LLeonart, M. E. (2013). Oxidative stress and cancer: an overview. *Ageing Research Reviews*, 12(1), 376-390.
- Stadlbauer, Vanessa, Peter Fickert, Carolin Lackner, Jutta Schmerlaib, Peter Krisper, Michael Trauner, and Rudolf E. Stauber. "Hepatotoxicity of NONI juice: report of two cases." *World Journal of Gastroenterology*, 11(30), 4758-4760.
- Steensma, D.P. and Witzig, T.E (2001). Elevated serum LDH in patients with non-Hodgkin's lymphoma: not always an ominous sign. *British Journal of Haematology*, 107(2), 463-464.
- Strati, P. and Shanafelt, T. D. (2015). Monoclonal B-cell lymphocytosis and early-stage chronic lymphocytic leukemia: diagnosis, natural history, and risk stratification. *Blood*, 126(4), 454-462.

- Suh, E.J., Kabir, M.H., Kang, U-B., Lee, J.W., Yu, J., Noh, D-Y. and Lee, C. (2012). Comparative profiling of plasma proteome from breast cancer patients reveals trombospondin-1 and BRWD3as serological biomarkers. *Experimental and Molecular Medicine*, 44, 36-44.
- Sumba, P.O., Kabiru, E.W, Namuyenga, E., Fiore, N., Otieno, R.O., Moormann, A.M., Orago, A.S., Rosenbaum, P.F. and Rochford, R. (2010). Microgeographic variations in Burkitt's lymphoma incidence correlate with difference in malnutrition, amalaria and Epstein-Barr virus. *British Journal of Cancer*, 103, 1736-1741.
- Sung, H. J., Kim, S. J., Lee, J. H., Lee, G., Lee, K. A., Choi, C. W., Kim, J. S. (2007). Persistent anemia in a patient with diffuse large B cell lymphoma: pure red cell aplasia associated with latent Epstein-Barr virus infection in bone marrow. *Journal of Korean Medical Science*, 22, 167-S170.
- Suthar, C., Rana, T., Singh, U.B., Singh, M., Ramesh, V., Sharma, V.K., Ramam, M., (2013). mRNA and DNA PCR tests in cutaneous tuberculosis. *Indian Journal of Dermatology, Venereology, and Leprology*, 79(1), 65-69.
- Svensk, A-M., Soini, Y., Paakko, P., Hirvikoski, P. and Kinnula, V.L. (2004). Differential expression of superoxide dismutases in lung cancer. *American Journal of Clinical Pathology*, 122, 395-404.
- Swann, F., P. and Magee, P., N. (1968). Nitrosamine induced carcinogen. The alkylation of nucleic acids of the rat by *N*-methyl-*N*-nitrosurea, dimethylnitrosamine, dimethyl sulphate and methyl methanesulphate. *Biochemical Journal*. 110(1), 39-47.
- Szumilak, D., Sułowicz, W. and Walatek, B. (1997). [Rhabdomyolysis: clinical features, causes, complications and treatment]. *Przegląd Lekarski*, 55(5), 274-279.
- Taib, I. S., Budin, S. B., Ain, S. M. S. N., Mohamed, J., Louis, S. R., Das, S., and Hidayatulfathi, O. (2009). Toxic effects of *Litsea elliptica* Blume essential oil on red blood cells of Sprague-Dawley rats. *Journal of Zhejiang University SCIENCE B*, 10(11), 813-819.
- Takahashi, Y., Kitadai, Y., Bucana, C.D., Clearly, K.R. and Ellis, M. (1995). Expression of vascular endothelial growth factor and its receptor, KDR, correlates with vascularity, metastasis, and proliferation of human colon cancer. *Cancer Research*, 55(18), 3964-3968.
- Takaoka, M., Sehata, S., Maejima, T., Imai, T., Torii, M., Satoh, H., Toyosawa, K., Tanakamaru, Z.Y., Adachi, T., Hisada, S., Ueda, M., Ogasawara, H., Matsumoto, M., Kobayashi, K., Mutai, M. and Usui, T. (2003). Interlaboratory comparison of short-term carcinogenicity studies using CB6F1-rasH2 transgenic mice. *Toxicologic Pathology*. 31(2), 191–199.

- Takayama, S., Thorgeirsson, U. P. and Adamson, R. H. (2008). Chemical carcinogenesis studies in nonhuman primates. *Proceedings of the Japan Academy. Series B, Physical and biological sciences*, 84(6), 176-188.
- Tan Bee Ling. (2015). Anti-colon cancer activity and mechanisms of action of brewer's rice in vitro and in vivo. Doctor of Philosophy's thesis, Universiti Putra Malaysia, Malaysia.
- Tandon, R., Bhakar, M., Pande, D., Karki, K., Negi, R. and Khana, H.D. (2012). Oxidative stress induced lipid peroxidation and DNA adduct formation in the pathogenesis of multiple myeloma and lymphoma. *Journal of Stress Physiology and Biochemistry*, 9(1), 106-112.
- Teo, S., Stirling, D., Thomas, S., Hoberman, A., Kiorpes, A. And Khetani, V. (2002) A 90-day oral gavage toxicity study of D-methylpenidate and D, L-methylpenidate in Sprague-Dawley rats. *Toxicology*, 179(3), 183-196.
- Thorpe, M. P., Valentine, R. J., Moulton, C. J., Wagoner Johnson, A. J., Evans, E. M. and Layman, D. K. (2011). Breast tumors induced by *N*-methyl-*N*-nitrosourea are damaging to bone strength, structure, and mineralization in the absence of metastasis in rats. *Journal of Bone and Mineral Research*, 26(4), 769-776.
- Thun, M.J., Namboodiri, M.M. and Health, C.W. (1991). Aspirin use and reduced risk of fatal of colon cancer. *The New England Journal of Medicine*, 325, 1593-1596.
- Toyota M, Shen L, Ohe-Toyota M, Hamilton SR, Sinicrope FA, Issa JP. (2000). Aberrant methylation of the cyclooxygenase 2 CpG island in colorectal tumors. *Cancer Research*, 60, 4044–4048.
- Trottmann, M., Becker, A. J., Stadler, T., Straub, J., Soljanik, I., Schlenker, B. and Stief, C. G. (2007). Semen quality in men with malignant diseases before and after therapy and the role of cryopreservation. *European Urology*, 52(2), 355-367.
- Tsubura, A., Lai, Y.C., Miki, H., Sasaki, T., Uehara, N., Yuri, T. and Yoshizawa, K. (2011). Review: animal models of *N*-methly-*N*-ntrosourea-induced mammary cancer and retinal degeneration with special emphasis on therapeutic trials. *International Journal of Experimental and Clinical Pathophysiology and Drug Research*, 25, 11-22.
- Tu, S-F., Liu, R-H., Cheng, Y-B., Hsu, Y-M., Du, Y-C., El-Shazly, M., Wu, Y-C. and Chang, F-R. (2014). Chemical constituents and bioactivities of *Clinacanthus nutans* aerial parts. *Molecules*, 19, 20382-20390.
- Tygstrup, N. (1990). Assessment of liver function: principles and practice. *Journal of Gastroenterology and Hepatology*, 5(4), 468-482.

- Tzoran, I., Rebibo-Sabbah, A., Brenner, B. and Aharon, A. (2015). Disease dynamics in patients with acute myeloid leukemia: New biomarkers. *Experimental Hematology*, 43(11), 936-943.
- Vaisheva, F., Delbes, G., Hales, B. F. and Robaire, B. (2007). Effects of the Chemotherapeutic Agents for Non-Hodgkin Lymphoma, Cyclophosphamide, Doxorubicin, Vincristine, and Prednisone (CHOP), on the Male Rat Reproductive System and Progeny Outcome. *Journal of Andrology*, 28(4), 578-587.
- Van der Loos, C. M., Meijer-Jorna, L. B., Broekmans, M. E., Ploegmakers, H. P., Teeling, P., de Boer, O. J. and van der Wal, A. C. (2010). Anti-human vascular endothelial growth factor (VEGF) antibody selection for immunohistochemical staining of proliferating blood vessels. *Journal of Histochemistry and Cytochemistry*, 58(2), 109-118.
- Verheul, H.M. and Pinedo, H.M. (2011). The role of vascular endothelial growth factor (VEGF) in tumour angiogenesis and early clinical development of VEGF-receptor kinase inhibitors. *Clinical Breast Cancer*, 1, 80-84.
- Vietti, T. J., Sullivan, M. P., Berry, D. H., Haddy, T. B., Haggard, M. E. and Blattner, R. J. (1965). The response of acute childhood leukemia to an initial and a second course of prednisone. *The Journal of Pediatrics*, 66(1), 18-26.
- VV Popp, J. A. (1990). Fibrosis, Spleen, Rat. In *Hemopoietic System*, Springer Berlin Heidelberg, 243-245.
- Waddell, W.R. and Loughry, R.W. (1983). Sulindac for polyposis of the colon. *Journal of Surgical Oncology*, 24(1), 83-87.
- Waldron, H.A. (1966). The anemia of lead poisoning: A review. *British Journal of Industrial Medicine*, 23(2), 83-100.
- Wang, F. X., Zhang, X. J., Pan, L., Qiao, S. K., Guo, X. L. and Dong, Z. R. (2007). Angioimmunoblastic T-cell lymphoma with autoimmune hemolytic anemia and pure red cell aplasia. *Journal of Experimental Hematology/Chinese Association of Pathophysiology*, 15(4), 862-865.
- Wang, M. Y., Nowicki, D., Anderson, G., Jensen, J. and West, B. (2008). Liver protective effects of *Morinda citrifolia* (Noni). *Plant Foods for Human Nutrition*, 63(2), 59-63.
- Wang, M. Y., Peng, L., Anderson, G. and Nowicki, D. (2013). Breast cancer prevention with *Morinda citrifolia* (noni) at the initiation stage. *Functional Foods in Health and Disease*, 3(6), 203-222.
- Wang, M-Y., Anderson, G., Nowicki, D., Jensen, J. (2008). Hepatic protection by noni fruit juice against CCl₄ induced chronic liver damage n female SD rats. *Plants Food for Human Nutrition*, 63(3), 141-145.

- Wanikiat, P., Panthong, A., Sujayanon, P., Yoosook, C., Rossi, A.G. and Reutrakul, V. (2008). The anti-inflammatory effects and the inhibition of neutrophil responsiveness by *Barleria lupulina* and *Clinacanthus nutans* extracts. *Journal of Ethnopharmacology*, 5, 116(2), 234-244.
- West, B.J., Jensen, C.J. and Wasterndorf, J. (2006). Noni juice is not hepatotoxic. *World Journal of Gastroenterology*, 12(22), 3616-3619.
- West, B.J., Jensen, J.C., Pal, A.K. and Deng, S. (2011). Toxicity and antioxidant test of *Morinda citrifolia* (noni) seed extract. *Advanced Journal of Food and Technology*, 3(4), 303-307.
- Who, J., and Consultation, F. E. (2003). Diet, nutrition and the prevention of chronic diseases. *World Health Organization*, 916, 1-8.
- Williams, C. S., Mann, M. and DuBois, R.N. (1999). The role of cyclooxygenase in inflammation, cancer and development. *Oncogene*, 18, 7908-7916.
- Williams, L. (2004). Comprehensive review of hematopoiesis and immunology: Implications for hematopoietic stem cell recipients. *Hematopoietic Stem Cell Transplantation: A Manual for Nursing Practice*, 1-12.
- Winter, M. C. and Hancock, B. W. (2009). Ten years of rituximab in NHL. *Expert Opinion on Drug Safety*, 8(2), 223-235.
- Wolf, P. L. (1999). Biochemical diagnosis of liver disease. *Indian Journal of Clinical Biochemistry*, 14(1), 59-90.
- Wu, G., Luo, J., Rana, J. S., Laham, R., Sellke, F. W. and Li, J. (2006). Involvement of COX-2 in VEGF-induced angiogenesis via P38 and JNK pathways in vascular endothelial cells. *Cardiovascular Research*, 69(2), 512-519.
- Wun, T., Mcknight, H. and Tuscano, J.M. (2014). Increase cyclooxygenase (cox-2): a potential role in pathogenesis. *Leukemia Research*, 28(2), 179-190.
- Yang, J., Shikata, N., Mizuoka, H. and Tsubura, A. (1996). Colon carcinogenesis in shrews by intrarectal infusion of N-methyl-N-nitrosourea. *Cancer Letters*, 110 (1), 105-112.
- Yang, T. C., Chen, Y. J., Chang, S. F., Chen, C. H., Chang, P. Y. and Lu, S. C. (2014). Malondialdehyde mediates oxidized LDL-induced coronary toxicity through the Akt-FGF2 pathway via DNA methylation. *Journal of Biomedical Science*, 21(1), 1.
- Yang, T., Singh, I., Pham, H., Sun, D., Smart, A., Schnermann, J. B. and Briggs, J. P. (1998). Regulation of cyclooxygenase expression in the kidney by dietary salt intake. *American Journal of Physiology-Renal Physiology*, 274(3), 481-489.
- Yen, H. C., Oberley, T. D., Gairola, C. G., Szweda, L. I. and Clair, D. K. S. (1999). Manganese superoxide dismutase protects mitochondrial complex I against

adriamycin-induced cardiomyopathy in transgenic mice. *Archives of Biochemistry and Biophysics*, 362(1), 59-66.

Yin, H., Xu, L. and Porter, N. A. (2011). Free radical lipid peroxidation: mechanisms and analysis. *Chemical Reviews*, 111(10), 5944-5972.

Ying, Z. T., Zheng, W., Wang, X. P., Xie, Y., Tu, M. F., Lin, N. J., Ping, L.Y., liu, W.P., Deng, L.J., Zhang, C. and Zhu, J. (2012). The clinical features, therapeutic responses, and prognosis of the patients with mantle cell lymphoma. *Chinese Journal of Cancer*, 31(7), 348-353.

Yong, Y.K., Tan, J.J., Teh. S.S., Mah, S.H., Cheng, Lian, Ea, G., Chiong, H.S. and Ahmad, Z. (2013). *Clinacanthus nutans* extracts are antioioxidant with antiproliferative effect on cultured human cancer cell lines. *Evidence Based Complementary Alternaitive Medicine*, 8-12.

Yoshizawa, K., Oishi, Y., Nambu, H., Yamamoto, D., Yang, J., Senzaki, H., and Tsubura, A. (2000). Cataractogenesis in neonatal Sprague-Dawley rats by N-methyl-N-nitrosourea. *Toxicologic Pathology*, 28(4), 555-564.

Yuann, J-M, P., Wang, J-S., Jian, H-L., Lin, C-C. and Liang, J-Y. (2012). Effects of *Clinacanthus nutans* (Burm.f) Lindau leaf extracts on protection of plasmid DNA from riboflavin photoreaction. *MC-Transaction of Biotechnology*, 4(1), 45-58.

Yüce, B., Gülberg, V., Diebold, J. and Gerbes, A. L. (2006). Hepatitis induced by Noni juice from *Morinda citrifolia*: a rare cause of hepatotoxicity or the tip of the iceberg?. *Digestion*, 73(2-3), 167-170.

Yusnaini Md Yusoff. (2013). Prevention effects of *Curcuma longa* L. crude on MNU induced leukaemia/lymphoma Sprague Dawley rats via assessment of Bcl-2 to Bax ratio using real time PCR assay. Master Thesis. Universiti Putra Malaysia.

Zainal Ariffin, O. and Nor Saleha, I. T. (2011). Malaysia Cancer Statistics Data and Figure 2007. *Putrajaya: National Cancer Registry, Ministry of Health, Malaysia*.

Zedan, H., Abdel-Motaleb, A. A., Kassem, N. M. A., Hafeez, H. A. A. and Hussein, M. R. A. (2015). Low glutathione peroxidase activity levels in patients with vitiligo. *Journal of Cutaneous Medicine and Surgery*, 19(2), 144-148.

Zelenetz, A.D., Abrasom, J.S., Advani, R.J. and *et al.* (2011). Non-Hodgkin's lymphoma. *Journal of National Comprehensive Cancer Network*, 9, 484-560.