

# **UNIVERSITI PUTRA MALAYSIA**

WHITE RICE-BASED CARBOHYDRATE DIETS AS DIET-INDUCED OBESITY MODEL IN RATS

JEEVETHA SUBRAMANIAM

FPSK(M) 2014 28



# WHITE RICE-BASED CARBOHYDRATE DIETS AS DIET-INDUCED OBESITY MODEL IN RATS



By

JEEVETHA SUBRAMANIAM

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfillment of the Requirements for the Degree of Master of Science

February 2014

# COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

# WHITE RICE-BASED CARBOHYDRATE DIETS AS DIET-INDUCED OBESITY MODEL IN RATS

By

### JEEVETHA SUBRAMANIAM

#### February 2014

#### Chair: Barakatun Nisak binti Mohd Yusof, PhD Faculty: Medicine and Health Sciences

High fat intake is a typical dietary pattern that is commonly consumed by Western populations. This diet is widely used in diet-induced obesity studies using animal models. On the other hand, in most Asian countries including Malaysia, dietary carbohydrate particularly white rice is a staple diet. Nevertheless, study on dietinduced obesity using white rice is limited. Therefore, this study aimed to investigate whether the Experimental Diet i.e. White Rice-based Carbohydrate Diets (WRBCD) can be used as a diet-induced obesity model in rats. It was hypothesized that, WRBCD were effective as a Corn-Based High Fat Diet (CBHFD) to induce obesity in rats. This study was divided into two phases; Phase I was a pre-experimental study and Phase II was an experimental study. In Phase I, nutrient and amylose content of selected white rice; namely fragrant white (FWR) and white rice, 5% broken (WR5%) were determined. White rice with lower amylose content was used at Phase II to prepare WRBCD. In Phase II, a total of 40 rats were divided equally into (i) Control group (Normal Purified Rat Diet (PD) and High Fat Diet (HFD)) and (ii) Experimental group (Normal (NCHO<sub>Rice</sub>) and High Carbohydrate Rice Diet (HCHO<sub>Rice</sub>)). A total of 8 rats died due to cardiac puncture procedure after the acclimatization period. Thus, there were only 32 rats included in the 8 weeks study. Obesity in rats was measured using body weight and body composition, while metabolic parameters including fasting plasma glucose, triglyceride and insulin levels were assessed. Rats were sacrificed at the end of study to measure weight of heart, kidney, liver, abdominal fat and thigh fat. In this study, the WR5% contained comparable amount of ash, fat, protein and available carbohydrate with the FWR (p>0.05). FWR had comparable amylose content with WR5% (p>0.05). Both WR5% and FWR were categorized as low amylose and therefore, WR5% or FWR rice can be used for Phase II. At Phase II, all the baseline parameters were comparable in all groups. After 8 weeks period, HFD (n=10) had comparable body weight (466.4 ±32.3 g) with NCHO<sub>Rice</sub> (n=7) (457.4 ±42.9 g) and HCHO<sub>Rice</sub> (n=7) (450.4 ±48.7 g) (p>0.05). In terms of body composition of the rats, rats consumed NCHO<sub>Rice</sub> (78.4  $\pm 13.0$  g) had the highest while rats in PD (59.3  $\pm 4.4$  g) had the lowest fat mass; the differences were significant (p<0.05). Total mass (378.0 ±16.4 g), lean body mass  $(308.6 \pm 12.7 \text{ g})$  and bone mineral content  $(8.6 \pm 0.5 \text{ g})$  were significantly highest in



HFD (p<0.05) as compared to the other three groups. At the end of the study, weight of heart, kidney, liver and abdominal fat of the rats were comparable except for thigh fat. The thigh fat weight of the rats in HCHO<sub>Rice</sub> (6.8 ±2.2 g) was significantly higher than NCHO<sub>Rice</sub> (2.6 ±1.3 g) (p<0.05). For the metabolic parameters, all baseline data were comparable. The HCHO<sub>Rice</sub> group rats exhibited the highest fasting plasma glucose (16.9 ±2.4 mmol/L) and the highest fasting triglyceride (1.3 ±0.0 mmol/L) (p<0.05) than the other 3 groups at the end of the study. The insulin level of HFD (41.9 ±2.5 mU/L) was significantly higher than PD (33.9 ±1.5 mU/L) and NCHO<sub>Rice</sub> (36.3 ±1.1 mU/L) groups (p<0.05) after 8 weeks of the study. This study shows that WRBCD were effective as HFD to induce obesity in rats. Despite having comparable body weight in all groups, rats in WRBCD had produced highest blood glucose level (HCHO<sub>Rice</sub>) and highest fat mass (NCHO<sub>Rice</sub>) than the HFD. Thus, it can be concluded that WRBCD can be used as a diet-induced obesity model in rats.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

### DIET KARBOHIDRAT BERASASKAN NASI PUTIH SEBAGAI MODEL PERKEMBANGAN OBESITI PADA TIKUS

Oleh

#### JEEVETHA SUBRAMANIAM

#### Februari 2014

#### Pengerusi: Barakatun Nisak binti Mohd Yusof, PhD Fakulti: Perubatan dan Sains Kesihatan

Diet tinggi lemak adalah cara pemakanan lazim populasi Barat. Diet ini digunakan secara meluas sebagai model obesiti haiwan. Tetapi, karbohidrat terutamanya nasi putih ialah makanan ruji kebanyakkan negara Asia termasuk Malaysia. Walau bagaimanapun, kajian mengenai model obesiti yang menggunakan nasi putih adalah terhad. Oleh itu, kajian ini adalah untuk menentukan sama ada Diet Karbohidrat berasaskan Beras Putih (WRBCD) boleh digunakan sebagai model obesiti untuk tikus. Ia telah dihipotesis bahawa, WRBCD adalah berkesan seperti diet tinggi lemak berasaskan jagung untuk menyebabkan obesiti pada tikus. Kajian ini terbahagi kepada dua fasa: Fasa I adalah kajian pra-eksperimen dan Fasa II adalah kajian eksperimen. Dalam Fasa I, kandungan nutrien dan amilosa beras putih wangi (FWR) dan beras putih hancur 5% (WR5%) telah ditentukan. Beras putih dengan kandungan amilosa yang lebih rendah telah digunakan pada Fasa II bagi membentuk diet WRBCD. Sebanyak 40 ekor tikus telah dibahagikan sama rata kepada empat kumpulan, iaitu Normal Purified Rat Diet (PD), High Fat Diet (HFD), Normal Carbohydrate Rice Diet (NCHO<sub>Rice</sub>) dan High Carbohydrate Rice Diet (HCHO<sub>Rice</sub>). Lapan ekor tikus telah mati semasa proses pengambilan darah dengan teknik *cardiac* puncture. Justeru, jumlah tikus untuk pengajian ini ialah 32. Tempoh masa kajian ini adalah selama 8 minggu. Obesiti pada tikus telah ditentukan dengan mengukur berat dan komposisi tubuh, manakala parameter metabolik yang diukur adalah glukosa, trigliserida dan paras insulin. Tikus telah dikorbankan pada akhir kajian untuk mengira berat organ seperti jantung, buah pinggang, hati, lemak abdominal dan lemak paha. Jumlah abu, lemak, protein dan karbohidrat tersedia beras putih hancur 5% (WR5%) adalah setara dengan beras putih wangi (FWR) (p>0.05). Kandungan amilosa FWR adalah setara dengan WR5% (p>0.05). WR5% dan FWR dikategorikan sebagai rendah amilosa dan salah satu daripadanya boleh dipilih untuk fasa seterusnya. Pada Fasa II, semua parameter asal adalah setara dalam semua kumpulan. Pada akhir kajian, tikus yang diberi makan dengan HFD (n=10) mempunyai berat badan yang setara (466.4 ±32.3 g) dengan tikus diberi makan dengan NCHO<sub>Rice</sub> (n=7) (457.4 ±42.9 g) dan tikus yang diberi makan dengan HCHO<sub>Rice</sub> (n=7) (450.4 ±48.7 g) (p>0.05). Bagi jisim lemak yang terenap, tikus dalam kumpulan NCHO<sub>Rice</sub> (78.4 ±13.0 g) mempunyai bacaan tertinggi manakala tikus dalam kumpulan PD (59.3 ±4.4 g) mempunyai lemak terenap paling rendah;



perbezaan adalah signifikan (p <0.05). Jumlah jisim (378.0  $\pm$ 6.4 g), jisim badan tanpa lemak (308.6  $\pm$ 12.7 g) dan kandungan mineral tulang (8.6  $\pm$  0.5g) adalah lebih tinggi dalam kumpulan HFD (p <0.05). Pada akhir kajian, berat organ seperti jantung, buah pinggang, hati dan perut lemak tikus adalah setara kecuali lemak bahagian paha. Berat lemak paha bagi kumpulan HCHO<sub>Rice</sub> (6.8 ±2.2 g) adalah jauh lebih tinggi daripada kumpulan NCHO<sub>Rice</sub> (2.6 ±1.3 g) (p <0.05). Pada permulaannya, semua parameter metabolik adalah sama bagi semua kumpulan. Tikus kumpulan HCHO<sub>Rice</sub> menunjukkan bacaan yang tertinggi bagi plasma glukosa berpuasa (16.9 ±2.4 mmol/L) dan trigliserida berpuasa (1.3 ±0.0 mmol/L) (p <0.05). Tahap insulin bagi kumpulan HFD (41.9 ±2.5 mU/L) adalah lebih tinggi daripada kumpulan PD (33.9  $\pm 1.5$  mU/L) dan kumpulan NCHO<sub>Rice</sub> (36.3  $\pm 1.1$  mU/L) (p <0.05) selepas 8 minggu kajian. Kajian ini menunjukkan WRBCD boleh menyebabkan obesiti setara dengan diet lemak tinggi pada tikus. Walaupun memiliki berat badan yang setara, tikus dalam kumpulan WRBCD mempunyai tahap glukosa darah yang tertinggi (HCHO<sub>Rice</sub>) dan jisim lemak tertinggi (NCHO<sub>RICE</sub>) berbanding dengan diet tinggi lemak. Kesimpulannya, WRBCD boleh digunakan sebagai model obesiti yang disebabkan oleh diet pada tikus.

#### ACKNOWLEDGEMENTS

First and foremost, I would like to express my deep appreciation to my main supervisor, Dr Barakatun Nisak binti Mohd Yusof for her endless guidance, patience, support, motivation and advice throughout my study. I specially thank my cosupervisors Professor Dr Amin bin Ismail and Associate Professor Dr Azrina binti Azlan for their motivation, support, advice, suggestions and comments. I would like to express my gratitude to Associate Professor Dr Goh Yong Meng from Faculty of Veterinary Medicine, Universiti Putra Malaysia for his guidance on study design and animal care.

I also would like to convey my thankfulness to the lab staffs of Department of Nutrition and Dietetics of Faculty of Medicine and Health Sciences, UPM especially, Mr Syed Hasbullah Syed Kamaruddin and Mr Simon Nawawi for their help and guidance throughout my study. I would also like to thank Mr Ramli, staff of Animal House, UPM for guiding me in rats handling methods. I would like to thank Mrs Rabizah, lab technician at Endocrine Unit, Universiti Kebangsaan Malaysia Medical Center (UKMMC) for assisting in the process of DEXA scan.

The highest gratitude appreciations to my beloved parents, Mr Subramaniam Sinnappan and Mrs Kogila Krishnan for their support, patience, love and care throughout this endeavor and over the years. Deepest appreciation goes to all my friends and lab mates for their encouragement and constant help throughout this journey.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science in Nutritional Sciences. The members of the Supervisory Committee were as follows:

#### Barakatun Nisak binti Mohd Yusof, PhD

Senior Lecturer Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Chairman)

#### Amin bin Ismail, PhD

Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Member)

### Azrina binti Azlan, PhD

Associate Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Member)

#### **BUJANG BIN KIM HUAT, PhD**

Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date:

# **DECLARATION**

### **Declaration by graduate student**

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature:	Date:	
0		

Name and Matric No.: Jeevetha Subramaniam (GS 27733)

# **Declaration by Members of Supervisory Committee**

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_\_ Name of Chairman of Supervisory Committee : DR BARAKATUN NISAK BINTI MOHD YUSOF

UPM

Signature: \_\_\_\_\_\_ Name of Member of Supervisory Committee : PROF. DR AMIN BIN ISMAIL

Signature: \_\_\_\_\_\_ Name of Member of Supervisory Committee : ASSOC. PROF. DR AZRINA BINTI AZLAN

# TABLE OF CONTENTS

		Page
ABSTRACT		ii
ABSTRAK		iv
ACKNOWLEDGEMENTS		
APPROVAL		VII in
LIST OF TA	ABLES	1X xiv
LIST OF FI	GURES	XV
LIST OF AI	BREVIATIONS	xvi
CHAPTER		
1	INTRODUCTION	1
2	LITERATURE REVIEW	5
	2.1 Obesity prevalence	5
	2.1.1 Obesity health-related issues	6
	2.1.2 Etiology of obesity	8
	2.2 Assessment of obesity	8
	2.2.1 Assessment of obesity in rats	9
	2.3 Obesity induction models in animals	11
	2.3.1 Diet-induced obesity in rats	12
	2.3.2 Dietary carbohydrate to induce obesity in rats	15
	2.4 Types of animal models used in obesity induction studies	17
	2.4.1 Rats	17
	2.5 White rice-based diet and impact to health	17
	2.6 Diet-induced obesity effect on metabolic parameters	18
3	METHODOLOGY	20
	3.1 Scope of study	20
	3.2 Phase I: Pre-experimental study	20
	3.2.1 Chemicals and reagents	21
	3.2.2 Sample selection	21
	3.2.3 Sample preparation	21

5.2.4 Sample analyses	22
3.2.4.1 Determination of nutritional composition	22
3.2.4.2 Amylose content determination and calculation	24
3.3 Phase II: Experimental study	25
3.3.1 Chemicals and reagents	25
3.3.2 Preparation of rats	26
3.3.3 Animal care	26
3.3.4 Study design	26
3.3.5 Diet preparation	28
3.3.5.1 Control diets	28
3.3.5.2 Experimental diets	31
3.3.6 Study outcome measurements	32
3.3.6.1 Primary outcome measurements	32
3.3.6.2 Secondary outcome measurements	35
3.3.6.3 Food intake	37
3.4 Statistical analysis	37
4 <b>RESULTS</b>	38
4.1 Phase I: Pre-experimental study	38
4.1.1 Proximate of white rice	38
4.1.2 Amylose content analysis	39
4.2 Phase II: Experimental study	
1.2 Thuse II. Experimental study	39
4.2.1 Changes in body weight and body composition	39 39
4.2.1 Changes in body weight and body composition 4.2.2 Changes in metabolic parameters	39 39 44
4.2.1 Changes in body weight and body composition 4.2.2 Changes in metabolic parameters 4.2.2.1 Changes in fasting plasma glucose	39 39 44 44
4.2.1 Changes in body weight and body composition 4.2.2 Changes in metabolic parameters 4.2.2.1 Changes in fasting plasma glucose 4.2.2.2 Changes in fasting triglyceride	39 39 44 44 46
4.2.1 Changes in body weight and body composition 4.2.2 Changes in metabolic parameters 4.2.2.1 Changes in fasting plasma glucose 4.2.2.2 Changes in fasting triglyceride 4.2.2.3 Changes in fasting insulin	<ol> <li>39</li> <li>39</li> <li>44</li> <li>44</li> <li>46</li> <li>47</li> </ol>
4.2.1 Changes in body weight and body composition 4.2.2 Changes in metabolic parameters 4.2.2.1 Changes in fasting plasma glucose 4.2.2.2 Changes in fasting triglyceride 4.2.2.3 Changes in fasting insulin 4.2.3 Food and energy intake	<ol> <li>39</li> <li>39</li> <li>44</li> <li>44</li> <li>46</li> <li>47</li> <li>50</li> </ol>
4.2.1 Changes in body weight and body composition 4.2.2 Changes in metabolic parameters 4.2.2.1 Changes in fasting plasma glucose 4.2.2.2 Changes in fasting triglyceride 4.2.2.3 Changes in fasting insulin 4.2.3 Food and energy intake 5 DISCUSSION	<ul> <li>39</li> <li>39</li> <li>44</li> <li>44</li> <li>46</li> <li>47</li> <li>50</li> <li>52</li> </ul>
4.2.1 Changes in body weight and body composition 4.2.2 Changes in metabolic parameters 4.2.2.1 Changes in fasting plasma glucose 4.2.2.2 Changes in fasting triglyceride 4.2.2.3 Changes in fasting insulin 4.2.3 Food and energy intake 5 DISCUSSION 5.1. Phase I: Pre-experimental study	<ul> <li>39</li> <li>39</li> <li>44</li> <li>44</li> <li>46</li> <li>47</li> <li>50</li> <li>52</li> <li>52</li> </ul>
<ul> <li>4.2.1 Changes in body weight and body composition</li> <li>4.2.2 Changes in metabolic parameters</li> <li>4.2.2.1 Changes in fasting plasma glucose</li> <li>4.2.2.2 Changes in fasting triglyceride</li> <li>4.2.2.3 Changes in fasting insulin</li> <li>4.2.3 Food and energy intake</li> </ul> 5 DISCUSSION 5.1. Phase I: Pre-experimental study 5.2. Phase II: Experimental study	<ul> <li>39</li> <li>39</li> <li>44</li> <li>46</li> <li>47</li> <li>50</li> <li>52</li> <li>52</li> <li>52</li> <li>53</li> </ul>
<ul> <li>4.2.1 Changes in body weight and body composition</li> <li>4.2.2 Changes in metabolic parameters</li> <li>4.2.2.1 Changes in fasting plasma glucose</li> <li>4.2.2.2 Changes in fasting triglyceride</li> <li>4.2.2.3 Changes in fasting insulin</li> <li>4.2.3 Food and energy intake</li> </ul> 5 DISCUSSION 5.1. Phase I: Pre-experimental study 5.2. Phase II: Experimental study 5.2.1 Changes in body weight and body composition	<ul> <li>39</li> <li>39</li> <li>44</li> <li>46</li> <li>47</li> <li>50</li> <li>52</li> <li>52</li> <li>53</li> <li>54</li> </ul>

xii

6	CONCLUSION, LIMITATION AND RECOMMEND	ATION 62
	6.1 Conclusion	62
	6.2 Limitations	63
	6.2 Recommendations	63

REFERENCES APPENDEICES BIODATA OF STUDENT 64

61

77 80

# LIST OF TABLES

	Table		Page
	2.1	Classification of overweight and obesity	8
	2.2	Advantages and disadvantages between carcass composition analyses and Dual X-Ray Absorptiometry (DEXA) in rats	11
	2.3	High Fat Diet Induced Obesity in Rats	14
	2.4	High Carbohydrate Diet Induced Obesity in Rats	16
	3.1	Amylose content (%) category	25
	3.2	Energy composition of the diets for each group (per 1000g)	29
	3.3	Diet ingredients for preparing the diets for each group (per 1000g)	30
	3.4	Time point of measurements	33
	4.1	Proximate analysis of the tested white rice	38
	4.2	Amylose content value of the tested white rice	39
	4.3	Body weight of the rats	40
	4.4	Lee Obesity Index of the rats	42
	4.5	Body composition of the rats at baseline	42
	4.6	Body composition of the rats at the end of 8 week study	43
	4.7	Weight of Organs	44
	4.8	Glucose level of experimental rats	45
	4.9	Triglyceride level of the rats	46
	4.10	Insulin level of the rats	48
	4.11	HOMAIR level of the rats	49
	5.1	Comparative Analysis of Amylose Content vs. GI of Rice	53
	5.2	Comparative Analysis of High Fat vs. High Carbohydrate Diets to Induce Obesity in Rats	55

# LIST OF FIGURES

Figure		
1.1	Conceptual framework of study	4
2.1	Comparison between Asian countries on prevalence of obesity	5
2.2	Prevalence of obesity in Malaysia	6
2.3	Health issues related with obesity	7
2.4	Prevalence of diabetes mellitus type 2 in Malaysia	7
2.5	Comparison between BMI and DEXA	9
2.6	Mechanism on upper body obesity leading to hyperglycemia and impaired glucose tolerance	19
3.1	Flowchart on phases of the study	20
3.2	Study Design	27
4.1	Changes in rats' body weight throughout the feeding period	41
4.2	Body weight gain of the rats in each group	41
4.3	Changes in glucose level of the rats	45
4.4	Changes in triglyceride level of experimental rats	47
4.5	Changes in insulin level of experimental rats	48
4.6	HOMA <sub>IR</sub> changes of experimental rats	49
4.7	Food intake of rats over 8 weeks of study	50
4.8	Energy intake of the rats over 8 weeks of study	51
5.1	Glucose hepatic metabolisms	59

# LIST OF ABBREVIATIONS

ACUC	Animal Care and Use Committee
ANOVA	Analysis of Variance
BMI	Body Mass Index
С	Control
CBHFD	Corn-Based High Fat Diet
СНО	Carbohydrate
CV	Coefficient Variation
DEXA	Dual Energy X-ray Absorptiometry
EDTA	Ethylenediaminetetraacetic acid
FWR	Fragrant White Rice
GI	Glycemic Index
HFCS	High Fructose Corn Syrup
HFD	High Fat Diet
HOMA-IR	Homeostasis Model Assessment-Estimated Insulin Resistance
HRP	Horseradish Peroxidase
HSD	High Sucrose Diet
ISO	International Organization for Standardization
kcal	kilocalories
mg	milligram
mmol/L	millimol per liter
mU/L	milliU per liter
NHLBI	The National Heart, Lung, and Blood Institute
NHMS	The National Health and Morbidity Survey

NIH	National Institutes of Health
Nm	nanometer
NR	Not Recorded
NS	Not Significant
OECD	Organization for Economic Co-operation and Development
PD	Purified Diet
RPM	Revolutions Per Minute
SD	Standard Deviation
UKMMC	Universiti Kebangsaan Malaysia Medical Center
VMH	Ventromedial Hypothalamic Nucleus
WHO	World Health Organization
WR5%	White Rice, 5% Broken
WRBCD	White Rice-Based Carbohydrate Diets

C

# **CHAPTER 1**

# INTRODUCTION

# 1.1 Background

Obesity is a substantial public health crisis around the world. It represents a state of excess storage of body fat that has an adverse effect on health. The prevalence of obesity is growing rapidly worldwide, particularly in developing countries (WHO, 2011). Malaysia has not been spared from this epidemic. The prevalence of obesity has increased rapidly by 175% over a period of 15 years between 1996 and 2011 (NHMS II, 1996; NHMS IV, 2011).

Obesity increases the likelihood of various chronic diseases particularly type 2 diabetes. This is because obesity is not solely related to the excess storage of body fat, but it also coexisted with other metabolic abnormalities including worsening of insulin resistance and elevating triglycerides level (Must et al., 1999). It comes as no surprise that the sharp increase in obesity prevalence has also contributed to the escalating prevalence of type 2 diabetes, especially in Asia, where it accounts for 60% of the world's' diabetic population (Hu et al., 2011). In Malaysia, diabetes prevalence has also increased by a staggering 83% within a period of 15 years between 1996 and 2011 (NHMS II, 1996; NHMS IV, 2011).

Obesity is a result from an imbalance between energy intake and energy expenditure (Johnson et al., 2008). In general, relative excessive intake of energy particularly dietary fat has been a major cause of dietary obesity (Swinburn et al., 2004). The relationship between excess fat consumption and obesity development has been supported by diet-induced-obesity in animal studies (Bray et al., 2004; Buettner et al., 2007). In addition, a high fat diet has an adverse effect on fat deposition especially at visceral areas and promoting insulin resistance (Wilcox et al., 2005).

 $\bigcirc$ 

Since diets high in fat are usually energy dense and palatable, it usually leads to an increase in energy intake (Rolls, 2009). In addition, it has been proposed that a high fat diet has appeared more efficiently to be stored as body fat resulting in a greater increase in adipose mass than iso-energetic portions of high carbohydrate or high protein diets (Brand-Miller et al., 2002).

High fat intake is a typical dietary pattern that is commonly consumed by Western populations. It is absolutely an undeniable fact that dietary fat consumption has increased over the years in Asian countries which could possibly explain the tremendous obesity rate increment. A similar trend has also been observed in Malaysia, where the total fat consumption of Malaysian adults derived from energy has increased by 4% from 23 to 27% within a decade (Kandiah et al., 2008). Nonetheless, the fat consumption pattern is not as high as compared to Western countries and it was still below the 30% limit recommended by the American Heart Association (Wylie-Rosett, 2002).

The low fat intake in Asian diets including in Malaysia will relatively increase dietary carbohydrates (Kandiah et al., 2008). In addition, Asian populations have a different dietary pattern which is mainly based on carbohydrate foods, especially rice. Rice, particularly in the form of white rice remains the major staple food and is unlikely to be displaced by any other staples in Asian countries. Recently, there has been considerable renewed interest on the significant positive association between rice consumption and risk of obesity (Kim et al., 2012) and type 2 diabetes (Nanri et al., 2010; Zhang et al., 2010; Hu et al., 2012) especially among Asian populations. This could be due to the high glycemic index of the polished white rice than those of the brown and whole grain rice (Foster-Powell, 2002).

The glycemic index (GI) is used to classify dietary carbohydrates based on their impact on the blood glucose response 2-hours after meals (Wolever, 2006). Foods that are high in GI such as white rice may result in a rapid rate of digestion and absorption, hence inducing rapid elevation of postprandial hyperglycemia as well as insulin concentration (Radulian et al., 2009). These characters of high GI foods can increase appetite and promote over consumption along with imbalance in energy partitioning, relatively increasing the fat storage over the long term (Brand-Miller et al., 2002; Hao et al., 2012). Thus, habitual consumption of high carbohydrate-high GI foods may initiate sequence of metabolic events that promote fat deposition and deteriorated metabolic parameters (Mc Neel and Mersmann, 2005; Claessens et al., 2009).

# **1.2** Problem statements

In Malaysia, white rice is consumed by almost 97% of the population with approximately  $2\frac{1}{2}$  plates (~ 450 g) daily (Norimah et al., 2008). The consumption of ~ 450 g white rice in a day is considered excessive in relation to diabetes risk as indicated in the Shanghai Women's Health Study, where women who consumed more than 300 g white rice/day had a 78% greater risk of developing type 2 diabetes mellitus compared to women who consumed less than 200 g white rice/day (Villegas et al., 2007). A traditional Japanese diet, with a high intake of white rice, has also shown to be significantly associated with an increased risk of obesity among Japanese women aged 18 to 20 years (Okubo et al., 2008). Similarly, in a traditional Korean diet, which includes white rice and kimchi was independently also associated

with increasing risk of obesity among Korean adults (Kim et al., 2012). There was also a significant relationship between rice intake and total and central obesity among Hispanic population (Lin et al., 2003).

Nevertheless, data relating the intake of white rice and its association with obesity and diabetes risk among Malaysian adults is scarce. As dietary compliance has always been an important bias in human studies and is it beyond the control of the investigator, the use of animal models is highly sought to investigate specific dietinduced obesity that primarily mimics the Asian-based human diet. Although studies using high carbohydrate diet in inducing obesity have been well conducted in animal models, most studies used either corn or sucrose as the primary source of carbohydrate (McNeel et al., 2005; Zhang et al., 2007; Claessens et al., 2009; Bocarsly et al., 2010).

The present study was conducted to identify whether the white rice-based carbohydrate diets can be used as a diet-induced obesity model in rats. The white rice-based carbohydrate diets were designed to mimic Malaysian food consumption pattern in terms of energy proportion from carbohydrates.

### **1.3** Significance of the study

The findings from this study can provide fundamental data in developing an appropriate dietary obesity induction model for animal study that represents a Malaysian-style diet using white rice-based diets. The effect induced by white rice to promote obesity and diabetes in animals can be translated to the human model in preventing obesity and diabetes development. Essentially, the diets are planned to resemble Malaysian trends in food intake and the results of the study would benefit in terms of improvised public health system.

# 1.4 Study objectives

The general objective of this study was to investigate whether the White Rice-based Carbohydrate Diets (WRBCD) can be used as a diet-induced obesity model in rats.

The specific objectives of the study were as follows:

1. To determine the nutritional composition, amylose content and amylose: amylopectin ratio of the two types of white rice.

- 2. To determine the effects of white rice-based carbohydrate diet on body weight and composition (body composition measured by DEXA and weight of organs) and metabolic parameters (fasting glucose, triglycerides, and insulin) of the rats after 8 weeks of study.
- 3. To compare the effects of the white rice-based carbohydrate diets against cornbased control diets, High Fat Diet and the Normal Purified Rat Diet on body weight gain and metabolic parameter in the rats after 8 weeks of study.

# **1.5** Study hypothesis

It was hypothesized that the White Rice-based Carbohydrate Diets (WRBCD) can induce obesity as effective as High Fat Diet.

#### 1.6 Conceptual framework

The etiology of obesity is based on increased energy intake and reduced physical activity. The increase in energy due to macronutrients, specifically fat and carbohydrate can lead to obesity; and carbohydrate can alter metabolic parameters. Those inside the dotted lines of conceptual frame work, Figure 1.1 are interest of current study.





Boxes inside the dotted lines are interest of current study

#### REFERENCES

Aekplakorn, W., Chaiyapong, Y., Neal, B., Chariyalertsak, S., Kunanusont, C., Phoolcharoen, W., Suriyawongpaisal, P., 2004, Prevalence and determinants of overweight and obesity in Thai adults: results of the Second National Health Examination Survey, *Journal of Medical Association of Thailand*, 87(6): pp. 685-93

Araujo-Farro, P.C., Amaral-Sobral, J.P., Menegalli, F.C., Comparison of starch pasting and retrogradation properties of Quinoa (*chenopodium quinoa* willd), rice, potato, cassava, wheat and corn starches, *ENPROMER*: 2nd Mercosur Congress on Chemical Engineering; 4th Mercosur Congress on Process Systems Engineering, 2005

Axen, K.V. and Kenneth, A., 2006, Very low-carbohydrate versus isocaloric highcarbohydrate diet in dietary obese rats, *Obesity*, 14: 1344–1352

Baynes, J.W. and Thorpe, S.R., 1999, Role of oxidative stress in diabetic complications: A new perspective on an old paradigm, *Diabetes*, 48: pp. 1–9

Bechmann, L.P., Hannivoort R.A., Gerken, G., Hotamisligil, G.S., Trauner, M., Canbay, A., 2012, The interaction of hepatic lipid and glucose metabolism in liver diseases, *Journal of Hepatology*, Volume 56(Issue 4): pp. 952-964

Behall, K.M. and Hallfrisch, J., 2002, Plasma glucose and insulin reduction after consumption of breads varying in amylose content, *European Journal of Clinical Nutrition*, 56: 913-920

Benmoussa, M., Moldenhauer, K.A.K. and Hamaker, B.R., 2007, Rice Amylopectin Fine Structure Variability Affects Starch Digestion Properties, *Journal of Agricultural Food Chemistry*, 55 (4): pp 1475–1479

Bernardis, L.L. and Patterson, B.D., 1968, Correlation between 'lee index' and carcass fat content in weanling and adult female rats with hypothalamic lesions, *Journal ofEndocrinology*, 40: pp. 527-528

Bernardis, L.L., 1970, Prediction of carcass fat, water and lean body mass from Lee's nutritive ratio in rats with hypothalamic obesity, *Experientia*, 26: 789–90

Bernardis, L.L., 1970, Sex and age differences in hyperthermia response to ether anesthesia in weanling rats with ventromedial hypothalamic lesions, *Experientia*, 26: pp. 1322-1324

Bernfeld, P., 1955, In amylase,  $\alpha$  and  $\beta$ , Chapter 17, Methods in Enzymology, pp. 149-158, New York: Academic Press

Berthoud, H.R., Lenard N.R. and ShinA.C., 2011, Food reward, hyperphagia, and obesity, *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, 300: R1266–R1277



Bessesen, D.H., 2001, The Role of Carbohydrates in Insulin Resistance, *Journal of Nutrition*, 131: pp. 2782S–2786S

Blake, G. M. and Fogelman, I., 2007, The role of DXA bone density scans in the diagnosis and treatment of osteoporosis, *Postgraduate Medical Journal*, 83: pp. 509 - 517

Bocarsly, M. E., Powell, E. S., Avena, N. M. and Hoebel, B. G., 2010, High-fructose corn syrup causes characteristics of obesity in rats: increased body weight, body fat and triglyceride levels, *Pharmacology Biochemistry and Behavior*, Vol.97(No.1): pp. 101–106

Brand-Miller, J., Foster-Powell, K., Wolever, T.M.S. and Calgiuri, S., 2003, *The new glucose revolution: The authoritative guide to the glycemic index*, New York: Marlowe & Company

Brand-Miller, J., Pang, E. and Bramall, L., 1992, Rice: a high or low glycemic index food, *American Journal of Clinical Nutrition*, 56: pp. 1034-6.

Brand-Miller, J.C., Holt, S.H.A., Pawlak, D.B. and McMillan, J., 2002, Glycemic index and obesity, *American Journal of Clinical Nutrition*, July: Vol. 76(No.1): pp. 281S-285S

Bray, G.A. and Popkin, B.M., 1998, Dietary fat intake does affect obesity, *American Journal of Clinical Nutrition*, 68: pp. 1157–73

Bray, G.A., 2004, Medical Consequences of Obesity, *Journal of Clinical Endocrinology Metabolism*, 89(6): pp. 2583–2589

Bucolo, G. and David, H., 1973, Quantitative determination of serum triglycerides by the use of enzymes, *Clinical Chemistry*, 19(5): 476-482

Buettner, R., Scholmerich, J., Bollheimer, L.C., 2007, High-fat diets: Modeling the metabolic disorders of human obesity in rodents, *Obesity*, 15: pp. 798-808

Byrnes, S.E., Brand-Miller, J.C. and Denver, G.S., 1995, Amylopectin Starch Promotes the Development ofInsulin Resistance in Rats, *Journal of Nutrition*, 125: pp. 1430-1437

Calpe, C., *Rice: International Commodity Profile*, Food and Agriculture Organization of the United Nations-Markets and Trade Division, 2007

Cameron, D.K., Wang, Y-J., Moldenhauer, K.A., 2007, Comparison of Starch Physicochemical Properties from Medium-Grain Rice Cultivars Grown in California and Arkansas, *Starch*, 59: pp. 600–608

Cao, L., Liu, X., Cao, H., Lv, Q. and Tong, N., 2012, Modified High-Sucrose Diet-Induced Abdominally Obese and Normal-Weight Rats Developed High Plasma Free Fatty Acid and Insulin Resistance, *Oxidative Medicine and Cellular Longevity*, Volume 2012: Article ID 374346

 $\bigcirc$ 

Capdevila, S., Giral, M., J L Ruiz de la Torre, Russell, R.J. and Kramer, K., 2007, Acclimatization of rats after ground transportation to a new animal facility, *Laboratory Animals*, 41: pp. 255-261

Carroll, J., 1970, A colorimetric serum glucose determination using hexokinase and glucose-6-phosphate dehydrogenase, *Biochemistry Medicine*, 4: pp. 171-180

Chinnaswamy, R. and Hanna, M.A., 1988, Relationship between Amylose Content and Extrusion-Expansion Properties of Corn Starches, *Cereal Chemistry*, 65(2): pp. 138-143

Claessens, M., van-Baak, M.A., Monsheimer, S., Saris, W.H., 2009, The effect of a low-fat, high-protein or high-carbohydrate ad libitum diet on weight loss maintenance and metabolic risk factors, *International Journal of Obesity*, March; 33(3): pp. 296-304

Clegg, K.M., 1956, The application of the anthrone reagent to the estimation of starch in cereals, *Journal of the Science of Food and Agriculture*, Volume 7, Issue 1, pp. 40–44

Cook, C.B., Shawar, L., Thompson, H. and Prasad, C., 1997, Caloric Intake and Weight Gain of Rats Depends on Endogenous Fat Preference, *Physiology & Behavior*, Vol.61 (No.5): pp. 743–748

Corbett, S.W., Stern, J.S., Keesey, R.E., 1986, Energy expenditure in rats with dietinduced obesity, *American Journal of Clinical Nutrition*, 44: pp. 173–80

Delvalle, D., Dumez, S., Wattebled, F., Planchot, I.R.V., Berbezy, P., Colonna, P., Vyas, D., Chatterjee, M., Ball, S., A' ngel Me' rida and D'Hulst, C., 2005, Soluble starch synthase I: a major determinant for the synthesis of amylopectin in Arabidopsis thaliana leaves, *The Plant Journal*, 43: pp. 398–412

Denardin, C.C., Boufleur, N., Reckziegel, P., Silva, L.P., Walter, M., 2012, Amylose content in rice (*Oryza sativa*) affects performance, glycemic and lipidic metabolism in rats, *Ciência Rural*, Vol.42 (No.2): pp.381-387

Despre's, J.P., Lemieux, I., Bergeron, J., Pibarot, P., Mathieu, P., Larose, E., Cabau, J.R., Bertrand, O.F., Poirier, P., 2008, Abdominal Obesity and the Metabolic Syndrome: Contribution to Global Cardiometabolic Risk, *Arteriosclerosis Thrombosis Vascular and Biology*, 28: pp. 1039-1049

Diaz, V.A., Mainous, A.G.III, Koopman, R.J., Geesey, M.E., 2004, Undiagnosed obesity: implications for undiagnosed hypertension, diabetes, and hypercholesterolemia, *Family Medicine*, 36(9): pp. 639-44

Diemen, V.V., Trindade, E.N, Trindade, M.B.M., 2006, Experimental model to induce obesity in rats, *Acta Cirúrgica Brasileira*, Vol. 21: No. 6

Egan, H., Kirk, R.S. and Sawyer, R., 1981(a), *Chemical Analysis of Food:* 8<sup>th</sup> *Edition*, pp. 8-11, Churchill Livingstone, London: Pearson



Egan, H., Kirk, R.S. and Sawyer, R., 1981(b), *Chemical Analysis of Food:* 8<sup>th</sup> *Edition*, pp. 11-12, Churchill Livingstone, London: Pearson

Egan, H., Kirk, R.S. and Sawyer, R., 1981(c), *Chemical Analysis of Food:* 8<sup>th</sup> *Edition*, pp. 15-20, Churchill Livingstone, London: Pearson

Ellis, K.J., 2000, Human Body Composition: In Vivo Methods, *Physiological Reviews*, Vol. 80: No.2

Epidemiology & Disease Control Division, *National Health Survey*, 2010, Singapore Ministry of Health,

Farley, C., Cook, J.A., Spar, B.D., Austin, T.M. and Kowalski, T.J., 2003, Meal pattern analysis of diet-induced obesity in susceptible and resistant rats, *Obesity Research*, 11: pp. 845–851

Fossati, P. and Prencipe, L., 1982, Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide, *Clinical Chemistry*, 28(10): pp. 2077-2080

Foster-Powell, K., Holt, S.H.A. and Brand-Miller, J.C., 2002, International table of glycemic index and glycemic loadvalues: 2002, *American Journal of Clinical Nutrition*, 76: pp. 5–56.

Frei, M., Siddhuraju, P. and Becker, K., 2003, Studies on the in vitro starch digestibility and the GI of six different indigenous rice cultivars from the Philippines, *Food Chemistry*, 83: pp. 395-402

Fukuchi, S., Hamaguchi, K., Seike, M., Himeno, K., Sakata, T. and Yoshimatsu, H., 2004, Role of Fatty Acid Composition in the Development of Metabolic Disorders in Sucrose-Induced Obese Rats, *Experimental Biology and Medicine* 2004, 229: pp. 486-493

Gauthier, M.S., Couturier, K., Latour, J.G. and Lavoie, J.M., 2003, Concurrent exercise prevents high-fat-diet-induced macrovesicular hepatic steatosis, *Journal of Applied Physiology*, 94: pp. 2127–2134

Ghibaudi, L., Cook, J., Farley, C., Heek, M., and Hwa, J.J., 2002, Fat intake affects adiposity, comorbidity factors, and energy metabolism of Sprague-Dawley rats. *Obesity Research*, 10: pp. 956–963

Granfeldt, Y., Drews, A. and Bjorck, I., 1995, Arepas Made from High Amylose Corn Flour Produce Favorably Low Glucose and Insulin Responses in Healthy Humans, *Journal of Nutrition*, 125: pp. 459-465

Gross, L.S., Li, L., Ford, E.S. and Liu, S., 2004, Increased consumption of refined carbohydrates and the epidemic of type 2 diabetes in the United States: an ecologic assessment, *American Journal of Clinical Nutrition*, 79: pp. 774–9



Ha, S.K., Kim, J. and Chae, C., 2011, Role of AMP-activated Protein Kinase and Adiponectin during Development of Hepatic Steatosis in High-fat Diet-induced Obesity in Rats, *Journal of Comparative Pathology*, Vol.145: 88e94

Hao, Q., Lillefosse, H.H., Fjaere, E., Myrmel, L.S., Midtbø, L.K., Jarlsby, R.H., Ma, T., Jia, B., Petersen, R.K., Sonne, S.B., Chwalibog, A., Frøyland, L., Liaset, B., Kristiansen, K. and Madsen L., 2012, High-glycemic index carbohydrates abrogate the anti-obesity effect of fish oil in mice, *American Journal of Physiology: Endocrinology and Metabolism*, May 15; 302(9): E1097-112.

Hariri, N. and Thibault, L., 2010, High-Fat Diet-Induced Obesity in Animal Models, *Nutrition Research Reviews*, 23: pp. 270-299

Hariri, N., Gougeon, R. and Thibault, L., 2010, A highly saturated fat-rich diet is more obesogenic than diets with lower saturated fat content, *Nutrition Research*, 30: pp. 632-643

Haslam, D.W. and James, W.P., 2005, Obesity, Lancet, 366: pp. 1197-1209

Hernan, M.A., Hernandez-Diaz, S. and Robins, J.M., 2004, A Structural Approach to Selection Bias, *Epidemiology*, 15: pp. 615–625

Honors, MA, Davenport, BM and Kinzig, KP, 2009, Effects of consuming a high carbohydrate diet after eight weeks of exposure to a ketogenic diet, *Nutrition & Metabolism*, Vol.6: No.46

Hoover, R. and Ratnayake, W.S., 2001, Current Protocols in Food Analytical Chemistry, *Starch and Starch Derivatives*, E2.3.1-E2.3.5

Hsu, I. R., Kim, S. P., Kabir, M. and Bergman, R. N, 2007 Metabolic syndrome, hyperinsulinemia, and cancer, *American Journal of Clinical Nutrition*, 86(suppl.): pp. 867S–71S

Hu, E.A., Pan, A., Malik, V. and Sun, Q., 2012, White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review, *British Medical Journal*, 344: e1454

Hu, F.B., 2010, Are refined carbohydrates worse than saturated fat? *American Journal of Clinical Nutrition*, 91: pp. 1541–2

Hu, F.B., 2011, Globalization of Diabetes: The role of diet, lifestyle, and genes, *Diabetes Care*, 34: pp. 1249–1257

Hu, P., Zhao, H., Duan, Z., Linlin, Z. and Wu, D., 2004, Starch digestibility and the estimated glycemic score of different types of rice differing in amylose contents, *Journal of Cereal Science*, 40: pp. 231-237

Huang, X.F., Liu, Y., Rahardjo, G.L., Mclennan, P.L., Tapsell, L.C. and Buttemer, W.A., 2008, Effects of diets high in whey, soy, red meat and milk protein on body

weight maintenance in diet induced obesity in mice, *Nutrition & Dietetics*,65 (Suppl. 3): pp. S53–S59

Imam, M.U., Musa, S.N.A., Azmi N.H. and Ismail M., 2012, Effects of White Rice, Brown Rice and Germinated Brown Rice on Antioxidant Status of Type 2 Diabetic Rats,*International Journal of Molecular Science*, 13: pp. 12952-12969

Institute for Public Health. *The Fourth National Health and Morbidity Survey* (*NHMS IV*) 2011, Ministry of Health, Malaysia,

Institute for Public Health. *The Second National Health and Morbidity Survey* (*NHMS II*) 1996, Ministry of Health, Malaysia.

Institute for Public Health. *The Third National Health and Morbidity Survey (NHMS III) 2006*, Ministry of Health, Malaysia.

Jalil, A.M.M., Ismail, A., Pei, C.P., Hamid, M. and Kamaruddin, S.H.S., 2008, Effects of Cocoa Extract on Glucometabolism, Oxidative Stress and Antioxidant Enzymes in Obese-Diabetic (Ob db) Rats, *Journal of Agricultural Food Chemistry*, 56: pp. 7877–7884

Johnson, L., Mander, A.P., Jones, L.R., Emmett, P.M. and Jebb S.A, 2008, Energydense, low-fiber, high-fat dietary pattern is associated with increased fatness in childhood, *American Journal of Clinical Nutrition*, 87: pp. 846–54

Johnson, P.D. and Besselsen, D.G., 2002, Practical Aspects of Experimental Design in Animal Research, *ILAR Journal*, Volume 43: Number 4

Joslyn, M.A., 1970, Methods in Food Analysis: 2<sup>nd</sup> Edition, pp. 109-121

Juan, G., Luis, A. and David, B., 2006, Isolation and molecular characterization of Makal (*Xanthosoma yucatanensis*) starch, *Starch*, 58: pp. 300-307

Juliano, B.O., 1971, A simplified assay for milled rice amylose, *Cereal Science Today*, 16: pp. 334-338

Juliano, B.O., Perez, C.M. and Blakeney, A.B., 1981, International Cooperative testing on the amylose content of milled rice, *Starch*,33(5): pp. 157-162

Juliano, B. O., Tuano, A. P. P., Monteroso, D. N., Aoki, N., Mestres, C., Duldulao, J. B. A., Bergonio, K. B., 2012, Replacement of Acetate with Ammonium Buffer to Determine Apparent Amylose Content of Milled Rice, *Cereal Foods World*, January-February, Vol. 57 (1)

Jung, T., 2008, Gaining Weight: How Media Exposure, Socioeconomic Status, Sedentary Lifestyle and Dietary Habits Work Together – A Secondary Analysis of a Nationally Representative Study, *The Open Communication Journal*, 2: pp. 67-73

Kahn, S.E., Prigeon, R.L., Mcculloch, D.K., Boyko, E.J., Bergman, R.N., Schwartz, M.W., Neifing, J.L, Ward, W.K., Beard, J.C., Palmer, J.P. and Porte-Jr, D., 1993,



Quantification of the Relationship between Insulin Sensitivity and p-Cell Function in Human Subjects: Evidence for a Hyperbolic Function, *Diabetes*, 42: pp. 1663-72

Kanarek, R.B. and Markskaufman, R., 1979, Developmental aspects of sucrose induced obesity in rats, *Physiology Behavior*, 23: pp. 881-855

Kanasaki, K. and Koya, D., 2011, Biology of Obesity: Lessons from Animal Models of Obesity, *Journal of Biomedicine and Biotechnology*, Volume 2011, Article ID 197636

Kandiah, M., Zalilah, M.S., Safiah, M.Y., Tahir, A., Siti Haslinda, M.D., Siti Rohana, D., Khairul Zarina, M.Y., Mohd Hasyami, S. and Normah, H., 2008, Energy and Nutrient Intakes: Findings from the Malaysian Adult Nutrition Survey (MANS), *Malaysia Journal of Nutrition*, 14(1): pp. 1-24

Kapoor, B. and Bhattacharya, M., 2001, Steady shear and transient properties of starch in dimethylsulfoxide, *Carbohydrate Polymers*, Vol.44 (3): pp. 217-231

Kim, J., Inho, Jo., Hyojee, Joung., 2012, A Rice-Based Traditional Dietary Pattern Is Associated with Obesity in Korean Adults, *Journal of the Academy of Nutrition and Dietetic*, 112: pp. 246-253

Klaus, S., 2005, Increasing the Protein: Carbohydrate Ratio in a High-Fat Diet Delays the Development of Adiposity and Improves Glucose Homeostasis in Mice, *Journal of Nutrition*, 135: pp. 1854–1858

Langendonk, J.G., Kok, P., Frolich, M., Pijl, H., Meinders, A.E., 2006 Decrease in visceral fat following diet-induced weight loss in upper body compared to lower body obese premenopausal women, *European Journal of International Medicine*, 17: pp. 465–469

Lee M.O., 1929, Determination of the surface area of the white rat with its application to the expression of metabolic results, *American Journal of Physiology*, 89: pp. 24-33.

Li, S., Zhang, H.Y., Hu, C.C., Lawrence, F., Gallagher, K.E., Surapaneni, A., Estrem, S.T., Calley, J.N., Varga, G., Dow, E.R. and Chen Y., 2008, Assessment of Diet-induced Obese Rats as an Obesity Model by Comparative Functional Genomics, *Obesity*, 16: pp. 811–818

Lin, H., Bermudez, O.I. and Tucker, K.L., 2003, Dietary Patterns of Hispanic Elders Are Associated with Acculturation and Obesity, *Journal of Nutrition*, 133: pp. 3651– 3657

Lomba, A., Martínez, J.A., García-Díaz, D.F., Paternain, L., Marti, A., Campión, J., Milagro, F.I., 2010, Weight gain induced by an isocaloric pair-fed high fat diet: A nutriepigenetic study on FASN and NDUFB6 gene promoters, *Molecular Genetics and Metabolism*, 101: pp. 273–278



Lomba, A., Milagro, F.I., García-Díaz, D.F., Campión, J., Marzo, F., Martínez, J.A., 2009, A High-Sucrose Isocaloric Pair-Fed Model Induces Obesity and Impairs NDUFB6 Gene Function in Rat Adipose Tissue, *Journal of Nutrigenetics and Nutrigenomics*, 2: pp. 267–272

London, E., Lala, G., Berger, R., Panzenbeck, A., Kohli, A.A., Renner, M., Jackson, A., Raynor, T., Loya, K. and Castonguay, T.W., 2007, Sucrose Access Differentially Modifies 11b-Hydroxysteroid Dehydrogenase-1 and Hexose-6-Phosphate Dehydrogenase Message in Liver and Adipose Tissue in Rats, *The Journal of Nutrition: Biochemical, Molecular, and Genetic Mechanisms*, 137: pp. 2616–2621

Lukaski, H.C., Hall, C.B., Marchello, M.J. and Siders, W.A.,2001, Validation of Dual X-Ray Absorptiometry for Body-Composition Assessment of Rats Exposed to Dietary Stressors, *Nutrition*, 17: pp. 607–613

Maggio, C.A. and Pi-Sunyer, F.X., 1997, The prevention and treatment of obesity: Application to type 2 diabetes, *Diabetes Care*, 20: pp. 1744-66

Maghraoui A. E. and Roux C., 2008, DXA scanning in clinical practice, *QJM: An international journal of medicine*, 101: pp. 605–617

Maritim, A.C., Sanders, R.A., Watkins, J.B.3<sup>rd</sup>, 2003, Diabetes, oxidative stress, and antioxidants: a review, *Journal of Biochemical and Molecular Toxicology*, 17(1): pp. 24-38

Matthews, D.R., Hosker, J.P., Rudenski, A.S., Naylor, B.A., Treacher, D.F., Turner, R.C., 1985, Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man, Diabetologia, 28: pp. 412–419.

McGowan, M.W., Artiss, J.D., Strandbergh, D.R. et al, 1983, A peroxidase-coupled method for the colorimetric determination of serum triglycerides, *Clinical Chemistry*, 29(3): pp. 538-542

McGrance, S.J., Cornell, H.J. and Rix, C.J., 1998, A simple and rapid colorimetric method for the determination of amylose in starch products, *Starch*, 50: pp. 158-163

McNeel, R.L. and Mersmann, H.J., 2005, Low- and High-Carbohydrate Diets: Body Composition Differences in Rats, *Obesity Research*, 13: pp. 1651–1660

Mendez, A.J., Cabeza, C. and Hsia, S.L., 1986, A fluorometric method for the determination of triglycerides in nanomolar quantities, *Analytical Biochemistry*, 156: pp. 386-389

Mercer, J.G. and Archer, Z.A., 2005, Diet-induced obesity in the Sprague-Dawley rat: dietary manipulations and their effect on hypothalamic neuropeptide energy balance systems, *Biochemical Society Transactions*, 33: pp. 1068–1072

Mickelsen O, Takahashi S, Craig C., 1955, Experimental obesity: Production of obesity in rats by feeding high-fat diets, *Journal of Nutrition*, 57: pp. 541-554

Mikkelsen, R., Baunsgaard, L. and Blennow, A., 2003, Functional characterization of α-glucan, water dikinase, the starch phosphorylating enzyme, *Biochemical Journal Immediate Publication*, BJ20030999

Miller, C.N., Kauffman, T.G., Cooney, P.T., Ramseur, K.R., Brown, L.M., 2011, *Physiology & Behavior*, Comparison of DEXA and QMR for assessing fat and lean body mass in adult rats. 103: pp. 117–121

Moussavi, N., Gavino, V., Receveur, O., 2008, Could the quality of dietary fat, and not just its quantity, be related to risk of obesity? *Obesity*, 16: pp: 7-15

Musa, A.S.N., Umar, I.M. and Ismail, M., 2011, Physicochemical properties of germinated brown rice (*Oryza sativa* L.) starch, *African Journal of Biotechnology*, Vol. 10(33): pp. 6281-6291

Must, A., Spadano, J., Coackley, E.H., Field, A.E., Colditz, G. and Dietz, W.H., 1999, The Disease Burden associated with Overweight and Obesity, *The Journal of the American Medical Association*, Vol. 282: No. 16

Nanri, A., Mizoue, T., Noda, M., Takahashi, Y., Kato, M., Inoue, M. and Tsugane, S., 2010, Rice intake and type 2 diabetes in Japanese men and women: The Japan Public Health Center-based Prospective Study, *American Journal of Clinical Nutrition*, 92: pp. 1468–77

National Heart, Lung, and Blood Institute (NHLBI), Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: The Evidence Report, National Institutes of Health (NIH) Publications, 1998

Ngan, H. B., N., Barakatun-Nisak, M.Y., Ismail, A., Faez-Bachok, M. and Jeevetha, S., 2012, Glycemic index values of selected commercialized rice in Malaysia. Poster presentation: Group C: Nutrients and Other Components in Food/Products, abstract No C07, Nutrition Society of Malaysia (NSM) 27<sup>th</sup> Conference, Malaysia, http://www.nutriweb.org.my/downloads/Program%20and%20abstract%20combined. pdf. Accessed on 7 November 2012

Nik-Shanita, S., Hasnah, H and Khoo, C.W., 2011, Amylose and Amylopectin in Selected Malaysian Foods and its Relationship to Glycemic Index, *Sains Malaysiana*, 40(8): pp. 865–870

Nilsson, C., Raun, K., Yan, F., O Larsen, M., Tang-Christensen, M., 2012, Laboratory animals as surrogate models of human obesity, *Acta Pharmacologica Sinica*, 33: pp. 173–181

Norazmir, M.N. and Ayub, M.Y., 2010, Beneficial Lipid-Lowering Effects of Pink Guava Puree in High Fat Diet Induced-Obese Rats, *Malaysian Journal of Nutrition*, 16(1): pp. 171 - 185

Norimah, A.K., Safiah, M., Jamal, K., Haslinda, S., Zuhaida, H., Rohida, S., Fatimah, S., Norazlin, S., Poh, B.K., Kandiah, M., Zalilah, M.S., Wan-Manan, W.M., Fatimah, S. and Azmi, M.Y., 2008, Food Consumption Patterns: Findings from the

Malaysian Adult Nutrition Survey (MANS), *Malaysian Journal of Nutrition*, 14(1): pp. 25 – 39

Ofei, F., 2005, Obesity-A preventable disease, *Ghana Medical Journal*, Volume 39: Number 3

Official methods of analysis: 17<sup>th</sup> Edition, 2000, Association of Official Analytical Chemists (AOAC) International, Gaithersburg, Maryland

Oh, K., Hu, F.B., Cho, E., Rexrode, K.M., Stampfer, M.J., Manson, J.E., Liu, S. and Willett, W.C., 2005, Carbohydrate Intake, Glycemic Index, Glycemic Load, and Dietary Fiber in Relation to Risk of Stroke in Women, *American Journal of Epidemiology*, Vol. 161: No. 2

Okubo, H., Sasaki, S., Murakami, K., Kim, M.K., Takahashi, Y., Hosoi, Y., Itabashi, M. and the Freshmen in Dietetic Courses Study II group, 2008, Three major dietary patterns are all independently related to the risk of obesity among 3760 Japanese women aged 18–20 years, *International Journal of Obesity*, 32: pp. 541–549

Organization for Economic Co-operation and Development (OECD), 2011, *Health Data*, Version November 2011

Parks, E. J., 2001, Effect of Dietary Carbohydrate on Triglyceride Metabolism in Humans, *Journal of Nutrition*, 131: pp. 2772S–2774S

Pawlak, D.B., Bryson, J.M., Denyer, G.S. and Brand-Miller J.C., 2001, High Glycemic Index Starch Promotes Hypersecretion of Insulin and Higher Body Fat in Rats without Affecting Insulin Sensitivity, *The Journal of Nutrition*, 131: pp. 99–104

Pi-Sunyer, F.X., 2002, The obesity epidemic: pathophysiology and consequences of obesity, *Obesity Research*, 10: pp. 97S–104S

Pi-Sunyer, F.X., 2009, The medical risks of obesity, *Postgraduate Medicine*, Nov; 121(6): pp. 21-33

Popkin,B.M., 2008, The Nutrition Transition and Obesity in the Developing World, *Journal of Nutrition*, 131: pp. 871S–873S

Porter, S.A., Massaro, J.M., Hoffmann, U., Vasan, R.S., O'Donnel, C.J. and Fox, C.S., 2009, Abdominal Subcutaneous Adipose Tissue: A Protective Fat Depot, *Diabetes Care: American Diabetes Association*, Vol.32 (No.6): pp. 1068-1075

Prunet-Marcassus, B., Desbazeille, M., Bros, A., Louhce, K., Delagrange, P., Renard, P., Casteilla, L. and Nicaud, L., 2003, Melatonin reduces body weight gain in Sprague Dawlwy rats with diet-induced obesity, *Endocrinology*, 144: pp. 5347-52

Radulian, G., Rusu, E., Dragomir, A. and Posea, M., 2009, Metabolic effects of low glycemic index diets, *Nutrition Journal*, Vol. 8: No. 5



Ramachandran, A. and Snehalatha, C., 2010, Rising Burden of Obesity in Asia, *Journal of Obesity*, Article ID 868573

Rogers, P. and Webb, G.P., 1980, Estimation of body fat in normal and obese mice. *British Journal of Nutrition*, 43: pp. 83-86

Rolls. B.J., 2009, The relationship between dietary energy density and energy intake, *Physiology Behavior*, 97(5): pp. 609-15

Rossmeisl, M., Rim, J.S., Koza, R.A. and Kozak, L.P., 2003, Variation in type 2 diabetes-related traits in mouse strains susceptible to diet-induced obesity, *Diabetes*, 52: pp. 1958-1966

Ryu, M.H. and Cha, Y.S., 2003, The Effects of a High-fat or High-sucrose Diet on Serum Lipid Profiles, Hepatic Acyl-CoA Synthetase, Carnitine Palmitoyltransferase-I, and the Acetyl-CoA Carboxylase mRNA Levels in Rats, *Journal of Biochemistry and Molecular Biology*, Vol.36(No.3): pp. 312-318

Sampey, B.P., Vanhoose, A.M., Winfield, H.M., Freemerman, A.J., Muehlbauer, M.J., Fueger, P.T., Newgard, C.B., Makowski, L., 2011, Cafeteria diet is a robust model of human metabolic syndrome with liver and adipose inflammation: comparison to high-fat diet. *Obesity (Silver Spring)*, 19: pp. 1109–1117

Saris, W.H.M., 2003, Sugars, energy metabolism, and body weight control, *American Journal of Clinical Nutrition*, 78(suppl.): pp. 850S–7S

Sclafani, A., 1987, Carbohydrate taste, appetite, obesity: an overview, *Neuroscience Biobehavior Review*, 11: pp. 131–153

Sclafani, A., 1991, Starch and sugar tastes in rodents: an update, *Brain Research Bulletin*, Sep-Oct; 27(3-4): pp. 383-6.

Self, R., 2005, Extraction of Organic Analytes from Food: A Manual of Methods, Cambridge: Royal Society of Chemistry

Shiwaku, K., Anuurad, E., Enkhmaa, B., Kitajima, K. and Yamane, Y., 2004, Appropriate BMI for Asian populations, *The Lancet*, March: Vol. 363

Sichieri, R., Moura, A.S., Genelhu, V., Hu, F. and Willett, W.C., 2007, An 18-mo randomized trial of a low glycemic-index diet and weight change in Brazilian women, *American Journal of Clinical Nutrition*, 86: pp. 707-713

Sompong, R., Siebenhandl-Ehn, S., Berghofer, E. and Schoenlechner, E., 2011, Extrusion cooking properties of white and colored rice varieties with different amylose content, *Starch*, 63: pp. 55–63

Son, B.K., Kim, J.Y., Lee, S.S., 2008, Effect of Adlay, Buckwheat and Barley on Lipid Metabolism and Aorta Histopathology in Rats Fed an Obesogenic Diet, *Annals of Nutrition and Metabolism*, 52: pp. 181–187



Stubbs, R.J., Mazlan, N. and Whybrow, S., 2001, Carbohydrates, Appetite and Feeding Behavior in Humans, *Journal of Nutrition*, 131: pp. 2775S–2781S

Sugiura, M. and Hirano, K., 1977, A new colorimetric method for determination of serum glucose, *Clinca Chimca Acta*, 75: pp. 387-391

Suzuki, H., Goshi, H. and Sugisawa, H., 1975, Effects of Previous Feeding of a High Carbohydrate or a High Fat Diet on Changes in Body Weight and Body Composition of Fasted Rats, *Journal of Nutrition*, 105: pp. 90-95

Swinburn, B.A., Caterson, I., Seidell, J.C. and James W.P.T., 2004, Diet, nutrition and the prevention of excess weight gain and obesity, *Public Health Nutrition*, 7(1A): pp. 123–146

Taraschenko, O.D., Maisonneuve, I.M., Glick, S.D., 2011, Sex differences in high fatinduced obesity in rats: Effects of 18 methoxycoronaridine, *Physiology and Behavior*, 103: pp. 308–314

Tataranni, P.A., 2002, Pathophysiology of obesity-induced insulin resistance and type 2 diabetes mellitus, *European Review for Medical and Pharmacological Sciences*, 6: pp. 27-32

Tecator, A.B., Soxtec System HT2 Manual, 1982

Vasanthan, T., 2001, Overview of Laboratory Isolation of Starch from Plant Materials, *Current Protocols in Food Analytical Chemistry*, pp. E2.1.1 - E2.1.6

Vento, P.J., Swartz, M.E., Martin, L.B.E. and Daniels, D., 2008, Food Intake in Laboratory Rats Provided Standard and Fenbendazole-supplemented Diets, *Journal of the American Association for Laboratory Animal Science*, Vol 47(No 6): pp. 46-50

Villegas, R., Liu, S., Gao, Y.T., Yang, G., Li, H., Zheng, W. and Shu, X.O., 2007, Prospective Study of Dietary Carbohydrates, Glycemic Index, Glycemic Load, and Incidence of Type 2 Diabetes Mellitus in Middle-aged Chinese Women, *JAMA Network: Archives of International Medicine*, 167(21): pp. 2310-2316

Wannamethee, S.G., Shaper, A.G., Walker M., 2005, Overweight and obesity and weight change in middle aged men: impact on cardiovascular disease and diabetes, *Journal of Epidemiology and Community Health*, 59: pp. 134–139

West, D. B. and York B., 1998, Dietary fat, genetic predisposition, and obesity: lessons from animal models, *American Journal of Clinical Nutrition*, 67(suppl): 505S–12S

Westerterp, K.R., 2002, Weight Loss and Bone Mineral Content, *Obesity Research*, June; Vol.10: No.6

Wilcox, G., 2005, Insulin and Insulin Resistance, Clinical Biochemistry Review, May: Vol. 26

 $\bigcirc$ 

Willett, W., Manson, J. and Liu, S., 2002, Glycemic index, glycemic load, and risk of type 2 diabetes, *American Journal of Clinical Nutrition*, 76(suppl.): pp. 274S–80S

Williams, S., *Official Methods of Analysis: 14th Edition*, Association of Official Analytical Chemists (AOAC), Virginia, USA, 1984

Wolden-Hanson, T., 2010, Changes in Body Composition in Response to Challenges during Aging in Rats, *Interdisciplinary Topics in Gerontology*, Vol.37: pp. 64–83

Wolever, T.M.S., *The glycemic index: A physiological classification of dietary carbohydrate*, CABI: United Kingdom, 2006

World Health Organization (WHO), 1998, Obesity: preventing and managing the global epidemic. Report of a WHO consultation of obesity, Geneva: 1997, *Obesity Research*, 6: pp. 51S–210S

World Health Organization (WHO), *Fact sheet: obesity and overweight*, 2011 www.who.int/mediacentre/factsheets/fs311/en/. Accessed 3<sup>rd</sup> December 2011

World Health Organization (WHO), *World Health Statistics*, Department of Health Statistics and Information Systems of Innovation, Information, Evidence and Research Cluster, 2012

World Health Organization, *Report of a Joint FAO/WHO Ad Hoc Expert Committee on Energy and Protein Requirements*; WHO Geneva, Report Series No.522, 1973

Wylie-Rosett, J., 2002, Fat Substitutes and Health: An Advisory from the Nutrition Committee of the American Heart Association (AHA), *Circulation: AHA*,105: pp. 2800-2804

Yang, W., Lu J., Weng, J., Jia, W., Ji, L., Xiao, J., Shan, Z., Liu, J., Tian, H., Ji, Q., Zhu, D., Ge, J., Lin, L., Chen, L., Guo, X., Zhao, Z., Li, Q., Zhou, Z., Shan, G. and He, J., 2010, Prevalence of Diabetes among Men and Women in China, *The New England Journal of Medicine*, March; 362: 12

Zaman, M. Q., Leray, V., Bloc'h, J. L., Thorin, C., Ouguerram, K. and Nguyen, P., 2011, Lipid profile and insulin sensitivity in rats fed with high-fat or high-fructose diets, *British Journal of Nutrition*, 106: pp. S206–S210

Zhang, G., Malik, V.S., Pan, A., Kumar, S., Holmes, M.D., Spiegelman, D., Lin, X. and Hu, F.B., 2010, Substituting Brown Rice for White Rice to Lower Diabetes Risk: A Focus-Group Study in Chinese Adults, *Journal of the American Dietetic Association*, 110: pp. 1216-1221

Zhang, X., Hua, J., Wang, S. and Sun, C., 2007, Post-weaning isocaloric hypersoybean oil versus a hyper-carbohydrate diet reduces obesity in adult rats induced by a high-fat diet, *Asia Pacific Journal of Clinical Nutrition*, 16 (Suppl. 1): pp. 368-373

