



UNIVERSITI PUTRA MALAYSIA

***PHASE-FITTED AND AMPLIFICATION-FITTED RUNGE-KUTTA TYPE
METHODS FOR SOLVING LINEAR DIFFERENTIAL EQUATIONS
WITH OSCILLATORY SOLUTIONS***

MOHAMMED MAHMOOD SALIH

FS 2016 4



**PHASE-FITTED AND AMPLIFICATION-FITTED RUNGE-KUTTA TYPE
METHODS FOR SOLVING LINEAR DIFFERENTIAL EQUATIONS
WITH OSCILLATORY SOLUTIONS**

By

MOHAMMED MAHMOOD SALIH

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

August 2016

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree of Doctor of Philosophy

PHASE-FITTED AND AMPLIFICATION-FITTED RUNGE-KUTTA TYPE METHODS FOR SOLVING LINEAR DIFFERENTIAL EQUATIONS WITH OSCILLATORY SOLUTIONS

By

MOHAMMED MAHMOOD SALIH

August 2016

Chairman : Professor Fudziah Binti Ismail, PhD
Faculty : Science

New phase-fitted and amplification-fitted Runge-Kutta methods (RK) based on the existing Runge-Kutta methods of order four and five were derived to solve second order ordinary differential equations with oscillatory solutions. The new method has the property of zero phase-lag and zero dissipation. The effects of phase-fitted and amplification-fitted relations are tested over a large interval on homogeneous and non-homogeneous problems which have oscillatory solutions and the numerical results proved that the new methods are more accurate compared to the existing methods. Then, the first order Fuzzy Differential Equations (FDEs) are solved using the RK methods with phase-fitted and amplification-fitted and the numerical results show that the methods are more accurate than the existing methods.

The technique of phase-fitted and amplification-fitted is then extended to diagonally implicit Runge-Kutta methods (DIRK) for solving second order ordinary differential equations (ODEs) with oscillatory solution. We derived the phase-fitted and amplification-fitted fourth DIRK method based on the fourth order existing DIRK methods. Numerical results show that the DIRK with phase-fitted and amplification-fitted is more accurate and efficient for solving oscillatory problems.

In the next part of the thesis, we derived the order conditions of Runge-Kutta Nystrom method purposely for solving linear second order ordinary differential equations. Based on the order conditions we derived the new fifth order four-stage and sixth order five-stage explicit Runge-Kutta-Nyström methods for linear ordinary differential equations (LODEs). Then the methods are phase-fitted and amplification-fitted so that they will have zero-dispersion and zero-dissipation. The fifth order four-stage RKN method for LODEs has the property of First Same As Last (FSAL). Numerical results proved that the methods with phase-fitted and amplification-fitted are much more efficient than the existing methods with the same algebraic order.

Next we used the RKN methods for solving Hyperbolic partial differential equations (PDEs) that is the second order wave equations. We very well know that the second order PDEs can be converted to second order linear ODEs using the methods of lines. Thus we applied the RKN methods for LODEs to solve the resulting second order linear ODEs. Numerical results show that the RKN methods for LODEs are accurate and reliable for solving the second order wave equation.

As a conclusion, in this thesis, we have derived phase-fitted and amplification-fitted RK and RKN methods for solving first and second order oscillatory problems. The phase-fitted and amplification-fitted RK method is also applied to first order fuzzy differential equations (FDEs). The non fitted RKN method for LODEs is also used for solving hyperbolic partial differential equations. Numerical results show that all the methods are more accurate than the existing methods in the scientific literature.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**SUAI-FASA DAN SUAI-AMPLIFIKASI JENIS KAEDAH RUNGE-KUTTA
UNTUK MENYELESAIKAN MASALAH BERAYUN BAGI
PERSAMAAN PEMBEZAAN LINEAR**

Oleh

MOHAMMED MAHMOOD SALIH

Ogos 2016

Pengerusi : Profesor Fudziah Binti Ismail, PhD
Fakulti : Sains

Kaedah Baru Runge-Kutta (RK) yang di suai-fasa dan suai-amplifikasi berdasarkan kaedah Runge-Kutta yang sedia ada peringkat keempat dan kelima diterbitkan untuk menyelesaikan persamaan pembezaan biasa dengan penyelesaian berayun. Kaedah baru ini mempunyai sifat serakan sifar dan lesapan sifar. Kesan hubungan suai-fasa dan suai-amplifikasi diuji untuk selang yang lebih besar keatas masalah homogen dan tak homogen yang mempunyai penyelesaian berayun dan keputusan berangka membuktikan bahawa kaedah baru tersebut adalah lebih tepat berbanding dengan kaedah yang sedia ada. Kemudian, Persamaan Pembezaan Kabur (PPK) Peringkat pertama diselesaikan dengan menggunakan kaedah RK dengan suai-fasa dan suai-amplifikasi tersebut dan keputusan berangka menunjukkan bahawa kaedah ini adalah lebih tepat daripada kaedah yang sedia ada.

Seterusnya teknik suai-fasa dan suai-amplifikasi diperluaskan pula kepada kaedah Runge-Kutta Pepenjuru tersirat (RKPT) untuk menyelesaikan persamaan pembezaan biasa (PPB) peringkat pertama dengan penyelesaian berayun. Kami terbitkan kaedah suai-fasa dan suai-amplifikasi berdasarkan kaedah RKPT peringkat empat yang sedia ada. Keputusan berangka menunjukkan bahawa kaedah RKPT dengan suai-fasa dan suai-amplifikasi adalah lebih tepat untuk menyelesaikan masalah berayun.

Dalam bahagian seterusnya tesis ini, kami terbitkan syarat peringkat kaedah Runge-Kutta Nystrom khas untuk menyelesaikan persamaan pembezaan biasa linear peringkat kedua. Berdasarkan syarat peringkat tersebut, kami terbitkan kaedah tak tersirat Runge-Kutta-Nystrom baharu peringkat kelima tahap-empat dan peringkat keenam tahap-lima untuk menyelesaikan persamaan pembezaan biasa linear (PPBL) peringkat kedua. Kemudian kaedah ini telah di suai-fasa dan suai-implifikasi agar ianya mempunyai serakan sifar dan lesapan sifar. Kaedah RKN peringkat kelima tahap-empat untuk PPBL ini mempunyai ciri yang pertama sama dengan yang terakhir (PSDA). Keputusan berangka membuktikan bahawa kaedah baharu dengan suai-fasa

dan suai-implifikasi ini adalah lebih cekap berbanding kaedah sedia ada dengan peringkat aljabar yang sama.

Seterusnya kami menggunakan kaedah RKN yang telah diterbitkan untuk menyelesaikan persamaan pembezaan separa (PPS) hiperbolik iaitu persamaan gelombang peringkat kedua. Kita sedia ketahui bahawa persamaan pembezaan separa peringkat kedua boleh diubah kepada persamaan pembezaan biasa linear peringkat kedua dengan menggunakan kaedah garis. Oleh itu kami gunakan kaedah RKN untuk PPBL bagi menyelesaikan persamaan pembezaan yang terhasil. Keputusan berangka menunjukkan bahawa kaedah RKN untuk PPBL adalah lebih jitu bagi menyelesaikan persamaan gelombang peringkat kedua.

Kesimpulannya, dalam tesis ini, kami telah menerbitkan kaedah RK dan RKN dengan suai-fasa dan suai-amplifikasi untuk menyelesaikan masalah berayun peringkat pertama dan kedua masing-masingnya. Kaedah RK dengan suai-fasa dan suai-amplifikasi juga digunakan untuk menyelesaikan persamaan pembezaan kabur peringkat pertama. Kaedah RKN tanpa suai-fasa untuk PPBL juga digunakan untuk menyelesaikan persamaan pembezaan separa hiperbola. Keputusan berangka menunjukkan bahawa semua kaedah ini adalah lebih jitu dari kaedah sedia ada dalam literatur saintifik.

ACKNOWLEDGEMENTS

First and foremost, I would like to show my deepest appreciation and gratitude to the Chairman of the Supervisor Committee, Professor Dr. Fudziah Binti Ismail for her invaluable assistance, advice and guidance throughout the duration of the studies. I also wish to express my sincere thank to Yang Berbahagia Professor Dato Dr. Mohamed Bin Suleiman and Dr. Norazak Senu for their guidance and assistance.

Great appreciation goes to my beloved family especially my parents for their conditional love, support and understanding throughout the course of my research. Last but not least, special thanks to all my friends and also colleagues who kindly provided valuable and helpful comments during my studies.



I certify that a Thesis Examination Committee has met on 16 August 2016 to conduct the final examination of Mohammed Mahmood Salih on his thesis entitled "Phase-Fitted and Amplification-Fitted Runge-Kutta Type Methods for Solving Linear Differential Equations with Oscillatory Solutions" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Mohamad Rushdan bin Md Said, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Zarina Bibi binti Ibrahim, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Leong Wah June, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Abduvali Khaldjigitov, PhD

Professor
National University of Uzbekistan
Uzbekistan
(External Examiner)



NOR AINI AB. SHUKOR, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 3 November 2016

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Fudziah Binti Ismail, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Mohamed Bin Suleiman, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Norazak Bin Senu, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No: Mohammed Mahmood Salih (GS34265)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Professor Dr. Fudziah Binti Ismail

Signature: _____
Name of Member
of Supervisory
Committee: Professor Dr. Mohamed Bin Suleiman

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Norazak Bin Senu

TABLE OF CONTENTS

		Page
ABSTRACT		i
ABSTRAK		iii
ACKNOWLEDGEMENTS		v
APROVAL		vi
DECLARATION		viii
LIST OF TABLES		xiii
LIST OF FIGURES		xvi
LIST OF ABBREVIATIONS		xix
CHAPTER		
1	INTRODUCTION	1
	1.1 Ordinary Differential Equations	1
	1.2 Fuzzy Differential Equations	3
	1.3 Partial Differential Equations	5
	1.4 Runge-Kutta (RK) methods	6
	1.4.1 Algebraic Condition for RK method	7
	1.4.2 Local Truncation Error for RK method	9
	1.4.3 Analysis of Absolute Stability for RK method	10
	1.5 Runge-Kutta-Nyström (RKN) methods	10
	1.5.1 Algebraic Condition for RKN method	11
	1.5.2 Local Truncation Error for RKN method	12
	1.6 Objectives of the Thesis	14
	1.7 Outline of Thesis	14
	1.8 Problem Statement	16
	1.9 Scope of Study	16
 2	LITERATURE REVIEW	 17
	2.1 Introduction	17
	2.2 Phase-Lag and Dissipation Errors	17
	2.3 Fuzzy Differential Equations	17
	2.4 Diagonally Implicit Runge-Kutta (DIRK) Methods	18
	2.5 Runge-Kutta-Nyström (RKN) Methods	18
	2.6 Hyperbolic Partial Differential Equations	18
 3	PHASE-FITTED AND AMPLIFICATION-FITTED EXPLICIT RUNGE-KUTTA METHODS FOR SOLVING OSCILLATORY PROBLEMS	 20
	3.1 Introduction	20
	3.1.1 Analysis of Dispersion (Phase-lag) and Dissipation (Amplification error) for RK method	20
	3.1.2 Analysis of Phase-fitted and Amplification-fitted for RK method	21
	3.2 Derivation of Phase-fitted and Amplification-fitted for Fourth Order Classical Runge-Kutta Method	21
	3.2.1 Problems Tested	23

	3.2.2 Numerical Results	25
	3.2.3 Discussion	35
	3.3 Derivation of Phase-fitted and Amplification-fitted for Fifth order Runge-Kutta Method for Linear Ordinary Differential Equations (LODEs)	35
	3.3.1 Numerical Results	37
	3.3.2 Discussion	46
	3.4 Conclusion	46
4	SOLVING FIRST ORDER FUZZY DIFFERENTIAL EQUATIONS USING PHASE-FITTED AND AMPLIFICATION-FITTED EXPLICIT RUNGE-KUTTA METHODS	47
	4.1 Introduction	47
	4.2 RK method for solving FDEs	47
	4.3 Problems Tested	48
	4.4 Numerical Results	49
	4.5 Discussion	56
5	PHASE-FITTED AND AMPLIFICATION-FITTED FOURTH ORDER DIAGONALLY IMPLICIT RUNGE-KUTTA METHOD FOR SOLVING OSCILLATORY PROBLEMS	57
	5.1 Introduction	57
	5.2 Derivation of Phase-fitted and Amplification-fitted for Fourth order Diagonally Implicit Runge-Kutta Method (DIRK4)	57
	5.3 Problems Tested	61
	5.4 Numerical Results	62
	5.5 Discussion	72
6	PHASE-FITTED AND AMPLIFICATION-FITTED RUNGE-KUTTA-NYSTRÖM METHODS FOR LINEAR ORDINARY DIFFERENTIAL EQUATIONS FOR SOLVING OSCILLATORY PROBLEMS	73
	6.1 Introduction	73
	6.1.1 Derivation of order conditions for RKN for LODEs	73
	6.1.2 Analysis of Absolute Stability for RKN method	75
	6.1.3 Analysis of Dispersion (Phase-lag) and Dissipation (Amplification error) for RKN method	76
	6.2 Derivation of Fifth order Four-Stage RKN Method for LODEs	77
	6.3 Derivation of Sixth order Five-Stage RKN Method for LODEs	78
	6.4 Derivation of Phase-fitted and Amplification-fitted for Fifth order Four-Stage RKN Method for LODEs	79
	6.4.1 Problems Tested	80
	6.4.2 Numerical Results	81
	6.4.3 Discussion	91
	6.5 Derivation of Phase-fitted and Amplification-fitted for sixth order Five-Stage RKN Method for LODEs	91
	6.5.1 Numerical Results	92
	6.5.2 Discussion	102

6.6	Conclusion	102
7	SOLVING SECOND ORDER WAVE EQUATION USING RUNGE-KUTTA-NYSTRÖM METHODS FOR LINEAR ORDINARY DIFFERENTIAL EQUATIONS	103
7.1	Introduction	103
7.2	Method of Lines	103
7.3	Problems Tested	105
7.4	Numerical Results	105
7.5	Discussion	110
8	CONCLUSION	111
8.1	Summary	111
8.2	Future Work	112
	BIBLIOGRAPHY	113
	BIODATA OF STUDENT	117
	LIST OF PUBLICATIONS	118

LIST OF TABLES

Table		Page
1.1	Butcher Table	7
3.1	4 th order Classical RK method	22
3.2	Comparison of New ERK4 PLDS and Classical RK4 methods when solving Problem 3.1	26
3.3	Comparison of New ERK4 PLDS and Classical RK4 methods when solving Problem 3.2	27
3.4	Comparison of New ERK4 PLDS and Classical RK4 methods when solving Problem 3.3	28
3.5	Comparison of New ERK4 PLDS and Classical RK4 methods when solving Problem 3.4	29
3.6	Comparison of New ERK4 PLDS and Classical RK4 methods when solving Problem 3.5	30
3.7	Comparison of New ERK4 PLDS and Classical RK4 methods when solving Problem 3.6	31
3.8	5 th order RK method for LODEs	35
3.9	Comparison of New ERK5 PLDS and RK5 Zingg methods when solving Problem 3.1	37
3.10	Comparison of New ERK5 PLDS and RK5 Zingg methods when solving Problem 3.2	38
3.11	Comparison of New ERK5 PLDS and RK5 Zingg methods when solving Problem 3.3	39
3.12	Comparison of New ERK5 PLDS and RK5 Zingg methods when solving Problem 3.4	40
3.13	Comparison of New ERK5 PLDS and RK5 Zingg methods when solving Problem 3.5	41
3.14	Comparison of New ERK5 PLDS and RK5 Zingg methods when solving Problem 3.6	42
4.1	Comparison of FRK4 PLDS and FRK4 methods when solving problem 4.1	50

4.2	Comparison of FRK4 PLDS and FRK4 methods when solving problem 4.2	50
4.3	Comparison of FRK4 PLDS and FRK4 methods when solving problem 4.3	51
4.4	Comparison of FRK4 PLDS and FRK4 methods when solving problem 4.4	52
4.5	Comparison of FRK4 PLDS and FRK4 methods when solving problem 4.5	53
5.1	4 th order 5-stage DIRK derived from Hairer	58
5.2	4 th order 4-stage DIRK derived from Che Jawias	60
5.3	Comparison of Hairer DIRK4 PLDS, Hairer DIRK4, Izzati DIRK4 PLDS, and Izaati DIRK4 methods when solving Problem 5.1	63
5.4	Comparison of Hairer DIRK4 PLDS, Hairer DIRK4, Izzati DIRK4 PLDS, and Izaati DIRK4 methods when solving Problem 5.2	64
5.5	Comparison of Hairer DIRK4 PLDS, Hairer DIRK4, Izzati DIRK4 PLDS, and Izaati DIRK4 methods when solving Problem 5.3	65
5.6	Comparison of Hairer DIRK4 PLDS, Hairer DIRK4, Izzati DIRK4 PLDS, and Izaati DIRK4 methods when solving Problem 5.4	66
5.7	Comparison of Hairer DIRK4 PLDS, Hairer DIRK4, Izzati DIRK4 PLDS, and Izaati DIRK4 methods when solving Problem 5.5	67
5.8	Comparison of Hairer DIRK4 PLDS, Hairer DIRK4, Izzati DIRK4 PLDS, and Izaati DIRK4 methods when solving Problem 5.6	68
6.1	The new 4-stage 5 th order RKN for LODEs	77
6.2	The new 5-stage 6 th order RKN for LODEs	78
6.3	Comparison of RKN5 PLDS, New RKN5, VRKN5, and SRKN5 methods when solving Problem 6.1	82
6.4	Comparison of RKN5 PLDS, New RKN5, VRKN5, and SRKN5 methods when solving Problem 6.2	83
6.5	Comparison of RKN5 PLDS, New RKN5, VRKN5, and SRKN5 methods when solving Problem 6.3	84
6.6	Comparison of RKN5 PLDS, New RKN5, VRKN5, and SRKN5 methods when solving Problem 6.4	85

6.7	Comparison of RKN5 PLDS, New RKN5, VRKN5, and SRKN5 methods when solving Problem 6.5	86
6.8	Comparison of RKN5 PLDS, New RKN5, VRKN5, and SRKN5 methods when solving Problem 6.6	87
6.9	Comparison of RKN6 PLDS, New RKN6, and VRKN6 methods when solving Problem 6.1	93
6.10	Comparison of RKN6 PLDS, New RKN6, and VRKN6 methods when solving Problem 6.2	94
6.11	Comparison of RKN6 PLDS, New RKN6, and VRKN6 methods when solving Problem 6.3	95
6.12	Comparison of RKN6 PLDS, New RKN6, and VRKN6 methods when solving Problem 6.4	96
6.13	Comparison of RKN6 PLDS, New RKN6, and VRKN6 methods when solving Problem 6.5	97
6.14	Comparison of RKN6 PLDS, New RKN6, and VRKN6 methods when solving Problem 6.6	98
7.1	Comparison between RKN5 LODEs and RKN6 LODEs for problem 7.1 when $t = 0.5$	106
7.2	Comparison between RKN5 LODEs and RKN6 LODEs for problem 7.2 when $t = 0.5$	107
7.3	Comparison between RKN5 LODEs and RKN6 LODEs for problem 7.3 when $t = 0.4$	108
7.4	Numerical results for New RKN5 LODEs and RKN6 LODEs with FCN and TIME for problems 7.1, 7.2 and 7.3	108

LIST OF FIGURES

Figure		Page
3.1	Accuracy curves for Classical RK4 method with phase-fitted and amplification-fitted for Problem 3.1 with $b=1000$	32
3.2	Accuracy curves for Classical RK4 method with phase-fitted and amplification-fitted for Problem 3.2 with $b=1000$	32
3.3	Accuracy curves for Classical RK4 method with phase-fitted and amplification-fitted for Problem 3.3 with $b=1000$	33
3.4	Accuracy curves for Classical RK4 method with phase-fitted and amplification-fitted for Problem 3.4 with $b=1000$	33
3.5	Accuracy curves for Classical RK4 method with phase-fitted and amplification-fitted for Problem 3.5 with $b=1000$	34
3.6	Accuracy curves for Classical RK4 method with phase-fitted and amplification-fitted for Problem 3.6 with $b=1000$	34
3.7	Accuracy curves for RK5 Zingg method with phase-fitted and amplification-fitted for Problem 3.1 with $b=1000$	43
3.8	Accuracy curves for RK5 Zingg method with phase-fitted and amplification-fitted for Problem 3.2 with $b=1000$	43
3.9	Accuracy curves for RK5 Zingg method with phase-fitted and amplification-fitted for Problem 3.3 with $b=1000$	44
3.10	Accuracy curves for RK5 Zingg method with phase-fitted and amplification-fitted for Problem 3.4 with $b=1000$	44
3.11	Accuracy curves for RK5 Zingg method with phase-fitted and amplification-fitted for Problem 3.5 with $b=1000$	45
3.12	Accuracy curves for RK5 Zingg method with phase-fitted and amplification-fitted for Problem 3.6 with $b=1000$	45
4.1	Results curve for Classical RK4 method with phase-fitted and amplification-fitted for Problem 4.1	54
4.2	Results curve for Classical RK4 method with phase-fitted and amplification-fitted for Problem 4.2	54
4.3	Results curve for Classical RK4 method with phase-fitted and amplification-fitted for Problem 4.3	55

4.4	Results curve for Classical RK4 method with phase-fitted and amplification-fitted for Problem 4.4	55
4.5	Results curve for Classical RK4 method with phase-fitted and amplification-fitted for Problem 4.5	56
5.1	Accuracy curves for Hairer DIRK4 and Izzati DIRK4 methods with phase-fitted and amplification-fitted for Problem 5.1	69
5.2	Accuracy curves for Hairer DIRK4 and Izzati DIRK4 methods with phase-fitted and amplification-fitted for Problem 5.2	69
5.3	Accuracy curves for Hairer DIRK4 and Izzati DIRK4 methods with phase-fitted and amplification-fitted for Problem 5.3	70
5.4	Accuracy curves for Hairer DIRK4 and Izzati DIRK4 methods with phase-fitted and amplification-fitted for Problem 5.4	70
5.5	Accuracy curves for Hairer DIRK4 and Izzati DIRK4 methods with phase-fitted and amplification-fitted for Problem 5.5	71
5.6	Accuracy curves for Hairer DIRK4 and Izzati DIRK4 methods with phase-fitted and amplification-fitted for Problem 5.6	71
6.1	Accuracy curves for New RKN5 with phase-fitted and amplification-fitted for Problem 6.1 with $b=1000$	88
6.2	Accuracy curves for New RKN5 with phase-fitted and amplification-fitted for Problem 6.2 with $b=1000$	88
6.3	Accuracy curves for New RKN5 with phase-fitted and amplification-fitted for Problem 6.3 with $b=1000$	89
6.4	Accuracy curves for New RKN5 with phase-fitted and amplification-fitted for Problem 6.4 with $b=1000$	89
6.5	Accuracy curves for New RKN5 with phase-fitted and amplification-fitted for Problem 6.5 with $b=1000$	90
6.6	Accuracy curves for New RKN5 with phase-fitted and amplification-fitted for Problem 6.6 with $b=1000$	90
6.7	Accuracy curves for New RKN6 with phase-fitted and amplification-fitted for Problem 6.1 with $b=1000$	99
6.8	Accuracy curves for New RKN6 with phase-fitted and amplification-fitted for Problem 6.2 with $b=1000$	99
6.9	Accuracy curves for New RKN6 with phase-fitted and amplification-fitted for Problem 6.3 with $b=1000$	100

6.10	Accuracy curves for New RKN6 with phase-fitted and amplification-fitted for Problem 6.4 with $b=1000$	100
6.11	Accuracy curves for New RKN6 with phase-fitted and amplification-fitted for Problem 6.5 with $b=1000$	101
6.12	Accuracy curves for New RKN6 with phase-fitted and amplification-fitted for Problem 6.6 with $b=1000$	101
7.1	Curves for New RKN5 LODEs, RKN6 LODEs for solving Problem 7.1	109
7.2	Curves for New RKN5 LODEs, RKN6 LODEs for solving Problem 7.2	109
7.3	Curves for New RKN5 LODEs, RKN6 LODEs for solving Problem 7.3	110

LIST OF ABBREVIATIONS

DEs	Differential Equations
ODEs	Ordinary Differential Equations
IVPs	Initial Value Problems
RK	Runge-Kutta method
RKN	Runge-Kutta-Nyström method
FDEs	Fuzzy Differential Equations
FIVP	Fuzzy Initial Value Problem
PDEs	Partial Differential Equations
HPDEs	Hyperbolic Partial Differential Equations
LTE	Local Truncation Error
FSAL	The First Same as Last
LODEs	Linear Ordinary Differential Equations
DIRK	Diagonally Implicit Runge-Kutta method
PL	Phase-lag or dispersion
DS	Dissipation
E	Absolute error
MAXE	Maximum of absolute error
FCN	Number of function evaluations
TIME	Execution time
New ERK4 PLDS	New Classical fourth order Runge-Kutta method with phase-fitted and amplification-fitted
Classical RK4	Classical Runge-Kutta method of order four
New ERK5 PLDS	New Runge-Kutta method of order five for Linear Ordinary Differential Equations proposed from Zingg (1999) with phase-fitted and amplification-fitted

RK5 Zingg	Fifth order Runge-Kutta method for Linear Ordinary Differential Equations proposed from Zingg (1999)
FRK4 PLDS	Classical Runge-Kutta method of order four with phase-fitted and amplification-fitted for solving Fuzzy Differential Equations
FRK4	Classical Runge-Kutta method of order four for solving Fuzzy Differential Equations
DIRK4	Fourth order Diagonally Implicit Runge-Kutta method
Hairer DIRK4 PLDS	Fourth order five-stage diagonally implicit Runge-Kutta method proposed from Hairer (2010) with phase-fitted and amplification-fitted
Hairer DIRK4	Fourth order five-stage diagonally implicit Runge-Kutta method proposed from Hairer (2010)
Izzati DIRK4 PLDS	Fourth order four-stage diagonally implicit Runge-Kutta method proposed from Che Jawias (2010) with phase-fitted and amplification-fitted
Izzati DIRK4	Fourth order four-stage diagonally implicit Runge-Kutta method proposed from Che Jawias (2010)
New RKN5	New four-stage fifth order Runge-Kutta-Nyström of Linear Ordinary Differential Equations
RKN5 PLDS	New four-stage fifth order Runge-Kutta-Nyström of Linear Ordinary Differential Equations with phase-fitted and amplification-fitted
VRKN5	Fifth order four-stage Runge-Kutta-Nyström proposed from Van der Houwen et.al (1987) with dispersion of order two and dissipation of order eight
SRKN5	Fifth order four-stage Runge-Kutta-Nyström proposed from Simos et.al (1994) with dispersion of order four and dissipation of order eight
New RKN6	New five-stage sixth order Runge-KuttaNyström method for Linear Ordinary Differential Equations
RKN6 PLDS	New five-stage sixth order Runge-Kutta-Nyström method for Linear Ordinary Differential Equations with phase-fitted and amplification-fitted

VRKN6	Sixth order five-stage Runge-Kutta-Nyström proposed from Van der Houwen et.al (1987) with dispersion of order two and dissipation of order eight
RKN5 LODEs	New four-stage fifth order Runge-Kutta-Nyström of Linear Ordinary Differential Equations
RKN6 LODEs	New five-stage sixth order Runge-KuttaNyström method for Linear Ordinary Differential Equations



CHAPTER 1

INTRODUCTION

Ordinary differential equations (ODEs) are equations that involve an unknown function with independent variable and one or more of its derivatives. ODEs arise in many contexts of engineering and science such as fluid dynamics, radioactive decay and population growth.

Many theoretical and numerical studies for such equations have appeared in literature. The analytical way to solve ODEs is via application of integration technique. However, the anti-derivatives for most realistic systems of ODEs are difficult or impossible to find. Thus, numerical methods for ODEs have attracted considerable attention.

The initial value problems (IVPs) of special second order ODEs are defined as follows:

$$y'' = f(x, y), \quad y(x_0) = y_0, \quad y'(x_0) = y'_0 \quad (1.1)$$

where $f: R \times R \rightarrow R^n$, and $y_0, y'_0 \in R^n$. The solution of (1.1) exhibits a pronounced oscillatory character. One way to solve oscillatory problems is by Runge-Kutta methods. In this study, we are focusing on solving problem (1.1) by using RK methods and RKN methods for oscillating problems.

1.1 Ordinary Differential Equations

Any research on ordinary differential equations gives a number of methods for explicitly finding solutions to first-order initial-value problems (IVPs). In practice, however, few of the problems originating from the study of physical phenomena can be solved exactly.

The n -th order ODEs can be written as:

$$y^{(n)} = f(x, y, \dots, y^{(n-1)}), \text{ where } n = 2, 3, 4$$

with initial conditions:

$$y(a) = y_0 \text{ and } y^{(i)}(a) = \eta_i, 0 < i < n - 1, x \in [a, b]$$

while the first order ODEs can be written as:

$$\frac{dy}{dx} = f(x, y(x)), y(a) = y_0 \text{ for } x \in [a, b] \quad (1.2)$$

In (1.2), the quantity being differentiated, y is named as the dependent variable, while the quantity with respect to which y is differentiated, x is named as independent variable.

Below, we state the existence and uniqueness theorem for finding the solution to the first order IVPs.

Theorem 1.1: (Existence and Uniqueness)

Let $f(x, y)$ be defined and continuous for all points (x, y) in the region D defined by $a \leq x \leq b, -\infty < y < \infty$, where a and b are finite, and let there exists a constant L such that for any $x \in [a, b]$ and any two numbers y_1 and y_2 ,

$$|f(x, y_1) - f(x, y_2)| \leq L|y_1 - y_2|$$

This condition is known as *Lipschitz condition*. Then there exists exactly one function $y(x)$ with the following three properties:

- i. $y(x)$ is continuous and differentiable for $x \in [a, b]$,
- ii. $y' = f(x, y(x)), x \in [a, b]$,
- iii. $y(a) = \eta$.

The proof is given by Henrici (1962).

Basically, the numerical methods for ODEs are classified as one-step method and multistep method. One-step method requires the information from only one previous point x_n to find the approximation at the mesh point x_{n+1} . On the other hand, multistep method requires the usage of information from more than one previous points to find the next approximation.

Many differential equations which appear in practice are systems of second order IVPs (1.1) in which the derivative does not appear explicitly. Such a system can be transformed into first order differential equations of doubled dimension by considering the vector (y, y') as the new variable. In this study, we are focusing on solving special second order equation (1.1) for which it is known in advance that their solution is oscillating. Consider the second order linear differential equation:

$$y'' = Ay \quad (1.3)$$

where A is a continuous real-valued function. A solution of (1.3) is said to be *oscillatory* if it has arbitrarily large zeros, and otherwise it is said to be *nonoscillatory*. Equation (1.3) is called oscillatory if all its solutions are oscillatory.

When dealing with the oscillatory problems of (1.1), we need to consider the dispersion (phase-lag) and dissipation (amplification error) properties of the methods developed.

1.2 Fuzzy Differential Equations

In many cases of the modeling of real world problem, knowledge or information about the behavior of the physical system is often incomplete, uncertain or vague. For example, values of parameter, functional relationship or initial conditions, may not be accurate. These uncertainties have to be considered for obtaining a more realistic model. Fuzzy Differential Equations (FDEs) are utilized for the purpose of the modeling of uncertainty and processing vague or incomplete information in mathematical models.

Most of the problems in science and engineering require the solution of a Fuzzy Differential Equation (FDE) which are satisfied in Fuzzy initial conditions. Due to the large potential of FDEs involved in various fields, a lot of research has been focused in this area. Sometimes it is too complicated to obtain the exact solution of (FDE) which models the science and engineering problems, hence numerical methods are used.

Below, we state some definitions and theorems for FDEs.

Definition 1.2.1:

An arbitrary fuzzy number is represented by ordered pair of functions $(\underline{u}(\alpha), \bar{u}(\alpha))$, $0 \leq \alpha \leq 1$, satisfying the following requirements:

- $\underline{u}(\alpha)$ is a bounded left continuous nondecreasing function over $[0,1]$.
- $\bar{u}(\alpha)$ is a bounded left continuous nonincreasing function over $[0,1]$.
- $\underline{u}(\alpha) \leq \bar{u}(\alpha)$, $0 \leq \alpha \leq 1$.

Definition 1.2.2:

Let I be a real interval. A mapping $y: I \rightarrow E$ is called a fuzzy process and its α -level set is denoted by:

$$[y(t)]^\alpha = [\underline{y}^\alpha(t), \bar{y}^\alpha(t)], t \in I, \alpha \in (0,1] \quad (1.4)$$

Definition 1.2.3:

Let R_F denote the class of fuzzy numbers and let $x, y \in R_F$. If there exists $z \in R_F$ such that $x = y + z$, then z is called the Hukuhara difference of x and y and it is denoted by $x \ominus y$.

Note: the " \ominus " sign stands always for Hukuhara difference and $x \ominus y \neq x + (-1)y$.

Definition 1.2.4:

Let $f: (a, b) \rightarrow R_F$ and $x_0 \in (a, b)$. We say that f is Hukuhara differentiable at x_0 , if there exists an element $f'(x_0) \in R_F$, such that for all $h > 0$ sufficiently small:

$\exists f(x_0 + h) \ominus f(x_0), f(x_0) \ominus f(x_0 - h)$ and the limits:

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) \ominus f(x_0)}{h} = \lim_{h \rightarrow 0} \frac{f(x_0) \ominus f(x_0 - h)}{h} = f'(x_0) \quad (1.5)$$

Theorem 1.2.1:

Let $F: (a, b) \rightarrow R_F$ be Hukuhara differentiable and denote $[F(t)]^\alpha = [\underline{F}^\alpha(t), \overline{F}^\alpha(t)]$.

Then the boundary functions $\underline{F}^\alpha(t)$ and $\overline{F}^\alpha(t)$ are differentiable and:

$$[F'(t)]^\alpha = [(\underline{F}^\alpha)'(t), (\overline{F}^\alpha)'(t)], \alpha \in [0, 1] \quad (1.6)$$

The proof is given by Kaleva (2006).

Definition 1.2.5:

Let us consider the fuzzy initial value problem (FIVP):

$$y' = f(t, y), y(t_0) = y_0 \quad (1.7)$$

where $f: [t_0, t_0 + \alpha] \times R_F \rightarrow R_F$ and $y_0 \in R_F$. Then the theorem (1.2.1) shows us a way how to translate FIVP (1.7) into a system of ODEs.

Definition 1.2.6:

Let $[y(t)]^\alpha = [\underline{y}^\alpha(t), \overline{y}^\alpha(t)]$. If $y(t)$ is Hukuhara differentiable then $[y'(t)]^\alpha = [(\underline{y}^\alpha)'(t), (\overline{y}^\alpha)'(t)]$. So equation (4.4) translates into the following systems of ODEs:

$$\begin{cases} ((\underline{y}^\alpha)'(t)) = \underline{f}^\alpha(t, \underline{y}^\alpha(t), \bar{y}^\alpha(t)) \\ ((\bar{y}^\alpha)'(t)) = \bar{f}^\alpha(t, \underline{y}^\alpha(t), \bar{y}^\alpha(t)) \\ \underline{y}^\alpha(t_0) = \underline{y}_0^\alpha \\ \bar{y}^\alpha(t_0) = \bar{y}_0^\alpha \end{cases} \quad (1.8)$$

where:

$$[f(t, y)]^\alpha = [\underline{f}^\alpha(t, \underline{y}^\alpha, \bar{y}^\alpha), \bar{f}^\alpha(t, \underline{y}^\alpha, \bar{y}^\alpha)] \quad (1.9)$$

Definition 1.2.7:

The Seikkala derivative $y'(t)$ of a fuzzy process y (defined by equation (1.9)) is defined by:

$$[y'(t)]^\alpha = [(\underline{y}^\alpha)'(t), (\bar{y}^\alpha)'(t)], 0 < \alpha \leq 1 \quad (1.10)$$

Remark 1.2.1:

If $y: I \rightarrow E$ is Seikkala differentiable and its Seikkala derivative y' is integrable over $[0,1]$, then:

$$y(t) = y(t_0) + \int_{t_0}^t y'(s) ds \quad (1.11)$$

for all values of t_0, t where $t_0, t \in I$.

1.3 Partial Differential Equations

Many problems in applied science, physics, and engineering can be modeled as partial differential equations (PDEs), that is differential equation which involves more than one independent variable.

A PDE for the function $u(x_1, \dots, x_n)$ is an equation of the form:

$$f\left(x_1, \dots, x_n, u, \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n}, \frac{\partial^2 u}{\partial x_1 \partial x_1}, \dots, \frac{\partial^2 u}{\partial x_n \partial x_n}, \dots\right) = 0 \quad (1.12)$$

Below we listed some of the common partial differential equations:

Elliptic equations:

The PDE involves $u_{xx}(x, y) + u_{yy}(x, y)$ is an *elliptic equation*. The particular elliptic equation we will consider in this thesis is known as the *Poisson equation*:

$$\frac{\partial^2 u}{\partial x^2}(x, y) + \frac{\partial^2 u}{\partial y^2}(x, y) = f(x, y) \quad (1.13)$$

This equation arise in the study various time-independent physical problems such as the steady-state distribution of heat in a plane region, the potential energy of a point in a plane acted on by gravitational forces in the plane and two dimensional steady-state problems involving incompressible fluids.

Parabolic equations:

Another type of PDE is the *parabolic* partial differential equation of the form:

$$\frac{\partial u}{\partial t}(x, t) - \alpha^2 \frac{\partial^2 u}{\partial x^2}(x, t) = 0 \quad (1.14)$$

The PDE concerns the flow of heat along a rod of length l , which has a uniform temperature within each cross-sectional element. The parameter α is determined by the heat conductivity of the rod.

Hyperbolic equations:

The hyperbolic equation is the one-dimensional *wave equation*. Suppose an elastic string of length l is stretched between two supports at the same horizontal level.

If the string is set to vibrate in a vertical plane, the vertical displacement $u(x, t)$ of a point x at time t satisfies the PDE:

$$\alpha^2 \frac{\partial^2 u}{\partial x^2}(x, t) - \frac{\partial^2 u}{\partial t^2}(x, t) = 0 \quad \text{for } 0 < x < 1 \text{ and } 0 < t \quad (1.15)$$

1.4 Runge-Kutta (RK) methods

An s-stage explicit Runge-Kutta method can be expressed by the following relations:

$$y_{n+1} = y_n + h \sum_{i=1}^s b_i k_i \quad (1.16)$$

where

$$k_i = f \left(x_n + c_i, y_n + h \sum_{j=1}^i a_{ij} k_j \right) \quad (1.17)$$

The method is used for the computation of the approximation of y_{n+1} , when y_n is known. The method shown above can also be presented using the Butcher table below:

Table 1.1 : Butcher Table

c_2	a_{21}			
c_3	a_{31}	a_{32}		
\vdots	\vdots	\vdots	\vdots	
c_s	a_{s1}	a_{s2}	\vdots	$a_{s,s-1}$
	b_1	b_2	\vdots	b_s

where the coefficients c_2, \dots, c_s must satisfy the equations:

$$c_i = \sum_{j=1}^{i-1} a_{ij}, \quad i = 2, \dots, s \quad (1.18)$$

1.4.1 Algebraic Order Condition for RK method

General order conditions for RK method can be attained from direct expansion of Taylor series by using Local Truncation Error (LTE). The s-stage up to order six RK methods are given as follows:

$$\text{order 1: } \sum b_i = 1 \quad (1.19)$$

$$\text{order 2: } \sum b_i c_i = \frac{1}{2} \quad (1.20)$$

$$\text{order 3: } \frac{1}{2} \sum b_i c_i^2 = \frac{1}{6} \quad (1.21)$$

$$\text{order 4: } \frac{1}{6} \sum b_i c_i^3 = \frac{1}{24} \quad (1.22)$$

$$\sum b_i a_{ij} c_j = \frac{1}{24} \quad (1.23)$$

$$\text{order 5: } \frac{1}{24} \sum b_i c_i^4 = \frac{1}{120} \quad (1.24)$$

$$\frac{1}{4} \sum b_i c_i a_{ij} c_j = \frac{1}{120} \quad (1.25)$$

$$\frac{1}{2} \sum b_i a_{ij} c_j^2 = \frac{1}{120} \quad (1.26)$$

$$\text{order 6: } \frac{1}{120} \sum b_i c_i^5 = \frac{1}{720} \quad (1.27)$$

$$\frac{1}{20} \sum b_i c_i^2 a_{ij} c_j = \frac{1}{720} \quad (1.28)$$

$$\frac{1}{10} \sum b_i c_i a_{ij} c_j^2 = \frac{1}{720} \quad (1.29)$$

$$\frac{1}{6} \sum b_i a_{ij} c_j^3 = \frac{1}{720} \quad (1.30)$$

$$\sum b_i a_{ij} a_{jk} c_k = \frac{1}{720} \quad (1.31)$$

where the coefficients may be made dependent by imposing the Butcher conditions (1963):

$$\sum_{i=1}^s b_i a_{ij} = b_j (1 - c_j), \quad j = 2, 3, \dots, s \quad (1.32)$$

Additionally, if $b_2 = 0$, we have:

$$\sum_{j=1}^{i-1} a_{ij} c_j = \frac{c_i^2}{2}, \quad i = 3, 4, \dots, s \quad (1.33)$$

These valuable relations actually reduce the total number of conditions arising in higher orders. So, the conditions (1.34) and (1.35) are supplemented by extra simplifying relations.

1.4.2 Local Truncation Error for RK method

Dormand (1996) proposed that having achieved a particular order of accuracy. The best strategy for practical purposes would be choosing the free RK parameters is to minimize the error norm:

$$A^{(p+1)} = \|\tau^{(p+1)}\|_2 = \sqrt{\left(\sum_{j=1}^{n_{p+1}} \tau_j^{(p+1)}\right)^2} \quad (1.34)$$

The error coefficients up to order six for RK methods are as follows:

$$\text{order 1: } \tau_1^{(1)} = \sum b_i - 1 \quad (1.35)$$

$$\text{order 2: } \tau_1^{(2)} = \sum b_i c_i - \frac{1}{2} \quad (1.36)$$

$$\text{order 3: } \tau_1^{(3)} = \frac{1}{2} \sum b_i c_i^2 - \frac{1}{6} \quad (1.37)$$

$$\text{order 4: } \tau_1^{(4)} = \frac{1}{6} \sum b_i c_i^3 - \frac{1}{24} \quad (1.38)$$

$$\tau_2^{(4)} = \sum b_i a_{ij} c_j - \frac{1}{24} \quad (1.39)$$

$$\text{order 5: } \tau_1^{(5)} = \frac{1}{24} \sum b_i c_i^4 - \frac{1}{120} \quad (1.40)$$

$$\tau_2^{(5)} = \frac{1}{4} \sum b_i c_i a_{ij} c_j - \frac{1}{120} \quad (1.41)$$

$$\tau_3^{(5)} = \frac{1}{2} \sum b_i a_{ij} c_j^2 - \frac{1}{120} \quad (1.42)$$

$$\text{order 6: } \tau_1^{(6)} = \frac{1}{120} \sum b_i c_i^5 - \frac{1}{720} \quad (1.43)$$

$$\tau_2^{(6)} = \frac{1}{20} \sum b_i c_i^2 a_{ij} c_j - \frac{1}{720} \quad (1.44)$$

$$\tau_3^{(6)} = \frac{1}{10} \sum b_i c_i a_{ij} c_j^2 - \frac{1}{720} \quad (1.45)$$

$$\tau_4^{(6)} = \frac{1}{6} \sum b_i a_{ij} c_j^3 - \frac{1}{720} \quad (1.46)$$

$$\tau_5^{(6)} = \sum b_i a_{ij} a_{jk} c_k - \frac{1}{720} \quad (1.47)$$

1.4.3 Analysis of Absolute Stability for RK method

Consider the standard test problem of differential equation:

$$y' = f(t, y) = \lambda y \text{ and } y(x_n) = y_n \quad (1.48)$$

which has the true solution:

$$y(x) = y_n e^{\lambda(x-x_n)} \quad (1.49)$$

Applying the test equation (1.48) to the RK formula (1.16) and by setting $v = \lambda h$, we obtain:

$$y_{n+1} = R(v)y_n, \text{ where } |R(v)| < 1 \quad (1.50)$$

$$y_{n+1} = [1 + vb^T(I - vA)^{-1}]y_n \quad (1.51)$$

where $b = b_i$ and $A = a_{ij}$, $i, j = 1, \dots, s$ are parameters of RK method and $R(v)$ is said to be the stability polynomial. The stability polynomial for RK method of order p is:

$$R(v) = 1 + v + \frac{v^2}{2!} + \frac{v^3}{3!} + \frac{v^4}{4!} + \dots + \frac{v^p}{p!} \quad (1.52)$$

1.5 Runge-Kutta-Nyström (RKN) methods

Nyström (1925) introduced RK methods for second order ODEs which has been called RKN methods. The form of s -stage RKN method of order p is:

$$y_{n+1} = y_n + hy'_n + h^2 \sum_{i=1}^s b_i k_i$$

$$y'_{n+1} = y_n + h \sum_{i=1}^s \bar{b}_i k_i \quad (1.53)$$

where

$$k_i = f\left(x_n + c_i h, y_n + c_i h y'_n + h^2 \sum_{j=1}^{s-1} a_{ij} k_j\right), i = 1, 2, \dots, s \quad (1.54)$$

1.5.1 Algebraic Condition for RKN method

The order condition for RKN methods can be attained from direct expansion of the Taylor series by using the local Truncation Error (LTE). The order conditions for RKN methods up to order six obtained from Dormand (1996) are given as follows:

for y:

$$\text{order 2: } \sum b_i = \frac{1}{2} \quad (1.55)$$

$$\text{order 3: } \sum b_i c_i = \frac{1}{6} \quad (1.56)$$

$$\text{order 4: } \frac{1}{2} \sum b_i c_i^2 = \frac{1}{24} \quad (1.57)$$

$$\text{order 5: } \frac{1}{6} \sum b_i c_i^3 = \frac{1}{120} \quad (1.58)$$

$$\sum b_i a_{ij} c_j = \frac{1}{120} \quad (1.59)$$

$$\text{order 6: } \frac{1}{24} \sum b_i c_i^4 = \frac{1}{720} \quad (1.60)$$

$$\frac{1}{2} \sum b_i c_i a_{ij} c_j = \frac{1}{720} \quad (1.61)$$

$$\frac{1}{2} \sum b_i a_{ij} c_j^2 = \frac{1}{720} \quad (1.62)$$

for y':

$$\text{order 1: } \sum \bar{b}_i = 1 \quad (1.63)$$

$$\text{order 2: } \sum \bar{b}_i c_i = \frac{1}{2} \quad (1.64)$$

$$\text{order 3: } \frac{1}{2} \sum \bar{b}_i c_i^2 = \frac{1}{6} \quad (1.65)$$

$$\text{order 4: } \frac{1}{6} \sum \bar{b}_i c_i^3 = \frac{1}{24} \quad (1.66)$$

$$\sum \bar{b}_i a_{ij} c_j = \frac{1}{24} \quad (1.67)$$

$$\text{order 5: } \frac{1}{24} \sum \bar{b}_i c_i^4 = \frac{1}{120} \quad (1.68)$$

$$\frac{1}{4} \sum \bar{b}_i c_i a_{ij} c_j = \frac{1}{120} \quad (1.69)$$

$$\frac{1}{2} \sum \bar{b}_i a_{ij} c_j^2 = \frac{1}{120} \quad (1.70)$$

$$\text{order 6: } \frac{1}{120} \sum \bar{b}_i c_i^5 = \frac{1}{720} \quad (1.71)$$

$$\frac{1}{20} \sum \bar{b}_i c_i^2 a_{ij} c_j = \frac{1}{720} \quad (1.72)$$

$$\frac{1}{10} \sum \bar{b}_i c_i a_{ij} c_j^2 = \frac{1}{720} \quad (1.73)$$

$$\frac{1}{6} \sum \bar{b}_i a_{ij} c_j^3 = \frac{1}{720} \quad (1.74)$$

$$\sum \bar{b}_i a_{ij} a_{jk} c_k = \frac{1}{720} \quad (1.75)$$

The Nyström row sum conditions that need to be satisfied are:

$$\sum_{i=1}^s a_{ij} = \frac{c_i^2}{2}, \quad i = 1, \dots, s \quad (1.76)$$

The simplifying assumption given by Butcher (2003) which is used in order to reduce the number of equations:

$$b_i = \bar{b}_i (1 - c_i) \quad (1.77)$$

The First Same as Last (FSAL) property where the last stage is evaluated at the same point as the first stage of the next step is used to reduce function evaluation. For RKN method to be FSAL, it has to satisfy:

$$c_1 = 0, c_s = 1 \text{ and } a_{sj} = b_j, j = 1, \dots, s - 1 \quad (1.78)$$

1.5.2 Local Truncation Error for RKN method

We need to use the quantity (1.36) for y and the quantity (1.81) for y' in order to find the norms of (LTE) coefficients for RKN method. The quantity for y' is known as:

$$\|\tau^{(p+1)}\|_2 = \sqrt{\left(\sum_{j=1}^{n_{p+1}} \tau_j^{(p+1)} \right)^2} \text{ for } y'_n \quad (1.79)$$

The error coefficients for RKN methods up to order six are:
for y:

$$\text{order 2: } \tau_1^{(2)} = \sum b_i - \frac{1}{2} \quad (1.80)$$

$$\text{order 3: } \tau_1^{(3)} = \sum b_i c_i - \frac{1}{6} \quad (1.81)$$

$$\text{order 4: } \tau_1^{(4)} = \frac{1}{2} \sum b_i c_i^2 - \frac{1}{24} \quad (1.82)$$

$$\text{order 5: } \tau_1^{(5)} = \frac{1}{6} \sum b_i c_i^3 - \frac{1}{120} \quad (1.83)$$

$$\tau_2^{(5)} = \sum b_i a_{ij} c_j - \frac{1}{120} \quad (1.84)$$

$$\text{order 6: } \tau_1^{(6)} = \frac{1}{24} \sum b_i c_i^4 - \frac{1}{720} \quad (1.85)$$

$$\tau_2^{(6)} = \frac{1}{2} \sum b_i c_i a_{ij} c_j - \frac{1}{720} \quad (1.86)$$

$$\tau_3^{(6)} = \frac{1}{2} \sum b_i a_{ij} c_j^2 - \frac{1}{720} \quad (1.87)$$

for y':

$$\text{order 1: } \tau_1^{(1)} = \sum \bar{b}_i - 1 \quad (1.88)$$

$$\text{order 2: } \tau_1^{(2)} = \sum \bar{b}_i c_i - \frac{1}{2} \quad (1.89)$$

$$\text{order 3: } \tau_1^{(3)} = \frac{1}{2} \sum \bar{b}_i c_i^2 - \frac{1}{6} \quad (1.90)$$

$$\text{order 4: } \tau_1^{(4)} = \frac{1}{6} \sum \bar{b}_i c_i^3 - \frac{1}{24} \quad (1.91)$$

$$\tau_2^{(4)} = \sum \bar{b}_i a_{ij} c_j - \frac{1}{24} \quad (1.92)$$

$$\text{order 5: } \tau_1^{(5)} = \frac{1}{24} \sum \bar{b}_i c_i^4 - \frac{1}{120} \quad (1.93)$$

$$\tau_2^{(5)} = \frac{1}{4} \sum \bar{b}_i c_i a_{ij} c_j - \frac{1}{120} \quad (1.94)$$

$$\tau_3^{(5)} = \frac{1}{2} \sum \bar{b}_i a_{ij} c_j^2 - \frac{1}{120} \quad (1.95)$$

$$\text{order 6: } \tau_1^{(6)} = \frac{1}{120} \sum \bar{b}_i c_i^5 - \frac{1}{720} \quad (1.96)$$

$$\tau_2^{(6)} = \frac{1}{20} \sum \bar{b}_i c_i^2 a_{ij} c_j - \frac{1}{720} \quad (1.97)$$

$$\tau_3^{(6)} = \frac{1}{10} \sum \bar{b}_i c_i a_{ij} c_j^2 - \frac{1}{720} \quad (1.98)$$

$$\tau_4^{(6)} = \frac{1}{6} \sum \bar{b}_i a_{ij} c_j^3 - \frac{1}{720} \quad (1.99)$$

$$\tau_5^{(6)} = \sum \bar{b}_i a_{ij} a_{jk} c_k - \frac{1}{720} \quad (1.100)$$

1.6 Objectives of the Thesis

In this study, we developed new and more efficient methods based on RK methods and RKN methods for solving oscillatory problems, first order fuzzy differential equations and second order PDE particularly the wave equation. The main objectives of the thesis are:

- i. To derive phase-fitted and amplification-fitted explicit RK methods for the fourth order (classical) method and fifth order method of Zingg (1999) for linear ordinary differential equations (LODEs) for solving ordinary differential equations with oscillatory solutions.
- ii. To discover the effect of the phase-fitted and amplification-fitted of explicit RK methods on the first order fuzzy differential equations (FDEs).
- iii. To derive phase-fitted and amplification-fitted fourth order five-stage diagonally implicit RK method (DIRK4) proposed from Hairer et.al (2010) and phase-fitted and amplification-fitted fourth order four-stage diagonally implicit RK method proposed from Che Jawias et.al (2010) for ordinary differential equations with oscillatory solutions.
- iv. To derive fifth order four-stage and sixth order five-stage RKN methods for LODEs for solving ordinary differential equations with oscillatory solutions.
- v. To derive the phase-fitted and amplification-fitted RKN methods for LODEs for solving ordinary differential equations with oscillatory solutions.
- vi. To discover the effect of RKN methods for LODEs on hyperbolic partial differential equations (PDEs).

1.7 Outline of Thesis

In Chapter 1, a brief introduction on ordinary differential equations, fuzzy differential equations and partial differential equations are given. The development of numerical

methods, basic theory on algebraic order of RK method and RKN method and LTE for RK and RKN methods are also discussed in this chapter.

Chapter 2 focused on the literature reviews on explicit RK methods, DIRK methods, RKN methods, (FDEs) and hyperbolic partial differential equations.

In Chapter 3, we discuss the phase-fitted and amplification-fitted for explicit RK methods. We derive phase-fitted and amplification-fitted for fourth order classical RK method and phase-fitted and amplification-fitted for fifth order RK method for LODEs proposed from Zingg (1999). The stability region of the methods will be determined. The results of the new methods have been compared with the existing methods for solving LODEs.

In Chapter 4, we used the methods derived in Chapter 3 for solving first order fuzzy differential equations (FDEs). The numerical results for method with phase-fitted and amplification-fitted have been compared with the method without phase-fitted and amplification-fitted for solving FDEs, which clearly shown the advantage of the method with phase-fitted and amplification-fitted.

In Chapter 5, we derived the phase-fitted and amplification-fitted fourth order five-stage diagonally implicit RK method proposed by Hairer et.al (2010) and fourth order four-stage diagonally implicit RK method for LODEs proposed by Che Jawias et.al (2010). The results of the new methods have been compared with the original methods without phase-fitted and amplification-fitted.

In Chapter 6, we derived the order conditions of Runge-Kutta-Nyström method purposely for solving linear second order ordinary differential equations. Based on the order conditions we derived the new fifth order four-stage and sixth order five-stage explicit Runge-Kutta-Nyström methods for linear ordinary differential equations (LODEs). Then the methods are phase-fitted and amplification-fitted so that they will have zero-dispersion and zero-dissipation. The fifth order four-stage RKN method for LODEs has the property of First Same As Last (FSAL). Numerical results proved that the methods with phase-fitted and amplification-fitted are much more efficient than the existing methods with the same algebraic order.

Chapter 7, focused on the application of RKN method for solving hyperbolic partial differential equations (PDEs) that is the second order wave equations. The PDE is first converted to a system of second order linear ODEs using the methods of lines. The resulting second order linear ODEs are then solved using the RKN method derived in Chapter 6. Numerical results show that the RKN methods for LODEs are accurate and reliable for solving the second order wave equation.

Finally, we summarized and conclude our studies in the last chapter that is Chapter 8. Future studies are also suggested in this chapter.

1.8 Problem Statement

Initial value problem (IVPs) for second order ODEs where the first derivative does not appear explicitly often arise in many scientific areas of engineering and applied science such as celestial mechanics, molecular dynamics and quantum mechanics. The solution of these IVPs often exhibits a pronounced oscillatory character and it is well known that it is always difficult to get the accurate numerical results if the initial value problems are oscillatory in nature. To address the problem we will focus on developing RK type methods with reduced phase-lag and dissipation errors. Phase-lag or dispersion error is the angle between the true and the approximated solution and dissipation is the distance of the approximate solution from the standard cyclic solution. The aim of this research is to derive methods with phase-fitted and amplification-fitted so that the methods have zero phase-lag and zero dissipation; hence they are suitable for solving oscillatory problems.

1.9 Scope of Study

The main purpose of this research is to solve the second order ordinary differential equations (1.1) in which the solutions exhibit a pronounced oscillatory character. In this study we are focusing on solving problem (1.1) by using Runge-Kutta methods and Runge-Kutta-Nyström methods for oscillatory problems with phase-fitted and amplification-fitted techniques. The research also includes numerical solutions of first order fuzzy differential equations by using the phase-fitted and amplification fitted explicit RK methods and numerical solutions of hyperbolic partial differential equations by using RKN methods for LODEs.

BIBLIOGRAPHY

- Abbasbandy S, Allahviranloo T, (2002), Numerical solution of fuzzy differential equations by Taylor method. *Computational Methods In Applied Mathematics*, Vol. 2, No. 2, 113–124.
- Abbasbandy S, Allahviranloo T, Lopez-Pouso O, Nieto JJ, (2004), Numerical methods for fuzzy differential inclusions. *Comput Math Appl* 48(10–11), 1633–1641.
- Ahmad S. Z, Ismail F., Senu N. and Suleiman M., (2013), Semi Implicit Hybrid Methods with Higher Order Dispersion for Solving Oscillatory Problems, *Abstract and Applied Analysis*, doi.org/10. 1155/2013/36961.
- Allahviranloo T, Ahmady N. E., (2007), Numerical solution of fuzzy differential equations by predictor-corrector method. *Inform Sci* 177/7, 1633–1647.
- Anastassi Z. A, Vlachos D. S, and Simos T. E., (2009), A family of Runge-Kutta methods with zero phase-lag and derivatives for the numerical solution of the Schrödinger equation and related problems, *J. Math Chem* 46, 1158-1171.
- Bede B, Gal SG, (2005), Generalizations of the differentiability of fuzzy-number-valued functions with applications to fuzzy differential equations. *Fuzzy Set Sys.* 151, 581–599.
- Bede B, Gnana Bhaskar T, Lakshmikantham V, (2007), Perspectives of fuzzy initial value problems, *Communications in Applied Analysis* 11, 339-358.
- Bettis D. G., (1973), A Runge-Kutta-Nyström Algorithm, *Celestial Mechanics*, 8, 229-233.
- Bouziani A., (1994), Mixed problem for certain nonclassical equations containing a small parameter, *Acad. Roy. Belg. Bull. CI, Sci.* 6 (5), 389–400.
- Bouziani A., (1997), Strong solution for a mixed problem with nonlocal condition for certain pluriparabolic equations, *Hiroshima Math. J.* 27 (3), 373–390.
- Brusa L. and Nigro L., (1980), A One-Step Method for Direct Integration of Structural Dynamic Equation, *Int. J. Numer. Method Engg*, 15, 685-699.
- Butcher J. C, (2003), Numerical Methods for Ordinary Differential Equation, *John Wiley. Sons Ltd.*
- Calvo M. P. and Sanz-Serna J. M., (1993), High-order symplectic Runge-Kutta-Nyström methods, *SIAM J. Sci. Comput.*, 14(5), 1237-1252.
- Carlson D.E., (1972), Linear thermoelasticity, in: *Encyclopedia of Physics, Vol. 2, Springer, Berlin.*

- Chalco-Cano Y, Román-Flores H, (2006), On new solutions of fuzzy differential equations, *Chaos, Solitons and Fractals*, doi:10.1016/j.chaos.2006.10043.
- Chawla M. M. and Rao P. S., (1985), High-accuracy P -stable methods for $y'' = f(x, y)$, *IMA Journal of Numerical Analysis*, vol. 5, no. 2, 215–220.
- Che Jawias N. Izzati, Ismail F., Suleiman M. and Jaafar A., (2010), Fourth order four-stage diagonally implicit Runge-Kutta method for linear ordinary differential equations, *Malaysian Journal of Mathematical Sciences* 4(1), 95-105.
- Che Jawias N. Izzati, Ismail F., Suleiman M. and Jaafar A., (2011), Singly diagonally implicit fifth order five-stage Runge-Kutta method for linear ordinary differential equations, *Proceedings of the 8th WSEAS International Conference on Applied Computer and Applied Computational Science*, 82-86.
- Cushman J.H., Deng H. Xu, F., (1995), Nonlocal reactive transport with physical and chemical heterogeneity: Localization error, *Water Resource Res.* 31, 2219–2237.
- Cushman J.H., Ginn T.R., (1993), Nonlocal dispersion irregular media with continuously evolving scales of heterogeneity, *Transport Porous Media* 13, 123–138.
- Dormand J. R., (1996), Numerical methods for differential equations a computational approach, *CRC Press, Inc.*
- Dormand J. R., El-Mikkawy M.E.A. and Prince P. J., (1987), Families of Runge-Kutta-Nyström Formulae, *IMA J. Numer. Anal.*, 7, 235-250.
- Ferracina L., Spijker M.N., (2008), Strong stability of singly-diagonally-implicit Runge–Kutta methods, *Applied Numerical Mathematics* 58, 1675-1686.
- Franco J. M., (2003), A 5(3) pair of explicit ARKN methods for the numerical integration of perturbed oscillators, *Journal of Computational Applied Mathematics* 161 (2), 283–293.
- Franco J.M., (2006), A class of explicit two-step hybrid methods for second-order IVPs, *Journal of Computational Applied Mathematics* 187, 41–57.
- Gordeziani I. S. and Avalishvili G.A., (2000), On the constructing of solutions of the nonlocal initial boundary problems for one-dimensional medium oscillation equations, *Matem. Modelirovanie* 12 (1), 94–103.
- Hairer E., Wanner G., (2010), Solving ordinary differential equations II: stiff and differential-algebraic problems, Second Revised Edition, *Springer-Verlag Berlin Heidelberg*.
- Henrici P., (1962), Discrete Variable Methods in Ordinary Differential Equations. *John Wiley, New York, USA*.

- Hussian E. A. and Suhhiem M. H., (2015), Numerical Solution of Fuzzy Differential Equations Based on Taylor Series by Using Fuzzy Neural Networks, *Journal of Advances in Mathematics*, Vol. 11, No. 3, 4080-4093.
- Ixaru L. Gr., (1984), Numerical Methods for Differential Equations and Applications. *Reidel, Dordrecht-Boston-Lancaster*.
- Kaleva O, (2006), A note on fuzzy differential equations, *Nonlinear Analysis* 64, 895-900.
- Kavalloris N.I. and Tzanetis D.S., (2002), Behaviour of critical solutions of a nonlocal hyperbolic problem in ohmic heating of foods, *Appl. Math. E-Notes* 2, 59–65.
- Lambert J. D. and Watson I. A., (1976), Symmetric multistep methods for periodic initial-value problems, *Journal of the Institute of Mathematics and its Applications* 18, 189–202.
- Ma M, Freidman M, Kandel A, (1999), Numerical Solutions of fuzzy differential equations, *Fuzzy sets and Systems* 105, 133-138.
- Mohamad M., (2013), Explicit Runge-Kutta-Nyström methods with high order dispersion and dissipation for solving Oscillatory second order ordinary differential equation, *MSc. Thesis, Universiti Putra Malaysia*.
- Okunbor D. and Skeel R. D., (1992b), Explicit Canonical methods for Hamiltonian Systems, *Math Comp.*, 59, 439-455.
- Papadopoulos D. F., Anastassi Z. A. and Simos T. E. (2010), A modified phase-fitted and amplification-fitted Runge-Kutta-Nyström method for numerical solution of radial Schrodinger equation, *J. Mol Model*, 16, 1339-1346.
- Papadopoulos D. F., Anastassi Z. A., and Simos T. E., (2009), A phase-fitted Runge-Kutta Nyström method for the numerical solution of initial value problems with oscillating solutions, *Journal of Computer Physics Communications*, vol. 180, 1839-1846.
- Samarskii A.A., (1980), Some problems in modern theory of differential equations, *Differents. Uravn.* 16, 1221–1228.
- Senu N., Suleiman M., Ismail F. and Othman M., (2010), A fourth-order diagonally implicit Runge-Kutta- Nyström method with dispersion of high order, *ASM'10 Proceedings of the 4th International Conference on Applied Mathematics, simulation, modeling*, 78-82.
- Shakeri F., Dehghan M., (2008), The method of lines for solution of the one-dimensional wave equation subject to an integral conservation condition, *Comput. Math. Appl.* 56 (9), 2175–2188.

- Simos T. E. and Vigo-Aguiar J., (2000), A new modified Runge-Kutta-Nyström method with phase-lag of order infinity for the numerical solution of Schrodinger equation and related problems, *International Journal of Modern Physics C*, 11(6), 1195-1208.
- Simos T. E. and Vigo-Aguiar J., (2001), A modified phase-fitted Runge-Kutta method for numerical solution of Schrodinger equation, *Journal of Mathematical Chemistry*, 30(1), 121-131.
- Simos T. E., (1990), Numerical Solution of Ordinary Differential Equations with Periodical Solution. *Doctoral Dissertation, National Technical University of Athens*.
- Simos T. E., (1997), Eighth order method with minimal phase-lag for accurate computations for the elastic scattering phase-shift problem. *J. Math. Chem.* 21, 359–372.
- Simos T. E., Dimas E. and Sideridis A. B., (1994), A Runge-Kutta-Nyström method for the numerical integration of special second-order periodic initial-value problems, *Journal of Computational and Applied Mathematics*, 51, 317-326.
- Stefanini L, (2007), On generalized LU-fuzzy derivative and fuzzy differential equations, in: *IEEE International Fuzzy Systems Conference, FUZZ-IEEE, London, UK, 23-26 July 2007*, pp. 1-6.
- Stiefel E., Bettis D. G., (1969), Stabilization of Cowell's methods, *Numerische Mathematik* 13, 154–175.
- Thomas R. M., (1984), Phase Properties of High Order, almost P-stable Formulae, *BIT Numerical Mathematics*, 24(2), 225-238.
- Van de Vyver H., (2007), A symplectic Runge-Kutta- Nyström method with minimal phase-lag, *Physics Letters A* 367, 16-24.
- Van der Houwen P. J. and Sommeijer NB. P., (1987), Explicit Runge-Kutta (-Nyström) methods with reduced phase errors for computing oscillating solutions, *SIAM Journal on Numerical Analysis*, Vol.24, No.3, 595-617.
- Zingg D. W. and Chisholm T. T., (1999), Runge–Kutta methods for linear ordinary differential equations, *Applied Numerical Mathematics* 31, 227-238.