



UNIVERSITI PUTRA MALAYSIA

***WEIGHTED BLOCK RUNGE-KUTTA METHODS FOR SOLVING STIFF
ORDINARY DIFFERENTIAL EQUATIONS***

SAUFIANIM BINTI JANA AKSAH

FS 2016 22



**WEIGHTED BLOCK RUNGE-KUTTA METHODS FOR SOLVING STIFF
ORDINARY DIFFERENTIAL EQUATIONS**

By

SAUFIANIM BINTI JANA AKSAH

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Master of
Science**

May 2016

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

WEIGHTED BLOCK RUNGE-KUTTA METHODS FOR SOLVING STIFF ORDINARY DIFFERENTIAL EQUATIONS

By

SAUFIANIM BINTI JANA AKSAH

May 2016

Chairman : Zarina Bibi Binti Ibrahim, PhD
Faculty : Science

Weighted Block Runge-Kutta (WBRK) methods are derived to solve first order stiff ordinary differential equations (ODEs). The proposed methods approximate solutions at two points concurrently in a block at each step. Three sets of weight are chosen and implemented to the WBRK methods. Stability regions of the WBRK methods with each set of weight are constructed by using MAPLE14. Stability properties of the proposed methods with each weight show that the methods are suitable for solving stiff ODEs.

Numerical results are presented and illustrated in the form of efficiency curves. Performances of the WBRK methods in terms of maximum error and computational time are compared with the third order Runge-Kutta (RK3) and the modified weighted RK3 method based on centroidal mean (MWRK3CeM). These methods are tested with problems of single and system of first order stiff ODEs. Comparison of the proposed methods between sets of weight is also analyzed. The numerical results are obtained by using MATLAB R2011a.

Numerical results generated show that the WBRK methods obtained better accuracy and less computational time than the RK3 and MWRK3CeM method.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KAEDAH BLOK RUNGE-KUTTA BERPEMBERAT UNTUK
MENYELESAIKAN PERSAMAAN PEMBEZAAN BIASA KAKU**

Oleh

SAUFIANIM BINTI JANA AKSAH

Mei 2016

Pengerusi : Zarina Bibi Binti Ibrahim, PhD
Fakulti : Sains

Kaedah Blok Runge-Kutta Berpemberat (BRKB) dibina untuk menyelesaikan persamaan pembezaan biasa (PPB) kaku peringkat pertama. Kaedah yang dicadangkan ini mengangalkan penyelesaian pada dua titik serentak dalam satu blok pada setiap langkah. Tiga set pemberat dipilih dan dilaksanakan pada kaedah BRKB. Rantau kestabilan kaedah BRKB dengan setiap set pemberat dibina dengan menggunakan MAPLE14. Sifat kestabilan kaedah yang dicadangkan dengan setiap pemberat menunjukkan bahawa kaedah tersebut sesuai untuk menyelesaikan PPB kaku.

Keputusan berangka diberikan dan diilustrasikan dalam bentuk lengkungan kecekapan. Prestasi kaedah BRKB dalam bentuk ralat maksima dan masa pengiraan dibandingkan dengan kaedah Runge-Kutta peringkat ketiga (RK3) dan kaedah RK3 berpemberat yang diubah suai berdasarkan min sentroid (RK3PDMS). Kaedah-kaedah ini diuji dengan masalah PPB kaku peringkat pertama tunggal dan sistem. Perbandingan antara set-set pemberat kaedah yang dicadangkan juga dianalisis. Keputusan berangka diperoleh dengan menggunakan MATLAB R2011a.

Keputusan yang dihasilkan menunjukkan bahawa kaedah BRKB memperoleh ketepatan yang lebih baik dan masa pengiraan yang lebih cepat daripada kaedah RK3 dan RK3PDMS.

ACKNOWLEDGEMENTS

In the name of Allah, the most Compassionate and the most Merciful

First and foremost, I would like to express my sincere and deepest gratitude to the chairman of the supervisor committee, Assoc. Prof. Dr. Zarina Bibi Ibrahim for the valuable advice, encouragement and inspiration for me to generate ideas throughout this research. This research cannot be done without her constructive comments and suggestions. I am very grateful to the member of supervisory committee, Dr. Siti Nur Iqmal Ibrahim and also Pn. Yong Faezah Rahim for their guidance, support and knowledge to improve the research.

Special acknowledgement is extended to Dr. Sarkhosh for his idea and suggestion in developing the programs. Special thanks to my lab mates and friends in Universiti Putra Malaysia for their advices, reminder and motivation in order for me to complete the research.

I would like to thank Universiti Putra Malaysia for the Graduate Research Fellowship (GRF) and the Ministry of Higher Education for the MyBrain scholarship that make it possible for me to pursue my studies in Master Degree.

Finally, my deepest appreciation to my parents, Hamida Ahamed and Faizal Moideen; siblings, Mohd Saufi and Saufieanis for the continuous love, devotion, prayers and support. To those who contribute indirectly, may Allah reward you all. Thank you.

I certify that a Thesis Examination Committee has met on 12 May 2016 to conduct the final examination of Saufianim binti Jana Aksah on her thesis entitled "Weighted Block Runge-Kutta Methods for Solving Stiff Ordinary Differential Equations" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Mohamad Rushdan b Md Said, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Fudziah binti Ismail, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Rokiah @ Rozita Ahmad, PhD

Associate Professor
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 June 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Zarina Bibi Binti Ibrahim, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Siti Nur Iqmal Binti Ibrahim, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Saufianim Binti Jana Aksah, GS37889

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____




TABLE OF CONTENTS

		Page
ABSTRACT		i
ABSTRAK		ii
ACKNOWLEDGEMENTS		iii
APPROVAL		iv
DECLARATION		vi
LIST OF TABLES		x
LIST OF FIGURES		xi
LIST OF ABBREVIATIONS		xiii
CHAPTER		
1	INTRODUCTION	1
	1.1 Preliminaries	1
	1.2 Differential Equation	1
	1.3 Problem Statement	2
	1.4 Numerical Methods	2
	1.5 Runge-Kutta Methods	3
	1.6 Stability of Runge-Kutta Methods	5
	1.7 Stiff Systems of Ordinary Differential Equations	6
	1.8 Mean	7
	1.9 Objective of the Thesis	8
	1.10 Planning of the Thesis	8
 2	LITERATURE REVIEW	 10
	2.1 Introduction	10
	2.2 Runge-Kutta Methods	10
	2.3 Mean	11
	2.4 Block Methods	11
	2.5 Stiffness	12
 3	WEIGHTED BLOCK RUNGE-KUTTA METHOD FOR SOLVING STIFF ORDINARY DIFFERENTIAL EQUATIONS	 13
	3.1 Introduction	13
	3.2 Derivation of Weighted Block Runge-Kutta Methods	16
	3.3 Implementation of Sets of Weight	21
	3.3.1 Set of Weight, $w_1 = \frac{1}{2}; w_2 = \frac{1}{2}$	21
	3.3.2 Set of Weight, $w_1 = \frac{1}{4}; w_2 = \frac{3}{4}$	22

3.3.3	Set of Weight, $w_1 = \frac{2}{5}; w_2 = \frac{3}{5}$	23
4	STABILITY OF WEIGHTED BLOCK RUNGE-KUTTA METHODS	25
4.1	Introduction	25
4.2	Stability of the WBRK Methods	25
4.2.1	WBRK of Weights, $w_1 = \frac{1}{2}; w_2 = \frac{1}{2}$	26
4.2.2	WBRK of Weights, $w_1 = \frac{1}{4}; w_2 = \frac{3}{4}$	29
4.2.3	WBRK of Weights, $w_1 = \frac{2}{5}; w_2 = \frac{3}{5}$	34
4.3	Comparison of Stability of the WBRK Methods between Sets of Weight	37
5	NUMERICAL RESULTS	39
5.1	Introduction	39
5.2	Test Problems	40
5.3	Numerical Results for Comparison of Methods	43
5.3.1	Discussion	58
5.4	Numerical Results for Comparison of Weights	59
5.4.1	Discussion	74
6	CONCLUSION	76
6.1	Summary	76
6.2	Future Studies	77
	REFERENCES	78
	APPENDICES	83
	BIODATA OF STUDENT	88

LIST OF TABLES

Table	Page
3.1 Parameters for each sets of weight	24
4.1 Stability interval for each sets of weight	38
5.1 Numerical Results between methods for Problem 5.1	43
5.2 Numerical Results between methods for Problem 5.2	43
5.3 Numerical Results between methods for Problem 5.3	44
5.4 Numerical Results between methods for Problem 5.4	44
5.5 Numerical Results between methods for Problem 5.5	45
5.6 Numerical Results between methods for Problem 5.6	45
5.7 Numerical Results between methods for Problem 5.7	46
5.8 Numerical Results between methods for Problem 5.8	46
5.9 Numerical Results between methods for Problem 5.9	47
5.10 Numerical Results between methods for Problem 5.10	47
5.11 Numerical Results between sets of weight for Problem 5.1	59
5.12 Numerical Results between sets of weight for Problem 5.2	60
5.13 Numerical Results between sets of weight for Problem 5.3	60
5.14 Numerical Results between sets of weight for Problem 5.4	61
5.15 Numerical Results between sets of weight for Problem 5.5	61
5.16 Numerical Results between sets of weight for Problem 5.6	62
5.17 Numerical Results between sets of weight for Problem 5.7	62
5.18 Numerical Results between sets of weight for Problem 5.8	63
5.19 Numerical Results between sets of weight for Problem 5.9	63
5.20 Numerical Results between sets of weight for Problem 5.10	64

LIST OF FIGURES

Figure	Page
1.1 Trapezoid with Means	7
3.1 Block Explicit RK Method of Constant Stepsize	13
4.1 Stability region of WBRK with set of weight, $w_1 = \frac{1}{2}; w_2 = \frac{1}{2}$	28
4.2 Stability region of WBRK with set of weight, $w_1 = \frac{1}{4}; w_2 = \frac{3}{4}$	33
4.3 Stability region of WBRK with set of weight, $w_1 = \frac{2}{5}; w_2 = \frac{3}{5}$	36
4.4 Stability regions of WBRK with various sets of weight	38
5.1 Efficiency curves between methods for Problem 5.1	48
5.2 Efficiency curves between methods for Problem 5.2	49
5.3 Efficiency curves between methods for Problem 5.3	50
5.4 Efficiency curves between methods for Problem 5.4	51
5.5 Efficiency curves between methods for Problem 5.5	52
5.6 Efficiency curves between methods for Problem 5.6	53
5.7 Efficiency curves between methods for Problem 5.7	54
5.8 Efficiency curves between methods for Problem 5.8	55
5.9 Efficiency curves between methods for Problem 5.9	56
5.10 Efficiency curves between methods for Problem 5.10	57
5.11 Efficiency curves between sets of weight for Problem 5.1	64
5.12 Efficiency curves between sets of weight for Problem 5.2	65
5.13 Efficiency curves between sets of weight for Problem 5.3	66
5.14 Efficiency curves between sets of weight for Problem 5.4	67

5.15	Efficiency curves between sets of weight for Problem 5.5	68
5.16	Efficiency curves between sets of weight for Problem 5.6	69
5.17	Efficiency curves between sets of weight for Problem 5.7	70
5.18	Efficiency curves between sets of weight for Problem 5.8	71
5.19	Efficiency curves between sets of weight for Problem 5.9	72
5.20	Efficiency curves between sets of weight for Problem 5.10	73



LIST OF ABBREVIATIONS

DEs	Differential equations
ODEs	Ordinary differential equations
PDEs	Partial differential equations
IVP	Initial value problem
LMM	Linear multistep method
RK	Runge-Kutta
BDF	Backward differentiation formula
RK3	Third order Runge-Kutta
AM	Arithmetic mean
GM	Geometric mean
HaM	Harmonic mean
HeM	Heronian mean
CoM	Contraharmonic mean
RMS	Root mean square
CeM	Centroidal mean
WBRK	Weighted block Runge-Kutta
MWRK3CeM	Modified weighted third order Runge-Kuta based on centroidal mean
BDIRK	Block diagonally implicit Runge-Kutta
BBDF	Block backward differentiation formulae
RK3CeM	Third order Runge-Kutta based on centroidal mean
MAXE	Maximum error
TIME	Computational time

WBRKI Weighted block Runge-Kutta method with weights,
 $w_1 = \frac{1}{2}; w_2 = \frac{1}{2}$

WBRKII Weighted block Runge-Kutta method with weights,
 $w_1 = \frac{1}{4}; w_2 = \frac{3}{4}$

WBRKIII Weighted block Runge-Kutta method with weights,
 $w_1 = \frac{2}{5}; w_2 = \frac{3}{5}$

VSVO Variable step variable order

DDEs Delay differential equations

FDEs Fuzzy differential equations

CHAPTER 1

INTRODUCTION

1.1 Preliminaries

Many problems in engineering, physical and social sciences are reduced to quantifiable form through the process of mathematical modeling that involved differential equations (DEs). For instance, the spread of epidemic in a population. This model enables researchers to prove the famous “Threshold Theorem of Epidemiology” which states that an epidemic will occur only if the number of people susceptible to the disease exceeds a certain threshold value (Ismail, 1999). Another example is the design of catenary curve shape of the Gateway Arch in St. Louis which is based on the solution graph of DEs (Becker, 2012). These problems are very difficult and complicated to be solved analytically, therefore researchers used numerical methods to solve them.

1.2 Differential Equation

A DE is an equation that involves one or more derivatives of some unknown function or functions. There are various types of DEs. Some of the commonly known are ordinary differential equations (ODEs) and partial differential equations (PDEs). ODEs involve ordinary derivatives which contains a single independent variable and one or more dependent variables as shown

$$f\left(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots, \frac{d^n y}{dx^n}\right) = 0.$$

Usually, the independent variable is denoted by x and the dependent variable is y .

On the other hand, PDEs involve partial derivatives with more than one independent variable and one or more dependent variables of the form

$$f\left(x_1, \dots, x_n, y, \frac{\partial y}{\partial x_1}, \dots, \frac{\partial y}{\partial x_n}, \dots, \frac{\partial^2 y}{\partial x_1 \partial x_1}, \dots, \frac{\partial^2 y}{\partial x_1 \partial x_n}, \dots\right) = 0.$$

One characteristic of DE is the order which is the highest derivative appearing in the equation represent by n as shown in the above functions.

1.3 Problem Statement

This research focuses in solving problems of first order stiff ODEs of the form

$$y'(x) = f(x, y), \quad y(a) = \eta \quad (1.1)$$

where $y = [y_1(x), y_2(x), \dots, y_n(x)]^T$ and $f = [f_1(x, y), f_2(x, y), \dots, f_n(x, y)]^T$ with $\eta = [\eta_1, \eta_2, \dots, \eta_n]^T$ is a known vector.

Throughout this research, we shall assume that $f(x, y)$ satisfies the following theorem that guarantees the existence of a unique solution of Eq. (1.1) as stated by Lambert (1973).

Theorem 1.1

Let $f(x, y)$ be defined and continuous for all points (x, y) in the region D defined by $a \leq x \leq b$, $-\infty < y < \infty$, a and b finite, and let there exist a constant L such that, for every x, y, y^* such that (x, y) and (x, y^*) are both in D ,

$$|f(x, y) - f(x, y^*)| \leq L|y - y^*|. \quad (1.2)$$

Proof: see Henrici (1962).

Then, if η is any given number, there exists a unique solution $y(x)$ of the initial value problem (IVP) (1.1), where $y(x)$ is continuous and differentiable for all (x, y) in D . The requirement (1.2) is known as Lipschitz condition, and the constant L as a Lipschitz constant.

1.4 Numerical Methods

The exact solution of IVP (1.1) can be approximated by using numerical method. Generally, there are two classes of methods; a one-step method and a linear multistep method (LMM). One-step method uses the solution of current point for example, y_n , as initial value to compute solution at the next point, y_{n+1} . On the other hand, multistep method uses information from the previous steps to calculate the next value.

Euler's method is the easiest and widely used numerical method for solving IVP (1.1). It is written as

$$y_{n+1} - y_n = hf(x_n, y_n) \equiv hf_n.$$

The Euler's method is an explicit one-step method that does not require any additional starting values. It also readily permits a change of step length during the computation. The most famous family of one-step methods is Runge-Kutta (RK) methods.

By referring to IVP (1.1), we indicate the solution by $y(x)$. Next, we consider the sequence points $\{x_n\}$ defined by $x_n = a + nh$, $n = 0, 1, 2, \dots$ where h is the step size. Let y_n be an approximation to the theoretical solution x_n , that is, to $y(x_n)$, and let $f_n \equiv f(x_n, y_n)$. If a computational method for determining the sequence $\{y_n\}$ takes the form of a linear relationship between y_{n+i} and f_{n+i} with $i = 0, 1, \dots, k$, we call the method as LMM of step number k .

The general LMM can be written as

$$\sum_{i=0}^k \alpha_i y_{n+i} = h \sum_{i=0}^k \beta_i f_{n+i}$$

where α_i and β_i are constants. The method is explicit when $\beta_k = 0$ and implicit when $\beta_k \neq 0$. Iterative methods such as Newton's method are often used to solve the implicit formula.

In certain cases, an explicit LMM is used to "predict" the value of y_{n+i} . That value is then used in an implicit formula to "correct" the value. The result is a predictor-corrector method. There are three families of LMM; Adams-Bashforth methods, Adams-Moulton methods, and the backward differentiation formula (BDF). Further information on LMM can be found in Lambert (1973).

1.5 Runge-Kutta Methods

RK methods assumed that correct value of the slope over the step can be written as a linear combination of function $f(x, y)$ evaluated at certain points at the step. RK methods may be regarded as a particular case of the general explicit one-step method of the form

$$y_{n+1} - y_n = h\phi(x_n, y_n, h). \quad (1.3)$$

Definition 1.1

Eq. (1.3) is said to have order p if p is the largest integer for which

$$y(x+h) - y(x) - h\phi(x, y(x), h) = O(h^{p+1})$$

holds, where $y(x)$ is the theoretical solution of the IVP shown in Eq. (1.1).

Definition 1.2

Eq. (1.3) is consistent with the IVP (1.1) if

$$\phi(x, y, 0) \equiv f(x, y).$$

Definition 1.3

Eq. (1.3) is convergent by the following properties:

- (i) Let the function $\phi(x, y, h)$ be jointly continuous as a function of its three arguments, in the region D defined by $x \in [a, b]$, $y \in (-\infty, \infty)$, and $h \in [0, h_0]$ with $h_0 > 0$.
- (ii) Let $\phi(x, y, h)$ satisfy a Lipschitz condition of the form $|\phi(x, y^*, h) - \phi(x, y, h)| \leq M|y^* - y|$ for all points (x, y^*, h) and (x, y, h) in D .

Eq. (1.3) is convergent if and only if it is consistent (Lambert, 1973).

In addition, Lambert also considers that conditions (i) and (ii) are satisfied if the function satisfies the condition stated in Theorem 1.1. It is also mentioned that there is no requirement regarding the zero stability, since no parasitic solutions can arise with a one-step method.

In this research, we are using explicit third order RK (RK3) methods as the basic for development of the proposed method. The formula below shows the general form of classical RK3 methods as shown in Lambert (1973).

$$y_{n+1} - y_n = \frac{h}{6}(k_1 + 4k_2 + k_3), \quad (1.4)$$

where

$$\begin{aligned} k_1 &= f(x_n, y_n), \\ k_2 &= f\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}hk_1\right), \\ k_3 &= f(x_n + h, y_n - hk_1 + 2hk_2). \end{aligned}$$

1.6 Stability of Runge-Kutta Methods

The stability theory in numerical analysis is concerned with the growth of numerical errors in computed solution of the DE. The linear stability and accuracy of explicit RK methods are characterized completely by the stability function of the method, which in turn dictates the acceptable stepsize as stated by Butcher (2008b). The following stability analysis for explicit RK methods and proof for the stated theorem are shown in Hairer and Wanner (1991).

Definition 1.4

The function $R(z)$ is called the stability function of the method. It can be interpreted as the numerical solution after one step for

$$y' = \lambda y, \quad y_0 = 1, \quad z = h\lambda,$$

the famous Dahlquist test equation. The set

$$S = \{z \in \mathbb{C}; |R(z)| \leq 1\}$$

is called the stability domain of the method.

Theorem 1.2

If RK method is of order p , then

$$R(z) = 1 + z + \frac{z^2}{2!} + \dots + \frac{z^p}{p!} + O(z^{p+1})$$

with $h\lambda = z$.

As referred in Rahim (2004), absolute stability can also be defined as follows.

Definition 1.5

By letting $z = h\lambda$, then $R(z)$ is known as the stability function of the method.

Hence, $y_n \rightarrow 0$ as $n \rightarrow \infty$ if and only if

$$|R(z)| < 1 \quad (1.5)$$

and the method is absolutely stable for those value of z for which Eq. (1.5) holds. The region \mathfrak{R}_A of the complex z -plane which Eq. (1.5) holds is the region of absolute stability of the method.

Burden *et al.* (2015) stated that a method can be applied effectively to a stiff equation only if $h\lambda$ is in the region of absolute stability of the method which will results in a restriction on the size of h . The condition also applies for the approximation to decay to zero and the growth of error to be under control. The absolute criterion forces h to remain small.

1.7 Stiff Systems of Ordinary Differential Equations

Many fields of application, notably chemical engineering and control theory yield IVPs that sometimes exhibit a phenomenon known as stiffness (Akanbi *et al.*, 2011). There has been various definition of stiffness used by former researchers as mention in the literature with respect to the linear system of first order equation which defined as below

$$\tilde{y}' = A\tilde{y} + \tilde{\phi}(x), \quad \tilde{y}(a) = \tilde{\eta}, \quad a \leq x \leq b \quad (1.6)$$

where $\tilde{y}^T = (y_1, y_2, \dots, y_m)$, $\tilde{\eta}^T = (\eta_1, \eta_2, \dots, \eta_m)$, and A is an $m \times m$ matrix with the eigenvalues $\lambda_i, i = 1, 2, \dots, m$. In this research, we used the most widely accepted definition of stiffness given by Lambert (1973).

Definition 1.6

The linear system (1.6) is said to be stiff if

- (i) $\text{Re}(\lambda_i) < 0, i = 1, 2, \dots, m$ and
- (ii) $\max_i |\text{Re}(\lambda_i)| \gg \min_i |\text{Re}(\lambda_i)|$, where λ_i are the eigenvalues of A , and the ratio

$$S = \frac{\max_i |\text{Re}(\lambda_i)|}{\min_i |\text{Re}(\lambda_i)|}$$

is called the stiffness ratio. By following the definition

above, stiff problem has $S \gg 1$.

Next, we will discuss on mean and introduce some of the means that commonly used in mathematics. These means are then represented in a trapezoid to show the connection between each means.

1.8 Mean

According to Umberger (2000), there are many types of mean in mathematics represent by different equations. However, most people memorize these equations without making a relation to geometrical form of the means. Umberger had presented seven types of mean –arithmetic mean (AM), geometric mean (GM), harmonic mean (HaM), heronian mean (HeM), contraharmonic mean (CoM), root mean square (RMS) and centroidal mean (CeM)- and shows how to construct them in a trapezoid.

Figure below is an example of trapezoid containing all seven means.




Figure 1.1: Trapezoid with Means.

As for this research, we are focusing on CeM. CeM is the length of a segment which is parallel to the bases a and b of the trapezoid, and also passes through the centroid area of the trapezoid as shown in Figure 1.1. CeM of any two real numbers a and b can be represented by the following equation.

$$CeM = \frac{2(a^2 + ab + b^2)}{3(a + b)}.$$

A detailed step-by-step construction of each means in a trapezoid can be found in Umberger (2000).

1.9 Objective of the Thesis

In this research, we develop a block method based on RK method for solving first order stiff ODEs. The aims in developing a block method are to reduce the computational time and improve accuracy of the proposed methods when compared with existing methods.

Objectives of this thesis are

- i. to develop weighted block RK (WBRK) methods for solving first order stiff ODEs,
- ii. to implement the new developed methods on stiff ODEs problems in order to test the performance of the methods,
- iii. to construct the stability regions of the WBRK methods with different sets of weight and analyze their stability properties to indicate that the methods are suitable for solving first order stiff ODEs, and
- iv. to determine the accuracy and efficiency of the WBRK methods between weights and between the existing methods.

1.10 Planning of the Thesis

In Chapter 1, we give some introductions on DE that related to the research. Then, a brief statement regarding the problem that need to be solved is also included. Numerical methods that are commonly used in solving ODEs problems are discussed. This leads to introduction on the RK method and its stability properties. Definition of stiff system for solving ODEs is presented. Means in a trapezoid is defined in this chapter. The objective and planning of the thesis are also discussed.

In Chapter 2, we review on the literature of the classical RK methods and stiffness. Some historical background of mean and block methods related to the research are also discussed.

Derivation of CeM by Pushpam & Dhayabaran (2011), and modified weighted RK3 based on CeM (MWRK3CeM) method by Sharmila & Amirtharaj (2011) are discussed in the beginning of Chapter 3. Then, we present the modification of MWRK3CeM to form the new proposed methods called the WBRK methods. The general form of the methods is presented in the chapter. Implementation of the chosen sets of weight to the WBRK methods can be found in the subsections of Chapter 3.

In Chapter 4, we study the stability of the WBRK methods with different sets of weight. In order to do so, we construct the stability regions of the methods for each weight from the stability polynomial of the proposed methods. Then, we analyze the stability properties of each set of weight.

Numerical results are presented in Chapter 5. In the chapter, we present the problems of single and system of first order stiff ODEs. These problems are tested with different stepsizes to see the relationship between the stepsizes and the accuracy of the proposed methods. Comparisons are made with the RK3 method and MWRK3CeM in terms of maximum error and computational time of the proposed methods with the comparing methods to analyze the performances of the methods. The comparison is also made between the sets of weight implemented. Results are presented in table form and illustrated in efficiency curve. The results are analyzed and discussed.

Finally, Chapter 6 will summarize the findings of the research and suggestions for related future studies.

REFERENCES

- Ababneh, O. Y. & Ahmad, R. R. (2009). New Third Order Runge-Kutta Based on Contraharmonic Mean for Stiff Problems. *Applied Mathematical Sciences*, 3(8): 365-376.
- Ahmad, R. R. & Yaacob, N. (2005). Third Order Composite Runge-Kutta Method for Stiff Problems. *International Journal of Computer Mathematics*, 2(10): 1221-1226.
- Akanbi, M. A., Okunuga, S. A., & Sofoluwe, A. B. (2011). Numerical Treatment of Kap's Equation using Fourth Order Method. *International Journal of the Physical Sciences*, 6(15): 3827-3833.
- Becker, L. C. (2012). *Ordinary Differential Equations: Concepts, Methods, and Models*. Tennessee: Christian Brothers University Memphis.
- Berland, H. (2007). Explicit Numerical Methods Experiencing Instability on a Stiff Equation.
- Burden, R.L. & Faires, J.D. (2001). *Numerical Analysis*. Brooks: Cole-Thomson Learning.
- Burden, R. L., Faires, J. D., & Burden, A. M. (2015). *Numerical Analysis (10th ed.)*. Brooks: Cole-Thomson Learning.
- Butcher, J. C. (1963). Coefficients for the Study of Runge-Kutta Integration Process. *Journal of Australia Mathematical Society*, 3(2): 185-201.
- Butcher, J. C. (2008). Order and Stability of Generalized Pade Approximations. *Applied Numerical Mathematics*, 59(3-4): 558-567.
- Butcher, J. C. (2008). *Numerical Methods for Ordinary Differential Equations*. England: John Wiley & Sons.
- Cash, J. R. (1975). A Class of Implicit Runge-Kutta Methods for the Numerical Integration of Stiff Ordinary Differential Equations. *Journal of Association for Computing Machinery*, 22(4): 504-511.
- Cash, J. R. (1983). Block Runge-Kutta Method for the Numerical Integration of IVP. *Mathematics of Computation*, 40(161): 175-191.
- Cash, J. R. (1985). Block Embedded Explicit Runge-Kutta Methods. *Computers and Mathematics with Applications*, 11(4): 395-409.
- Cash, J. R. & Vieira, I. (1995). Block 6(5) and 7(6) Explicit Runge-Kutta Formulae. *Journal of Applied Numerical Mathematics*, 17(3): 223-234.

- Cash, J. R. (2003). Efficient Numerical Methods for the Solution of Stiff Initial-Value Problems and Differential Algebraic Equations. *Proceedings of the Royal Society a Mathematical Physical and Engineering Sciences*, 459(2032): 797-815.
- Cheney, W. & Kincaid, D. (1999). *Numerical Mathematics and Computing*. Brooks: Cole-Thomson Learning.
- Enright, W. H., Higham, D. J., Owren, B., & Sharp, P. W. *A Survey of the Explicit Runge-Kutta Method (1994)*. Technical Report, 291/94, Department of Computer Science, University of Toronto: Toronto.
- Evans, D. J. & Yaacob, N. (1995). A Fourth Order Runge-Kutta Method Based on the Heronian Mean. *International Journal of Computer Mathematics*, 58: 103-115.
- Evans, D. J. & Yaakub, A. R. (1998). Weighted Fifth Order Runge-Kutta Formulas for Second Order Differential Equations. *International Journal of Computer Mathematics*, 70(2): 233-239.
- Faradj, M. K. (2004). *Which Do You Mean? An Exposition on Means*, Master Thesis, Louisiana State University.
- Gear, C. W. (1971). *Numerical Initial Value Problems in Ordinary Differential Equations*. New Jersey: Prentice Hall, Inc.
- Gear, C. W. (1980). Runge-Kutta Starter for Multistep Method. *ACM Transactions on Mathematical Software*, 6(3): 263-279.
- Hairer, E. & Wanner, G. (1974). On The Butcher Group and General Multi-Value Methods, *Computing*, 13(1): 1-15.
- Hairer, E. & Wanner, G. (1991). *Solving Ordinary Differential Equations II: Stiff and Differential Algebraic Problems*. US: Springer.
- Henrici, P. (1962). *Discrete Variable Methods in Ordinary Differential Equations*. New York: John Wiley and Sons.
- Heun, K. (1900). Neue Methoden zur approximativen Integration der Differentialgleichungen einer unabhängigen Veränderlichen. *Zeitschrift für angewandte Mathematik und Physik*, 45: 23–38.
- Huang, S. J. Y. (2005). *Implementation of General Linear Methods for Stiff ODEs*, PhD Thesis, University of Auckland.
- Ibrahim, Z. B. (2006). *Block Multistep Methods for Solving Ordinary Differential Equations*, PhD Thesis, Universiti Putra Malaysia.

- Ibrahim, Z. B., Othman, K. I., & Suleiman, M. B. (2007). Variable Step Size Block Backward Differentiation Formula for Solving Stiff ODEs. *Proceedings of World Academy of Science, Engineering and Technology*, 2: 785-789.
- Ismail, F. (1999). *Numerical Solution of Ordinary and Delay Differential Equations by Runge-Kutta Type Methods*, PhD Thesis, Universiti Putra Malaysia.
- Jackiewicz, Z., Renaut, R. A., & Zennaro, M. (1995). Explicit Two-Step Runge-Kutta Methods. (English). *Applications of Mathematics*, 40(6): 433-456.
- Johnson, A. I. & Barney, J. R. (1976). Numerical Solution of Large Systems of Stiff Ordinary Differential Equations in a Modular Simulation Framework. In Lapidus L. and Schiesser W.E. (Ed.), *Numerical Methods for Differential Systems: Recent Developments in Algorithms, Software and Applications*. New York: Academic Press Inc.
- Kutta, W. (1901). Beitrag zur näherungsweise Integration totaler Differentialgleichungen. *Zeitschrift für angewandte Mathematik und Physik*, 46: 435–453.
- Lambert, J. D. (1973). *Computational Methods in Ordinary Differential Equations*. England: John Wiley & Sons.
- LeVeque, R. J. (2007). Absolute Stability for Ordinary Differential Equations [Electronic version], Society for Industrial and Applied Mathematics. <https://www.siam.org/books/ot98>.
- Majid, Z. A., Suleiman, M. B., & Omar, Z. (2006). *Bulletin of the Malaysian Mathematical Sciences Social*, 29(1): 23-31.
- Milne, W. E. (1953). *Numerical Solution of Differential Equations*. New York: John Wiley and Sons.
- Murugesan, K., Dhayabaran, D. P., Amirtharaj, E. C. H., & Evans, D. J. (2001). A Comparison of Extended Runge-Kutta Formulae based on Variety of Means to Solve System of IVPs. *International Journal of Computational Mathematics*, 78(2): 225-252.
- Musa, H., Suleiman, M. B., & Senu, N. (2012). A-stable 2-Point block extended backward differentiation formulas for stiff ordinary differential equations. *Proceedings of the 5th International Conference on Research and education in Mathematics*, 1450: 254–258.
- Nasir, N. A. A. M. (2011). *Multiblock Backward Differentiation Formulae for Solving First Order Ordinary Differential Equations*, Master Thesis, Universiti Putra Malaysia.

- Novati, P. (2003). An Explicit One-Step Method for Stiff Problems. *Computing*, 71(2003): 133-151.
- Piché, R. Numerical Differential Equation Solvers [Electronic version], Tampere University of Technology. <http://icosym-nt.cvut.cz/odl/partners/tut/unit2/>.
- Pushpam, A. E. K. & Dhayabaran, D. P. (2011). Comparison of Single Term Walsh Series Technique and Extended RK Methods Based on Variety of Means to Solve Stiff Non-linear Systems. *Recent Research in Science and Technology*, 3(9): 22-30.
- Rahim, Y. F. (2004). *Block Diagonally Implicit Runge-Kutta Method for Solving Ordinary Differential Equations*, Master Thesis, Universiti Putra Malaysia.
- Rosser, J. B. (1967). A Runge-Kutta for All Season. *SIAM Review*, 9(3): 417-452.
- Runge, C. (1895). Über die numerische Auflösung von Differentialgleichungen. *Mathematische Annalen*, 46(2): 167-178.
- Sanugi, B. B. & Evans, D. J. (1994). A New Fourth Order Runge-Kutta Method Based on Harmonic Mean, *International Journal of Computer Mathematics*, 50(1-2): 113-118.
- Sanugi, B. B. & Yaacob, N. A. (1995). New Fifth Order Five-Stage Runge-Kutta Method for Initial Value Type Problems in ODEs. *International Journal of Computer Mathematics*, 59(3-4): 187-207.
- Sarafyan, D. *Multistep Methods for the Numerical Solution of Ordinary Differential Equations Made Self-Starting (1965)*. Technical Report No. 495, Mathematics Research Centre: Madison.
- Shampine, L. F. & Gear, C. W. (1979). A User's View of Solving Stiff Ordinary Differential Equations. *SIAM Review*, 21 (1): 1-17.
- Sharmila, R. G. & Amirtharaj, E. C. H. (2011). Implementation of A New Third Order Weighted Runge-Kutta Formula Based on Centroidal Mean for Solving Stiff Initial Value Problems. *Recent Research in Science and Technology*, 3(10): 91-97.
- Sharp P. W. & Verner J. H. (1998). Generation of High-Order Interpolants for Explicit Runge-Kutta Pairs. *ACM Transactions on Mathematical Software*, 24(1): 13-29.
- Thanh, N. V. (2009). Approximation Theory [Electronic version], Institute of Physics. http://iop.vast.ac.vn/~nvthanh/cours/numerical_methods.
- Umberger, S. (2000). Some "Mean" Trapezoids [Electronic version], The University of Georgia. <http://jwilson.coe.uga.edu>.

Wazwaz, A. M. (1990). A Modified Third Order Runge-Kutta Method. *Applied Mathematics Letter*, 3(3): 123-125.

Wu, X. Y. (1998). A Sixth-Order A-Stable Explicit One-Step Method for Stiff Systems. *Computers and Mathematics with Application*, 35(9): 59-64.

Zainuddin, N., Ibrahim, Z. B., Suleiman, M., & Othman, K. I. (2011). *Two-Point Block Backward Differentiation Formula for Solving Higher Order Ordinary Differential Equations*, Master Thesis, Universiti Putra Malaysia.

