



UNIVERSITI PUTRA MALAYSIA

***OPTIMISATION ON BIODEGRADATION OF GLYPHOSATE BY
LOCALLY ISOLATED BACTERIA***

MOTHARASAN MANOGARAN

FBSB 2018 34



**OPTIMISATION ON BIODEGRADATION OF GLYPHOSATE BY LOCALLY-
ISOLATED BACTERIA**

By

MOTHARASAN MANOGARAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirement for the Degree of Master of Science**

November 2017

COPYRIGHT

All materials contained within the thesis including without limitation text, logos, icons, photographs and all other artworks are copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from copyright holder. Commercial use of materials may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This thesis is dedicated to my parents, farmers, pioneers and fellow researchers.



© COPYRIGHT UPM

Abstract of thesis presented to Senate of Universiti Putra Malaysia in fulfillment
of the requirements for the degree of Master of Science

OPTIMISATION ON BIODEGRADATION OF GLYPHOSATE BY LOCALLY- ISOLATED BACTERIA

By

MOTHARASAN MANOGARAN

November 2017

Chairman: Siti Aqlima Binti Ahmad, PhD
Faculty : Biotechnology and Biomolecular Sciences

N-Phosphonomethylglycine (glyphosate) is often used to control weeds in agriculture land. In Malaysia alone, thousands of tonnes of glyphosate formulation is used annually. Although this low-cost herbicide is able to effectively in killing weeds, its extensive use has been negatively linked to the human health due to its toxicity. Moreover, many past studies have reported its negative effects on aquatic animals and vertebrates. Thus, there is a need for an eco-friendly method to manage this environmental contaminant. Furthermore, it is crucial to find glyphosate-degrading microorganisms in the soil of interest for local applications. The objective of this study is to isolate local glyphosate-degrading bacteria and optimisation of culture medium condition to improve the degradation rate. Two bacteria isolated from agriculture site located in Kedah, Malaysia were identified by physical, biochemical and 16S rRNA sequencing techniques as *Burkholderia vietnamiensis* strain AQ5-12 and *Burkholderia* sp. strain AQ5-13. These strains were found with the ability to tolerate up to 12 mL/L Roundup concentration and were successfully used to degrade glyphosate. Factors affecting glyphosate biodegradation such as carbon and nitrogen sources, pH of the medium, glyphosate concentration and temperature were optimised using one factor at time (OFAT) and response surface method (RSM) using free cells. Initial free cells of strain AQ5-12 and AQ5-13 were able to degrade 79.7% and 40.67% of 50 ppm glyphosate, respectively, within 24 h incubation. Using these optimisation processes, free cells of AQ5-12 were able to degrade 94% of 100 ppm glyphosate, whereas strain AQ5-13 degraded 94% of 50 ppm glyphosate under optimal condition. The results illustrated fructose at 8.62 g/L, ammonium sulphate at 0.5 g/L, pH 5.41, 100 ppm of glyphosate concentration and 32°C as the optimum biodegradation conditions required by *Burkholderia vietnamiensis* strain AQ5-12, whereas the optimum biodegradation conditions for *Burkholderia* sp. strain AQ5-13 were sucrose at 8.0 g/L, ammonium sulphate at 0.5 g/L, pH 6.0, 50 ppm of glyphosate concentration and temperature at 32°C. The optimised condition for free cells resulted in significant improvement in degradation rate. The bacteria were immobilised in gellan gum gelling agent with its conditions optimised. The results presented a degradation rate of 87.2% 100 ppm glyphosate with gellan gum concentration of 0.55 g, 285 number of beads and bead size of 0.48 cm for immobilised cells of AQ5-12.

Meanwhile, immobilised cells of AQ5-13 illustrated a degradation rate of 96.68% 50 ppm glyphosate with gellan gum concentration of 0.55 g, 280 number of beads and bead size of 0.45 cm. It was seen that immobilised form of bacteria showed better biodegradation in terms of duration as it degrades the glyphosate within 12 h compared to free cells that require 24 h degradation process in optimised media. In conclusion, these strains possess the potential of being used in the management of glyphosate contamination. Furthermore, the success of isolating bacteria from local soils in Malaysia has shown prominent ability in glyphosate degradation rate, which can be applied for glyphosate treatment in agricultural land.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Sarjana Sains

BIOPENURUNAN DAN PENGOPTIMUMAN GLIFOSAT OLEH PEMENCILAN BAKTERIA TEMPATAN

Oleh

MOTHARASAN MANOGARAN

November 2017

Pengerusi : Siti Aqlima Binti Ahmad, PhD
Fakulti : Bioteknologi dan Sains Biomolekul

N-Phosphonomethylglycine (glyphosate) sering digunakan untuk mengawal rumpai di tanah pertanian. Di Malaysia sahaja, beribu-ribu tan formulasi glifosat digunakan setiap tahun. Walaupun racun herba murah ini berkeupayaan berkesan dalam membunuh rumpai, penggunaannya yang luas telah dikaitkan secara negatif dengan kesihatan manusia akibat keracunannya. Selain itu, banyak kajian lepas telah melaporkan kesan negatif ke atas haiwan akuatik dan vertebrata. Oleh itu, terdapat keperluan untuk kaedah yang mesra alam untuk menguruskan pencemaran alam sekitar ini. Tambahan pula, adalah penting untuk mencari mikroorganisma yang menurunkan glifosat di tanah yang menarik untuk aplikasi tempatan. Oleh demikian, objektif kajian ini adalah untuk mendapatkan bakteria tempatan yang boleh mendegradasikan glifosat dan pengoptimuman media untuk meningkatkan kadar degradasi. Dua bakteria yang dipencilkan dari tapak pertanian yang terletak di Kedah, Malaysia telah dikenal pasti dengan teknik penjujukan fizikal, biokimia dan 16S rRNA sebagai *Burkholderia vietnamiensis* strain AQ5-12 dan *Burkholderia* sp. strain AQ5-13. Strain ini didapati dengan keupayaan untuk bertolak ansur sehingga kepekatan 12 mL/L Roundup dan berjaya digunakan untuk merendahkan glifosat. Faktor-faktor yang mempengaruhi biopenurunan glifosat seperti sumber karbon dan nitrogen, pH medium, kepekatan glifosat dan suhu dioptimumkan dengan menggunakan satu faktor pada waktu (OFAT) dan kaedah permukaan tanggapan (RSM) menggunakan sel bebas. Sel-sel bebas awal strain AQ5-12 dan AQ5-13 mampu menurunkan 79.7% dan 40.67% glifosat 50 ppm, masing-masing, dalam pengeraman 24 jam. Dengan menggunakan proses pengoptimuman ini, sel-sel bebas AQ5-12 mampu menurunkan 94% daripada 100 ppm glifosat, manakala strain AQ5-13 menurunkan 94% daripada 50 ppm glifosat dalam keadaan optimum. Keputusan menggambarakan fruktosa pada 8.62 g/L, ammonium sulfat pada 0.5 g/L, pH 5.41, 100 ppm kepekatan glyphosate dan 32°C sebagai syarat biopenurunan optimum yang diperlukan oleh *Burkholderia vietnamiensis* strain AQ5-12, manakala keadaan biodegradasi optimum untuk *Burkholderia* sp. strain AQ5-13 adalah sukrosa pada 8.0 g/L, ammonium sulfat pada 0.5 g/L, pH 6.0, 50 ppm kepekatan glikosat dan suhu pada 32°C. Keadaan yang dioptimumkan untuk sel-sel bebas menghasilkan peningkatan yang ketara dalam kadar penurunan. Bakteria telah diabadikan dalam agen gel gellan gum dengan keadaannya dioptimumkan. Hasilnya

memperlihatkan kadar penurunan 87.2% 100 ppm glifosat dengan kepekatan gellan gum 0.55 g, 285 bilangan manik dan saiz manik 0.48 cm untuk sel sekat gerak AQ5-12. Sementara itu, sel sekat gerak AQ5-13 menggambarkan kadar kemerosotan 96.68% 50 ppm glyphosate dengan kepekatan gellan gum 0.55 g, 280 bilangan manik dan saiz manik 0.45 cm. Ia telah memperlihatkan bahawa bentuk sekat gerak bakteria menunjukkan biopenurunan yang lebih baik dari segi tempoh kerana ia menurunkan glifosat dalam masa 12 jam berbanding dengan sel-sel bebas yang memerlukan 24 jam proses penurunan dalam media yang dioptimumkan. Kesimpulannya, strain ini mempunyai potensi untuk digunakan dalam pengurusan pencemaran glifosat. Tambahan pula, kejayaan memencilkan bakteria dari tanah tempatan di Malaysia telah menunjukkan keupayaan menonjol dalam kadar penurunan glifosat, yang boleh digunakan untuk rawatan glifosat di tanah pertanian.



ACKNOWLEDGEMENTS

The completion of this work was made possible with the help, assistance and aids from many important people. First foremost, I would like express my special appreciation and profound gratitude to my main supervisor, Dr. Siti Aqlima Ahmad for her kindness, hands-on support, willingness to teach and supervision during my period of study. I was also indebted to the management of Universiti Putra Malaysia for selecting and sponsoring me to conduct my M.Sc. Special thanks to the government of Malaysia, for providing MyBrain15 scholarship. My special thanks also goes to co-supervisors Associate Prof. Dr. Mohd Yunus Abd Shukor, Dr. Adeela Yasid, Dr. Khalilah Abdul Khalil and Dr. Wan Lutfi Wan Johari who gave me all the needed support, guide, hints and prompt attention that keep me on track throughout my research.

I am highly grateful to my parents, Manogaran Ratnam and Janakei Sangunni Nair for their love, support and prayers during the course of this research. My appreciation to my beloved sisters, Kokilah and Baarathyee for their moral support. Thanks to Mdm. Hasdianty Abdullah for her guidance during my degree time. Without their advice and support, my path for master would be incomplete.

I also wish to express my appreciation to all my colleagues not forgetting Dr. Ibrahim Yusof, Dr. Kabiru Karamba, Dr. Abubakar, Syahir Habib, Siti Zeera, Shakirah Sha'arani, Ain Aqilah and members of Bioremediation group for the enormous contributions they rendered during the conduct of this thesis. No amount of thanks is comparable with their efforts.

Last and not the least my brothers from another mother, Logesh Chinnatambhi, Danesh Anathan and Vinod Selavadurai for always keeping my spirits up. They all kept me going, and this thesis would not been possible without them.

"It's the job that's never started as takes longest to finish."

Motharasan Manogaran, 2017

I certify that a Thesis Examination Committee has met on 3 November 2017 to conduct the final examination of Motharasan a/l Manogaran on his thesis entitled "Optimisation on Biodegradation of Glyphosate by Locally-Isolated Bacteria" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Suhaimi bin Napis, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Phang Lai Yee, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Siti Aisah Alias, PhD

Associate Professor
University of Malaya
Malaysia
(External Examiner)



NOR AINI AB. SHUKOR, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 29 January 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Siti Aqlima Ahmad, PhD

Senior Lecturer,
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairperson)

Mohd Yunus Abd. Shukor, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Nur Adeela Yasid, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Wan Lutfi Wan Johari, PhD

Senior Lecturer
Faculty of Environmental Studies
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writing, seminar paper, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Motharasan a/l Manogaran, GS43841

Declaration by Members of Supervisory committee

This is to certify that:

- the research conducted and the writing of the thesis was under our supervision
- supervision of responsibilities as slated in Rule 41 in Rules 2003 (Revision 2012 – 2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Dr. Siti Aqlima Ahmad

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Nur Adeela Yasid

Signature: _____
Name of Member
of Supervisory
Committee: Associate Professor
Dr. Mohd Yunus Shukor

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Wan Lutfi Wan Johari

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENT	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xvi
LIST OF FIGURES	xix
LIST OF ABBREVIATIONS	xxiii
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	3
2.1 History of glyphosate	3
2.2 Properties of glyphosate	3
2.3 Mode of action	4
2.4 Uptake and translocation	6
2.5 Glyphosate resistance in Malaysia	6
2.6 Degradation and persistence in soil	8
2.7 Toxicity of polyethoxylated tallow amine	8
2.8 Effects on non-target aquatic plants and algae	8
2.9 Effects on non-target aquatic vertebrates	11
2.10 Glyphosate metabolism	11
2.11 Microbial degradation	13
2.12 Chemical degradation	15
2.13 Bioremediation	15
2.14 Methods of optimisation in glyphosate degradation using experimental design and statistical approach	16
2.15 Application of immobilised cells for glyphosate degradation	18
3 MATERIALS AND METHODS	22
3.1 Experimental design	22
3.2 Chemicals, reagents and equipment	23
3.3 Analytical method	23
3.3.1 Preparation of standard phosphate assay	23
3.3.2 Preparation of ascorbic reagent	24
3.3.3 Procedure for glyphosate assay	25
3.4 Isolation of glyphosate-degrading bacteria	25

3.4.1	Sampling	25
3.4.2	Handling and growth of bacterial isolates	26
3.5	Screening for glyphosate-degrading bacteria	27
3.5.1	Primary screening	27
3.5.2	Secondary screening	27
3.5.3	Tertiary screening	27
3.5.4	Quaternary screening	28
3.5.5	Identification of glyphosate-degrading bacteria	28
3.5.5.1	Gram staining	28
3.5.5.2	Oxidase test	28
3.5.5.3	Catalase test	28
3.5.5.4	Voges-Proskauer test	29
3.5.5.5	Nitrate reduction test	29
3.5.5.6	Citrate test	29
3.5.5.7	Indole production test	30
3.5.5.8	Urease test	30
3.5.5.9	Sugar tests	30
3.5.5.10	Molecular identification	30
3.6	Preparation of resting cells	32
3.7	Optimisation of degradation and growth conditions using OFAT approach by free cells	32
3.7.1	Effect of carbon source	33
3.7.2	Effect of nitrogen source	33
3.7.3	Effect of pH	33
3.7.4	Effect of temperature	33
3.7.5	Effect of glyphosate concentration	34
3.8	Optimisation of degradation and growth conditions using RSM by free cells	34
3.8.1	Plackett-Burman design	34
3.8.2	Central composite design	36
3.8.3	Verification of RSM for free cells	38
3.9	Optimisation of degradation using OFAT approach by immobilised cells	38
3.9.1	Characterisation of immobilised cells protocols	38
3.9.2	Effect of gellan gum concentration	38
3.9.3	Effect of beads size	39
3.9.4	Effect of number of beads	39
3.10	Optimisation of degradation using RSM by immobilised cells	39
3.10.1	Plackett-Burman design	39
3.10.2	Central composite design	41

	3.10.3	Verification of RSM for immobilised cells	42
	3.10.4	Reusability of beads	42
	3.11	Statistical analysis	43
4		RESULT AND DISCUSSION	44
	4.1	Isolation of glyphosate-degrading bacteria	44
	4.2	Biodegradation of glyphosate	44
	4.3	Screening of glyphosate-degrading bacteria	45
	4.3.1	Primary screening	45
	4.3.2	Secondary screening	46
	4.3.3	Tertiary screening	47
	4.3.4	Quaternary screening	49
	4.4	Identification of glyphosate-degrading bacteria	50
	4.4.1	Gram staining	50
	4.4.2	16S rRNA analysis	51
		4.4.2.1 Genomic extraction	51
		4.4.2.2 Polymerase chain reaction	52
	4.4.3	Phylogenetic analysis	52
	4.4.4	Biochemical test	59
	4.5	Optimisation via OFAT using free cells	61
	4.5.1	Effects of carbon source	61
		4.5.1.1 Effects of different carbon sources on biodegradation of 50 ppm glyphosate by <i>Burkholderia vietnamiensis</i> strain AQ5-12	61
		4.5.1.2 Effects of different fructose concentration on biodegradation of 50 ppm glyphosate by <i>Burkholderia vietnamiensis</i> strain AQ5-12	62
		4.5.1.3 Effects of different carbon sources on biodegradation of 50 ppm glyphosate by <i>Burkholderia</i> sp. strain AQ5-13	63
		4.5.1.4 Effects of different sucrose concentration on biodegradation of 50 ppm glyphosate by <i>Burkholderia</i> sp. strain AQ5-13	64
	4.5.2	Effects of nitrogen source	65
		4.5.2.1 Effects of different nitrogen sources on biodegradation of 50 ppm glyphosate by <i>Burkholderia vietnamiensis</i> strain AQ5-12	66
		4.5.2.2 Effects of different ammonia sulphate concentration on	67

	biodegradation of 50 ppm glyphosate by <i>B. vietnamiensis</i> strain AQ5-12	
4.5.2.3	Effects of different nitrogen sources on biodegradation of 50 ppm glyphosate by <i>Burkholderia</i> sp. strain AQ5-13	68
4.5.2.4	Effects of different ammonia sulphate concentration on biodegradation of 50 ppm glyphosate by <i>Burkholderia</i> sp. strain AQ5-13	69
4.5.3	Effects of pH	70
4.5.3.1	Effects of pH on biodegradation of 50 ppm glyphosate by <i>B. vietnamiensis</i> strain AQ5-12	70
4.5.3.2	Effects of pH on biodegradation of 50 ppm glyphosate by <i>Burkholderia</i> sp. strain AQ5-13	72
4.5.4	Effect of temperature	73
4.5.4.1	Effects of different temperature on biodegradation of 50 ppm glyphosate by <i>Burkholderia vietnamiensis</i> strain AQ5-12	73
4.5.4.2	Effects of different temperature on biodegradation of 50 ppm glyphosate by <i>Burkholderia</i> sp. strain AQ5-13	74
4.5.5	Effect of glyphosate concentration	75
4.5.5.1	Effects of different glyphosate concentration on biodegradation of glyphosate by <i>Burkholderia vietnamiensis</i> strain AQ5-12	75
4.5.5.2	Effects of different glyphosate concentration on biodegradation of 50 ppm glyphosate by <i>Burkholderia</i> sp. strain AQ5-13	77
4.6	Optimisation via RSM using free cells	78
4.6.1	Plackett-Burman	78
4.6.1.1	Plackett-Burman design for <i>Burkholderia vietnamiensis</i> strain AQ5-12	78
4.6.1.2	Plackett-Burman for <i>Burkholderia</i> sp. AQ5-13	80
4.6.2	Central composite design	82
4.6.2.1	Central composite design for <i>Burkholderia vietnamiensis</i> strain AQ5-12	82
4.6.2.2	Central composite design for <i>Burkholderia</i> sp. strain AQ5-13	88
4.6.3	Verification of RSM experiment	95

	4.6.3.1	Verification of RSM experiment for <i>Burkholderia vietnamiensis</i> strain AQ5-12	95
	4.6.3.2	Verification of RSM experiment for <i>Burkholderia</i> sp. strain AQ5-13	96
4.7		Immobilisation of bacterial cells	98
4.8		Optimisation via OFAT using immobilised cells	98
	4.8.1	Effects of gelling components composition	98
	4.8.1.1	Effect of different gellan gum concentrations on biodegradation of 100 ppm glyphosate using immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12	98
	4.8.1.2	Effect of different gellan gum concentrations on biodegradation of 50 ppm glyphosate using immobilised <i>Burkholderia</i> sp. strain AQ5-13	99
	4.8.2	Effects of beads size	100
	4.8.2.1	Effect of different beads sizes on biodegradation of 100 ppm glyphosate using immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12	101
	4.8.2.2	Effects of different beads sizes biodegradation of 50 ppm glyphosate using immobilised <i>Burkholderia</i> sp. strain AQ5-13	102
	4.8.3	Effects of cell loading (number of beads)	102
	4.8.3.1	Effect of different beads number on biodegradation of 100 ppm glyphosate using immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12	103
	4.8.3.2	Effect of different beads number biodegradation of 50 ppm glyphosate using immobilised <i>Burkholderia</i> sp. strain AQ5-13	104
4.9		Optimisation via RSM using immobilised cells	105
	4.9.1	Plackett-Burman	105
	4.9.1.1	Plackett-Burman design for immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12	105
	4.9.1.2	Plackett-Burman design for immobilised <i>Burkholderia</i> sp. strain AQ5-13	106
	4.9.2	Central composite design	108

4.9.2.1	Central composite design for immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12	108
4.9.2.2	Central composite design for immobilised <i>Burkholderia</i> sp. strain AQ5-13	114
4.9.2.3	Limitation in bead parameters	119
4.9.3	Verification of RSM experiment	121
4.9.3.1	Verification of RSM experiment for immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	121
4.9.3.2	Verification of RSM experiment for immobilised beads of <i>Burkholderia</i> sp. strain AQ5-13	123
4.10	Reusability of immobilised beads	124
4.11	Summary on the overall study on biodegradation of glyphosate by locally- isolated bacteria	126
5	CONCLUSION AND RECOMMENDATIONS	128
	REFERENCES	130
	APPENDICES	143
	BIODATA OF STUDENT	148
	LIST OF PUBLICATIONS	149

LIST OF TABLES

Table	Page
2.1 Glyphosate properties.	4
2.2 Known herbicide resistant weeds in Malaysia.	7
2.3 Effects of glyphosate, different commercial formulations of glyphosate and POEA on algae and aquatic plants.	10
2.4 List of known glyphosate-degrading bacteria.	14
2.5 Advantage and disadvantage of materials used as carrier for cell immobilisation.	18
3.1 Calculation for 1 g of PO ₄ ³ in glyphosate, monopotassium phosphate and disodium phosphate.	24
3.2 Preparation of standard phosphate solution in ppm.	24
3.3 Global positioning system location of obtained samples.	26
3.4 Range and level of variables that affects glyphosate degradation using Plackett-Burman design for free cells.	34
3.5 Plackett-Burman experimental design for glyphosate degradation by <i>Burkholderia vietnamiensis</i> AQ5-12.	35
3.6 Plackett-Burman experimental design for glyphosate degradation by <i>Burkholderia</i> sp. AQ5-13.	35
3.7 Central composite experimental design for glyphosate degradation by <i>Burkholderia vietnamiensis</i> AQ5-12.	36
3.8 Central composite experimental design for glyphosate degradation by <i>Burkholderia</i> sp. AQ5-13.	37
3.9 Range and level of variables that affects glyphosate degradation using Plackett-Burman design for immobilised cells	39
3.10 Plackett-Burman experimental design for glyphosate degradation by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	40
3.11 Plackett-Burman experimental design for glyphosate degradation by immobilised <i>Burkholderia</i> sp. strain AQ5-13.	40
3.12 Central composite experimental design for glyphosate degradation by immobilised cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12.	41
3.13 Central composite experimental design for glyphosate degradation by immobilised <i>Burkholderia</i> sp. strain AQ5-13.	42
4.1 Effect of different glyphosate formula, ROUNDUP® concentration on bacterial growth of isolates.	45
4.2 Foremost 19 sequences producing important alignment with <i>Burkholderia vietnamiensis</i> isolate AQ5-12 from NCBI blast.	53

4.3	Foremost 18 sequences producing important alignment with <i>Burkholderia</i> sp. isolate AQ5-13 from NCBI blast.	56
4.4	Morphological and biochemical properties of the different isolates. (+), strains positive; (-), strains negative.	60
4.5	Experimental design and results of Plackett-Burman on glyphosate degradation by <i>Burkholderia vietnamiensis</i> strain A5-12.	79
4.6	Analysis of parameters for glyphosate degradation for Plackett-Burman for <i>Burkholderia vietnamiensis</i> strain A5-12.	80
4.7	Experimental design and results of Plackett-Burman on glyphosate degradation by <i>Burkholderia</i> sp. strain AQ5-13.	81
4.8	Analysis of parameters for glyphosate degradation for Plackett-Burman for <i>Burkholderia</i> sp. strain AQ5-13.	82
4.9	Experimental design and results of central composite design on glyphosate degradation by <i>Burkholderia vietnamiensis</i> strain A5-12.	83
4.10	Analysis of parameters for glyphosate degradation for central composite design for <i>Burkholderia vietnamiensis</i> strain A5-12.	84
4.11	Experimental design and results of central composite design on glyphosate degradation by <i>Burkholderia</i> sp. strain AQ5-13.	88
4.12	Analysis of parameters for glyphosate degradation for central composite design for <i>Burkholderia</i> sp. strain AQ5-13.	90
4.13	Suggested parameter for each variables for optimum degradation of 100 ppm glyphosate by free cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12 based on CCD.	95
4.14	Comparison of optimum conditions and results obtained between OFAT and RSM for <i>Burkholderia vietnamiensis</i> strain AQ5-12.	96
4.15	Suggested parameter for each variables for optimum degradation of 50 ppm glyphosate by free cells of <i>Burkholderia</i> sp. strain AQ5-13 based on CCD.	97
4.16	Comparison of optimum conditions and results obtained between OFAT and RSM for <i>Burkholderia</i> sp. strain AQ5-13.	98
4.17	Experimental design and results of Plackett-Burmann on glyphosate degradation by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	105
4.18	Analysis of parameters for glyphosate degradation for Plackett-Burman for immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	106
4.19	Experimental design and results of Plackett-Burman on glyphosate degradation by immobilised <i>Burkholderia</i> sp. strain AQ5-13.	107
4.20	Analysis of parameters for glyphosate degradation for central composite design for immobilised <i>Burkholderia</i> sp. strain AQ5-13.	107

4.21	Experimental design and results of CCD on glyphosate degradation by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	108
4.22	Analysis of parameters for glyphosate degradation for central composite design for immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	110
4.23	Experimental design and results of central composite design on glyphosate degradation by immobilised <i>Burkholderia</i> sp. strain AQ5-13.	114
4.24	Analysis of parameters for glyphosate degradation for central composite design for immobilised <i>Burkholderia</i> sp. strain AQ5-13	115
4.25	Suggested parameter for each variables for optimum degradation of 50 ppm glyphosate by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12 based on CCD.	122
4.26	Comparison of optimum conditions and results obtained between OFAT and RSM by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	122
4.27	Suggested parameter for each variables for optimum degradation of 50 ppm glyphosate by immobilised <i>Burkholderia</i> sp. strain AQ5-13 based on CCD.	123
4.28	Comparison of optimum conditions and results obtained between OFAT and RSM by immobilised <i>Burkholderia</i> sp. strain AQ5-13.	124
4.29	Summary of isolates tolerance towards Roundup concentration and identification of high glyphosate tolerating isolates AQ5-12 and AQ5-13.	126
4.30	Summary of optimisation using OFAT and RSM by free cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12 and <i>Burkholderia</i> sp. strain AQ5-13.	127
4.31	Summary of optimisation using OFAT and RSM by immobilised cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12 and <i>Burkholderia</i> sp. strain AQ5-13.	127

LIST OF FIGURES

Figure		Page
2.1	Structure of glyphosate molecule.	4
2.2	The shikimate pathway that leads to the formation of aromatic amino acids and the inhibition of EPSPS by glyphosate.	5
2.3	Glyphosate metabolic pathways.	13
2.4	3D contour for RSM design	17
3.1	Experimental design of glyphosate by locally-isolated bacteria.	22
4.1	Effect of 3 mL/L glyphosate formula, ROUNDUP® on bacterial growth of isolates.	46
4.2	Effect of 6 mL/L glyphosate formula, ROUNDUP® on bacterial growth of isolates.	47
4.3	Effect of 9 mL/L glyphosate formula, ROUNDUP® on bacterial growth of isolates.	48
4.4	Effect of 12 mL/L glyphosate formula, ROUNDUP® on bacterial growth of isolates.	49
4.5	Gram stain smear of (A) Isolate AQ5-12 and (B) Isolate AQ05-13	50
4.6	Agarose gel electrophoresis of 16S rRNA genomic of isolate AQ5-12 and AQ5-13.	51
4.7	Agarose gel electrophoresis of PCR product of 16S rRNA gene of isolate AQ5-12 and AQ5-13.	52
4.8	This cladogram was carried out based on neighbour joining technique illustrating the phylogenetic relationship among isolate AQ5-12 and other interconnected analogous bacteria based on 16S rRNA gene sequence examination.	54
4.9	The accession number of 16S rRNA sequence of isolate AQ5-12 deposited in GeneBank.	55
4.10	This cladogram was carried out based on neighbour joining technique illustrating the phylogenetic relationship among isolate AQ5-13 and other interconnected analogous bacteria based on 16S rRNA gene sequence examination.	57
4.11	The accession number of 16S rRNA sequence of isolate AQ5-13 deposited in GeneBank.	58
4.12	Effects of different carbon source on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12.	62

4.13	Effects of different sucrose concentration on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12.	63
4.14	Effects of different carbon source on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia</i> sp. strain AQ5-13.	64
4.15	Effects of different sucrose concentration on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia</i> sp. strain AQ5-13.	65
4.16	Effects of nitrogen source on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia vietnamiensis</i> isolate AQ5-12.	67
4.17	Effects of different ammonia sulphate concentration on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12.	68
4.18	Effects of nitrogen source on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia</i> sp. strain AQ5-13.	69
4.19	Effects of different ammonia sulphate concentration on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia</i> sp. strain AQ5-13.	70
4.20	Effects of pH on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12.	71
4.21	Effects of pH on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia</i> sp. strain AQ5-13.	72
4.22	Effects of temperature on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12.	74
4.23	Effects of temperature on biodegradation of 50 ppm glyphosate by resting cells of <i>Burkholderia</i> sp. strain AQ5-13.	75
4.24	Effects of glyphosate concentration on biodegradation of glyphosate by resting cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12.	76
4.25	Effects of glyphosate concentration on biodegradation of glyphosate by resting cells of <i>Burkholderia</i> sp. strain AQ5-13.	77
4.26	Response surface 3D plot showing the effect of pH and carbon concentration on glyphosate degradation.	85
4.27	Similarity plot between predicted and actual value for free cells of <i>Burkholderia vietnamiensis</i> strain AQ5-12.	86
4.28	Effects of higher fructose concentration on biodegradation of 100 ppm glyphosate by resting cells of <i>Burkholderia vietnamiensis</i> strain AQ5-13.	87
4.29	Response surface 3D plot showing the effect of pH and nitrogen concentration on glyphosate degradation.	91

4.30	Response surface 3D plot showing the effect of carbon and glyphosate concentration on glyphosate degradation.	92
4.31	Response surface 3D plot showing the effect of nitrogen and glyphosate concentration on glyphosate degradation.	93
4.32	Response surface 3D plot showing the effect of pH and glyphosate concentration on glyphosate degradation.	94
4.33	Similarity plot between predicted and actual value for free cells of <i>Burkholderia</i> sp. strain AQ5-13.	94
4.34	Validation of experiment (RSM) conducted to ascertain the optimum solution obtained through the conduct of the experiment by <i>Burkholderia vietnamiensis</i> strain AQ5-12.	96
4.35	Validation of experiment (RSM) conducted to ascertain the optimum solution obtained through the conduct of the experiment.	97
4.36	Effect of gellan gum concentration in the bioremoval of 100 ppm glyphosate by immobilised cells <i>Burkholderia vietnamiensis</i> strain AQ5-12.	99
4.37	Effect of gellan gum concentration in the bioremoval of 50 ppm glyphosate by immobilised cells <i>Burkholderia</i> sp. strain AQ5-13.	100
4.38	Effect of beads sizes in the bioremoval of 100 ppm glyphosate by immobilised cells <i>Burkholderia vietnamiensis</i> strain AQ5-12.	101
4.39	Effect of beads sizes in the bioremoval of 50 ppm glyphosate by immobilised <i>Burkholderia</i> sp. strain AQ5-13.	102
4.40	Effect of beads numbers in the bioremoval of 100 ppm glyphosate by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	103
4.41	Effect of beads numbers in the bioremoval of 50 ppm glyphosate by immobilised <i>Burkholderia</i> sp. strain AQ5-13.	104
4.42	Response surface 3D plot showing the effect of beads size and number of beads on glyphosate degradation by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	111
4.43	Response surface 3D plot showing the effect of number of beads and gellan gum concentration on glyphosate degradation by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	112
4.44	Response surface 3D plot showing the effect beads size and gellan gum concentration on glyphosate degradation by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	113
4.45	Similarity plot between predicted and actual value.	113
4.46	Response surface 3D plot showing the effect of number of beads and gellan gum concentration on glyphosate degradation	116
4.47	Response surface 3D plot showing the effect of beads size and gellan gum concentration on glyphosate degradation.	117

4.48	Response surface 3D plot showing the effect of beads size and number of beads on glyphosate degradation by immobilised <i>Burkholderia</i> sp. strain AQ5-13.	118
4.49	Similarity plot between predicted and actual value for immobilised <i>Burkholderia</i> sp. strain AQ5-13.	119
4.50	Validation of experiment (RSM) conducted to ascertain the optimum solution obtained through the conduct of the experiment by immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12.	122
4.51	Validation of experiment (RSM) conducted to ascertain the optimum solution obtained through the conduct of the experiment by immobilised <i>Burkholderia</i> sp. strain AQ5-13.	124
4.52	Repeated usage of immobilised <i>Burkholderia vietnamiensis</i> strain AQ5-12 in 100 ppm glyphosate; <i>Burkholderia</i> sp. strain AQ5-13 in 50 ppm glyphosate.	125

LIST OF ABBREVIATIONS

%	Percent
<	Less than
>	Greater than
Abs	Absorbance
AMPA	Aminomethylphosphonic acid
ANOVA	The analysis of variance
CCD	Central composite design
cm	Centimetre
Co	Cobalt
Cr	Chromium
Cu	Copper
Da	Dalton
dH ₂ O	Distilled water
EDTA	Ethylene diamine tetra acetic acid
et al	and friends
EC	Half maximal effective concentration
EPSPS	5-enolpyruvyl-shikimate-3-phosphate-synthase
Fe	Iron
FFD	Fractional factorial design
G	Gram
h	Hour
IPA	Isopropylamine
Hg	Mercury
kb	Kilo base
kDa	Kilo dalton
kg	Kilogram
L	Liter
LC	Lethal concentration
M	Molar
min	Minute
mL	Mililitre
MSM	Minimal salt medium
MW	Molecular weight
NA	Nutrient agar
°C	Degree celcius
OD	Optical density
OFAT	One-factor-at-a-time
POEA	Polyethoxylated tallow amine
RPM	Rotation per minute
RSM	Response surface methodology
SDS	Sodium dodecyl sulphate
SEM	Scanning electron microscope
μL	Microlitre
μm	Micrometre
μM	Micromolar

CHAPTER 1

INTRODUCTION

Herbicides are used as a means of controlling a variety of weed species. Newly formulated herbicides are fast acting and possess unique actions against different environmental conditions. Herbicides can help to increase crop yield by reducing crop loss due to uncontrolled weed growth. Meanwhile, glyphosate is a well-known herbicide in the agricultural field. The rise of glyphosate is mainly due to its fast action in controlling weed number. It is the only herbicide that targets 5-enolpyruvyl-shikimate-3-phosphate synthase (EPSPS) (Yu et al., 2015). Hence, there are no rival herbicide analogues or classes for this herbicide (Duke and Powles, 2008). By inhibiting this crucial enzyme, plants will wither and die. Glyphosate herbicides were first introduced as Roundup brand by Monsanto Company. Since then, more than 172 glyphosate-based products can be found in Malaysia alone.

Herbicides are mainly used to kill and destroy unwanted terrestrial weeds. However, during raining seasons, applied Roundup formulations on agricultural land are washed away and ended up in an aquatic medium. Aquatic plants and algae are consequently the most sensitive group of aquatic nontarget organisms. Moreover, they play significant role in aquatic ecosystems. Destruction of these aquatic plants reduces the stability of sediments in the lakes and running water, thus destroying the habitat where juvenile fishes and crustaceans take shelter (Gurnell et al., 2012). Besides, exposure to glyphosate on amphibians resulted in abnormalities. A high percentage of morphology alterations was observed in sharp-snouted tree frog (*Ssinax nassicus*) incubated with 3 - 7 mg/L of glyphosate, which is the exact amount used in the sub-agricultural field (Sviridov et al., 2014).

Increased awareness on the profound effects of environmental problems such as traces of glyphosate in drinking waters and accepts of herbicide usage has encouraged necessary investigations to avoid, reduce or eliminate these problems. Applications of microbial inoculants for in-situ treatment of contamination of soil include the enhancement of associative N₂ fixation and symbiotic, prevention of soil-borne plant pathogens, biological control of frost injury on the tubular plant and biodegradation of xenobiotic compounds (Elsas and Heijnen, 1990). This microorganism has been introduced into contaminated sites in the form of liquid suspensions or adsorbed into different physical carriers (Sparrow and Ham, 1983). Nevertheless, the efficiency of glyphosate biodegradations depends on the success in isolating glyphosate-degrading strains.

A number of glyphosate degrading microorganisms with varying inherent capability to utilise glyphosate as carbon or phosphorus source have been isolated and reported from different environmental sources. However, high tolerance glyphosate degrading bacteria were frequently isolated from agricultural contaminated sites. To date, *Bacillus subtilis*

Bs-15 is one of the best glyphosate degraders with the maximum concentration tolerated by this bacterium reaches as high as 40,000 mg/L isolated from glyphosate exposed soil (Fan et al., 2012). The growth of these bacteria and their ability to effectively degrade glyphosate are significantly influenced by various physical and nutritional factors. Optimisation of these factors is therefore crucial to maximise their degradation ability. One factor at a time (OFAT) optimisation technique is often deployed to achieve optimum glyphosate degradation. However, this method results in some limitations; some of them are the lack of capability to address the interactive effects between factors and time-consuming as the parameters cannot be simultaneously run (Okoroma et al., 2012). Statistical optimisation approach using response surface methodology (RSM) is used to correct the limitations of the OFAT, which has successfully optimised many biological processes (Karamba et al., 2016; Nawawi et al., 2016; Fakhfakh-Zouari et al., 2010)

Most of these bacteria are isolated using glyphosate alone without introducing any herbicide formulation in the medium. Although it has been reported that *Bacillus subtilis* Bs-15 able to degrade up to 40,000 ppm of glyphosate, the application in situ is still questionable (Fan et al., 2012). This is because these bacteria need to withstand the toxic compounds that usually come together with glyphosate formulation. Chemicals such as polyethoxylated tallow amine, acrylamide and urea, which can be found in major glyphosate formulation, can significantly reduce bacterial growth or kill them (Mesnage et al., 2014). Therefore, it is important to isolate bacteria using Roundup than using glyphosate alone as glyphosate alone isolated bacteria may not perform as expected in the application of bioremediation. Furthermore, only two glyphosate degrading bacteria were isolated from Malaysia (Nourouzi et al., 2011). Moreover, no reports or studies were found on glyphosate degradation using gellan gum for immobilisation. The advantage encapsulation of bacterial cells in gellan gum using immobilisation technique provides the ability to be used in real life application with fewer risks. Beads, made of gellan gum has the ability to be repeatedly used and the composition of the beads are made of non-toxic materials which is safer towards the environment (Karamba et al., 2016). Moreover, this beads prevents the target bacterial cells leaked out which could be pathogenic to native strains. These gaps need to be filled as data from this study could provide significant understanding on the effects of immobilisation using native strains in Malaysia.

Based on the statements above, this study aims at isolating a native bacterium from Malaysia with a potential of withstanding the toxicity of Roundup and ability to degrade glyphosate. Thus, the objectives of this study are:

1. To isolate, characterise, identify and select glyphosate-degrading bacterium from the soil.
2. To determine optimise nutritional and physical conditions of the bacterium for a maximum glyphosate degradation using OFAT approach and RSM by free cells.
3. To study the glyphosate degradation rate using immobilised bacterium, optimise the immobilisation conditions and investigate the degradation ability of the immobilised cells via OFAT approach and RSM.

REFERENCES

- Adelowo, F. E., Olu-Arotiowa, O. A., & Amuda, O. S. (2014). Biodegradation of glyphosate by fungi species. *Advances in Bioscience and Bioengineering*, 2(1), 104–118
- Adjei, M. D., & Ohta, Y. (2000). Factors affecting the biodegradation of cyanide by *Burkholderia cepacia* strain C-3. *Journal of Bioscience and Bioengineering*, 89(3), 274–277.
- Ahmad, S.A. (2011). Biodegradation of phenol by locally isolated *Acinetobacter* sp. strain AQ5NOL 1 and purification of phenol hydroxylase (Doctoral thesis, Universiti Putra Malaysia).
- Ahmad, S. A., Shamaan, N. A., Arif, N. M., Koon, G. B., Shukor, M. Y. A., & Syed, M. A. (2012). Enhanced phenol degradation by immobilized *Acinetobacter* sp. strain AQ5NOL 1. *World Journal of Microbiology and Biotechnology*, 28(1), 347–352.
- Ahmed, S. A. (2008). Invertase production by *Bacillus macerans* immobilized on calcium alginate beads. *Journal of Applied Science Research*, 4(12), 1777–1781.
- Ayoola, S. O. (2008). Toxicity of glyphosate herbicide on Nile tilapia (*Oreochromis niloticus*) juvenile. *African Journal of Agricultural Research*, 3(12), 825–834.
- Baek, I., Seo, B., Lee, I., Lee, K., Park, S.-C., Yi, H., & Chun, J. (2015). *Burkholderia megalo-chromosomata* sp. nov., isolated from grassland soil. *International Journal of Systematic and Evolutionary Microbiology*, 65(3), 959–964.
- Baerson, S. R., Rodriguez, D. J., Tran, M., Feng, Y., Biest, N. A., & Dill, G. M. (2002). Glyphosate-resistant goosegrass. Identification of a mutation in the target enzyme 5-enolpyruvylshikimate-3-phosphate synthase. *Plant Physiology*, 129(3), 1265–1275.
- Balthazor, T. M., & Hallas, L. E. (1986). Glyphosate-degrading microorganisms from industrial activated sludge. *Applied and Environmental Microbiology*, 51(2), 432–434.
- Bazot, S., & Lebeau, T. (2008). Simultaneous mineralization of glyphosate and diuron by a consortium of three bacteria as free-and/or immobilized-cells formulations. *Applied Microbiology and Biotechnology*, 77(6), 1351–1358.
- Ben Khedher, S., Jaoua, S., & Zouari, N. (2013). Application of statistical experimental design for optimisation of bioinsecticides production by sporeless *Bacillus thuringiensis* strain on cheap medium. *Brazilian Journal of Microbiology*, 44(3), 927–933.
- Benachour, N., & Seralini, G.-E. (2009). Glyphosate formulations induce apoptosis and necrosis in human umbilical, embryonic, and placental cells. *Chemical Research in Toxicology*, 22(1), 97–105.
- Benjaminson, M. A., de Guzman, B. C., & Weil, A. J. (1964). Voges-Proskauer test: Expeditious techniques for routine use. *Journal of Bacteriology*, 87(1), 234–235.

- Benslama, O., & Boulahrouf, A. (2013). Isolation and characterization of glyphosate-degrading bacteria from different soils of Algeria. *African Journal of Microbiology Research*, 7(49), 5587–5595.
- Beshay, U. (2003). Production of alkaline protease by *Teredinobacter turnirae* cells immobilized in Ca-alginate beads. *African Journal of Biotechnology*, 2(3), 60–65.
- Beveridge, T. J. (2001). Use of the Gram stain in microbiology. *Biotechnic and Histochemistry*, 76(3), 111–118.
- Binupriya, A. R., Sathishkumar, M., Ku, C. S., & Yun, S.-I. (2010). Sequestration of Reactive Blue 4 by free and immobilized *Bacillus subtilis* cells and its extracellular polysaccharides. *Colloids and Surfaces B: Biointerfaces*, 76(1), 179–185.
- Boontawan, A., & Boontawan, P. (2011). Isolation and characterization of Jatropha oil degradation by *Enterococcus faecalis* and *Burkholderia cenocepacia* W-1 under anaerobic condition. *African Journal of Biotechnology*, 10(63), 13841–13851.
- Borggaard, O. K., & Gimsing, A. L. (2008). Fate of glyphosate in soil and the possibility of leaching to ground and surface waters: A review. *Pest Management Science*, 64(4), 441–456.
- Boyd, C. E., & Massaut, L. (1999). Risks associated with the use of chemicals in pond aquaculture. *Aquacultural Engineering*, 20(2), 113–132.
- Box, G., Hunter, W. and Hunter, J. (1978). *Statistics for experimenters*. John Wiley and Sons, New York. pp.144.
- Brena, B., González-Pombo, P., & Batista-Viera, F. (2013). Immobilization of enzymes: A literature survey. In *Immobilization of Enzymes and Cells* (pp. 15–31). Humana Press, Totowa, NJ.
- Carlisle, S. M., & Trevors, J. T. (1988). Glyphosate in the environment. *Water, Air, and Soil Pollution*, 39(4), 409–420.
- Cavalcante, V. A., & Dobereiner, J. (1988). A new acid-tolerant nitrogen-fixing bacterium associated with sugarcane. *Plant and Soil*, 108(1), 23–31.
- Cedergreen, N., & Streibig, J. C. (2005). The toxicity of herbicides to non-target aquatic plants and algae: Assessment of predictive factors and hazard. *Pest Management Science*, 61(12), 1152–1160.
- Campbell, C. D., Grayston, S. J., & Hirst, D. J. (1997). Use of rhizosphere carbon sources in sole carbon source tests to discriminate soil microbial communities. *Journal of Microbiological Methods*, 30(1), 33–41.
- Chatterjee, S., Sau, G. B., & Mukherjee, S. K. (2011). Bioremediation of Cr(VI) from chromium-contaminated wastewater by free and immobilized cells of *Cellulosimicrobium cellulans* KUCr3. *Bioremediation Journal*, 15(3), 173–180.

- Chee, J. Y., Tan, Y., Samian, M. R., & Sudesh, K. (2010). Isolation and characterization of a *Burkholderia* sp. USM (JCM15050) capable of producing polyhydroxyalkanoate (pha) from triglycerides, fatty acids and glycerols. *Journal of Polymers and the Environment*, 18(4), 584–592.
- Chibata, I., Tosa, T., & Sato, T. (1986). Biocatalysis: Immobilized cells and enzymes. *Journal of Molecular Catalysis*, 37(1), 1–24.
- Choi, S. H., Moon, S. H., & Gu, M. B. (2002). Biodegradation of chlorophenols using the cell-free culture broth of *Phanerochaete chrysosporium* immobilized in polyurethane foam. *Journal of Chemical Technology and Biotechnology*, 77(9), 999–1004.
- Christensen, W. B. (1946). Urea decomposition as a means of differentiating proteus and paracolon cultures from each other and from *Salmonella* and *Shigella* Types 1. *Journal of Bacteriology*, 52(4), 461–466.
- Coupland, D., & Caseley, J. C. (1979). Presence of 14c activity in root exudates and guttation fluid from agropyron repens treated with 14c-labelled glyphosate. *New Phytologist*, 83(1), 17–22.
- Coutinho, D. F., Sant, S. V., Shin, H., Oliveira, J. T., Gomes, M. E., Neves, N. M., & Reis, R. L. (2010). Modified gellan gum hydrogels with tunable physical and mechanical properties. *Biomaterials*, 31(29), 7494–7502.
- Crestani, M., Menezes, C., Gluszczak, L., Santos Miron, D. dos, Spanevello, R., Silveira, A., Goncalves, F. F., Zanella, R., & Loro, V. L. (2007). Effect of clomazone herbicide on biochemical and histological aspects of silver catfish (*Rhamdia quelen*) and recovery pattern. *Chemosphere*, 67(11), 2305–2311.
- Debbabi, F., Gargoubi, S., Hadj Ayed, M. A., & Abdessalem, S. B. (2017). Development and characterization of antibacterial braided polyamide suture coated with chitosan-citric acid biopolymer. *Journal of Biomaterials Applications*, 32(3), 384–398.
- Della-Cioppa, G., Bauer, S. C., Klein, B. K., Shah, D. M., Fraley, R. T., & Kishore, G. M. (1986). Translocation of the precursor of 5-enolpyruvylshikimate-3-phosphate synthase into chloroplasts of higher plants in vitro. *Proceedings of the National Academy of Sciences of the United States of America*, 83(18), 6873–6877.
- Dick, R. E., & Quinn, J. P. (1995). Glyphosate-degrading isolates from environmental samples: occurrence and pathways of degradation. *Applied Microbiology and Biotechnology*, 43(3), 545–550.
- Dotson, S. B., Smith, C. E., Ling, C. S., Barry, G. F., & Kishore, G. M. (1996). Identification, characterization, and cloning of a phosphonate monoester hydrolase from *Burkholderia caryophilli* PG2982. *Journal of Biological Chemistry*, 271(42), 25754–25761.
- Duke, S. O., & Powles, S. B. (2008). Glyphosate: A once-in-a-century herbicide. *Pest Management Science*, 64(4), 319–325.
- Dyhrman, S. T., Ammerman, J. W., & Van Mooy, B. A. (2007). Microbes and the marine phosphorus cycle. *Oceanography*, 20(2), 110–116.

- van den Berg, L., van Vliet, T., van der Linden, E., van Boekel, M. A. J. S., & van de Velde, F. (2007). Serum release: The hidden quality in fracturing composites. *Food Hydrocolloids*, 21(3), 420–432.
- Van Elsas, J.D., & Heijnen, C. E. (1990). Methods for the introduction of bacteria into soil: A review. *Biology and Fertility of Soils*, 10(2), 127–133.
- El-Naas, M. H., Mourad, A. H. I., & Surkatti, R. (2013). Evaluation of the characteristics of polyvinyl alcohol (PVA) as matrices for the immobilization of *Pseudomonas putida*. *International Biodeterioration and Biodegradation*, 85(1), 413-420.
- Eman, A., Abdel-Megeed, A., Suliman, A. M. A., Sadik, M. W., & Sholkamy, E. N. (2013). Biodegradation of glyphosate by fungal strains isolated from herbicides polluted-soils in Riyadh area. *British Journal of Environmental Sciences*, 1(1), 7–29.
- Ermakova, I. T., Kiseleva, N. I., Shushkova, T., Zharikov, M., Zharikov, G. A., & Leontievsky, A. A. (2010). Bioremediation of glyphosate-contaminated soils. *Applied Microbiology and Biotechnology*, 88(2), 585–594.
- Fakhfakh-Zouari, N., Haddar, A., Hmidet, N., Frikha, F., & Nasri, M. (2010). Application of statistical experimental design for optimization of keratinases production by *Bacillus pumilus* A1 grown on chicken feather and some biochemical properties. *Process Biochemistry*, 45(5), 617–626.
- Fan, J., Yang, G., Zhao, H., Shi, G., Geng, Y., Hou, T., & Tao, K. (2012). Isolation, identification and characterization of a glyphosate-degrading bacterium, *Bacillus cereus* CB4, from soil. *The Journal of general and applied microbiology*, 58(4), 263-271.
- Farh, M. E.-A., Kim, Y.-J., An, H. V., Sukweenadhi, J., Singh, P., Huq, M. A., & Yang, D.-C. (2015). *Burkholderia ginsengiterrae* sp. nov. and *Burkholderia panaciterrae* sp. nov., antagonistic bacteria against root rot pathogen *Cylindrocarpon destructans*, isolated from ginseng soil. *Archives of Microbiology*, 197(3), 439–447.
- Fiske, C. H., & Subbarow, Y. (1925). The colorimetric determination of phosphorus. *Journal of Biological Chemistry*, 66(2), 375–400.
- Fitzgibbon, J., & Braymer, H. D. (1988). Phosphate starvation induces uptake of glyphosate by *Pseudomonas* sp. strain PG2982. *Applied and Environmental Microbiology*, 54(7), 1886–1888.
- Folmar, L. C., Sanders, H. O., & Julin, A. M. (1979). Toxicity of the herbicide glyphosate and several of its formulations to fish and aquatic invertebrates. *Archives of Environmental Contamination and Toxicology*, 8(3), 269–278.
- Franz, J. E., Mao, M. K., & Sikorski, J. A. (1997). *Glyphosate: A unique global herbicide*. (Vol. 189). American Chemical Society, Washington D.C.
- Gardner, S. C., Grue, C. E., Grassley, J. M., Lenz, L. A., Lindenauer, J. M., & Seeley, M. E. (1997). Single species algal (*Ankistrodesmus*) toxicity tests with Rodeo and Garlon 3A. *Bulletin of Environmental Contamination and Toxicology*, 59(1), 492–499.

- Gibson, F. (1964). Chorismic acid: purification and some chemical and physical studies. *Biochemical Journal*, 90(2), 256–261.
- Gimsing, A. L., Borggaard, O. K., & Sestoft, P. (2004). Modeling the kinetics of the competitive adsorption and desorption of glyphosate and phosphate on goethite and gibbsite and in soils. *Environmental Science & Technology*, 38(6), 1718–1722.
- Glass, R. L. (1981). Colorimetric determination of glyphosate in water after oxidation to orthophosphate. *Analytical Chemistry*, 53(6), 921–923.
- Goldsborough, L. G., & Brown, D. J. (1988). Effect of glyphosate (Roundup® formulation) on periphytic algal photosynthesis. *Bulletin of Environmental Contamination and Toxicology*, 41(2), 253–260.
- Guanxi, L. I., Xiaoqin, W. U., & Jianren, Y. E. (2013). The characterization of glyphosate degradation by *Burkholderia multivorans* WS-FJ9. *Acta Ecologica Sinica*, 33(21), 6885–6894.
- Gurnell, A. M., Bertoldi, W., & Corenblit, D. (2012). Changing river channels: The roles of hydrological processes, plants and pioneer fluvial landforms in humid temperate, mixed load, gravel bed rivers. *Earth-Science Reviews*, 111(1), 129–141.
- Hadi, F., Mousavi, A., Noghabi, K. A., Tabar, H. G., & Salmanian, A. H. (2013). New bacterial strain of the genus *Ochrobactrum* with glyphosate-degrading activity. *Journal of Environmental Science and Health*, 48(3), 208–213.
- Hallas, L. E., Adams, W. J., & Heitkamp, M. A. (1992). Glyphosate degradation by immobilized bacteria: Field studies with industrial wastewater effluent. *Applied and Environmental Microbiology*, 58(4), 1215–1219.
- Halter, S. (2007). A brief history of Roundup. Proceedings of the First International Symposium on Glyphosate. Agronomical Sciences College of the University of the State of Sao Paulo, Sao Paulo, Brazil, October 15 – 19.
- Hance, R. J. (1976). Adsorption of glyphosate by soils. *Pesticide Science*, 7(4), 363–366.
- Herrmann, K. M. (1995). The shikimate pathway as an entry to aromatic secondary metabolism. *Plant Physiology*, 107(1), 7–12.
- Hove-Jensen, B., Zechel, D. L., & Jochimsen, B. (2014). Utilization of glyphosate as phosphate source: Biochemistry and genetics of bacterial carbon-phosphorus lyase. *Microbiology and Molecular Biology Reviews*, 78(1), 176–197.
- Hui, L., Sheng, G., Johnston, C. T., & Boyd, S. A. (2003). Sorption and desorption of pesticides by clay minerals and humic acid-clay complexes. *Soil Science Society of America Journal*, 67(1), 122–131.
- Ibrahim, S., Muhammad, A., Tanko, A. S., Abubakar, A., Ibrahim, H., Shukor, M. Y., & Ahmad, S. A. (2015). Studies of action of heavy metals on caffeine degradation by immobilised *Leifsonia* sp. strain SIU. *Bayero Journal of Pure and Applied Sciences*, 8(2), 138–144.

- Imandi, S. B., Bandaru, V. V. R., Somalanka, S. R., Bandaru, S. R., & Garapati, H. R. (2008). Application of statistical experimental designs for the optimization of medium constituents for the production of citric acid from pineapple waste. *Bioresource Technology*, 99(10), 4445–4450.
- Kah, M., Beulke, S., & Brown, C. D. (2007). Factors influencing degradation of pesticides in soil. *Journal of Agricultural and Food Chemistry*, 55(11), 4487–4492.
- Kamjumphol, W., Chareonsudjai, P., Taweechaisupapong, S., & Chareonsudjai, S. (2015). Morphological alteration and survival of *Burkholderia pseudomallei* in soil microcosms. *The American Journal of Tropical Medicine and Hygiene*, 93(5), 1058–1065.
- Karamba, K. I., Ahmad, S. A., Zulkharnain, A., Syed, M. A., Khalil, K. A., Shamaan, N. A., Dahalan, F. A., & Shukor, M. Y. (2016). Optimisation of biodegradation conditions for cyanide removal by *Serratia marcescens* strain AQ07 using one-factor-at-a-time technique and response surface methodology. *Rendiconti Lincei*, 27(3), 533–545.
- Kare, L. H., & Hakon O., F. (1986). Photodegradation of the herbicides glyphosate in water. *Bulletin of environmental contamination and toxicology*, 36(1), 723–729.
- Karigar, C. S., & Rao, S. S. (2011). Role of microbial enzymes in the bioremediation of pollutants: A Review. *Enzyme Research*, 2011. <http://dx.doi.org/10.4061/2011/805187>
- Kertesz, M., Elgorriaga, A., & Amrhein, N. (1991). Evidence for two distinct phosphonate-degrading enzymes (C-P lyases) in *Arthrobacter* sp. GLP-1. *Biodegradation*, 2(1), 53–59.
- Khuri, A. I., & Mukhopadhyay, S. (2010). Response surface methodology. *Wiley Interdisciplinary Reviews: Computational Statistics*, 2(2), 128-149.
- Kiertiburanakul, S., Sungkanuparph, S., Kositchiwat, S., & Vorachit, M. (2002). *Burkholderia pseudomallei*: Abscess in an unusual site. *Journal of Postgraduate Medicine*, 48(2), 124–126.
- Kim, T. J., Lee, E. Y., Kim, Y. J., Cho, K.-S., & Ryu, H. W. (2003). Degradation of polyaromatic hydrocarbons by *Burkholderia cepacia* 2A-12. *World Journal of Microbiology and Biotechnology*, 19(4), 411–417.
- King, A. (2001). Recommendations for susceptibility tests on fastidious organisms and those requiring special handling. *Journal of Antimicrobial Chemotherapy*, 48(1), 77–80.
- Kishore, G. M., & Jacob, G. S. (1987). Degradation of glyphosate by *Pseudomonas* sp. PG2982 via a sarcosine intermediate. *The Journal of Biological Chemistry*, 262(25), 12164–12168.
- Kowalska, M., Bodzek, M., & Bohdziewicz, J. (1998). Biodegradation of phenols and cyanides using membranes with immobilized microorganisms. *Process Biochemistry*, 33(2), 189–197.

- Kryuchkova, Y. V., Burygin, G. L., Gogoleva, N. E., Gogolev, Y. V., Chernyshova, M. P., Makarov, O. E., Fedorov, E.E., & Turkovskaya, O. V. (2014). Isolation and characterization of a glyphosate-degrading rhizosphere strain, *Enterobacter cloacae* K7. *Microbiological Research*, 169(1), 99–105.
- Krzyśko-Lupicka, T., & Orlik, A. (1997). The use of glyphosate as the sole source of phosphorus or carbon for the selection of soil-borne fungal strains capable to degrade this herbicide. *Chemosphere*, 34(12), 2601–2605.
- Kuklinsky-Sobral, J., Araújo, W. L., Mendes, R., Pizzirani-Kleiner, A. A., & Azevedo, J. L. (2005). Isolation and characterization of endophytic bacteria from soybean (*Glycine max*) grown in soil treated with glyphosate herbicide. *Plant and Soil*, 273(1–2), 91–99.
- Kumar, P., & Satyanarayana, T. (2007). Optimization of culture variables for improving glucoamylase production by alginate-entrapped *Thermomucor indicae-seudaticae* using statistical methods. *Bioresource Technology*, 98(6), 1252–1259.
- Lee, L. J., & Ngim, J. (2000). A first report of glyphosate-resistant goosegrass (*Eleusine indica* (L) Gaertn) in Malaysia. *Pest Management Science*, 56(4), 336–339.
- Leong, K. H., Tan, L. L. B., & Mustafa, A. M. (2007). Contamination levels of selected organochlorine and organophosphate pesticides in the Selangor River, Malaysia between 2002 and 2003. *Chemosphere*, 66(6), 1153–1159.
- Lerbs, W., Stock, M., & Parthier, B. (1990). Physiological aspects of glyphosate degradation in *Alcaligenes* spec. strain GL. *Archives of Microbiology*, 153(2), 146–150.
- Magrí, A., Vanotti, M. B., & Szögi, A. A. (2012). Anammox sludge immobilized in polyvinyl alcohol (PVA) cryogel carriers. *Bioresource Technology*, 114(2), 231–240.
- Malik, J., Barry, G., & Kishore, G. (1989). The herbicide glyphosate. *BioFactors* (Oxford, England), 2(1), 17–25.
- Mann, R. M., & Bidwell, J. R. (1999). The toxicity of glyphosate and several glyphosate formulations to four species of southwestern Australian frogs. *Archives of Environmental Contamination and Toxicology*, 36(2), 193–199.
- Margesin, R., Moertelmaier, C., & Mair, J. (2013). Low-temperature biodegradation of petroleum hydrocarbons (n-alkanes, phenol, anthracene, pyrene) by four actinobacterial strains. *International Biodeterioration and Biodegradation*, 84(1), 185–191.
- Maule, A., & Wright, S. J. L. (1984). Herbicide effects on the population growth of some green algae and cyanobacteria. *Journal of Applied Bacteriology*, 57(2), 369–379.
- McAuliffe, K. S., Hallas, L. E., & Kulpa, C. F. (1990). Glyphosate degradation by *Agrobacterium radiobacter* isolated from activated sludge. *Journal of Industrial Microbiology*, 6(3), 219–221.

- Mesnage, R., Defarge, N., Vend, S. de, Mois, J., & Ralini, G. E. (2014). Major pesticides are more toxic to human cells than their declared active principles. *BioMed Research International*, 2014, e179691. <http://dx.doi.org/10.1155/2014/179691>.
- Miller, J. M., & Wright, J. W. (1982). Spot indole test: evaluation of four reagents. *Journal of Clinical Microbiology*, 15(4), 589–592.
- Mohanty, S. S., & Jena, H. M. (2017). Biodegradation of phenol by free and immobilized cells of a novel *Pseudomonas* sp. NBM11. *Brazilian Journal of Chemical Engineering*, 34(1), 75–84.
- Molin, W. T., & Hirase, K. (2005). Effects of surfactants and simulated rainfall on the efficacy of the Engame formulation of glyphosate in johnsongrass, prickly sida and yellow nutsedge. *Weed Biology and Management*, 5(3), 123–127.
- Moneke, A. N., Okpala, G. N., & Anyanwu, C. U. (2010). Biodegradation of glyphosate herbicide in vitro using bacterial isolates from four rice fields. *African Journal of Biotechnology*, 9(26), 4067–4074.
- Morgan, M. J., & Kiceniuk, J. W. (1992). Response of rainbow trout to a two month exposure to Vision®, a glyphosate herbicide. *Bulletin of Environmental Contamination and Toxicology*, 48(5), 772–780.
- Morris, E. R., Nishinari, K., & Rinaudo, M. (2012). Gelation of gellan—a review. *Food Hydrocolloids*, 28(2), 373–411.
- Muralidhar, R. V., Chirumamila, R. R., Marchant, R., & Nigam, P. (2001). A response surface approach for the comparison of lipase production by *Candida cylindracea* using two different carbon sources. *Biochemical Engineering Journal*, 9(1), 17–23.
- Nandula, V. K. (2010). *Glyphosate Resistance in Crops and Weeds: History, Development, and Management*. John Wiley & Sons.
- Nawawi, N. M., Ahmad, S. A., Shukor, M. Y., Syed, M. A., & Khalil, K. A. (2016). Statistical optimisation for improvement of phenol degradation by *Rhodococcus* sp. NAM 81. *Journal of Environmental Biology*, 37(3), 443–451.
- Nešković, N. K., Poleksić, V., Elezović, I., Karan, V., & Budimir, M. (1996). Biochemical and histopathological effects of glyphosate on carp, *Cyprinus carpio* L. *Bulletin of Environmental Contamination and Toxicology*, 56(2), 295–302.
- Neve, P., Diggle, A. J., Smith, F. P., & Powles, S. B. (2003). Simulating evolution of glyphosate resistance in *Lolium rigidum* II: past, present and future glyphosate use in Australian cropping. *Weed Research*, 43(6), 418–427.
- Ngigi, A. N., Getenga, Z. M., Boga, H. I., & Ndalut, P. K. (2012). Biodegradation of s-triazine herbicide atrazine by *Enterobacter cloacae* and *Burkholderia cepacia* sp. from long-term treated sugarcane-cultivated soils in Kenya. *Journal of Environmental Science and Health, Part B*, 47(8), 769–778.
- Nomura, N. S., & Hilton, H. W. (1977). The adsorption and degradation of glyphosate in five Hawaiian sugarcane soils. *Weed Research*, 17(2), 113–121.

- Norris, M. H., Kang, Y., Lu, D., Wilcox, B. A., & Hoang, T. T. (2009). Glyphosate resistance as a novel select-agent-compliant, non-antibiotic-selectable marker in chromosomal mutagenesis of the essential genes *asd* and *dapB* of *Burkholderia pseudomallei*. *Applied and Environmental Microbiology*, 75(19), 6062–6075.
- Nourouzi, M. M., Chuah, T. G., Choong, T. S. Y., & Lim, C. J. (2011). Glyphosate utilization as the source of carbon: Isolation and identification of new bacteria. *Journal of Chemistry*, 8(4), 1582–1587.
- Nussinovitch, A. (2010). *Bead Formation, Strengthening, and Modification*. In *Polymer Macro- and Micro-Gel Beads: Fundamentals and Applications* (pp. 27–52). New York : Springer.
- Obojska, A., Ternan, N. G., Lejczak, B., Kafarski, P., & McMullan, G. (2002). Organophosphonate utilization by the thermophile *Geobacillus caldxylosilyticus* T20. *Applied and Environmental Microbiology*, 68(4), 2081–2084.
- Oh, S., Yun A. and Park, D. 2011. Effects of physicochemically hydrolyzed human hairs on the soil microbial community and growth of the hot pepper plant. *Biotechnology and Bioprocess Engineering*, 16(4): 746-754
- Okoroma, E. A., Garelick, H., Abiola, O. O., & Purchase, D. (2012). Identification and characterisation of a *Bacillus licheniformis* strain with profound keratinase activity for degradation of melanised feather. *International Biodeterioration and Biodegradation*, 74, 54–60.
- Oliveira, J. T., Martins, L., Picciochi, R., Malafaya, P. B., Sousa, R. A., Neves, N. M., Mano, J.F., & Reis, R. L. (2010). Gellan gum: A new biomaterial for cartilage tissue engineering applications. *Journal of Biomedical Materials Research Part A*, 93A(3), 852–863.
- Ory, I. de, Cabrera, G., Ramirez, M., & Blandino, A. (2006). Immobilization of cells on polyurethane foam. In *Immobilization of Enzymes and Cells* (pp. 357–365). Humana Press, New York City, New York, United States.
- Paerl, H. (2008). Nutrient and other environmental controls of harmful cyanobacterial blooms along the freshwater–marine continuum. *Advances in Experimental Medicine and Biology*, 37(1), 217–237.
- Paungfoo-Lonhienne, C., Lonhienne, T. G. A., Yeoh, Y. K., Webb, R. I., Lakshmanan, P., Chan, C. X., Lim, P.E., Ragan, M.A., Schmidt, S., & Hugenholtz, P. (2014). A new species of *Burkholderia* isolated from sugarcane roots promotes plant growth. *Microbial Biotechnology*, 7(2), 142–154.
- Perelo, L. W. (2010). Review: In situ and bioremediation of organic pollutants in aquatic sediments. *Journal of Hazardous Materials*, 177(1), 81–89.
- Pipke, R., & Amrhein, N. (1988). Degradation of the phosphonate herbicide glyphosate by *Arthrobacter atrocyaneus* ATCC 13752. *Applied and Environmental Microbiology*, 54(5), 1293–1296.
- Pollegioni, L., Schonbrunn, E., & Siehl, D. (2011). Molecular basis of glyphosate resistance: Different approaches through protein engineering. *The FEBS Journal*, 278(16), 2753–2766.

- Powles, S. B., & Preston, C. (2006). Evolved glyphosate resistance in plants: Biochemical and genetic basis of resistance. *Weed Technology*, 20(2), 282–289.
- Quan, C. S., Zheng, W., Liu, Q., Ohta, Y., & Fan, S. D. (2006). Isolation and characterization of a novel *Burkholderia cepacia* with strong antifungal activity against *Rhizoctonia solani*. *Applied Microbiology and Biotechnology*, 72(6), 1276–1284.
- Quan, L. M., Khanh, D. P., Hira, D., Fujii, T., & Furukawa, K. (2011). Reject water treatment by improvement of whole cell anammox entrapment using polyvinyl alcohol/alginate gel. *Biodegradation*, 22(6), 1155–1167.
- Ren, L., Wang, P., Wang, C., Chen, J., Hou, J., & Qian, J. (2017). Algal growth and utilization of phosphorus studied by combined mono-culture and co-culture experiments. *Environmental Pollution*, 220(1), 274–285.
- Roberts, T. R., Hutson, D. H., Lee, P. W., Nicholls, P. H., Plimmer, J. R., Roberts, M. C., & Croucher, L. (Eds.). (1998). *Metabolic Pathways of Agrochemicals: Part 1: Herbicides and Plant Growth Regulators* (1 edition). Cambridge: Royal Society of Chemistry.
- Roshon, R. D., McCann, J. H., Thompson, D. G., & Stephenson, G. R. (1999). Effects of seven forestry management herbicides on *Myriophyllum sibiricum*, as compared with other nontarget aquatic organisms. *Canadian Journal of Forest Research*, 29(7), 1158–1169.
- Rueppel, M. L., Brightwell, B. B., Schaefer, J., & Marvel, J. T. (1977). Metabolism and degradation of glyphosate in soil and water. *Journal of Agricultural and Food Chemistry*, 25(3), 517–528.
- Saeki, D., Nagao, S., Sawada, I., Ohmukai, Y., Maruyama, T., & Matsuyama, H. (2013). Development of antibacterial polyamide reverse osmosis membrane modified with a covalently immobilized enzyme. *Journal of Membrane Science*, 428, 403–409.
- Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4(4), 406–425.
- Sala, G., Stieger, M., & van de Velde, F. (2010). Serum release boosts sweetness intensity in gels. *Food Hydrocolloids*, 24(5), 494–501.
- Sánchez, D., Graça, M. A. S., & Canhoto, J. (2007). Testing the use of the water milfoil (*Myriophyllum spicatum* L.) in laboratory toxicity assays. *Bulletin of Environmental Contamination and Toxicology*, 78(6), 421–426.
- Santos, P. E.-D. L., Bustillos-Cristales, R., & Caballero-Mellado, J. (2001). *Burkholderia*, a genus rich in plant-associated nitrogen fixers with wide environmental and geographic distribution. *Applied and Environmental Microbiology*, 67(6), 2790–2798.
- Sass, A. M., Schmerk, C., Agnoli, K., Norville, P. J., Eberl, L., Valvano, M. A., & Mahenthalingam, E. (2013). The unexpected discovery of a novel low-oxygen-activated locus for the anoxic persistence of *Burkholderia cenocepacia*. *The ISME Journal*, 7(8), 1568–1581.

- Schaffer, J. D., & Sebetich, M. J. (2004). Effects of aquatic herbicides on primary productivity of phytoplankton in the laboratory. *Bulletin of Environmental Contamination and Toxicology*, 72(5), 1032–1037.
- Schnurer, Y., Persson, P., Nilsson, M., Nordgren, A., & Giesler, R. (2006). Effects of surface sorption on microbial degradation of glyphosate. *Environmental Science and Technology*, 40(13), 4145–4150.
- Selvapandiyan, A., & Bhatnagar, R. K. (1994). Cloning of genes encoding for C-P lyase from *Pseudomonas* isolates PG2982 and GLC11: Identification of a cryptic allele on the chromosome of *P. aeruginosa*. *Current Microbiology*, 29(5), 255–261.
- Sharma, S. D., & Singh, M. (2001). Environmental factors affecting absorption and bio-efficacy of glyphosate in Florida beggarweed (*Desmodium tortuosum*). *Crop Protection*, 20(6), 511–516.
- Singh, B. K., & Walker, A. (2006). Microbial degradation of organophosphorus compounds. *FEMS Microbiology Reviews*, 30(3), 428–471.
- Sobrero, M. C., Rimoldi, F., & Ronco, A. E. (2007). Effects of the glyphosate active ingredient and a formulation on *Lemma gibba* L. at different exposure levels and assessment end-points. *Bulletin of Environmental Contamination and Toxicology*, 79(5), 537–543.
- Soletto, D., Binaghi, L., Lodi, A., Carvalho, J. C. M., & Converti, A. (2005). Batch and fed-batch cultivations of *Spirulina platensis* using ammonium sulphate and urea as nitrogen sources. *Aquaculture*, 243(1), 217–224.
- Sparrow, S.D. & Ham, G. E. (1983). Survival of *Rhizobium phaseoli* in six carrier materials. *Agronomy Journal*, 75(2), 181–184
- Sprankle, P., Meggitt, W. F., & Penner, D. (1975). Adsorption, mobility, and microbial degradation of glyphosate in the soil. *Weed Science*, 23(3), 229–234.
- Steel, K. J. (1961). The Oxidase reaction as a taxonomic tool. *Microbiology*, 25(2), 297–306.
- Sviridov, A. V., Shushkova, T. V., Ermakova, I. T., Ivanova, E. V., & Leontievsky, A. A. (2014). Glyphosate: Safety risks, biodegradation, and bioremediation. In G. Cao & R. Orrù (Eds.), *Current Environmental Issues and Challenges* (pp. 183–195). New York: Springer
- Talbot, H. W., Johnson, L. M., & Munnecke, D. M. (1984). Glyphosate utilization by *Pseudomonas* sp. and *Alcaligenes* sp. isolated from environmental sources. *Current Microbiology*, 10(5), 255–259.
- Tanyildizi, M. S., Özer, D., & Elibol, M. (2005). Optimization of α -amylase production by *Bacillus* sp. using response surface methodology. *Process Biochemistry*, 40(7), 2291–2296.
- Tavichakorntrakool, R., Prasongwattana, V., Sungkeeree, S., Saisud, P., Sribenjalux, P., Pimratana, C., Bovornpadungkitti, S., Sriboonlue, P., & Thongboonkerd, V. (2012). Extensive characterizations of bacteria isolated from catheterized urine and stone matrices in patients with nephrolithiasis. *Nephrology Dialysis Transplantation*, 27(11), 4125–4130.

- Taylor, W. I., & Achanzar, D. (1972). Catalase test as an aid to the identification of *Enterobacteriaceae*. *Applied Microbiology*, 24(1), 58–61.
- Thapa, B., Kc, A. K., & Ghimire, A. (2012). A Review on bioremediation of petroleum hydrocarbon contaminants in soil. *Kathmandu University Journal of Science, Engineering and Technology*, 8(1), 164–170.
- Thys, R. C. S. and Brandelli, A. 2006. Purification and properties of a keratinolytic metalloprotease from *Microbacterium* sp. *Journal of Applied Microbiology*, 101(6): 1259-1268
- Tsui, M. T. K., & Chu, L. M. (2003). Aquatic toxicity of glyphosate-based formulations: comparison between different organisms and the effects of environmental factors. *Chemosphere*, 52(7), 1189–1197.
- Tzin, V., & Galili, G. (2010). The biosynthetic pathways for shikimate and aromatic amino acids in *Arabidopsis thaliana*. *The Arabidopsis Book* e0132. doi.org/10.1199/tab.0132 .
- van Pelt, C., Verduin, C. M., Goessens, W. H. F., Vos, M. C., Tümmler, B., Segonds, C., Reubsæet, F., Verbrugh, H., & van Belkum, A. (1999). Identification of *Burkholderia* spp. in the clinical microbiology laboratory: Comparison of conventional and molecular methods. *Journal of Clinical Microbiology*, 37(7), 2158–2164.
- Vendrell, E., Ferraz, D., Sabater, C., & Carrasco, J. M. (2009). Effect of glyphosate on growth of four freshwater species of phytoplankton : A microplate bioassay. *Bulletin of Environmental Contamination and Toxicology*, 82, 538–542.
- Vera, M. S., Lagomarsino, L., Sylvester, M., Pérez, G. L., Rodríguez, P., Mugni, H., Sinistro, R., Ferraro, Marcela., Bonetto, C., Zagarese, H., & Pizarro, H. (2010). New evidences of Roundup (glyphosate formulation) impact on the periphyton community and the water quality of freshwater ecosystems. *Ecotoxicology*, 19(4), 710–721.
- Wackett, L. P., Shames, S. L., Venditti, C. P., & Walsh, C. T. (1987). Bacterial carbon-phosphorus lyase: Products, rates, and regulation of phosphonic and phosphinic acid metabolism. *Journal of Bacteriology*, 169(2), 710–717.
- Waclawovsky, A. J., Sato, P. M., Lembke, C. G., Moore, P. H., & Souza, G. M. (2010). Sugarcane for bioenergy production: an assessment of yield and regulation of sucrose content. *Plant Biotechnology Journal*, 8(3), 263–276.
- Wang, X., Gai, Z., Yu, B., Feng, J., Xu, C., Yuan, Y., ... Xu, P. (2007). Degradation of carbazole by microbial cells immobilized in magnetic gellan gum gel beads. *Applied and Environmental Microbiology*, 73(20), 6421–6428.
- Ware, G. W. (2000). *Reviews of Environmental Contamination and Toxicology: Continuation of Residue Reviews*. New York: Springer Science & Business Media.
- Wetzel, R. G. (2001). *Limnology: Lake and River Ecosystems*. Houston: Gulf Professional Publishing.

- White, A. K., & Metcalf, W. W. (2004). Two C—P lyase operons in *Pseudomonas stutzeri* and their roles in the oxidation of phosphonates, phosphite, and hypophosphite. *Journal of Bacteriology*, *186*(14), 4730–4739.
- Witek-Krowiak, A., Chojnacka, K., Podstawczyk, D., Dawiec, A., & Pokomeda, K. (2014). Application of response surface methodology and artificial neural network methods in modelling and optimization of biosorption process. *Bioresource technology*, *160*, 150–160.
- Wösten, J. H. M., Ismail, A. B., & van Wijk, A. L. M. (1997). Peat subsidence and its practical implications: A case study in Malaysia. *Geoderma*, *78*(1), 25–36.
- Yadzir, M., & Hani, Z. (2007). Characterization, identification and application of *Acinetobacter baumannii* SERDANG 1 for phenol biodegradation (Doctoral dissertation, Universiti Teknologi PETRONAS).
- Yu, J., Zhang, X., & Tan, T. (2007). An novel immobilization method of *Saccharomyces cerevisiae* to sorghum bagasse for ethanol production. *Journal of Biotechnology*, *129*(3), 415–420.
- Yu, Q., Jalaludin, A., Han, H., Chen, M., Sammons, R. D., & Powles, S. B. (2015). Evolution of a double amino acid substitution in the 5-enolpyruvylshikimate-3-phosphate synthase in *Eleusine indica* conferring high-level glyphosate resistance. *Plant Physiology*, *167*(4), 1440–1447.
- Zeng, L., Huang, J., Zhang, Y., Qiu, G., Tong, J., Chen, D., Zhou, J., & Luo, X. (2008). An effective method of DNA extraction for bioleaching bacteria from acid mine drainage. *Applied Microbiology and Biotechnology*, *79*(5), 881.
- Zhao, H., Tao, K., Zhu, J., Liu, S., Gao, H., & Zhou, X. (2015). Bioremediation potential of glyphosate-degrading *Pseudomonas* spp. strains isolated from contaminated soil. *The Journal of General and Applied Microbiology*, *61*(5), 165–170.
- ZoBell, C. E. (1932). Factors influencing the reduction of nitrates and nitrites by bacteria in semisolid media. *Journal of Bacteriology*, *24*(4), 273–281.