



UNIVERSITI PUTRA MALAYSIA

***ROLE OF yqiG PSEUDOGENE IN Escherichia coli TOWARDS
BIOHYDROGEN PRODUCTION***

MUHAMMAD AZMAN BIN ZAKARIA

FBSB 2018 36



**ROLE OF *yqiG* PSEUDOGENE IN *Escherichia coli* TOWARDS
BIOHYDROGEN PRODUCTION**

By

MUHAMMAD AZMAN BIN ZAKARIA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

June 2017

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

**ROLE OF *yqiG* PSEUDOGENE IN *Escherichia coli*
TOWARDS BIOHYDROGEN PRODUCTION**

By

MUHAMMAD AZMAN BIN ZAKARIA

June 2017

Supervisor : Mohd Zulkhairi Mohd Yusoff, PhD
Faculty : Biotechnology and Biomolecular Sciences

Biohydrogen gas has a great potential as alternative energy and it is considered as environmental friendly energy source. Extensive studies have been carried out on the production of biohydrogen from *Escherichia coli* (*E. coli*). *E. coli* was used due to its accessibility of full genome and well characterized proteins. Interestingly, there are some of pseudogenes present in *E. coli* genome. The pseudogenes were considered as non-functional genes as result of genetic variability. In this study, we found that *yqiG* is one of the pseudogenes in *E. coli* that is essential for biohydrogen production. The deletion of *yqiG* has reduced the biohydrogen productivity in comparison to the *E. coli* wild type as control. The objective of this study is to investigate the role of *yqiG* pseudogene of *E. coli* during hydrogen production. The *yqiG* single gene deletion strain and ASKA strains were used throughout the study. The *yqiG* mutant strain has significantly interrupted biohydrogen productivity and suppressed glucose consumption up to 50% compared to wild type, respectively. In addition, overexpressed *yqiG* protein in the wild type via pCA24N-YqiG had increased the hydrogen productivity to 56 $\mu\text{mol H}_2/\text{mg}$ protein from glucose compared to control which only 38 $\mu\text{mol H}_2/\text{mg}$ protein. Similar pattern was observed from formate fermentation whereas wild type producing *yqiG* protein dominated the biohydrogen productivity about 141.8 $\mu\text{mol H}_2/\text{mg}$ protein. Higher expression of formate hydrogen lyase components (*fdhF*, *hycE* and *fhla*) were observed in mutant *yqiG* strain. In contrast, the *yqiG* mutation causes down regulation of *pflB* and *pykF* of glycolysis as well as formic acid accumulation in the fermentation broth compared to the wild type strain. The *pflB* and *pykF* encode pyruvate kinase and pyruvate formate lyase, respectively. We conclude that the *yqiG* protein is important in the conversion of phosphoenolpyruvate to pyruvate and utilization of pyruvate to formate for hydrogen production. Thus, *yqiG* pseudogene makes an important contribution during hydrogen metabolism in *E. coli*.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PERANAN PSEUDOGENE *yqiG* DALAM *Escherichia coli*
KE ARAH PENGELUARAN BIOHIDROGEN**

Oleh

MUHAMMAD AZMAN BIN ZAKARIA

Jun 2017

Penyelia : Mohd Zulkhairi Mohd Yusoff, PhD
Fakulti : Bioteknologi dan Sains Biomolekul

Gas biohidrogen berpotensi besar sebagai tenaga alternatif dan ia dianggap sebagai sumber tenaga mesra alam. Kajian yang menyeluruh sudah dijalankan terhadap pengeluaran biohidrogen daripada *Escherichia coli* (*E.coli*). *E. coli* diguna disebabkan oleh kebolehcapaian terhadap genomnya yang menyeluruh dan protein-proteinnnya yang sudah dicirikan. Menariknya, terdapat beberapa pseudogene wujud di dalam genom *E. coli*. Pseudogene dianggap sebagai gen yang tidak berfungsi akibat daripada kebolehubahan genetik. Di dalam kajian ini, kami mendapati bahawa *yqiG* merupakan salah satu pseudogene dalam *E. coli* yang mana penting untuk pengeluaran biohidrogen. Penghapusan *yqiG* telah mengurangkan produktiviti biohidrogen dalam perbandingan kepada *E. coli* jenis liar sebagai kawalan. Objektif kajian ini adalah untuk menyiasat peranan pseudogene *yqiG* daripada *E.coli* semasa penghasilan biohidrogen. Strain tanpa satu gen *yqiG* dan strain ASKA telah digunakan sepanjang kajian ini dijalankan. Strain mutan *yqiG* dengan ketara mengganggu produktiviti biohidrogen dan mengurangkan penggunaan glukosa sehingga 50% berbanding strain jenis liar, masing-masing. Di samping itu, ekspresi berlebihan YqiG dalam jenis liar melalui pCA24N-YqiG telah meningkatkan produktiviti hidrogen kepada 56 $\mu\text{mol H}_2/\text{mg}$ protein daripada glukosa berbanding strain kawalan, 38 $\mu\text{mol H}_2/\text{mg}$ protein. Bentuk yang sama diperhatikan daripada penapaian yang mana strain jenis liar yang menghasilkan protein *yqiG* mendominasi produktiviti biohidrogen sebanyak 141.8 $\mu\text{mol H}_2/\text{mg}$ protein. Keputusan yang diperolehi adalah selaras dengan keputusan analisis transkripsi dengan ekspresi yang lebih tinggi kepada komponen format hidrogen lyase (*fdhF*, *hycE* dan *fhlA*). Sebaliknya, mutasi *yqiG* menyebabkan regulasi negatif kepada glycolysis *pflB* dan *pykF* serta pengumpulan asid formik dalam medium penapaian berbanding strain jenis liar. *pflB* dan *pykF* mengekod kepada pyruvate kinase dan pyruvate format lyase masing-masing. Kami menyimpulkan bahawa protein *yqiG* ini adalah penting dalam penukaran phosphoenolpyruvate ke pyruvate dan penggunaan pyruvate ke formate untuk pengeluaran biohidrogen. Oleh itu, pseudogen *yqiG* menyumbang kepada metabolisme hidrogen dalam *E.coli*.

ACKNOWLEDGEMENTS

I would like to express my deepest appreciation to my main supervisor, Dr. Mohd Zulkhairi Mohd Yusoff and my supervisory committee members, Prof. Dr. Mohd Ali Hassan, Dr. Mohd Rafein Zakaria, and Assoc. Prof. Dr. Toshinari Maeda for their guidance, encouragement, motivation, and advice throughout my studies. With their knowledge, experience and guidance, I was able to solve any problems whenever I had encountered it during my research. Special thanks to Ministry of Higher Education, Malaysia, School of Graduate Studies, Universiti Putra Malaysia (UPM), Kyushu Institute of Technology (KYUTECH) and Japan Student Services Organization (JASSO) for the scholarship during my study.

I would like to thank Environmental Biotechnology (EB) group members and Maeda laboratory members such as Dr. Nazlina Haiza Mohd Yassin, Dr. Yee Lian Ngit, and others for the moral support and help whenever I needed. I really appreciated the guidelines and ideas from you.

Last but not least, I am gratefully thanked my beloved parent, Zakaria Bin Abdullah and Latipah Binti Ahmad and siblings for their understanding and moral support. Their endless love, support and encouragement have made me a better person.

I certify that a Thesis Examination Committee has met on 9 June 2017 to conduct the final examination of Muhammad Azman bin Zakaria on his thesis entitled "Role of *yqiG* Pseudogene in *Escherichia coli* Towards Biohydrogen Production" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Phang Lai Yee, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Mohd Shukuri bin Mohamad Ali, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Khanom Simarani, PhD

Senior Lecturer
University of Malaya
Malaysia
(External Examiner)



RUSLI HAJI ABDULLAH, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 27 September 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Mohd Zulkhairi Mohd Yusoff, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Ali Hassan, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Mohd Rafein Zakaria, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Toshinari Maeda, PhD

Associate Professor
Graduate School of Life Science and Systems Engineering
Kyushu Institute of Technology
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Muhammad Azman Bin Zakaria (GS36199)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature : _____
Name of Chairman of
Supervisory Committee : Dr. Mohd Zulkhairi Mohd Yusoff

Signature : _____
Name of Member of
Supervisory Committee : Professor Dr. Mohd Ali Hassan

Signature : _____
Name of Member of
Supervisory Committee : Dr. Mohd Rafein Zakaria @ Mamat

Signature : _____
Name of Member of
Supervisory Committee : Associate Professor Dr. Toshinari Maeda

TABLE OF CONTENTS

ABSTRACT	Page
ABSTRAK	i
ACKNOWLEDGEMENTS	ii
APPROVAL	iii
DECLARATION	v
LIST OF TABLES	vi
LIST OF FIGURES	x
LIST OF ABBREVIATIONS	xi
	xii

CHAPTER

1 INTRODUCTION	1
1.1 Research overview	1
1.2 Problem statements	2
1.3 Objectives	2
2 LITERATURE REVIEW	3
2.1 Hydrogen production	3
2.1.1 Hydrocarbon reforming	4
2.1.2 Water electrolysis	4
2.1.3 Microbial fuel cell (MFC)	5
2.1.4 Biological hydrogen production	6
2.2 Biohydrogen through dark fermentation	6
2.2.1 Hydrolysis	7
2.2.2 Acidogenesis	8
2.2.3 Acetogenesis	9
2.2.4 Methanogenesis	9
2.3 Biohydrogen from <i>Escherichia coli</i>	9
2.3.1 Hydrogenases in <i>E. coli</i>	11
2.3.1.1 Hydrogenase 1 and hydrogenase 2	13
2.3.1.2 Hydrogenase 3	14
2.3.1.3 Hydrogenase 4	14
2.4 Pseudogene	14
2.4.1 Types of pseudogenes	15
2.4.1.1 Processed pseudogene	16
2.4.1.2 Duplicated pseudogene	17
2.4.1.3 Unitary pseudogene	17
2.4.2 <i>yqiG</i> pseudogene in <i>E. coli</i>	17
2.5 Glycolysis pathway	19
2.6 Metabolic engineering of <i>E. coli</i> for biohydrogen production	23
2.7 Factor affecting biohydrogen production	25
2.7.1 Organic acid and pH	25

2.7.2	Macro and micronutrients	27
2.7.3	Temperature	28
2.7.4	Effect of substrate	29
3	MATERIALS AND METHODS	30
3.1	Bacteria and strain maintenance	30
3.2	Mutant verification	30
3.3	Biohydrogen assay analysis	31
3.4	Chemical analysis	32
3.5	Complementation analysis test	33
3.6	Effect of IPTG	33
3.7	Pellet and RNA extraction	33
3.7.1	Integrity of RNA sample	34
3.7.2	Quantitative real time polymerase chain reaction qRT-PCR	34
3.8	Protein hydrophobicity profile	36
3.9	Statistical method	36
4	RESULTS	37
4.1	Strains verification	37
4.2	Biohydrogen production	38
4.2.1	Biohydrogen assay and specific growth rate	38
4.2.2	Glucose consumption and organic acid	41
4.3	Effect of IPTG on biohydrogen production	43
4.3.1	Glucose consumption and organic acid	43
4.4	RNA extraction and transcriptional analysis	46
4.4.1	Integrity of RNA sample	46
4.4.2	Transcriptional analysis	47
4.5	Protein hydrophobicity profile	49
5	DISCUSSIONS, CONCLUSIONS AND RECOMMENDATIONS	52
5.1	Discussions	52
5.1	Conclusions	56
5.2	Recommendations for future research	56
	REFERENCES	57
	APPENDICES	74
	BIODATA OF STUDENT	87
	LIST OF PUBLICATIONS	88

LIST OF TABLES

Table		Page
2.1	The pure bacterial cultures for biohydrogen fermentation	10
2.2	Biohydrogen production from variants of <i>E. coli</i> strain	12
2.3	General information of <i>yqiG</i> pseudogene	18
2.4	Strategies in metabolic engineering for biohydrogen production	25
2.5	The effect of initial pH for biohydrogen production	26
2.6	The effect of temperature on biohydrogen production	28
3.1	<i>E. coli</i> strains and plasmids used in this study	30
3.2	Primer used for strain verification	31
3.3	Primers designed for gene expression analysis	35
3.4	Chemical reaction for qRT-PCR	36
3.5	Thermal cycling condition for qRT-PCR	36
4.1	Specific growth rate of strains used in 100 mM glucose	40
4.2	Distribution of organic acids in fermentation broth after 24h fermentation with complex glucose	43
4.3	Effect of IPTG on organic acid accumulation	45
4.4	Fold change in promoter activity during transcriptional analysis of targeted genes	48
4.5	Protein features of <i>yqiG</i>	50

LIST OF FIGURES

Figure	Page
2.1	5
2.2	8
2.3	15
2.4	16
2.5	18
2.6	19
2.7	20
2.8	22
2.9	24
3.1	32
4.1	37
4.2	38
4.3	39
4.4	40
4.5	41
4.6	42
4.7	44
4.8	44
4.9	46
4.10	49
4.11	50

LIST OF ABBREVIATIONS

APS	Ammonium Persulfate
Bp	Base pair
CBB	Coomassie Brilliant Blue
cDNA	Complementary DNA
CM	Chloramphenicol
CU	Chaperone Usher
DNA	Deoxyribonucleic Acid
dNTP	Deoxynucleotide
<i>E. coli</i>	<i>Escherichia coli</i>
EDTA	Ethylenediaminetetraacetic acid
FDH _N	Formate dehydrogenase-N
FDH _O	Formate dehydrogenase-O
FHL	Formate Hydrogen lyase System
FUP	Fimbrial Usher Protein
GC	Gas Chromatography
H ₂	Hydrogen
HPLC	High Performance Liquid Chromatography
HCl	Hydrochloric acid
Hyd	Hydrogenase
IS	Insertion Sequence
IPTG	Isopropyl β-D-1-Thiogalactopyranoside
Km ^R	Kanamycin Resistance Gene
LB	Luria Bertani
mRNA	Messenger RNA
MFC	Microbial Fuse Cell
MgSO ₄	Magnesium Sulphate
NaOCl	Sodium hypochlorite
NaCl	Sodium Chloride
Ni-NTA Agarose	Nickel-charged Affinity resin
OMP	Outer Membrane Protein
psig	Pounds per Square Inch gauge
PCR	Polymerase Chain Reaction
PMSF	Phenylmethylsulfonyl Fluoride
PEP	Phosphoenolpyruvate
PTS	Phosphotransferase system
qPCR	Quantitative Polymerase Chain Reaction
RNA	Ribonucleic Acid
rpm	Rotation per minute
SDS-PAGE	Sodium dodecyl sulfate Polyacrylamide gel Electrophoresis
TAE	Tris-acetate
TEMED	Tetramethylethylenediamine

CHAPTER 1

INTRODUCTION

1.1 Research overview

Biohydrogen is an attractive energy source due its combustion only produces water vapour as by-products. It is known as environmental friendly fuel since it can be produced from renewable sources such as biomass and organic materials. Compared to other energy fuel such as fossil fuel and methane gas, biohydrogen contain higher energy content (Mohd Yasin *et al.*, 2011). Hydrogen energy research became worldwide phenomena for clean energy production (Patterson *et al.*, 2013; Pawar and van Niel, 2013). Photosynthetic hydrogen production is one of the biological methods for biohydrogen production. It will convert solar energy into hydrogen in the presence of sunlight by photosynthetic bacteria. However, due to low efficiency of light consumption and reactor design difficulties make it not suitable for large scale application (Das and Veziroglu, 2008; Hawkes *et al.*, 2007). Fermentative hydrogen production is the best option of sustainable technology whereas it can consume organic feedstocks, waste biomass or lignocellulosic materials. The fermentation process can be carried out at ambient temperature and lower atmospheric pressure (Hallenbeck and Ghosh, 2009; Hallenbeck, 2014). A lot of studies were reported on biohydrogen production from *Escherichia coli* (*E. coli*). The modification of *E. coli* through metabolic engineering approaches have delivered a great impact on the enhancement of biohydrogen production (Maeda *et al.*, 2007a; Seol *et al.*, 2014; Tran *et al.*, 2014). Metabolic engineering has increased the biohydrogen production by eliminating some of pathways that are not contributed to hydrogen production (Ghosh *et al.*, 2013).

In recent research, pseudogenes in *E. coli* have been found to influence hydrogen production (Mohd Yusoff *et al.*, 2013). Pseudogenes are usually considered as non-functional. They are derived from an active gene which has evolutionarily lost its function as a result of a defect in transcription and translation. Hence, pseudogenes are considered junk genes due to the presence of inappropriate stop codons, repetitive element frame shifts and lack of transcription elements (Balakirev and Ayala, 2003; Harrison *et al.*, 2001).

There are more than 4500 genes in the *E. coli* genome and 178 of the genes are known as pseudogenes (Zhou and Rudd, 2013). According to Rouchka and Cha (2009), pseudogenes can be classified as processed and unprocessed pseudogenes. Processed pseudogenes are from retrotransposable events from mRNA and lack of intronic regions, flanking repeat regions and poly-A tracts at the carboxyl 3' end (Pavlicek *et al.*, 2006). This type of pseudogene can be identified by its 3' end by the presence of polyadenine (Esnault *et al.*, 2000). Unprocessed pseudogenes emerge during the gene duplication process and the genetic sequences can go through several changes such as

mutations, frame shifts, insertions and deletions (Rouchka and Cha, 2009). Any obvious changes during the transcriptional or translational process can affect the functionality of the gene. Failure of duplication of the genes resulted in a new sequence that could be a pseudogenes (Vanin, 1985). In addition, functional genes have high possibilities to turn into pseudogenes without duplication process when their selective constraints are rapidly removed or diminished due to the changes of genetic background or environment (Sakai *et al.*, 2008).

The uncorrupted *yqiG* encodes for an 821 amino acid (aa) protein (Keseler *et al.*, 2010). However, in *E. coli*, *yqiG* is characterized as a pseudogene due to internal insertion-sequences (IS) insC9 and insD9 (Zhou and Rudd, 2013; Hayashi *et al.*, 2006). IS elements being a segment of bacterial DNA that can be shifted from one part of the chromosome to other parts of the same chromosome or another chromosome. The presence of IS elements in the middle of a gene interrupts the coding sequence, thus inactivating the protein product. Furthermore, the presence of an IS element near the promoter of an operon can block the expression of other genes in the same operon (Griffiths *et al.*, 2000). Hence, the presence of IS elements suggests that *yqiG* is a non-functional gene in *E. coli* and results its classification as a pseudogene (Korea *et al.*, 2010).

1.2 Problem statements

The function of the *yqiG* pseudogene in *E. coli* is unknown; however, it is necessary to consider pseudogenes in hydrogen metabolism since deleting some of pseudogenes in *E. coli* has delivered a positive response on the hydrogen production (Mohd Yusoff *et al.*, 2013). Most of the researchers conceive pseudogenes as junk or non-functional genes and almost no report has mentioned a relation of pseudogenes towards hydrogen metabolism. This consideration is supported since the metabolic pathway of biohydrogen production is well understood which neglect the presence of pseudogenes in *E. coli*. Other studies carried out by Watanabe *et al.* (2008) and Tam *et al.* (2008) found that pseudogenes involved in biological interaction especially in gene regulation. In this study, thorough investigations of *yqiG* characteristics were carried out to elucidate *yqiG* role during hydrogen metabolism. Hence, fermentation and molecular biotechnology approaches were considered to elaborate the role of *yqiG* protein towards biohydrogen production in *E. coli*.

1.3 Objectives

The objectives of this research are:

1. To confirm the presence of *yqiG* pseudogene in wild type *E. coli* BW25113 genome through molecular biotechnology technique.
2. To study the role of *yqiG* pseudogene in *E. coli* for biohydrogen production.

REFERENCES

- Abdel-Hamid, A. M., Attwood, M. M., and Guest, J. R. (2001). Pyruvate oxidase contributes to the aerobic growth efficiency of *Escherichia coli*. *Microbiology*, *147*, 1483–1498.
- Abo-Hashesh, M., Wang, R., and Hallenbeck, P. C. (2011). Metabolic engineering in dark fermentative hydrogen production; theory and practice. *Bioresource Technology*, *102*(18), 8414–8422.
- Abreu, A. A., Danko, A. S., Costa, J. C., Ferreira, E. C., and Alves, M. M. (2009). Inoculum type response to different pHs on biohydrogen production from l-arabinose, a component of hemicellulosic biopolymers. *International Journal of Hydrogen Energy*, *34*(4), 1744–1751.
- Adams, M. W., and Hall, D. O. (1979). Purification of the membrane-bound hydrogenase of *Escherichia coli*. *The Biochemical Journal*, *183*(1), 11–22.
- Adav, S. S., Lee, D.-J., Wang, A., and Ren, N. (2009). Functional consortium for hydrogen production from cellobiose: concentration-to-extinction approach. *Bioresource Technology*, *100*(9), 2546–2550.
- Ahuja, S. K., Ferreira, G. M., and Moreira, A. R. (2004). Utilization of enzymes for environmental applications. *Critical Reviews in Biotechnology*, *24*(2-3), 125–154.
- Akutsu, Y., Li, Y.-Y., Harada, H., and Yu, H.-Q. (2009). Effects of temperature and substrate concentration on biological hydrogen production from starch. *International Journal of Hydrogen Energy*, *34*(6), 2558–2566.
- Andrews, S. C., Berks, B. C., McClay, J., Ambler, A., Quail, M. A., Golby, P., and Guest, J. R. (1997). A 12-cistron *Escherichia coli* operon (*hyf*) encoding a putative proton-translocating formate hydrogenlyase system. *Microbiology*, *143*, 3633–3647.
- Argaman, L., and Altuvia, S. (2000). *fhlA* repression by OxyS RNA: kissing complex formation at two sites results in a stable antisense-target RNA complex. *Journal of Molecular Biology*, *300*(5), 1101–1112.
- Argun, H., Kargi, F., Kapdan, I., and Oztekin, R. (2008). Biohydrogen production by dark fermentation of wheat powder solution: Effects of C/N and C/P ratio on hydrogen yield and formation rate. *International Journal of Hydrogen Energy*, *33*(7), 1813–1819.
- Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Baba, M., Datsenko, K. A., Tomita, M., Wanner, B. L., and Mori, H. (2006). Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Molecular Systems Biology*, *2*, 2006–2008.
- Baffert, C., Demuez, M., Cournac, L., Burlat, B., Guigliarelli, B., Bertrand, P., Girbal, L., and Léger, C. (2008). Hydrogen-Activating Enzymes: Activity Does Not Correlate with Oxygen Sensitivity. *Angewandte Chemie*, *120*(11), 2082–2084.

- Baghchehsaraee, B., Nakhla, G., Karamanev, D., Margaritis, A., and Reid, G. (2008). The effect of heat pretreatment temperature on fermentative hydrogen production using mixed cultures. *International Journal of Hydrogen Energy*, 33(15), 4064–4073.
- Bagramyan, K., Mnatsakanyan, N., Poladian, A., Vassilian, A., and Trchounian, A. (2002). The roles of hydrogenases 3 and 4, and the F₀F₁-ATPase, in H₂ production by *Escherichia coli* at alkaline and acidic pH. *FEBS Letters*, 516(1-3), 172–178.
- Bagramyan, K., and Trchounian, A. (2003). Structural and functional features of formate hydrogen lyase, an enzyme of mixed-acid fermentation from *Escherichia coli*. *Biochemistry*, 68(11), 1159–1170.
- Bakonyi, P., Nemestóthy, N., and Bélafi-Bakó, K. (2012). Comparative study of various *E. coli* strains for biohydrogen production applying response surface methodology. *The Scientific World Journal*, 2012, 793–819.
- Balakirev, E. S., and Ayala, F. J. (2003). Pseudogenes: are they “junk” or functional DNA? *Annual Review of Genetics*, 37, 123–151.
- Ballantine, S. P., and Boxer, D. H. (1986). Isolation and characterisation of a soluble active fragment of hydrogenase isoenzyme 2 from the membranes of anaerobically grown *Escherichia coli*. *European Journal of Biochemistry*, 156(2), 277–284.
- Bertero, M. G., Rothery, R. A., Palak, M., Hou, C., Lim, D., Blasco, F., Weiner, J. H., and Strynadka, N. C. J. (2003). Insights into the respiratory electron transfer pathway from the structure of nitrate reductase A. *Nature Structural Biology*, 10(9), 681–687.
- Bisaillon, A., Turcot, J., and Hallenbeck, P. (2006). The effect of nutrient limitation on hydrogen production by batch cultures of *Escherichia coli*. *International Journal of Hydrogen Energy*, 31(11), 1504–1508.
- Blattner, F. R., Plunkett, G., Bloch, C. A., Perna, N. T., Burland, V., Riley, M., Collado-Vides, J., Glasner, J. D., Rode, C. K., Mayhew, G. F., Gregor, J., Davis, N.W., Kirkpatrick, H. A., Goeden, M. A., Rose, D. J., Mau, B., and Shao, Y. (1997). The complete genome sequence of *Escherichia coli* K-12. *Science*, 277(5331), 1453–1462.
- Bleijlevens, B., Buhrke, T., van der Linden, E., Friedrich, B., and Albracht, S. P. J. (2004). The auxiliary protein HypX provides oxygen tolerance to the soluble [NiFe]-hydrogenase of *Ralstonia eutropha* H16 by way of a cyanide ligand to nickel. *The Journal of Biological Chemistry*, 279(45), 46686–46691.
- Blokesch, M., Magalon, A., and Böck, A. (2001). Interplay between the specific chaperone-like proteins HybG and HypC in maturation of hydrogenases 1, 2, and 3 from *Escherichia coli*. *Journal of Bacteriology*, 183(9), 2817–2822.
- Böck, A., and Sawers, G. (1996). Cellular and molecular biology. In Neidhardt, F.C., Curtiss, J. R., II, Ingraham, J. L., Lin, E. C. C., Low, K. B., Magasanik, B. (eds), *Escherichia coli* and *Salmonella* (pp. 262–282). Washington, ASM Press.

- Böhm, R., Sauter, M., and Böck, A. (1990). Nucleotide sequence and expression of an operon in *Escherichia coli* coding for formate hydrogenlyase components. *Molecular Microbiology*, 4(2), 231–243.
- Bond, D. R., Holmes, D. E., Tender, L. M., and Lovley, D. R. (2002). Electrode-reducing microorganisms that harvest energy from marine sediments. *Science*, 295(5554), 483–485.
- Bouallagui, H., Touhami, Y., Ben Cheikh, R., and Hamdi, M. (2005). Bioreactor performance in anaerobic digestion of fruit and vegetable wastes. *Process Biochemistry*, 40(3-4), 989–995.
- Brøndsted, L., and Atlung, T. (1994). Anaerobic regulation of the hydrogenase 1 (*hya*) operon of *Escherichia coli*. *Journal of Bacteriology*, 176(17), 5423–5428.
- Cakır, A., Ozmihci, S., and Kargi, F. (2010). Comparison of bio-hydrogen production from hydrolyzed wheat starch by mesophilic and thermophilic dark fermentation. *International Journal of Hydrogen Energy*, 35(24), 13214–13218.
- Casjens, S., Palmer, N., Van Vugt, R., Mun Huang, W., Stevenson, B., Rosa, P., Lathigra, R., Sutton, G., Peterson, J., Dodson, R. J., Haft, D., Hickey, E., Gwinn, M., White, O., and Fraser, C. (2002). A bacterial genome in flux: the twelve linear and nine circular extrachromosomal DNAs in an infectious isolate of the Lyme disease spirochete *Borrelia burgdorferi*. *Molecular Microbiology*, 35(3), 490–516.
- Chairattanamakorn, P., Penthamkeerati, P., Reungsang, A., Lo, Y.-C., Lu, W.-B., and Chang, J.-S. (2009). Production of biohydrogen from hydrolyzed bagasse with thermally preheated sludge. *International Journal of Hydrogen Energy*, 34(18), 7612–7617.
- Chang, C. Y. and Lin, F. Y. (2006). Calcium effect on fermentative hydrogen production in an anaerobic up-flow sludge blanket system. *Water Science and Technology*, 54, 105–112.
- Chen, R., Wang, Y.-Z., Liao, Q., Zhu, X., and Xu, T.-F. (2013). Hydrolysates of lignocellulosic materials for biohydrogen production. *BMB reports*, 46(5), 244–251.
- Chen, S.-D., Sheu, D.-S., Chen, W.-M., Lo, Y.-C., Huang, T.-I., Lin, C.-Y., and Chang, J.-S. (2007). Dark hydrogen fermentation from hydrolyzed starch treated with recombinant amylase originating from *Caldimonas taiwanensis* On1. *Biotechnology progress*, 23(6), 1312–1320.
- Chen, W., Chen, S., Kumarkhanal, S., and Sung, S. (2006). Kinetic study of biological hydrogen production by anaerobic fermentation. *International Journal of Hydrogen Energy*, 31(15), 2170–2178.
- Chen, W., Tseng, Z., Lee, K., and Chang, J. (2005). Fermentative hydrogen production with CGS5 isolated from anaerobic sewage sludge. *International Journal of Hydrogen Energy*, 30(10), 1063–1070.
- Cherepanov, P. P., and Wackernagel, W. (1995). Gene disruption in *Escherichia coli*: TcR and KmR cassettes with the option of F1p-catalyzed excision of the antibiotic-resistance determinant. *Gene*, 158(1), 9–14.

- Chong, M., Rahim, R., Shirai, Y., and Hassan, M. (2009a). Biohydrogen production by *Clostridium butyricum* EB6 from palm oil mill effluent. *International Journal of Hydrogen Energy*, 34(2), 764–771.
- Chong, M.-L., Abdul Rahman, N. A., Rahim, R. A., Aziz, S. A., Shirai, Y., and Hassan, M. A. (2009b). Optimization of biohydrogen production by *Clostridium butyricum* EB6 from palm oil mill effluent using response surface methodology. *International Journal of Hydrogen Energy*, 34(17), 7475–7482.
- Cole, S. T., Eiglmeier, K., Parkhill, J., James, K. D., Thomson, N. R., Wheeler, P. R., Honoré, N., Garnier, T., Churcher, C., Harris, D., Mungall, K., Basham, D., Brown, D., Chillingworth, T., Connor, R., Davies, R. M., Devlin, K., Duthoy, S., Feltwell, T., Fraser, A., Hamlin, N., Holroyd, S., Hornsby, T., Jagels, K., Lacroix, C., Maclean, J., Moule, S., Murphy, L., Oliver, K., Quail, M. A., Rajandream, M. A., Rutherford, K. M., Rutter, S., Seeger, K., Simon, S., Simmonds, M., Skelton, J., Squares, R., Squares, S., Stevens, K., Taylor, K., Whitehead, S., Woodward, J. R., and Barrell, B. G. (2001). Massive gene decay in the *leprosy bacillus*. *Nature*, 409(6823), 1007–1011.
- Das, D. (2001). Hydrogen production by biological processes: a survey of literature. *International Journal of Hydrogen Energy*, 26(1), 13–28.
- Das, D., and Veziroglu, T. (2008). Advances in biological hydrogen production processes. *International Journal of Hydrogen Energy*, 33(21), 6046–6057.
- Datsenko, K. A., and Wanner, B. L. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proceedings of the National Academy of Sciences of the United States of America*, 97(12), 6640–6645.
- de Souza, R. F., Padilha, J. C., Gonçalves, R. S., de Souza, M. O., and Rault-Berthelot, J. (2007). Electrochemical hydrogen production from water electrolysis using ionic liquid as electrolytes: Towards the best device. *Journal of Power Sources*, 164(2), 792–798.
- Doberenz, C., Zorn, M., Falke, D., Nannemann, D., Hunger, D., Beyer, L., Ihling, C. H., Meiler, J., Sinz, A., and Sawers, R. G. (2014). Pyruvate formate-lyase interacts directly with the formate channel FocA to regulate formate translocation. *Journal of Molecular Biology*, 426(15), 2827–2839.
- Domka, J., Lee, J., Bansal, T., and Wood, T. K. (2007). Temporal gene-expression in *Escherichia coli* K-12 biofilms. *Environmental Microbiology*, 9(2), 332–3346.
- Dong-Jie, N., Jing-Yuan, W., Bao-Ying, W., and You-Cai, Z. (2011). Effect of Mo-containing additives on biohydrogen fermentation from cassava's stillage. *International Journal of Hydrogen Energy*, 36(9), 5289–5295.
- Dubini, A. (2002). How bacteria get energy from hydrogen: a genetic analysis of periplasmic hydrogen oxidation in *Escherichia coli*. *International Journal of Hydrogen Energy*, 27(11-12), 1413–1420.
- Esnault, C., Maestre, J., and Heidmann, T. (2000). Human LINE retrotransposons generate processed pseudogenes. *Nature Genetics*, 24(4), 363–367.

- Feng, X., Wang, H., Wang, Y., Wang, X., and Huang, J. (2010). Biohydrogen production from apple pomace by anaerobic fermentation with river sludge. *International Journal of Hydrogen Energy*, 35(7), 3058–3064.
- Forzi, L., and Sawers, R. G. (2007). Maturation of [NiFe]-hydrogenases in *Escherichia coli*. *Biometals*, 20(3-4), 565–578.
- Francis, K., Patel, P., Wendt, J. C., and Shanmugam, K. T. (1990). Purification and characterization of two forms of hydrogenase isoenzyme 1 from *Escherichia coli*. *Journal of Bacteriology*, 172(10), 5750–5757.
- Frey, M. (2002). Hydrogenases: hydrogen-activating enzymes. *ChemBiochem*, 3(2-3), 153–160.
- Gadow, S. I., Li, Y.-Y., and Liu, Y. (2012). Effect of temperature on continuous hydrogen production of cellulose. *International Journal of Hydrogen Energy*, 37(20), 15465–15472.
- Ghosh, D., Bisailon, A., and Hallenbeck, P. C. (2013). Increasing the metabolic capacity of *Escherichia coli* for hydrogen production through heterologous expression of the *Ralstonia eutropha* SH operon. *Biotechnology for biofuels*, 6(1), 122.
- Gibson, L. J. (1994). Pseudogene and Origins. *Origins*, 21, 91-108.
- Goncalves, I. (2000). Nature and Structure of Human Genes that Generate Retroseudogenes. *Genome Research*, 10(5), 672–678.
- Griffiths, A. J., Miller, J. H., Suzuki, D. T., Lewontin, R. C., and Gelbart, W. M. (2000). Bacterial insertion sequences. W. H. Freeman.
- Grot S. A., Logan, B. E., H. Liu, S., and Mallouk, T. (2010). Bio-electrochemically assisted microbial reactor that generates hydrogen gas and methods of generating hydrogen gas, 113.
- Hakobyan, M., Sargsyan, H., and Bagramyan, K. (2005). Proton translocation coupled to formate oxidation in anaerobically grown fermenting *Escherichia coli*. *Biophysical Chemistry*, 115(1), 55–61.
- Hallenbeck, C. P., and Benemann, J. R. (2002). Biological hydrogen production; fundamentals and limiting processes. *International Journal of Hydrogen Energy*, 27(11-12), 1185–1193.
- Hallenbeck, P. C. (2009). Fermentative hydrogen production: Principles, progress, and prognosis. *International Journal of Hydrogen Energy*, 34(17), 7379–7389.
- Hallenbeck, P. C. (2011). *Microbial Technologies in Advanced Biofuels Production*. Springer Science & Business Media.
- Hallenbeck, P. C. (2014). Microbial paths to renewable hydrogen production. *Biofuels*, 2, 285–302.
- Hallenbeck, P. C., and Ghosh, D. (2009). Advances in fermentative biohydrogen production: the way forward? *Trends in Biotechnology*, 27(5), 287–97.

- Harrison, P. M. (2001). Digging for dead genes: an analysis of the characteristics of the pseudogene population in the *Caenorhabditis elegans* genome. *Nucleic Acids Research*, 29(3), 818–830.
- Hawkes, F., Hussy, I., Kyazze, G., Dinsdale, R., and Hawkes, D. (2007). Continuous dark fermentative hydrogen production by mesophilic microflora: Principles and progress. *International Journal of Hydrogen Energy*, 32(2), 172–184.
- Hayashi, K., Morooka, N., Yamamoto, Y., Fujita, K., Isono, K., Choi, S., Ohtsubo, E., Baba, T., Wanner, B. L., Mori, H., and Horiuchi, T. (2006). Highly accurate genome sequences of *Escherichia coli* K-12 strains MG1655 and W3110. *Molecular Systems Biology*, 2(1), 2006–2007.
- Herzberg, M., Kaye, I. K., Peti, W., and Wood, T. K. (2006). YdgG (TqsA) controls biofilm formation in *Escherichia coli* K-12 through autoinducer 2 transport. *Journal of Bacteriology*, 188(2), 587–598.
- Holladay, J. D., Hu, J., King, D. L., and Wang, Y. (2009). An overview of hydrogen production technologies. *Catalysis Today*, 139(4), 244–260.
- Holt, P. K., Barton, G. W., Wark, M., and Mitchell, C. A. (2002). A quantitative comparison between chemical dosing and electrocoagulation. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 211(2-3), 233–248.
- Hopp, T. P., and Woods, K. R. (1981). Prediction of protein antigenic determinants from amino acid sequences. *Proceedings of the National Academy of Sciences of the United States of America*, 78(6), 3824–3828.
- Hube, M., Blokesch, M., and Böck, A. (2002). Network of hydrogenase maturation in *Escherichia coli*: role of accessory proteins HypA and HybF. *Journal of Bacteriology*, 184(14), 3879–3885.
- Jaapar, S. Z. S., Ali, E., Kalil, M. S., and Anuar, N. (2011). Effects of Different Initial pH, Argon Gas and Nitrogen Gas on Cell Growth and Hydrogen Production using *Rhodobacter sphaeroides*. *Bacteriology Journal*, 1(1), 8–15.
- Janssen, H. J. C., Bringmann, B., Emontsa, V. S. (2004). Safety-related studies on hydrogen production in high-pressure electrolyzers. *International Journal of Hydrogen Energy*, 29(7), 759–770.
- Kamalaskar, L. B., Dhakephalkar, P. K., Meher, K. K., and Ranade, D. R. (2010). High biohydrogen yielding *Clostridium* sp. DMHC-10 isolated from sludge of distillery waste treatment plant. *International Journal of Hydrogen Energy*, 35(19), 10639–10644.
- Kapdan, I. K., and Kargi, F. (2006). Bio-hydrogen production from waste materials. *Enzyme and Microbial Technology*, 38(5), 569–582.
- Kelleher, B., Leahy, J., Henihan, A., O'Dwyer, T., Sutton, D., and Leahy, M. (2002). Advances in poultry litter disposal technology – a review. *Bioresource Technology*, 83(1), 27–36.

- Keseler, Collado-Vides, J., Santos-Zavaleta, A., Peralta-Gil, M., Gama-Castro, S., Muniz-Rascado, L., César B. M., Suzanne, P., Markus, K., Tomer A., Pallavi, K., Aaron, S., John, P., Mario, L., Carol, F., Malabika, S., Alexander, G. S., Amanda, M., Ian, P., Robert P. G., and Karp, P. D. (2010). EcoCyc: a comprehensive database of *Escherichia coli* biology. *Nucleic Acids Research*, *39*, 583–590.
- Keseler, I. M., Collado-Vides, J., Gama-Castro, S., Ingraham, J., Paley, S., Paulsen, I. T., Peralta-Gil, M., and Karp, P. D. (2005). EcoCyc: a comprehensive database resource for *Escherichia coli*. *Nucleic acids research*, *33*, 334–337.
- Khanal, S. (2003). Biological hydrogen production: effects of pH and intermediate products. *International Journal of Hydrogen Energy*, *29*(11), 1123–1131.
- Khemis, M., Leclerc, J.-P., Tanguy, G., Valentin, G., and Lapicque, F. (2006). Treatment of industrial liquid wastes by electrocoagulation: Experimental investigations and an overall interpretation model. *Chemical Engineering Science*, *61*(11), 3602–3609.
- Kim, D.-H., Kim, S.-H., Jung, K.-W., Kim, M.-S., and Shin, H.-S. (2011a). Effect of initial pH independent of operational pH on hydrogen fermentation of food waste. *Bioresource Technology*, *102*(18), 8646–8652.
- Kim, D.-H., Kim, S.-H., Kim, H.-W., Kim, M.-S., and Shin, H.-S. (2011b). Sewage sludge addition to food waste synergistically enhances hydrogen fermentation performance. *Bioresource Technology*, *102*(18), 8501–8506.
- Kim, J. Y. H., Jo, B. H., and Cha, H. J. (2010). Production of biohydrogen by recombinant expression of [NiFe]-hydrogenase 1 in *Escherichia coli*. *Microbial Cell Factories*, *9*(1), 54.
- King, P. W., and Przybyla, A. E. (1999). Response of *hya* Expression to External pH in *Escherichia coli*. *Journal of Bacteriology*, *181*(17), 5250–5256.
- Kirkpatrick, C., Maurer, L. M., Oyelakin, N. E., Yoncheva, Y. N., Maurer, R., and Slonczewski, J. L. (2001). Acetate and formate stress: opposite responses in the proteome of *Escherichia coli*. *Journal of Bacteriology*, *183*(21), 6466–6477.
- Kitagawa, M., Ara, T., Arifuzzaman, M., Ioka-Nakamichi, T., Inamoto, E., Toyonaga, H., and Mori, H. (2005). Complete set of ORF clones of *Escherichia coli* ASKA library (a complete set of *E. coli* K-12 ORF archive): unique resources for biological research. *DNA Research*, *12*(5), 291–299.
- Kline, K. A., Fälker, S., Dahlberg, S., Normark, S., and Henriques-Normark, B. (2009). Bacterial adhesins in host-microbe interactions. *Cell Host & Microbe*, *5*(6), 580–592.
- Korea, C.-G., Badouraly, R., Prevost, M.-C., Ghigo, J.-M., and Beloin, C. (2010). *Escherichia coli* K-12 possesses multiple cryptic but functional chaperone-usher fimbriae with distinct surface specificities. *Environmental Microbiology*, *12*(7), 1957–1977.

- Korea, C.-G., Ghigo, J.-M., and Beloin, C. (2011). The sweet connection: Solving the riddle of multiple sugar-binding fimbrial adhesins in *Escherichia coli*: Multiple *E. coli* fimbriae form a versatile arsenal of sugar-binding lectins potentially involved in surface-colonisation and tissue tro. *BioEssays*, 33(4), 300–311.
- Korsa, I., and Böck, A. (1997). Characterization of *fhlA* mutations resulting in ligand-independent transcriptional activation and ATP hydrolysis. *Journal of Bacteriology*, 179(1), 41–45.
- Kotay, S. M., and Das, D. (2007). Microbial hydrogen production with *Bacillus coagulans* IIT-BT S1 isolated from anaerobic sewage sludge. *Bioresource Technology*, 98(6), 1183–1190.
- Laurinavichene, T. V., and Tsygankov, A. A. (2001). H₂ consumption by *Escherichia coli* coupled via hydrogenase 1 or hydrogenase 2 to different terminal electron acceptors. *FEMS Microbiology Letters*, 202(1), 121–124.
- Laurinavichene, T. V., Zorin, N. A., and Tsygankov, A. A. (2002). Effect of redox potential on activity of hydrogenase 1 and hydrogenase 2 in *Escherichia coli*. *Archives of Microbiology*, 178(6), 437–442.
- Laurinavichene, T. V., Chanal, A., Wu, L. F., and Tsygankov, A. A. (2001). Effect of O₂, H₂ and redox potential on the activity and synthesis of hydrogenase 2 in *Escherichia coli*. *Research in Microbiology*, 152(9), 793–798.
- Lay, J.-J. (2000). Modeling and optimization of anaerobic digested sludge converting starch to hydrogen. *Biotechnology and Bioengineering*, 68(3), 269–278.
- Lay, J.-J., Fan, K.-S., Hwang, J.-I., Chang, J.-I., and Hsu, P.-C. (2005). Factors Affecting Hydrogen Production from Food Wastes by *Clostridium*-Rich Composts. *Journal of Environmental Engineering*, 131(4), 595–602.
- Lee, J., Maeda, T., Hong, S. H., and Wood, T. K. (2009). Reconfiguring the quorum-sensing regulator SdiA of *Escherichia coli* to control biofilm formation via indole and N-acylhomoserine lactones. *Applied and Environmental Microbiology*, 75(6), 1703–1716.
- Levin, D. (2004). Biohydrogen production: prospects and limitations to practical application. *International Journal of Hydrogen Energy*, 29(2), 173–185.
- Levin, D. B., Zhu, H., Beland, M., Cicek, N., and Holbein, B. E. (2007). Potential for hydrogen and methane production from biomass residues in Canada. *Bioresource Technology*, 98(3), 654–660.
- Li, C., and Fang, H. H. P. (2007a). Fermentative Hydrogen Production From Wastewater and Solid Wastes by Mixed Cultures. *Critical Reviews in Environmental Science and Technology*, 37(1), 1–39.
- Li, C., and Fang, H. H. P. (2007b). Inhibition of heavy metals on fermentative hydrogen production by granular sludge. *Chemosphere*, 67(4), 668–673.
- Li, Q., Xing, D., Ren, N., Zhao, L., and Song, Y. (2006). Hydrogen production and enzyme activity of acidophilic strain X-29 at different C/N ratio. *Huan Jing Ke Xue*, 27(4), 810–814.

- Liebott, P.-P., Leroux, F., Burlat, B., Dementin, S., Baffert, C., Lautier, T., Fourmond, Vincent Ceccaldi, P., Cavazza, C., Meynial-Salles, I., Soucaille, P., Fontecilla-Camps, J. C., Guigliarelli, B., Bertrand, P., Rousset, M., and Léger, C. (2010). Relating diffusion along the substrate tunnel and oxygen sensitivity in hydrogenase. *Nature Chemical Biology*, 6(1), 63–70.
- Lin, C. (2008). Heavy metal effects on fermentative hydrogen production using natural mixed microflora. *International Journal of Hydrogen Energy*, 33(2), 587–593.
- Liu, H., Grot, S., and Logan, B. E. (2005). Electrochemically Assisted Microbial Production of Hydrogen from Acetate. *Environmental Science & Technology*, 39(11), 4317–4320.
- Liu, H., Ramnarayanan, R., and Logan, B. E. (2004). Production of Electricity during Wastewater Treatment Using a Single Chamber Microbial Fuel Cell. *Environmental Science & Technology*, 38(7), 2281–2285.
- Liu, Y., and Whitman, W. B. (2008). Metabolic, phylogenetic, and ecological diversity of the methanogenic archaea. *Annals of the New York Academy of Sciences*, 1125, 171–189.
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*, 25(4), 402–408.
- Ma, S., Wang, H., Wang, Y., Bu, H., and Bai, J. (2011). Bio-hydrogen production from cornstarch wastes by orthogonal design method. *Renewable Energy*, 36(2), 709–713.
- Maeda, T., Sanchez-Torres, V., and Wood, T. K. (2007a). Enhanced hydrogen production from glucose by metabolically engineered *Escherichia coli*. *Applied Microbiology and Biotechnology*, 77(4), 879–890.
- Maeda, T., Sanchez-Torres, V., and Wood, T. K. (2007b). *Escherichia coli* hydrogenase 3 is a reversible enzyme possessing hydrogen uptake and synthesis activities. *Applied Microbiology and Biotechnology*, 76(5), 1035–1042.
- Maeda, T., Sanchez-Torres, V., and Wood, T. K. (2008a). Metabolic engineering to enhance bacterial hydrogen production. *Microbial Biotechnology*, 1(1), 30–9.
- Maeda, T., Sanchez-Torres, V., and Wood, T. K. (2008b). Protein engineering of hydrogenase 3 to enhance hydrogen production. *Applied Microbiology and Biotechnology*, 79(1), 77–86.
- Maeda, T., Vardar, G., Self, W. T., and Wood, T. K. (2007). Inhibition of hydrogen uptake in *Escherichia coli* by expressing the hydrogenase from the *Cyanobacterium Synechocystis* sp. PCC 6803. *BMC Biotechnology*, 7(1), 25.
- Mathews, J., and Wang, G. (2009). Metabolic pathway engineering for enhanced biohydrogen production. *International Journal of Hydrogen Energy*, 34(17), 7404–7416.
- Meherkotay, S., and Das, D. (2008). Biohydrogen as a renewable energy resource—Prospects and potentials. *International Journal of Hydrogen Energy*, 33(1), 258–263.

- Menon, N. K., Chatelus, C. Y., Dervartanian, M., Wendt, J. C., Shanmugam, K. T., Peck, H. D., and Przybyla, A. E. (1994). Cloning, sequencing, and mutational analysis of the *hyb* operon encoding *Escherichia coli* hydrogenase 2. *Journal of Bacteriology*, *176*(14), 4416–4423.
- Meza, E., Becker, J., Bolivar, F., Gosset, G., and Wittmann, C. (2012). Consequences of phosphoenolpyruvate:sugar phosphotransferase system and pyruvate kinase isozymes inactivation in central carbon metabolism flux distribution in *Escherichia coli*. *Microbial Cell Factories*, *11*(1), 127.
- Mighell, A. J., Smith, N. R., Robinson, P. A., and Markham, A. F. (2000). Vertebrate pseudogenes. *FEBS letters*, *468*(2-3), 109–114.
- Mnatsakanyan, N., Bagramyan, K., and Trchounian, A. (2004). Hydrogenase 3 but not hydrogenase 4 is major in hydrogen gas production by *Escherichia coli* formate hydrogenlyase at acidic pH and in the presence of external formate. *Cell Biochemistry and Biophysics*, *41*(3), 357–366.
- Mohd Yasin, N. H., Abdul Rahman, N. A., Ismail, F., Yusof, M. Z. M., and Hassan, M. A. (2009). Effect of different temperature, initial pH and substrate composition on biohydrogen production from food waste in batch fermentation. *Asian Journal of Biotechnology*, *1*, 42–50.
- Mohd Yasin, N. H., Rahman, N. A., Man, H. C., Mohd Yusoff, M. Z., and Hassan, M. A. (2011). Microbial characterization of hydrogen-producing bacteria in fermented food waste at different pH values. *International Journal of Hydrogen Energy*, *36*(16), 9571–9580.
- Mohd Yasin, N. H., Toshinari, M., Fukuzaki, M., Miyazaki, T., Hakiman Che Maail, C. M., Ariffin, H., and Wood, T. K. (2013). Biohydrogen production from oil palm frond juice and sewage sludge by a metabolically engineered *Escherichia coli* strain. *International Journal of Hydrogen Energy*, *38*(25), 10277–10283.
- Mohd Yusoff, M. Z., Hashiguchi, Y., Maeda, T., and Wood, T. K. (2013). Four products from *Escherichia coli* pseudogenes increase hydrogen production. *Biochemical and Biophysical Research Communications*, *439*(4), 576–579.
- Mohd Yusoff, M. Z., Maeda, T., Sanchez-Torres, V., Ogawa, H. I., Shirai, Y., Hassan, M. A., and Wood, T. K. (2012). Uncharacterized *Escherichia coli* proteins YdjA and YhjY are related to biohydrogen production. *International Journal of Hydrogen Energy*, *37*(23), 17778–17787.
- Mu, Y., Zheng, X., Yu, H., and Zhu, R. (2006). Biological hydrogen production by anaerobic sludge at various temperatures. *International Journal of Hydrogen Energy*, *31*(6), 780–785.
- Mudhoo, A., Forster-Carneiro, T., and Sánchez, A. (2011). Biohydrogen production and bioprocess enhancement: A review. *Critical Reviews in Biotechnology*, *31*(3), 250–263.
- Nandi, R., and Sengupta, S. (1998). Microbial Production of Hydrogen: An Overview. *Critical Reviews in Microbiology*, *24*(1), 61–84.
- Nath, K., and Das, D. (2004). Improvement of fermentative hydrogen production: various approaches. *Applied Microbiology and Biotechnology*, *65*(5), 520–529.

- Nishihara, H., Miyashita, Y., Aoyama, K., Kodama, T., Igarashi, Y., and Takamura, Y. (1997). Characterization of an extremely thermophilic and oxygen-stable membrane-bound hydrogenase from a marine hydrogen-oxidizing bacterium *Hydrogenovibrio marinus*. *Biochemical and Biophysical Research Communications*, 232(3), 766–770.
- Nuccio, S.-P., and Bäumlér, A. J. (2007). Evolution of the chaperone/usher assembly pathway: fimbrial classification goes Greek. *Microbiology and Molecular Biology Reviews*, 71(4), 551–575.
- Ogata, H., Audic, S., Renesto-Audiffren, P., Fournier, P. E., Barbe, V., Samson, D., Roux, V., Cossart, P., Weissenbach, J., Claverie, J. M., and Raoult, D. (2001). Mechanisms of evolution in *Rickettsia conorii* and *R. prowazekii*. *Science*, 293(5537), 2093–2098.
- Ogino, H., Miura, T., Ishimi, K., Seki, M., and Yoshida, H. (2005). Hydrogen production from glucose by anaerobes. *Biotechnology Progress*, 21(6), 1786–1788.
- Oh, Y. K., Seol, E. H., Kim, J. R., and Park, S. (2003). Fermentative biohydrogen production by a new chemoheterotrophic bacterium *Citrobacter* sp. Y19. *International Journal of Hydrogen Energy*, 28(12), 1353–1359.
- Oleszkiewicz, J. A., and Sharma, V. K. (1990). Stimulation and inhibition of anaerobic processes by heavy metals—A review. *Biological Wastes*, 31(1), 45–67.
- Ostrem, K. M., Millrath, K., and Themelis, N. J. (2004). Combining Anaerobic Digestion and Waste-to-Energy. In *12th Annual North American Waste-to-Energy Conference* (pp. 265–271). ASME.
- Pakarinen, O. (2008). Batch dark fermentative hydrogen production from grass silage: The effect of inoculum, pH, temperature and VS ratio. *International Journal of Hydrogen Energy*, 33(2), 594–601.
- Parkhill, J., Wren, B. W., Mungall, K., Ketley, J. M., Churcher, C., Basham, D., hillingsworth, T Davies, R. M., Feltwell, T., Holroyd, S., Jagels, K., Karlyshev, A. V., Moule, S., Pallen, M. J., Penn, C. W., Quail, M. A., Rajandream, M. A., Rutherford, K. M., van Vliet, A. H., Whitehead, S., and Barrell, B. G. (2000). The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hypervariable sequences. *Nature*, 403(6770), 665–668.
- Parkhill, J., Wren, B. W., Thomson, N. R., Titball, R. W., Holden, M. T., Prentice, M. B., Sebahia, M., James, K. D., Churcher, C., Mungall, K. L., Baker, S., Basham, D., Bentley, S. D., Brooks, K., Cerdeño-Tárraga, A. M., Chillingworth, T., Cronin, A., Davies, R. M., Davis, P., Dougan, G., Feltwell, T., Hamlin, N., Holroyd, S., Jagels, K., Karlyshev, A. V., Leather, S., Moule, S., Oyston, P. C., Quail, M., Rutherford, K., Simmonds, M., Skelton, J., Stevens, K., Whitehead, S., and Barrell, B. G. (2001). Genome sequence of *Yersinia pestis*, the causative agent of plague. *Nature*, 413(6855), 523–527.
- Patterson, T., Esteves, S., Dinsdale, R., Guwy, A., and Maddy, J. (2013). Life cycle assessment of biohydrogen and biomethane production and utilisation as a vehicle fuel. *Bioresource Technology*, 131, 235–245.

- Pavlicek, A., Gentles, A. J., Paces, J., Paces, V., and Jurka, J. (2006). Retroposition of processed pseudogenes: the impact of RNA stability and translational control. *Trends in Genetics*, 22(2), 69–73.
- Pavliček, A., Pačes, J., Zika, R., and Hejnar, J. (2002). Length distribution of long interspersed nucleotide elements (LINEs) and processed pseudogenes of human endogenous retroviruses: implications for retrotransposition and pseudogene detection. *Gene*, 300(1-2), 189–194.
- Pawar, S. S., and van Niel, E. W. J. (2013). Thermophilic biohydrogen production: how far are we? *Applied Microbiology and Biotechnology*, 97(18), 7999–8009.
- Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research*, 29(9), 45.
- Pinske, C., McDowall, J. S., Sargent, F., and Sawers, R. G. (2012). Analysis of hydrogenase 1 levels reveals an intimate link between carbon and hydrogen metabolism in *Escherichia coli* K-12. *Microbiology*, 158(3), 856–866.
- Laura Poliseno* (2012). *Pseudogenes: Newly Discovered Players in Human Cancer*, Sci Signal 5(242)
- Qian, C., Chen, L., Rong, H., and Yuan, X. (2011). Hydrogen production by mixed culture of several facultative bacteria and anaerobic bacteria. *Progress in Natural Science: Materials International*, 21(6), 506–511.
- Rachman, M. A., Furutani, Y., Nakashimada, Y., Kakizono, T., and Nishio, N. (1997). Enhanced hydrogen production in altered mixed acid fermentation of glucose by *Enterobacter aerogenes*. *Journal of Fermentation and Bioengineering*, 83(4), 358–363.
- Ray, S., Chowdhury, N., Lalman, J. A., Seth, R., and Biswas, N. (2008). Impact of Initial pH and Linoleic Acid (C18:2) on Hydrogen Production by a Mesophilic Anaerobic Mixed Culture. *Journal of Environmental Engineering*, 134(2), 110–117.
- Redwood, M. D., Mikheenko, I. P., Sargent, F., and Macaskie, L. E. (2008). Dissecting the roles of *Escherichia coli* hydrogenases in biohydrogen production. *FEMS Microbiology Letters*, 278(1), 48–55.
- Richard, D. J., Sawers, G., Sargent, F., McWalter, L., and Boxer, D. H. (1999). Transcriptional regulation in response to oxygen and nitrate of the operons encoding the [NiFe] hydrogenases 1 and 2 of *Escherichia coli*. *Microbiology*, 145, 2903–2912.
- Rosales-Colunga, L. M., Razo-Flores, E., Ordoñez, L. G., Alatraste-Mondragón, F., and De León-Rodríguez, A. (2010). Hydrogen production by *Escherichia coli* ΔhycA ΔlacI using cheese whey as substrate. *International Journal of Hydrogen Energy*, 35(2), 491–499.
- Rossmann, R., Sawers, G., and Böck, A. (1991). Mechanism of regulation of the formate-hydrogenlyase pathway by oxygen, nitrate, and pH: definition of the formate regulon. *Molecular Microbiology*, 5(11), 2807–2814.
- Rostrup-Nielsen, J., and Christiansen, L. J. (2011). *Concepts in Syngas Manufacture*. World Scientific.

- Rostrup-Nielsen, J. R., Sehested, J., and Nørskov, J. K. (2002). Hydrogen and synthesis gas by steam- and CO₂ reforming. *Advances in Catalysis*, 47, 65–139.
- Rouchka, E. C., and Cha, I. E. (2009). Current Trends in Pseudogene Detection and Characterization. *Current Bioinformatics*, 4(8), 112–119.
- Sakai, H., Itoh, T. and Gojobori, T. (2008). *Processed Pseudogenes and Their Functional Resurrection in the Human and Mouse Genomes*. John Wiley & Sons, Ltd Chichester, UK.
- Salih, O., Remaut, H., Waksman, G., and Orlova, E. V. (2008). Structural analysis of the Saf pilus by electron microscopy and image processing. *Journal of Molecular Biology*, 379(1), 174–187.
- Sanchez-Torres, V., Maeda, T., and Wood, T. K. (2009). Protein engineering of the transcriptional activator FhlA To enhance hydrogen production in *Escherichia coli*. *Applied and Environmental Microbiology*, 75(17), 5639–5646.
- Sanchez-Torres, V., Mohd Yusoff, M. Z., Nakano, C., Maeda, T., Ogawa, H. I., and Wood, T. K. (2013). Influence of *Escherichia coli* hydrogenases on hydrogen fermentation from glycerol. *International Journal of Hydrogen Energy*, 38(10), 3905–3912.
- Sauer, F. G., Remaut, H., Hultgren, S. J., and Waksman, G. (2004). Fiber assembly by the chaperone-usher pathway. *Biochimica et Biophysica Acta*, 1694(1-3), 259–267.
- Sauter, M., Böhm, R., and Böck, A. (1992). Mutational analysis of the operon (hyc) determining hydrogenase 3 formation in *Escherichia coli*. *Molecular Microbiology*, 6(11), 1523–1532.
- Sawers, G., and Suppmann, B. (1992). Anaerobic induction of pyruvate formate-lyase gene expression is mediated by the ArcA and FNR proteins. *Journal of Bacteriology*, 174(11), 3474–3478.
- Sawers, R. G., Ballantine, S. P., and Boxer, D. H. (1985). Differential expression of hydrogenase isoenzymes in *Escherichia coli* K-12: evidence for a third isoenzyme. *Journal of Bacteriology*, 164(3), 1324–1331.
- Sawers, R. G., and Boxer, D. H. (1986). Purification and properties of membrane-bound hydrogenase isoenzyme 1 from anaerobically grown *Escherichia coli* K12. *European Journal of Biochemistry*, 156(2), 265–275.
- Schink, B. (2002). Synergistic interactions in the microbial world. *Antonie van Leeuwenhoek*, 81(1-4), 257–261.
- Seifert, K., Waligorska, M., Wojtowski, M., and Laniecki, M. (2009). Hydrogen generation from glycerol in batch fermentation process. *International Journal of Hydrogen Energy*, 34(9), 3671–3678.
- Self, W. T., Hasona, A., and Shanmugam, K. T. (2001). N-terminal truncations in the FhlA protein result in formate- and MoeA-independent expression of the hyc (formate hydrogenlyase) operon of *Escherichia coli*. *Microbiology*, 147(11), 3093–3104.

- Self, W. T., Hasona, A., and Shanmugam, K. T. (2004). Expression and regulation of a silent operon, *hyf*, coding for hydrogenase 4 isoenzyme in *Escherichia coli*. *Journal of Bacteriology*, 186(2), 580–587.
- Sen, K., and Ghosh, T. C. (2013). Pseudogenes and their composers: delving in the “debris” of human genome. *Briefings in Functional Genomics*, 12(6), 536–547.
- Seol, E., Ainala, S. K., Sekar, B. S., and Park, S. (2014). Metabolic engineering of *Escherichia coli* strains for co-production of hydrogen and ethanol from glucose. *International Journal of Hydrogen Energy*, 39(33), 19323–19330.
- Shi, X., Kim, D.-H., Shin, H.-S., and Jung, K.-W. (2013). Effect of temperature on continuous fermentative hydrogen production from *Laminaria japonica* by anaerobic mixed cultures. *Bioresource Technology*, 144, 225–231.
- Shin, J., Hyunyon, J., Kim, M., Junsim, S., and Park, T. (2007). Fermentative hydrogen production by the newly isolated *Enterobacter asburiae* SNU-1. *International Journal of Hydrogen Energy*, 32(2), 192–199.
- Slonczewski, J. L., Fujisawa, M., Dopson, M., and Krulwich, T. A. (2009). Cytoplasmic pH measurement and homeostasis in bacteria and archaea. *Advances in Microbial Physiology*, 55, 1–79.
- Steinsiek, S., and Bettenbrock, K. (2012). Glucose transport in *Escherichia coli* mutant strains with defects in sugar transport systems. *Journal of bacteriology*. 194(21), 5897-5908.
- Tam, O. H., Aravin, A. A., Stein, P., Girard, A., Murchison, E. P., Cheloufi, S., Hodges, E., Anger, M., Sachidanandam, R., Schultz, R. M., and Hannon, G. J. (2008). Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature*, 453(7194), 534–538.
- Tanisho, S., and Ishiwata, Y. (1994). Continuous hydrogen production from molasses by the bacterium *Enterobacter aerogenes*. *International Journal of Hydrogen Energy*, 19(10), 807–812.
- TeSlaa, T., and Teitell, M. A. (2014). *Conceptual Background and Bioenergetic/Mitochondrial Aspects of Oncometabolism. Methods in Enzymology* (Vol. 542). Elsevier.
- Tran, K. T., Maeda, T., and Wood, T. K. (2014). Metabolic engineering of *Escherichia coli* to enhance hydrogen production from glycerol. *Applied Microbiology and Biotechnology*, 98(10), 4757–4770.
- Trchounian, K., Pinske, C., Sawers, R. G., and Trchounian, A. (2012a). Characterization of *Escherichia coli* [NiFe]-hydrogenase distribution during fermentative growth at different pHs. *Cell Biochemistry and Biophysics*, 62(3), 433–440.
- Trchounian, K., Poladyan, A., Vassilian, A., and Trchounian, A. (2012b). Multiple and reversible hydrogenases for hydrogen production by *Escherichia coli*: dependence on fermentation substrate, pH and the F(0)F(1)-ATPase. *Critical Reviews in Biochemistry and Molecular Biology*, 47(3), 236–249.

- Trchounian, K., Sargsyan, H., and Trchounian, A. (2014). Hydrogen production by *Escherichia coli* depends on glucose concentration and its combination with glycerol at different pHs. *International Journal of Hydrogen Energy*, 39(12), 6419–6423.
- Trchounian, K., and Trchounian, A. (2013). *Escherichia coli* hydrogenase 4 (hyf) and hydrogenase 2 (hyb) contribution in H₂ production during mixed carbon (glucose and glycerol) fermentation at pH 7.5 and pH 5.5. *International Journal of Hydrogen Energy*, 38(10), 3921–3929.
- Turcot, J., Bisaillon, A., and Hallenbeck, P. (2008). Hydrogen production by continuous cultures of *Escherichia coli* under different nutrient regimes. *International Journal of Hydrogen Energy*, 33(5), 1465–1470.
- Ust'ak, S., Havrland, B., Muñoz, J. O. J., Fernández, E. C., and Lachman, J. (2007). Experimental verification of various methods for biological hydrogen production. *International Journal of Hydrogen Energy*, 32(12), 1736–1741.
- Vanin, E. F. (1985). Processed pseudogenes: characteristics and evolution. *Annual Review of Genetics*, 19, 253–72.
- Vardar-Schara, G., Maeda, T., and Wood, T. K. (2008). Metabolically engineered bacteria for producing hydrogen via fermentation. *Microbial Biotechnology*, 1(2), 107–25.
- Venkata Mohan, S., Lalit Babu, V., and Sarma, P. N. (2007). Anaerobic biohydrogen production from dairy wastewater treatment in sequencing batch reactor (AnSBR): Effect of organic loading rate. *Enzyme and Microbial Technology*, 41(4), 506–515.
- Verger, D., Bullitt, E., Hultgren, S. J., and Waksman, G. (2007). Crystal structure of the P pilus rod subunit PapA. *PLoS Pathogens*, 3(5), 73.
- Verhaart, M. R. A., Bielen, A. A. M., van der Oost, J., Stams, A. J. M., and Kengen, S. W. M. (2010). Hydrogen production by hyperthermophilic and extremely thermophilic bacteria and archaea: mechanisms for reductant disposal. *Environmental Technology*, 31(8-9), 993–1003.
- Vignais, P. M., and Billoud, B. (2007). Occurrence, classification, and biological function of hydrogenases: an overview. *Chemical Reviews*, 107(10), 4206–4272.
- Vignais, P. M., Billoud, B., and Meyer, J. (2001). Classification and phylogeny of hydrogenases. *FEMS Microbiology Reviews*, 25(4), 455–501.
- Wang, B., Wan, W., and Wang, J. (2009). Effect of ammonia concentration on fermentative hydrogen production by mixed cultures. *Bioresource Technology*, 100(3), 1211–1213.
- Wang, G. M. Y., and Yu, H. Q. (2005). Response surface analysis to evaluate the influence of pH, temperature and substrate concentration on the acidogenesis of sucrose-rich wastewater. *Biochem Eng J*, 23, 175–184.
- Wang, J., and Wan, W. (2008a). Influence of Ni²⁺ concentration on biohydrogen production. *Bioresource Technology*, 99(18), 8864–8868.

- Wang, J., and Wan, W. (2008b). Effect of temperature on fermentative hydrogen production by mixed cultures. *International Journal of Hydrogen Energy*, 33(20), 5392–5397.
- Wang, W., Xie, L., Chen, J., Luo, G., and Zhou, Q. (2011). Biohydrogen and methane production by co-digestion of cassava stillage and excess sludge under thermophilic condition. *Bioresource Technology*, 102(4), 3833–3839.
- Watanabe, T., Totoki, Y., Toyoda, A., Kaneda, M., Kuramochi-Miyagawa, S., Obata, Y., Chiba, H., Kohara, Y., Kono, T., Nakano, T., Surani, M. A., and Sakaki, Y. H. (2008). Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature*, 453(7194), 539–543.
- Wawrzynczyk, J., Recktenwald, M., Norrlöw, O., and Dey, E. S. (2008). The function of cation-binding agents in the enzymatic treatment of municipal sludge. *Water Research*, 42(6-7), 1555–1562.
- Waygood, E. B., Mort, J. S., and Sanwal, B. D. (1976). The control of pyruvate kinase of *Escherichia coli*. Binding of substrate and allosteric effectors to the enzyme activated by fructose 1,6-bisphosphate. *Biochemistry*, 15(2), 277–282.
- Wei, J., Liu, Z.-T., and Zhang, X. (2010). Biohydrogen production from starch wastewater and application in fuel cell. *International Journal of Hydrogen Energy*, 35(7), 2949–2952.
- Wiechert, W. (2002). Modeling and simulation: tools for metabolic engineering. *Journal of Biotechnology*, 94(1), 37–63.
- Wu, K.-J., and Chang, J.-S. (2007). Batch and continuous fermentative production of hydrogen with anaerobic sludge entrapped in a composite polymeric matrix. *Process Biochemistry*, 42(2), 279–284.
- Xiao, B., and Liu, J. (2009). Biological hydrogen production from sterilized sewage sludge by anaerobic self-fermentation. *Journal of Hazardous Materials*, 168(1), 163–167.
- Xue, Y., Daly, A., Yngvadottir, B., Liu, M., Coop, G., Kim, Y., Sabeti, P., Chen, Y., Stalker, J., Huckle, E., Burton, J., Leonard, S., Rogers, J., and Tyler-Smith, C. (2006). Spread of an inactive form of caspase-12 in humans is due to recent positive selection. *American Journal of Human Genetics*, 78(4), 659–670.
- Y.-K., Oh, E.-H., Seola, J. R. Kimb, S., P. (2003). Fermentative biohydrogen production by a new chemoheterotrophic bacterium *Citrobacter* sp. Y19. *International Journal of Hydrogen Energy*, 28(12), 1353–1359.
- Yokoi, H., Maki, R., Hirose, J., and Hayashi, S. (2002). Microbial production of hydrogen from starch-manufacturing wastes. *Biomass and Bioenergy*, 22(5), 389–395.
- Yoshida, A., Nishimura, T., Kawaguchi, H., Inui, M., and Yukawa, H. (2005). Enhanced hydrogen production from formic acid by formate hydrogen lyase-overexpressing *Escherichia coli* strains. *Applied and Environmental Microbiology*, 71(11), 6762–6768.
- Yoshida, A., Nishimura, T., Kawaguchi, H., Inui, M., and Yukawa, H. (2007). Efficient induction of formate hydrogen lyase of aerobically grown *Escherichia coli* in a

- three-step biohydrogen production process. *Applied Microbiology and Biotechnology*, 74(4), 754–760.
- Yossan, S., O-Thong, S., and Prasertsan, P. (2012). Effect of initial pH, nutrients and temperature on hydrogen production from palm oil mill effluent using thermotolerant consortia and corresponding microbial communities. *International Journal of Hydrogen Energy*, 37(18), 13806–13814.
- Yuan, Z., Yang, H., Zhi, X., and Shen, J. (2010). Increased performance of continuous stirred tank reactor with calcium supplementation. *International Journal of Hydrogen Energy*, 35(7), 2622–2626.
- Zavialov, A. V, and Knight, S. D. (2007). A novel self-capping mechanism controls aggregation of periplasmic chaperone Caf1M. *Molecular Microbiology*, 64(1), 153–164.
- Zavialov, A. V, Tischenko, V. M., Fooks, L. J., Brandsdal, B. O., Aqvist, J., Zav'yalov, V. P., ... Knight, S. D. (2005). Resolving the energy paradox of chaperone/usher-mediated fibre assembly. *The Biochemical Journal*, 389(3), 685–694.
- Zeng, K., and Zhang, D. (2010). Recent progress in alkaline water electrolysis for hydrogen production and applications. *Progress in Energy and Combustion Science*, 36(3), 307–326.
- Zhai, Y., and Saier, M. H. (2002). The beta-barrel finder (BBF) program, allowing identification of outer membrane beta-barrel proteins encoded within prokaryotic genomes. *Protein Science*, 11(9), 2196–2207.
- Zhang, Z. D., Cayting, P., Weinstock, G., and Gerstein, M. (2008). Analysis of nuclear receptor pseudogenes in vertebrates: how the silent tell their stories. *Molecular Biology and Evolution*, 25(1), 131–143.
- Zhang, Z. D., Frankish, A., Hunt, T., Harrow, J., and Gerstein, M. (2010). Identification and analysis of unitary pseudogenes: historic and contemporary gene losses in humans and other primates. *Genome Biology*, 11(3), R26.
- Zhao, J.-F., Song, W.-L., Cheng, J., and Zhang, C.-X. (2009). Heterologous expression of a hydrogenase gene in *Enterobacter aerogenes* to enhance hydrogen gas production. *World Journal of Microbiology and Biotechnology*, 26(1), 177–181.
- Zhou, J., and Rudd, K. E. (2013). EcoGene 3.0. *Nucleic Acids Research*, 41, 613–624.
- Zhu, J., Sanborn, J. Z., Diekhans, M., Lowe, C. B., Pringle, T. H., and Haussler, D. (2007). Comparative genomics search for losses of long-established genes on the human lineage. *PLoS Computational Biology*, 3(12), 247.
- Zoetemeyer, R. J., Matthijsen, A. J. C. M., Cohen, A., and Boelhouwer, C. (1982). Product inhibition in the acid forming stage of the anaerobic digestion process. *Water Research*, 16(5), 633–639.