



**UNIVERSITI PUTRA MALAYSIA**

***TRANSCRIPTOMIC ANALYSIS OF MALAYSIAN RICE SEEDLINGS  
(Oryza sativa L. ssp. indica) IN EARLY RESPONSES TO SALT-SHOCK***

**NOR MUSTAIQAZAH BINTI MOHAMAD JURI**

**FBSB 2018 37**



**TRANSCRIPTOMIC ANALYSIS OF MALAYSIAN RICE SEEDLINGS (*Oryza sativa* L. ssp. *indica*) IN EARLY RESPONSES TO SALT-SHOCK**

By

**NOR MUSTAIQAZAH BINTI MOHAMAD JURI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in  
Fulfilment of the Requirements for the Degree of Master of Science**

**November 2017**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artworks, is copyright material of the Universiti Putra Malaysia unless otherwise stated. Use may be of the material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright© Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**TRANSCRIPTOMIC ANALYSIS OF MALAYSIAN RICE SEEDLINGS (*Oryza sativa* L. ssp. *indica*) IN EARLY RESPONSES TO SALT-SHOCK**

By

**NOR MUSTAIQAZAH BINTI MOHAMAD JURI**

**November 2017**

**Chairman : Noor Azmi Bin Shaharuddin, PhD**  
**Faculty : Biotechnology and Biomolecular Sciences**

Salinization of rice cultivation land is progressively enlarged thus negatively impair the world's rice bowl. Due to the polygenic nature and complexity of salinity tolerance mechanisms in plants, the development of new rice varieties with better adaptation to salinity has become a great challenge. Regarding this, transcriptomic profiling was seen as a promiseable technology for a holistic understanding of salinity tolerance mechanisms in rice. Here, by using Illumina HiSeq 2000 sequencing platform, transcriptomes of salt-tolerant Malaysian rice variety MR211 and salt-sensitive MR220 were analyzed after nine hours of severe salinity stress (12 dS/m) treatment, labeled as S211 and S220, respectively. After trimming, a total of 76,456,236 (S211) and 57,323,996 (S220) high-quality reads were obtained. The assembly of these reads resulting a total of 20,853 (S211) and 19,315 (S220) genes. Through comparative expression between both samples, 252 significant genes were differentially expressed and were dominated by variety-induced genes (n=235; 93.3%) with majority of them (n=221; 88%) were categorized as uniquely expressed in salt-tolerant MR211. Further pathway based analysis on the DEGs that were categorized as "uniquely" and "higher expressed" in S211 when compared to S220 had assigned them to 33 KEGG pathways with the highest number of DEGs were accounted in purine metabolism and thiamine metabolism pathways. The functional annotation of these group of DEGs also revealed the presence of regulatory genes such as transcription factors (TFs), protein kinases and protein phosphatases, as well as functional genes that involves in various adaptation mechanisms such as mechanical support, ROS-scavenging system, ion exclusion and intracellular compartmentalization thus suggest how this salt tolerant genotype (MR211) gains its salt adaptation trait. The expression accuracy and reproducibility of the 252 DEGs identified from the RNA-seq experiment were further verified through RT-PCR followed by qRT-PCR analysis. Nine genes were selected as the representative with 4 of them namely *FER2*, *Thaumatococcus*, *VI* and *UBC* were in line with data generated from the RNA-seq analysis. The other 2 (*MT* and *HOX16*) showed a contradict trend of expression as compared to the RNA-seq data, whereas the other three candidate genes (*PSII*, *SAPK6* and *PAO*) had exhibited a similar (no difference)

level of expressions between S211 and S220. Next, the incorporation of control (untreated) cDNA samples (C211 and C220) in the expression analyses had revealed the expression of the genes in untreated plants as compared to after being subjected to salt stress. The expression analyses had highlighted *UBC* and *SPK6* genes as the most responsive towards salinity stress in MR211 and MR220, respectively thus might represent their uniqueness in response to salinity stress.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Sarjana Sains

**ANALISIS TERHADAP RESPON AWAL TRANSKRIPTOM VARIATI PADI MALAYSIA (*Oryza sativa* L. ssp. *indica*) TERHADAP KEJUTAN KEMASINAN**

Oleh

**NOR MUSTAIQAZAH BINTI MOHAMAD JURI**

**November 2017**

**Pengerusi : Noor Azmi Bin Shaharuddin, PhD**  
**Fakulti : Bioteknologi dan Sains Biomolekul**

Proses pemasinan tanah berlaku dengan progresif di kawasan-kawasan penanaman padi dan telah menjejaskan bekalan beras dunia. Walaubagaimanapun, penghasilan varieti padi yang mempunyai sifat ketahanan yang tinggi terhadap kemasinan telah menjadi satu cabaran yang besar disebabkan oleh mekanisme ketahanan tumbuhan yang bersifat poligenik dan kompleks. Oleh itu, pemprofilan transkriptom dilihat sebagai satu teknologi baru yang menjanjikan pemahaman terhadap mekanisme pertahanan padi secara holistik. Dalam kajian ini, transkriptom dua varieti padi Malaysia, MR211 yang mempunyai ketahanan yang tinggi terhadap kemasinan dan MR220 yang sensitif terhadap kemasinan telah dianalisa selepas diberikan rawatan kemasinan yang tinggi (12 dS/m) selama 9 jam dan masing-masing dilabel sebagai S211 dan S220. Sejumlah 76,456,236 (S211) dan 57,323,996 (S220) bacaan berkualiti tinggi telah diperolehi, dan perhimpunan bacaan ini menghasilkan sejumlah 20,853 (S211) dan 19,315 (S220) gen. Melalui perbandingan pengekspresan yang dibuat di antara S211 dan S220, sebanyak 252 gen yang signifikan telah diekspres secara berbeza (DEGs). DEGs ini secara keseluruhannya telah didominasi oleh gen-gen yang dikspres secara spesifik pada satu varieti sahaja (n=235, 93.3%) dengan majoriti daripadanya diekspres secara unik hanya pada MR211. Analisis lanjutan yang dibuat terhadap DEGs yang diekspres secara "unik" dan "lebih tinggi" pada S211 apabila dibandingkan dengan S220 telah menunjukkan bahawa kumpulan gen ini terlibat di dalam 33 laluan KEGG, dengan bilangan gen yang tertinggi terlibat di laluan metabolisme purin dan tiamin. Anotasi fungsian kumpulan DEGs ini juga menunjukkan kehadiran gen-gen pengawalaturan seperti faktor transkripsi (TFs), protein kinase dan protein fosfatase, serta gen berfungsi yang terlibat dalam pelbagai mekanisme penyesuaian terhadap tekanan seperti sokongan mekanikal, sistem penghapus ROS, pengeksklusion ion dan pemetakan intrasel, yang dengan itu memberi petunjuk bagaimana pokok MR211 memperoleh ciri kerintangan terhadap tekanan kemasinan. Ketepatan dan kebolehlungan pengekspresan 252 DEGs yang diperolehi dari analisis RNA-seq disahkan melalui analisis RT-PCR dan qRT-PCR. Sembilan gen wakil dipilih dengan 4 daripadanya iaitu *FER2*, *Thaumatin*, *VI* dan *UBC* menunjukkan pengekspresan yang sejajar dengan

data yang dihasilkan dari analisis RNA-seq. *MT* dan *HOX16* walaupun bagaimanapun memperlihatkan trend pengekspresan yang bercanggah berbanding dengan data RNA-seq, manakala tiga lagi gen (*PSII*, *SAPK6* dan *PAO*) memperlihatkan tahap ekspresi yang sama (tidak ada perbezaan) antara S211 dan S220. Seterusnya, penggabungan sampel cDNA yang tidak dikenakan rawatan kemasinan (kawalan) (C211 dan C220) dalam analisis pengekspresan melalui RT-PCR dan qRT-PCR telah menonjolkan *UBC* dan *SPK6* masing-masing sebagai gen yang paling responsif terhadap tekanan kemasinan oleh MR211 dan MR220, sekali gus mewakili keunikan tindak balas kedua-dua varieti padi Malaysia ini terhadap tekanan kemasinan.



## ACKNOWLEDGEMENTS

All praises and thanks are to ALLAH SWT, the magnificent and merciful. I would like to thank Almighty ALLAH for His unfailing love, protection, guidance along the journey.

I avail myself of this opportunity to express my gratitude to my supervisor, Dr Noor Azmi Shaharuddin, chairman of my supervisory committee, for his dedicated effort, support, invaluable advice, intellectual guidance and encouragement in the conduct of my research and in the preparation of thesis. Grateful thanks are also to the members of the supervisory committee, for their endless guidance and advice throughout my study. I truly appreciate their patience, passionate, and encouragement within the process of research and until finished writing this thesis.

Besides that, I would like to thank for Ministry of Education (KPM) and Graduate Research Fellowship (GRF) for sponsored my postgraduate studies in Universiti Putra Malaysia. Other than that, a deepest gratitude to Fundamental Research Grant Scheme (FRGS) that providing the grant for my research work and Malaysian Agriculture Research and Development Institute (MARDI Serdang) for providing me with complete facilities in the laboratory and materials for this research.

Not to forget, an endless thank to my family for their love, moral and financial supports. Thank you for being supportive.



I certify that a Thesis Examination Committee has met on 23 November 2017 to conduct the final examination of Nor Mustaiqazah binti Mohamad Juri on her thesis entitled "Transcriptomic Analysis of Malaysian Rice Seedlings (*Oryza sativa* L. ssp. *indica*) in Early Responses to Salt-Shock" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

**Nik Mohd Afizan Nik Abdul Rahman, PhD**

Senior Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

**Mohd. Puad bin Abdullah, PhD**

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

**Nik Marzuki Sidik, PhD**

Professor

Universiti Malaysia Kelantan

Malaysia

(External Examiner)



**RUSLI HAJI ABDULLAH, PhD**

Professor and Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 27 September 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of Supervisory Committee were as follows:

**Noor Azmi Bin Shaharuddin, PhD**

Associate Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Chairman)

**Noor Baity Saidi, PhD**

Senior Lecturer  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Member)

**Yahya Bin Awang, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Maziah Mahmood, PhD**

Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by Graduate Student

I hereby confirm that:

- This thesis is my original work;
- Quotations, illustrations, and citations have been referenced;
- This thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- Intellectual property from thesis and copyright of thesis are fully owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- Written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books journals, modules, proceedings, popular writings, seminar papers, manuscript, posters, reports, lecture notes, learning modules, or any materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/ fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Nor Mustaiqazah Binti Mohamad Juri (GS37054)

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of Chairman of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of Member of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of Member of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of Member of  
Supervisory  
Committee: \_\_\_\_\_

## TABLE OF CONTENTS

		Page
	<b>ABSTRACT</b>	i
	<b>ABSTRAK</b>	iii
	<b>ACKNOWLEDGEMENTS</b>	v
	<b>APPROVAL</b>	vi
	<b>DECLARATION</b>	vii
	<b>LIST OF TABLES</b>	xiv
	<b>LIST OF FIGURES</b>	xv
	<b>LIST OF APPENDICES</b>	xix
	<b>LIST OF ABBREVIATIONS</b>	xx
	<b>CHAPTER</b>	
<b>1</b>	<b>INTRODUCTION</b>	<b>1</b>
	1.1 General Introduction	1
	1.2 Research Objectives	3
<b>2</b>	<b>LITERATURE REVIEW</b>	<b>4</b>
	2.1 <i>Oryza sativa</i> L.	4
	2.1.1 Taxon, Origin and Domestication	4
	2.1.2 Importance of Rice	5
	2.2 Rice Production in Malaysia	5
	2.2.1 Rice Cultivation Areas in Malaysia	6
	2.2.2 Malaysian Rice Varieties	6
	2.2.2.1 MR211	6
	2.2.2.2 MR220	7
	2.3 Salinity Stress	7
	2.3.1 Categories of Salt-affected Soils: Salinity, Saline and Sodic Soil	7
	2.3.2 Causes of Land Salinization	8
	2.3.2.1 Primary Salinization (Natural-Caused)	8
	2.3.2.2 Secondary Salinization (Human Activity)	9
	2.3.3 Progression of Rice-cultivation Land Salinization in Malaysia	9
	2.3.4 Effect of Salinity Stress on Plants	9
	2.3.5 Adaptation Strategies of Plants to Salinity Stress	10
	2.4 Application of Transcriptomics Study for Rice Improvement	11
<b>3</b>	<b>IDENTIFICATION OF EARLY RESPONSES SALT-SHOCK DIFFERENTIALLY EXPRESSED GENES (DEGs) IN MR211 AND MR220 VIA COMPARATIVE RNA-SEQ TRANSCRIPTOMIC ANALYSIS</b>	<b>13</b>
	3.1 Introduction	13
	3.2 Materials and Methods	13
	3.2.1 Supply and Selection of Plant Materials	13

3.2.2	Seeds Germination	14
3.2.3	Preparation of Nutrient Solution	15
	3.2.3.1 Stock Nutrient Solution Preparation	15
	3.2.3.2 Preparation of Half- and Full-strength Working Nutrient Solutions	15
3.2.4	Plants Growth Conditions	16
3.2.5	Salt Shock Treatments and Plants Sampling	16
3.2.6	Total RNA Extraction	16
3.2.7	Quantity and Quality Assessments of S211 and S220 Total RNA	17
3.2.8	cDNA Library Construction for Next-Generation Sequencing (NGS)	18
	3.2.8.1 Normalized and Pooled Libraries	22
3.2.9	Clustering of S211 and S211 Libraries	22
3.2.10	Paired-end Sequencing	22
3.2.11	Quality Check and Preprocessing of Raw Sequencing Data	23
3.2.12	Reads Mapping and Transcripts Assembly	23
3.2.13	Identification of Differentially Expressed Genes (DEGs)	24
3.2.14	Functional Annotation of DEGs	24
3.3	Results and Discussions	24
3.3.1	Variety Selection and Shoot Part as Plant Sample	24
3.3.2	Strength/Concentration of Salinity Treatment	25
3.3.3	Duration of Salinity Treatment	26
3.3.4	Quality Assessment of Total RNA	26
3.3.5	Quality Assessment of Sequenced Reads	29
3.3.6	Reads Mapping and Transcripts Assembly	30
3.3.7	Global Transcripts Abundance and Annotation	30
3.3.8	Identification of Differentially Expressed Genes (DEGs)	31
3.3.9	Gene Ontology (GO) and Pathway-based Analysis of DEGs	38
3.3.10	Involvement of DEGs in Salt-stress Adaptation Mechanism in MR211	42
	3.3.10.1 Genes Involved in Plants Mechanical Support via Cell Wall Organization	42
	3.3.10.2 Genes Involved in ROS-scavenging Mechanism	43
	3.3.10.3 Genes Involved in Salt Ion Exclusion and Intracellular Compartmentalization	45
	3.3.10.4 Transcription Factors (TFs)	46
	3.3.10.5 Protein Kinases and Phosphatase	46
	3.3.10.6 Photosynthetic Related Genes	49
	3.3.10.7 DNA Methylation	51
3.4	Conclusion	51

<b>4</b>	<b>EXPRESSION VALIDATION ANALYSIS OF THE SALT-SHOCK RESPONSIVE DEGs BY REVERSE TRANSCRIPTION PCR (RT-PCR) AND REAL-TIME QUANTITATIVE PCR (qRT-PCR)</b>	<b>52</b>
4.1	Introduction	52
4.2	Materials and Methods	52
4.2.1	Plant Samples Preparation	52
4.2.2	Extraction, Quality and Quantity Assessment of Total RNA	53
4.2.3	cDNA Synthesis	53
4.2.4	Selection of Candidate Genes and Primer Design for Expression Profiling via RT-PCR and qRT-PCR	53
4.2.5	Semi-quantitative RT-PCR	55
4.2.5.1	Standard RT-PCR Program and Components	55
4.2.5.2	Optimization of RT-PCR Cycling Parameter	55
4.2.5.3	Purification and Sequencing of PCR Product	55
4.2.5.4	Expression Profiling of Candidate Genes via RT-PCR and Data Analysis	56
4.2.6	Real-time PCR (qRT-PCR)	57
4.2.6.1	Standard qRT-PCR Program and Components	57
4.2.6.2	Primer Efficiency Test and Melting Curve Analysis	57
4.2.6.3	Validation of Reference Genes	57
4.2.6.4	Expression Profiling of Candidate Genes via qRT-PCR and Data Analysis	57
4.3	Results and Discussions	58
4.3.1	Quantity and Quality Assessment of Extracted Total RNA	58
4.3.2	Expression Profiling of DEGs via RT-PCR	59
4.3.2.1	Optimization of Cycling Parameter for DEGs Profiling via RT-PCR	59
4.3.2.2	Sequencing and Analyses of RT-PCR Product	60
4.3.2.3	qRT-PCR Primer Efficiency and Melting Curve Analysis	61
4.3.3	DEGs Expression Profiling via RT-PCR and qRT-PCR for RNA-Seq Data Validation	64
4.3.4	Profiling of Candidate Genes in Control and Salt-treated Samples	67
4.3.4.1	Expression Profiling of Candidate Genes in Salt-sensitive Variety, MR220	68
4.3.4.2	Expression Profiling of Candidate Genes in Salt-tolerant Variety,	69

	MR211	
4.4	Conclusion	71
<b>5</b>	<b>SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE STUDIES</b>	<b>72</b>
5.1	Summary and Conclusion	72
5.2	Recommendation for Future Research	73
	<b>REFERENCES</b>	<b>74</b>
	<b>APPENDICES</b>	<b>95</b>
	<b>BIODATA OF STUDENT</b>	<b>100</b>





## LIST OF TABLES

<b>Table</b>		<b>Page</b>
3.1	Constitution of Yoshida's stock solutions	15
3.2	Shoots samples labeling	16
3.3	The classification of several accessions of Malaysia rice varieties in response to salinity (sodium chloride) stress at different growth stages	25
3.4	Quality assessment of reads, mapping and assembly of the sequenced data	29
3.5	List of non-coding RNA (ncRNA) that differentially expressed in S211 and S220 transcriptome	35
3.6	List of DEGs transcripts that were involved in providing the mechanical support for salt-stress adaptation in MR211	43
3.7	List of DEGs transcripts that were involved in ROS-scavenging mechanism for salt-stress adaptation in MR211	44
3.8	List of DEGs transcripts that were involved in salt ion exclusion and intracellular compartmentalization	45
3.9	List of DEGs transcripts that were encoded as transcription factor	46
3.10	List of protein kinases that were found to be uniquely expressed ( $\text{Log}_2\text{FC} = +\text{INF}$ ) salt-treated MR211 (S211)	49
3.11	List of photosynthesis-related transcripts that might contribute in salinity stress adaptation mechanism in rice	50
3.12	List of DEGs transcripts that were encoded as retrotransposon	51
4.1	List of candidate genes to be validated via semi-quantitative RT-PCR and real-time PCR analysis	54
4.2	The primer sets used in RT-PCR and qRT-PCR	54
4.3	BLAST nucleotide (BLASTn) hit for the sequenced PCR product	61
4.4	Efficiency of the primer sequences of reference and candidate genes	62
4.5	Gene expression stability values (M) and coefficient of variance (CV) of the housekeeping genes	64

## LIST OF FIGURES

Figure		Page
2.1	Schematic illustration of the effect of salinity stress to the plant and their corresponding adaptive responses (Horie <i>et al.</i> , 2012). Bold boxes and arrows point out the damaging effects of salinity stress; dash boxes and arrows point out the adaptation mechanisms adapted by plants to cope with damaging effect of salinity stress	10
3.1	Three days old seedlings of (A) MR211 and (B) MR220 readied to be transferred into the hydroponic growth system	14
3.2	The setup of hydroponic cultivation of rice seedlings	15
3.3	Quality assessments of S211 and S220 total RNA. (A) The gel image of RNA in 1% agarose gel. The table showed the concentration and purity of RNA, measured by NanoDrop 1000 Spectrophotometer. (B) The image of RNA in 1.5% denatured RNA gel. M <sub>1</sub> : RNA ladder (Promega)	27
3.4	Electropherogram (left) and pseudo-gel image (right) generated by Agilent 2100 Bioanalyzer showed two distinct ribosomal peaks corresponding to 18S and 25S rRNA. RNA concentration and integrity (rRNA ratio 25S/18S and RIN) were also presented in the figure above	28
3.5	Bar graph of the division of annotated (known) and un-annotated (novel) global transcripts of S221 and S220. Blue bar represents genes that are annotated in RAP-DB and red bar represents genes with no hit (un-annotated) in the database	31
3.6	Volcano plots representing estimated fold changes (log <sub>2</sub> , x-axis) and statistically significant differences (-log <sub>10</sub> , p-value, y-axis) of genes in S211 and S220. Each dot represents a gene. Fold changes >0 indicated genes with higher expression in S211, whereas fold changes <0 indicated genes with lower expression in S211 compared to S220. Red dots: significant differentially expressed genes. Black dots: non-significant expressed genes. DEGs with FPKM=0 were not shown in this figure	32
3.7	Clustered heat map of the statistically significant expressed genes identified between S211 and S220. Left and right column represent S211 and S220, respectively. Each horizontal line refers to one gene, which made up of one or multiple number of transcripts. The color bar at the right side of the figure represent the expression level of the genes expression in FPKM values; those in the lightest orange have lower expression relative to the geometrical means, while dark orange indicates genes with	33

	higher expression	
3.8	The categorization of 252 DEGs into known (197, blue) and novel genes (55, red). The categorization was made based by referring to gene annotation from RAP-DB database	34
3.9	The pie chart represents the classification of known DEGs transcripts into their respective function based on RAP-DB database	34
3.10	Species distribution of the top BLAST hits for un-annotated DEGs transcripts. Hits shown for 101 transcripts. The most matches are to <i>Oryza sativa</i> , followed by other monocot plant species	36
3.11	The pie chart represents the classification of novel (un-annotated in RAP-DB) DEGs transcripts into their respective categories based on NCBI non-redundant (Nr) database. (A) Through BLASTX analyses of 101 DEG that are un-annotated in RAP-DB, 62% of them showed a significant homology with sequences in NCBI Nr database (blue), whereas no any significant hit showed for other 38% (red). (B) The 62% of DEGs were made up of unknown protein, hypothetical protein and well-annotated in other organism	37
3.12	The Gene Ontology (GO) terms demonstrated by DEGs transcripts. The GO terms were categorized into three main categories (Biological process: BP, molecular function: MF and cellular component: CC) which composed of 28 subcategories with GO classifications at level 2. Top 10 most presented categories and the number of transcripts belong to each category are shown	39
3.13	The KEGG classification demonstrated by the DEGs transcripts. (A) DEGs category that were uniquely and higher expressed in S211. (B) DEGs category that were not or very lowly expressed in S211	40
4.1	Quality assessment of total RNA on 1% agarose gel. Intact 25S and 18S total RNA bands were observed in all samples, indicating a good integrity of the total RNA. C1, C2 and C3 are the three biological replicates of untreated samples, whereas S1, S2 and S3 are the three biological replicated of salt-treated samples of MR211 and MR220, respectively	59
4.2	Exponential ranges of candidate and housekeeping genes amplification via RT-PCR. The density (%) of band in each cycle was quantified by ImageJ and was presented in bar graph form. The red arrows indicated the selected number of cycle used in the sqRT-PCR expression profile analysis	60

- 4.3 Melting curve of (A) representative DEGs (*PAO*, *UBC*, *HOX16*, *Thaumatococcus* and *MT*) and (B) housekeeping genes (*EP* and *HNR*). The red arrow indicates single-peak of melt curve whereas the blue arrow indicates negative control (NTC) without any amplification detected 63
- 4.4 Validation of the expression of nine representative DEGs in S211 and S220 cDNA via semi-quantitative RT-PCR. (A) The PCR products were run in 2.0% gel electrophoresis. (B) Analysis of quantified band intensity. Data are expressed relative to the S220 levels of expression and was normalized to *HNR* (housekeeping gene). Blue bar represent the expression of target gene in S220 cDNA sample, whereas the red bar represent the expression of target gene in S211 cDNA. Asterisks (\*) denote genes with consistent expression with the RNA-seq data 66
- 4.5 Quantitative real-time PCR analyses of *PAO* and *UBC* genes in S211 and S220. *HNR* and *EP* genes were used as reference genes, whereas S220 was used as the control sample. Blue bar represent the expression of target gene in S220 whereas the red bar represent the expression of target gene in S211 66
- 4.6 Validation of the expression of nine representative DEGs in C220 and S220 cDNA via semi-quantitative RT-PCR. (A) The image of PCR products analyzed in 2.0% gel electrophoresis. (B) Analysis of quantified band intensity. Data are expressed relative to the C220 levels of expression and was normalized to *HNR* (reference gene). White bar represent the expression of target gene in C220 cDNA template whereas the blue bar represent the expression of target gene in S220 cDNA. Asterisks (\*) denote genes with significant fold of expression between C220 and S220 68
- 4.7 Quantitative qRT-PCR analyses of *PAO* and *UBC* in C220 and S220. *HNR* and *EP* were used as reference genes, whereas C220 was used as the control sample. White bar represent the expression of target gene in C220 cDNA template whereas the blue bar represent the expression of target gene in S220 cDNA 69
- 4.8 Validation of the expression of nine representative DEGs in C211 and S211 cDNA via semi-quantitative RT-PCR. (A) The image of PCR products analyzed in 2.0% gel electrophoresis. (B) Analysis of quantified band intensity. Data are expressed relative to the C211 levels of expression and was normalized to *HNR* (reference gene). White bar represent the expression of target gene in C211 cDNA template whereas the red bar represent the expression of target gene in S211 cDNA. Asterisks (\*) denote genes with significant fold of expression between C211 and S211 70
- 4.9 Quantitative real-time PCR analyses of *PAO* and *UBC* in C211 70

and S211. *HNR* and *EP* were used as internal control, whereas C211 was used as the control sample. White bar represent the expression of target gene in C211 cDNA template whereas the red bar represent the expression of target gene in S211 cDNA



## LIST OF APPENDICES

<b>Appendix</b>		<b>Page</b>
A	Nanodrop reading for MR211 and MR220 RNA samples	95
B	Generation of standard curve to assess reaction optimization. A standard curve was generated using a 10-fold dilution of the template cDNA	96
C	The analysis of band intensity of candidate genes in salt-treated MR211 (S211) and salt-treated MR220 (S220) sample cDNA	97
D	The analysis of band intensity of candidate genes in control (C220) and salt-treated (S220) sample of MR220 cDNA	98
E	The analysis of band intensity of candidate genes in control (C211) and salt-treated (S211) sample of MR211 cDNA	99

## LIST OF ABBREVIATIONS

NGS	Next Generation Sequencing
DNA	Deoxyribonucleic Acid
RNA	Ribonucleic Acid
PCR	Polymerase Chain Reaction
RT-PCR	Reverse Transcription Polymerase Chain Reaction
qRT-PCR	Real-time Reverse Transcription Polymerase Chain Reaction
$\epsilon$	Extinction Coefficient
EDTA	Ethylenediamine Tetraacetic Acid
TAE	Tris-Acetate-EDTA
Tris	Tris(Hydroxymethyl) Aminomethane
$\times g$	Gravitational Constant
kDa	KiloDalton
NCBI	National Centre for Biotechnology Information
nt	Nucleotide
QC	Quality Control



## CHAPTER 1

### INTRODUCTION

#### 1.1 General Introduction

Rice (*Oryza sativa* L.) is a critical crop for global food security. It is ranked as one of the most important crops of the world, primarily in Asian and certain part of Latin America and African countries (Muthayya *et al.*, 2014). It feeds almost half of world population which represented by more than 3.5 billion of peoples, as well as an important source of employment and incomes for the rural farmers. Over the years, a massive growth of world population has led to the increment of global rice demand. It has been estimated that an additional of 114 million tons of rice is required in order to meet global rice demand in 2035 (FAO, 2016; Mohanty *et al.*, 2013).

Unfortunately, rice is prone to a wide range of environmental constraint. A stagnation of rice yield in many Asian countries has been recorded due to various abiotic stress factors (Redfern *et al.*, 2012). Among them, salt stress has been represented as a significant threat for rice growth and development (Munns 2011; Kumar and Khare, 2016), accounted in 30% to 50% (Islam *et al.*, 2007; Joseph and Mohanan, 2013) or even higher losses of rice yield (Michael *et al.*, 2004; Zeng and Shannon, 2000).

Soil is categorized as saline once its Electrical Conductivity (EC) reading reached 4 dS/m or higher, with pH less than 8 and Exchangeable Sodium Percentage (ESP) less than 15 (Allotey *et al.*, 2008). A continuous salinization process is naturally occurring through release of soluble salts from parental rocks and deposition of oceanic salts carried in wind and rain in coastal areas. In agricultural land, application of chemical fertilizer and irrigation of brackish water itself are the common source of salinization (FAO, 2005; Petronia 2011; Munns and Tester, 2008). More than that, the effect of climate change such as increasing of annual temperature, decreasing of rainfall and rise in global sea level have led to a tremendous effect of soil salinization (Brinkman 1980; Hakim *et al.*, 2013).

In a scientific assessment conducted by Yuen & Kong in 2009, the coastlines of Southeast Asia that made up of the regions which serve as 'world rice bowl' are highly vulnerable to the effects of climate change. In Malaysia, Indonesia and Vietnam for example, thousands square kilometer ( $\text{km}^2$ ) loss of land were expected due to the 30-50 cm increment of sea level (Wassmann *et al.*, 2004; UNEP, 2006; IPCC, 2001). For granary lands which especially located on delta and coastal areas (Nguyen *et al.*, 2014; Brinkman 1980), rising of sea level will lead into sea water intrusion and submersion of the fertile fields under sea thus makes it unsuitable for rice planting. Depending on its concentration and duration, salinity stress invokes various changes in physiological and



metabolic events in plants, in which, ultimately inhibit their production and survival (Rahnama *et al.*, 2010; Rozema & Flowers, 2008).

The destructive effect of soil salinity is caused by two major factors known as osmotic and ionic effects (Kosová *et al.*, 2013). Osmotic effect occurs when high levels of salt ion in soil around plant root are triggered into the decreasing of cellular osmotic potential and ultimately lead to cell dehydration. Meanwhile ionic effect occurs when the salt ion penetrate and accumulate in cell cytoplasm via plasma membrane (James *et al.*, 2011; Rahnama *et al.*, 2010; Kosová *et al.*, 2013). These effects will result in the interruption of intracellular enzyme activities, disruption of membrane structures and functions, nutrient imbalance, accumulation of reactive oxygen species (ROS), decreased photosynthetic activity, decrease in stomatal aperture and reduction of cell division and expansion (Munns, 2002; Rahnama *et al.*, 2010; Zhu, 2007).

As in other plant species, responses of rice plant to salt stress vary with varieties and growth stages. It has been stated that rice is very sensitive to salinity throughout young seedling growth stage (Heenan *et al.*, 1988; Lutts *et al.*, 1996). Based on standard evaluating score (SES) in rating the visual symptoms of salt toxicity (IRRI, 2002; Gregorio *et al.*, 1997), varieties of rice can be differentiated as highly tolerant, tolerant, moderately tolerant and susceptible. As being applied by Hakim *et al.*, (2010), the screening of eight Malaysian rice varieties using the SES had identified MR211 as the most tolerant while MR220 as the most susceptible variety among others varieties.

Salinity tolerance is a quantitative trait controlled by many genes that involve in different pathways. Therefore, a full understanding on the molecular responses of rice plants to varying conditions and identification of genes that involved in salinity stress response is crucial to serve as a foundation in developing rice with better adaptation to salinity. In this prospect, identification of salt stress related genes is a promising approach in crop improvement program through development of rice varieties with higher harvestable yield during environmental stresses (Amudha and Balalubramani, 2011). Although the conventional breeding has been playing a crucial role in rice improvement, it is somewhat a slow process as it is time consuming (Miah *et al.*, 2012).

Plant biotechnology through genetic engineering and molecular breeding approaches offers much rapid development in a crop improvement program under stressful environments. Recently, the attention on utilizing modern high-throughput genetic approach such as transcriptomics and proteomics has extensively grown. As plants vary in their response to stresses, the application of these “omicss” method in comparative studies between related plant species enables the identification of various functional genes, their transcript and protein products, including the novel ones that are responsible for stress responses and adaptation (Kosová *et al.*, 2013).

Here in present work, using Illumina RNA-Seq method, transcriptomes of two contrasting Malaysian rice varieties, salt-tolerant MR211 and salt-sensitive MR220 in responses to salt stress treatment were analysed. Transcriptomes analysis using Tuxedo

package enabled the identification of significant genes that were differentially expressed between both varieties where further analyses had identified candidate genes related to salt tolerant.

The data generated from this study will serve as an invaluable genomic reference to further our knowledge on the molecular and cellular events that specifically occur in rice seedlings during their early response to severe salinity stress. Not only constricted as the candidate genes for rice improvements through genetic engineering, the potential salt-tolerant genes can also be utilized as DNA markers to fasten the selection process in conventional breeding.

## **1.2 Research objectives**

The main aim of this project is to study salt-shock adaptation pathways exhibited by Malaysian rice varieties, MR211 and MR220 in response to severe salinity stress treatment. Therefore the specific objectives of this work were:

1. To identify the early responses salt-shock differentially expressed genes (DEGs) of MR211 and MR220 seedlings via comparative RNA-seq transcriptomic analysis.
2. To validate the expression profile of salt-shock DEGs in MR211 and MR220 identified from the RNA-seq analysis through semi-quantitative reverse transcription-PCR and quantitative real-time PCR analyses.

## REFERENCES

- Akinbile, C.O., Abd El-Latif K.M., Abdullah R. and Yusoff, M.S. 2011. Rice production and water use efficiency for self-sufficiency in Malaysia: a review. *Trends in Applied Sciences Research*. 6 (10): 1127–1140.
- Alamillo, J.M., Díaz-Leal, J.L., Sánchez-Moran, M.V. and Pineda, M. 2010. Molecular analysis of ureide accumulation under drought stress in *Phaseolus vulgaris* L. *Plant Cell Environment*. 33:1828–1837.
- Albacete, A., Cantero-Navarro, E., Großkinsky, D.K. et al. 2015. Ectopic overexpression of the cell wall invertase gene CIN1 leads to dehydration avoidance in tomato. *Journal of Experimental Botany*. 66:863–878.
- Alias, I., Othman, O., Mohamad, H., Saad, A., Habibuddin, H., Abd Rahman, A.B. and Azlan, S. 2005. New rice variety MR220. *Buletin Teknologi Tanaman*. 2: 7-13.
- Allotey DFK, Asiamah RD, Dedzoe CD, Nyamekye AL. 2008. Physico-Chemical Properties of Three Salt-Affected Soils in the Lower Volta Basin and Management Strategies for their Sustainable Utilization. *West African Journal of Applied Ecology*. 12:163-182.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. 1990. Basic local alignment search tool. *Journal of Molecular Biology*. 215 (3): 403-410.
- Altschul, S.F., Madden, L.T., Schffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. 1997. Gapped blast and psi-blast: a new generation of protein database search programs. *Nucleic Acids Research*. 25(17): 3389-3402.
- Alvarado, M.C., Zsigmond, L.M., Kovacs, I., Cseplo, A., Koncz, C. and Szabados, L.M. 2004. Gene trapping with firefly luciferase in *Arabidopsis*. Tagging of stress-responsive genes. *Plant Physiology*. 134:18-27.
- Ambawat, S., Sharma, P., Yadav, N. R., & Yadav, R. C. 2013. MYB transcription factor genes as regulators for plant responses: an overview. *Physiology and Molecular Biology of Plants*. 19(3): 307–321.
- Amudha, J. and Balalubramani, G. 2011. Recent molecular advances to combat abiotic stress tolerance in crop plants. *Biotechnology, Molecular and Biological Review*. 6(2): 31-58.
- Anders, S. and Huber, W. 2010. Differential expression analysis for sequence count data. *Genome Biology*. 11:R106.
- Andersen, C.L., Jensen, J.L. and Ørntoft, T.F. 2004. Normalization of real-time quantitative reverse transcription-PCR data: A model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Research*. 64:5245–5250.
- André, S., Siebert, H.C., Nishiguchi, M., Tazaki, K. and Gabius, H.J. 2005. Evidence for lectin activity of a plant receptor-like protein kinase by application of neoglycoproteins and bioinformatic algorithms. *Biochimica et Biophysica Acta*. 1725(2):222-32.
- Andrea, K., Werner, and Witte, C.P. 2011. The biochemistry of nitrogen mobilization: purine ring catabolism. *Trends in Plant Science*. 16(7): 381-387.
- Antiabong, J.F., Ngoepe, M.G. and Abechi, A.S. 2016. Semi-quantitative digital analysis of polymerase chain reaction electrophoresis gel: Potential applications in low-income veterinary laboratories. *Veterinary World*. 9(9): 935-939.
- Apse, M.P. and Blumwald, E. 2002. Engineering salt tolerance in plants. *Current Opinion in Biotechnology*. 13: 146-150.

- Asai, T., Tena, G., Plotnikova, J., Willmann, M.R., Chiu, W.L., Gomez-Gomez, L., Boller, T., Ausubel, F.M. and Sheen, J. 2002. MAP kinase signaling cascade in *Arabidopsis* innate immunity. *Nature*. 415:977–983.
- Auer, P.L. and Doerge, R.W. 2011. A Two-stage Poisson model for testing RNA-Seq data. *Statistical Applications in Genetics and Molecular Biology*. 10:1–26.
- Azizi, P., Rafii, M.Y., Mahmood, M., Abdullah, S.N.A., Hanafi, M.M., Latif, M.A., Sahebi, M. and Ashkani, S. 2017. Evaluation of RNA extraction methods in rice and their application in expression analysis of resistance genes against Magnaporthe oryzae. *Biotechnology & Biotechnological Equipment*. 31(1): 75-84.
- Azman, E.A., Jusop, S., Ishak, C.F. and Ismail, R. 2014. Increasing Rice Production Using Different Lime Sources on an Acid Sulphate Soil in Merbok, Malaysia. *Pertanika Journal of Tropical Agricultural Science*. 37 (2): 223 – 247.
- Baloglu, M.C. and Patir, M.G. 2014. Molecular characterization, 3D model analysis, and expression pattern of the CmUBC gene encoding the melon ubiquitin-conjugating enzyme under drought and salt stress conditions. *Biochemical Genetics*. 52: 90–105.
- Barre, A., Herve, C., Lescure, B. and Rouge, P. 2002. Lectin receptor kinases in plants. *Critical Reviews in Plant Sciences*. 21: 379–399.
- Basir, E.K. 2016. Bad Weather During King Tide Phenomenon Can Cause Severe Flooding – Experts. *BERNAMA*. Retrieved at 20 June 2017. <http://www.bernama.com/bernama/v8/ge/newsperspectives.php?id=1284607>.
- Baxter, C.J., Redestig, H., Schauer, N., Reipsilber, D., Patil, K.R., Nielsen, J., Selbig, J., Liu, J., Fernie, A.R. and Sweetlove, L.J. 2007. The metabolic response of heterotrophic *Arabidopsis* cells to oxidative stress. *Plant Physiology*. 143:312–325.
- Begley, T.P., Kinsland, C. and Strauss, E. 2001. The biosynthesis of coenzyme A in bacteria. *Vitamins and Hormones*. 61: 157–171.
- BERNAS. 2016. <http://www.bernas.com.my/index.php/127-ricepedia/rice-type-in-malaysia/local-white-rice>.
- Bettendorff, L. and Wins, P. 2009. Thiamin diphosphate in biological chemistry: new aspects of thiamin metabolism, especially triphosphate derivatives acting other than as cofactors. *FEBS Journal*. 276:2917-2925.
- Bird, A. 2002. DNA methylation patterns and epigenetic memory. *Genes & Development*. 16: 6–21.
- Bokszczanin, K.L., Krezdorn, N., Fragkostefanakis, S., Müller, S., Rycak, L., Chen, Y., Hoffmeier, K., Kreutz, J., Paupière, M.J., Chaturvedi, P., Iannacone, R., Müller, F., Bostan, H., Chiusano, M.L., Scharf, K.D., Rotter, B., Schleiff, E. and Winter, P. 2015. Identification of novel small ncRNAs in pollen of tomato. *BMC Genomics*. 16:714.
- Bolger, A.M., Lohse, M., Usadel, B. 2014. Trimmomatic: A flexible trimmer for Illumina Sequence Data. *Bioinformatics*. 30(15): 2114-20.
- Bouwmeester, K. and Govers, F. 2009. Arabidopsis L-type lectin receptor kinases: phylogeny, classification, and expression profiles. *Journal of Experimental Botany*. 60 (15): 4383-4396.
- Briat, J.F., Duc, C., Ravet, K., Gaymard, F. 2010. Ferritins and iron storage in plants. *Biochim Biophys Acta*. 1800:806–14.
- Brinkman. 1980. Saline and sodic soils, In: Land reclamation and water management. International Institute for Land Reclamation and Improvement (ILRI), Wageningen, the Netherlands. pp 62-68.

- Brychkova G, Alikulov Z, Fluhr R, Sagi M. 2008. A critical role for ureides in dark and senescence-induced purine remobilization is unmasked in the *Atxdh1 Arabidopsis* mutant. *The Plant Journal*. 54(3):496–509.
- Bustin, S.A. and Nolan, T. 2004. Pitfalls of quantitative real-time reverse-transcription polymerase chain reaction. *Journal of Biomolecular Techniques*. 15:155–166.
- Caldana, C., Scheible, W.R. and Mueller-Roeber, B. 2007. A quantitative RT-PCR platform for high-throughput expression profiling of 2500 rice transcription factors. *Plant Methods*. 3: 7–15
- Chamhuri, S., A. Mahmudul, M. Wahid and A.Q. Al-Amin. 2009. Climate Change, Agricultural Sustainability, Food Security and Poverty in Malaysia. *International Review of Business Research Paper*. 5(6): 309-321.
- Chan, Y.S. 2017. Our kids must learn to swim. *The Star Malaysia*. Retrieved in 23 June 2017. <http://www.pressreader.com/malaysia/the-star-malaysia/20170320/282063391784420>
- Chang H, Siegel BZ, Siegel SM. 1984. Salinity induced changes in isoperoxidase in taro, *Colocasia esculenta*. *Phytochemistry* 23: 233–235.
- Chaudhary, S., Chaudhary, P.S. and Vaishnani, T.A. 2016. Small RNA Extraction using Fractionation Approach and Library Preparation for NGS Platform. *Journal of Advanced Research in Biotechnology*. 1(1):7.
- Chen, L., Zhong, H.Y. and Kuang, J.F. 2011. Validation of reference genes for RT-qPCR studies of gene expression in banana fruit under different experimental conditions. *Planta*. 234: 377–390.
- Chun, J., Li, F.-S., Ma, Y., Wang, S.-H. and Chen, F. 2014. Cloning and characterization of a SnRK2 gene from *Jatropha curcas* L. *Genetics and Molecular Research*. 13 (4): 10958-10975.
- Chung, E., Cho, C.W., So, H.A., Kang, J.S., Chung, Y.S. and Lee, J.H. 2013. Overexpression of VrUBC1, a Mung Bean E2 Ubiquitin-Conjugating Enzyme, Enhances Osmotic Stress Tolerance in *Arabidopsis*. *PLOS ONE*. 8: e66056. PMID: 23824688.
- Clement M, Lambert A, Herouart D, Boncompagni E. Identification of new up-regulated genes under drought stress in soybean nodules. *Gene*. 2008;426:15–22.
- Conley, T.R., Peng, H.P. and Shih, M.C. 1999. Mutations affecting induction of glycolytic and fermentative genes during germination and environmental stress in *Arabidopsis*. *Plant Physiology* 119:599–607.
- Cotsaftis, O., Plett, D., Johnson, A.A., Walia, H., Wilson, C., Ismail, A.M., Close, T.J., Tester, M. and Baumann, U. 2011. Root-specific transcript profiling of contrasting rice genotypes in response to salinity stress. *Mol Plant*. 4(1):25-41.
- Coué, I., Sulmon, C., Gouesbet, G., El Amrani, A. 2006. Involvement of soluble sugars in reactive oxygen species balance and responses to oxidative stress in plants. *Journal of Experimental Botany*. 57:449–459.
- Cramer GR, Epstein E, Läuchli A. 1989. Na-Ca interactions in barley seedlings: relationship to ion transport and growth. *Plant, cell and environment*. 12: 551-558.
- Crouzet, J., Roland, J. and Peeters, E. 2013. *Plant Molecular Biology*. 82: 181.
- Cui, F., Liu, L., Zhao, Q., Zhang, Z., Li, Q., Lin, B. 2012. *Arabidopsis* ubiquitin conjugase UBC32 is an ERAD component that functions in brassinosteroid-mediated salt stress tolerance. *Plant Cell*. 24: 233–244.
- Cuin TA, Miller AJ, Laurie SA, Leigh RA. 2003. Potassium activities in cell compartments of salt-grown barley leaves. *Journal of experimental botany*. 54: 657-661.



- De Diego, N., Sampedro, M.C., Barrio, R.J., Saiz-Fernandez, I., Moncalean, P. and Lacuesta, M. 2013. Solute accumulation and elastic modulus changes in six *radiata pine* breeds exposed to drought. *Tree Physiology*. 33:69–80.
- De Leon, T. B., Linscombe, S., Gregorio, G., & Subudhi, P. K. 2015. Genetic variation in Southern USA rice genotypes for seedling salinity tolerance. *Frontiers in Plant Science*. 6: 374.
- De Lorenzo, L., Merchan, F., Laporte, P., Thompson, R., Clarke, J., Sousa, C. and Crespi, M. 2009. A Novel Plant Leucine-Rich Repeat Receptor Kinase Regulates the Response of *Medicago truncatula* Roots to Salt Stress. *Plant Cell*. 21:668-680.
- Deak, M., Horvath, G.V., Davletova, S., Torok, K., Sass, L., Vass, I., Barna, B., Kiraly, Z. and Dudits, D. 1999. Plants ectopically expressing the iron-binding protein, ferritin, are tolerant to oxidative damage and pathogens. *Nature Biotechnology*. 17: 192–196.
- Deinlein, U., Stephan, A.B., Horie, T., Luo, W., Xu, G. and Schroeder, J.I. 2014. Plant salt-tolerance mechanisms. *Trends in plant science*. 19(6): 371-379.
- Deivanai, S.R., Xavier, V., Vinod, K.T. and Lim, O.F. 2011. Role of Exogenous Proline in Ameliorating Salt Stress at Early Stage in Two Rice Cultivars. *Journal of Stress Physiology & Biochemistry*. 7(4):157-174.
- Deng, K., Wang, Q., Zeng, J., Guo, X., Zhao, X., Tang, D., and Liu, X. 2009. A lectin receptor kinase positively regulates ABA response during seed germination and is involved in salt and osmotic stress response. *Journal of Plant Biology*. 52: 493–500.
- Department of Agriculture Peninsular Malaysia. 2014. Paddy Statistics of Malaysia 2014. [http://www.doa.gov.my/index/resources/aktiviti\\_sumber/sumber\\_awam/maklumat\\_pertanian/perangkaan\\_tanaman/perangkaan\\_padi\\_2014.pdf](http://www.doa.gov.my/index/resources/aktiviti_sumber/sumber_awam/maklumat_pertanian/perangkaan_tanaman/perangkaan_padi_2014.pdf)
- Dievart, A., Perin, C., Hirsch, J., Bettembourg, M., Lanau, N. and Artus, F. 2016. The phenome analysis of mutant alleles in leucine-rich repeat receptor-like kinase genes in rice reveals new potential targets for stress tolerant cereals. *Plant Science*. 242:240–249.
- Dreher, K., and Callis, J. 2007. Ubiquitin, hormones and biotic stress in plants. *Annals of Botany*. 99:787–822.
- Drew, M.C. 1997. Oxygen deficiency and root metabolism: injury and acclimation under hypoxia and anoxia. *Annual Review of Plant Physiology and Plant Molecular Biology*. 48:223–250.
- Duque AS, Almeida AM de, Silva AB da, Silva JM da, Farinha AP, Santos D, Fevereiro P, Araújo S de S. 2013. Abiotic Stress Responses in Plants: Unraveling the Complexity of Genes and Networks to Survive. *InTech*. <http://dx.doi.org/10.5772/52779>.
- FAO STAT online database. 2016. <http://www.fao.org/about/meetings/agroecology-symposium-china/en/>. Retrieved on March 2017.
- FAO webpage <http://www.fao.org/soils-portal/soil-management/management-of-some-problem-soils/salt-affected-soils/more-information-on-salt-affected-soils/en/>. Accessed on July 21, 2017.
- FAO, Food and Agricultural Organization. 2005. Salt-affected soils from sea water intrusion: Strategies for rehabilitation and management. Report of the regional workshop. Bangkok, Thailand, 62 pp. Accessed June 29, 2017.
- FAO, Food and Agricultural Organization. 2012. <http://www.fao.org/publications/sofi/2012/en/>. Accessed June 29, 2017.

- FAO, Food and Agriculture Organization. 2004. Rice is Life. Rome: FAO: Food and Agriculture Organization. June 29, 2013. <http://www.fao.org/newsroom/en/focus/200436887/index.html>.
- Fei, G., YiJun, Z., LingYun, H., DaCheng, H.E. and GenFa, Z. 2008. Proteomic analysis of long-term salinity stress responsive proteins in *Thellungiella halophila* leaves. *Chinese Science Bulletin*. 53: 3530–3537.
- Ferreira, L.J., Azevedo, V., Maroco, J., Oliveira, M.M. and Santos, A.P. 2015. Salt Tolerant and Sensitive Rice Varieties Display Differential Methylome Flexibility under Salt Stress. *PLoS ONE*. <http://dx.doi.org/10.1371/journal.pone.0124060>.
- Fleige, S. and Pfaffl, M.W. 2006. RNA integrity and the effect on the real-time qRT-PCR performance. *Molecular Aspects of Medicine*. 27:126–139.
- Flowers, T.J., Troke, P.F. and Yeo, A.R. 1977. The Mechanism of Salt Tolerance in Halophytes. *Annual Review of Plant Physiology*. 28: 89–121.
- Fox TC, Guerinot ML. 1998. Molecular biology of cation transport in plants. *Annu. Rev. Plant Physiol. Plant Molecular Biology*. 49: 669–696.
- Gallego Romero, I., Pai, A.A., Tung, J. and Gilad, Y. 2014. RNA-seq: impact of RNA degradation on transcript quantification. *BMC Biology*. 12(1):42.
- Garcia-Hernandez, M., Berardini, T.Z., Chen, G., Crist, D., Doyle, A., Huala, E., Knee, E., Lambrecht, M., Miller, N., Mueller, L.A. 2002. TAIR: a resource for integrated *Arabidopsis* data. *Functional & Integrative Genomics*. 2: 239–253.
- Garg R, Shankar R, Thakkar B, Kudapa H, Krishnamurthy L, Mantri N, Varshney RK, Bhatia S, Jain M. 2016. Transcriptome analyses reveal genotype- and developmental stage-specific molecular responses to drought and salinity stresses in chickpea. *Scientific Reports* 6:19228.
- Garg, R., Sahoo, A. and Tyagi, A.K. 2010. Validation of internal control genes for quantitative gene expression studies in chickpea (*Cicer arietinum* L.). *Biochemical and Biophysical Research Communications*. 396: 283–288.
- Ge, X., Chen, W., Song, S., Wang, W., Hu, S., Yu, J. 2008. Transcriptomic profiling of mature embryo from an elite super-hybrid rice LYP9 and its parental lines. *BMC Plant Biology*. 8:114.
- Geng, Y., Wu, R., Wee, C.W., Xie, F., Wei, X., Chan, P.M., Tham, C., Duan, L., Dinneny, J.R. 2013. A spatio-temporal understanding of growth regulation during the salt stress response in *Arabidopsis*. *Plant Cell*. 25: 2132–2154.
- GIEWS and FAO. 2016. <http://www.fao.org/giews/countrybrief/country.jsp?code=MYS> Reference Date: 20 December 2016.
- Glenn, E., Miyamoto, M., Moore, D., Brown, J. J., Thompson, T. L. and Brown, P. 1997. Water requirements for cultivating *Salicornia bigelovii* Torr. with seawater on sand in a coastal desert environment. *Journal of Arid Environments*. 36:711–730.
- Gracey, A.Y. and Cossins, A.R. 2003. Application of microarray technology in environmental and comparative physiology. *Annual Review of Physiology*. 65: 231–259.
- Gregorio GB, Senadhira D, Mendoza RD. 1997. Screening rice for salinity tolerance. *Plant Breeding, Genetics, and Biochemistry Division, IRRI*. IRRI discussion paper series No. 22.
- GRiSP (Global Rice Science Partnership). 2013. Rice almanac, 4th edition. Los Baños (Philippines): *International Rice Research Institute*. 283 p.
- Gururani, M. A., Venkatesh, J. and Tran, L.S. P. 2015. Regulation of photosynthesis during abiotic stress-induced photoinhibition. *Mol. Plant*. 8: 1304–1320.

- Hai, N.X., Trang, T.N. and Long, N.H. 2014. Paddy salinization process in Tien Hai District, Thai Binh province under sea level rise impacts. *ARPN Journal of Agricultural and Biological Science*. 9:11.
- Hakim M A, Juraimi AS, Razi Ismail M, Hanafi MM, Selamat A. 2013. A survey on weed diversity in coastal rice fields of Sebarang Perak in peninsular Malaysia. *Journal of Animal and Plant Sciences*. 23(2): 534–542.
- Hakim, M.A., Juraimi, A.S., Begum, M., Hanafi, M.M., Ismail, M.R. and Selamat, A. 2010. Effect of salt stress on germination and early seedling growth of rice (*Oryza sativa* L.). *African Journal of Biotechnology*. 9(13): 1911-1918.
- Hakim, M.A., Juraimi, A.S., Hanafi, M.M., Ali, E., Ismail, M.R., Selamat, A. and Karim, S.M. 2014. Effect of salt stress on morpho-physiology, vegetative growth and yield of rice. *Journal of Environment Biology*. 35(2):317-26.
- Hamid, N.H. 2016. Sawah padi dimasuki air masin. *SINAR*. <http://www.sinarharian.com.my/ultras/liga-malaysia/sawah-padi-dimasuki-air-masin-1.568470>. Retrieved on March 2017.
- Hartmann, M.D., Mendler, C.T., Bassler, J., Karamichali, I., Ridderbusch, O., Lupas, A.N., and Hernandez Alvarez, B. 2016.  $\alpha/\beta$  coiled coils. *eLife* 5. e11861.
- Hasegawa, P.M. 2013. Sodium ( $\text{Na}^+$ ) homeostasis and salt tolerance of plants. *Environmental and Experimental Botany*. 92: 19–31.
- Hashim, G. M. 2003. Salt-Affected Soils of Malaysia in Proceedings of the Workshop on Soil Science in Malaysia towards 2020” Malaysian Society of Soil Science. (Unpublished). [Date Accessed: 30 November 2016] Source: <ftp://ftp.fao.org/agl/agll/ladadocs/malaysia.doc>.
- He, X.J., Zhang, Z.G., Yan, D.Q., Zhang, J.S., and Chen, S.Y. 2004. A salt-responsive receptor-like kinase gene regulated by the ethylene signaling pathway encodes a plasma membrane serine/threonine kinase. *Theoretical and Applied Genetics*. 109: 377–383.
- He, Z.H., Fujiki, M. and Kohorn, B.D. 1996. A cell wall-associated, receptor-like protein kinase. *Journal of Biological Chemistry*. 271(33):19789-93.
- Heenan, D.P., Lewin, L.G., McCaffery, D.W. 1988. Salinity tolerance in rice varieties at different growth stages. *Australian Journal of Experimental Agriculture*. 28:343-349.
- Herman, T., Murchie, E.H. and Warsi, A.A. 2015. Rice Production and Climate Change: A Case Study of Malaysian Rice. *Pertanika Journal of Tropical Agricultural Science*. 38 (3): 321 – 328.
- Hesberg, C., Hänsch, R., Mendel, R.R. and Bittner, F. 2004. Tandem orientation of duplicated xanthine dehydrogenase genes from *Arabidopsis thaliana*: differential gene expression and enzyme activities. *Journal of Biological Chemistry*. 279(14):13547-54.
- Ho, C.L., Tan, Y.C., Yeoh, K.A., Ghazali, A.K., Yee, W.Y. and Hoh, C.C. 2016. De novo transcriptome analyses of host- fungal interactions in oil palm (*Elaeis guineensis* Jacq.). *BMC Genomics*. 17:66 DOI 10.1186/s12864-016-2368-0.
- Hoang, T.M.L., Tran, T.N., Nguyen, T.K.N., Williams, B., Wurm, P. and Bellairs, S. 2016. Improvement of salinity stress tolerance in rice: challenges and opportunities. *Agronomy*. 6(4):54.
- Horie, T. and Karahara, I. 2012. Salinity tolerance mechanisms in glycophytes: An overview with the central focus on rice plants. *Rice*5.
- Hou, X., Tong, H., Selby, J., DeWitt, J., Peng, X., & He, Z.-H. 2005. Involvement of a Cell Wall-Associated Kinase, WAKL4, in *Arabidopsis* Mineral Responses. *Plant Physiology*. 139(4): 1704–1716.



- <http://english.vietnamnet.vn/fms/special-reports/146141/vietnam-s--rice-bowl-is-sinking.html>. 13/11/2015.
- Htwe, N.N., Maziah, M., Ling, H.C., Zaman, F.Q. and Zain, A.M. 2011. Responses of some selected Malaysian rice genotypes to callus induction under *in vitro* salt stress. *African Journal of Biotechnology*. 10(3): 350-362.
- Hu, M.L., Chen, Y.K. and Lin, Y.F. 1995. The antioxidant and prooxidant activity of some B vitamins and vitamin-like compounds. *Chemico-Biological Interactions*. 97:63–73.
- Huang, X., Kurata, N., Wei, X., Wang, Z.X., Wang, A. and Zhao, Q. 2012. A map of rice genome variation reveals the origin of cultivated rice. *Nature*. 490: 7421.
- Huse, Huber, J.A. and Morrison, H.G. 2007. Accuracy and quality of massively parallel DNA pyrosequencing. *Genome Biology*. 8:7:143.
- Illumina. 2011. Technical note: Sequencing. Quality scores for next-generation sequencing: assessing sequencing accuracy using Phred quality scoring. Illumina [http://www.illumina.com/documents/products/technotes/technote\\_Q-Scores.pdf](http://www.illumina.com/documents/products/technotes/technote_Q-Scores.pdf).
- International Fund for Agricultural Development, World Food Programme, Food and Agricultural Organization. 2012. The State of Food Insecurity in the World 2012. Economic growth is necessary but not sufficient to accelerate reduction of hunger and malnutrition. Rome.
- IPCC. 2001. Climate Change 2001: Impacts, Adaptation and Vulnerability. McCarthy J. *et al.* (editors) Contribution of Working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change, World Meteorological Organisation and United Nations Environment Programme. Cambridge University Press.
- IRRI. 2002. Standard evaluating score for rice (SES).
- IRRI. 2012. Rice Knowledge Bank. Accessed 15 March 2016 at <http://www.knowledgebank.irri.org/qualityseedcourse/index.php/module-2-production-of-rice-seed-in-irrigated-mainmenu-103/3-sowing-seeds-and-managing-the-seed-bed-mainmenu-48/187>.
- Ishikawa, T. 1992. The ATP-dependent glutathione S-conjugate export pump. *Trends Biochem Science*. 17(11):463-8.
- Islam, M.Z., Baset Mia, M.A., Islam, M.R. and Akter, A. 2007. Effect of Different Saline Levels on Growth and Yield Attributes of Mutant Rice. *Journal of soil and nature*. 1(2): 18-22.
- Jain, M., Nijhawan, A., Tyagi, A.K. 2006. Validation of housekeeping genes as internal control for studying gene expression in rice by quantitative real-time PCR. *Biochemical and Biophysical Research Communications*. 345: 646–651.
- James RA, Blake C, Byrt CS, Munns R. 2011. Major genes for Na<sup>+</sup> exclusion, Nax1 and Nax2 (wheat HKT1;4 and HKT1;5), decrease Na<sup>+</sup> accumulation in bread wheat leaves under saline and waterlogged conditions. *Journal of Experimental Botany*. 62(8):2939–2947.
- Jeon, E.H., Pak, J.H., Kim, M.J., Kim, H.J., Shin, S.H., Lee, J.H. 2012. Ectopic expression of ubiquitin-conjugating enzyme gene from wild rice, OgUBC1, confers resistance against UV-B radiation and Botrytis infection in *Arabidopsis thaliana*. *Biochemical and Biophysical Research Communications*. 427: 309–314.
- Jiyanan, H., Xiaobo, Z., Huihui, Y., Yidan, O., Lei, W. and Qifa, Z. 2009. The ankyrin repeat gene family in rice: genome-wide identification, classification and expression profiling. *Plant Molecular Biology*. 71:207–226.

- Jones, D., Crowe, E., Stevens, T.A. and Candido, E.P.M. 2001. Functional and phylogenetic analysis of the ubiquitylation system in *Caenorhabditis elegans*: ubiquitin-conjugating enzymes, ubiquitin-activating enzymes, and ubiquitin-like proteins. *Genome Biol.* 3: research0002.1—research0002.15.
- Jones, P., Binns, D., Chang, H.Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.Y., Lopez, R. and Hunter, S. 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics.* 30 (9): 1236-1240.
- Joseph, E.A. and Mohanan, K.V. 2013. A Study on the Effect of Salinity Stress on the Growth and Yield of Some Native Rice Cultivars of Kerala State of India. *Agriculture, Forestry and Fisheries.* 2(3): 141-150. doi: 10.11648/j.aff.20130203.14.
- Joshi, A., Hung, D.Q., Vaid, N., and Tuteja, N. 2010. Pea lectin receptor-like kinase promotes high salinity stress tolerance in bacteria and expresses in response to stress in planta. *Glycoconjugate Journal.* 27: 133–150.
- Kakumanu A, Ambavaram MM, Klumas C, Krishnan A, Batlang U, Myers E. 2012. Effects of drought on gene expression in maize reproductive and leaf meristem tissue revealed by RNA-Seq. *Plant Physiology.* 160(2): 846–67.
- Kam, S.P., Tawang, A., Hoanh, C.T., Hamzah, A.R., Rala, A. and Villano, L. 2002. Rice supply and demand scenario for Malaysia. In: “*Developments in the Asian Rice Economy*” (M Sombilla, M Hossain & B Hardy (eds)). IRRI, Philippines. p 261-88.
- Kamaruddin, H. 1998. Coastal Zone Management In Malaysia- Pollution Control. *Malaysian Journal of Law Society.* 2: 46-55.
- Kang, J., Park, J., Choi, H., Burla, B., Kretzschmar, T., Lee, Y., & Martinoia, E. 2011. Plant ABC Transporters. The Arabidopsis Book. *American Society of Plant Biologists*, 9, e0153. <http://doi.org/10.1199/tab.0153>
- Karan, R., DeLeon, T., Biradar, H. and Subudhi, P.K. 2012. Salt stress induced variation in DNA methylation pattern and its influence on gene expression in contrasting rice genotypes. *PLoS One.* 7(6): e40203.
- Kawasaki, S., Borchert, C., Deyholos, M., Wang, H., Brazille, S., Kawai, K., Galbraith, D. and Bohnert, H.J. 2001. Gene expression profiles during the initial phase of salt stress in rice. *Plant Cell.* 13(4):889-905.
- Khan, Y.J., Tyagi, H., Singh, A.K. and Magadum, S. 2016. Abiotic Stress Tolerance in Soybean : Regulated by ncRNA. *Journal of AgriSearch.* 3(1): 1-6.
- Khan, Y.S., Tyagi, H., Sigh, A.K. and Magadum, S. 2016. Abiotic Stress Tolerance in Soybean : Regulated by ncRNA. *Journal of AgriSearch* 3(1): 1-6.
- Kobayashi, Y., Yamamoto, S., Minami, H., Kagaya, Y. and Hattori, T. 2004. Differential activation of the rice sucrose nonfermenting1-related protein kinase2 family by hyperosmotic stress and abscisic acid. *Plant Cell.* 16,1163–1177.
- Koenig D, Jimenez-Gomez JM, Kimura S, Fulop D, Chitwood DH, et al. 2013. Comparative transcriptomics reveals patterns of selection in domesticated and wild tomato. *Proceedings of the National Academy of Sciences.* 110: E2655–2662.
- Kohorn, B.D. 2000. Plasma membrane-cell wall contacts. *Plant Physiology.* 124(1): 31-8.
- Korf, I. 2013. Genomics: the state of the art in RNA-seq analysis. *Nature Methods.* 10: 1165–1166.

- Kosová K, Prášil IT, Vitámvás P. 2013. Protein contribution to plant salinity response and tolerance acquisition. *International Journal of Molecular Sciences*. 14:6757-6789 DOI:10.3390/ijms14046757.
- Kovach, M.J., Sweeney, M.T. and McCouch, S.R. 2007. New insights into the history of rice domestication. *Trends Genetics*. 23:578–587.
- Kreuz, K., Tommasini, R. and Martinoia, E. 1996. Old Enzymes for a New Job (Herbicide Detoxification in Plants). *Plant Physiology*. 111(2):349-353.
- Kursteiner, O., Dupuis, I. and Kuhlemeier, C. 2003. The pyruvate decarboxylase1 gene of *Arabidopsis* is required during anoxia but not other environmental stresses. *Plant Physiology*. 132:968–978.
- Kulik, A., Wawer, I., Krzywińska, E., Bucholc, M., & Dobrowolska, G. 2011. SnRK2 Protein Kinases—Key Regulators of Plant Response to Abiotic Stresses. *OMICS: A Journal of Integrative Biology*. 15(12): 859–872.
- Kumar, V. and Khare, T. 2016. Differential growth and yield responses of salt-tolerant and susceptible rice cultivars to individual (Na<sup>+</sup> and Cl<sup>-</sup>) and additive stress effects of NaCl. *Acta Physiologiae Plantarum*. 38:170.
- Lally, D., Ingmire, P., Tong, H.Y. and He, Z.H. 2001. Antisense expression of a cell wall-associated protein kinase, WAK4, inhibits cell elongation and alters morphology. *Plant Cell*. 13(6):1317-31.
- Langmead, B., Trapnell, C., Pop, M. and Salzberg, S.L. 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biology*. 10:R25.
- Laporte P, Merchan F, Amor BB, Wirth S, Crespi M. 2007. Riboregulators in plant development. *Biochemical Society Transactions*. 35(6):1638-42.
- Ledergerber, C. and Dessimoz, C. 2011. Base-calling for next-generation sequencing platforms. *Briefings Bioinformatics*. 12 (5): 489-497.
- Li, S., Fan, C., Li, Y., Zhang, J., Sun, J., Chen, Y., Tian, C., Su, X., Lu, M., Liang, C. and Hu, Z. 2016. Effects of drought and salt-stresses on gene expression in *Caragana korshinskii* seedlings revealed by RNA-seq. *BMC Genomics*. 17: 200.
- Liepmann, A.H., Wightman, R., Geshi, N., Turner, S.R. and Scheller, H.V. 2010. *Arabidopsis* a powerful model system for plant cell wall research. *Plant Journal*. 61:1107–1121.
- Linh, T.H., Xuan, T.D., Ham, L.H., Ismail, A.M. and Khanh, T.D. 2012. Molecular breeding to improve salt tolerance of rice (*Oryza sativa* L.) in the Red River Delta of Vietnam. *International Journal of Plant Genomics*. Article ID 949038, 10.1155/2012/949038.
- Liu, F., Jenssen, T.K., Trimarchi, J., Punzo, C., Cepko, C.L., Ohno-Machado, L. 2007. Comparison of hybridization-based and sequencing-based gene expression technologies on biological replicates. *BMC Genomics*. 20(8): 153.
- Liu, J., Wang, H. and Chua, N.H. 2015. Long noncoding RNA transcriptome of plants. *Plant Biotechnology Journal*. 13(3): 319-328.
- Liu, X., Hao, L., Li, D., Zhu, L. and Hu, S. 2015. Long Non-coding RNAs and Their Biological Roles in Plants. *Genomics, Proteomics & Bioinformatics*. 3: 137–147.
- Long, X.Y., Liu, Y.X. and Rocheleau, H. 2011. Identification and validation of internal control genes for gene expression in wheat leaves infected by strip rust. *Int J Plant Breeding Genet*. 5: 255–267.
- Long, X.Y., Wang, J.R. and Ouellet, T. 2010. Genome-wide identification and evaluation of novel internal control genes for q-PCR based transcript normalization in wheat. *Plant Mol Biol*. 74: 307–311.

- Los Baños (Philippines): International Rice Research Institute. 14 p
- Lukienko, P.I., Mel'nichenko, N.G., Zverinskii, I.V. and Zabrodskaya, S.V. 2000. Antioxidant properties of thiamin. *Bull Exp Biol Med.* 130: 874–876.
- Lupas, A., Van Dyke, M. and Stock, J. 1991. Predicting coiled coils from protein sequences. *Science.* 252:1162–1164.
- Lutts S, Kinet JM, Bouharmont J. 1996. NaCl-induced senescence in leaves of rice (*Oryza sativa* L.) varieties, differing in salinity resistance. *Ann Bot* 78:389–398.
- Ma, N., Ma, X., Li, A., Cao, X. and Kong, L. 2012. Cloning and Expression Analysis of Wheat Pheophorbide a Oxygenase Gene *TaPaO*. *Plant Mol Biol Rep.* 30:1237–1245.
- Mackill, D.J. 1995. Classifying japonica rice cultivars with RAPD markers. *Crop Sci.* 35:889–894.
- Maischak, H., Zimmermann, M.R., Felle, H.H., Boland, W. and Mithofer, A. 2010. Alamethicin-induced electrical long distance signaling in plants. *Plant Signal Behav.* 5:988-990.
- Maksup, S., Supaibulwatana, K. and Selvaraj, G. 2013. High-quality reference genes for quantifying the transcriptional responses of *Oryza sativa* L. (ssp. *indica* and *japonica*) to abiotic stress conditions. *Chin Sci Bull.* 58:1919-1930.
- Mardis, E.R. 2008. Next-generation DNA sequencing methods. *Annu Rev Genomics Hum Genet.* 9:387-402.
- Marone, M., Mozzetti, S., De Ritis, D., Pierelli, L. and Scambia, G. 2001. Semiquantitative RT-PCR analysis to assess the expression levels of multiple transcripts from the same sample. *Biol. Proced. Online.* 3(1): 19-25.
- Martínez, J.P., Silva, H., Ledent, J.F. and Pinto, M. 2007. Effect of drought stress on the osmotic adjustment, cell wall elasticity and cell volume of six cultivars of common beans (*Phaseolus vulgaris* L.). *Eur. J. Agron.* 26:30–38.
- Martinoia, E., Grill, E., Tommasini, R., Kreuz, K. and Amrhein, N. 1993. ATP-dependent glutathione S-conjugate export pump in the vacuolar membrane of plants. *Nature.* 364:247–249.
- Mary-Huard, T., Daudin, J.J., Baccini, M., Biggeri, A. and Bar-Hen, A. 2007. Biases induced by pooling samples in microarray experiments. *Bioinformatics.* 23:i313–318.
- Mäser, P., Gierth M, Schroeder JI. 2002. Molecular mechanisms of potassium and sodium uptake in plants. *Plant and Soil* 247, 43-54.
- Matić, S., Bagnaresi, P., Biselli, C., Orru', L., Amaral Carneiro, G., Siciliano, I., Valé, G., Gullino, M.L. and Spadaro, D. 2016. Comparative transcriptome profiling of resistant and susceptible rice genotypes in response to the seedborne pathogen *Fusarium fujikuroi*. *BMC Genom.* 17:608.
- Matsui, A., Ishida, J., Morosawa, T., Okamoto, M., Kim, J.M., Kurihara, Y., Kawashima, M., Tanaka, M., To, T.K. and Nakaminami, K. 2010. *Arabidopsis* tiling array analysis to identify the stress-responsive genes. *Methods Mol. Biol.* 639: 141–155.
- Matsui, A., Nguyen, A.H., Nakaminami, K. and Seki, M. 2013. *Arabidopsis* non-coding RNA regulation in abiotic stress responses. *Int J Mol Sci.* 14(11): 22642–22654.
- McGuire, S. 2013. WHO, world food programme, and international fund for agricultural development. 2012. The State of Food Insecurity in the World 2012. Economic growth is necessary but not sufficient to accelerate reduction of hunger and malnutrition. Rome,



- Menguer, P.L., Sperotto, R.A. and Ricachenevsky, F.K. 2017. A walk on the wild side: *Oryza* species as source for rice abiotic stress Tolerance. *Genetics and Molecular Biology*. 40(1): 238-252.
- Menshanov, P.N. and Dygalo, N.N. 2016. The Specific Problems of the Experimental Design in the High-Throughput Sequencing Studies of Transcriptome published in Vavilovskii Zhurnal Genetiki i Selektzii. 20(2): 247–254.
- Miah, G., Rafii, M.Y., Ismail, M.R., Puteh, A.B., Rahim, H.A., Asfaliza, R. and Latif, M.A. 2012. Blast resistance in rice: a review of conventional breeding to molecular approaches. *Mol Biol Rep*. 40:2369–2388.
- Michael D, Peel B, Waldron L, Kevin B. 2004. Screening for salinity tolerance in Alfalfa. *Crop Sci*. 44: 2049-2053.
- Mirouze, M. and Paszkowski, J. 2011. Epigenetic contribution to stress adaptation in plants. *Curr Opin Plant Biol*. 14: 267–274.
- Misra RC, Sandeep, Kamthan M, Kumar S, Ghosh S. 2016. A thaumatin-like protein of *Ocimum basilicum* confers tolerance to fungal pathogen and abiotic stress in transgenic Arabidopsis. *Sci Rep*. 6:25340
- Misra, R.C., Sandeep, Kamthan, M., Kumar, S. and Ghosh, S. 2016. A thaumatin-like protein of *Ocimum basilicum* confers tolerance to fungal pathogen and abiotic stress in transgenic Arabidopsis. *Sci Rep*. 6: 25340.
- Misratia, K.M., Ismail, M. R., Hakim, M.A., Musa, M.H. and Puteh, A. 2013. Effect of salinity and alleviating role of gibberellic acid (GA3) for improving the morphological, physiological and yield traits of rice varieties. *Aust. J. Crop Sci*. 7: 1682-1692.
- Mittler, R. 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci*. 7:405–410.
- Mittler, R., Vanderauwera, S., Gollery, M. and Breusegem, V. 2004. Reactive oxygen gene network of plants. *Trends Plant Sci*. 9:490–498.
- Mittler, R., Vanderauwera, S., Suzuki, N., Miller, G., Tognetti, V.B., Vandepoele, K., Gollery, M, Shulaev, V, Van, Breusegem, F. 2011. ROS signaling: the new wave? *Trends Plant Sci*. 16:300-309.
- Mizuno H, Kawahara Y, Sakai H, Kanamori H, Wakimoto H, Yamagata H. 2010. Massive parallel sequencing of mRNA in identification of unannotated salinity stress-inducible transcripts in rice (*Oryza sativa* L.). *BMC Genomics*. 11: 683.
- Mohanty S, Wassmann R, Nelson A, Moya P, and Jagadish SVK. 2013. Rice and climate change: significance for food security and vulnerability. IRRI Discussion Paper Series No. 49.
- Mohanty, S. 2013. Rice facts: Trends in global rice consumption. Social Sciences Division at IRRI. Rice Today January-March 2013.
- Morillo, S.A. and Tax, .F.E. 2006. Functional analysis of receptor-like kinases in monocots and dicots. *Curr. Opin. Plant Biol*. 9:460–469.
- Munns R. 2002. Comparative physiology of salt and water stress. *Plant Cell and Environment* 25(2):239-250. DOI: 10.1046/j.0016-8025.2001.00808.x
- Munns, R. 2011. Plant adaptations to salt and water stress: Differences and commonalities. In *Advances in Botanical Research*; Ismail, T., Ed.; Elsevier: Wyman Street, Waltham, MA, USA. 57: 1–32.
- Munns, R. and Tester, M. 2008. Mechanisms of Salinity Tolerance. *Annual Review of Plant Biology* 59:651-681.
- Munns, R., James, R.A. and Lauchli, A. 2006. Approaches to increasing the salt tolerance of wheat and other cereals. *Journal of Experimental Botany* 57, 1025-1043.

- Muthayya, S., Sugimoto, J.D., Montgomery, S. and Maberly, G.F. 2014. An overview of global rice production, supply, trade, and consumption. *New York Academy of Sciences*. 1324:7–14.
- Najim, M.M.M., Lee, T.S., Haque, M.A. and Esham, M. 2007. Sustainability of rice production: a Malaysian perspective. *J Agric Sci*. 3(1):1e12.
- Nam, M.H., Huh, S.M., Kim, K.M., Park, W.J., Seo, J.B. and Cho, K. 2012. Comparative proteomic analysis of early salt stress-responsive proteins in roots of *SnRK2* transgenic rice. *Proteome Sci*. 10: 25–44. pmid:22462395.
- Narsai, R., Ivanova, A. and Ng, S. 2010. Defining reference genes in *Oryza sativa* using organ, development, biotic and abiotic transcriptome datasets. *BMC Plant Biol*. 10: 56.
- Nelson, N. and Yocum, C. Structure and function of photosystems I and II. 2006. *Annu. Rev. Plant Biol*. 57: 521–565 .
- Nguyen, C.D., Doan, Q.T., Chen, Y.C. and Pawan, K.M. 2014. Modeling the influence of river flow and salinity intrusion in the Mekong river estuary, Vietnam. *Lowland Technology International*. 16(1):14-25.
- Nishiguchi, M., Yoshida, K., Sumizono, T., and Tazaki, K. 2002. A receptor-like protein kinase with a lectin-like domain from Lombardy poplar: gene expression in response to wounding and characterization of phosphorylation activity. *Mol. Genet. Genomics*. 267: 506–514.
- Nouri, M.Z., Moumenu, A. and Komatsu, S. 2015. Abiotic Stresses: Insight Into Gene Regulation and Protein Expression in Photosynthetic Pathways of Plants. *Int J Mol Sci*. 16 (9): 20392-20416.
- Nozulaidi M, Khairi M, and Jahan MS. 2015. Effects of different salinity levels on rice production. 2015. *Australian Journal of Basic and Applied Sciences*. 9(5): 524-530.
- OCED, Organisation for Economic Co-operation and Development. 1999. Consensus document on the biology of *Oryzasativa* (rice). Report No. ENV/JM/MONO(99)26. OCED Environmental health and Safety Publications. Paris: Organisation for Economic Co-operation and Development. Accessed June 29, 2013. <http://www.oecd.org/env/ehs/biotrack/46815658.pdf>
- Oliveira, A.B.D., Alencar, N.L.M. and Gomes-Filho, E. 2013. Comparison between the water and salt stress effects on plant growth and development. In: Sener A (ed) Responses of organisms to water stress. *InTech*. ISBN 978-953-51-0933-4. doi:10.5772/5422.
- Oono Y, Kawahara Y, Kanamori H, Mizuno H, Yamagata H, Yamamoto M. 2011. mRNA-Seq Reveals a Comprehensive Transcriptome Profile of Rice under Phosphate Stress. *Rice*. 4(2): 50–65.
- Ouyang B, Yang T, Li H, Zhang L, Zhang Y, Zhang J, Fei Z, Ye Z. 2007. Identification of early salt stress response genes in tomato root by suppression subtractive hybridization and microarray analysis. *J Exp Bot*. 58(3):507–520.
- Ouyang, S.Q., Liu, Y.F., Liu, P., Lei, G., He, S.J., Ma, B., Zhang, W.K., Zhang, J.S. and Chen, S.Y. 2010. Receptor-like kinase OsSIK1 improves drought and salt stress tolerance in rice (*Oryza sativa*) plants. *Plant J*. 62(2):316-329.
- Panikashvili, D., Shi, J.X., Bocobza, S., Franke, R.B., Schreiber, L. and Aharoni, A. 2010. The *Arabidopsis* DSO/ABCG11 transporter affects cutin metabolism in reproductive organs and suberin in roots. *Mol. Plant*. 3:563-575.
- Park, S.J., Huang, Y., Ayoubi, P. 2006. Identification of expression profiles of sorghum genes in response to greenbug phloem-feeding using cDNA subtraction and microarray analysis. *Planta*. 223(5): 932–47.

- Park, Y.W., Baba, K., Furuta, Y., Iida, I., Sameshima, K., Arai, M. and Hayashi, T. 2004. Enhancement of growth and cellulose accumulation by overexpression of xyloglucanase in poplar. *FEBS Lett.* 564:183–187.
- Park, Y.W., Tominaga, R., Sugiyama, J., Furuta, Y., Tanimoto, E., Samejima, M., Sakai, F. and Hayashi, T. 2003. Enhancement of growth by expression of poplar cellulase in *Arabidopsis thaliana*. *Plant J.* 33:1099–1106.
- Parveen, S., Gupta, D.B., Dass, S., Kumar, A., Pandey, A., Chakraborty, S. and Chakraborty, N. 2016. Chickpea Ferritin CaFer1 Participates in Oxidative Stress Response, and Promotes Growth and Development. *Scientific reports.* 6:31218.
- Patishtan, J., Hartley, T.N., Fonseca de Carvalho, R. and Maathuis, F.J. 2017. Genome Wide Association Studies To Identify Rice Salt-Tolerance Markers. *Plant Cell Environ.* doi: 10.1111/pce.12975.
- Paulin L, Beritognolo I, Harfouche A, Brillì F, Prosperini G, Gaudet M, Brosché M, Salani F, Kuzminsky E, Auvinen P. 2011. Comparative study of transcriptional and physiological responses to salinity stress in two contrasting *Populus alba* L. genotypes. *Tree Physiol* 31:1335–1355.
- Peng, Z., He, S., Gong, W., Sun, J., Pan, Z., Xu, F. 2014. Comprehensive analysis of differentially expressed genes and transcriptional regulation induced by salt stress in two contrasting cotton genotypes. *BMC Genomics.* 15:760.
- Pessaraki, M. and Szabolcs, I. 2010. Soil salinity and sodicity as particular plant/crop stress factors. In “Handbook of plant and crop stress, 3<sup>rd</sup> ed., Revised and expanded (Ed. Pessaraki, M.), CRC Press, Taylor and Francis Publishing Company, Florida. 3-21.
- Peterson, T.A., Lovatt, C.J. and Nieman, R.C. 1988. Salt Stress Causes Acceleration of Purine Catabolism and Inhibition of Pyrimidine Salvage in *Zea mays* Root Tips. *Journal of Experimental Botany.* 39(207):1389-1395.
- Petronia, C. 2011. Abiotic Stress in Plants - Mechanisms and Adaptations (A Shanker, Ed.). *InTech.*
- Pfaffl, M.W., Tichopad, A. and Prgomet, C. 2004. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: Bestkeeper-excel-based tool using pair-wise correlations. *Biotechnol Lett.* 26: 509–515.
- Pratt, L.H., Liang, C., Shah, M., Sun, F., Wang, H., Reid, S.P. 2005. Sorghum expressed sequence tags identify signature genes for drought, pathogenesis, and skotomorphogenesis from a milestone set of 16,801 unique transcripts. *Plant Physiol.* 139(2): 869–84.
- Qin D, Wu H, Peng H, Yao Y, Ni Z, Li Z, *et al.* Heat stress-responsive transcriptome analysis in heat susceptible and tolerant wheat (*Triticum aestivum* L.) by using Wheat Genome Array. *BMC Genomics.* 2008;9:432.
- Qin, D., Wu, H., Peng, H., Yao, Y., Ni, Z., Li, Z., Zhou, C. and Sun, Q. 2008. Heat stress-responsive transcriptome analysis in heat susceptible and tolerant wheat (*Triticum aestivum* L.) by using Wheat Genome Array. *BMC Genomics.* 9: 432.
- Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R. and Lopez, R. 2005. InterProScan: protein domains identifier. *Nucleic Acids Res.* 33 (suppl\_2).
- Rafii, M.Y., Zakiah, M.Z., Asfaliza, R., Iffah Haifaa, M.D., Latif, M.A. and Malek, M.A. 2014. Grain quality performance and heritability estimation in selected F1 rice genotypes. *Sains Malaysiana.* 43 (1):1-7. ISSN 0126-6039

- Rahaie, M., Xue, G.P. and Schenk, P.M. 2013. The role of transcription factors in wheat under different abiotic stress. In: Vahdati K, Leslie C (eds) Abiotic stress—plant responses and applications in agriculture. *Intech*. doi:10.5772/54795.
- Rahnama A, James RA, Poustini K, Munns R. 2010. Stomatal conductance as a screen for osmotic stress tolerance in durum wheat growing in saline soil. *Functional Plant Biology*. 37(3): 255–263.
- Rajamoorthy, Y., Rahim, K.B.A. and Munusamy, S. 2015. Rice industry in Malaysia: challenges, policies and implications. *Procedia Economics and Finance*. 31: 861–867.
- Rajkumar, A.P., Qvist, P., Lazarus, R., Lescai, F., Ju, J., Nyegaard, M., Mors, O., Borglum, A.D., Li, Q. and Christensen, J.H. 2015. Experimental validation of methods for differential gene expression analysis and sample pooling in RNA-seq. *BMC Genomics*. 16: 548.
- Rapala-Kozik, M., Kowalska, E. and Ostrowska, K. 2008. Modulation of thiamine metabolism in *Zea mays* seedlings under conditions of abiotic stress. *Journal of Experimental Botany*. 59(15):4133–4143.
- Rapala-Kozik, M., Wolak, N., Kudja, M. and Banas, A.K. 2012. The upregulation of thiamine (vitamin B<sub>1</sub>) biosynthesis in *Arabidopsis thaliana* seedlings under salt and osmotic stress conditions is mediated by abscisic acid at the early stages of this stress response. *BMC Plant Biology*. 12(2).
- Rayon, C., Gall, H.L., Philippe, F., Domon, J., Gillet, F. and Pelloux, J. 2015. Cell wall metabolism in response to abiotic stress. *Plants*. 4:112-166.
- Reddy MP, Sanish S, Iyengar ERR. 1992. Photosynthetic studies and compartmentation of ions in different tissues of *Salicornia brachiata* Roxb. under saline conditions. *Photosynthetica* 26: 173–179.
- Reddy, A.S., Ali, G.S., Celesnik, H. and Day, I.S. 2011. Coping with stresses: roles of calcium- and calcium/calmodulin-regulated gene expression. *Plant Cell*. 23(6): 2010-2032. doi:tpc.111.084988 [pii]
- Reddy, I.N.B.L., Kim, B.K., Yoon, I.S., Kim, K.H., and Kwon, T.R. 2017. Salt tolerance in rice: Focus on mechanisms and approaches. *Rice Science*. 24: 123–144.
- Redfern, S.K., Azzu, N., Binamira, J.S. 2012. Rice in Southeast Asia: facing risks and vulnerabilities to respond to climate change. Build Resilience Adapt Climate Change Agri Sector. 23:295.
- Reinders, J., Wulff, B.B.H., Mirouze, M., Mari-Ordonez, A. and Dapp, M. 2009. Compromised stability of DNA methylation and transposon immobilization in mosaic *Arabidopsis* epigenomes. *Genes Dev*. 23: 939–950.
- Rengasamy P. 2002. Transient salinity and subsoil constraints to dryland farming in Australian sodic soils: an overview. *Aust. J. Exp. Agric*. 42:351–61.
- Rennenberg, H. and Poll, A. 1994. Protection from oxidative stress in transgenic plants. *Biochem. Soc. Trans*. 22:936–940.
- Revathi, S. and Pillai, M.A. 2015. Novel approach in screening rice genotype for tolerance to salt stress under hydroponic culture. *Agric. Sci. Digest*. 35(4): 308-310.
- Reynolds M, and Tuberosa R, 2008. Translational research impacting on crop productivity in drought-prone environments. *Curr. Opin. Plant Biol*. 11:171-179.
- Ribeiro DT, Farias LP, de Almeida JD, Kashiwabara PM, Ribeiro AF, Silva-Filho MC, Menck CF, Van Sluys MA. 2005. Functional characterization of the th1 promoter region from *Arabidopsis thaliana*. *J Exp Bot* 2005, 56:1797-1804.



- Richards L.A. 1954. Diagnosis and improvement of saline and alkali soils. United States Department of Agriculture, Washington, DC.
- Riou, C., Hervé, C., Pacquit, V., Dabos, P., and Lescure, B. 2002. Expression of an *Arabidopsis* lectin kinase receptor gene, *lecRK-a1*, is induced during senescence, wounding and in response to oligogalacturonic acids. *Plant Physiol. Biochem.* 40: 431–438.
- Roberts, A., Pimentel, H., Trapnell, C. and Pachter, L. 2011. Identification of novel transcripts in annotated genomes using RNA-seq. *Bioinformatics* 27:2325–2329.
- Robinson, M.D., McCarthy, D.J. and Smyth, G.K. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics.* 26:139–40.
- Robles JA, Qureshi SE, Stephen SJ, Wilson SR, Burden CJ, Taylor JM. 2012. Efficient experimental design and analysis strategies for the detection of differential expression using RNA-Sequencing. *BMC Genomics.* 13: 484-10.1186/1471-2164-13-484.
- Rouhier, N., Lemaire, S.D. and Jacquot, J.P. 2008. The role of glutathione in photosynthetic organisms: emerging functions for glutaredoxins and glutathionylation. *Annu. Rev. Plant Biol.* 59:143-166.
- Roy, B. and Bhadra, S. 2014. Effects of Toxic Levels of Aluminium on Seedling Parameters of Rice under Hydroponic Culture. *Rice Science.* 21(4): 217–223.
- Roy, S. and Chakraborty, U. 2014. Salt tolerance mechanisms in Salt Tolerant Grasses (STGs) and their prospects in cereal crop improvement. *Botanical Studies.* 55: 31.
- Rozema J, Flowers T. 2008. Ecology: crops for a salinized world. *Science.* 322(5907): 1478–1480.
- Rubio, S., Whitehead, L., Larson, T.R., Graham, I.A. and Rodriguez, P.L. 2008. The Coenzyme A Biosynthetic Enzyme Phosphopantetheine Adenylyltransferase Plays a Crucial Role in Plant Growth, Salt/Osmotic Stress Resistance, and Seed Lipid Storage Plant Physiology. 148: 546–556.
- Rymarquis, L.A., Kastenmayer, J.P., Hüttenhofer, A.G. and Green, P.J. 2008. Diamonds in the rough: mRNA-like non-coding RNAs. *Trends Plant Sci.* 13(7):329-34.
- Sadanandom, A., Bailey, M., Ewan, R., Lee, J., Nelis, S. 2012. The ubiquitin-proteasome system: central modifier of plant signalling. *New Phytol.* 196: 13–28. doi: 10.1111/j.1469-8137.2012.04266.x PMID: 22897362.
- Sakai, H., Lee, S.S., Tanaka, T., Numa, H., Kim, J., Kawahara, Y., Wakimoto, H., Yang, C.C., Iwamoto, M. and Abe, T. 2013. Rice Annotation Project Database (RAP-DB): an integrative and interactive database for rice genomics. *Plant Cell Physiol.* 54:e6.
- Sakamoto, H., Tochimoto, J., Kurosawa, S., Suzuki, M. and Oguri, S. 2014. A Novel Gene Encoding Ankyrin-Repeat Protein, *SHG1*, is Indispensable for Seed Germination under Moderate Salt Stress. *International Journal of Biological, Biomolecular, Agricultural, Food and Biotechnological Engineering* 8(3).
- Sani, K., 1991. The management of saline soils for paddy cultivation (in Malay ). *Teknologi Kejuruteraan Pertanian.*, 20: 33-36.
- Schachtman D, Liu WH. 1999. Molecular pieces to the puzzle of the interaction between potassium and sodium uptake in plants. *Trends Plat Sci.* 4, 282-287.
- Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. 2012. “NIH Image to ImageJ: 25 years of image analysis”. *Nature Methods.* 671.

- Schroeder, A., Mueller, O., Stocker, S., Salowsky, R., Leiber, M. and Gassmann, M. 2006. The RIN: an RNA integrity number for assigning integrity values to RNA measurements. *BMC Mol Biol.* 7:3.
- Schurch N. J., Schofield P., Gierliński M., Cole C., Sherstnev A., Singh V. 2016. How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use?. *RNA.* 22: 839–851. 10.1261/rna.053959.115
- Selamat A, Ismail MR. 2009. Deterministic model approaches in identifying and quantifying technological challenges in rice production and research, and in predicting population, rice production and consumption in Malaysia. *Pertanika J of Trop Agric Sci.* 32:267-291.
- Sengupta, S. and Majumder, A.L. 2009. Insight into the salt tolerance factors of a wild halophytic rice, *Porteresia coarctata*: A physiological and proteomic approach. *Planta.* 229:911–929.
- Seong, E.S., Cho, H.S., Choi, D., Joung, Y.H., Lim, C.K., Hur, J.H. and Wang, M.H. 2007a. Tomato plants overexpressing CaKR1 enhanced tolerance to salt and oxidative stress. *Biochem Biophys Res Commun* 363:983–988.
- Seong, E.S., Choi, D., Cho, H.S., Lim, C.K., Cho, H.J. and Wang, M.H. 2007b. Characterization of a stress-responsive ankyrin repeat-containing zinc finger protein of *Capsicum annuum* (CaKR1). *J. Biochem. Mol. Biol.* 40(6).
- Shankar, R., Bhattacharjee, A., & Jain, M. 2016. Transcriptome analysis in different rice cultivars provides novel insights into desiccation and salinity stress responses. *Scientific Reports.* 6: 23719.
- Shaoliang, Zhiqin, L., Deyi, G., Ailian, Q., Yan, L. and Shuilin, H. 2013. Functional Analysis and Expressional Characterization of Rice Ankyrin Repeat-Containing Protein, OsPIANK1, in Basal Defense against *Magnaporthe oryzae* Attack. *PLoS One.* 8(3): e59699.
- Shavrukov, Y. 2012. Salt stress or salt shock: which genes are we studying? *Journal of Experimental Botany Advance Access.* DOI:10.1093/jxb/ers316.
- Shinozaki, K., Yamaguchi-Shinozaki, K. and Sekiz, M. 2003. Regulatory network of gene expression in the drought and cold stress responses. *Curr. Opin. Plant Biol.* 6:410-417.
- Shiu, S.H. and Bleecker, A.B. 2001. Receptor-like kinases from *Arabidopsis* form a monophyletic gene family related to animal receptor kinases. *PNAS.* 98: 10763–10768.
- Singh, A., Kushwaha, H.R., Soni, P., Gupta, H., Singla-Pareek S.L., and Pareek, A. 2015. Tissue specific and abiotic stress regulated transcription of histidine kinases in plants is also influenced by diurnal rhythm. *Front. Plant Sci.* 6:711. 10.3389/fpls.2015.00711.
- Sita Devi, K, Ponnarasi. T. 2009. An economic analysis of modern rice production technology and its adoption behaviour in Tamil Nadu. *Agric. Econ. Res. Rev.* 22: 341–347.
- Sivaguru, M., Ezaki, B., He, Z-H., Tong, H., Osawa, H., Baluska, F., Volkmann, D. and Matsumoto, H. 2003. Aluminum induced gene expression and protein localization of a cell wall-associated receptor protein kinase in *Arabidopsis thaliana*. *Plant Physiol.* 132: 2256–2266.
- Sohrabi, M., Rafii, M.Y., Hanafi, M.M., Siti Nor Akmar, A. and Latif, M.A. 2012. Genetic diversity of upland rice germplasm in Malaysia based on quantitative traits. *Sci World J.* 416291.
- Soneson, C and Delorenzi, M. 2013. A comparison of methods for differential expression analysis of RNA-seq data. *BMC Bioinformatics.* 14:91.

- Song, X., Yu, X., Hori, C., Demura, T., Ohtani, M., & Zhuge, Q. 2016. Heterologous Overexpression of Poplar SnRK2 Genes Enhanced Salt Stress Tolerance in *Arabidopsis thaliana*. *Frontiers in Plant Science*. 7: 612.
- Sonon, L.S., Saha, U. and Kissel, D.E. 2015. Soil Salinity Testing, Data Interpretation and Recommendations. *Agricultural and Environmental Services Laboratories*. UGA Cooperative Extension Circular 1019. <http://extension.uga.edu/publications/detail.cfm?number=C1019>.
- Spargo, J. 2013. Interpreting Your Soil Test Results. UMass Soil And Plant Nutrient Testing Laboratory; Tracy Allen, Laboratory Manager; Solomon Kariuki, Laboratory Program Coordinator. Ref. No.SPTTL\_2.
- Stasolla C, Katahira R, Thorpe TA, Ashihara H. 2003. Purine and pyrimidine nucleotide metabolism in higher plants. *J Plant Physiol*. 160(11):1271-95.
- Sun, X.-L., Yu, Q.-Y., Tang, L.-L., Ji, W., Bai, X., Cai, H., Liu, X.-F., Ding, X.-D., and Zhu, Y.-M. 2012. GsSRK, a G-type lectin S-receptor-like serine/threonine protein kinase, is a positive regulator of plant tolerance to salt stress. *J. Plant Physiol*.
- Suzuki, M.M. and Bird, A. 2008. DNA methylation landscapes: provocative insights from epigenomics. *Nat Rev Genet*. 9: 465–476.
- Sweeney, M. and McCouch, S. 2007. The Complex History of the Domestication of Rice. *Ann Bot*. 100 (5): 951-957.
- Szabolcs I. 1989. *Salt-Affected Soils*. Boca Raton, FL: CRC Press.
- T H P, Friedlander MR, Almlöf J, Sammeth M, Pulyakhina I, Anvar SY. 2013. Reproducibility of high-throughput mRNA and small RNA sequencing across laboratories. *Nat Biotechnol*. 31(11): 1015–22.
- Takagi, H., Ishiga, Y., Watanabe, S., Konishi, T., Egusa, M., Akiyoshi, N., Matsuura, T., Mori, I.C., Hirayama, T., Kaminaka, H., Shimada, H. and Sakamoto, A. 2016. Allantoin, a stress-related purine metabolite, can activate jasmonate signaling in a MYC2-regulated and abscisic acid-dependent manner. *J Exp Bot*. 67(8): 2519-2532.
- Tanaka, H., Osakabe, Y., Katsura, S., Mizuno, S., Maruyama, K., Kusakabe, K., Mizoi, J., Shinozaki, K., Yamaguchi-Shinozaki, K. 2012. Abiotic stress-inducible receptor-like kinases negatively control ABA signaling in *Arabidopsis*. *Plant J*. 70: 599–613.
- Tanaka, T., Antonio, B.A., Kikuchi, S., Matsumoto, T., Nagamura, Y., Numa, H., Sakai, H., Wu, J. and Itoh T. 2008. The Rice Annotation Project Database (RAP-DB): 2008 update. *Nucleic Acids Res*. 36:D1028–D1033.
- Tavakkoli, E., Rengasamy, P., and McDonald, G. K. 2010. High concentrations of Na<sup>+</sup> and Cl<sup>-</sup> ions in soil solution have simultaneous detrimental effects on growth of faba bean under salinity stress. *Journal of Experimental Botany*, 61(15): 4449–4459.
- Teh, S.Y. 2016. Climate change and soil salinization: impact on agriculture, water and food security. *International Journal of Agriculture, Forestry and Plantation*. Vol 2.
- Ten Hove, C.A., Bochdanovits, Z. and Jansweijer, V.M.A. 2011. *Plant Mol Biol*. 76: 69.
- Thiemann, A., Fu, J., Schrag, T.A., Melchinger, A.E., Frisch, M. and Scholten S. 2010. Correlation between parental transcriptome and field data for the characterization of heterosis in *Zea mays* L. *Theor Appl Genet*. 120(2): 401–13.

- Thompson GA, Goggin FL. 2006. Transcriptomics and functional genomics of plant defence induction by phloem-feeding insects. *Journal of Experimental Botany* 57(4): 755-766.
- Toews-Shimizu, J. 2012. Rice in Japan: beyond 3.11. *Rice Today*. 11(5): 20–23, 2012.
- Trapnell, C., Hendrickson, D.G., Sauvageau, M., Goff, L., Rinn, J.L. and Pachter, L. 2013. Differential analysis of gene regulation at transcript resolution with RNA-seq. *Nat Biotechnol*. 31:46–53.
- Trapnell, C., Pachter, L. and Salzberg, S.L. 2009. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics (Oxford, England)*. 25: 1105–1111.
- Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D.R., Pimentel, H., Salzberg, S.L., Rinn, J.L. and Pachter, L. 2012. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat. Protoc*. 7(3):562–578 DOI: 10.1038/nprot.2012.016.
- Trapnell, C., Williams, B.A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M.J., Salzberg, S.L., Wold, B.J. and Pachter, L. 2010. Transcript assembly and quantification by RNA-seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol* 28: 511–515.
- Tuteja, N. 2007. Mechanisms of high salinity tolerance in plants. *Methods in enzymology*. 428: 419–38.
- Twine, N.A., Janitz, K., Wilkins, M.R. and Janitz, M. 2011. Whole transcriptome sequencing reveals gene expression and splicing differences in brain regions affected by Alzheimer’s disease. *PLoS One* 6(1): e16266.
- Ueno, K. and Miyoshi, K. 2005. Difference of optimum germination temperature of seeds of intact and dehusked japonica rice during seed development. *Springer nature*. 143: 271-275.
- Umezawa, T., Sugiyama, N., Mizoguchi, M., Hayashi, S., Myouga, F. and Yamaguchi-Shinozaki, K. 2009. Type 2C protein phosphatases directly regulate abscisic acid-activated protein kinases in *Arabidopsis*. *Proc Natl Acad Sci USA*. 106:17588–17593.
- UNEP. 2006. Geo Year Book 2006: An Overview of Our Changing Environment. Early Warning and Assessment, United Nations Environment Programme, Nairobi, Kenya, [http://www.unep.org/geo/yearbook/yb2006/PDF/Complete\\_pdf\\_GYB\\_2006.pdf](http://www.unep.org/geo/yearbook/yb2006/PDF/Complete_pdf_GYB_2006.pdf).
- USSL. 2005. George E. Brown, Jr Salinity Laboratory. Riverside, CA, USA: USDA-ARS. <http://www.ussl.ars.usda.gov>.
- Vaid, N., Pandey, P.K. and Tuteja, N. 2012. Genome-wide analysis of lectin receptor-like kinase family from *Arabidopsis* and rice. *Plant Mol. Biol*. 80: 365–388.
- Vandesompele, J., De Preter, K. and Pattyn, F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol*. 3:RESEARCH0034.
- Vaughan, D.A., Morishima, H. and Kadowaki, K. 2003. Diversity in the *Oryza* genus. *Curr. Opin. Plant Biol*. 6: 139–146.
- Verdoy, D., Lucas, M.M., Manrique, E., Covarrubias, A.A., De Felipe, R. and Pueyo, J.J. 2004. Differential organ-specific response to salt stress and water deficit in nodulated bean (*Phaseolus vulgaris*). *Plant, Cell & Environment*. 27:757–767.
- Verica, J.A., Chae, L., Tong, H., Ingmire, P. and He, Z.H. 2003. Tissue-specific and developmentally regulated expression of a cluster of tandemly arrayed cell wall-associated kinase-like genes in *Arabidopsis*. *Plant Physiol*. 133(4):1732-46.



- Vierstra, R.D. 1996. Proteolysis in plants: mechanisms and functions. *Plant Mol Biol.* 32: 275–302. PMID: 8980483.
- Vlad, F., Rubio, S., Rodrigues, A., Sirichandra, C., Belin, C. and Robert, N. 2009. Protein phosphatases 2C regulate the activation of the Snf1-related kinase OST1 by abscisic acid in *Arabidopsis*. *Plant Cell.* 21: 3170–3184.
- Wagner, T. and Kohorn, B. 2001. Wall-associated kinases are expressed throughout plant development and are required for cell expansion. *Plant Cell.* 13: 303–318.
- Wailes, E.J. and Chavez, E.C. 2012. ASEAN and Global Rice Situation and Outlook. ADB Sustainable Development Working Paper Series. Asian Development Bank. No. 22.
- Wan, X., Mo, A., Liu, S., Yang, L. and Li, L. 2011. Constitutive expression of a peanut ubiquitin-conjugating enzyme gene in *Arabidopsis* confers improved water-stress tolerance through regulation of stress-responsive gene expression. *Journal of Bioscience and Bioengineering.* 111: 478–484. doi: 10.1016/j.jbiosc.2010.11.021 PMID: 21193345
- Wang, H., Wang, H., Shao, H. and Tang, X. 2016. Recent advances in utilizing transcription factors to improve plant abiotic stress tolerance by transgenic technology. *Front. Plant Sci.* 7:67.
- Wang, J., Li, B., Meng, Y., Ma, X., Lai, Y., Si, E., Yang, K., Ren, P., Shang, X. and Wang, H. 2015. Transcriptomic profiling of the salt-stress response in the halophyte *Halogeton glomeratus*. *BMC Genomics* 16:169.
- Wang, J., Tian, L., Lee, H.S., Wei, N.E., Jiang, H., Watson, B. 2006. Genomewide nonadditive gene regulation in *Arabidopsis* allotetraploids. *Genetics.* 172(1): 507–17.
- Wang, L., Xie, W. and Chen, Y. 2010. A dynamic gene expression atlas covering the entire life cycle of rice. *Plant J.* 61: 752–766.
- Wang, W.S., Zhao, X.Q., Li, M., Huang, L.Y., Xu, J.L., Zhang, F., Cui, Y.R., Fu, B.Y., and Li, Z.K. 2016. Complex molecular mechanisms underlying seedling salt tolerance in rice revealed by comparative transcriptome and metabolomic profiling. *J. Exp. Bot.*, 67: 405-419.
- Wang, X., Liu, Z., & He, Y. 2008. Responses and tolerance to salt stress in bryophytes. *Plant Signaling & Behavior.* 3(8), 516–518.
- Wang, Z., Gerstein, M., Snyder, M. 2009. RNA-Seq: a revolutionary tool for transcriptomics. *Nat Rev Genet.* 10(1): 57–63.
- Wassmann F, Hien NX, Hoanh CT, Tuong TP. 2004. Sea level rise affecting the Vietnamese Mekong Delta: Water elevation in the flood season and implications for rice production. *Climatic Change* 66:89-107.
- Watanabe, S., Matsumoto, M., Hakomori, Y., Takagi, H., Shimada, H. and Sakamoto, A. 2014. The purine metabolite allantoin enhances abiotic stress tolerance through synergistic activation of abscisic acid metabolism. *Plant Cell Environ.* 37(4):1022-36.
- Watanabe, S., Nakagawa, A., Izumi, S., Shimada, H. and Sakamoto A. 2010. RNA interference-mediated suppression of xanthine dehydrogenase reveals the role of purine metabolism in drought tolerance in *Arabidopsis*. *FEBS Lett.* 584:1181–1186.
- Waziri, A., Kumar, P. and Purty, R.S. 2016. Saltol QTL and Their Role in Salinity Tolerance in Rice. *Austin J Biotechnol Bioeng.* 3(3): 1067.
- Welchman, R., Gordon, C., Mayer, J. 2005. Ubiquitin and ubiquitin-like proteins as multifunctional signals. *Nat Rev Mol Cell Biol* 6:599–609. PMID: 16064136.

- Werner AK, Witte C. 2011. The biochemistry of nitrogen mobilization: purine ring catabolism. *Trends in Plant Science* 16(7): 381-387.
- Wilhelm J, Pingoud A. 2003. Review Real-time polymerase chain reaction. *Chembiochem*. 4(11):1120-8.
- Wong CE, Li Y, Labbe A, Guevara D, Nuin P, Whitty B, Diaz C, Golding GB, Gray GR, Weretilnyk EA, Griffith M, Moffatt BA. 2006. Transcriptional profiling implicates novel interactions between abiotic stress and hormonal responses in *Thellungiella*, a close relative of *Arabidopsis*. *Plant Physiol* 140:1437-1450.
- Yamamoto, N., Takano, T., Tanaka, K., Ishige, T., Terashima, S., Endo, C. and Tada, Y. 2015. Comprehensive analysis of transcriptome response to salinity stress in the halophytic turf grass *Sporobolus virginicus*. *Frontiers in Plant Science*. 6: 241.
- Yan, J., Wang, J. and Zhang, H. 2002. An ankyrin repeat-containing protein plays a role in both disease resistance and antioxidation metabolism. *Plant J*. 29: 193-202.
- Yang, T.H. and Chang, P.L. 2012. Determination of RNA degradation by capillary electrophoresis with cyan light-emitted diode-induced fluorescence. *Journal of Chromatography A*. 1239:78–84.
- Yang-Rong C., Hao-Wei, C., Zhi-Gang, L., Jian-Jun, T., Biao, M., Wan-Ke, Z., Shou-Yi, C. and Jin-Song, Z. 2015. Tobacco Ankyrin Protein NEIP2 Interacts with Ethylene Receptor NTHK1 and Regulates Plant Growth and Stress Responses. *Plant Cell Physiol*. 56(4).
- Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S. and Madden, T.L. 2012. Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*. 13:134.
- Yesbergenova, Z., Yang, G., Oron, E., Soffer, D., Fluhr, R. and Sagi, M. 2005. The plant Mo-hydroxylases aldehyde oxidase and xanthine dehydrogenase have distinct reactive oxygen species signatures and are induced by drought and abscisic acid. *Plant J*. 42(6):862-76.
- Yoshida, S., Forno, D.A., Cook, J.H. and Gomez, K.A. 1976. Laboratory Manual for Physiological Studies of Rice, International Rice Research Institute, Manila: 61–65.
- Yuan JS, Reed A, Chen F, Stewart CN Jr. 2006. Statistical analysis of real-time PCR data. *BMC Bioinformatics*. 7():85.
- Yuen, B. and Kong., L. 2009. Climate Change and Urban Planning in Southeast Asia. *S.A.P.I.E.N.S* 2(3).
- Zeng L and Shannon MC. 2000. Effect of salinity on grain yield and yield components of rice at different seeding densities. *Agron. J*. 92:418-423.
- Zeng, L.H. 2005. Exploration of relationships between physiological parameters and growth performance of rice (*Oryza sativa* L.) seedlings under salinity stress using multivariate analysis. *Plant Soil* 268: 51–59.
- Zhang T, Cao C, Tang X, Cai Y, Yang C, Pan Y (2017) Enhanced peroxidase activity and tumour tissue visualization by cobalt-doped magnetoferritin nanoparticles. *Nanotechnology* 28: 045704. doi: 10.1088/1361-6528/28/4/045704.
- Zhang T, Zhao X, Wang W, Huang L, Liu X, Zong Y. 2014. Deep transcriptome sequencing of rhizome and aerial-shoot in *Sorghum propinquum*. *Plant Mol Biol*. 84(3): 315–27.
- Zhang X, Zhen J, Li Z, Kang D, Yang Y, Kong J, Hua J. 2011. Expression profile of early responsive genes under salt stress in upland cotton (*Gossypium hirsutum* L.). *Plant Molecular Biology Reporter*. 29: 626–637.

- Zhang, H., Mao, X., Jing, R., Chang, X., & Xie, H. 2011. Characterization of a common wheat (*Triticum aestivum* L.) TaSnRK2.7 gene involved in abiotic stress responses. *Journal of Experimental Botany*. 62(3): 975–988.
- Zhang, H., Mao, X., Wang, C., & Jing, R. 2010. Overexpression of a Common Wheat Gene TaSnRK2.8 Enhances Tolerance to Drought, Salt and Low Temperature in *Arabidopsis*. *PLoS ONE*. 5(12): e16041.
- Zhang, H.X., Hodson, J.N., Williams, J.P. and Blmwald, E. 2001. Engineering salt-tolerant Brassica plants: Characterization of yield and seed oil quality in transgenic plants.
- Zhang, J.L. and Shi, H. 2013. Physiological and molecular mechanisms of plant salt tolerance. *Photosynthesis research*. 115: 1–22.
- Zhang, L., Zhang, C., Wu, P., Chen, Y., Li, M., Jiang, H. and Wu, G. 2014. Global analysis of gene expression profiles in physic nut (*Jatropha curcas* L.) seedlings exposed to salt stress. *PLoS One* 9. e97878.
- Zhang, S., An, S., Li, Z., Wu, F., Yang, Q., Liu, Y., Cao, J., Zhanga, H., Zhanga, Q. and Liu, X. 2015. Identification and validation of reference genes for normalization of gene expression analysis using qRT-PCR in *Helicoverpa armigera* (Lepidoptera: Noctuidae). 555 (2): 393–402.
- Zhang, X., Rerksiri, W., Liu, A., Zhou, X., Xiong, H. and Xiang, J. 2013. Transcriptome profile reveals heat response mechanism at molecular and metabolic levels in rice flag leaf. *Gene*. 530: 185–192. doi: 10.1016/j.gene.2013.08.048 PMID: 23994682
- Zhang, Z.S., Li, X.R., Liu, L.C., Jia, R.L., Zhang, J.G. and Wang, T. 2009. Distribution, biomass, and dynamics of roots in a revegetated stand of *Caragana korshinskii* in the Tengger Desert, northwestern China. *J Plant Res*. 122(1):109-19.
- Zhao, J., He, Q., Chen, G., Wang, L., Jin, B. 2016. Regulation of Non-coding RNAs in Heat Stress Responses of Plants. *Front. Plant Sci.* 7: 1213.
- Zhao, Q., Suo, J., Chen, S., Jin, Y., Ma, X., Yin, Z., Zhang, Y., Wang, T., Luo, J., Jin, W., Zhang, X., Zhou, Z. and Dai, S. 2016a. Na<sub>2</sub>CO<sub>3</sub>-responsive mechanisms in halophyte *Puccinellia tenuiflora* roots revealed by physiological and proteomic analyses. *Scientific Reports*. 6: 32717.
- Zhinguo, E., Zhang, Y., Li, T., Wang, L. and Zhao, H. 2015. Characterization of the Ubiquitin-Conjugating Enzyme Gene Family in Rice and Evaluation of Expression Profiles under Abiotic Stresses and Hormone Treatments. *PLoS ONE*. 10(4): e0122621. doi:10.1371/journal.pone.0122621.
- Zhou, G.A., Chang, R.Z. and Qiu, L.J. 2010. Overexpression of soybean ubiquitin-conjugating enzyme gene GmUBC2 confers enhanced drought and salt tolerance through modulating abiotic stress-responsive gene expression in *Arabidopsis*. *Plant Mol Biol*. 72: 357–367. doi: 10.1007/s11103-009-9575-x PMID: 19941154
- Zhu JK. 2003. Regulation of ion homeostasis under salt stress. *Current Opinion in Plant Biology* 6(5): 441–445.
- Zhu, J.K. 2007. Plant salt stress. Encyclopedia of Life Sciences. *John Willey and Sons*. 1-7.
- Zok, A., Oláh, R., Hideg, É. 2010. Effect of *Medicago sativa* ferritin gene on stress tolerance in transgenic grapevine. *Plant Cell Tiss Organ Cult*. 100:339–344.
- Zrenner, R., Stitt, M., Sonnewald, U. and Boldt, R. 2006. Pyrimidine and Purine Biosynthesis and Degradation in Plants. *Annu. Rev. Plant Biol.* 57:805–36