



UNIVERSITI PUTRA MALAYSIA

**EXPRESSION OF HAEMAGGLUTININ-NEURAMINIDASE ENVELOPE
PROTEIN OF NEWCASTLE DISEASE VIRUS STRAIN AF2240 IN
CENTELLA ASIATICA (PEGAGA) EMBRYOGENIC CALLI THROUGH
OPTIMIZED PARTICLE BOMBARDMENT METHOD**

LAI KOK SONG

FBSB 2008 18



**EXPRESSION OF HAEMAGGLUTININ-NEURAMINIDASE
ENVELOPE PROTEIN OF NEWCASTLE DISEASE VIRUS STRAIN
AF2240 IN *CENTELLA ASIATICA* (PEGAGA) EMBRYOGENIC CALLI
THROUGH OPTIMIZED PARTICLE BOMBARDMENT METHOD**

LAI KOK SONG

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2008



**EXPRESSION OF HAEMAGGLUTININ-NEURAMINIDASE
ENVELOPE PROTEIN OF NEWCASTLE DISEASE VIRUS STRAIN
AF2240 IN *CENTELLA ASIATICA* (PEGAGA) EMBRYOGENIC CALLI
THROUGH OPTIMIZED PARTICLE BOMBARDMENT METHOD**

By

LAI KOK SONG

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirement for the Degree of Master of Science**

NOVEMBER 2008



Abstract of thesis presented to the Senate of University Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**EXPRESSION OF HAEMAGGLUTININ-NEURAMINIDASE
ENVELOPE PROTEIN OF NEWCASTLE DISEASE VIRUS STRAIN
AF2240 IN *CENTELLA ASIATICA* (PEGAGA) EMBRYOGENIC CALLI
THROUGH OPTIMIZED PARTICLE BOMBARDMENT METHOD**

By

LAI KOK SONG

June 2008

Chairman: Professor Maziah Mahmood, PhD

Faculty: Biotechnology and Biomolecular Sciences

Centella asiatica is a locally important medicinal plant. It is non-toxic, high in medicinal values, and can serve as a good candidate for genetic manipulation. However, to date no transformation protocol has been developed to fully utilize the potential of this plant. Therefore, this research is to establish an efficient particle bombardment transformation protocol for *C. asiatica* embryogenic calli. In addition, an attempt to express the haemagglutinin-neuraminidase (HN) protein from Newcastle disease virus (NDV) strain AF2240 in *C. asiatica* embryogenic calli were carried out using the developed transformation system. The HN protein can serve as a potential vaccine candidate for Newcastle disease (ND) in poultry. The induced embryogenic calli revealed the presence of extracellular matrix layer (ECM) during the microscopy studies. Particle bombardment transformation protocol was developed using the green fluorescent protein (GFP) as reporter. A total of eight parameters mainly different target distance, helium pressure, gold particles size, chamber vacuum



pressure, number of bombardment, precipitation agents, post-bombardment incubation time, and plasmid DNA concentration were identified and successfully optimized. Based on the established protocol, transformations of *C. asiatica* embryogenic calli were performed using the constructed recombinant pMDC32'HN and HBT95:sGFP(S65T)-NOS'HN plasmids. Genomic PCR analysis revealed the presence of HN transgene in the transformed lines. Unfortunately no protein bands were detected during SDS-PAGE and western blotting, indicating low or no HN protein expression. Transformation using recombinant HBT95:sGFP(S65T)-NOS'HN plasmid resulted in very low GFP expression as compared to the positive control. Nonetheless, the mRNA transcripts were detected in the RT-PCR analysis. Positive signal from the dot blot assay further confirmed the presence of the HN protein expression in the transformed lines.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Master Sains

**EXPRESI PROTEIN HAEMAGGLUTININ-NEURAMINIDASE DARI
VIRUS PENYAKIT NEWCASTLE STRAIN AF2240 UNTUK
KALUS EMBRYOGENIK *CENTELLA ASIATICA* (PEGAGA) MELALUI
KAEDAH MIKROPROJEKTIL BEDILAN YANG DIOPTIMASIKAN**

Oleh

LAI KOK SONG

June 2008

Pengerusi: Profesor Maziah Mahmood, PhD

Fakulti: Bioteknologi dan Sains Biomolekul

Centella asiatica merupakan tumbuhan ubatan tempatan yang penting. Ia tidak bertoksik, kaya dengan nilai ubatan, dan boleh bertindak sebagai calon manipulasi genetik yang baik. Malangnya, hingga kini tiada protokol transformasi yang dihasilkan dalam memaksimumkan potensi tumbuhan ini. Justeru itu, penyelidikan ini bermatlamat untuk menghasilkan protokol mikroprojektil bedilan transformasi yang cekap untuk kallus embryogenik *C. asiatica*. Tambahan pula, cubaan untuk mengekspreskan protein haemagglutinin-neuraminidase (HN) strain virus penyakit Newcastle (NDV) AF2240 dalam kallus embryogenik *C. asiatica* juga dijalankan berdasarkan sistem protokol yang dihasilkan. Protein HN boleh digunakan sebagai calon vaksin yang berpotensi untuk penyakit Newcastle (ND) dalam perternakan. Kallus embryogenik yang diinduksikan menunjukkan kehadiran lapisan matrik luaran (ECM) semasa kajian mikroskop. Protokol mikroprojektil bedilan transformasi telah dihasilkan dengan menggunakan



green fluorescent protein (GFP) sebagai sistem pelapor. Sejumlah lapan parameter yang terdiri dari jarak tisu sasaran, tekanan helium, saiz partikel emas, tekanan ruang vakum, bilangan bedilan, bahan pengikatan, masa pos-bedilan, dan kepekatan DNA plasmid telah dikenalpasti dan berjaya dioptimasikan. Berdasarkan protokol yang dihasilkan, transformasi dijalankan pada kalus embryogenik *C. asiatica* dengan menggunakan plasmid pMDC32'HN dan HBT95:sGFP(S65T)-HN. Analisis PCR genomik menunjukkan kehadiran transgen HN dalam transforman. Malangnya, tiada produk protein yang dikesan semasa SDS-PAGE dan blot western menunjukkan rendah atau ketiadaan ekspresi protein HN. Transformasi menggunakan plasmid HBT95:sGFP(S65T)-HN memberi ekspresi GFP yang rendah berbanding pada kawalan positif. Walaubagaimanapun, transkrip mRNA berjaya dikesan dalam analisis RT-PCR. Keputusan positif dari pemblotan titikan memastikan kehadiran ekspresi protein HN pada transforman.

ACKNOWLEDGEMENTS

I wish to express my deepest thanks and appreciation to my supervisor, Professor Dr. Maziah Mahmood for her guidance, time, and encouragement throughout this research project. Special thanks to my co-supervisor Professor Datin Paduka Dr. Khatijah Yusoff for her invaluable advice, support, and opportunity to carry out some stages of my research in her laboratory. My sincere appreciation is also extended to Dr. Mohd. Puad Abdullah, Dr. Janna Ong Abdullah, Dr. Parameswary Namasivayam, Associate Professor Dr. Tan Wen Siang, and Associate Professor Dr. Siti Nor Akmar Abdullah for their assistance and helpful suggestions during the research. Special appreciation goes to KUOK Foundation for providing the scholarship during my study in UPM.

I wish to thank to all my ex- and present labmates: Tee, CY, Rosli, Chew, Wai Sum, Wilson, Wai San, Noorman, Azmeer, Cecillia, Uncle Thong, Azreena, Haiza, Kah Fai, Nicole, Eddie, Erin, Hamzah, Jaffri, Rajik, Hidayah, Halimi, Hassan, Fariq, Somayeh, Nagmeh, Sara, Samira, Firoozeh, Swee Tin, Michelle, Ah Ho, Lip Nam, Wei Boon, Jason, Fui Chin, Senthil, Kie Hie and Vahid for the friendship and assistance.

Appreciation is extended to Dr. Jen Sheen (Boston, USA) for the GFP construct and Dr. Emma Knee (Ohio, USA) for the pMD C32 construct. Last but not least to my parents and brothers for their endless love, care and support.



I certify that an Examination Committee has met on 25 September 2008 to conduct the final examination the of Lai Kok Song on his Master of Science thesis entitled “Expression of Haemagglutinin-neuraminidase Envelope Protein from Newcastle Disease Virus Strain AF2240 in *Centella asiatica* (Pegaga) Embryonic Calli Through Optimized Particle Bombardment Method” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the Master of Science.

Members of the Examination Committee were as follows:

Noorjahan Banu Mohamed Alitheen, PhD

Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Puad Abdullah, PhD

Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

Ho Chai Ling, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

Salmijah Surif, PhD

Professor

Faculty of Science and Technology

Universiti Kebangsaan Malaysia

(External Examiner)

HASANAH MOHD. GHAZALI, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 27 November 2008



This thesis was submitted to Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Maziah Mahmood, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Khatijah Yusoff, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

HASANAH MOHD. GHAZALI, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 27 November 2008



DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledge. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institutions.

(LAI KOK SONG)

Date:

TABLE OF CONTENTS

	Page
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	ix
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xvii
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 Newcastle Disease	4
2.2 Newcastle Disease Virus	5
2.2.1 Classification	5
2.2.2 Virion Structure	6
2.2.3 Viral Genome Organization	9
2.2.4 Haemagglutinin-neuraminidase Protein	11
2.3 Plant Tissue Culture	14
2.3.1 <i>Centella asiatica</i>	14
2.3.2 Embryogenic Tissue	15
2.3.3 Organogenesis	17
2.4 Plant Genetic Transformation	19
2.4.1 DNA Delivery Methods	19
2.4.2 Particle Bombardment	21
2.5 Plant protein Expression System	25
2.5.1 Strategies for Protein Expression	25
2.5.2 Why Plant the Better Expression System?	28
3 MATERIALS & METHODS	32
3.1 Materials	32
3.1.1 Chemicals, Biological Reagents, Kits, Enzymes, and Markers	32
3.1.2 Virus	32
3.1.3 Bacteria	32
3.1.4 Plant Material	33
3.1.5 Plasmid Constructs	33
3.2 Methods	34
3.2.1 General Procedures	34
3.3 Plant Tissue Culture and Microscopy Studies	35
3.3.1 Preparation of Tissue Culture Media	35
3.3.2 Leaf Explant Sterilization	35
3.3.3 Calli Initiation and Maintenance	36
3.3.4 Growth Measurement of the Calli Culture	36



3.3.5	Organogenesis of <i>Centella asiatica</i> Calli Culture	37
3.3.6	Histology: Light Microscopy	37
3.3.7	Scanning Electron Microscopy (SEM)	38
3.3.8	Transmission Electron Microscopy (TEM)	39
3.4	Molecular Cloning Methods	40
3.4.1	Virus Cultivation and Purification	40
3.4.2	Haemagglutinin (HA) Test	41
3.4.3	Viral RNA Extraction	42
3.4.4	Primer Design	43
3.4.5	Reverse Transcription-Polymerase Chain Reaction (RT-PCR)	43
3.4.6	DNA Agarose Gel Purification	45
3.4.7	Isolation of DNA from Agarose Gel	45
3.4.8	Preparation of <i>Escherichia Coli</i> Competent Cells	45
3.4.9	Cloning of HN Gene into pCR [®] 8/GW/TOPO Vector	46
3.4.10	Subcloning of HN Gene into pMDC32'HN	46
3.4.11	Cloning of HN Gene into pGEM [®] -T Easy Vector	48
3.4.12	Subcloning of HN Gene into HBT95:sGFP(S65T)-NOS	48
3.4.13	Transformation of <i>Escherichia Coli</i> Competent Cells	49
3.4.14	Small Scale Plasmid Extraction	49
3.4.15	Restriction Enzyme Digestion	51
3.4.16	Polymerase Chain Reaction (PCR)	51
3.4.17	DNA Sequencing	52
3.5	Particle Bombardment Transformation Parameters Optimization	52
3.5.1	Consumables for Particle Bombardment (PDS/He 1000 Bio-Rad System)	52
3.5.2	Target Tissue Preparation for Transformation	53
3.5.3	Large Scale Plasmid Extraction	53
3.5.4	Preparation of Gold Microcarriers and Coating of Plasmid DNA onto Gold Microcarriers	53
3.5.5	Particle Bombardment	54
3.5.6	GFP Monitoring	56
3.6	Transformation of pMDC32'HN and HBT95:sGFP(S65T)-NOS'HN	56
3.7	Selection and Molecular Analyses of the Transformants	57
3.7.1	Minimal Hygromycin Concentration Determination	57
3.7.2	Selection of Putative Transformants	57
3.7.3	Plant Genomic DNA Extraction	58
3.7.4	Plant Genomic PCR Analysis	58
3.7.5	Total Plant RNA Extraction	59
3.7.6	Total Soluble Protein Extraction	60
3.7.7	Bradford Assay	60
3.7.8	SDS-Polyacrylamide Gel Electrophoresis (SDS-PAGE)	61
3.7.9	Western Blot Analysis	62

3.7.10	Dot Blot Analysis	63
4	RESULTS AND DISCUSSION	65
4.1	Plant Tissue Culture	65
4.1.1	<i>Centella asiatica</i> Callus Induction	65
4.1.2	The Growth Curve of Callus	66
4.1.3	Organogenesis of <i>Centella asiatica</i> Callus	68
4.2	Microscopy Studies	70
4.2.1	Histology: Light Microscopy	70
4.2.2	Scanning Electron Microscopy (SEM)	73
4.2.3	Transmission Electron Microscopy (TEM)	76
4.3	Minimal Hygromycin Concentration Determinations	81
4.4	Particle Bombardment Parameters Optimisation	83
4.4.1	Effect of Different Target Distance	83
4.4.2	Effect of Different Helium Pressure (psi)	84
4.4.3	Effect of Gold Particle Size	86
4.4.4	Effect of Chamber Vacuum Pressure	87
4.4.5	Effect of Bombardment Number	89
4.4.6	Effect of Spermidine and Calcium Chloride Precipitation Agents	90
4.4.7	Effect of Post-Bombardment Incubation Time	92
4.4.8	Effect of Total Plasmid DNA Concentration	93
4.5	RT-PCR and Construction of pMDC32'HN	96
4.6	Selection of the Putative Transformants	100
4.7	Genomic PCR Analyses of the Transformants	101
4.8	Protein Analysis with SDS-PAGE and Western Blotting	103
4.9	PCR and Construction of HBT:sGFP(S65T)-NOS'HN	107
4.10	Transient GFP Expression Monitoring	111
4.11	RT-PCR and Dot Blot Analysis	111
5	CONCLUSIONS	114
	REFERENCES	118
	APPENDICES	135
	LIST OF PUBLICATIONS	143
	BIODATA OF THE AUTHOR	144



LIST OF TABLES

Table		Page
3.1	Oligonucleotide primers used in amplification of the HN gene	44
3.2	The bombardment parameters were studied independently in the co-bombardment event	56

LIST OF FIGURES

Figure	Page	
2.1	Virion structure of NDV	8
2.2	TEM of NDV particles purified from allantoic fluid	8
2.3	NDV genome organization and viral transcripts	10
2.4	Schematic diagram of important features and domains of HN protein from NDV	13
3.1	Flow chart of the construction of the recombinant pMDC32'HN plasmid	47
4.1	<i>C. asiatica</i> calli derived from leaf explants	67
4.2	Growth curve measurement of <i>C. asiatica</i>	67
4.3	Organogenesis of <i>C. asiatica</i> callus	69
4.4	Light micrographs of embryogenic and non-embryogenic callus	71
4.5	SEM of embryogenic and non-embryogenic calli	74
4.6	SEM of embryogenic calli	75
4.7	TEM of embryogenic and non-embryogenic cells	77
4.8	The ultrastructure of ECM layer of the embryogenic cells	79
4.9	The hygromycin killing level for <i>C. asiatica</i> embryogenic calli	82
4.10	Effect of different target distance on transient GFP expression in <i>C. asiatica</i> embryogenic calli	85
4.11	Effect of different helium pressure (psi) on transient GFP expression in <i>C. asiatica</i> embryogenic calli	85
4.12	Effect of different gold particle size (μm) on transient GFP expression <i>C. asiatica</i> embryogenic calli	88
4.13	Effect of different chamber vacuum pressure (mm Hg) on transient GFP expression in <i>C. asiatica</i> embryogenic calli	88
4.14	Effect of different number of bombardments on transient GFP expression in <i>C. asiatica</i> embryogenic calli	91

4.15	Effect of different treatments on transient GFP expression in <i>C. asiatica</i> embryogenic calli.	91
4.16	Effect of different post bombardment time on transient GFP expression in <i>C. asiatica</i> embryogenic calli	94
4.17	Effect of different plasmid DNA concentration (μg) on transient GFP expression in <i>C. asiatica</i> embryogenic calli	94
4.18	Transient GFP expression on <i>C. asiatica</i> embryogenic calli	95
4.19	Construction of recombinant pMDC32'HN plasmid	98
4.20	Morphology of the survive transformants after 2 months subculturing in 15 mg/L hygromycin medium	102
4.21	Genomic PCR analyses of the transformants	102
4.22	SDS-12% PAGE and coomassie blue staining of total soluble protein from transformed and non-transformed lines	106
4.23	Western blot analysis of total soluble protein from transformed and non-transformed lines	106
4.24	Construction of recombinant HBT95:sGFP(S65T)-NOS'HN plasmid	109
4.25	Transient GFP expression analysis	113
4.26	RT-PCR and dot blot analysis of transformants	113

LIST OF ABBREVIATIONS

2'4 D	2,4-dichlorophenoxy acetic acid
5'-UTL	5' untranslated leader
Ads	adenine sulfate
APS	ammonium persulfate
BA	6-benzyladenine
BDMA	benzyl dimethylamine
BSA	bovine serum albumin
°C	Celsius
CaCl ₂	calcium chloride
cDNA	complementary deoxyribonucleic acid
cm ²	centimetre square
DDSA	dodecenylsuccinic anhydride
DEPC	diethyl pyrocarbonate
dH ₂ O	distilled water
ECM	extracellular matrix
EDTA	ethylenediaminetetraacetic acid
ELISA	enzyme-linked immunosorbent assay
ER	endoplasmic reticulum
F	fusion (glycoprotein)
FAA	formaldehyde-acetic acid-ethanol
GFP	green fluorescent protein
GE	gene end
g	gram
GS	gene start

h	hour
HN	haemagglutinin-neuraminidase (glycoprotein)
IBA	indole-3-butyric acid
Ig G	immunoglobulin G
kb	kilobase
kDa	kilo dalton
Kn	kinetin
kPa	kilopascal
L	litre
LB	Luria-Bertani
M	molar
mA	milliampere
MgCl ₂	magnesium chloride
mg	milligram
min	minute
mL	millilitre
mm	millimeter
mM	millimolar
mm Hg	millimetre of mercury
MNA	2-methyl-4-nitro-aniline
MS	Murashige and Skoog
mRNA	messenger ribonucleic acid
NAA	naphthalene acetic acid
NaOH	sodium hydroxide
NDV	Newcastle disease virus

ng	nanogram
Nm	nanometer
NP	nucleoprotein
P	phosphoprotein
PBS	phosphate buffered saline
PCR	polymerase chain reaction
pmol	picomol
s	second (time)
scFv	single-chain variable fragment
SCW	silicon carbide whisker
SDS-PAGE	sodium dodecyl sulphate-PAGE
SEM	scanning electron microscope
RBC	red blood cell
RE	restriction enzyme
RNA	ribonucleic acid
rpm	rotations per minute
RT	room temperature
RT-PCR	reverse transcriptase polymerase chain reaction
TAE	40 mM Tris-Cl(pH 7.4), 20mM sodium acetate, 1 mM EDTA
T-DNA	transfer deoxyribonucleic acid
TDZ	thidiazuran
TE	10 mM Tris-Cl (pH 8.0), 1 mM EDTA
TEM	transmission electron microscope
TEMED	N,N,N',N'-tetramethylethylenediamine
TPBS	Tris phosphate buffer saline

μg	microgram
μL	microlitre
μm	micrometer
μM	micro Molar
v/v	volume /volume
w/v	weight/volume

CHAPTER 1

INTRODUCTION

The term plant genetic engineering has long conveyed a highly efficient and precise process for the manipulation of plant genomes (Ow, 2007). This technology has become a versatile platform for cultivar improvement, expression and production of valuable proteins, and as well as studying gene function in plant. Recently, much effort has been channeled to develop the transformation system for medicinal and aromatic plants. In our country, *Centella asiatica* (*C. asiatica*) is an important medicinal plant that is grown for commercialization purposes. *C. asiatica* is non-toxic, easily grown, high in medicinal values, and can serve as a good platform for genetic engineering. Unfortunately, to date no efficient transformation protocol has been developed and genetic manipulation been carried out to fully exploit this essential medicinal herb. The aims of this research study are to develop an efficient particle bombardment transformation protocol for *C. asiatica* embryogenic calli and to express HN protein from Newcastle disease virus strain AF2240 based on the establish transformation protocol.

Newcastle disease (ND) is a worldwide economic problem in poultry industry caused by NDV. The continued presence of ND causes economic losses to the industry in terms of production (such as in death, loss of body weight and impaired egg production) and control (such as quarantine, mass slaughter and disinfection) (Wong, 2004). The current protocol for NDV vaccination utilizes



both inactivated and live viruses for the control of ND in poultry. Although both types of vaccination produce excellent immunity among the flocks, the drawback of these vaccines still remain the major obstacle in the industry. Live vaccine will cause disease in the presence of the complicating infections, while it is often difficult to ensure complete inactivation of the virus which could remain as risk in inactivated vaccines (Wong, 2004). Studies had shown that one of the enveloped proteins of NDV known as haemagglutinin-neuraminidase (HN) protein is the potential candidate as immunogen for the development of NDV subunit vaccine. Passive immunization with either polyclonal or monoclonal antibodies to this protein is able to confer protection in chickens against NDV challenge (Reynolds and Maraqa, 2000). Thus many attempts had been applied to express the HN protein in different systems such as baculovirus and animal systems for the production of NDV subunit vaccine. Despite the successful expression and promising immunogenicity of the HN protein, they are still too costly for commercial production. As a result, a more feasible host protein expression system is needed to ensure the continuous supply of subunit vaccine for the control of NDV.

Extensive research over the past two decades has shown that plant can provide a new platform for the expression of recombinant proteins. Its protein synthesis pathway appears to be well conserved especially in glycosylation and post-translation modification enabling them for the production of various pharmacokinetics and biological active recombinant proteins. Moreover, plant is considered to be much safer than both animals and microbes because they generally lack human pathogens, endotoxin, and oncogenic sequences

(Commandeur, 2003). This system also provides lower cost of production and rapid scale up unlike the current available systems. Hence, plant can serve as a better and more feasible alternative system for the expression of recombinant proteins.

Thus the specific objectives of this study are:

1. to study the histochemical and morphological features of the target tissue, *C. asiatica* embryogenic calli;
2. to optimize transformation conditions for particle bombardment of *C. asiatica* embryogenic calli and to construct recombinant plasmids carrying the HN gene of NDV strain AF2240;
3. to bombard, select, verify the integration of transgene using PCR and to express the HN protein of NDV strain AF2240 in *C. asiatica* embryogenic calli.

CHAPTER 2

LITERATURE REVIEW

2.1 Newcastle Disease

Newcastle disease (ND) is a worldwide economic problem in poultry industry caused by Newcastle disease virus (NDV), a prototype *Avulavirus* in the subfamily *Paramyxovirinae* (De Leeuw and Peeters, 1999). Outbreaks of ND were first reported in poultry from Java, Indonesia and Newcastle-upon-Tyne in 1926 (Bruce et al., 2000). NDV has a wide host range with 27 of the 50 orders of birds reported to be infected by the virus (Kaleta and Baldauf, 1988). The NDV isolates can be divided into three main pathotypes: lentogenic, mesogenic, and velogenic depending on the severity of disease produced by the isolate.

Mildly virulent lentogenic strain usually causes mild respiratory infection in young chicks but not in adult birds. Mesogenic strain is more virulent and can induce mild disease with mortality accruing primarily in young chickens. Meanwhile the highly virulent velogenic strain induces severe diseases and mortality to birds of all ages. Differential diagnosis of NDV involves electron microscopic identification, hemagglutination inhibition with polyclonal NDV specific antisera, use of the ELISA, oligonucleotide probes, and viral genomic RNA fingerprint analysis (Bruce et al., 2000) The current protocol for NDV vaccination utilizes both inactivated and live viruses for the control of ND in poultry. Various routes of vaccination were applied such as injection, inhalation