



UNIVERSITI PUTRA MALAYSIA

***BIODEGRADATION OF DIESEL FUEL BY TWO PSYCHROTOLERANT
STRAINS ISOLATED FROM SOUTHERN VICTORIA ISLAND,
ANTARCTICA***

NUR MUHAMAD SYAHIR BIN ABDUL HABIB

FBSB 2018 41



**BIODEGRADATION OF DIESEL FUEL BY TWO PSYCHROTOLERANT
STRAINS ISOLATED FROM SOUTHERN VICTORIA ISLAND,
ANTARCTICA**

By

NUR MUHAMAD SYAHIR BIN ABDUL HABIB

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirement for the Degree of Master of Science**

November 2017

COPYRIGHT

All materials contained within the thesis including without limitation text, logos, icons, photographs and all other artworks are copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from copyright holder. Commercial use of materials may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This thesis is dedicated to my parents.



Abstract of thesis presented to Senate of Universiti Putra Malaysia in fulfillment
of the requirements for the degree of Master of Science

**BIODEGRADATION OF DIESEL FUEL BY TWO PSYCHROTOLERANT
STRAINS ISOLATED FROM SOUTHERN VICTORIA ISLAND,
ANTARCTICA**

By

NUR MUHAMAD SYAHIR BIN ABDUL HABIB

November 2017

Chairman: Nur Adeela Binti Yasid, PhD

Faculty: Biotechnology and Biomolecular Sciences

Hydrocarbon contamination in Antarctica poses a great threat to the delicate and unique ecosystems of this continent. Bioremediation of hydrocarbon pollutants via utilisation of the indigenous hydrocarbon-degrading bacteria, has been proposed as an environmentally friendly method to clean-up contaminated soils in Antarctica. This study focused on diesel-degrading *Pseudomonas* and *Rhodococcus* species isolated from pristine soils located at the Southern Victoria Island, Antarctica. Isolates were assessed for their ability to grow on diesel as a sole carbon source on solid media at 4°C. Nine isolates showed significant growth in enriched agar after 14 days of incubation. Isolates were then screened to obtain the most promising diesel-degrading strains through colourimetric assay. Two potent isolates that possess rapid utilisation of 0.5% (v/v) diesel were selected and identified as *Pseudomonas* sp. strain ADL15 and *Rhodococcus* sp. strain ADL36. The factors that contribute to the growth of both strains were characterised initially using the conventional 'one-factor-at-a-time' approach. During this stage, the optimal condition for the growth of both ADL15 and ADL36 were at pH 7.0, 20°C, 1.0% (w/v) NaCl, and 1.0 g/L NH₄NO₃. However, strain ADL36 favoured a higher amount of diesel (2.0% (v/v)) for bacterial growth by comparison to ADL15 (1.0% (v/v)). Percentage of dodecane mineralisation was used as the mean to indicate diesel reduction through gas chromatographic analysis. While strain ADL36 showed 83.75% of dodecane mineralisation, the reduction of dodecane by ADL15 is merely at 22.39%. Response surface methodology (RSM) based on central composite design (CCD) was used to improve and optimise the effect of significant factors toward the biodegradation of diesel. RSM proved to enhance the reduction of experimented hydrocarbon (dodecane) with a 15% and 16% increment of mineralisation for isolate ADL15 (38.32%) and ADL36 (99.89%), respectively. The results also demonstrated that addition of salt to culture media was the limiting factor in hydrocarbon degradation. Whole genome sequencing showed that ADL15 and ADL36 were closely related to the *Pseudomonas fluorescens* and *Rhodococcus erythropolis* grouping, respectively. Metagenomic analyses revealed the presence of alkane hydroxylases systems which was responsible for alkane degradation in ADL36 but not in ADL15. This finding corresponds to the

gas chromatographic analysis in which ADL36 proved to be a better alkane degrader than ADL15. Detection of the complete pathway of aromatic compound degradation in the latter strain indicates a stronger inclination of the strain to utilise aromatic components in diesel as the carbon source. The presence of putative monooxygenases may also suggest that this strain may utilise specific alkane for their growth. The results from this study showed that strain ADL15 and ADL36 have an excellent potential in bioremediation of aromatics and aliphatics, respectively.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Sarjana Sains

**BIODEGRADASI MINYAK DIESEL OLEH DUA STRAIN PSIKROTOLERAN
YANG DIPENCILKAN DARIPADA PULAU VICTORIA SELATAN,
ANTARTIKA**

Oleh

NUR MUHAMAD SYAHIR BIN ABDUL HABIB

November 2017

Pengerusi : Nur Adeela Binti Yasid, PhD
Fakulti : Bioteknologi dan Sains Biomolekul

Pencemaran hidrokarbon di Antartika menyebabkan ancaman yang besar kepada ekosistem yang unik dan rapuh di benua ini. Bioremediasi yang menggunakan bakteria yang mampu menguraikan hidrokarbon, telah dicadangkan sebagai kaedah yang mesra alam bagi membersihkan kawasan tercemar di Antartika. Kajian ini memberi tumpuan kepada pengurai minyak diesel daripada spesis *Pseudomonas* dan *Rhodococcus* yang diasingkan daripada tanah yang suci yang terletak di Pulau Victoria Selatan, Antartika. Pencilan dinilai melalui keupayaan mereka untuk menumbuh dengan menggunakan diesel sebagai sumber karbon tunggal pada media pepejal pada 4°C. Sembilan pencilan menunjukkan pertumbuhan yang jelas di dalam agar yang diperkaya selepas 14 hari inkubasi. Pencilan kemudian disaring untuk mendapatkan pencilan pengurai minyak diesel yang paling bagus melalui ujian berwarna. Dua pencilan kekar yang menggunakan 0.5% (v/v) minyak diesel secara pantas telah dipilih dan dikenalpasti sebagai *Pseudomonas* sp. strain ADL15 dan *Rhodococcus* sp. strain ADL36. Faktor-faktor yang menyumbang kepada pertumbuhan kedua-dua strain dicirikan pada mulanya menggunakan pendekatan konvensional 'satu-faktor-pada-satu-masa'. Pada peringkat ini, keadaan optimum pertumbuhan ADL15 dan ADL36 berada pada pH 7.0, 20°C, 1.0% (w/v) NaCl, dan 1.0 g/L NH₄NO₃. Walau bagaimanapun, strain ADL36 menyukai jumlah minyak diesel yang lebih tinggi (2.0% (v/v)) bagi pertumbuhan bakteria berbanding ADL15 (1.0% (v/v)). Peratusan penguraian dodekana digunakan sebagai tanda bagi menunjukkan pengurangan diesel melalui analisis kromatografi gas. Walaupun strain ADL36 menunjukkan 83.75% penguraian dodekana, pengurangan dodekana oleh ADL15 adalah hanya pada 22.39%. Pengkaedahan tindakbalas permukaan (RSM) berdasarkan reka bentuk komposit pusat (CCD) digunakan untuk meningkatkan dan mengoptimumkan kesan faktor-faktor penting kearah penguraian diesel. RSM terbukti dapat meningkatkan pengurangan hidrokarbon (dodekana) yang diuji dengan peningkatan sebanyak 15% dan 16% untuk penurunan dodekana bagi pencilan ADL15 (38.32%) dan ADL36 (99.89%), masing-masing. Keputusan yang diperoleh juga menunjukkan bahawa penambahan garam ke media kultur adalah faktor yang mengurangkan penguraian hidrokarbon. Penjujukan keseluruhan genom menunjukkan bahawa ADL15 dan ADL36 mempunyai kaitan rapat dengan kelompok *Pseudomonas fluorescens* dan *Rhodococcus erythropolis*. Analisis metagenomik mendedahkan

kehadiran sistem alkane hidroksilase yang bertanggungjawab terhadap penguraian alkana di dalam ADL36 tetapi tidak di dalam ADL15. Penemuan ini bersesuaian dengan analisis kromatografi gas dimana ADL36 terbukti menjadi pengurai alkana yang lebih baik daripada ADL15. Pengesanan laluan penguraian hidrokarbon aromatik yang lengkap didalam ADL15 mungkin menunjukkan kecenderungan ADL15 yang lebih kuat untuk menggunakan bahagian aromatik didalam diesel sebagai sumber karbon. Kehadiran monooksigenase putatif mungkin juga menunjukkan bahawa strain ini menggunakan alkana yang khusus untuk pertumbuhan mereka. Keputusan daripada kajian ini menunjukkan bahawa strain ADL15 dan ADL36 mempunyai potensi yang baik didalam bioremediasi hidrokarbon aromatik dan alifatik.



ACKNOWLEDGEMENTS

BISMILLAHIRRAHMANIRRAHIM

First of all, I would like to take this opportunity to express my appreciation to every person who motivates, helps and contributes in completing this research project directly or indirectly. I would like to give my earnest gratefulness to my supervisor, Dr. Nur Adeela Yasid for her constant involvement, useful opinion and her supervision throughout the project.

Furthermore, I would also like to acknowledge Dr. Siti Aqlima Ahmad, Assoc. Prof. Dr. Mohd Yunus Shukor, Dr. Wan Lutfi Wan Johari and all the staffs in Bioremediation and general laboratory who contributed in giving brilliant suggestions and constant engagement in aiding me to complete the project. A million thanks also to Dr. Khalilah, Dr. Azham, Dr. Khalizan, Dr. Radzi, Dr. Azah, Dr. Ain, Dr. Noor Azmi, Dr. Syahida, Dr. Grace, Dr. Kenneth, Sook Yee, Basyar and Arief for their guidance and wisdom throughout this project.

Last but not least, I would like to thank my family especially to my loving parents, Abdul Habib Alapitchay and Zaharah Dahlan, and my siblings, Afzan, Aisyah, Hafiz and Bahri who had shown me their continuous support and encouragement. I would also like to appreciate all my laboratory partners and members – Motharasan, Nadzirah, Shakirah, Ain, Allamin, Hafiz, Fadhil, Umar, Aisami, Maryam, Gillian, Nadia Izzati, Aida and Fatin – and all my friends who have been my inspiration, motivation and support all these years.

Finally, special thanks are given to the Faculty of Biotechnology and Biomolecular Sciences and Faculty of Environmental Sciences for giving me a good environment and facilities. I doubt that this project will be possible without the support from the people that I have mentioned above. Thank you very much.

Syahir Habib, 2017

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Nur Adeela Yasid, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairperson)

Siti Aqlima Ahmad, PhD

Senior Lecturer
Faculty Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Mohd Yunus Shukor, PhD

Associate Professor
Faculty of Biotechnology
Universiti Putra Malaysia
(Member)

Wan Lutfi Wan Johari, PhD

Senior Lecturer
Faculty of Environmental Sciences
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writing, seminar paper, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Nur Muhamad Syahir Bin Abdul Habib, GS44253

Declaration by Members of Supervisory committee

This is to certify that:

- the research conducted and the writing of the thesis was under our supervision
- supervision of responsibilities as slated in Rule 41 in Rules 2003 (Revision 2012 – 2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Dr. Nur Adeela Yasid

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Siti Aqlima Ahmad

Signature: _____
Name of Member
of Supervisory
Committee: Associate Professor
Dr. Mohd Yunus Shukor

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Wan Lutfi Wan Johari

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENT	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xvi
 CHAPTER	
 1 INTRODUCTION	 1
 2 LITERATURE REVIEW	 3
2.1 Hydrocarbon pollution in Antarctica	3
2.2 Petroleum hydrocarbons	3
2.2.1 Composition of petroleum hydrocarbons	3
2.2.2 Diesel oil	4
2.2.3 Fate of petroleum hydrocarbons	5
2.2.3.1 Soil system	5
2.2.3.2 Aqueous and marine system	6
2.3 Petroleum hydrocarbon toxicity	6
2.4 Removal techniques of hydrocarbon	8
2.4.1 Physical and chemical methods	9
2.4.2 Biological methods	9
2.4.2.1 Biostimulation	10
2.4.2.2 Bioaugmentation	10
2.5 Microbial degradation of petroleum hydrocarbons	11
2.6 Petroleum hydrocarbons degradation pathways	13
2.6.1 Bacterial aerobic degradation	14
2.6.2 Bacterial anaerobic degradation	20
2.7 Factors affecting petroleum hydrocarbon degradation	21
2.7.1 Temperature	21
2.7.2 Oxygen availability	22
2.7.3 pH	22
2.7.4 Nutrient availability	22
2.7.5 Salinity	23
2.8 Bacterial whole genome sequencing (WGS)	23
2.8.1 First-generation sequencing	23
2.8.2 Second-generation sequencing	24
2.8.3 Illumina (Solexa) sequencer	25
2.8.3.1 Illumina MiSeq	26
 3 MATERIALS AND METHODS	 27
3.1 Chemicals and equipment	27
3.2 Preparation of culture growth medium	28

3.2.1	Nutrient agar (NA)	28
3.2.2	Nutrient broth (NB)	28
3.2.3	Bushnell Haas (BH) medium	28
3.2.4	Bushnell Haas (BH) agar with diesel	28
3.2.5	Phosphate-buffered saline (PBS)	28
3.3	Isolation and screening of diesel-degrading strains	29
3.3.1	Collection of soil samples	29
3.3.2	Strain isolation and maintenance	29
3.3.3	2,6-dichlorophenolindophenol (DCPIP) assay	31
3.4	Identification of diesel-degrading strains	32
3.4.1	Morphological characterisation	32
3.4.2	Biochemical characterisation	32
3.4.3	Molecular identification	33
3.4.3.1	Genomic extraction	33
3.4.3.2	Quantitative and qualitative assessment of DNA extract	33
3.4.3.3	Amplification of 16S rRNA gene via PCR	33
3.4.3.4	Sequence analysis	34
3.4.3.5	Phylogenetic analysis	34
3.5	Physiological characterisation	34
3.5.1	Cell suspension preparation and experimental design	35
3.5.2	Effect of substrate concentration	35
3.5.3	Effect of pH	35
3.5.4	Effect of temperature	35
3.5.5	Effect of salt concentration	36
3.5.6	Effect of nitrogen sources	36
3.5.6.1	Effect of different concentration of selected nitrogen sources	36
3.6	Optimisation by statistical approach	36
3.6.1	Plackett-Burman (PB) design	36
3.6.2	Central composite design (CCD)	37
3.7	Gas chromatographic analysis	38
3.8	Whole genome sequencing for isolated strain	39
3.8.1	Preparation of genomic DNA template	39
3.8.2	Library preparation	39
3.8.2.1	Tagmentation of gDNA and amplification	40
3.8.2.2	Library cleanup and normalisation	40
3.8.3	Cluster generation and <i>de novo</i> sequencing	40
3.8.4	Data analysis	40
3.8.4.1	FastQ files analysis	41
3.8.4.2	Reads assembly, annotation and analysis	41
4	RESULT AND DISCUSSION	42
4.1	Isolation and screening of diesel-degrading strains	42
4.2	Identification of diesel-degrading strains	46
4.2.1	Morphological characterisation	46
4.2.2	Molecular identification	49

4.2.2.1	Genomic extraction	49
4.2.2.2	Isolation of 16S rRNA gene sequence	50
4.2.3	16S rRNA sequence analysis	52
4.2.4	Phylogenetic analysis	55
4.3	Physiological characterisation	58
4.3.1	Effect of substrate concentration	58
4.3.2	Effect of pH	60
4.3.3	Effect of nitrogen sources	62
4.3.3.1	Effect of different concentration of NH_4NO_3	63
4.3.4	Effect of salinity	64
4.3.5	Effect of temperature	66
4.4	Evaluation and assessment of residual hydrocarbon components	68
4.5	Statistical optimisation by response surface methodology (RSM)	71
4.5.1	Plackett-Burman (PB) design	71
4.5.1.1	Viable plate count (ADL15)	71
4.5.1.2	Viable plate count (ADL36)	73
4.5.2	Central composite design (CCD)	75
4.5.2.1	Justification of Central composite design (CCD)	76
4.5.2.2	<i>Pseudomonas</i> sp. ADL15	76
4.5.2.3	<i>Rhodococcus</i> sp. ADL36	80
4.5.3	Model prediction and validation	84
4.5.3.1	<i>Pseudomonas</i> sp. ADL15	84
4.5.3.2	<i>Rhodococcus</i> sp. ADL36	86
4.5.4	Verification of optimisation experiment	89
4.6	Bacterial whole genome sequencing	90
4.6.1	Preparation of genomic DNA template	91
4.6.2	Validation of gDNA library and sequencing run	91
4.6.2.1	Quantitative and qualitative validation	91
4.6.3	Data analysis	93
4.6.3.1	MiSeq run analysis	93
4.6.3.2	FastQ files analysis	93
4.6.4	Reads assembly, annotation and analysis	94
4.6.4.1	<i>Pseudomonas</i> sp. ADL15	96
4.6.4.2	<i>Rhodococcus</i> sp. ADL36	100
4.7	Summary on the utilisation of hydrocarbons by ADL15 and ADL36	104
5	CONCLUSIONS AND RECOMMENDATION	106
	REFERENCES	108
	BIODATA OF STUDENT	136
	LIST OF PUBLICATIONS	137

LIST OF TABLES

Table	Page
2.1 Toxic effects of different hydrocarbons on human health	8
2.2 Toxic effects of different hydrocarbons on animal and plants	8
2.3 List of several Antarctic hydrocarbon-degrading bacteria	12
2.4 List of strain used for plasmid-related hydrocarbon degradation	13
3.1 List of chemicals	27
3.2 List of equipment	27
3.3 List of soil samples with exact sampling location and description	29
3.4 Experimental values for variables tested on bacterial growth	37
3.5 Experimental design for independent factors on bacterial growth	37
3.6 Experimental values selected for CCD optimisation	37
3.7 Experimental CCD on dodecane mineralisation	38
3.8 Index adaptor used for gDNA sample from isolate 15 and 36	40
4.1 List of pure bacterial isolates with growth on different media	43
4.2 Purity and genomic DNA concentration of ADL15 and ADL36	50
4.3 PB experimental design matrix with levels of bacterial growth	72
4.4 Analysis of variance (ANOVA) for bacterial growth of ADL15	72
4.5 PB experimental design matrix with levels of bacterial growth	73
4.6 Analysis of variance (ANOVA) for bacterial growth of ADL36	74
4.7 Experimental CCD on dodecane degradation by ADL15	77
4.8 ANOVA for dodecane mineralisation by ADL15 in CCD	78
4.9 Experimental CCD on dodecane degradation by ADL36	81
4.10 ANOVA for dodecane mineralisation by ADL36 in CCD	82
4.11 Suggestions for maximum dodecane mineralisation by ADL15	84
4.12 Suggestions for maximum dodecane mineralisation by ADL36	86
4.13 Differences of optimised state of dodecane mineralisation by ADL15 from different optimisation method	89
4.14 Differences of optimised state of dodecane mineralisation by ADL36 from different optimisation method	89
4.15 Purity and gDNA concentration of ADL15 and ADL36	91
4.16 The final Qubit concentration and average library size obtained	92
4.17 The summary of genomic analysis with features of both strains	96
4.18 General characteristics of the genomes of <i>P. fluorescens</i> strains Pf0-1, SBW25 and <i>Pseudomonas</i> sp. strain ADL15 and GM60	98
4.19 General characteristics of the genomes of <i>R. erythropolis</i> strains PR4 and CCM2925 along with ADL36	102

LIST OF FIGURES

Figure		Page
2.1	Examples of aliphatic and aromatic hydrocarbons	4
2.2	Peripheral pathways of alkane mineralisation	15
2.3	The initial attack on <i>n</i> -alkane and aromatic compound benzene by oxygenases	18
2.4	The schematic illustration for the divergent pathways for aerobic degradation of toluene	20
4.1	The growth of ADL15 and ADL36 on BH agar added with 0.5% (v/v) diesel	42
4.2	The structural changes of DCPIP redox indicator from blue (oxidised) to colourless (reduced)	44
4.3	The DCPIP assay for primary screening	45
4.4	The DCPIP assay for secondary screening	45
4.5	The results of colourimetric assay on discolouration of DCPIP with different hydrocarbons	46
4.6	The gram staining of a 24 hours culture of isolate ADL15 and ADL36 grown on nutrient agar	46
4.7	The oxidase test results on ADL15 and ADL36	48
4.8	The catalase test results on ADL15 and ADL36	48
4.9	Gel image of genomic extracts strain ADL15 and ADL36 on 1.0% (w/v) agarose gel	49
4.10	Gel image of PCR products for ADL15 and ADL36 on 1.0% (w/v) agarose gel	51
4.11	The accession number of 16S rRNA gene sequence of isolate ADL15 as deposited in GenBank	53
4.12	The accession number of 16S rRNA gene sequence of isolate ADL36 as deposited in GenBank.	54
4.13	Neighbour-joining method cladogram showing the phylogenetic relationship between strain ADL15 and other 20 closely related reference microorganisms based on 16S rRNA gene sequence	56
4.14	Neighbour-joining method cladogram showing the phylogenetic relationship between strain ADL36 and other 20 closely related reference microorganisms based on 16S rRNA gene sequence	57
4.15	The effect of substrate concentration on bacterial growth of ADL15	59

4.16	The effect of substrate concentration on bacterial growth of ADL36	60
4.17	The effect of pH on bacterial growth of ADL15 using three overlapping buffers	61
4.18	The effect of pH on bacterial growth of ADL36 using three overlapping buffers	61
4.19	The effect of different inorganic nitrogen source on bacterial growth of ADL15 and ADL36	62
4.20	The effect of different concentration of NH_4NO_3 on growth of ADL15	63
4.21	The effect of different concentration of NH_4NO_3 on growth of ADL36	63
4.22	The effect of salt (NaCl) on bacterial growth of ADL15	65
4.23	The effect of salt (NaCl) on bacterial growth of ADL36	65
4.24	The effect of temperature on bacterial growth of ADL15	67
4.25	The effect of temperature on bacterial growth of ADL36	67
4.26	GC-FID profile of hydrocarbons obtained from an optimised ADL15 culture at 0 day and after 7 days of incubation	69
4.27	GC-FID profile of hydrocarbons obtained from an optimised ADL36 culture at 0 day and after 7 days of incubation	70
4.28	The 3D response surface plots showing the effect of interactions between significant factors of ADL15	80
4.29	The 3D response surface plots showing the effect of interactions between significant factors of ADL36	83
4.30	GC-FID profile of hydrocarbons obtained from RSM-optimised ADL15 culture at 0 (top) and after 7 (bottom) days of incubation	85
4.31	The relationship between the dodecane mineralisation and bacterial growth of ADL15	86
4.32	GC-FID profile of hydrocarbons obtained from RSM-optimised ADL36 culture at 0 (top) and after 7 (bottom) days of incubation	87
4.33	The relationship between the dodecane mineralisation and bacterial growth of ADL36	88
4.34	The schematic diagram of bacterial whole genome sequencing	90
4.35	The Krona chart showing distribution of ADL15 sequencing reads based on metagenomics at the domain and species level	97
4.36	The Krona chart showing distribution of ADL36 sequencing reads based on metagenomics at the domain and species level	101

LIST OF ABBREVIATIONS

%	Percent
% (v/v)	Percent concentration volume / volume
% (w/v)	Percent concentration weight / volume
°C	Degree celsius
µl	Microlitre
µm	Micrometre
x g	Relative centrifugal force
bp	Base pair
CCD	Central composite design
CFU	Colony forming unit
DNA	Deoxyribonucleic acid
dNTP	Deoxyribonucleotide phosphate
EDTA	Ethylenediaminetetraacetic acid
EtBr	Ethidium bromide
et al.,	And friends
g	Gram
GC-FID	Gas chromatography- flame ionisation detector
gDNA	Genomic DNA
g/L	Gram per Litre
HCl	Hydrochloric acid
K	Kelvin
kb	Kilobase
L	Litre
M	Molar
min	Minute
mg/ml	Milligram per millilitre
mM	Milimolar
mm	Millimetre
ng/µl	Nanogram per microlitre
nM	Nanomolar
nm	Nanometer
OD	Optical density
OFAT	One-factor-at-a-time
PAHs	Polycyclic aromatic hydrocarbons
RNA	Ribonucleic acid
rpm	Revolution per minute
rRNA	Ribosomal RNA
RSM	Response surface methodology
sp.	Species (singular)
TAE	Tris-acetate-EDTA

CHAPTER 1

INTRODUCTION

Oil and its refined products represent a significant fraction of the pollution found in the Antarctic region, an area considered as the most pristine in the world (Raymond et al., 2017). The occurrences of pollution are more clustered near former military and industrial spots, scientific research stations, rural communities, and remote airfields, while recent spills and discharges tend to be linked with resource development and mishaps in transportation (Aislabie et al., 2004). Fuel spills are recognised as a potential threat as contamination can cause risks for humans and other living organisms if oil and fuel reach groundwater reservoirs and water bodies (Wang and Bartha, 1990; Jesus et al., 2015). Hydrocarbon contamination in these ecosystems is perceived as damaging as they are more sensitive, being profoundly adapted to extreme conditions (McDonald and Knox, 2014; Yang et al., 2009). Besides, due to the slow natural attenuation rates in cold climates, hydrocarbons can persist for longer periods of time than their temperate counterparts causing a stunted ecosystem recovery (Snape et al., 2008). Although the crude oil extraction from the polar region is declining and a strict legislation such as Antarctic Treaty was introduced, contamination of hydrocarbons in the Antarctic region may still be introduced by the booming numbers of tourists during the austral summer (November to March) period. According to the recent tourism statistics recorded by the International Association of Antarctica Tour Operators (IAATO), there is a 16% increase in the number of tourists landed on Antarctic from the 2015-2016 to 2016-2017 tour (IAATO 2016, 2017). While tourism cannot be solely blamed for their chances of introducing contamination, any small contamination can cause great risks to the environment for a remote and almost pristine land such as Antarctica.

Soil remediation in Antarctica is driven by several critical factors such as cost, strict environmental policy, and remediation constraints (Filler et al., 2006). Attempts to clean-up Antarctic polluted sites using both physical and chemical methods have been done but considered as a minor success. Several methods which highly practical in temperate environments such as thermal incineration, is banned from the Antarctic environment while soil excavation and removal of contaminated soils are often impractical, for the reason of high cost and risks of further damage from excavation (Snape et al., 2008). Bioremediation is widely proposed to remove pollutants from the contaminated Antarctic environment due to the increased interest in using the eco-friendly method as a process of remediating diesel fuel polluted sites (Aislabie et al., 2006; Jesus et al., 2015; Rayner et al., 2007). Bioremediation aids remediation activities being carried out either near or on site, which can be appealing in an isolated contamination spot. However, the effectiveness of this approach depends on strong limitations in temperature, bioavailability, oxygen, toxicity, and soil freeze-thaw cycle (Yang et al., 2009; Delille and Coulon, 2008). Among the factors, temperature plays a significant role in determining the rate and degree of microbial hydrocarbon biodegradation while affecting the volatilisation and viscosity of hydrocarbons (Delille and Coulon, 2008).

Biodegradation of varied components of hydrocarbons at low temperatures in Antarctic soils (Baraniecki et al., 2002; Bej et al., 2000) has been reported and is a result of the degradation capacity of indigenous cold-adapted microorganisms. Cold-adapted microorganisms are able to grow at temperatures around 0°C and have adapted their metabolism to function optimally at low temperatures. These microorganisms play a substantial role in the *in situ* biodegradation of hydrocarbons in cold environments, where ambient summer temperatures often correspond with their growth temperature range. As the Antarctic Treaty prohibits the introduction of non-native organisms, microbes that are indigenous to the Antarctic soil were required for the application of bioremediation (Aislabie et al., 2000). Among microbes, bacterial species play a key role in degrading hydrocarbon pollutants. A large number of hydrocarbon-degrading bacteria from cold soils have been identified, including representatives of gram-negative and gram-positive genera (Aislabie et al., 2000; Ruberto et al., 2005; Shukor et al., 2009). Although large numbers of hydrocarbon-degrading bacteria were isolated from contaminated soils, Margesin et al. (2003) and Stallwood et al. (2005) have observed the occurrence of bacterial species with hydrocarbon-degradative ability in the pristine soil. The addition of indigenous bacteria isolated from Antarctic pristine soil that possess a high competency to degrade diesel fuel may speed up the mineralisation process of petroleum hydrocarbons by several folds when favourable condition is maintained and the regulation of genetic diversity in the bacteria is acknowledged.

Thus, a study was carried out with the following objectives:

1. To isolate and identify bacterial species with hydrocarbon-degrading capacity from the Antarctic pristine soil.
2. To determine the optimum condition for bacterial growth and dodecane mineralisation for isolated strain using statistical analysis and model prediction.
3. To observe the residual hydrocarbon compounds in culture-optimised state qualitatively using gas chromatographic analysis.
4. To analyse the responsible alkane pathways, hydrocarbon-degrading enzymes and their respective genes through bacterial whole genome sequencing.

REFERENCES

- Abdel-Shafy, H. I., & Mansour, M. S. M. (2016). A review on polycyclic aromatic hydrocarbons: source, environmental impact, effect on human health and remediation. *Egyptian Journal of Petroleum*, 25(1), 107–123.
- Abd-Elsalam, K. A. (2003). Bioinformatic tools and guideline for PCR primer design. *African Journal of Biotechnology*, 2(5), 91–95.
- Abedon, S. T. (2012). Bacterial 'immunity' against bacteriophages. *Bacteriophage*, 2(1), 50-54.
- ACGIH (American Conference of Governmental Industrial Hygienists). (2005). Polycyclic aromatics hydrocarbons (PAHs) biologic exposure indices (BEI) Cincinnati.
- Aggarwal, R. K., Dawar, C., Phanindranath, R., Mutnuri, L., & Dayal, M. (2016). Draft genome sequence of a versatile hydrocarbon-degrading bacterium, *Rhodococcus pyridivorans* strain KG-16, collected from oil fields in India. *Genome Announcements*, 4(1), 1704–1715.
- Aghaie, E., Pazouki, M., Hosseini, M. R., Ranjbar, M., & Ghavipanjeh, F. (2009). response surface methodology (RSM) analysis of organic acid production for kaolin beneficiation by *Aspergillus niger*. *Chemical Engineering Journal*, 147, 245-251.
- Aguilera, F., Mendez, J., Pasaro, E., & Laffon, B. (2010). Review on the effects of exposure to spilled oils on human health. *Journal of Applied Toxicology*, 30(4), 291–301.
- Aislabie, J., McLeod, M., & Fraser, R. (1998). Potential of biodegradation of hydrocarbons in soil from the Ross Dependency, Antarctica. *Applied Microbiology and Biotechnology*, 49, 210–214.
- Aislabie, J., Foght, J., & Saul, D. (2000). Aromatic hydrocarbon-degrading bacteria from soil near Scott Base, Antarctica. *Polar Biology*, 23, 183–188.
- Aislabie, J., Balks, M., Foght, J., & Waterhouse, E. (2004). Hydrocarbon spills on Antarctic soils: effects and management. *Environmental Science and Technology*, 38(5), 1266–1274.
- Aislabie, J., Saul, D., & Foght, J. (2006). Bioremediation of hydrocarbon-contaminated polar soils. *Extremophiles*, 10, 171–179.
- Akpe, A. R., Ekundayo, A. O., & Esumeh, F. I. (2013). Degradation of crude oil by bacteria: a role for plasmid borne genes. *Global Journal of Science Frontier Research Biological Science*, 13(6), 1-7.
- Alexander, M. (1999). *Biodegradation and Bioremediation*. 2nd ed. New York: Academic Press.

- Alkan, C., Sajjadian, S., & Eichler, E. E. (2011). Limitations of next-generation genome sequence assembly. *Nature Methods*, 8(1), 61-65.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lioman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, 215, 403-410.
- Alvarez, L. M. M., Lo Balbo, A., Mac Cormack, W. P., & Ruberto, L. A. M. (2015). Bioremediation of a petroleum hydrocarbon-contaminated Antarctic soil: optimisation of a biostimulation strategy using response surface methodology (RSM). *Cold Regions Science and Technology*, 119, 61-67.
- Anderson, M. J., & Whitcomb, P. J. (2004). RSM Simplified: Optimizing Processes using response surface methods for design of experiments. *Business and Economics*, CRC Press, 304 pp.
- Ansorge, W., Sproat, B. S., Stegemann, J., & Schwager, C. (1986). A non-radioactive automated method for DNA sequence determination. *Journal of Biochemical and Biophysical Methods*, 13(6), 315-323.
- Ansorge, W., Sproat, B., Stegemann, J., Schwager, C., & Zenke, M. (1987). Automated DNA sequencing: ultrasensitive detection of fluorescent bands during electrophoresis. *Nucleic Acids Research*, 15(11), 4593-4602.
- Armstrong, B. G., Hutchinson, E., Unwin, J., & Fletcher, T. (2004). Lung cancer risk after exposure to polycyclic aromatic hydrocarbons; a review and meta-analysis. *Environmental Health Perspectives*, 112(9), 970-978.
- Atlas, R. M. (1991). Microbial hydrocarbon degradation - Bioremediation of oil spills. *Journal of Chemical Technology and Biotechnology*, 52, 149-156.
- Atlas, R. M. (1986). Fate of petroleum pollutants in Arctic ecosystems. *Water Science and Technology*, 18(2), 59-67.
- Atlas, R. M., & Bartha, R. (1998). *Microbial ecology: fundamentals and applications*. Benjamin/Cummings Publishing Company, pp 281-324.
- Atlas, R., & Bragg, J. (2009). Bioremediation of marine oil spills: when and when not - the Exxon Valdez experience. *Microbial Biotechnology*, 2(2), 213-221.
- ATCM (1998). *Final report of the twenty-second Antarctic Treaty Consultative Meeting*, Antarctic Treaty Consultative Meeting: Tromso, Norway.
- ATSDR (Agency for Toxic Substances and Disease Registry) (1995). Toxicological profile for polycyclic aromatic hydrocarbons (PAHs). Atlanta, GA, US Department of Health and Human Services, Public Health Service.
- ATSDR (Agency for Toxic Substances and Disease Registry) (1999). Toxicological profile for total petroleum hydrocarbons (TPH). Atlanta, GA, US Department of Health and Human Services, Public Health Service.

- Austin, R. N., Chang, H-K., Zylstra, G. J., & Groves, J. T. (2000). The non-heme diiron alkane monooxygenase of *Pseudomonas oleovorans* (AlkB) hydroxylates via a substrate radical intermediate. *Journal of American Chemical Society*, 122, 11747–11748.
- Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., Formsma, K., et al. (2008). The RAST server: rapid annotations using subsystems technology. *BMC Genomics*, 9,75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
- Bader, R. F. W., Tang, T. H., Tal, Y., & Biegler-Koenig, F. W. (1982). Molecular structure and its change: hydrocarbons. *Journal of the American Chemical Society*, 104(4), 940-945. doi: 10.1021/ja00368a003.
- Balasubramanian, S. (2011). Sequencing nucleic acids: from chemistry to medicine. *Chem Commun (Camb)*, 47(26), 7281–7286.
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M., et al. (2012). SPAdes: a new genome assembly algorithm and its application to single cell sequencing. *Journal of Computational Biology*, 19(5), 455-477. doi: 10.1089/cmb2012.0021..
- Baraniecki, C. A., Aislabie, J., & Foght, J. M. (2002). Characterization of *Sphingomonas* sp. Ant 17, an aromatic hydrocarbon-degrading bacterium isolated from Antarctic soil. *Microbial Ecology*, 43, 22–54.
- Baş, D., & Boyaci, I. H. (2007). Modelling and optimization I: usability of response surface methodology. *Journal of Food Engineering*, 78(3), 836-845. doi: 10.1016/j.foodeng.2005.11.024.
- Bej, A. K., Saul, D., & Aislabie, J. (2000). Cold-tolerant alkane-degrading *Rhodococcus* species from Antarctica. *Polar Biology*, 23, 100–105.
- Bell, K. S., Philip, J. C., Aw, D. W., & Christofi, N. (1998). The genus *Rhodococcus*. *Journal of Applied Microbiology*, 85, 195-210. <https://doi.org/10.1046/j.1365-2672.1998.00525.x>.
- Bosset, I. & Bartha, R. (1984). The fate of petroleum in soil ecosystems. In Atlas, R. M. (Ed), *Petroleum Microbiology* (pp. 435-476). New York, NY: Macmillan Publishing Company.
- Belhaj, A., Desnoues, N., & Elmerich, C. (2002). Alkane biodegradation in *Pseudomonas aeruginosa* strains isolated from a polluted zone: identification of *alkB* and *alkB*-related genes. *Research in Microbiology*, 153(6), 339–344.
- Bentley, D. R. (2006). Whole genome re-sequencing. *Current Opinion in Genetics and Development*, 16(6), 545-552.

- Bentley, D. R., Balasubramanian, S., Swerdlow, H. P., Smith, G. P., Milton, J., Brown, C. G., Hall, K. P., et al. (2008). Accurate whole human genome sequencing using reversible terminator chemistry. *Nature*, 456(7218), 53-59. doi: 10.1038/nature07517.
- Bockheim, J. G., & McLeod, M. (2006). Soil formation in Wright Valley, Antarctica since the late Neogene. *Geoderma*, 137(1), 109–116. <https://doi.org/10.1016/j.geoderma.2006.08.028>
- Box, G. E. P., & Wilson, K. B. (1951). On the Experimental Attainment of Optimum Conditions. *Journal of the Royal Statistical Society. Series B (Methodological)*, 13(1), 1–45.
- Brennerova, M. V., Josefiova, J., Brenner, V., Pieper, D. H., & Junca, H. (2009). Metagenomics reveals diversity and abundance of *meta*-cleavage pathways in microbial communities from soil highly contaminated with jet fuel under air-sparging bioremediation. *Environmental Microbiology*, 11(9), 2216–2227.
- Buchanan, B. B. (1991). Regulation of CO₂ assimilation in oxygenic photosynthesis: the ferredoxin thioredoxin system: perspective on its discovery, present status, and future development. *Archive of Biochemistry and Biophysics*, 288, 1-9.
- Bushnell, L. D., & Haas, H. F. (1941). The utilization of certain hydrocarbons by microorganisms. *Journal of Bacteriology*, 41(5), 653-673.
- Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S. M., et al. (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *The ISME Journal*, 6(8), 1621-1624.
- Cappuccino, J. G., & Sherman, N. (2005). *Microbiology: a laboratory manual* (7th ed.) San Francisco: Pearson/Benjamin Cummings
- Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madeen, T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*, 10(421). doi: 10.1186/1471-2105-10-421.
- Camenzuli, D., Freidman, B. L., Statham, T. M., Mumford, K. A., & Gore, D. B. (2013). On-site and *in situ* remediation technologies applicable to petroleum hydrocarbon contaminated sites in the Antarctic and Arctic. *Polar Research*, 33, doi: <http://dx.doi.org/10.3402/polar.v33.21522>.
- Camenzuli, D. & Freidman, B. L. (2015). On-site and *in situ* remediation technologies applicable to petroleum hydrocarbon contaminated sites in the Antarctic and Arctic. *Polar Research*, 34, <http://dx.doi.org/10.3402/polar.v34.24492>.
- Campbell, I. B., Claridge, G. G. C., Campbell, D. I., & Balks, M. R. (1998). The soil environments of the Mcmurdo Dry Valleys, Antarctica, in ecosystem dynamics in a polar desert: the Mcmurdo Dry Valleys, Antarctica. In Prisco, J. C., *American Geophysical Union*, Washington , D. C. doi: 10.1029/AR072p0297.

- Canard, B., & Sarfati, R. S. (1994). DNA polymerase fluorescent substrates with reversible 3'-tags. *Gene*, 148(1), 1-6.
- Carmichael, L. M., & Pfaender, F. K. (1997). The effect of inorganic and organic supplements on the microbial degradation of phenanthrene and pyrene in soils. *Biodegradation*, 8(1), 1-13.
- Case, R. J., Boucher, Y., Dahllöf, I., Homstrom, C., Doolittle, W. F., & Kjelleberg, S. (2007). Use of 16S rRNA and *ropB* genes as molecular markers for microbial ecology studies. *Applied and Environmental Microbiology*, 73(1), 278-288.
- Chaineau, C. H., Morel, J., Dupont, J., Bury, E., & Oudot, J. (1999). Comparison of the fuel oil biodegradation potential of hydrocarbon-assimilating microorganisms isolated from a temperate agricultural soil. *Science of Total Environment*, 227(2-3), 237-247.
- Chaineau, C. H., Rougeux, G., Yepremian, C., & Oudot, J. (2005). Effect of nutrient concentration on the biodegradation of crude oil and associated microbial populations in the soil. *Soil Biology and Biochemistry*, 20, 1-8.
- Chuvilin, E. M., Naletova, N. S., Miklyaeva, E. C., Kozlova, E. V., & Instanes, A. (2001). Factors affecting spreadability and transportation of oil in regions of frozen ground. *Polar Record*, 37, 229-238.
- Compeau, G., Al-achi, B. J., Platsouka, E., & Levy, S. B. (1988). Survival of rifampin-resistant mutants of *Pseudomonas fluorescens* and *Pseudomonas putida* in soil systems. *Applied and Environmental Microbiology*, 54(10), 2432-2438.
- Cookson, J. T. (1995). Bioremediation engineering design and application. McGraw Hill, New York.
- Cooley, R. B., Dubbels, B. L., Sayavedra-Soto, L. A., Bottomley, J. & Arp, D. J. (2009). Kinetic characterization of the soluble butane monooxygenase from *Thauera butanivorans*, formerly '*Pseudomonas butanovora*'. *Microbiology*, 155(6), 2086-2096.
- Coulon, F., & Delille, D. (2003). Effects of biostimulation on growth of indigenous bacteria in sub-Antarctic soil contaminated with oil hydrocarbons. *Oil and Gas Science and Technology*, 58(4), 469-479.
- Coulon, F., Pelletier, E., St. Louis, R., Gourhant, L. & Delille, D. (2004). Degradation of petroleum hydrocarbons in two sub-Antarctic soils: influence of an oleophilic fertilizer. *Environmental Toxicology and Chemistry*, 23(8), 1893-1901.
- Cripps, G. C., & Priddle, J. (1991). Hydrocarbons in the Antarctic marine environment. *Antarctic Science*, 3(3), 233-250. doi: 10.1017/S0954102091000299
- Curtosi, A., Pelletier, E., Vodopivec, C. L., & Mac Cormack, W. P. (2007). Polycyclic aromatic hydrocarbons in soil and surface marine sediments near Jubany Station (Antarctica) Role of permafrost as a low-permeability barrier. *Science of the Total Environment*, 383, 193-204.

- Czitrom, V. (1999). One-Factor-at-a-Time versus Designed Experiments. *The American Statistician*, 53(2), 126–131. <https://doi.org/10.1080/00031305.1999.10474445>
- Das, N. & Chandran, P. (2011). Microbial degradation of petroleum hydrocarbon contaminants: an overview. *Biotechnology Research International*, doi: 10.4061/2011/941810.
- Date, A. W. (2011). *Analytical Combustion: With Thermodynamics, Chemical Kinetics and Mass Transfer (Google eBook)*. Cambridge University Press. ISBN 1-107-00286-9.
- Dawson, R. M. C., Elliott, D. C., Elliott, W. H., & Jones, K. M. (1986). Data for biochemical research. *Oxford Science*, Oxford.
- Dear, P. H. (2005). Genome Mapping. eLS. doi: 10.1038/npg.els.0005353.
- de Carvalho, C. C.R., & da Fonseca, M. M. R. (2005). Degradation of hydrocarbons and alcohols at different temperatures and salinities by *Rhodococcus erythropolis* DCL14. *FEMS Microbiology Ecology*, 51(3), 389–399.
- de Carvalho, C. C. R., da Cruz, A. A. R. L., Pons, M-N., Pinheiro, H. M. R. V., Cabral, J. M. S., da Fonseca, M. M. R., Ferreira, B. S. & Fernandes, P. (2004). *Mycobacterium* sp., *Rhodococcus erythropolis*, and *Pseudomonas putida* behaviour in the presence of organic solvents. *Microscopy Research and Technique*, 64(3), 215–222.
- Decker, A., & Solomon, E. I. (2005). Dioxygen activation by copper, heme and non-heme iron enzymes: comparison of electronic structure and reactivities. *Current Opinion in Chemical Biology*, 9(2), 152–163.
- De Domenico, M., Lo Giudice, A., Michaud, L., Saitta, M., & Bruni, V. (2004). Diesel and PCB-degrading psychrotrophic bacteria isolated from Antarctic seawaters (Terra Nova Bay, Ross Sea). *Polar Research*, 23(2), 141–146.
- Delille, D., & Coulon, F. (2008). Comparative mecosystem study of biostimulation efficiency in two different oil-amended sub-Antarctic soils. *Microbial Ecology*, 56(2), 243–252.
- Delille, D., Pelletier, E., Delille, B. & Coulon, F. (2003). Effect of nutrient enrichments on the bacterial assemblage of Antarctic soils contaminated by diesel or crude oil. *Polar Research*, 39, 1–10.
- Dennett, G. V., & Blamey, J. M. (2016). A new thermophilic nitrilase from an Antarctic hyperthermophilic microorganism. *Frontiers in Bioengineering and Biotechnology*, 4(5). doi: 10.3389/fbioe.2016.00005.
- Desjardins, P., & Conklin, C. (2010). NanoDrop microvolume quantitation of nucleic acids. *Journal of Visualized Experiments*, 45, e2565, 1–4.

- Dias, R. L., Ruberto, L., Hernandez, E., Vazquez, S. C., Lo Balbo, A., Del Panno, M. T., et al. (2012). Bioremediation of an aged diesel oil-contaminated Antarctic soil: evaluation of the "on site" biostimulation strategy using different nutrient sources. *International Biodeterioration and Biodegradation*, 75, 96–103.
- Diks, R. M. M., Ottengraf, S. P. P., & van den Oever, A. H. C. (1994). The influence of NaCl on the degradation rate of dichloroethane by *Hyphomicrobium* sp. *Biodegradation*, 5, 129-141.
- Dojka, M. A., Hugenholtz, P., Haack, S. K., & Norman, R. P. (1998). Microbial diversity in a hydrocarbon- and chlorinated-solvent-contaminated aquifer undergoing intrinsic bioremediation. *Applied and Environmental Microbiology*, 64(10), 3869–3877.
- Donkor, E. S., Dayie, N. T. K. D., & Adiku, T. K. (2014). Bioinformatics with basic local alignment search tool (BLAST) and fast alignment (FASTA). *Journal of Bioinformatics and Sequence Analysis*, 6(1), 1–6.
- Durán, N., & Esposito, E. (2000). Potential applications of oxidative enzymes and phenoloxidase-like compounds in wastewater and soil treatment: a review. *Applied Catalysis B: Environmental*, 28, 83-99.
- Eddy, S. R. (1998). Profile Hidden Markov Models. *Bioinformatics*, 14, 755-763.
- Emtiazi, G., Mridamadian, S., & Habibi, M. H. (2005). Instability of petroleum oil degradation by induction of mutation in 45kb and 60kb plasmids. *International Journal of Environmental Studies*, 62(4), 467–472.
- Engelbrektson, A., Kunin, V., Wrighton, K. C., Zvenigorodsky, N., Chen, F., Ochman, H., & Hugenholtz, P. (2010). Experimental factors affecting PCR-based estimates of microbial species richness and evenness. *The ISME Journal*, 4(5), 642–647. <https://doi.org/10.1038/ismej.2009.153>
- Eweis, J. B., Ergas, S. J., Chang, D. P. Y., & Schroeder, E. D. (1998). *Bioremediation Principles*, McGraw-Hill, Boston.
- Fasan, R., Mehareenna, Y. T., Snow, C. D., Poulos, T. L., & Arnold, F. H. (2008). Evolutionary history of a specialized P450 propane monooxygenase. *Journal of Molecular Biology*, 383(5), 1069-1080.
- Fedorco, M., Romieu, A., Williams, S., Lawrence, I., & Turcatti, G. (2006). BTA, a novel reagent for DNA attachment on glass and efficient generation of solid-phase amplified DNA colonies. *Nucleic Acids Research*, 34(3).
- Felsenstein, J. (1985). Confidence limits on the phylogenies: an approach using the bootstrap. *Evolution*, 39(4), 783-791.

- Feng, L., Wang, W., Cheng, J., Ren, Y., Zhao, G., Gao, C., et al. (2007). Genome and proteome of long-chain alkane degrading *Geobacillus thermodenitrificans* NG80-2 isolated from a deep-subsurface oil reservoir. *Proceedings of the National Academy of Sciences of USA*, 104(13), 5602-5607.
- Ferguson, S. H., Franzmann, P. D., Revill, A. T., Snape, I., & Rayner, J. L. (2003). The effects of nitrogen and water on mineralisation of hydrocarbons in diesel-contaminated terrestrial Antarctic soils. *Cold Regions Science and Technology*, 37, 197-212.
- Filler, D. M., Reynolds, C. M., Snape, I., Dagulis, A. J., Barnes, D. L., & Williams, P. J. (2006). Advances in engineered remediation for use in the Arctic and Antarctic. *Polar Record*, 42, 111-120.
- Fleischmann, R., Adams, M., White, O., Clayton, R., Kirkness, E., Kerlavage, A., Bult, C., et al. (1995). Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science*, 269(5223), 496-512. doi: 10.1126/science.7542800.
- Fletcher, G. L., Hew, C. L., & Davies, P. L. (2001). Antifreeze proteins of teleost fishes. *Annual Review of Physiology*, 63, 359-390.
- Florou, A. B., Prodromidis, M. I., Tzouvara-Karayanni, S. M., & Karayannis, M. I. (2000). Fabrication and voltammetric study of lanthanum 2,6-dichlorophenolindophenol chemically modified screen printed electrodes: application for the determination of ascorbic acid. *Analytica Chimica Acta*, 423, 107-114.
- Fox, G. E., Magrum, L. J., Balch, W. E., Wolfe, R. S., & Woese, C. R. (1977). Classification of methanogenic bacteria by 16S ribosomal RNA characterization. *Proceedings of the National Academy of Science USA*, 74(10), 4537-4541.
- Franzmann, P. D. (1996). examination of Antarctic prokaryotic diversity through molecular comparisons. *Biodiversity and Conservation*, 85(11), 1295-1305.
- Fritsche, W., & Hofrichter, M. (2008). Aerobic degradation by microorganisms. *Biotechnology Set, Second Edition*, pp. 144-167.
- Gallagher, P. E., Weiss, R. B., Brent, T. P., & Duker, N. J. (1989). Wavelength dependence of DNA incision by a human ultraviolet endonuclease. *Photochemistry and Photobiology*, 49(3), 363-367. doi: 10.1111/j.1751-1097.1989.tb04120.x.
- Gallego, J. L. R., Loredó, J., Llamas, J. F., Vázquez, F., & Sánchez, J. (2001). Bioremediation of diesel-contaminated soils: evaluation of potential *in situ* techniques by study of bacterial degradation. *Biodegradation*, 12, 325-335.

- Gesheva, V., Stackebrandt, E. & Vasileva-Tonkova, E. (2010). Biosurfactant production by halotolerant *Rhodococcus fascians* from Casey Station, Wilkes Land, Antarctica. *Current Microbiology*, 61, 112–117.
- Gibson, G., & Muse, S. V. (2009). A Primer of Genome Science (3rd ed). Sinauer Associates, pp 84.
- Gibson, D. T., & Parales, R. E. (2000). Aromatic hydrocarbon dioxygenases in environmental biotechnology. *Current Opinion in Biotechnology*, 11(3), 236–243.
- Gibson, J., Shokralla, S., Porter, T. M., King, I., van Konynenburg, S., Janzen, D. H., Hallwachs, W., & Hajibabaei, M. (2014). Simultaneous assessment of the macrobiome and microbiome in a bulk sample of tropical arthropods through DNA metasytemics. *Proceedings of the National Academy of Sciences of USA*, 111(22), 8007–8012.
- Giorgio, M., Trinei, M., Migliaccio, E., & Pelicci, P. G. (2007). Hydrogen peroxide: a metabolic by-product or a common mediator of ageing signals? *Nature Reviews Molecular Cell Biology*, 8, 722–728. doi: 10.1038/nrm2240.
- Goodsell, D. S. (2009). Molecule of the month: antifreeze proteins. *Worldwide Protein Data Bank*, New jersey, USA.
- Goordial, J., Raymond-Bouchard, I., Ronholm, J., Shapiro, N., Woyke, T., Whyte, L., et al. (2015). Improved-high-quality draft gene sequence of *Rhodococcus* sp. JG-3, a eurypsychrophilic Actinobacteria from Antarctic Dry Valley permafrost. *Standard in Genomic Sciences*, 10(61), doi: 10.1186/s40793-015-0043-8.
- Greer, C. W., Whyte, L. G., Niederberger, T. D., & Timmis, K. N. (2010). Microbial communities in hydrocarbon contaminated temperate, tropical, alpine, and polar soils. *Handbook of Hydrocarbon and Lipid Microbiology*. Berlin Springer, 2313–2328.
- Greenshields, J. B., & Rossini, F. D. (1958). Molecular structure and properties of hydrocarbons and related compounds, *The Journal of Physical Chemistry*, 62(3), 271–280.
- Gao, B., & Gupta, R. S. (2012). Phylogenetic framework and molecular signatures for the main clades of the phylum Actinobacteria. *Microbiology and Molecular Biology Reviews*, 76(1), 66–112.
- Goody, R. S. (2013). How Bacteria Choose Phosphate. *Angewandte Chemie International Edition*, 52(9), 2406–2407. <https://doi.org/10.1002/anie.201209376>
- Haas, D., & Defago, G. (2005). Biological control of soil-borne pathogens by fluorescent pseudomonads. *Nature Reviews: Microbiology*, 3(4), 307–319.

- Haas, D., & Keel, C. (2003). Regulation of antibiotic production in root-colonizing *Pseudomonas* spp. and relevance for biological control of plant disease. *Annual Review of Phytopathology*, 41, 117-153.
- Haddock, J. D. (2010). Aerobic degradation of aromatic hydrocarbon: enzyme structures and catalytic mechanisms. In Timmis, K. N. (ed.), *Handbook of Hydrocarbon and Lipid Microbiology*, doi: 10.1007/978-3-540-77587-4_74, Springer-Verlag Berlin Heidelberg.
- Hamamura, N., Storfa, R. T., Semprini, L. & Arp, D. J. (1999). Diversity in butane monooxygenase among butane-grown bacteria. *Applied and Environmental Microbiology*, 65(10), 4586-4593.
- Hanlon, D. W., & Ordal, G. W. (1994). Cloning and characterization of genes encoding methyl-accepting chemotaxis proteins in *Bacillus subtilis*. *Journal of Biological Chemistry*, 269(19), 14038-14046.
- Hanlon, D. W., Carpenter, P. B., & Ordal, G. W. (1992). Influence of attractants and repellents on methyl group turnover on methyl-accepting chemotaxis proteins of *Bacillus subtilis* and role of CheW. *Journal of Bacteriology*, 174(13), 4218-4222.
- Hanson, K. G., Desai, J. D., & Desai, A. J. (1993). A rapid and simple screening technique for a potential crude oil degrading microorganism. *Journal of Chemical Technology and Biotechnology*, 7, 745-748.
- Harayama, S., Kok, M. & Neidle, E. L. (1992). Functional and evolutionary relationships among diverse oxygenases. *Annual Review of Microbiology*, 46, 565-601.
- Harayama, S., Kishira, H., Kasai, Y. & Shutsubo, K. (1999). Petroleum biodegradation in marine environments. *Journal of Molecular Microbiology and Biotechnology*, 1(1), 63-70.
- Harwood, C. S., & Ornston, L. N. (1984). TOL plasmid can prevent induction of chemotactic responses to aromatic acids. *Journal of Bacteriology*, 160(2), 797-800.
- Hathaway, L. J., Brugger, S., Martynova, A., Aebi, S., & Mühlemann, K. (2007). Use of the Agilent 2100 Bioanalyzer for Rapid and Reproducible Molecular Typing of *Streptococcus pneumoniae*. *Journal of Clinical Microbiology*, 45(3), 803-809. <https://doi.org/10.1128/JCM.02169-06>
- Hazan, R., Que, Y.-A., Maura, D., & Rahme, L. G. (2012). A method for high throughput determination of viable bacteria cell counts in 96-well plates. *BMC Microbiology*, 12, 259.
- Hazen, T. C. (1994). Chemotactic selection of pollutants degrading soil bacteria, In U.S. Patent 5324661.

- Hazen, T. C., & Lopez-de-Victoria, G. (1994). Method of degrading pollutants in soil, In U.S. Patent 5236703
- Head, I. M., & Swannell, R. P. (1999). Bioremediation of petroleum hydrocarbon contaminants in marine habitats. *Current Opinion in Biotechnology*, 10(3), 234-239.
- Heather, J. M., & Chain, B. (2016). The sequence of sequencers: the history of sequencing DNA. *Genomics*, 107(1), 1-8.
- Heider, J., Spormann, A. M., Beller, H. R., & Widdel, F. (1998). Anaerobic bacterial metabolism of hydrocarbons. *FEMS Microbiology Reviews*, 22(5), 459–473.
- Hunkapiller, T., Kaiser, R. J., Koop, B. F., & Hood, L. (1991). Large-scale and automated DNA sequence determination. *Science*, 254(5028), 59-67.
- Hyatt, D., Chen, G. L., Locascio, P. F., Land, M. L., Larimer, F. W., & Hauser, L. J. (2010). Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*, 11(119). doi: 10.1186/1471-2105-11-119.
- Hyman, E. D. (1988). A new method of sequencing DNA. *Analytical Biochemistry*, 174(2), 423–436.
- ITRC (Interstate Technology and Regulatory Council). (2014). *Petroleum Vapor Intrusion; Fundamentals of Screening, Investigation, and Management*. PVI-1. Washington, D.C.: Interstate Technology and Regulatory Council, Petroleum Vapor Intrusion Team. www.itrcweb.org/PetroleumVI-Guidance.
- Ismail, W., & Gescher, J. (2012). Epoxy coenzyme a thioester pathway for degradation of aromatic compounds. *Applied and Environmental Microbiology*, 78(15), 5043–5051.
- Jain, P. K., Gupta, V. K., Gaur, R. K., Lowry, M., Jaroli, D. P., & Chauhan, U. K. (2011). Bioremediation of petroleum oil contaminated soil and water. *Research Journal of Environmental Toxicology*, 5, 1–26. doi: 10.3923/rjet.2011.1.26.
- Janda, J. M., & Abbott, S. L. (2007). 16S rRNA gene sequencing for bacterial identification in diagnostic laboratory: pluses, perils, and pitfalls. *Journal of Clinical Microbiology*, 45, 2761–2764.
- Jansson, M. (1988). Phosphate uptake and utilization by bacteria and algae. *Hydrobiologia*, 170(1), 177–189. <https://doi.org/10.1007/BF00024904>
- Jesus, H. E., Peixoto, R. S., & Rosado, A. S. (2015). Bioremediation in Antarctic soils. *Environmental Biotechnology*, 6(6), 1-11.
- John, R. C., & Okpokwasili, G. C. (2012). Crude oil-degradation and plasmid profile of nitrifying bacteria isolated from oil-impacted mangrove sediment in the Niger Delta of Nigeria. *Bulletin of Environmental Contamination and Toxicology*, 88, 1020-1026.

- Johnson, E. L., & Hyman, M. R. (2006). Propane and *n*-butane oxidation by *Pseudomonas putida* GPo1. *Applied and Environmental Microbiology*, 72(1), 950-952.
- Johnson, G. R., & Olsen, R. H. (1997). Multiple pathways for toluene degradation in *Burkholderia* sp. strain JS150. *Applied and Environmental Microbiology*, 63, 4047-4052.
- Jones, O. A. H., Spurgeon, D. J., Svendsen, C., & Griffin, J. L. (2008). A metabolomics based approach to assessing the toxicity of the polycyclic aromatic hydrocarbon pyrene to the earthworm *Lumbricus rubellus*. *Chemosphere*, 71(3), 601-609.
- Joseph, S. J., Hugenholtz, P., Sangwan, P., Osborne, C. A., & Janssen, P. H. (2003). Laboratory cultivation of widespread and previously uncultured soil bacteria. *Applied Environmental Microbiology*, 69(12), 7210-7215.
- Jouanneau, Y., Meyer, C., Jakoncic, J., Stojanoff, V., & Gaillard, J. (2006). Characterization of a naphthalene dioxygenase endowed with an exceptionally broad substrate specificity toward polycyclic aromatic hydrocarbons. *Biochemistry*, 45, 12380-12391.
- Jukes, T. H., & Cantor, C. R. (1969). *Evolution of protein molecules*. In: Munro, H. N. (Ed). *Mammalian Protein Metabolism III*. Academic Press, New York, pp 21-132.
- Kaczorek, E., & Olszanowski, A. (2011). Uptake of hydrocarbon by *Pseudomonas fluorescens* (P1) and *Pseudomonas putida* (K1) strains in the presence of surfactants: a cell surface modification. *Water, Air and Soil Pollution*, 214(1-4), 451-459.
- Kambara, H., Nishikawa, T., Katayama, Y., & Yamaguchi, T. (1988). Optimization of parameters in a DNA sequenator using fluorescence detection. *Nature Biotechnology*, 6, 816-821.
- Kalman, B. & Johnson, R. (2007). *Endangered Penguins*: New York: Crabtree Publishing and Cooperation.
- Kauppi, S., Sinkkonen, A., & Romantschuk, M. (2011). Enhancing bioremediation of diesel-fuel-contaminated soil in a boreal climate: comparison of biostimulation and bioaugmentation. *International Biodeterioration and Biodegradation*, 65(2), 359-368. doi: 10.1016/j.ibiod.2010.10.011.
- Kennicutt, M. C. (1990). Oil spillage in Antarctica: initial report of the National Science Foundation-sponsored Quick Response Team on the grounding of the Bahia Paraiso. *Environmental Science and Technology*, 24(5), 620-624.
- Kennicutt, M. C., McDonald, T. J., Denoux, G. J., & McDonald, S. J. (1992). Hydrocarbon contamination on the Antarctic Peninsula: I. Arthur harbor-subtidal sediments. *Marine Pollution Bulletin*, 24(10), 499-506. [https://doi.org/10.1016/0025-326X\(92\)90474-K](https://doi.org/10.1016/0025-326X(92)90474-K).

- Kim, K., Jahan, S. A., Kabir, E., & Brown, R. J. C. (2013). A review of airborne polycyclic aromatic hydrocarbons (PAHs) and their human health effects. *Environmental International*, 60, 71-80.
- Kinawy, A. A. (2009). Impact of gasoline inhalation on some neurobehavioural characteristics of male rats. *BMC Physiology*, 9(21). doi: 10.1186/1472-6793/9/21
- Knothe, G. (2010). Biodiesel and renewable diesel: a comparison. *Progress in Energy and Combustion Sciences*, 36, 364-373.
- Komukai-Nakamura, S., Sugiura, K., Yamauchi-Inomata, Y., Toki, H., Venkateswaran, K., Yamamoto, S., Tanaka, H., & Harayama, S. (1996). Construction of bacterial consortia that degrade Arabian light crude oil. *Journal of Fermentation and Bioengineering*, 382(6), 570-574.
- Konidari, C. N., Tzouware, S. M., Bowman, L. E., & Karayannis, M. I. (1992). Kinetic and mechanistic study of the reaction of 2,6-dichlorophenolindophenol and cysteine. *Talanta*, 39, 863-868.
- Kosina, M., Barták, M., Mašlaňová, I., Pascutti, A. V., Sedo, O., Lexa, M., & Sedláček, I. (2013). *Pseudomonas prosekii* sp. nov., a novel psychrotrophic bacterium from Antarctica. *Current Microbiology*, 67(6), 637-646. <https://doi.org/10.1007/s00284-013-0406-6>
- Kotani, T., Yamamoto, T., Yurimoto, H., Sakai, Y., & Kato, N. (2003). Propane monooxygenase and NAD⁺ dependent secondary alcohol dehydrogenase in propane metabolism by *Gordonia* sp. strain TY-5. *Journal of Bacteriology*, 185, 7120-7128.
- Krastanov, A., Alexieva, Z., & Yemendzhiev, H. (2013). Microbial degradation of phenol and phenolic derivatives. *Engineering in Life Sciences*, 13(1), 76-87.
- Kuhn, E., Bellicanta, G. S., & Pellizari, V. H. (2009). New *alk* genes detected in Antarctic marine sediments. *Environmental Microbiology*, 11(3), 669-673.
- Kuo, C. Y., Hsu, Y. W. & Lee, H. S. (2003). Study of human exposure to particulate PAHs using personal air samplers. *Archive of Environmental Contamination and Toxicology*, 44, 454-459.
- Labinger, J. A., & Bercaw, J. E. (2002). Understanding and exploiting C-H bond activation. *Nature*, 417(6888), 507-514.
- Laczi, K., Kis, A., Horvath, B., Maroti, G., Hegedus, B., Perei, K., & Rakhely, G. (2015). Metabolic responses of *Rhodococcus erythropolis* PR4 grown on diesel oil and various hydrocarbons. *Applied Microbiology and Biotechnology*, 99(22), 9745-9759.

- Lagesen, K., Hallin, P., Rodland, E. A., Staerfeldt, H. H., Rognes, T., & Ussery, D. W. (2007). RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Research*, 35(9), 3100-3108.
- Laslett, D. & Bjorn, C. (2004). ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Research*, 32(1), 11-16.
- Laurence, M., Hatzis, C., & Brash, D. E. (2014). Common contaminants in next-generation sequencing that hinder discovery of low-abundance microbes. *PLoS One*. 9(5), e97876. doi: 10.1371/journal.pone.0097876.
- Leahy, J. G., Batchelor, P. J., & Morcomb, S. M. (2003). Evolution of the soluble diiron monooxygenases. *FEMS Microbiology Reviews*, 27(4), 449-714.
- Lee, C-H., Liang, S., Haumann, T., Boese, R., & de Meijere, A. (1993). p -[3²,5⁶] octahedrane, the (CH)₁₂ hydrocarbon with D_{3d} symmetry. *Angewandte Chemie International Edition*, 32(4), 559-561.
- Liang, J-L., JiangYang, J-H., Nie, Y. & Wu, X-L. (2016). Regulation of alkane hydroxylase CYP153 gene in a gram-positive alkane-degrading bacterium, *Dietzia* sp. strain DQ12-45-1b. *Applied and Environmental Microbiology*, 82(2), 608-619.
- Likhoshvay, A., Lomakina, A., & Grachev, M. (2014). The complete *alk* sequences of *Rhodococcus erythropolis* from Lake Baikal. *SpringerPlus*, 39(621). doi: 10.1186/2193-1801-3-621.
- Lo Giudice, A., Bruni, V., De Domenico, M. & Michaud, L. (2010). Psychrophiles - cold-adapted hydrocarbon-degrading microorganisms. In Timmis, K. N. (Ed), *Handbook of Hydrocarbon and Lipid Microbiology*, doi: 10.1007/978-3-540-77587-4, 1897-1921.
- Loman, N. J., Constantinidou, C., Chan, J. Z. M., Halachev, M., Sergeant, M., Penn, C. W., Robinson, E. R., & Pallen, M. J. (2012). High-throughput bacterial genome sequencing: an embarrassment of choice, a world of opportunity. *Nature Reviews Microbiology*, 10, 499-606.
- Long, J., Ma, J., Luo, C., Mo, X., Sun, L., Zang, W., & Liu, J. (2009). Comparison of two methods for assaying complex I activity in mitochondrial isolated from rat liver, brain and heart. *Life Science*, 85, 276-280.
- Luckey, J. A., Drossman, H., Kosticjka, A. J., Mead, D. A., D'Cunha, J., Norris, T. B., & Smith, L. M. (1990). High speed DNA sequencing by capillary electrophoresis. *Nucleic Acids Research*, 18(15), 4417-4421.
- Ludwig, B., Akundi, A., & Kendall, K. (1995). A long-chain secondary alcohol dehydrogenase from *Rhodococcus erythropolis* ATCC 4277. *Applied and Environmental Microbiology*, 61(10), 3729-3733.

- Lugtenberg, B., & Kamilova, F. (2009). Plant-growth-promoting rhizobacteria. *Annual Review of Microbiology*, 63, 541-546.
- Ma, Y., Wang, L., & Shao, Z. (2006). *Pseudomonas*, the dominant polycyclic aromatic hydrocarbon-degrading bacteria isolated from Antractic soils and the role of large plasmids in horizontal gene transfer. *Environmental Microbiology*, 8(3), 455-465.
- Mac Cormack, W. P., & Fraile, E. R. (1997). Characterization of a hydrocarbon degrading psychrotrophic Antarctic bacterium. *Antarctic Science*, 9(2), 150-155.
- Macoustra, G. K., King, C. K., Wasley, J., Robinson, S. A., & Jolley, D. F. (2015). Impact of hydrocarbons from a diesel fuel on the germination and early growth of sub-Antarctic plants. *Environmental Science Processes and Impacts*, 17(7), 1238-1248.
- Malavenda, R., Rizzo, C., Michaud, L., Gerçe, B., Bruni, V., Syltatk, C., et al. (2015). Biosurfactant production by Arctic and Antarctic bacteria growing on hydrocarbons. *Polar Biology*, 38, 1565-1574.
- Marchal, R., Penet, S. Solano-Serena, F., & Vandecasteele, J. P. (2003). Gasoline and diesel oil biodegradation. *Oil and Gas Science and Technology*, 58(4), 441-448.
- Mardis, E. R. (2008). Next-generation DNA sequencing methods. *Annual Review of Genomics and Human Genetics*, 9, 387-402.
- Margesin, R., & Schinner, F. (1999). Biological decontamination of oil spills in cold environments. *Journal of Chemical Technology and Biotechnology*, 74, 381-389.
- Margesin, R., Gander, S., Zacke, G., Gounot, A. M., & Schinner, F. (2003). Hydrocarbon degradation and enzyme activities of cold-adapted bacteria and yeasts. *Extremophiles*, 7(6), 451-458.
- Marguiles, M., Egholm, M., Altman, W. E., Attiya, S., Bader, J. S., Bemben, L. A., Berka, J., et al. (2005). Genome sequencing in microfabricated high-density picolitre reactors. *Nature*, 437(7057), 376-380.
- Mariano, A. P., Tomasella, R. C., de Oliveira, L. M., Contiero, J., & de Angelis, D. F. (2008). Biodegradability of diesel and biodiesel blends. *African Journal of Biotechnology*, 7(9), 1323-1328.
- Marshall, B., Robleto, E. A., Wetzler, R., Kulle, P., Casaz, P., & Levy, S. B. (2001). The *adnA* transcriptional factor affects persistence and spread of *Pseudomonas fluorescens* under natural field conditions. *Applied Environmental Microbiology*, 67, 852-857.

- Martinez, Salette, & Hausinger, R. P. (2015). Catalytic mechanisms of Fe (II)- and 2-oxoglutarate-dependent oxygenases. *The Journal of Biological Chemistry*, 290(34), 20702-20711.
- Martinez-Alvarez, L. M., Ruberto, L. A. M., Lo Balbo, A., & Mac Cormack, W. P. (2017). Bioremediation of hydrocarbon-contaminated soils in cold regions: development of a pre-optimized biostimulation biopile-scale field assay in Antarctica, *Science of the Total Environment*, 90-91, 194-203. doi: 10.1016/j.scitotenv.2017.02.204.
- Masotti, A., & Preckel, T. (2006). Analysis of small RNAs with the Agilent 2100 Bioanalyzer. *Nature Methods| Application Notes*.
- McDonald, R., & Knox, G. G. O. (2014). Cold region bioremediation of hydrocarbon contaminated soils: do we know enough?. *Environmental Science and Technology*, 48(17), 9980-9981.
- Meliani, A., & Bensoltane, A. (2014). Review of *Pseudomonas* attachment and biofilm formation in food industry. *Poultry, Fisheries and Wildlife Sciences*, 2(126), doi: 10.4172/2375-446X.1000126.
- Mengel, K., & Kirkby, E. A. (1987). Principles of Plant Nutrition. International Potash Institute, Worblaufen-Bern, Switzerland.
- Meyer, M., & Kircher, M. (2010). Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Springs Harbor protocols*. doi: 10.1101/pdb.prot5448.
- Meyer, J-M., Stintzi, A., Coulanges, V., Shivaji, S., Voss, J. A., Taraz, K., & Budzikiewicz (1998). Siderotyping of fluorescent pseudomonads: characterization of pyoverdines of *Pseudomonas fluorescens* and *Pseudomonas putida* strains from Antarctica. *Microbiology*, 144, 3119-3126.
- Michaud, L., Lo Giudice, A., Saitta, M., De Domenico, M., & Bruni, V. (2004). The biodegradation efficiency on diesel oil by two psychrotrophic Antarctic marine bacteria during a two-long experiment. *Marine Pollution Bulletin*, 49, 405-409.
- Mille, G., Almallah, M., Bianchi, M., van Wambeke, F., & Bertrand, J. C. (1991). Effect of salinity on petroleum biodegradation, *Fresenius Journal of Analytical Chemistry*, 339, 788-791.
- Molavian, H. R., Kohandel, M., & Sivaloganathan, S. (2016). High concentrations of H₂O₂ make aerobic glycolysis energetically more favorable for cellular respiration. *Frontiers in Physiology*, 7, 362. doi: 10.3389/fphys.2016.00362.
- Montagnolli, R. N., Lopes, P. R. M., & Bidoia, E. D. (2015). Screening the toxicity and biodegradability of petroleum hydrocarbons by a rapid colorimetric method. *Archives of Environmental Contamination and Toxicology*, 68, 342-353.

- Morgan, P., & Watkinson, R. J. (1994). Biodegradation of components of petroleum. In Rathledge, C. (Ed), *Biochemistry of Microbial Degradation* (pp. 1-31). Dordrecht, Netherlands: Kluwer Academic Publishers.
- Morita, R. Y. (1975). Psychrophilic bacteria. *Bacteriological Reviews*, 37, 144–167.
- Muangchinda, C., Chavanich, S., Viyakara, V., Watanabe, K., Imura, S., Vangnai, A. S., et al. (2015). Abundance and diversity of functional genes involved in the degradation of aromatic hydrocarbons in Antarctic soils and sediments around Syowa Station. *Environmental Science and pollution Research*, 22, 4725–4735.
- Muhammad, S., Muller, T., & Jorgensen, R. G. (2008). Relationships between soil biological and other soil properties in saline and alkaline arable soils from the Pakistani Punjab. *Journal of Arid Environments*, 72, 448–457.
- Mukherjee, S., Huntemann, M., Ivanova, N., Kyrpides, N. C., & Pati, A. (2015). Large-scale contamination of microbial isolate genomes by Illumina PhiX control. *Standards in Genomic Sciences*, 10(18). doi: 10.1186/1944-3277-10-18.
- Mulder, H., Breure, A. M., Van Andel, J. G., Grotenhuis, J. T., & Rulkens, W. H. (1998). Influence of hydrodynamic conditions on naphthalene dissolution and subsequent biodegradation. *Biotechnology and Bioengineering*, 57(2), 145–154.
- Müller, T., Walter, B., Wirtz, A., & Burkovski, A. (2006). Ammonium toxicity in bacteria. *Current Microbiology*, 52(5), 400–406.
- Munoz, P. A., Marquez, S. L., Gonzalez-Nilo, F. D., Marquez-Miranda, V., & Blamey, J. M. (2017). Structure and application of antifreeze proteins from Antarctic bacteria. *Microbial Cell Factories*, 16(138), 1–13.
- Myers, R. H., & Montgomery, D. C. (1995). *Response surface methodology: process and product optimization using designed experiments*, John Wiley and Sons, New York.
- Nakamura, H., Kobayashi, S., Hirata, Y., Suzuki, K., Mogi, Y., & Karube, I. (2007). A spectrophotometric biochemical oxygen demand determination method using 2,6-dichlorophenolindophenol as the redox color indicator and the eukaryote *Saccharomyces cerevisiae*. *Analytical Biochemistry*, 369, 168–174.
- Nakamura, F. M., Germano, M. G., & Tsai, S. M. (2014). Capacity of aromatic compound degradation by bacteria from Amazon Dark Earth. *Diversity*, 6, 339–353. doi: 10.3390/d6020339.
- Naseby, D. C., Way, J. A., Bainton, N. J., & Lynch, J. M. (2001). Biocontrol of *Phytophthora* in the pea rhizosphere by antifungal metabolite producing and non-producing *Pseudomonas* strains. *Journal of Applied Microbiology*, 90(3), 421–429.

- Nie, Y., Chi, C. Q., Fang, H., Liang, J. L., Lu, S. L., Lai, G. L., et al. (2014). Diverse alkane hydroxylase genes in microorganisms and environments. *Scientific Reports*, 4, 4968.
- Nkem, J. N., Virginia, R. A., Barrett, J. E., Wall, D. H., & Li, G. (2006). Salt tolerance and survival thresholds for two species of Antarctic soil nematodes. *Polar Biology*, 29(8), 643–651. <https://doi.org/10.1007/s00300-005-0101-6>
- NRC (National Research Council) (2003). Oil in the sea III: inputs, fates and effects. Washington: The National Academic Press.
- Nydahl, A. C., King, C. K., Wasley, J., Jolley, D. F., & Robinson, S. A. (2015). Toxicity of fuel-contaminated soil to Antarctic moss and terrestrial algae. *Environmental Toxicology and Chemistry*, 34(9), 2004–2012.
- Nyren, P. (1987). Enzymatic method for continuous monitoring of DNA polymerase activity. *Analytical Biochemistry*, 167(2), 235–238.
- Nyren, P., & Lundin, A. (1985). Enzymatic method for continuous monitoring of inorganic pyrophosphate synthesis. *Analytical Biochemistry*, 151(2), 504–509.
- Obidike, I. R., Maduabuchi, I. U., & Olumuyiwa, S. S. O. (2007). Testicular morphology and cauda epididymal sperm reserves of male rats exposed to Nigerian Qua Iboe Brent crude oil. *Oil and Chemical Pollution*, 3, 117–129.
- Oboh, B. O., Adeyinka, Y., Awonuga, S., & Akinola, M. O. (2007). Impact of soil types and petroleum effluents on the earthworm *Eudrillus eugeniae*. *Journal of Environmental Biology*, 28(2), 209–212.
- OEHHA (Office of Environmental Health Hazard Assessment) (2001). Health effects of diesel exhaust. Sacramento, CA. California Environmental Protection Agency.
- Okpokwasili, G. C., Somerville, C. C., Sullivan, M., Grimes, D. J., & Colwell, R. R. (1986). Plasmid-mediated degradation of hydrocarbons by estuarine bacteria. *Research Journal of Environmental Toxicology*, 3(1), 9–23.
- Ondov, B. D., Bergman, N. H., & Phillippy, A. M. (2011). Interactive metagenomic visualization in a Web browser. *BMC Bioinformatics*, 12(385). doi: 10.1186/1471-2105-12-385.
- Oudot, J., Merlin, F. X., & Oinvidic, P. (1998). Weathering rates of oil components in a bioremediation experiment in estuarine sediments. *Marine Environmental Research*, 45(2), 113–125.
- Overbeek, R., Olson, R., Pusch, G. D., Olsen, G. J., Davis, J. J., Disz, T., Edwards, R. A., et al. (2014). The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acid Research*, 42, 206–214. <https://doi.org/10.1093/nar/gkt1226>.

- Pacheco, G. J., Ciapina, E. M., Gomes, E. B., & Junior, N. P. (2010). Biosurfactant production by *Rhodococcus erythropolis* and its application to oil removal. *Brazilian Journal of Microbiology*, 41(3), 685-693. doi: 10.1590/S1517-83822010000300019.
- Pacwa-Plociniczak, M., Plaza, G. A., Poliwoda, A., & Piotrowska-Seget, Z. (2014). Characterization of hydrocarbon-degrading and biosurfactant-producing *Pseudomonas* sp. P-1 strain as a potential tool for bioremediation of petroleum-contaminated soil. *Environmental Science and Pollution Research*, 21, 9385–9395.
- Paixao, J. F., Nascimento, I. A., Pereira, S. A., Leite, M. B. L., Carvalho, G. C., Silveira Jr, J. S. C., et al. (2007). Estimating the gasoline component and formulations toxicity to microalgae (*Tetramis chuii*) and oyster (*Crassostrea rhizophorae*) embryos: an approach to minimize environmental pollution risk. *Environmental Research*, 103(3), 365-374.
- Pandey, G., & Jain, R. K. (2002). Bacterial chemotaxis toward environmental pollutants: role in bioremediation. *Applied and Environmental Microbiology*, 68(12), 5789-5795.
- Panke, S., Sanchez-Romero, J. M., & de Lorenzo, V. (1998). Engineering of quasi-natural *Pseudomonas putida* strains for toluene metabolism through an ortho-cleavage degradation pathway. *Applied and Environmental Microbiology*, 64(2), 748–751.
- Parales, R. E., & Harwood, C. S. (2002). Bacterial chemotaxis to pollutants and plant-derived aromatic molecules. *Current Opinion in Microbiology*, 5(3), 266-273.
- Parales, R. E., Parales, J. V., Pelletier, D. A., & Ditty, J. L. (2008). Diversity of microbial toluene degradation pathways. *Advances in Applied Microbiology*, 64, 1-73.
- Perelo, L. W. (2010). Review: *in situ* and bioremediation of organic pollutants in aquatic sediments. *Journal of Hazardous Materials*, 177, 81–89.
- Peressutti, S. R., Alvarez, A. M., & Pucci, O. H. (2003). Dynamics of hydrocarbon-degrading bacteriocenosis of an experiment oil pollution in Patagonian soil. *International Biodeterioration and Biodegradation*, 52, 21–30.
- Perez-de-Mora, A., Engel, M., & Schlöter, M. (2011). Abundance and diversity of *n*-alkane degrading bacteria in a forest soil co-contaminated with hydrocarbons and metals: a molecular study on *alkB* homologous gene. *Microbial Ecology*, 62(4), 959–972.
- Pérez-Pantoja, D., Gonzalez, B., & Pieper, D. H. (2010). Aerobic degradation of aromatic hydrocarbons. In Timmis, K. N. (Ed), *Handbook of Hydrocarbon and Lipid Microbiology*. doi: 10.1007/978-3-540-77587-4_60, Springer-Verlag Berlin Heidelberg.

- Peterson, D. R. (1994). Calculating the aquatic toxicity of hydrocarbon mixtures. *Chemosphere*, 29(12), 2493–2506.
- Pieper, D. H. (2005). Aerobic degradation of polychlorinated biphenyls. *Applied Microbiology and Biotechnology*, 67(2), 170–191.
- Powell, S. M., Ferguson, S. H., Snape, I., & Siciliano, S. D. (2006). Fertilization stimulates anaerobic fuel degradation of Antarctic soils by denitrifying microorganisms. *Environmental Science and Technology*, 40(15), 2011–2017.
- Powell, S. M., Harvey, P. M., Stark, J. S., Snape, I., & Riddle, M. J. (2007). Biodegradation of petroleum products in experimental plots in Antarctic marine sediments is location dependent. *Marine Pollution Bulletin*, 54(4), 434–440.
- Peng, X., Taki, H., Komukai, S., Sekine, M., Kanoh, K., Kasai, H., Choi, S-K., et al. (2006). Characterization of four *Rhodococcus* alcohol dehydrogenase genes responsible for the oxidation of aromatic alcohols. *Applied Microbiology and Biotechnology*, 71, 824–832.
- Perry, J. J. (2007). Microbial metabolism of cyclic alkanes. In Atlas, R. M. *Petroleum Microbiology*. New York, Macmillan Publishing Co., 61–97.
- Plackett, R. L., & Burman, J. P. (1946). The Design of Optimum Multifactorial Experiments. *Biometrika*, 33(4), 305–325. <https://doi.org/10.2307/2332195>
- Poland, J. S., Riddle, M. J., & Zeeb, B. A. (2003). Contaminants in the Arctic and Antarctic: a comparison of sources, impacts, and remediation options. *Polar Record*, 39, 369–384.
- Prosser, J. I., & Embley, T. M. (2002). Cultivation-based and molecular approaches to characterisation of terrestrial and aquatic nitrifiers. *Antonie Van Leeuwenhoek*, 81(1–4), 165–179.
- Provin, T., & Pitt, J. L. (2001). Managing soil salinity. Texas A and M University, Texas: Texas Agricultural Extension Service, pp 5.
- Rabinowitz, J. D., Vacchino, J. F., Beeson, C., & McConnell, H. M. (1998). Potentiometric measurements of intracellular redox activity. *Journal of the American Chemical Society*, 120, 2464–2473.
- Rabus, R., Boll, M., Heider, J., Meckenstock, R. U., Buckel, W., Einsle, O., Ermler, U., et al (2016). Anaerobic microbial degradation of hydrocarbons: from enzymatic reactions to the environment. *Journal of Molecular Microbiology and Biotechnology*, 26(1–3), 5–28. doi: 10.1159/000443997.
- Rainey, F. A., Burghardt, J., Kroppenstedt, R. M., Klatte, S., & Stackbrandt, E. (1995). Phylogenetic analysis of the genera *Rhodococcus* and *Nocardia* and evidence for the evolutionary origin of the genus *Nocardia* from within the radiation of *Rhodococcus* species. *Microbiology*, 141, 523–528.

- Rainwater, F. H., & Thatcher, L. L. (1960). *Methods for Collection and Analysis of Water Samples* (Water Supply Paper No. 1454) (p. 301). U.S.: U.S. Government Printing Office.
- Rajendhran, J., & Gunasekaran, P. (2011). Microbial phylogeny and diversity: small subunit ribosomal RNA sequence analysis and beyond. *Microbiological Research*, 166(2), 99-110.
- Raymond, T., King, C. K., Raymond, B., Stark, J. S., & Snape, I. (2017). Chapter 14 - Oil Pollution in Antarctica. In M. Fingas (Ed.), *Oil Spill Science and Technology (Second Edition)* (pp. 759–803). Boston: Gulf Professional Publishing.
- Rayner, J. L., Snape, I., Walworth, J. L., Harvey, P. M., & Ferguson, S. H. (2007). Petroleum-hydrocarbon contamination and remediation by microbioventing at sub-Antarctic Macquarie Island. *Cold Regions Science and Technology*, 48, 139-153.
- Reddy, G. S., Matsumoto, G. I., Schumann, P., Stackerbrandt, E., & Shivaji, S. (2004). Psychrophilic pseudomonads from Antarctica: *Pseudomonas antarctica* sp. nov., *Pseudomonas meridiana* sp. nov. and *Pseudomonas proteolytica* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, 54(Pt 3), 713-719.
- Richard, J. Y., & Vogel, T. M. (1999) Characterization of a soil bacterial consortium capable of degrading diesel fuel. *International Biodeterioration and Biodegradation*, 44(2-3), 93-100.
- Richardson, E. L., King, C. K., & Powell, S. M. (2014) The use of microbial gene abundance in the development of fuel remediation guidelines in polar soils. *Integrated Environmental Assessment and Management*, 11(2), 235-241.
- Rhykerd, R. I., Weaver, R. W., & McInnes, K. J. (1995). Influence of salinity on bioremediation of oil in soil. *Environmental Pollution*, 90, 127-130.
- Rietz, D. N., & Haynes, R. J. (2003). effects of irrigation-induced salinity and sodicity on soil microbial activity. *Soil Biology and Biochemistry*, 35(6), 845-854.
- Rodriguez-Fernandez, J., Pereiro, R., & Sanz-Mendel, A. (2002). Optical fibre sensor for hydrogen sulphide monitoring in mouth air. *Analytica Chimica Acta*, 471, 13-23.
- Rodriguez-Trigo, G., Zock, J., Pozo-Rodriguez, F., Gomez, F. P., Monyarch, G., Bouso, L., et al. (2010). Health changes in fishermen 2 years after clean-up of the Prestige oil spill. *Annals of Internal Medicine*, 153(8), 489-498.
- Rojo, F. (2009). Degradation of alkanes by bacteria. *Environmental Microbiology*, 11(10), 2477–2490.

- Ruberto, L., Vazquez, S., Lo Balbo, A., & Mac Cormack, W. P. (2005). Psychrotolerant hydrocarbon-degrading *Rhodococcus* strains isolated from polluted Antarctic soils. *Antarctic Science*, 17(1), 47-56.
- Ruberto, L., Vazquez, S. C., & Mac Cormack, W. P. (2003). Effectiveness of the natural bacterial flora, biostimulation and bioaugmentation on the bioremediation of the hydrocarbon contaminated Antarctic soil. *International Biodeterioration and Biodegradation*, 52(2), 115-125.
- Ronaghi, M., Karamohamed, S., Pettersson, B., Uhlen, M., & Nyren, P. (1996). Real-time DNA sequencing using detection of pyrophosphate release. *Analytical Biochemistry*, 242(1), 84-89.
- Ronaghi, M., Uhlen, M., & Nyren, P. (1998). A sequencing method based on real-time pyrophosphate. *Science*, 281(5375), 363-365.
- Saito, A., Iwabuchi, T., & Harayama, S. (1999). Characterization of genes for enzymes involved in the phenanthrene degradation in *Nocardioides* sp. KP7, *Chemosphere*, 38, 1331-1337.
- Saitou, N., & Nei, M. (1987). The neighbour-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4(4), 406-425.
- Sanger, F., Nicklen, S., & Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors, *Proceedings of the National Academy Sciences USA*, 74, 5463-5467.
- Santa Anna, L. M., Sebastian, G. V., Menezes, E. P., Alves, T. L. M., Santos, A. S., Junior, N. P., et al. (2002) Production of biosurfactants from *Pseudomonas aeruginosa* PA1 isolated in oil environments. *Brazilian Journal of Chemical Engineering*, 19(2), 159-166.
- Saul, D. J., Aislabie, J. M., Brown, C. E., Harris, L., & Foght, J. M. (2005). Hydrocarbon contamination changes the bacterial diversity of soil from around Scott Base, Antarctica. *FEMS Microbiology Ecology*, 53, 141-155.
- SCAR (Scientific Committee on Antarctic Research) (1993). Protocol on environmental protection to the Antarctic Treaty. *Polar Record*, 29, 256-275.
- Schweitzer, C., & Scaiano, J. C. (2003). Selective binding and local photophysics of the fluorescent cyanine dye PicoGreen in double-stranded and single-stranded DNA. *Physical Chemistry Chemical Physics*, 5(21), 4911-4917.
- Sekine, M., Tanikawa, S. S., Omata, S., Saito, M., Fujisawa, T., Tsukatani, N., et al. (2006). Sequence analysis of three plasmids harboured in *Rhodococcus erythropolis* strain PR4. *Environmental Microbiology*, 8(2), 334-346.
- Sierra-Garcia, I. N., & de Oliveira, V. M. (2013). Microbial hydrocarbon degradation: efforts to understand biodegradation in petroleum reservoirs. *Biodegradation - Engineering and Technology*, doi: 10.5772/55920.

- Seo, J-S., Keum, Y-S., & Li, Q. X. (2009). Bacterial degradation of aromatic compounds. *International Journal of Environmental Research and Public Health*, 6(1), 278-309.
- Silby, M. W., & levy, S. B. (2004). Use of in vivo expression technology to identify genes important in growth and survival of *Pseudomonas fluorescens* Pf0-1 in soil: discovery of expressed sequences with novel genetic organization. *Journal of Bacteriology*, 186(21), 7411-7419.
- Silby, M. W., & Levy, S. B. (2008). Overlapping protein-encoding genes in *Pseudomonas fluorescens* Pf0-1. *PLoS Genetics*, 4(6), e1000094.
- Silby, M. W., Nicoll, J. S., & Levy, S. B. (2009). Requirement of polyphosphate by *Pseudomonas fluorescens* Pf0-1 for competitive fitness and heat tolerance in laboratory media and sterile soil. *Applied and Environmental Microbiology*, 75(12), 3872-3881.
- Singh, P., DeMarini, D. M., Dick, C. A. J., Tabor, D. G., Ryan, J. V., Linak, W. P., et al. (2004). Sample characterisation of automobile and forklift diesel exhaust particles and comparative pulmonary toxicity in mice. *Environmental Health Perspectives*, 112(8), 820-825.
- Singh, A., & Ward, O. P. (2004). Biotechnology and bioremediation- an overview. In: Singh, A., Ward, O. P. (ed.) Biodegradation and Bioremediation. *Soil Biology*, vol. 2. Springer-Verlag, Berlin, Heidelberg.
- Shimizu, K. (2013). Metabolic Regulation of a Bacterial Cell System with Emphasis on *Escherichia coli* Metabolism. *International Scholarly Research Notices*. Retrieved from <https://www.hindawi.com/journals/isrn/2013/645983/>
- Shukor, M. Y., Hassan, N. A., Jusoh, A. Z., Perumal, N., Shamaan, N. A., Mac Cormack, W. P., et al. (2009). Isolation and characterisation of a *Pseudomonas* diesel-degrading strain from Antarctica. *Journal of Environmental Biology*, 30(1), 1-6.
- Siqueira, J. F., Fouad, A. F., & Rôças, I. N. (2012). Pyrosequencing as a tool for better understanding of human microbiomes. *Journal of Oral Microbiology*, 4. <https://doi.org/10.3402/jom.v4i0.10743>
- Snape, I., Morris, C. E., & Cole, C. M. (2001). The use of permeable reactive barriers to control contaminants dispersal during site remediation in Antarctica. *Cold Regions Science and Technology*, 32, 157-174.
- Snape, I., Acomb, L., Barnes, D. I., Bainbridge, S., Eno, R., Filler, D. M., Plato, N., et al. (2008). Contamination, regulation and remediation: an introduction to bioremediation of petroleum hydrocarbons in cold regions. In Filler, D. M., Snape, I., & Barnes, D. L. *Bioremediation of petroleum hydrocarbons in cold regions*, Cambridge University Press.
- Staden, R. (1980). A new computer method for the storage and manipulation of DNA gel reading data. *Nucleic Acids Research*, 8(16), 3673-3694.

- Stallwood, B., Shears, J., Williams, P. A., & Hughes, K. A. (2005). Low temperature bioremediation of oil-contaminated soil using biostimulation and bioaugmentation with a *Pseudomonas* sp. from maritime Antarctica. *Journal of Applied Microbiology*, 99(4), 794-802.
- Stancu, M. M. (2015). Response of *Rhodococcus erythropolis* strain IBB_{P01} to toxic organic solvents. *Brazilian Journal of Microbiology*, 46(4), 1009-1018.
- Stevens, V., Thijs, S., McAmmond, B., Langill, T., Van Hamme, J., Weyens, N., & Vangronsveld, J. (2017). Draft genome sequence of *Rhodococcus erythropolis* VSD3, a diesel fuel-degrading and plant growth-promoting bacterium isolated from *Hedera helix* leaves. *Genome Announcements*, 5(8). doi: 10.1128/genomeA.01680-16.
- Strong, M. J., Xu, G., Morici, L., Bon-Durant, S. S., Baddoo, M., Lin, Z., Fewell, C., Taylor, C. M., & Flemington, E. K. (2014). Microbial contamination in next generation sequencing: implications for sequence-based analysis of clinical samples. *PLoS Pathogen*, 10(11): e1004437. doi: 10.1371/journal.ppat.1004437.
- Suyama, A., Iwakiri, R., Kimura, N., Nishi, A., Nakamura, K., & Furukawa, K. (1996). Engineering hybrid pseudomonads capable of utilizing a wide range of aromatic hydrocarbons and of efficient degradation of trichloroethylene. *Journal of Bacteriology*, 178(14), 4039-4046.
- Swerdlow, H., & Gesteland, R. (1990). Capillary gel electrophoresis for rapid, high resolution DNA sequencing. *Nucleic Acids Research*, 18(6), 1415-1419.
- Taketani, R. G., Zucchi, T. D., de Melo, I. S., & Mendes, R. (2013). Whole-genome shotgun sequencing of *Rhodococcus erythropolis* strain P27, a highly radiation-resistant actinomycete from Antarctica, *Genome Announcement*, 1(5), e00763-13.
- Takihara, H., Ogihara, J., Yoshida, T., Okuda, S., Nakajima, M., Iwabuchi, N., & Sunain, M. (2014). Enhanced translocation and growth of *Rhodococcus erythropolis* PR4 in the alkane phase of aqueous-alkane two phase cultures were mediated by GroEL2 overexpression. *Microbes and Environment*, 29(4), 346-352.
- Tanco, M., Viles, E., & Pozueta, L. (2009). Comparing Different Approaches for Design of Experiments (DoE). In *Advances in Electrical Engineering and Computational Science* (pp. 611-621). Springer, Dordrecht
- Tang, J., Wang, M., Wang, F., Sun, Q., & Zhou, Q. (2011). Eco-toxicity of petroleum hydrocarbon contaminated soil. *Journal of Environmental Sciences*, 23(5), 845-851.
- Tawfik, D. S., & Giffiths, A. D. (1998). Man-made cell-like compartments for molecular evolution. *Nature Biotechnology*, 16(7), 652-656.

- Teare, J. M., Islam, R., Flanagan, R., Gallagher, S., Davies, M. G., & Grabeau, C. (1997). Measurement of nucleic acid concentrations using the DyNA Quant™ and the GeneQuant™. *Biotechniques*, 22, 1170-1174.
- Tomei, M. C., & Daugulis, A. J. (2013). *Ex situ* bioremediation of contaminated soils: an overview of conventional and innovative technologies. *Critical Reviews in Environmental Science and Technology*, 43, 2107-2139.
- Tourova, T. P., Sokolova, D. S., Semenova, E. M., Shumkova, E. S., Korhunova, A. V., Babich, T. L., Poltarau, A. B., & Nazina, N. (2016). Detection of *n*-alkane biodegradation genes *alkB* and *ladA* in thermophilic hydrocarbon-oxidizing bacteria of the genera *Aeribacillus* and *Geobacillus*. *Microbiology*, 85(6), 693-707.
- Tripathi, S., Kumari, S., Chakraborty, A., Gupta, A., Chakrabarti, K., & Bandyapadhyay, B. K. (2006). Microbial biomass and its activities in salt-affected coastal soils. *Biology and Fertility of Soils*, 42, 273-277.
- Turcatti, G., Romieu, A., Fedurco, M., & Tairi, A. P. (2008). A new class of cleavable fluorescent nucleotides: synthesis and optimization as reversible terminators for DNA sequencing by synthesis. *Nucleic Acids Research*, 36(4), e25. doi: 10.1093/nar/gkn021.
- Tyagi, M., da Fonseca, M. M. R., & de Carvalho, C. C. C. R. (2011). Bioaugmentation and biostimulation strategies to improve the effectiveness of bioremediation processes. *Biodegradation*, 22(2), 231-241.
- Ulrich, A. C., Guigard, S. E., Foght, J. M., Semple, K. M., Pooley, K., Armstrong, J. E., & Biggar, K. W. (2006). Effect of salt on aerobic biodegradation of petroleum hydrocarbons in contaminated groundwater. *Biodegradation*, 20, 27-38.
- Urai, M., Yoshizaki, H., Anzai, H., Ogihara, J., Iwabuchi, N., Harayama, S., Sunairi, M., & Nakajima, M. (2007). Structural analysis of an acidic, fatty acid ester-bonded extracellular polysaccharides produced by pristane-assimilating marine bacterium, *Rhodococcus erythropolis* PR4. *Carbohydrate Research*, 342, 933-942.
- van Beilen, J. B., Li, Z., Duetz, W. A., Smits, T. H. M., & Witholt, B. (2003). Diversity of alkane hydroxylase systems in the environment. *Oil and Gas Science and Technology*, 58, 427-440.
- van Beilen, J. B., & Funhoff, E. G. (2007). Alkane hydroxylases involved in microbial alkane degradation. *Applied Microbiology and Biotechnology*, 74(1), 13-21.
- Van Gestel, K., Mergaert, J., Swings, J., Coosemans, J., & Ryckeboer, J. (2003). Bioremediation of diesel oil-contaminated soil by composting with biowaste. *Environmental Pollution*, 125, 361-368. doi: 10.1016/S0269-7491(03)00109-X.
- Van Hamme, J. D., Singh, A., & Ward, O. P. (2003). Recent advances in petroleum microbiology. *Microbiology and Molecular Biology Review*, 67(4), 503-549.

- Van Sluys, M. A., Monteiro-Vitorello, C. B., Camargo, L. E., Menck, C. F., Da Silva, A. C., Ferro, J. A., Oliveira, M. C., Setubal, J. C., Kitajima, J. P., & Simpson, A. J. (2002). Comparative genomic analysis of plant-associated bacteria. *Annual Review of Phytopathology*, 40, 169-189.
- Varivarn, K., Champa, L. A., Silby, M. W., & Robleto, E. A. (2013). Colonization strategies of *Pseudomonas fluorescens* Pf0-1: activation of soil-specific genes important for diverse and specific environments. *BMC Microbiology*, 13(92). doi: 10.1186/1471-2180-13-92.
- Vasudevan, N., Bharathi, S., & Arulazhagan, P. (2007). Role of plasmid in the degradation of petroleum hydrocarbons by *Pseudomonas fluorescens* NS1. *Journal of Environmental Science and Health*, 42(8), 1141–1146.
- Vazquez, S., Nogales, B., Ruberto, L., Mestre, C., Christie-Oleza, J., Ferrero, M., Bosch, R., & Mac Cormack, W. P. (2013). Characterization of bacterial consortia from diesel-contaminated Antarctic soils: towards the design of tailored formulas for bioaugmentation. *International Biodeterioration and Biodegradation*, 77, 22–30.
- Venkatesh, S., & Dayananda, C. (2008). Properties, potentials, and prospects of antifreeze proteins. *Critical Reviews in Biotechnology*, 28, 57-82.
- Vilchez-Vargas, R., Junca, H., & Pieper, D. H. (2010). Metabolic networks, microbial ecology and 'omics' technologies: towards understanding *in situ* biodegradation processes. *Environmental Microbiology*, 12(12), 3089-3104.
- Voelkerding, K. V., Dames, S., & Durtschi, J. D. (2009). Next-generation-sequencing for clinical diagnostic-principles and application to targeted resequencing for hypertrophic cardiomyopathy. *The Journal of Molecular Diagnostics*, 12(5), 539–551.
- von Wirèn, N., Lauter, F-R., Ninnermann, O., Gillissen, B., Walch-Liu, P., Engels, C., Jost, W., & Frommer, W. B. (2000). Differential regulation of three functional ammonium transporter genes by nitrogen in root hairs and by light in leaves of tomato. *The Plant Journal*, 21(2), 167-175.
- von Wirèn, N., & Merrick, M. (2004). Regulation and function of ammonium carriers in bacteria, fungi, and plants. In: *Molecular mechanisms controlling transmembrane transport. Current Genetics*, vol 9. Springer, Berlin, Heidelberg.
- Walworth, J., Harvey, P., & Snape, I. (2013) Low temperature soil petroleum hydrocarbon degradation at various oxygen levels. *Cold Regions Science and Technology*, 96, 117-121.
- Walworth, J., Pond, A., Snape, I., Rayner, J., Ferguson, S., & Harvey, P. (2007) Nitrogen requirements for maximizing petroleum bioremediation in a sub-Antarctic soil. *Cold Regions Science and Technology*, 48, 84-91.

- Wang, X., & Bartha, R. (1990). Effects of bioremediation on residues, activity and toxicity in soil contaminated by fuel soils. *Soil Biology and Biochemistry*, 22(4), 501-505.
- Wang, W., & Shao, Z. (2013). Enzymes and genes involved in aerobic alkane degradation. *Frontiers in Microbiology*, 4(116).
- Warhurst, A. M., & Fewson, C. A. (1994). Biotransformations catalysed by the genus *Rhodococcus*, *Critical Reviews in Biotechnology*, 14(1).
- Ward, D. M., & Brock, T. D. (1978). Hydrocarbon biodegradation in hypersaline environments. *Applied Environmental Microbiology*, 35(2), 353-359.
- Webster, J., Webster, K., Nelson, P., & Waterhouse, E. (2003). The behaviour of residual contaminants at a former station site, Antarctica. *Environmental Pollution*, 123, 163-179.
- Wentzel, A., Ellingsen, T. E., Kotlar, H. K., Zotchev, S. B., & Throne-Holst, M. (2007). Bacterial metabolism of long-chain *n*-alkanes. *Applied Microbiology and Biotechnology*, 76, 1209-1221.
- White, D. A., Hird, L. C., & Ali, S. T. (2013). Production and characterization of a trehalolipid biosurfactant produced by the novel marine bacterium *Rhodococcus* sp., strain PML026. *Journal of Applied Microbiology*, 115(3), 744-755.
- Whyte, L. G., Slagman, S. J., Pietrantonio, F., Bourbonniere, L., Koval, S. F., Lawrence, J. R., Inniss, W. E., & Greer, C. W. (1999). Physiological adaptations involved in alkane assimilation at a low temperature by *Rhodococcus* strain Q15. *Applied and Environmental Microbiology*, 65(7), 2961-2968.
- Whyte, L. G., Smits, T. H., Labbè, D., Witholt, B., Greer, C. W., & Van Beilen, J. B. (2002) Gene cloning and characterisation of multiple alkane hydroxylase systems in *Rhodococcus* strain Q15 and NRRL B-16531. *Applied and Environmental Microbiology*, 68(12), 5933-5942
- Woese, C. R., & Fox, G. E. (1977). Phylogenetic structure of prokaryotic domain: the primary kingdom. *Proceedings of the National Academy of Sciences of the USA*, 74(11), 5088-5090.
- Woolfenden, E. N. M., Hince, G., Powell, S. M., Stark, S. C., Snape, I., Stark, J. S., & George, S. C. (2011). The rate of removal and the compositional changes of diesel in Antarctic marine sediment. *Science of Total Environment*, 410-411, 205-216.
- Wu, L., Wen, C., Qin, Y., Yin, H., Tu, Q., Van Nostrand, J. D., Yuan, T., et al. (2015). Phasing amplicon sequencing on Illumina MiSeq for robust environmental microbiology community analysis. *BMC Microbiology*, 15(25).

- Xu, X., & Zhu, X. (2004). Treatment of refractory oily wastewater by electro-coagulation process. *Chemosphere*, 56, 889-894.
- Yari, A., & Kargosha, K. (2006). Simple photometric determination of free cyanide ion in aqueous solution with 2,6-dichlorophenolindophenol. *Central European Journal of Chemistry*, 4(2), 329-337.
- Yang, S. Z., Jin, H. J., Wei, Z., He, R. X., Ji, Y. J., Li, X. M., & Yu, S. P. (2009). Bioremediation of oil spills in cold environments: a review. *Pedosphere*, 19(3), 371-381.
- Yeung, C. W., Van Stempoot, D. R., Spoelstra, J., Bickerton, G., Voralek, J., & Greer, C. W. (2013). Bacterial community evidence for anaerobic degradation of petroleum hydrocarbons in cold climate groundwater. *Cold Regions Science and Technology*, 86, 55-68.
- Yoshida, N., Hoashi, J., Morita, T., McNiven, S. J., Nakamura, H., & Karube, I. (2001). Improvement of a mediator-type biochemical oxygen demand sensor for on-site measurement. *Journal of Biotechnology*, 88, 269-275.
- Young, E., Stewart, F., & Dimalanta, E. (2017). The quantitation question: how does accurate library quantitation influence sequencing? *New England BioLabs*, Feature article.
- Zanaroli, G., Di Toro, S., Todaro, D., Varese, G. C., Bertolotto, A., & Fava, F. (2010). Characterization of two diesel fuel degrading microbial consortia enriched from a non acclimated, complex source of microorganisms. *Microbial Cell Factories*, 9(10). doi: 10.1186/1475-2759-9-10.
- Zerbino, D. R. (2010). Using the velvet *de novo* assembler for short-read sequencing technologies. *Current Protocols in Bioinformatics*, 11(5), 1-13.
- Zerbino, D. R., & Birney, E. (2008). Velvet: algorithm for the *de novo* short read assembly using de Bruijn graphs. *Genome Research*, 18(5), 821-829.
- Zock, J. P., Rodriguez-Trigo, G., Pozo-Rodriguez, F., Barbera, J. A., Bouso, L., Torralba, Y., et al. (2007). Prolonged respiratory symptoms in clean-up workers of the *Prestige* oil spill. *American Journal of Respiratory and Critical Care Medicine*, 176(6), 610-616.
- Zock, J. P., Rodriguez-Trigo, G., Rodriguez-Rodriguez, E., Pozo-Rodriguez, F., Gomez, F. P., Fuster, C., et al. (2009). Long-term health effects of the *Prestige* oil spill (Galicia, Spain). *Epidemiology*, 20(6), 242-243.
- Zock, J. P., Rodriguez-Trigo, G., Rodriguez-Rodriguez, E., Espinosa, A., Pozo-Rodriguez, F., Gomez, F. P., et al. (2012). Persistent respiratory symptoms in clean-up workers 5 years after the *Prestige* oil spill. *Occupational and Environmental Medicine*, 69(7), 508-513.