



UNIVERSITI PUTRA MALAYSIA

**MOLECULAR CLONING AND EXPRESSION OF A THERMOSTABLE
 α -AMYLASE FROM *GEOBACILLUS* SP.**

ELIAS KEBEDE KASSAYE

FBSB 2009 23



**MOLECULAR CLONING AND EXPRESSION OF A THERMOSTABLE
 α -AMYLASE FROM *GEOBACILLUS* SP.**

By

ELIAS KEBEDE KASSAYE

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Master of
Science**

September 2009



Abstract of thesis presented to the senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree Master of Science

**MOLECULAR CLONING AND EXPRESSION OF A THERMOSTABLE
 α -AMYLASE FROM *GEOBACILLUS* SP.**

By

Elias Kebede Kassaye

September 2009

Chairman: Abu Bakar Salleh, PhD

Faculty: Biotechnology and Biomolecular Sciences

Starch degrading enzymes like amylase have received great deal of attention because of their perceived technological significance and economic benefits. Selection of a suitable strain is the most significant factor in the amylase production process. On the other hand the screening for a single amylase is difficult because one strain can produce different amylases with different specificities or the amount of amylase produced may be very low. Thus the cloning of one gene directing the synthesis of the desired amylase in a well characterized host like *E. coli* should help greatly in the characterization of new amylases and also allow a significant yield increase. Studies on the screening of amylase producing bacterial strains were carried out on soil and water samples collected from a hot spring located in Slim River, Perak, Malaysia. The bacteria were cultivated in a mineral medium containing soluble starch as the sole carbon source. Three of the isolates namely SR37, SR41, SR74 have demonstrated good activity based on the assay performed using DNS method at 60°C and pH 7.0. The



isolates showed activity of 2.44 U/ml, 4.5 U/ml, 2.05 U/ml for amylopectin and 1.78 U/ml, 3.54 U/ml, 1.65 U/ml for soluble starch, respectively. None of the isolates except SR74 (1.65 U/ml) showed activity at 70°C. Since it showed activity at 70°C, further study was conducted on the isolate SR74 for identification, gene cloning, sequencing and expression for the α -amylase enzyme. Gram staining and morphological studies revealed the isolate was a Gram positive *Bacillus*. Molecular characterization using the 16S rDNA for the isolate SR74 revealed the organism closely related to the members of the genus *Geobacillus*. The fatty acid methyl ester analysis using the Sherlock system also resulted in a typical fatty acid profile of a thermophilic *Geobacillus* and other bacilli. Among them the iso-branched pentadecanoic acid (iso-15:0), hexadecanoic acid (iso-16:0) and heptadecanoic acids (iso-17:0) accounted for 82.26% of the total fatty acids. Iso-15:0 and iso-17:0 were especially abundant. This isolate exhibited anteiso-15:0 (1.05%) and anteiso-17:0 (6.5%) as minor components (7.55% of the total). The isolate was identified as *Geobacillus* sp. SR74. The gene coding for a thermostable α -amylase from *Geobacillus* sp. SR74 was isolated, sequenced and expressed in *Escherichia coli* BL21 (DE3) pLysS. Gene sequencing showed that the enzyme secreted by this isolate shared 98% similarity with *Geobacillus stearothermophilus* α -amylase gene. The ORF of the gene codes for 549 amino acids. The signal peptide comprised 34 amino acids and the remaining 515 amino acids belong to the mature polypeptide. The region encoding the mature α -amylase was heterogeneously expressed in *E. coli* BL21 (DE3) pLysS cells using the pET-32b expression system under the control of the T7 promoter. The mature enzyme had a theoretical molecular weight of 58,547 Daltons and a theoretical *pI* of 5.61. Optimization studies revealed that the highest enzyme activity was



obtained at 16h post induction (32.414 U/ml). The optimum inducer concentration was found to be 0.15mMol L⁻¹ IPTG (39.73 U/ml). With regard to production media, LB (49.53 U/ml) and 0.75YT (51.06 U/ml) were found to be best for optimum production of the recombinant enzyme, while the A_{600nm} of 0.75 (58.3 U/ml) being the best microbial density for inducing the production of the enzyme.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGLONAN MOLEKUL DAN PENGEKSPRESAN α -AMILASE
TERMOSTABIL DARIPADA *GEOBACILLUS SP.***

Oleh

Elias Kebede Kassaye

September 2009

Pengerusi: Professor Dr. Abu Bakar Salleh

Fakulti: Bioteknologi dan Sains Biomolekul

Enzim pendegradasi kanji seperti amylase adalah, merupakan antara enzim yang mendapat perhatian ramai penyelidik memandangkan kepentingannya dalam bidang sains, teknologi dan ekonomi. Oleh itu, pemencilan dan penyaringan strain yang bersesuaian merupakan faktor yang amat penting dalam proses penghasilan amilase. Pada masa yang sama, pemencilan amilase tunggal merupakan satu proses yang sukar berdasarkan kepada kemampuan satu strain untuk menghasilkan pelbagai jenis amilase yang mempunyai spesifikasi berlainan dan ditambah dengan penghasilan enzim yang amat rendah. Oleh yang demikian, pengklonan satu gen yang mengkodkan amilase yang dikehendaki di dalam *E. coli* akan membantu dalam pencirian enzim tersebut di samping meningkatkan penghasilannya. Kajian mengenai pencarian strain bakteria yang menghasilkan amilase telah dijalankan ke atas sampel tanah dan air yang telah diambil daripada



kawasan air panas di Slim River, Perak. Bakteria telah ditumbuhkan di dalam medium mineral yang mengandungi kanji terlarut sebagai sumber utama karbon. Tiga pencilan iaitu SR37, SR41, SR74 telah menunjukkan aktiviti yang baik berdasarkan asai yang dijalankan menggunakan kaedah DNSA pada suhu 60°C dan pH 7.0. Pencilan tersebut masing – masing menunjukkan aktiviti sebanyak 2.44 U/ml, 4.5 U/ml, 2.05 U/ml untuk amilopektin dan 1.78 U/ml, 3.54 U/ml, 1.65 U/ml untuk kanji terlarut. Memandangkan ia telah menunjukkan aktiviti pada suhu 70°C, kajian lanjut telah dijalankan pada pencilan SR74 untuk pengenalpastian (identifikasi), pengklonan gen, penjujukan dan penghasilan untuk enzim α -amilase. Pewarnaan Gram dan ciri morfologi menunjukkan yang pencilan ini adalah *Bacillus* yang mempunyai dinding sel gram positif. Pencirian molekular yang menggunakan 16S rDNA untuk pencilan SR74 menunjukkan organisma ini mempunyai hubungan/pertalian yang rapat dengan genus dari kumpulan *Geobacillus*. Analisis ester metil asid lemak menggunakan sistem Sherlock juga menunjukkan profil asid lemak yang sama dengan *Geobacillus* yang termofilik dan lain – lain *bacillus*. Dalam profil tersebut, cabang asid iso-pentadekanoik (iso-15:0), asid heksadekanoik (iso-16:0) dan asid heptadekanoik (iso-17:0) menyumbang sebanyak 82.26% daripada keseluruhan asid lemak. Iso-15:0 dan iso-17:0 paling banyak didapati. Pencilan ini mempamerkan anteiso-15:0 (1.05%) dan anteiso-17:0 (6.5%) sebagai komponen minor (7.55% daripada keseluruhan). Pencilan ini dinamakan sebagai *Geobacillus* sp. SR74. Gen yang mengkodkan α -amilase termostabil daripada *Geobacillus* sp. SR74 ini telah dipencilkan, dijujukan dan diekspreskan di dalam *Escherichia coli* BL21 (DE3) pLysS. Penjujukan gen menunjukkan enzim yang dirembeskan oleh pencilan ini berkongsi 98% persamaan dengan gen α -amilase daripada *Geobacillus*



stearothermophilus. Gen α -amilase daripada ORF gen ini mengkodkan 549 asid amino. Peptida isyarat terdiri daripada 34 asid amino dan selebihnya iaitu 515 asid amino adalah polipeptida matang. Gen yang mengkodkan α -amilase matang telah diekspreskan secara heterogenus di dalam sel *E. coli* BL21 (DE3) pLysS menggunakan vektor pengekspresan, pET 32-b yang dikawal oleh promoter T7. Secara teori, enzim matang ini mempunyai berat molekul sebanyak 58,547 Daltons dan *pI* 5.61. Tempoh aruhan yang optimum dapat diperhatikan pada 16 jam selepas aruhan (39.734 U/ml). Aruhan optimum menggunakan IPTG adalah pada kepekatan 0.15 mmol/L (39.73 U/ml). Medium terbaik adalah dengan menggunakan 0.75YT (51.06 U/ml) dan LB (49.53 U/ml), sementara ketumpatan mikrob yang terbaik untuk aruhan penghasilan enzim ialah pada A_{600nm} 0.75 (58.3 U/ml).



ACKNOWLEDGEMENTS

First and foremost I would like to thank my supervisory committee chairperson, Professor Dr. Abu Bakar Salleh. His experienced opinion and sound advice have guided the course of this research to its successful conclusion. I wish to express my sincere appreciation to members of the supervisory committee, Professor Dr. Raja Noor Zaliha Raja Abdul Rahman and Professor Dr. Mahiran Basri for their invaluable guidance, encouragement, help and patience through the course of this thesis.

Special thanks are extended to Dr. Adam Leow Thean Chor and Madam Suriana Sabri for the invaluable help (both Technical and Social); Shukuri, Chee Fah, Salsabil and Hisham for making my Malaysian experience possible. I am grateful to all of my colleagues at Enzyme and Microbial Technology Laboratory – both past and present – which have assisted me with insightful comments, advice, technical support, helpful discussions, and for the friendship during the course of this work.

A huge thank you to my parents, for providing me with every opportunity they can to undergo my study, for their many sacrifices, and enduring encouragements. I wish to extend my sincere gratitude to all my siblings for you encouraged me throughout the course of my study.

Finally, I would like to thank the Faculty of Veterinary Medicine (Gondar University) for letting me pursue my Masters Study and Nuffic for the sponsorship.



I certify that a Thesis Examination Committee has met on September 16 2009 to conduct the final examination of Elias Kebede Kassaye on his thesis entitled “Molecular Cloning and Expression of a Thermostable α -Amylase from *Geobacillus* sp.” in accordance with Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 march 1998 The Committee recommends that the candidate be awarded the relevant degree.

Members of the Thesis Examination Committee were as follows:

Suraini Abdul Aziz, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairperson)

Shuhaimi Mustafa, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Sieo Chin Chin, PhD

Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Darah Ibrahim, PhD

Professor
Institute of Life Sciences
Universiti Sains Malaysia
(External Examiner)

Bujang Kim Huat, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 24 December 2009



This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Abu Bakar Salleh, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia (Chairman)

Raja Noor Zaliha Raja Abdul Rahman, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia (Member)

Mahiran Basri, PhD

Professor
Faculty of Science
Universiti Putra Malaysia (Member)

HASANAH MOHD GHAZALI, PhD

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia
Date: 14 January 2010



DECLARATION

I declare that this thesis is the result of my own work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or any other institution.

ELIAS KEBEDE KASSAYE

Date: 10 February 2010



TABLE OF CONTENTS

		Page
ABSTRACT		ii
ABSTRAK		v
ACKNOWLEDGEMENTS		viii
APPROVAL		ix
DECLARATION		xi
LIST OF TABLES		xv
LIST OF FIGURES		xvi
LIST OF APPENDICES		xviii
LIST OF ABBREVIATIONS		xix
CHAPTER		
1	Introduction	1
2	Literature Review	5
	2.1. Glycoside hydrolases	5
	2.1.1. The α -amylase family	6
	2.1.1.1. Characteristics of the α -amylase family	6
	2.1.1.2. The catalytic mechanism of the α -amylase family	8
	2.2. Sources of α -amylases	10
	2.3. Characteristics of α -amylases	11
	2.3.1. Substrate specificity	11
	2.3.1.1 Substrate source	12
	2.3.2. pH optima and pH stability of α -amylases	12
	2.3.3. Temperature optima and temperature stability	13
	2.3.4. Effect of metal ions	16
	2.3.5. Effect of activators/stabilizers	17
	2.3.6. Inhibitors	18
	2.4. Industrial applications of α -amylases	20
	2.5. Production of α -amylases	21
	2.6. Molecular biology of α -amylases	23
	2.6.1. Enzyme engineering	23
	2.6.2. Cloning of α -amylase genes	23
	2.7. Strategies for heterologous expression of recombinant enzymes	25
	2.7.1. Overview of expression systems for the production of recombinant enzymes	25
	2.7.1.1 Eukaryotic expression systems	25
	2.7.1.2. Prokaryotic expression systems	26
	2.7.1.3. <i>Escherichia coli</i> as a host	27
	2.7.3. Construction of recombinant production strains	29
3	Materials and Methods	32
	3.1 Materials	32
	3.2. Sample collection	32
	3.3. Isolation of Bacteria	32
	3.4. Qualitative determination of α -amylase activity through plate	



iodine test	33
3.5. Enzyme production	34
3.6. Quantitative determination of amylase activity	34
3.6.1. The dinitrosalicylic method for estimation of amylase activity	34
3.6.2 Determination of the type of amylase	35
3.7. Identification of microorganism	36
3.7.1. Morphological and physiological study	36
3.7.2. Cellular fatty acids analysis	36
3.7.3. 16S rDNA analysis	37
3.7.4. DNA base composition	38
3.8. Cloning of the α -amylase gene from <i>Geobacillus</i> sp. SR74	39
3.8.1. Preparation and Quantitation of genomic DNA	39
3.8.2. Polymerase chain reaction (PCR) amplification	40
3.8.2.1. Purification of PCR products	40
3.8.3. Genetic transformation of <i>Escherichia coli</i>	42
3.8.3.1. Culture medium	42
3.8.3.2. Preparation of the competent cells	42
3.8.3.3. Transformation of <i>Escherichia coli</i> cells	42
3.8.3.4. Screening for the transformants	43
3.8.3.5. Freezing of stock cells for future use	44
3.8.4. Preparation of plasmid DNA	44
3.8.5. Analysis of transformants	45
3.8.6. Sequencing and comparison	45
3.9. Transformation of Expression Host	46
3.9.1 Preparation for induction	46
3.10. Optimization of intracellular expression of SR74 α -amylase	46
3.10.1. Effect of post induction time on expression of SR74 α -amylase	47
3.10.2. Effect of inducer concentration on SR74 α -amylase expression	47
3.10.3. Effect of media on SR74 α -amylase expression	47
3.10.4. Effect of A_{600nm} on SR74 α -amylase expression	48
4 Results and Discussion	49
4.1. Isolation of a thermophilic bacterium producing α -amylase	49
4.2. Identification of the isolate SR74	52
4.2.1. Identification of the isolate SR74 using the 16S rDNA	56
4.2.2. Phylogenetic tree construction of the isolate SR74	60
4.2.3. Cellular Fatty acid composition of <i>Geobacillus</i> sp. SR74	62
4.3. Genomic DNA extraction	65
4.3.1. Genomic GC content of <i>Geobacillus</i> sp. SR74	68
4.4. Cloning of the α -amylase encoding gene	69
4.4.1. GC content of SR74 α -amylase gene	74
4.4.2. Phylogenetic relationships SR74 α -amylase with other α -amylases	76
4.4.3. Signal peptide prediction	79
4.4.4. Amino acid composition and stability indices of the SR74 α -amylase	82



4.4.5. Conserved regions in the <i>Geobacillus</i> sp. SR74 α -amylase as compared with others	86
4.5. Cloning and Expression of the mature fragment of α -amylase	90
6.5.1. PCR on plasmid using vector and insert primers	90
6.5.2. Sequencing and Restriction Digest analysis	92
4.6. Optimization for the expression of SR74 α -amylase	93
4.6.1. Effect of post induction time on α -amylase production	93
4.6.2. Effect of inducer concentration on the efficiency of induction	95
4.6.3. Effect of growth medium composition on α -amylase production	98
4.6.4. Effect of microbial density on α -amylase production	101
5 Conclusion and Recommendations	105
REFERENCES	108
BIODATA OF STUDENT	132



LIST OF TABLES

Table		Page
1	Known activities of Glycoside hydrolase family 13 enzymes	7
2	Source microorganisms and properties of thermostable starch hydrolyzing enzymes	14
3	Calcium requirement of industrially important starch degrading enzymes	17
4	Concentration effect of metal ions on the activity of α -amylases	19
5	Different fields of application of enzymes belonging to the α -amylase family	20
6	<i>E. coli</i> strains frequently used for heterologous protein production and their key features	28
7	Physiological and Morphological characteristics of the isolate SR74	54
8	Cellular fatty acid composition of <i>Geobacillus</i> sp. SR74 and its phylogenetic neighbour	63
9	The GC content of <i>Geobacillus</i> sp. SR74 α -amylase when compared with others	75
10	The amino acid composition and molecular weight of α -amylases from different sources	84



LIST OF FIGURES

Figure		Page
1	Schematic active site cleft subsite arrangement for glycosyl hydrolases	8
2	The double displacement mechanism and the formation of a covalent intermediate by which retaining glycosyl hydrolases act	9
3	Schematic presentation of the salient features and sequence elements of a prokaryotic expression vector	30
4	Slim river hot spring located in Perak Malaysia	33
5	Activity profile (U/ml) of the selected isolates for starch and amylopectin	50
6	The gel photo for PCR amplified 16s rDNA gene of the isolate SR74	57
7	The sequence of the 16s rDNA gene for the isolate SR74	59
8	Evolutionary relationships of <i>Geobacillus</i> sp. SR74 with other 15 <i>Geobacillus</i> organisms	61
9	Agarose gel photo of purified genomic DNA from <i>Geobacillus</i> sp. SR74	66
10	The gel photo of PCR amplified α -amylase gene from <i>Geobacillus</i> sp. SR74	71
11	The ORF of thermostable α -amylase gene from <i>Geobacillus</i> sp. SR74	73
12	Phylogenetic analysis of thermostable SR74 α -amylase from <i>Geobacillus</i> sp. SR74	78
13	Signal peptide prediction of SR74 α -amylase gene	81



14	The four conserved regions I-IV of <i>Geobacillus</i> sp. SR74 α -amylase	87
15	Gel photo for the PCR amplified thermostable α -amylase gene product	90
16	PCR product showing the presence of the insert in the recombinant plasmid	91
17	Restriction digest of the recombinant plasmid pET-32b/SR74	92
18	Expression of SR74 α -amylase gene in <i>E. coli</i> BL21 (DE3) pLysS harboring recombinant plasmid pET-32b/SR74 α -amylase when induced using 1mM L ⁻¹ IPTG at A _{600nm} ~0.5 up to 24h	94
19	Expression of SR74 α -amylase gene in <i>E. coli</i> BL21 (DE3) pLysS harboring recombinant plasmid pET-32b/SR74 when induced using different concentrations of IPTG (mM L ⁻¹) at A _{600nm} ~0.5	96
20	Effect of media on the expression of SR74 α -amylase gene in <i>E. coli</i> BL21 (DE3) pLysS harboring recombinant plasmid pET-32b/SR74 when induced using 0.15mmol L ⁻¹ IPTG at A _{600nm} ~0.5	99
21	Effect of microbial density on the Expression of SR74 α -amylase gene in <i>E. coli</i> BL21 (DE3) pLysS harboring recombinant plasmid pET-32b/SR74 when induced using 0.15mmol L ⁻¹ IPTG at different A _{600nm}	102



LIST OF APPENDICES

Appendix	Page
I	122
II	123
III	124
IV	125
V	126
VI	127
VII	128



LIST OF ABBREVIATIONS

A: Adenine

α : Alpha

β : Beta

C: Cytosine

CAZy: Carbohydrate active enzymes

Da: Daltons

dH₂O: distilled water

DNA: Deoxyribonucleic acid

DNSA: Dinitrosalicylic acid

DSMZ: Deutsche Sammlung Von Microorganismen

EDTA: Ethylene di-amine tetrachloro acetate

g/L: Gram per Litre

g :Gram

G: Guanine

GH: Glycosyl hydrolases

GOC: Groundnut oil cake

GTE: Glucose/Tris-Hcl/EDTA

h :Hour

IPTG: Isopropyl- β -D-galactoside

IUB-MB: International union for biochemistry and molecular biology

KDa: Kilo Daltons

L: Litre

LB: Luria Bertani

M: Molar

μ g: Microgram

μ l: Microliter

μ m: Micrometer

mg: milligram



min: Minute
ml: milliliter
mM: millimolar
mRNA: messenger RNA
NB: Nutrient broth
nm: nanometer
OD: Optical density
Ori: Origin of replication
ORF: Open Reading Frame
PCI: Phenol/Chloroform/Isoamyl alcohol
PCR: Polymerase Chain Reaction
RBS: Ribosomal Binding Site
RNA: Ribonucleic acid
rRNA: ribosomal RNA
SD : Shine Dalgarno
SDS–PAGE: Sodium dodecyl sulfate–polyacrylamide gel electrophoresis
Smf : submerged fermentation
SOC: Super optimal broth with catabolite repression
SP: Signal peptide
SR: Slim River
SSF: Solid state fermentation
T: Thiamine
TIM: Triose phosphate isomerase type barrels
TIR: Translation initiation region
U/mg: Unit per milligram
U/ml: Unit per milliliter
UV: Ultra violet
v/v :Volume per volume
WB: Wheat bran



w/v : Weight per volume

YT: Yeast Trypt



CHAPTER 1

Introduction

Enzymes or biocatalysts are the key components of industrial biotechnology. Their abilities to convert the substrates to desired products at mild conditions and in greatly specific manners are essential for sustainable development of various industries including food, textile and pharmaceuticals. In recent years the new potential of using microorganisms as biotechnological sources of industrially relevant enzymes has stimulated renewed interest in the exploration of extra-cellular enzymatic activity in several microorganisms (Gupta *et al.*, 2003; Omemu *et al.*, 2005; Yamabhai *et al.*, 2008).

Starch degrading enzymes like amylase have received great deal of attention because of their perceived technological significance and economic benefits. In terms of energy utilization and process simplicity, amylase conversion of raw starches is believed to be superior to the conventional method that makes use of pre-gelatinized starch as substrate. Amylases are among the most important enzymes having a great significance with extensive biotechnological application in a number of industrial processes such as in the food, fermentation, textiles, paper industries and in detergent producing processes comprising about 30% of the world's enzyme production. They would be potentially useful in the pharmaceutical and fine chemicals industries if enzymes with suitable properties can be prepared (Omemu *et al.*, 2005; Hmidet *et al.*, 2008).

Due to the increasing demand for amylase enzymes in various industries, there is enormous interest in developing enzymes with novel properties such as raw starch degrading amylases suitable for industrial applications and their cost effective



production techniques. Amylases can be derived from several sources such as plants, animals and micro-organisms; however the enzymes from microbial sources generally meet industrial demands. Microbial amylases have successfully replaced the chemical hydrolysis of starch in starch processing industries. The major advantages of using microorganisms for the production of amylases is the cost effective bulk production capacity, consistency, less time and space required for production, ease of process modification and optimization and microbes are relatively easy to manipulate to obtain enzymes of desired characteristics (Pandey *et al.*, 2000; Gupta *et al.*, 2003; Asgher *et al.*, 2007).

Genetic engineering has been used extensively for cloning of amylase producing strains, mainly on α -amylases and glucoamylases in order to achieve desirable characteristics in the cloned host. The purpose of gene cloning can be amongst others, the expression of thermo stable enzymes, higher enzyme productivity and co-expression of two enzymes by the same organism (Pandey *et al.*, 2000).

Escherichia coli is one of the most useful bacteria as a host cell in the field of gene engineering because of high-level expression and synthesis of gene products by utilizing *E. coli*. This organism has been used extensively for cloning of exogenous gene and expression of exogenous proteins. *E. coli* has been the “factory” of choice for the expression of many proteins because its genome has been fully mapped and the organism is easy to handle; grows rapidly; requires an inexpensive, easy-to-prepare medium for growth; and secretes protein into the medium which facilitates recovery of the protein (Coronado *et al.*, 2000; Kang *et al.*, 2004; Shiina *et al.*, 2004).



Demand for novel amylases worldwide is increasing day by day, as these enzyme application spectra are spreading in various industrial sectors. The classical approach is the isolation of microbial species, which produce novel enzyme from exotic environments and would offer a competitive advantage over the existing products. Subsequent characterization of these microbes under fermentation conditions to optimize the enzyme production properties plays vital role in evaluation of economic significance (Prakasham *et al.*, 2007).

Although amylases can be produced by several micro-organisms, it remains a challenging task to obtain a strain capable of producing commercially acceptable yields. Selection of a suitable strain is the most significant factor in the amylase production process. Each application of amylases requires unique properties with respect to specificity, stability, temperature and pH dependence. Screening of microorganisms with higher amylase activities could therefore, facilitate the discovery of novel amylases suitable to new industrial applications (Pandey *et al.*, 2000; Asgher *et al.*, 2007). On the other hand the screening for a single amylase is difficult because one strain can produce different amylases with different specificities or the amount of amylase produced may be very low. Thus the cloning of one gene directing the synthesis of the desired amylase in a well characterized host like *E. coli* should help greatly in the characterization of new amylases and also allow a significant yield increase (Ozcan *et al.*, 2001).

The general objective of the current study was therefore to isolate thermostable α -amylase producing bacteria, to clone and express the thermostable α -amylase gene.

