

UNIVERSITI PUTRA MALAYSIA

PHYSIOLOGICAL AND PSYCHOLOGICAL HEALTH BENEFITS OF URBAN GREEN SPACE

MOHAMED DANIEL BIN MOHAMED MOKHTAR

FH 2018 8



PHYSIOLOGICAL AND PSYCHOLOGICAL HEALTH BENEFITS OF URBAN GREEN SPACE



Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

PHYSIOLOGICAL AND PSYCHOLOGICAL HEALTH BENEFITS OF URBAN GREEN SPACE

Ву

MOHAMED DANIEL BIN MOHAMED MOKHTAR

October 2017

Chair : Nor Akmar Abdul Aziz, PhD

Faculty: Forestry

This study explores the physiological and psychological effects of urban green space by using stress as a determinant of well-being. This study also aims to provide scientific evidence supporting the use of urban green space as a method to relieve stress. The subjects of this study were 20 male students which their homogeneity was controlled. The study area chosen for this study was Taman Botani Perdana, Kuala Lumpur and Jalan Bukit Bintang was used as a control area. The first day the participants went to Taman Botani Perdana and on the second day to Jalan Bukit Bintang. Participants were asked to walk along a given route in both areas in which the time was set similarly in both areas (20 minutes). Saliva samples were collected before and after walking in both areas along with blood pressure measurements. In addition, psychometric tests namely the Profile of Mood States (POMS), Positive and Negative Affect Schedule (PANAS), Perceived Stress Scale (PSS) and the Restoration Outcome Scale were done to identify the psychological effects of urban green space use. The results indicated that salivary cortisol concentration levels among participants in the urban green space is significantly lower than the participants in the city. Blood pressure readings also showed a significant decrease in value among participants in UGS for diastolic pressure. In the Restoration Outcome Scale (ROS), participants in the UGS setting reported a higher score for all six statements in the scale compared to the City setting with a significance of p < 0.01. The POMS test indicated a clear distinction between the restorative effects of urban green space and city as the Total Mood Disturbance (TMD) of participants in the urban green space showed negative in value while participants in the city showed positive TMD. Perceived Stress scores in the urban green space is significantly higher in mean value compared to the city setting in both before and after period. When summing the scores of positive and negative subscales of the PANAS test, a significant difference can be found at the city setting for both positive and negative subscales. Positive attitudes decreased while negative attitudes increased after walking in the city. In the UGS setting however, only positive attitudes increased.

FAEDAH KESIHATAN FISIOLOGI DAN PSIKOLOGI KAWASAN HIJAU BANDAR

Oleh

MOHAMED DANIEL BIN MOHAMED MOKHTAR

Oktober 2017

Pengerusi : Nor Akmar Abdul Aziz, PhD

Fakulti : Perhutanan

Kajian ini meneroka kesan fisiologi dan psikologi kawasan hijau bandar dengan menggunakan tekanan (stress) sebagai penentu kesihatan. Kajian ini juga bertujuan untuk mengemukakan bukti saintifik yang menyokong penggunaan kawasan hijau bandar sebagai kaedah untuk mengurangkan tekanan. Subjek kajian ini adalah 20 pelajar lelaki yang dikendalikan secara homogen. Tempat yang dipilih untuk kajian ini ialah Taman Botani Perdana, Kuala Lumpur dan Jalan Bukit Bintang yang digunakan sebagai kawasan kawalan. Pada hari pertama peserta bergerak ke Taman Botani Perdana dan pada hari kedua ke Jalan Bukit Bintang. Peserta diminta untuk berjalan di sepanjang laluan yang ditetapkan di kedua-dua kawasan di mana masa ditetapkan sama dalam kedua-dua kawasan (20 minit). Sampel air liur dikumpulkan sebelum dan selepas berjalan di kedua-dua kawasan bersama dengan pembacaan tekanan darah. Di samping itu, ujian psikometrik jaitu Profil Keadaan Mood (POMS), Jadual Kesan Positif dan Negatif (PANAS), Skala Tekanan Dipahami (PSS) dan Skala Hasil Pemulihan (ROS) telah dilakukan untuk mengenal pasti kesan psikologi penggunaan ruang hijau bandar. Keputusan menunjukkan bahawa tahap kepekatan kortisol di kalangan peserta di ruang hijau bandar adalah jauh lebih rendah daripada peserta di bandar. Bacaan tekanan darah juga menunjukkan pegurangan nilai yang signifikan di kalangan peserta di UGS untuk tekanan diastolik. Hasil ujian Skala Output Pemulihan (ROS), peserta dalam kawasan hijau bandar melaporkan skor yang lebih tinggi untuk semua enam kenyataan pada skala berbanding dengan peserta di kawasan bandar dengan signifikan p < 0.01. Ujian POMS menunjukkan perbezaan yang jelas antara kesan pemulihan kawasan hijau bandar dan bandar kerana Jumlah Gangguan Mood (TMD) peserta di ruang hijau bandar menunjukkan nilai yang negatif manakala peserta di bandar menunjukkan nilai yang positif. Skor Tekanan yang dialami di kawasan hijau bandar jauh lebih tinggi dalam nilai min berbanding dengan bandar pada kedua-dua tempoh sebelum dan selepas. Dengan menjumlahkan skor subskala positif dan negatif ujian PANAS, perbezaan yang signifikan boleh didapati di kawasan bandar untuk kedua-dua subskala positif dan negatif. Nilai min sikap positif menurun manakala sikap negatif meningkat selepas berjalan di bandar. Walaubagaimanapun, pada kawasan hijau bandar, hanya sikap positif yang meningkat.



ACKNOWLEDGEMENTS

This thesis would have been impossible if not for a special circle of people whom I am motivated and inspired by. I would like to express my greatest of gratitudes to Dr. Nor Akmar Abdul Aziz, my supervisor for being patient and insightful, my co-supervisor, Assoc. Prof. Dr. Manohar Mariapan, with his great knowledge and guidance, my father Mokhtar Shah for his countless advices, my beloved Cindy Usun Sigau for being there to support me, and all the people that have helped throughout this journey.



I certify that a Thesis Examination Committee has met on 31 October 2017 to conduct the final examination of Mohamed Daniel bin Mohamed Mokhtar on his thesis entitled "Physiological and Psychological Health Benefits of Urban Green Space" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Kamziah bt Abd Kudus, PhD

Associate Professor Faculty of Forestry Universiti Putra Malaysia (Chairman)

Azlizam bin Aziz, PhD

Associate Professor Faculty of Forestry Universiti Putra Malaysia (Internal Examiner)

Nurhayati binti Abdul Malek, PhD

LAr. Senior Lecturer Universiti Teknologi MARA Malaysia (External Examiner)

> RUSLI HAJI ABĎULLAH, PhD Professor and Deputy Dean School of Graduate Studies

Universiti Putra Malaysia

Date: 30 August 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Nor Akmar Abdul Aziz, PhD

Senior Lecturer Faculty of Forestry Universiti Putra Malaysia (Chairman)

Manohar Mariapan, PhD

Associate Professor Faculty of Forestry Universiti Putra Malaysia (Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature:		Date:	
•			

Name and Matric No.: MOHAMED DANIEL BIN MOHAMED MOKHTAR (GS44397)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Nor Akmar Abdul Aziz	
PM	
1 20 PT 0 2 PT 0 2	
Manohar Mariapan	

TABLE OF CONTENTS

				Page
ABSTRA ABSTRA ACKNOW APPROW DECLAR LIST OF LIST OF	K VLEDGE AL ATION TABLES FIGURE	S ES		i iv v vii xi xii
СНАРТЕ	R			
1	1.1 1.2 1.3 1.4	Proble Objec Signif	ground of the Study em Statement	1 1 2 3 4 4
2		Urban The Be 2.2.1 2.2.2 2.2.3 2.2.4	Green Space enefits of Urban Green Space Psychological Benefits Physiological Benefits Environmental Benefits Social Benefits Economic Benefits	5 5 5 6 7 7 8
	2.3	Measu	rements of Physiological Health ts of Urban Green Space Salivary Cortisol as Biomarker for Stress Levels Circadian Rhythm and Normal Values Systolic, Diastolic Blood	9 9 10 11
	2.4	Benefi 2.4.1 2.4.2 2.4.3 2.4.4	Pressure and Pulse Rate frements of Psychological Health ts of Urban Green Space Profile of Mood States Restoration Outcome Scale Perceived Stress Scale Positive and Negative Affect Schedule ptual Framework	11 12 12 12 13
3	-	HODOI Study	_OGY	15 15

	3.2	Participants	15
	3.3	Consent and Ethics	16
	3.4	Study Design and Protocol	16
	3.5	Data Collection	17
		3.5.1 Physiological Measurements	17
		3.5.2 Psychological Measurements	18
	3.6		18
		3.6.1 Paired T-test	18
		3.6.2 Correlation	18
	3.7	Potential Risks of the Study	19
4	RES	ULTS AND DISCUSSION	20
	4.1	Blood Pressure Changes	20
	4.2	Salivary Cortisol	22
	4.3	Psychological Indices	23
		4.3.1 Restoration Outcome Scale	23
		4.3.2 Profile of Mood States	24
		4.3.3 Perceived Stress Scale	25
		4.3.4 Positive and Negative Affect Schedule	26
	4.4	Relationship between Psychological and	29
		Physiological Indices	
5	IMPI	LICATIONS OF STUDY,	32
	RECOMMENDATION AND CONCLUSION		
	5.1	Conclusion	32
	5.2	Recommendation for future research	32
	5.3	Implications of study	33
REFERENC	CES		34
APPENDIC	ES		39
BIODATA	OF ST	UDENT	50
LIST OF PUBLICATIONS			51

LIST OF TABLES

Table		Page
4.1	Descriptive of ROS statements and the p-value comparison between urban green space and city outcomes. $M = \text{mean}$, $SD = \text{standard deviation}$. Significance at $p < .05$.	23
4.2	Relationship between ROS and other psychological indices in urban green space.	30
4.3	Relationship between ROS and other	30

LIST OF FIGURES

Figure		Page
2.1	Benefits of urban green space adapted from	6
2.2	Schipperijn (2010). Normal values of salivary cortisol during the day of a population adapted from Kirschbaum and Hellhammer (2000).	10
2.3 3.1	Conceptual framework. A view on Jalan Bukit Bintang (left) and Taman Botani Perdana (right).	14 15
3.2 4.1	Study design. Mean comparison of systolic blood pressure, diastolic blood pressure and pulse rate in both urban green space and city. * indicates significance at <i>p</i> < .05 verified by paired t-test. Values are means ± standard error.	17 21
4.2	Comparison of the mean values of salivary cortisol levels between urban green space and the city before and after walking. * indicates significance at p < .05 verified by paired t-test. Values are means ± standard error.	22
4.3	Comparison between the mean values of Profile of Mood States subscales at two measurement periods in both urban green space and city. TMD; Total mood disturbance. (*) indicates significance at p < .05.	24
4.4	Comparison of mean values of the PSS scores at different time periods in both urban green space and city. Values are mean ± standard error; n = 20; ** p < .01; significant differences via paired t-test.	25
4.5	Mean value comparison of positive subscales at two measurement periods in both urban green space and city. Values are mean ± standard error, n = 20; ** p < .01; * p < .05; significant differences via paired t-test.	27
4.6	Mean value comparison of negative subscales at two measurement periods in both urban green space and city. Values are mean ± standard error, n = 20; ** p < .01; * p < .05; significant differences via paired t-test.	28
4.7	Comparison mean values between positive and negative attitudes scores in both urban green space and city. Mean \pm standard error; $n = 20$; ** $p < .01$; * $p < .05$; significant differences via paired t-test.	29

LIST OF ABBREVIATIONS

ART Attention Restoration Theory

EEG Electroencephalogram

ELISA Enzyme-linked immunosorbent assay

HPA Hypothalamus-pituitary-adrenal activation

mmHg Milimieters of mercury

nmol/l Nanomoles per litre

PANAS Positive and Negative Affect Schedule

POMS Profile of Mood States

PSS Perceived Stress Scale

ROS Restoration Outcome Scale

TMD Total mood disturbance

UGS Urban green space



CHAPTER 1

INTRODUCTION

1.1 Background of the study

Urban green space is defined as all publicly owned and publicly accessible open space with a high degree of cover by vegetation e.g. parks, woodlands, nature areas and other green spaces (Schipperijn et al., 2010). The use of urban green space is defined in general as any sort of visit to an urban green space, without looking at the duration of stay, the motivation of visit or the activity done; e.g. passing through on the way to a destination is also counted as use (Schipperijn et al., 2010). Any sort of interaction with an urban green space is considered as use as the many benefits of an urban green are passive or in other words intangible. These passive benefits are of or closely related to emotional or psychological responses of people. Chiesura (2004) states; despite their intangible and immaterial nature, these services (of urban parks) provide clear benefits to people, whose loss can have serious socio-economic consequences. Urban green space is often considered to be essential for urban dwellers as the benefits provided by them are extensive.

Green spaces within urban settings are believed to be experiencing pressures and threats as a result of urban growth. Therefore, there is a prevalent concern that urban sprawl and rapid expansion of cities occurring all over the globe can isolate urban dwellers from direct contact with nature (Willson, 1984). These concerns are more dominant in developing countries where cities lack proper development planning especially in terms of landscapes. The United Nations (UN) has estimated that more than 50% of the world's population are already living in urban areas and this percentage is forecasted to rise rapidly over the coming years (United Nations, 2014). This leads to more expansion of cities to provide housing, employment opportunities, roads and other infrastructures that may degrade the natural environment.

Willson's (1984) "biophillia" hypothesis states that humans possess a deep-seated biological need for connections and contact with nature, which can be understood as living systems other than that of human beings. This is where the role of urban green spaces comes into hand. As more areas are given way for urbanization and development, humans are starting to lose contact with nature. In addition, from a social and political context, the role of urban green spaces, particularly park and recreation settings, have also been related to other goals such as environmental preservation, community and economic development, rather than focused on public health.

In today's society, people are faced with work pressures, noise pollution, and other stressors particularly in urban areas. This phenomenon drives people to seek out relief and physical activity through outdoor recreation in restorative environments. Urban sprawl has also been linked directly to increment of obesity rates (Ewing et al., 2008). With obesity, risk of other diseases such as cardiovascular diseases, hypertension, diabetes and certain cancers increases many folds (Ramachandran & Snehalatha, 2010). For urbanites, urban green spaces provide the most ready access to restorative environments (Maller et al., 2006). Urban green spaces also act as a platform for people to exercise and be inspired to be more physically active. In an urban environment, the main contribution to the enhancement of quality of life in terms of perceived health conditions and environmental quality, as well as the affective and cognitive attachment to the place of residence, is thus provided by urban green spaces and their availability and accessibility (Conedera et al., 2015).

In spite of the recognised benefits of urban green spaces, the frequency of park use amongst Malaysians is still low (Malaysian Quality of Life, 2002). Only 20.2% of Malaysian families are engaged in recreation activities once a week; less than half of Malaysian families do so once a month. Most Malaysian cities lack a well-functioning green infrastructure; for example, poor accessibility to parks (National Urbanization Policy, 2006). According to a study by Sreetheran & Adnan (2007), Kuala Lumpur does not have a proper green network that links all existing open spaces. This is a problem, as accessibility and connectivity are important prerequisites for green space usage. Contrary to the belief that shade is much appreciated in a tropical hot climate, the availability of green space in large cities such as Kuala Lumpur is still lacking, thus, forcing people to stay indoors in air-conditioned rooms. This eventually would lead to low levels of physical activity which is one of the key reasons of the high obesity rate in the country. The boom in studies concerning physical activity fostered through urban green spaces and the built environment should be used as a basis to develop urban green spaces in large cities of Malaysia such as Penang and Johor Bahru.

Moreover, Malaysia is well known as a multi-racial nation consisting of various different races and ethnic groups such as Malay, Indian, Chinese, Kadazan, Iban, and Dayak. Previous studies for instance by Nor Akmar (2012) reported different use patterns of users depending on their demographic profile. For example, Malay users and Indian users reported different levels of usage of urban green spaces and also different use patterns. However, the studies on urban green space use in Malaysia also have little objective data on physical health as most of the studies use self-reported data.

1.2 Problem Statement

Urban green space such as parks are underutilized and somewhat neglected by Malaysians. With the low frequency of park users, it is difficult to justify the existence and the need to have an urban park. Hence, by studying the physiological and psychological restorative outcomes of urban green space through objective data and self-reported data accentuates the importance of it.

Existing studies on the restorative effects of urban green spaces have been extensive throughout the years. However, there is little objective or experimental data that could further provide solid backing in regard to the physical health of an individual after using urban green spaces. Most research on physical health benefits in the best only studied on self-reported data (Nor Akmar et al. 2012; Schipperijn, 2010; Pietila et al., 2015). The hypothesis of psycho-physiological benefits of natural environments has only hardly faced experimental tests (Hartig et al., 2003). (Bell et al., 2008) has also stated that only limited numbers of studies use objective measurements such as blood pressure monitoring, to assess physical health benefits. Majority of other studies utilize self-reported data.

Although it has been hypothesized by many researchers that level of physical activity increases well-being, there is no clear indication as how beneficial physical activity in urban green spaces is based on objective data as it is lacking. The gap in this knowledge needs to be addressed to better understand the benefits of urban green spaces which may be useful in public health and urban green space planning and management. Furthermore, regional studies on stress levels has always been limited to questionnaires whether in the form of open-ended questions or ratings and scales. While subjective ratings on stress may be viable, experimental tests on stress needs to be done in order to provide solid objective data. Therefore, a common biomarker for stress i.e., cortisol, must be measured to further understand the physiological effects of urban green space.

In addition, in order to objectively understand whether urban green space or natural environments (e.g. a park) has more benefits physiologically and psychologically, research needs to be done on the difference between engaging in the same physical activity in a natural environment and a more built or in other words, synthetic environment such as the city. With the aforementioned issues, the following research questions can be forwarded;

- 1. Is there any difference between walking in the city and walking in an urban park physiologically?
- 2. What are the perceived stress and other psychological parameters experienced by users in both city and urban park settings?

1.3 Objective

The main objective of this experimental research is to discover whether use of urban green space affaects physical health based on objective measurements

and also how it affects mental health and well-being by comparing it with the environment of a city. In order to meet this aim, the following research objectives were formulated;

- To determine the stress levels of users by measuring salivary cortisol concentration.
- To explore the effect of urban green space on blood pressure reading of users.
- To identify the restorative psychological effects of urban green space on users.

1.4 Significance of the Study

The results of this study could provide insight in the health promotion that urban green space offers. In light of the urbanization phenomenon that is occurring in Malaysia, stress, which is a well-known determinant of health, and its relation to use of urban green space can be further examined. The results would also justify on the need on preserving urban green spaces, particularly urban parks, as it is the easiest form of natural environment access for urbanites. In a broader and general outlook, this study is in line with the Malaysian National Landscape Policy (2010), which strives for developing urban landscapes that is conducive and healthy for the nation. This study's results could also be beneficial in the field of environmental psychology which plays a big role in helping urbanites' mental well-being.

1.5 Limitation of the Study

Due to a limited budget and a complex field study design, this study utilizes a small sample (N=20) in which the participants are controlled. The participants consist of male students of a similar age range and from the same course to ensure consistent background and IQ level. The age of the participants were in average 23.1 years old. Female samples were excluded as other outlying variables will need to be considered particularly menstrual cycle stages. Therefore, the results of this study is much more suitable to be extrapolated to young adult males.

REFERENCES

- Agyemang, C., van Hooijdonk, C., Wendel-Vos, W., Ujcic-Voortman, J. K., Lindeman, E., Stronks, K., & Droomers, M. (2007). Ethnic differences in the effect of environmental stressors on blood pressure and hypertension in the Netherlands. *BMC Public Health*, 7, 118-128.
- Akbari, H., Pomerantz, M., & Taha, H. (2001). Cool surfaces and shade trees to reduce energy use and improve air quality in urban areas. *Solar energy*, 70(3), 295-310.
- American Heart Association (2015, July). All about heart rate (pulse) [Web log post]. Retrieved January 2017, from http://www.heart.org
- American Heart Association (2016, October). Understanding blood pressure [Web log post]. Retrieved January 2017, from http://www.heart.org
- Bedimo-Rung, A. L., Mowen, A. J., & Cohen, D. A. (2005). The significance of parks to physical activity and public health: a conceptual model. *American Journal of Preventive Medicine*, 28(2), 159-168.
- Bell, S., Hamilton, V., Montarzino, A., Rothnie, H., Travlou, P., & Alves, S. (2008). Greenspace and quality of life: a critical literature review. *Greenspace Scotland*, (August), 75.
- Bolund, P., & Hunhammar, S. (1999). Ecosystem services in urban areas. *Ecological Economics*, *29*(2), 293–301.
- Chiesura, A. (2004). The role of urban parks for the sustainable city. Landscape and Urban Planning, 68(1), 129–138.
- Conedera, M., Biaggio, A. Del, Seeland, K., Moretti, M., & Home, R. (2015). Urban Forestry & Urban Greening Residents' preferences and use of urban and peri-urban green spaces in a Swiss mountainous region of the Southern Alps. *Urban Forestry and Urban Greening*, *14*(1), 139–147.
- Cohen, D. A., McKenzie, T. L., Sehgal, A., Williamson, S., Golinelli, D., & Lurie, N. (2007). Contribution of public parks to physical activity. *American Journal of Public Health*, *97*(3), 509–514.
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24(4), 385-396.
- de Vries, S., Verheij, R. A., Groenewegen, P. P., & Spreeuwenberg, P. (2003). Natural environments Healthy environments? An exploratory analysis of the relationship between greenspace and health. *Environment and Planning A*, *35*(10), 1717–1731.

- Ewing, R., Schmid, T., Killingsworth, R., Zlot, A., & Raudenbush, S. (2008). Relationship between urban sprawl and physical activity, obesity, and morbidity. *Urban Ecology: An International Perspective on the Interaction Between Humans and Nature*, 18(1), 567–582.
- Fukuda, Y., Nakamura, K., & Takano, T. (2004). Wide range of socioeconomic factors associated with mortality among cities in Japan. *Health Promotion International*, 19(2), 177–187.
- Giles-Corti, B., Broomhall, M. H., Knuiman, M., Collins, C., Douglas, K., Ng, K., Donovan, R. J. (2005). Increasing walking: how important is distance to, attractiveness, and size of public open space? *American Journal of Preventive Medicine*, *28*(2), 169–176.
- Hansmann, R., Hug, S. M., & Seeland, K. (2007). Restoration and stress relief through physical activities in forests and parks. *Urban Forestry and Urban Greening*, 6(4), 213–225.
- Hartig, T., Evans, G. W., Jamner, L. D., Davis, D. S., & Gärling, T. (2003). Tracking restoration in natural and urban field settings. *Journal of Environmental Psychology*, 23(2), 109–123.
- Hartig, T., Lindblom, K., & Ovefelt, K. (1998). The home and near-home area offer restoration opportunities differentiated by gender. *Scandinavian Housing and Planning Research*, *15*(4), 283-296.
- Hillsdon, M., Panter, J., Foster, C., & Jones, A. (2006). The relationship between access and quality of urban green space with population physical activity. *Public Health*, *120*(12), 1127–1132.
- Hodgkison, S., Hero, J. M., & Warnken, J. (2007). The efficacy of small-scale conservation efforts, as assessed on Australian golf courses. *Biological Conservation*, 135(4), 576-586.
- Jim, C. Y., & Chen, W. Y. (2006). Recreation-amenity use and contingent valuation of urban greenspaces in Guangzhou, China. *Landscape and Urban Planning*, 75(1–2), 81–96.
- Kaczynski, A., & Henderson, K. (2007). Environmental correlates of physical activity: a review of evidence about parks and recreation. *Leisure Science*, 29 (March), 315–354.
- Kaplan, S. (1995). The Restorative Benefits of Nature: Toward an Integrative Framework. *Journal of Environmental Psychology*, *15*, 169–182.
- Kaplan, R., & Kaplan, S. (1989). *The experience of nature: A psychological perspective*. CUP Archive.

- Kawakami, K., Kawamoto, M., Nomura, M., Otani, H., Nabika, T., & Gonda, T. (2004). Effects of phytoncides on blood pressure under restraint stress in SHRSP. *Clinical and Experimental Pharmacology and Physiology*, 31(s2), 27–28.
- Kirschbaum, C., & Hellhammer, D. H. (2000). Salivary Cortisol. *Encyclopedia of Stress*.
- Korpela, K. M., Ylén, M., Tyrväinen, L., & Silvennoinen, H. (2008). Determinants of restorative experiences in everyday favorite places. *Health and Place*, *14*(4), 636–652.
- Kweon, B. S., Sullivan, W. C., & Wiley, A. R. (1998). Green Common Spaces and the Social Integration of Inner-City Older Adults. *Environment and Behavior*, 30(6), 832-858.
- Lee, J., Park, B. J., Tsunetsugu, Y., Ohira, T., Kagawa, T., & Miyazaki, Y. (2011). Effect of forest bathing on physiological and psychological responses in young Japanese male subjects. *Public Health*, *125*(2), 93–100.
- Li, Q., Morimoto, K., Nakadai, A., Inagaki, H., Katsumata, M., Shimizu, T., ... Kawada, T. (2007). Forest bathing enhances human natural killer activity and expression of anti-cancer proteins. *International Journal of Immunopathology and Pharmacology*, 20(s2), 3–8.
- Maas, J., Verheij, R. A., Groenewegen, P. P., De Vries, S., & Spreeuwenberg, P. (2006). Green space, urbanity, and health: how strong is the relation?. *Journal of Epidemiology and Community Health*, 60(7), 587-592.
- MacKenzie, B. (2001) *Profile of Mood States (POMS)* [Web log post] Available from: https://www.brianmac.co.uk/poms.htm [Accessed 23/5/2015].
- Malaysian Quality of Life Report (2002). Retrieved September 15 2016 from http://www.epu.gov.my/en/kualiti-hidup-malaysia-2002.
- Maller, C., Townsend, M., Pryor, A., Brown, P., & St Leger, L. (2006). Healthy nature healthy people: "contact with nature" as an upstream health promotion intervention for populations. *Health Promotion International*, 21(1), 45–54.
- Mansor, M., Said, I., & Mohamad, I. (2012). Experiential Contacts with Green Infrastructure's Diversity and Well-being of Urban Community. *Procedia - Social and Behavioral Sciences*, 49, 257–267.
- McNair, D. M., Lorr, M., & Droppleman, L. F. (1971). *Manual for the Profile of Mood States*. San Diego, CA: Educational and Industrial Testing Services

- National Landscape Policy (2010). National Landscape Department, Ministry of Housing and Local Government.
- National Urbanization Policy (2006). Ministry of Urban Wellbeing, Housing and Local Government.
- Nor Akmar, A. A. (2012). *Green space use and management in Malaysia*. Forest & Landscape, University of Copenhagen.
- Nutsford, D., Pearson, A. L., & Kingham, S. (2013). An ecological study investigating the association between access to urban green space and mental health. *Public Health*, *127*(11), 1005–1011.
- Park, B. J., Tsunetsugu, Y., Kasetani, T., Hirano, H., Kagawa, T., Sato, M., & Miyazaki, Y. (2007). Physiological effects of shinrin-yoku (taking in the atmosphere of the forest)—using salivary cortisol and cerebral activity as indicators—. *Journal of Physiological Anthropology*, 26(2), 123-128.
- Pietilä, M., Neuvonen, M., Borodulin, K., Korpela, K., Sievänen, T., & Tyrväinen, L. (2015). Relationships between exposure to urban green spaces, physical activity and self-rated health. *Journal of Outdoor Recreation and Tourism*, 10, 44-54.
- Ramachandran, A., & Snehalatha, C. (2010). Rising burden of obesity in Asia. *Journal of Obesity*, 2010.
- Schipperijn, J., Stigsdotter, U. K., Randrup, T. B., & Troelsen, J. (2010). Influences on the use of urban green space A case study in Odense, Denmark. *Urban Forestry and Urban Greening*, *9*(1), 25–32.
- Sreetheran, M., & Adnan, M. R. (2007). Green Network Development of Kuala Lumpur City: From the perspective of Kuala Lumpur Structure Plan. *FORUM*, 7, 38–41.
- Staats, H., Kieviet, A., & Hartig, T. (2003). Where to recover from attentional fatigue: An expectancy-value analysis of environmental preference. *Journal of Environmental Psychology*, 23(2), 147–157.
- Takano, T., Nakamura, K., & Watanabe, M. (2002). Urban residential environments and senior citizens' longevity in megacity areas: the importance of walkable green spaces. *Journal of Epidemiology and Community Health*, *56*(12), 913–918.
- Thompson Coon, J., Boddy, K., Stein, K., Whear, R., Barton, J., & Depledge, M. H. (2011). Does participating in physical activity in outdoor natural environments have a greater effect on physical and mental wellbeing than physical activity indoors? A systematic review. *Environmental science and technology*, *45*(5), 1761-1772.

- Tyrväinen, L., Ojala, A., Korpela, K., Lanki, T., Tsunetsugu, Y., & Kagawa, T. (2014). The influence of urban green environments on stress relief measures: a field experiment. *Journal of Environmental Psychology*, 38, 1–9.
- United Nations, Department of Economic and Social Affairs, Population Division (2014). World Urbanization Prospects: The 2014 Revision, Highlights (ST/ESA/SER.A/352).
- Ward Thompson, C., Roe, J., Aspinall, P., Mitchell, R., Clow, A., & Miller, D. (2012). More green space is linked to less stress in deprived communities: evidence from salivary cortisol patterns. *Landscape and Urban Planning*, 105(3), 221–229.
- Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures of positive and negative affect: the PANAS scales. *Journal of Personality and Social Psychology*, *54*(6), 1063–1070.
- Wilson, E. O. (1984). Biophilia, Harvard University Press. Cambridge, Massachusetts.