



**UNIVERSITI PUTRA MALAYSIA**

***PRODUCTION OF RHAMNOSE FROM POMELO PEEL USING  
LOCALLY-ISOLATED FUNGI *Trametes sp. IP3****

**NUR HAZIQAH ANIYAH BINTI SALIHAN**

**FBSB 2016 47**



**PRODUCTION OF RHAMNOSE FROM POMELO PEEL USING  
LOCALLY-ISOLATED FUNGI *Trametes* sp. IP3**

**By**

**NUR HAZIQAH ANIYAH BINTI SALIHAN**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfillment of the Requirements for Master of Science**

**February 2016**

## **COPYRIGHT**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artworks, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia.



## **DEDICATION**

This work is dedicated to my beloved family, who has always given me courage and support to carry out my studies. Thanks to their everlasting love and care.



Abstract of thesis presented to Senate of Universiti Putra Malaysia in fulfilment of the requirement for degree of Master of Science

**PRODUCTION OF RHAMNOSE FROM POMELO PEEL USING  
LOCALLY-ISOLATED FUNGI *Trametes* sp. IP3**

By

**NUR HAZIQAH ANIYAH BINTI SALIHAN**

**February 2016**

**Chairman : Helmi Wasoh @ Mohamad Isa, PhD**  
**Faculty : Biotechnology and Biomolecular Sciences**

Pomelo fruit (*limau bali*) is very popular due to its huge flesh compared to the typical citrus fruit, orange. Malaysia produce in average around 8 000 tons of pomelo each year. However, because pomelo has a very thick skin, it was not highly demand by the industry. Despite its thick skin (peel), the peel contains a rich source of flavonoid; naringin. Naringin, a bitter taste of flavonoid can be biodegraded into rhamnose and glucose with the presence of naringinase enzyme;  $\beta$ -glucosidase and  $\alpha$ -L-rhamnosidase. Rhamnose is a typical 6-deoxyhexose similar to fucose and exhibits a taste similar to D-mannose and usually found in plant as a constituent of glycosides such as naringin. Many researches use chemical hydrolysis to obtain rhamnose from naringin. Rhamnose can be used to produce food flavour such as 4-hydroxy-2,5-dimethyl-furanone (HDMF) that was used abundantly in food flavor production for example Nestle as caramel flavour. Due to the abundance of acid used for the production of the food flavor, the process is not suitable for environmental friendly food production. As an alternative, rhamnose was obtained through microbial hydrolysis using enzyme from fungi. Therefore, this study was done to produce rhamnose and glucose using microbial (isolated fungi) process using the pomelo waste product (peel) as the main substrate with the minimal usage of hazardous chemical. In this study, 84 strains were isolated and it was found that strain IP 3 was able to produce 18.15 g/L rhamnose from 100 g/L dry treated pomelo peel, and 8.96 g/L of rhamnose from untreated pomelo peel. *Trametes* sp. IP3 has the potential to produce high naringinase enzyme for the bioconversion of naringin into rhamnose. However, the production of glucose was not reported with the production of rhamnose. Thus, a hydrolysate containing high activity (53.4 U/mL) of  $\beta$ -glucosidase enzyme from fermentation of *Ganoderma lucidum* (compost OPEFB as the substrate) was added to the fermentation media at day 6. Additional sugars were detected at the end of experiment; xylose, mannose, and rhamnose however, glucose was not detected. Nevertheless, *Trametes* sp. IP3 possesses the ability to produce sugars besides rhamnose with the presence of high activity of  $\beta$ -glucosidase enzyme.

Keywords: Rhamnose, naringin, pomelo, naringinase,  $\alpha$ -L-rhamnosidase

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Sarjana Sains

**PENGHASILAN RHAMNOSE MENGGUNAKAN KULIT LIMAU BALI  
YANG DIREMBESKAN OLEH KULAT *Trametes* sp. IP3**

Oleh

**NUR HAZIQAH ANIYAH BINTI SALIHAN**

**Februari 2016**

**Pengerusi : Helmi Wasoh @ Mohamad Isa, PhD**  
**Fakulti : Bioteknologi dan Sains Biomolekul**

*Pomelo* (limau bali) sangat popular disebabkan oleh saiz isinya yang lebih besar berbanding buah-buahan sitrus biasa seperti buah oren. Malaysia menghasilkan kira-kira 8 000 tan limau bali setiap tahun. Walau bagaimanapun, limau bali tidak menjadi tarikan dan pilihan industri makanan untuk tujuan pengkomersilan disebabkan oleh kulitnya yang sangat tebal. Walaubagaimanapun, kulit limau bali mengandungi sumber yang kaya dengan flavonoid iaitu naringin. Naringin, flavonoid yang agak pahit boleh di uraikan kepada rhamnose dan glukosa dengan kehadiran enzim naringinase;  $\beta$ -glucosidase dan  $\alpha$ -L-rhamnosidase. Rhamnose adalah sejenis 6-deoxyhexose seperti fucose dan mempunyai rasa yang sama seperti D-mannose. Rhamnose biasanya ditemui dalam tumbuhan sebagai satu daripada komponen glikosida seperti naringin. Banyak kajian dijalankan menggunakan hidrolisis kimia untuk mendapatkan rhamnose daripada naringin. Rhamnose boleh digunakan untuk menghasilkan rasa makanan seperti 4-hydroxy-2,5-dimetil-furanone (HDMF) dan banyak digunakan industri makanan seperti Nestle sebagai rasa karamel. Oleh kerana banyak asid yang digunakan dalam proses pembuatan perasa makanan tersebut, proses itu tidak sesuai untuk pengeluaran makanan mesra alam sekitar. Sebagai alternatif, rhamnose diperolehi melalui isolasi mikrob daripada limau bali untuk menghasilkan enzim  $\beta$ -glucosidase dan  $\alpha$ -L-rhamnosidase. Oleh itu, kajian ini dilakukan untuk menghasilkan dan glukosa rhamnose dengan bantuan mikrob (kulat) dengan menggunakan limau bali (kulit) sebagai substrat utama dengan penggunaan bahan kimia secara minimum. Dalam kajian ini, daripada 84 kulat, kulat jenis IP 3 dapat menghasilkan 18.15 g/L rhamnose dari 100 g/L kulit limau bali yang di rawat, dan 8.96 g/L rhamnose dari kulit limau bali tidak dirawat. *Trametes* sp. IP3 dikenalpasti mempunyai potensi untuk menghasilkan enzim naringinase tinggi untuk menukarkan naringin kepada rhamnose. Walau bagaimanapun, pengeluaran glukosa tidak dilaporkan dengan pengeluaran rhamnose. Oleh itu, hidrolisat yang mengandungi aktiviti yang tinggi (53.4 U/mL) daripada  $\beta$ -glucosidase enzim dari penapaian *Ganoderma lucidum* (kompos OPEFB sebagai substrat) telah ditambah kepada media penapaian pada hari ke enam. Gula tambahan

telah dikesan pada akhir eksperimen; xylosa, mannososa dan rhamnososa bagaimanapun, glukosa tidak dapat dikesan. Walau bagaimanapun, *Trametes* sp. IP3 mempunyai keupayaan untuk menghasilkan gula selain rhamnose dengan kehadiran aktiviti yang tinggi  $\beta$ -glucosidase enzim.

Kata kunci: Rhamnose, naringin, limau bali, naringinase,  $\alpha$ -L-rhamnosidase



## ACKNOWLEDGEMENT

Assalamualaikum wt wb.,

Thank you Allah for giving me strength and patience to finish my Master study. I would like to acknowledge my supportive supervisor, Dr. Helmi Wasoh @ Mohamad Isa for the endless guidance and care throughout my master studies. Thank you for showing the right way of doing research especially polishing my writing skills.

I would also like to acknowledge my co-supervisor Dr. Mohd Rafein Zakaria and Prof. Dr. Suraini Abd-Aziz for their guidance. Million thanks to our caring lab assistance Mr. Rosli Aslim, Mrs. Aluyah Marzuki, Mrs. Renuga Panjamurji, Mrs. Norazlina Mohamad and Mrs. Noraishah Abd Rahman for their kind assistance and suggestion throughout my study.

I would also like to acknowledge Ministry of Higher Education for funding this project through Fundamental Research Grant Scheme (FRGS).

Thank you to my dearest friends; Nordiyana Nordin, Faiqah Abd Rahim and all my friends, whom are very supportive, understanding, and comment. A special thanks to my parents and my husband, Mohd Rafidyy bin Miswan for their endless support and encouragement.



I certify that a Thesis Examination Committee has met on 24 February 2016 to conduct the final examination of Nur Haziqah Aniyah binti Salihan on her thesis entitled "Production of Rhamnose from Pomelo Peel using Locally-Isolated Fungi *Trametes* sp. IP3" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

**Noor Azmi Shaharuddin, PhD**

Senior Lecturer  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Chairman)

**Phang Lai Yee, PhD**

Associate Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Wan Mohtar Wan Yusoff, PhD**

Professor  
National University of Malaysia  
Malaysia  
(External Examiner)



---

**NOR AINI AB. SHUKOR, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 3 November 2016

Thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment for the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Helmi Wasoh @ Mohamad Isa, PhD**

Senior Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

**Suraini Abd-Aziz, PhD**

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

**Mohd Rafein Zakaria, PhD**

Senior Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

---

**BUJANG KIM HUAT, PhD**

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

## Declaration by graduate student

I declare that the thesis is:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No. Nur Haziqah Aniyah binti Salihan, GS 30650

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of  
Chairman of  
Supervisory  
Committee: Dr. Helmi Wasoh @ Mohamad Isa

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: Professor Dr. Suraini Abd-Aziz

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: Dr. Mohd Rafein Zakaria

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	ii
<b>ACKNOWLEDGEMENT</b>	iv
<b>APPROVAL</b>	v
<b>DECLARATION</b>	vii
<b>LIST OF TABLES</b>	xii
<b>LIST OF FIGURES</b>	xiii
<b>LIST OF ABBREVIATIONS</b>	xv
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	
1.1 Pomelo and alternative source of rare sugar (rhamnose)	1
<b>2 LITERATURE REVIEW</b>	
2.1 Advance in the use of pomelo in production of rhamnose	3
2.2 Naringin as the main component in pomelo peel	5
2.2.1 Biosynthesis pathway of naringin	6
2.3 Enzymatic hydrolysis of naringin by naringinase enzyme	7
2.3.1 $\alpha$ -L-rhamnosidase	8
2.3.2 $\beta$ -glucosidase	9
2.4 Rhamnose as the main product of naringin hydrolysis	9
2.5 General application of rhamnose	11
2.6 The enzymatic assistance of oil palm empty fruit bunch (OPEFB) compost substrate in producing glucose	12
2.6.1 OPEFB Nutrient Content	13
2.7 Diversity of fungi	13
2.7.1 Basidiomycetes	16
<i>Ganoderma lucidum</i>	16
2.7.2 Agaricomycetes	18
<i>Trametes</i> sp.	19
<b>3 MATERIALS AND METHODS</b>	
3.1 Experimental design	20
3.2 Source of pomelo peel	22
3.3 Naringin extraction	
3.3.1 Preparation of dry methalonic extraction (DME)	22
3.3.2 Method 1: Dichloromethane treatment	22
3.3.3 Method 2: Without dichloromethane	22
3.3.4 Method 3: Hot water treatment	23
3.3.5 Method 4: Reflux treatment	23
3.3.6 Method 5: Ultrasonic extraction	23
3.3.7 Method 6: Reflux and Soxhlet extraction	23

3.3.8	Albedo physical observation preparation using Scanning Electron microscope (SEM)	24
3.4	Isolation and screening of rhamnose producing fungi	24
3.4.1	Observation using SEM	24
3.4.2	DNA extraction and identification of rhamnose producing fungi	25
3.5	Rhamnose identification medium	25
3.6	Cultivation of <i>Ganoderma lucidum</i> for $\beta$ -glucosidase production	
3.6.1	Organism and cultivation medium	25
3.6.2	Fermentation of <i>Ganoderma lucidum</i> for $\beta$ -glucosidase production	26
3.7	Growth and production medium for <i>Trametes</i> sp. IP3	
3.7.1	Inoculum preparation	27
3.8	Fermentation medium for sugar production	
3.8.1	Effect of different media	27
3.8.2	Effect of different carbon sources	28
3.8.3	Effect of carbon source concentration	28
3.9	Analytical procedures	
3.9.1	Naringin assay	29
3.9.2	Naringinase assay	29
3.9.3	High performance liquid chromatography (HPLC) analysis	30
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	
4.1	Pomelo peel	31
4.2	Naringin extraction	32
4.2.1	Pre-treatment methods for naringin extraction from albedo	32
4.2.2	Physical observation of albedo using SEM	35
4.3	Isolation and Screening of Rhamnose Producing Fungi	
4.3.1	Rhamnose-producing fungi	38
4.3.2	Observation using SEM	42
4.3.3	DNA extraction and identification of fungi	44
4.4	Production of rhamnose from naringin biodegradation	45
4.4.1	Production of rhamnose	45
4.5	Naringinase enzyme	54
4.5.1	Production of $\alpha$ -L-rhamnosidase	58
4.5.2	Production of high $\beta$ -glucosidase	60
4.6	Effects of enzyme hydrolysate for the production of multiple rare sugar	63
4.6.1	Effect of enzyme hydrolysate (from <i>Ganoderma lucidum</i> fermentation to the end fermentation product of <i>Trametes</i> sp. IP3	63
<b>5</b>	<b>CONCLUSION AND RECOMMENDATION</b>	<b>65</b>

<b>REFERENCES</b>	66
<b>APPENDICES</b>	77
<b>BIODATA OF STUDENT</b>	91
<b>LIST OF PUBLICATIONS</b>	92



## LIST OF TABLES

Table		Page
2.1	Major components of OPEFB	13
2.2	Nutrient content of empty fruit bunch (EFB)	13
2.3	Biologically active compound in <i>Ganoderma lucidum</i>	18
3.1	Series of media used for selection of rhamnose producing media	28
4.1	Naringin extracted from dried albedo (10.0 g) using different extraction methods.	32
4.2	List of fungi that grow in naringin selective media and the production of rhamnose by isolated fungi using technical grade naringin.	41
4.3	A summary of rhamnose production (g/L) using different production media.	47
4.4	Comparison of rhamnose production based on different carbon source by isolate IP3 and PP5.	49
4.5	Comparison of rhamnose production from the previous studies using different microorganism and substrate with the current study.	53
4.6	Production of $\alpha$ -L-rhamnosidase and $\beta$ -glucosidase from different substrates and microorganisms	58



## LIST OF FIGURES

Figure		Page
2.1	Schematic diagram of pomelo fruit	4
2.2	Structure of naringin	5
2.3	Hydrolysis of naringin into prunin, rhamnose, naringenin and glucose by naringinase expressing $\alpha$ -l-rhamnosidase and $\beta$ -d-glucosidase activities	7
2.4	Rhamnose chemical structure	10
2.5	Sexual reproduction of <i>Basidiomycete</i> .	16
3.1	Overall experimental design	21
4.1	Component in pomelo fruit	31
4.2	Sample of albedo under SEM for (a) raw; (b) treated with ethanol; (c) treated with ethanol followed by Soxhlet extraction	37
4.3	Isolated fungi from pomelo after 7 days incubation at 27°C by using media added with naringin	40
4.4	Isolate IP 3 and PP 5 produced the highest rhamnose from naringin as substrate	41
4.5	Isolate PP5 and isolate IP3 viewed under SEM	43
4.6	Phylogenetic tree of isolate IP3	44
4.7	Production of rhamnose using potato dextrose broth media added with technical grade naringin	46
4.8	Production of rhamnose (g/L) using different concentration of dried albedo as carbon source by strain IP3	48
4.9	Production of rhamnose (g/L) of fungi IP3 using different concentration of dried albedo as carbon source	50
4.10	Naringin and rhamnose concentration within 10 d fermentation period for strain PP5	52
4.11	Hydrolysis of naringin to prunin by $\alpha$ -L-rhamnosidase	54
4.12	Enzyme activity of $\alpha$ -L-rhamnosidase within 9 days period	55
4.13	<i>Trametes</i> sp. strain IP3 fermentation results of enzymatic activity and production of rhamnose using extracted naringin as sole carbon source	56
4.14	Hydrolysis of naringin to prunin by $\alpha$ -L-rhamnosidase. (b) Hydrolysis of prunin to naringenin by $\beta$ -glucosidase.	57
4.15	Effects of different carbon sources on the production of $\beta$ -glucosidase activity.	60

4.16	Effect of composted OPEFB concentration, g/L on the production of $\beta$ -glucosidase by <i>Ganoderma lucidum</i>	61
4.17	Effect of temperatures on the synthesis of $\beta$ -glucosidase by <i>Ganoderma lucidum</i>	62
4.18	Sugar production, g/L of xylose (■), mannose (▲), and rhamnose (◆) from <i>Trametes</i> sp. IP3 with the mixture of <i>Ganoderma lucidum</i> hydrolysate	64



## LIST OF ABBREVIATIONS

g	Gram
L	Litre
Min	Minutes
mL	Milliliter
nm	Nanometer
°C	Degree Celsius
Rpm	Rotation per minute
OPEFB	Oil palm empty fruit bunch
g/L	Gram per liter
HPLC	High performance liquid chromatography
B	Beta
$\alpha$	Alpha
U/mL	Unit per milliliter
SEM	Scanning electron microscope
dTDP	deoxythymidine diphosphate
DME	Dry methalonic extraction
UDP	Uridine diphosphate glucose
$\mu\text{m}$	Micrometer
EC	Enzyme commission number
PD	Potato dextrose
DNA	Deoxyribonucleic acid

## CHAPTER 1

### INTRODUCTION

#### 1.1 Pomelo and Alternative Source of Rare Sugar (Rhamnose)

Pomelo (*Citrus maxima*) is a native citrus fruit to southern East Asia. In Malaysia, this Rutaceae family plant is widely grown in Perak, Kedah, Melaka and Johor. Pomelo peel consists of two major parts; flavedo and albedo. Flavedo, a green layer located at the outer side of fruit and albedo is a white layer with a spongy peel texture located between juice sacs and a flavedo (Panmanas and Charoonpong, 2012).

Every year massive peel is disposed as waste due to high consumption of pomelo flesh and causes a severe environmental issues (Hameed *et al.*, 2008). However, Dong-Mei *et al.*, (2011) stated that naringin, the main component in pomelo peel albedo can be utilized to produce rhamnose (rare sugar) and glucose by using naringinase enzyme ( $\alpha$ -l-rhamnosidase and  $\beta$ -d-glucosidase). Therefore, this has opened an opportunity for the agriculture waste industry to maximize the application of waste and use it to produce other useful products. L-rhamnose (6-deoxy-mannose) is a rare sugar that had been used as a starting material in synthesis of organic compound. It is considered as a high potential substrate in producing flavor. A highly quality flavor, furaneol (trademark of Firmenich SA, Geneva) was using rhamnose as its precursor. It acts as a chiral building block in chemical synthesis, the deoxysaccharide that can only be obtained with great effort by chemical means. It is produced industrially from a variety of vegetable raw materials, for example from rutin or from hesperidin and naringin (Lang and Wullbrandt, 1999). In chemical process, production of rhamnose involves large quantities of potentially toxic and dangerous waste products. The process of waste management cost the industry sometimes more than the production cost since waste need to be treated before it can be properly disposed.

Rhamnose had been produced by several bacteria from rhamnolipid. However, rhamnose that produced by pathogenic bacteria is not favourable in food industries (Giraud and Naismith, 2000) since it might cause harm to the consumers. Therefore, using the abundance of inexpensive substrate (albedo from pomelo peel) as the main carbon source is potentially used to produce high amount of rhamnose. In this study, the effects of different extraction process and the production of rhamnose using pomelo peel were also studied.

Naringinase however is one of the expensive enzyme and by culturing excellent strains, high yield naringinase can be produced and decrease production cost. Some researchers had suggested strain *Aspergillus niger* (Bram and Solomons, 1965), however Tselenis-Kotsowilis *et al.* (1982) reported that there have been some

defects which restricted the application of the naringinase involving immobilized cost, poor stability, and equipment used. Therefore, new strains which are safe, easy to maintain and convenient for fermentation process have been isolated.

As stated before, naringin can be further hydrolyzed to glucose by hydrolysis of  $\beta$ -d-glucosidase (Vila Real *et al.*, 2007). However, Hakasson *et al.* (2011) stated that the presence of  $\alpha$ -L-rhamnosidase will only inhibit the activity of  $\beta$ -glucosidase enzyme. Therefore, in this study, high  $\beta$ -glucosidase enzyme was extracted from other fermentation and was added into the rhamnose producing media to study whether the enzyme might help in producing glucose from naringin from hydrolysate.

It is generally known that cellulose can be converted into glucose by cellulase such as  $\beta$ -glucosidase. Cellulase has a great potential in industrial sector which can be used in textile, paper and poultry industry. The enzymatic breakdown of substrates by  $\beta$ -glucosidase brings the most promising technology for the conversion of lignocellulosic waste biomass (Baharuddin *et al.*, 2009a). In this work, further degradation of pomelo peel was carried out using cellulase from *Ganoderma lucidum*, a well-known fungi having the ability to grow well on oil plum tree. The hydrolysate of the final product was monitored using HPLC in order to check the quality of rhamnose. It is hope that this work will provide useful knowledge on the usage of pomelo peel as potential substrate for rhamnose production in the future.

The objectives of this study are:

- a) To isolate and identify rhamnose producing fungi from rotten pomelo peel.
- b) To produce rhamnose using albedo from pomelo peel through liquid state fermentation.

## REFERENCES

- Aneet Kaur, Satbir Singh, Ram S Singh, W H Schwarz and Munish Puri. (2010). Hydrolysis of citrus peel naringin by recombinant  $\alpha$ -L-rhamnosidase from *Clostridium stercorarium*. *Journal of Chemical Technology and Biotechnology* 85: 1419–1422.
- Aphylophorales in the Atlantic Rain Forest in Northeast Brazil. (2004). *Fungal Diversity*. 25: 49-67.
- Ariffin, D., Idris, A. S., and Singh, G. (2000). Status of Ganoderma in oil palm in Diseases of Perennial Crops. *CAB International*. 1: 49-68.
- Babitskaya, V.G., Shcherba, V.V., Puchkova. T.A., and Smirnov, D.A. (2005) Polysaccharides of *Ganoderma Lucidum*: Factors Affecting Their Production. *Applied Biochemistry and Microbiology*. 41(2): 169-173.
- Baharuddin, A. S., Kazunori, N., Abd-Aziz, S., Tabatabaei, M., Abdul Rahman, N. A., Hassan, M. A., Wakisaka, M., Sakai, K., and Shirai, Y. (2009a). Characteristics and Microbial Succession in Co-Composting of Oil Palm Empty Fruit Bunch and Partially Treated Palm Oil Mill Effluent. *The Open Biotechnology Journal*. 3:87-95.
- Baharuddin, A. S., Wakisaka, M., Shirai, Y., Abd-Aziz, S., Rahman, N. A., and Hassan, M. A. (2009b). Co-Composting of Empty Fruit Bunches and Partially Treated Palm Oil Mill Effluents in Pilot Scale. *International Journal Agriculture Resources*. 4(2): 69-78.
- Baldrian Petr, Kolařík Miroslav, Štursová Martina, Kopecký Jan, Valášková Vendula, Větrovský Tomáš, Žifčáková Lucia, Šnajdr Jaroslav, Rídl Jakub, Vlček Čestmír and Voříšková Jana. (2011). Active and total microbial communities in forest soil are largely different and highly stratified during decomposition. *The ISME Journal*. 6(2): 248–258.
- Busto, M. D., Meza, V., and Mateos, N. P. (2007) Immobilization of naringinase from *Aspergillus niger* CECT 2088 in poly (vinyl alcohol) cryogels for the debittering of juices. *Food Chemistry*. 104: 1177–1182.
- Campbell, L. A., and Palmer, G. H. (1978). Pectin, in: G.A. Spiller, R.J. Amen (Eds.), Topics in Dietary Fiber Research. *Plenum Press, New York*. 105–115.
- Cao, Y., and Tan, H. (2002) Effects of cellulase on the modification of cellulose. *Carbohydrate Resources*. 337:1291–1296.
- Capaldo, A., Walker, M. E., Ford, C. M., and Jiranek, V. (2011).  $\beta$ -Glucoside metabolism in *Oenococcus oeni*: cloning and characterisation of the phospho-b-glucosidase bglD. *Food Chemistry*. 125:476–482.

- Capaldo, A., Walker, M. E., Ford, C. M., and Jiranek, V. (2011).  $\beta$ -Glucoside metabolism in *Oenococcus oeni*: cloning and characterization of the phosphor- $\beta$ -glucosidase CelD. *Journal Molecular Catalysis*. 69: 27–34.
- Cartwright, K., St., G., and Findlay, P. K. (1951). Decay of Timber and Its Prevention. Scott S. Pauley. *The Quarterly Review of Biology*. 269(3): 296-297.
- Chang-Su P., Soo-Jin Y., Yu-Ri L., Yeong-Su K., and Deok-Kun O. (2010). Characterization of a recombinant thermostable L-rhamnose isomerase from *Thermotoga maritima* ATCC 43589 and its application in the production of L-lyxose and L-mannose. *Biotechnology Letters*. 32(22): 1947-1953.
- Cheong, M. W., Loke, X. Q., Liu, S. Q., Pramudya, K., Curran, P., and Yu, B. (2011). Characterization of volatile compounds and aroma profiles of Malaysian pomelo (*Citrus grandis* (L.) Osbeck) blossom and peel. *Journal Essential Oil Resources*. 23(2): 34-44.
- Chevenet, F., Brun, C., Banuls, A. L., Jacq, B., and Chisten, R. (2006). TreeDyn: towards dynamic graphics and annotations for analyses of trees. *BMC Bioinformatics*. 10(7): 439.
- Chien, P. J., Sheu, F., and Shiyu, Y. T. (2001). Monitoring enzymatic debittering in grapefruit juice by high performance liquid chromatography. *Journal Food and Drug Analysis*. 9: 115-120.
- Cock, Ian Edwin. (2013). Antimicrobial activity of *Backhousia citriodora* (lemon myrtle) methanolic extracts. *Pharmacognosy Communications*. 3(2): 58-63.
- Cui B. K., and Dai Y. C. (2011). "*Fomitiporia ellipsoidea* has the largest fruiting body among the fungi". *Fungal Biology*. 115(9): 813–814.
- Daniel Sliva. (2003). *Ganoderma lucidum* (Reishi) Cancer Treatment Integrative Cancer Therapies 2(4): 358-364.
- Davis, D. W. (1947). Determination of flavonones in citrus juice. *Analysis Chemistry*. 19: 46–48.
- de Lima, C. J. B., Ribeiro, E. J., Sérvulo, E. F. C., Resende, M. M., and Cardoso, V. L. (2009) Biosurfactant Production by *Pseudomonas aeruginosa* Grown in Residual Soybean Oil. *Applied Biochemistry Biotechnology*. 152: 156–168.
- Department of Agriculture Malaysia. (2009). Pomelo: Varieties / clones, [www.doa.gov.my/main.php/Content=sections&SubSectionID=390&SectionID=5&CurLocation=315&IID=](http://www.doa.gov.my/main.php/Content=sections&SubSectionID=390&SectionID=5&CurLocation=315&IID=).
- Deshmukh, V. V., Raut, B. T., Mane, S. S., Ingle, R.W., and Joshi, M. S., (2014). Compatibility of *Bradyrhizobium japonicum* isolates with agrochemicals.

*American International Journal of Research in Formal, Applied & naturam Sciences* 14-228.

Feng, B., Ma, B., Kang, L., Xiong, C., and Wang, S. (2005). The microbiological transformation of steroidal saponins by *Curvularia lunata*. *Tetrahedron* 61: 11758–11763.

Ferraria, L., Afonso, C., VilaReal, H., Alfaia, A., and Ribeiro, M. H. L. (2008). Evaluation of effect of high pressure on naringin hydrolysis in grapefruit juice with naringinase immobilized in calcium alginate beads. *Food Technology Biotechnology*. 46: 146-150.

Francesco Trotta, Enrico Driolib, Claudio Baggianic, and Domenico Lacopo. (2002). Molecular imprinted polymeric membrane for naringin recognition, *Journal of Membrane Science*. 201: 77–84.

Gao, Z., Hop, D. V., Yen, L. T. H., Ando, K., Hiyamuta, S., and Kondo, R. (2012). The production of beta-glucosidases by *Fusarium proliferatum* NBRC109045 isolated from Vietnamese forest.

Garcia-Garrido, J. M., Tribak, M., Reion-Palomares, A., Ocampo, J. A., and Garcia-Romera, I. (2000). Hydrolytic Enzymes and Ability of Arbuscular Mycorrhizal Fungi to Colonize Roots. *Journal of Experimental Botany*. 51(349): 1443-1448.

Gavin Towler and Sinnott, R. K. (2007). *Chemical Engineering Design: Principles, Practice and Economics of Plant and Process Design*. Butterworth-Heinemann.

Giannuzzo N. Amelia, He'ctor J. Boggetti, Mo'nica A. Nazareno and Horacio T. Mishima. (2003) Supercritical Fluid Extraction of Naringin from the Peel of *Citrus paradisi* *Phytochem. Analysis*. 14, 221–223.

Giannuzzo, A. N., Nazareno, M. A., Mishima, H. T., and Lopez de Mishima, B. A. (2010). Extraction of naringin from citrus paradise L. studies comparison and optimization of optimization technique. *Journal Science Technology*. 20: 257-251.

Gonzalez-Barrio, R., Trinidale, L. M., Manzanares, P., deGraaff, L. H., Tomas-Barberan, F. A., and Espin, J. C. (2004). *Journal of Agriculture and Food Chemistry*. 52: 6136–6142.

Habijani, J., Berovi, M., Wraber, B., Hodzar, D., and Boh, B. (2001). Immunostimulatory Effects of Fungal Polysaccharides from *Ganoderma lucidum* Submerged Biomass Cultivation. *Food Technology Biotechnology*. 39: 327-331.

Hakansson, A. E., Ameijde, J., Horne, G., Nash, R. J., Wormald, M. R., and Kato, A. (2008). Synthesis of the naringinase inhibitors l-swainsonine and related 6-Cmethyl-l-swainsonine analogues: (6R)-C-methyl-l-swainsonine is a more



potent inhibitor of l-rhamnosidase by an order of magnitude than lswainsonine. *Tetrahedron Letter*. 49: 179–84.

Hashimoto, W., Miyake, O., Nankai, H., and Murat, K. (2003). Molecular identification of an [alpha]-rhamnosidase from *Bacillus* sp. strain GL1 as an enzyme involved in complete metabolism of gellan. *Archieve Biochemistry Biophysics*. 415: 235–244.

Hashimoto, W., Nankai, H., Sato, N., Kawai, S. and Murata, K. (2000). Characterization of  $\alpha$ -L-rhamnosidase of *Bacillus* sp. GL1 responsible for the complete depolymerization of gellan. *Archives of Biochemistry and Biophysics*. 368: 56–60.

Hawksworth, D. L., Kirk, P. M., Sutton, B. C., and Pegler, D. N. (1995). Ainsworth and Bisby's Dictionary of the Fungi. *CAB International, Wallingford, UK*.

Heightman, T. D., Vasella, A., and Angew, T. (1999) Chemistry. *International Education*. 38: 750–770.

Helder Vila-Reala, António Alfaia, J., Robert Phillips, S., António Caladoa, R., and Maria Ribeiroa, H.L. (2010). Pressure-enhanced activity and stability of  $\alpha$ -l-rhamnosidase and  $\beta$ -d-glucosidase activities expressed by naringinase. *Journal of Molecular Catalysis B: Enzymatic* 65: 102–109.

Hibbett, D. S., and Thorn, R. G. (2001). Systematics and Evolution. *Springer-Verlag*. 7: 121–168.

Hibbett, D.S. (2007). "A higher level phylogenetic classification of the Fungi". *Mycological Research*. 111(5): 509–547.

Hollman, P. C. H. (2004). Absorption, bioavailability, and metabolism of flavonoids. *Pharmaceutical Biology*. 42: 74–83.

Hsu, T.A. (1996). Pretreatment of biomass, in *Handbook on Bioethanol: Production and Utilization*, ed by Wyman CE. *Taylor & Francis, Washington*.

Ibrahim M. F., Abd-Aziz S., Abdul Razak M. N., Phang L. Y. and Hassan M.A. (2012). Oil palm empty fruit bunch as alternative substrate for acetone–butanol–ethanol production by *Clostridium butyricum* EB6. *Applied Biochemistry and Biotechnology*. 166(7): 1615–1625

Jabalpurwala, F. A., Smoot, J. M., and Rouseff, R. L. (2009) A comparison of citrus blossom volatiles. *Phytochemistry*. 70: 1428–1434.

James, R., Ketudat Cairns, and Asim Esen. (2010).  $\beta$ -Glucosidases. *Cell Molecular Life Sciences*. 67: 3389–3405.

Kamal Bahrin, E., Seng, P. Y., and Abd-Aziz, S. (2011). Effect of oil Palm Empty Fruit Bunch Particle Size on Cellulase Production by *Botryosphaeria* sp. Under Solid State Fermentation. *Australian Journal of Basic and Applied Sciences*. 5(3): 276–280.

- Kanokpan Thammawat, Pongsuda Pongtanya, Vitchuporn Juntharasri and Patjaraporn Wongvithoonyaporn. (2008). Isolation, Preliminary Enzyme Characterization and Optimization of Culture Parameters for Production of Naringinase Isolated from *Aspergillus niger* BCC 25166 Kasetsart. *Journal of Natural Sciences*. 42: 61 – 72.
- Kara Rogers. (2011). Fungi, Algae and Protists. *Biochemistry, Cells and Life, 1<sup>st</sup> ed*, Britannica Educational Publishing. ISBN 978-1-61530-463-9 (eBook)
- Kirk, P. M., Cannon, P. F., Minter, D. W., and Stalpers, J. A. (2008). Dictionary of the Fungi. *Wallingford, UK: CAB International*. 10: 695.
- Tselenis-Kotsowilis A.D., Koliomichalis M.P., Papavassiliou J.T. (1982) Acute pyelonephritis caused by *Staphylococcus xylosus*. *Journal Clinic Microbiology* 16:593–594
- Krishna, C. (1999). Production of Bacterial Cellulase by Solid State Bioprocessing of Banana Waste. *Bioresource Technology*. 69: 231-239.
- Kumar, Atul, and Ram Awatar Maurya. (2008). Efficient Synthesis of Hantzsch Esters and Polyhydroquinoline Derivatives in Aqueous Micelles. *Synlett*. 1: 883–885.
- Lei, Z., Wang, H., Zhou, and R., Duan, Z. (2002). "Influence of salt added to solvent on extractive distillation". *Chemistry Engineering Journal*. 87(2): 149.
- Lide, D. R. (2000). CRC Handbook of Chemistry and Physics 81st edition. *CRC press*.
- Lindequist, U., Niedermeyer, T. H. J., and Jülich, W. D. (2005). "The pharmacological potential of mushrooms". *Evidence-based Complementary and Alternative Medicine*. 2(3): 285–299.
- Ma, S. J., Leng, B., Xu, X. Q., Zhu, X. Z., Shi, Y., Tao, Y. M., Chen, S. X., Long, M. N., and Chen, Q. X. (2011). Purification and Characterization of  $\beta$ -1,4-Glucosidase from *Aspergillus Glaucus*. *African Journal of Biotechnology*. 10(84): 19607-19614.
- Magan, N. (2008). Ecophysiology: Impact of Environment on Growth, Synthesis of Compatible Solutes and Enzyme Production in *Ecology of Saprotrophic Basidiomycetes: British Mycological Society Symposia Series*. Elsevier Ltd. 1: 64-77.
- Mannfred Rossberg, Wilhelm Lendle, Gerhard Pfeleiderer, Adolf Tögel, Theodore, R., Torkelson, Eckhard Löser, Klaus, K., and Beutel. (2011). "Chlorinated Hydrocarbons". *Ullmann's Encyclopedia of Industrial Chemistry*, Wiley-VCH, Weinheim.

- Manzanares, P., Broeck, H.C.V., Graaff, L.H.D. and Visser, J. (2001). Purification and characterization of two different  $\alpha$ -L-rhamnosidases, RhaA and RhaB from *Aspergillus aculeatus*. *Applied and Environmental Microbiology*. 67: 2230–2234.
- Manzanares, P., Orejas, M., Ibanez, E., Valles, S. and Ramon, D. (2000). Purification and characterization of an alpha-L-rhamnosidase from *Aspergillus nidulans*. *Letters in Applied Microbiology*. 31: 198–202.
- Manzanares, P., Orejas, M., Ibanez, E., Valles, S., and Ramon, D. (2003). Construction of a genetically modified wine yeast strain expressing the *Aspergillus aculeatus rhaA* gene, encoding an  $\alpha$ -L-rhamnosidase of enological interest. *Applied Environmental Microbiology*. 69: 7558–7562.
- Manzanares, P., Orejas, M., Ibanez, E., Valles, S., and Ramon, D. (2000). Purification and characterization of an alpha-l-rhamnosidase from *Aspergillus nidulans*. *Letter Applied Microbiology*. 31: 198–202.
- Matthews, A. (2004). Lactic acid bacteria as a potential source of enzymes for use in vinification. *Applied Environmental Microbiology*. 70: 5715–5731.
- Md Shah, U. K., Ariff, A. B., Zulkifli, H. S., Tong, C. C., Hassan, M. A., and Karim, M. I. A. (1997). The Treatment of oil Palm Empty Fruit Bunch Fibre for Subsequent Use as Substrate for Cellulase Production by *Chaetomium Globosum* Kunze. *Bioresource Technology*. 62: 1-9.
- Ming-Hua Yang, Yi-Ju Lin, Chang-Hsin Kuo and Kuo-Lung Ku. (2010). Medicinal Mushroom *Ganoderma lucidum* as a Potent Elicitor in Production of *t*-Resveratrol and *t*-Piceatannol in Peanut Calluses. *Journal Agriculture Food Chemistry*. 58(17): 9518–9522
- Minhee Han, Yule Kim, Seung Wook Kim and Gi-Wook Choi. (2011). High efficiency bioethanol production from OPEFB using pilot pretreatment reactor. *Journal Chemical Technology Biotechnology*. 86: 1527–1534.
- Miura, T., Yuan, L., Sun, B., Fujui, H., Yoshida, M., Wakame, K., and K, K. (2002). Isoflavone Aglycon Produced by Culture of Soybean Extracts with Basidiomycetes and its Anti-angiogenic Activity. *Bioscience, Biotechnology and Biochemistry*. 66(12): 2626-2631.
- Miyata, T., Kashige, N., Satho, T., Yamaguci, T., Aso, Y., and Miake, F. (2005). *Current Microbiology*. 51:105–109
- Moiser, N., Wyman, C., Dale, B., Elander, R., Lee, Y. Y., and Holtzapple, M. (2005). Features of promising technologies for pretreatment of lignocellulosic biomass. *Bioresource Technology*. 96: 673–686.
- Monti, D., Pisvejcova, A., Kren, V., Lama, M. and Riva, S. (2004). Generation of  $\alpha$ -L- rhamnosidase library and its application for the selective

derhamnosylation of natural products. *Biotechnology and Bioengineering*. 87: 763–771.

- Munish Puri, Aneet Kaur, Wolfgang, H., Schwarz, Satbir Singh, and Kennedy, J. F. (2011). Molecular characterization and enzymatic hydrolysis of naringin extracted from kinnow peel waste. *International Journal of Biological Macromolecules*. 48(1): 58–62.
- Mun-Wai Cheong , Xiu-Qing Loke , Shao-Quan Liu , Kiki Pramudya , Philip Curran and Bin Yu. (2011). Characterization of Volatile Compounds and Aroma Profiles of Malaysian Pomelo (*Citrus grandis* (L.) Osbeck) Blossom and Peel. *Journal of Essential Oil Research*. 23(2): 34-44.
- Mun-Wai Cheong, Xiu-Qing Loke, Shao-Quan Liu, Kiki Pramudya, Philip Curran and Bin Yu. (2011). Characterization of Volatile Compounds and Aroma Profiles of Malaysian Pomelo(*Citrus grandis* (L.) Osbeck) Blossom and Peel, *Journal of Essential Oil Research*. 23(2): 34-44.
- Murray, R. K., Granner, D. K., Mayes, P. A., and Rodwell, V. W., Harper. (1994). Bioquímica. *Atheneu: São Paulo*. 1: 628–646.
- Natalia Dudareva, Antje Klempien, Joelle K. Muhlemann, and Ian Kaplan. (2013). Biosynthesis, function and metabolic engineering of plant volatile organic compounds. *New Phytologist*. 198(1): 16–32.
- Njoroge, S. M., Koaze, H., Karanja, P. N., and Sawamura M. (2005). Volatile constituents of Redblush grapefruit (*Citrus paradisi*) and pummel (*Citrus grandis*) peel essential oils from Kenya. *Journal Agriculture Food Chemistry*. 53: 9790-9794.
- Norouzian, D., Hosseinzadeh, A., Inanlou, D. N., and Moazami, N. (1999). Various techniques used to immobilize naringinase produced by *Penicillium decumbens* PTCC 5248. *World Journal Microbiology Biotechnology*. 15: 501-502.
- Norouzian, D., Hosseinzadeh, A., Inanlou, D.N., and Moazami, N. (2000). Production and partial purification of naringinase by *Penicillium decumbens* PTCC 5248. *World Journal Microbiology Biotechnology*. 16: 471-473.
- Office of Environmental Health Hazard Assessment. (2000). "Dichloromethane". Public Health Goals for Chemicals in Drinking Water. *California Environmental Protection Agency*.
- Orejas, M., Ibanez, E., and Ramon, D. (1999). The filamentous fungus *Aspergillus nidulans* produces an  $\alpha$ -L-rhamnosidase of potential oenological interest. *Letters in Applied Microbiology*. 28: 383-388.
- Panmanas Sirisomboon and Charoonpong Theamprateep. (2012). Physicochemical and Textural Properties of Pomelo (*Citrus maxima* Merr. cv. Kao Nam

Pueng) Fruit at Preharvest, Postharvest and During the Commercial Harvest Period. *Philipp Agricultural Scientist*. 95(1): 43-52.

- Park, S., Kim, J. and Kim, D. (2005). Purification and characterization of quercitrin-hydrolyzing alpha-L-rhamnosidase from *Fusobacterium* K-60, a human intestinal bacterium. *Journal of Microbiology and Biotechnology*. 15: 519–524.
- Paterson, R. R. (2006). "Ganoderma therapeutic fungal biofactory". *Phytochemistry*. 67(18):1985-2001.
- Paudyal, K.P. and Haq, N. (2008). Variation of pomelo (*Citrus grandis* (L.) Osbeck) in Nepal and participatory selection of strains for further improvement. *Agroforest System*. 72: 195-204.
- Penelope M. Drake<sup>1</sup>, Wonryeon Cho, Bensheng Li, Akraporn Prakobphol, Eric Johansen, N. Leigh Anderson, Fred E. Regnier, Bradford W. Gibson and Susan J. Fisher. (2009). Sweetening the Pot: Adding Glycosylation to the Biomarker Discovery Equation. *Clinical Chemistry*. 56(2): 223-236.
- Prillinger, H., Lopandic, K., Schweigkofler, W., Deak, R., Aarts, H. J. M., Bauer, R., Sterflinger, K., Kraus, G. F., and Maraz, A. (2002). Phylogeny and Systematics of the Fungi with Special Reference to the Ascomycota and Basidiomycota Breitenbach M, Cramer R, Lehrer SB (eds): Fungal Allergy and Pathogenicity. *Chemical Immunology*. 81: 207-295.
- Puri, M., and Banerjee, U. C. (2000). Production purification and characterization of debittering enzyme naringinase. *Advanced Biotechnology*. 18: 207-217.
- Puri, M., and Kalra, S. (2005) Purification and characterization of naringinase from a newly isolated strain of *Aspergillus niger* 1344 for the transformation of flavonoids. *World Journal Microbiology Biotechnology*. 21: 753–758.
- Puri, M., Banerjee, A., and Banerjee, U.C. (2005). Optimization of process parameters for the production of naringinase by *Aspergillus niger* MTCC 1344. *Process Biochemistry*. 40: 195-201.
- Puri, M., Marhawa, S.S., and Kothari, R.M. (1996). Studies on the applicability of alginateentrapped naringinase for the debittering of kinnow juice. *Enzyme Microbiology Technology*. 18: 281-285
- Qian, S., Yu, H., Zhang, C., Lu, M., Wang, H., and Jin, F. (2005). Purification and characterization of dioscin- $\alpha$ -l-rhamnosidase from pig liver. *Chemical Pharmaceutical Bulletin*. 53(8): 911–914.
- Rahman, S. H. A., Choudhury, J. P., Ahmad, A. L., and Kamruddin, A. H. (2007). Optimization studies on acid hydrolysis of oil palm empty fruit bunch fiber for production of xylose. *Bioresource Technology*. 98: 554–559.

- Rahman, S. H. A., Choudhury, J. P., Ahmad, A. L., and Kamaruddin, A. H. (2006). Optimization studies on acid hydrolysis of oil palm empty fruit bunch fiber for production of *xylose*. *Bioresource Technology*. 8(3): 554-559.
- Rao, Y. K., Lub, S. C., Liub, B., and Tzeng, Y. M. (2006). Enhanced production of an extracellular protease from *Beauveria bassiana* by optimization of cultivation processes. *Biochemical Engineering Journal*. 28: 57–66.
- Rayner, A. J. R., Beeching, J. D., Crower, J. C., and Watkins, Z. R. (1999). Defining Individual Fungal Boundaries in *Structure and Dynamics of Fungal Populations* ed by Worrall JJ, Luwer Academic, Norwell, MA. 1: 19-24.
- Renee Miller, Simon J. Owens, and Bjørn Rørslett. (2011). Plants and colour: Flowers and pollination. *Optics & Laser Technology*. 43(2): 282–294.
- Rosa Margesin, Georges Feller, Charles Gerday, and Nick J. Russell. (2003). Cold-Adapted Microorganisms: Adaptation Strategies and Biotechnological Potential. *Encyclopedia of Environmental Microbiology*.
- Rouseff, R. and Perez-Cacho, P. R. (2007). Citrus flavour. In: Flavours and Fragrances: Chemistry, Bioprocessing and Sustainability. *Springer*. 1: 117-134.
- Schell, D. J., Farmer, J. J., Newman, M., and Mcmillan, J. D. (2003). Dilute-sulfuric acid pretreatment of corn stover in pilot-scale reactor: investigation of yields, kinetics, and enzymatic digestibilities of solids. *Applied Biochemistry Biotechnology*. 105: 69–85.
- Schols, H. A., and Voragen, A. G. J. (1994). Occurrence of pectic hairy regions in various plant cell wall materials and their degradability by rhamnogalacturonase. *Carbohydrate Resources*. 256: 83-95.
- Selosse, M.A., Bauer. R., and Moyersoer, B. (2002). Basal hymenomycetes belonging to Sebacinaceae are ectomycorrhizal on temperate deciduous trees. *New Phytology*. 155: 183-195.
- Serpil Yalim, Yuksel Ozdemir and Ibrahim Ekiz H. (2004). Naringin in turkish orange juices and its reduction by naringinase. *Journal of Food and Drug Analysis*. 3(2): 273 – 276
- Shahriarinour, M., Abd Wahab, M. N., Mustafa, S., Mohamed, R., and Ariff, A. (2011). Effect of Various Pretreatments of Oil Palm Empty Fruit Bunch Fibres for Subsequent Use as Substrate on The Performance of Cellulase Production by *Aspergillus terreus*. *Bioresource*. 6(1): 291-307.
- Shaw, P. E., Goodner, K. L., Moshonas, M. G., and Hearn, C. J. (2001). Comparison of grapefruit hybrid fruit with parent fruit based on composition of volatile components. *Sci. Hortic*. 91: 71-80.

- Shimkets, L. J. (2013). Prokaryotic Life Cycles. *The Prokaryotes 2013*. 1: 317-336. DOI:10.1007/978-3-642-30123-0\_54.
- Silverstein, R. A., Chen, Y., Sharma-Shivappa, R. R., Boyette, M. D., and Jason Osborne. (2007). A comparison of chemical pretreatment methods for improving saccharification of cotton stalks. *Bioresource Technology*. 98: 3000–3011.
- Smith, M., Bruhn, J. H., and Anderson, J. B. (1992). "The fungus *Armillaria bulbosa* is among the largest and oldest living organisms". *Nature*. 35(6368):428-431.
- Smith, S.E., and Read, D.J. (2008). Mycorrhizal Symbiosis. 3: 11-13.
- Somogyi, M. (2000). Notes on sugar determination. *Journal of Biological Chemistry*. 1951: 19–23.
- Song, M., Kim, N., Lee, S., and Hwang, S. (2007). Use of Whey Permeate for Cultivating *Ganoderma lucidum* Mycelia, *Journal Dairy Science*. 90: 2141–2146.
- Soxhlet, F. (1879). Die gewichtsanalytische Bestimmung des Milchfettes. *Polytechnisches J. (Dingler's)*. 232: 461.
- Strobel, H. J. and Russell, J. B. (1987). Regulation of  $\beta$ -Glucosidase in *Bacteroides rumenicola* by a Different Mechanism: Growth Rate-Dependent Derepression Herbert. *Applied and Environmental Microbiology*. 53 (10): 2505-2510.
- Swann, E. C. & Taylor, W. (1993). Higher Taxa of Basidiomycetes: an 18s rRNA Gene Perspective. *Mycologia* 85, 923-936.
- Thirkettle, J. (2000). B-253514 and Analogues; Novel Inhibitors of Lipoprotein Associated Phospholipase A2 Produced by *Pseudomonas fluorescens* DSM11579 III. Biotransformation Using Naringinase. *Journal of Antibiotic*. 53: 733–735.
- Torget, R., Walter, P., Himmel, M., and Grohmann, K. (1991). Dilute-acid pretreatment of corn residues and short-rotation woody crops. *Applied Biochemistry Biotechnology*. 28: 75–86.
- Ugliano, M., and Moio, L. (2006). The influence of malolactic fermentation and *Oenococcus oeni* strain on glycosidic aroma precursors and related volatile compounds of red wine. *Journal of Science Food and Agriculture*. 86: 2468–2476.
- Van Wyk, B. E. (2005). Tribe Podalyrieae. In: L. Lewis, B. Schrire, B. Mackinder and M. Lock (eds), *Legumes of the World*, pp. 267 – 271. Royal Botanic Gardens, Kew. ISBN 1900347806

- Vinoth Kumar V. (2010). Comparative studies on inducers in the production of naringinase from *Aspergillus niger* MTCC 1344. *African Journal of Biotechnology*. 9(45): 7683-7686.
- Wan Razali, W.A., Baharuddin, A.S., Talib, A.T., Sulaiman, A., Naim, M.N., Hassan, M.A., and Shirai, Y. (2012). Degradation of Oil Palm Empty Fruit Bunch (OPEFB) Fibre during Composting Process using in-vessel Composter. *Bioresources*. 7(4): 4786-4805.
- William B. Jensen. (2007). The Origin of the Soxhlet Extractor. *Journal of Chemical Education*. 84: 1913.
- Wood, T.M., and Bhat, K.M. (1988). Methods for measuring cellulase activities in *Methods in Enzymology*, eds. by Wood TM, Kellogg ST, London, Academic Press, Inc.
- Yadav, S. and Yadav, K.D.S. (2000). Secretion of  $\alpha$ -L-rhamnosidases by *Aspergillus terreus* and its role debittering of orange juice. *Journal of Scientific & Industrial Research*. 59: 1032–1037.
- Yadav, V., Yadav, P.K., Yadav, S., and Yadav, K.D.S. (2010).  $\alpha$ -l-rhamnosidase : A review. *Process Biochemistry*. 45: 1226–1235.
- Yadav, V., Yadav, S., Yadav, S. and Yadav, K.D.S. (2011).  $\alpha$ -l-rhamnosidase from *Aspergillus flavus* MTCC- 9606 isolated from lemon fruit peel. *Food Science and Technology International*. 46: 350–357.
- Yanai, T., and Sato, M. (2000). Purification and characterization of  $\alpha$ -L-rhamnosidases from *Pichia Angusta* X349. *Bioscience Biotechnology and Biochemistry*. 64: 2179–2185.
- Yang, F. C., and Liao, C. B. (1998). Effects of cultivating conditions on the mycelia growth of *Ganoderma lucidum* submerged flask cultures. *Bioprocess Engineering*. 19(3): 233- 236.
- Yang, F. C., Ke, Y. F., and Kuo, S. S. (2000). Effect of fatty acids on the mycelial growth and polysaccharide formation by *Ganoderma lucidum* in shake flask cultures. *Enzyme and Microbial Technology*. 27: 295-301.
- Yang, F. C., Ke, Y. F., and Kuo, S. S. (2000). Effect of fatty acids on the mycelial growth and polysaccharide formation by *Ganoderma lucidum* in shake flask cultures. *Enzyme and Microbial Technology*. 27: 295-30.
- Yu, H., Gong, J., Zhang, C., and Jin, F. (2002). Purification and characterization of ginsenoside - $\alpha$  - L- rhamnosidase. *Chemical and Pharmaceutical Bulletin*. 52: 175–178.
- Yusof, S., Ghazali, H. M., and King, G. S. (1990) Naringin content in local citrus fruits. *Food Chemistry*. 37: 113–21.