

UNIVERSITI PUTRA MALAYSIA

OPTIMIZATION OF PROCESSING PARAMETERS AND ACCELERATED SHELF LIFE OF KERISIK

NAEMAA BINTI MOHAMAD

FSTM 2014 35



OPTIMIZATION OF PROCESSING PARAMETERS AND ACCELERATED SHELF LIFE OF *KERISIK*



NAEMAA BINTI MOHAMAD

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science

December 2014

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

OPTIMIZATION OF PROCESSING PARAMETERS AND ACCELERATED SHELF LIFE OF *KERISIK*

By

NAEMAA BINTI MOHAMAD

December 2014

Chair: Prof. Russly Bin A. Rahman, PhD Faculty: Food Science and Technology

The study was conducted since there is no information about optimization of kerisik (roasted coconut paste) processing parameters and accelerated shelf life study of the optimized processing parameters of kerisik to date. Therefore, the objectives of the study were to use response surface methodology (RSM) to optimize roasting temperature and roasting time to produce kerisik and consequently to use accelerated shelf life study to determine the shelf life of the kerisik. The optimization of kerisik processing parameters such as roasting temperature (130-150 °C) and roasting time (210-240 minutes) was achieved by using central composite design (CCD) via Design Expert software. Responses measured were color (L and b values) and furan compound (measured at 280 and 290 nm). Subsequently, the kerisik were packed in nylon and polypropylene packaging films and then were stored at two storage temperatures namely 38 °C for 38 days and 48 °C for 30 days for each type of packaging film. The analyses for accelerated shelf life study included moisture content (%), total color different (TCDH), peroxide value (PV) (meq kg⁻¹), acid value and p-anisidine value. Accelerated shelf life study employed PV as it main parameter in shelf life estimation with PV of 10 meg per kg fat was considered as the threshold limit for oxidative rancidity. Result showed that the optimum processing conditions for kerisik (laboratory scale) by using oven were found to be 130 °C (roasting temperature) for 227 minutes (roasting time). Predicted value of responses (color and furan) obtained using model equations were in good agreement with the experimental values. L (lightness), b (yellowness) and furan measured at 280 and 290 nm were significantly affected by the processing parameters. Finally, the shelf life of kerisik in nylon packaging film was predicted to be 46-49 days and 41-43 days for kerisik in polypropylene film. Kerisik producers may consider the generated optimized processing parameters and accelerated shelf life information from this study for kerisik product development in order to find the best method to replace the conventional way of kerisik production.

Abstrak tesis yang dikemukan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah sarjana sains

PENGOPTIMUMAN PARAMETER PEMPROSESAN DAN TEMPOH SIMPAN YANG DIPERCEPAT BAGI KERISIK

Oleh

NAEMAA BINTI MOHAMAD

Disember 2014

Pengerusi: Prof. Russly Bin A. Rahman, PhD Fakulti: Fakulti Sains dan Teknologi Makanan

Kajian ini dijalankan kerana masih tiada maklumat yang boleh didapati mengenai pengoptimuman parameter pemprosesan bagi kerisik dan kajian tempoh penyimpanan yang dipercepat bagi kerisik yang dihasilkan daripada proses pengoptimuman. Objektif-objektif bagi kajian ini adalah penggunaan kaedah rangsangan permukaan bagi proses pengoptimuman suhu pemanggangan dan masa pemanggangan bagi penghasilan kerisik dan seterusnya menggunakan kajian tempoh penyimpanan yang dipercepat bagi menentukan tempoh penyimpanan bagi kerisik yang dihasilkan. Pengoptimuman parameter pemprosesan bagi kerisik seperti suhu pemanggangan (130-150 °C) dan masa pemanggangan (210-240 minit) dicapai dengan menggunakan rekaan komposit tengah melalui perisian Design Expert. Respon-respon yang diuji adalah warna (nilai L dan b) dan kandungan furan (yang diukur pada 280 dan 290 nm). Kerisik dibungkus dalam plastik nylon dan plastik polypropylene disimpan pada dua suhu berbeza iaitu 38 °C selama 38 hari dan 48 °C selama 30 hari. Analisis yang dijalankan untuk kajian tempoh simpan yang dipercepat adalah kandungan kelembapan (%), perbezaan warna keseluruhan, nilai peroksida (meq kg⁻¹), nilai asid dan nilai p-anisidine. Had nilai peroksida 10 meq bagi setiap kg lemak telah dipilih sebagai parameter utama untuk penentuan jangka hayat kerisik. Suhu pemanggangan 130 °C dan masa pemanggangan 227 minit telah dipilih sebagai parameter yang optimum bagi pemprosesan kerisik dalam skala makmal. Nilai respon yang dijangka untuk warna dan furan didapati selari dengan nilai yang didapati daripada ujikaji di makmal. Parameter pemprosesan terbukti mempengaruhi nilai L, b dan furan yang diukur pada 280 dan 290 nm. Tempoh simpan bagi kerisik dalam plastik nylon didapati adalah selama 46-49 hari manakala 41-43 hari adalah tempoh simpan bagi kerisik dalam plastik polypropylene. Akhir sekali pengusaha kerisik boleh mempertimbang maklumat yang didapati daripada kajian ini bagi tujuan pembangunan produk kerisik dalam usaha mencari pendekatan terbaik bagi menggantikan cara lama dalam penghasilan kerisik.



ACKNOWLEDGEMENTS

In the name of Allah, the Most Gracious and the Most Merciful. I am grateful to Allah for the good health and wellbeing that were necessary to complete this study.

First and foremost, I would like to take this opportunity to convey my greatest gratitude and appreciation to all the people who had assisted me throughout this project. Firstly, I would like to express my deepest gratitude to my supervisor, Yang Berbahagia Profesor Dr. Russly Bin Abd. Rahman who has supervised me from the beginning until completion of this project. This project cannot be completed without his advice, comments, suggestions and invaluable guidance throughout the planning and execution of this project. I could not have imagined having a better advisor and mentor for my study. I also would like to extend my special thank you to the member of my supervisory committee; Professor Dr. Jamilah Bakar and Assoc. Prof. Dr. Roselina Karim for sharing their knowledge and experience in analytical work. I also would like to thank Dr. Boo Huey Chern who had assisted me in statistical aspects in my thesis.

Special thanks and appreciation to the staffs of the Faculty Food Science and Technology, Universiti Putra Malaysia, Serdang, Malaysia for their guidance, encouragement and advice throughout this project especially to En. Azman, Pn. Lina, Pn. Pija, Pn. Suraya and Pn. Liza and also special thanks to a former UPM staff En. Azman. The insightful information and help is duly acknowledged.

I'm also feeling thankful to the scholarship of Young Lecturer Scheme, Universiti Teknologi MARA, Malaysia which funded me financially and other supportive aspect during my study. I am feeling so grateful that I could not wait to repay back all the good deeds that UiTM has done to me. I would like to express my deepest gratitude to my advisor from Faculty of Applied Science, UiTM, Dr. Zainal Samicho for his excellent guidance, caring and patience. I attribute the level of my master degree to his encouragement and effort and without him this thesis, too, would not have been completed or written. One simply could not wish for a better advisor. Thank you for believe in me.

I also would like to thank my colleagues Hanisah Hamzah, Nurul Aini Adnan, Suganya, Sook Wah, Yap Yuh Ming, Mahsa, Wan Farhana Yasmeen and other lab mates for their stimulating discussions, kind help, opinion sharing and friendly attitude and also for those who had involved directly or indirectly for their support and generous assistant throughout this project. My research would not have been possible without their helps. You all will remain my best fellows and thank you for making this study enjoyable to me. I really appreciate the kindness of all of them. Thank you very much.

Last but not least, my deepest thanks to my dearest parents Mohamad Harun and Norsimah Mohd for their unconditional support and unceasing encouragement, patience, helps, advice, attention and love they have given me. There is no way to express my gratitude in words for them. I also would like to thank my sisters Manisah and Noor Hayati and my brothers Mohd Yusri and Shahrul Nizam that were always supporting me and encouraging me with their best wishes. I am also grateful to my partner who supported me through this venture and was always there cheering me up and stood by me through the good times and bad.



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Russly Bin Abdul Rahman, PhD

Professor Faculty of Food Science and Technology Universiti Putra Malaysia (Chairman)

Jamilah Binti Bakar, PhD

Professor Faculty of Food Science and Technology Universiti Putra Malaysia (Member)

Roselina Binti Karim, PhD

Associate Professor Faculty of Food Science and Technology Universiti Putra Malaysia (Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from the supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature:

Date: _____

Name and Matric No.: Naemaa Binti Mohamad GS 26889

Declaration by Members of Supervisory Committee

This is to confirm that

C,

- the research conducted and the writing of this thesis was under our supervision
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to

Signature: Name of Chairman of Supervisory Committee:	Signature: Name of Member of Supervisory Committee:
Signature: Name of Member of Supervisory Committee:	

TABLE OF CONTENTS

		Page
ABSTRACT ABSTRAK ACKNOWLI APPROVAL DECLARAT LIST OF TA LIST OF FIC LIST OF AB	EDGEMENTS ON BLES URES BREVIATIONS	i iii v vii xii xiv xv
CHAPTER		
1	INTRODUCTION	1
2	LITERATURE REVIEW	6
	2.1 Coconut (<i>Cocos mucifera</i> L.)	6
	2.1.1 Coconut production in South East Asia	6
	2.1.2 Coconut kernel and its application	7
	2.1.3 Kerisik (roasted coconut paste) processing	7
	2.1.4 Coconut kernel rancidity	8
	2.2 Roasting	9
	2.2.1 Introduction	9
	2.2.2 Roasting parameters range	9
	2.2.3 The effects of roasting parameters on oxidation and furan compound	12
	2.2.3.1 Oxidation test methods	13
	2.2.3.2 Furan compounds methods	13
	2.2.4 The effects of roasting parameters on	13
	2.2.5 The effects of roasting parameters	1/
	on other attributes	14
	2.3 Response Surface Methodology (RSM)	14
	2.3.1 Introduction	14
	2.3.2 The optimization of processing parameters using RSM	15
	2.4 Shelf life study	16
	2.4.1 Accelerated shelf life testing (ASLT)	16
	2.4.2 Accelerating factors	16
	2.4.2.1 Temperature	16
	2.4.2.2 Light	20
	2.4.2.3 Water activity and relative humidity	20
	2.4.3 Arrhenius equation	20
	2.4.4 Modeling other quality deterioration	21
	2.4.5 Arrhenius and linear model	22
	2.5 Packaging	23

2.5 Packaging2.6 Lipid oxidation

26

2.7	Moisture content	28
2.8	Color	29

3 OPTIMIZATION OF KERISIK (ROASTED COCONUT PASTE) PROCESSING PARAMETERS USING RESPONSE SURFACE METHODOLOGY (RSM) 2.1 Introduction 20

3.1	Introd	uction	30
3.2	Mater	ials and methods	30
	3.2.1	Raw material and chemicals	30
	3.2.2	Preparation of samples	31
	3.2.3	Experimental design and statistical analysis	32
	3.2.4	Optimization and verification of model	33
	3.2.5	Moisture content by Karl Fischer titration	33
	3.2.6	Furan determination	34
	3.2.6.	1 Kerisik oil extraction	34
	3.2.6.2	2 Furan extraction in kerisik oil	34
	3.2.6.3	3 UV-VIS spectra of oils	35
	3.2.7	Color determination	35
3.3	Result	s and Discussions	36
	3.3.1	Fitting the response surface model to	36
		significant independent variables	
	3.3.2	Effects of roasting temperature and	41
		roasting time on color	
	3.3.3	Effects of roasting temperature and roasting	43
		Time on furan	
	3.3.4	Optimization for kerisik roasting parameters	45
	3.3.5	Verification of the final model	45
	3.3.6	Conclusion	46

SHELF LIFE TESTING OF KERISIK (ROASTED COCONUT PASTE) STORED UNDER DIFFERENT CONDITIONS

4

4.1	Introdu	action	47
4.2	Materia	als and Methods	47
	4.2.1	Raw material and chemicals	47
	4.2.2	Preparation of sample	47
	4.2.3	Moisture content by Karl Fischer titration	48
	4.2.4	Color measurement	48
	4.2.5	Kerisik oil extraction	48
	4.2.6	Peroxide value determination	49
	4.2.7	Acid value determination	49
	4.2.8	p-anisidine value determination	50
	4.2.9	Experimental design	50
	4.2.10	Model estimation (kinetics of PV formation)	51
	4.2.11	Statistical analysis	52
4.3	Results	s and Discussions	52
	4.3.1	Moisture content	52
	4.3.2	Color analysis	55

4.3.3	Peroxide value	58
4.3.4	Acid value and p-anisidine value analysis	62
4.3.5	Accelerated shelf life modeling	68
4.3.6	Arrhenius and linear model	70
4.3.7	Conclusion	74

75 76

77

89

100

101

CONCLUSION AND RECOMMENDATION FOR FUTURE 5 **STUDIES**

- 5.1
- Summary and conclusion Recommendation for future research 5.2

REFERENCES/BIBLIOGRAPHY APPPENDICES **BIODATA OF STUDENT PUBLICATION**

LIST OF TABLES

Table		Page
1.1	Grade specification for coconut based on FAMA standard	2
1.2	Size classification for coconut based on FAMA standard	2
1.3	Minimum requirement for coconut based on FAMA standard	2
1.4	Maturity index for coconut based on FAMA standard	3
1.5	Qualities of coconut based on FAMA standard	3
2.1	Top coconut producing countries in South East Asia in 2011	6
2.2	Top coconut exporters in 2011	6
2.3	Published studies that employed temperature as the accelerating Factor	17
2.4	Comparison of nylon and polypropylene properties	25
3.1	Coding levels of independent variables used in optimization of kerisik processing parameters	31
3.2	The central composite experimental design (in coded level of	32
	two variables) utilized in optimizing processing parameters for kerisik	
3.3	Experimental data obtained for the dependent variables	37
3.4	Regression coefficients (R^2) and probability (p) values of four dependent variables for kerisik roasting parameters optimization in term of coded factors	38
3.5	Analysis of variance showing significance of the variables on Responses	40
3.6	L, b and the values of furan (280 nm) and (290 nm)	45
3.7	Predicted and experimental values of roasting temperature at 130 °C and roasting time of 227 minutes	46
4.1	Unstandardized and standardized coefficients and probability (p) values for moisture content	54

4.2	Unstandardized coefficients and probability (p) values for L, a, b and total color difference	56
4.3	Unstandardized and standardized coefficients and probability (p) values for peroxide value	59
4.4	Unstandardized and probability (p) values for acid value and p-anisidine value	63
4.5	Values of the oxidation rate constants (k) and the regression coefficient (R^2) for PV at the experimental temperatures	70
4.6	Activation energy and Q10 values of kerisik stored under different conditions	71
4.7	In ts and 1/T (1/K)	72
4.8	Shelf life prediction by using Arrhenius equation (PV critical limit 10meq/kg fat)	73
4.9	Q ₁₀ values for roasted coconut paste under different conditions using Arrhenius and linear model	73
4.10	Estimated shelf life of kerisik under different conditions using Arrhenius and linear model at 25 °C	73

C

LIST OF FIGURES

Figur	e	Page
3.1	Response surface for L of kerisik as a function of roasting temperature and roasting time	42
3.2	Response surface for b of kerisik as a function of roasting temperature and roasting time	42
3.3	Response surface for furan (280 nm) of kerisik as a function of roasting temperature and roasting time	44
3.4	Response surface for furan (290 nm) of kerisik as a function of roasting temperature and roasting time	44
4.1	Moisture content observed at 38 and 48 °C.	53
4.2	Moisture content observed in nylon and polypropylene films	53
4.3	Peroxide value observed at 38 and 48 °C	61
4.4	Peroxide value observed in nylon and polypropylene films	61
4.5	Acid value observed at 38 and 48 °C	64
4.6	Acid value observed in nylon and polypropylene films	64
4.7	p-anisidine value observed at 38 and 48 °C	67
4.8	P-anisidine value observed in nylon and polypropylene films	67
4.9	Zero order reaction of peroxide value of kerisik in nylon packaging during storage at 38 and 48 °C	68
4.10	Zero order reaction of peroxide value of kerisik in polypropylene packaging during storage at 38 and 48 °C	69
4.11	Arrhenius shelf life plot of zero order PV of of kerisik stored in nylon and polypropylene film exposed to increased temperatures. Points are from experimental data. Lines are result of best fitting.	70
4.12	Arrhenius shelf life plot and equation for kerisik	72

LIST OF ABBREVIATIONS

0	Degree
μ	Micro
<	Less than
%	Percentage
AOCS	American Oil Chemists' Society
AV	Acid Value
ASLT	Accelerated Shelf Life Testing
b	Yellowness Value
°C	Degree Celcius
сс	Penetrant concentration
CDA	Conjugated Dienoic Acid
CL	Chemiluminescence
CODEX	Codex Alimentarius
CV	Coefficient of variation
DPPH	(2,2-diphenyl-1-picrylhydrazyl)
Ea	Activation Energy
FAMA	Federal Agricultural and Marketing Authority
FAOSTAT	Statistic from Food and Agriculture Organization of the United Nations
FDA	Food and Drug Administration
FFA	Free Fatty Acid
g	Gram
HMF	Hydroxymethyl Furfural
in	Inch
IP	Induction Period

k	Rate Constant
k	Thermal conductivity value
К	Kelvin
k _{ref}	Rate Constant at Reference Temperature
kg	Kilogram
L	Lightness Value
m	Metre
meq	Milliequivalents
ml	Milliliter
mg	Milligram
MPa	Millipascal
O ₂	Oxygen
OTR	Oxygen transmission rate
р	Probability
p-Av	Para-Anisidine Value
PV	Peroxide Value
R	Universal Gas Constant
RCP	Roasted Coconut Paste
RPM	Revolution Per Minute
RSM	Response Surface Methodology
T _{ref}	Reference Temperature
TBARS/TBA	Thiobarbituric Acid Reactive Substance
TCDH	Total Color Different
Tg	Glass transition temperature
T _m	Crystalline melting temperature
ΤΟΤΟΧ	Total Antioxidant

WWattWVTRWater vapor transmission rate



CHAPTER ONE

INTRODUCTION

Coconut palm (*cocos nucifera* L.) is one of the important plantation crops in tropical countries owing to its various parts that possess multiple benefits both for human and stock usage (Chan and Elevitch, 2006 and Guarte et al., 1996). Coconut kernel and coconut oil are two products of coconut palm that were important in this study since kerisik (roasted coconut paste) was prepared from roasting of the grated coconut kernel and then the roasted grated coconut kernel was ground by mechanical action and producing oily paste. Coconut oil is believed to have many medicinal properties such as anticoagulant, antiviral and antiseptic effects (Mandal and Mandal, 2011). The low unsaturated fatty acids in coconut oil are proven to increase the oxidative stability of coconut oil against oxidation and therefore it is suitable to be used in many products with the objective to prolong the shelf life of the products that are using it as the ingredient (Guarte et al., 1996).

There is growing trend of making various food products from different kernel based of coconut for human consumption. A mature kernel inside the coconut nut can be used in various ways such as eaten raw or being processed. Usually the mature kernel would be dried and grated for the purpose to be used in many dishes and delicacies (Chan and Elevitch, 2006). In kerisik making for this study, it was important to be consistent in choosing the coconut based on basic requirements such as the coconut must be free from defects, free from damages and uniform in maturity (color index). Federal Agricultural and Marketing Authority (FAMA) standard has been used in the coconut selection for this study and the standards used are listed in table 1 to table 5. In summary, coconut grade number two (Table 1.1), medium size coconut (Table 1.2), all requirements listed in Table 1.3, coconut with maturity index 2 and 3 (Table 1.4) and coconut with acceptable quality in the market (Table 1.5) were selected to be used in this study (Federal Agricultural and Marketing Authority 2011)

Grade	Specification
Premium	1. Same variety and clean
	2. Uniform in size (\leq 5%) and maturity (\leq 5%)
	3. Free from defects ($\leq 5\%$)
	4. Free from damages ($\leq 5\%$)
1	1 Same variety and clean
1	2. Uniform in size (<10%) and maturity (<5%)
	3. Almost free from defects ($\leq 5\%$)
	4. Almost free from damages ($\leq 10\%$)
2	1. Same variety and clean
	2. Uniform in size ($\leq 10\%$) and maturity ($\leq 10\%$)
	3. Rather free from defects (\leq 5%)
1 a 1	4. Rather free from damages ($\leq 10\%$)

Table 1.1. Grade specification for coconut based on FAMA standard (FederalAgricultural and Marketing Authority, 2011)

Table 1.2. Size	classification	for	coconut	based	on	FAMA	standard	(Federal
Agricultural and 1	Markething Au	thor	ity, 2011)					

0	3 ·	
Size	Code	Weight (g)
Extra large	XL	>2000
Large	L	>1500-2000
Medium	М	>1000-1500
Small	S	800-1000

Table 1.3. Minimum requirement for coconut based on FAMA standard (Federal Agricultural and Marketing Authority, 2011)

(1 odorar Agricultur and Marketing Authority, 2011)		
	Requirements for coconut	
1.	Has reach maturity as required by market	
2.	Rather free from defects and fit for human consumption	
3.	Still has coconut husk on top of coconut shell	
4.	Clean	

Index	Description
1	1. Immature kernel
	2. Light colored coconut shell
	3. No sound produces when shaken
2	1. Mature kernel
	2. Light brown colored coconut shell
	3. Produces sloshing sound when shaken
3	1. Mature kernel
	2. Light brown colored coconut shell
	3. Produces sloshing sound when shaken
	4. Budding
4	1. Too mature kernel
	2. Dark brown colored coconut shell
	3. No sound produces when shaken
	4. Bud

Table 1.4. Maturity index for coconut based on FAMA standard (Federal Agricultural and Marketing Authority, 2011)

Table 1.5. Qualities of coconut based on FAMA standard (Federal Agriculturaland Marketing Authority, 2011)

Quality	Description
Best	1. Mature kernel
	2. Clean
	3. Free from defects and damages
Accepted in	1. Mature kernel
the market	2. Clean
	3. Minimum defects and rather free from damage
Not accepted	1. Immature kernel
in the market	2. Dirty
	3. Major defects (pests and diseases) and damage
	(mechanical damages that affecting quality and shell
	life)

Kerisik is a semi convenient food product that usually being used in cooking. Semi convenient food products are to be used with raw or semi-cooked materials and followed by minimal preparations afterward. Minimal preparation includes heating in microwave or frying in wok or short cooking in a container which does not need a lot of time in preparing the meal. Thus kerisik is categorized under semi-prepared ingredient. The process to make kerisik is usually simple and does not need costly materials. Kerisik is being used in many dishes in Malaysia to add flavour. It can make dishes more tasty and appealing. It is specially prepared for dishes with beef,

chicken, fishes and seafood. Kerisik also suitable to be used in *kerabu* (salads) and *rendang* (spicy meat dish) served with aubergines or pineapple (Nazrul, 2010). The fresh coconut kernel will first be grated by a coconut grinder. After that, the grated coconut kernel will be roasted until it turns crispy and flaky. Soon after it cooled, it will be blended until the fried grated coconut kernel particles become fine. At this level, the kerisik will excretes oil. Some manufacturers separate the oil from kerisik and the kerisik texture becomes harder than the kerisik which oil is not separated from the kerisik. It is depend on the consumer to choose which one they prefer (Mohammad, 2010).

The acceptable characteristics of kerisik include it must be in paste form, the color of the paste could vary from light brown to brown and have roasted or fried coconut aroma. The aroma of kerisik is unacceptable when burnt coconut aroma could be detected due to high temperature used during frying or roasting of grated coconut kernel, or the time taken to roast or fry the grated coconut kernel was too long. The grated coconut kernel will be heat treated by the action of toasting (Johan et al., 2007), frying (Hansen, 1993) or roasting (Tan, 2013). Afterward, to make the heat treated grated coconut kernel becomes an oily paste, it will be pounded in a mortar (Hansen, 1993) or ground (Tan, 2013). By varying the method to produce kerisik, the kerisik producers may increase the kerisik production by replacing the traditional way to produce kerisik that is time consuming and requires a lot of commitments with an easier method to produce the kerisik. Young generations nowadays prefer instant foods that are easy to prepare. Good response has been received from them towards convenience traditional food like kerisik and this trend could boost up the selling of the kerisik and benefiting small medium enterprise (Mohd et al., 2013).

Unfortunately most kerisik producers' lack of knowledge about the determination of optimum conditions for roasting the grated coconut kernel brought variation in quality of kerisik. By controlling the roasting parameters especially roasting temperature, the development of color and furan can be controlled (Beliz et al., 2009; Demir et al., 2002 and Özdemir and Devres, 2000). The maillard reaction happens by the interaction of carbonyls with amines during thermal processing (Eichner, 1981) and gives rise to brown pigments (Saklar et al., 2001) and these brown pigments increase as the browning and caramelization reactions progress (Moss and Otten, 1989; Mayer, 1985 and Buckholz et al., 1980). Several authors have found a good correlation between color parameters and the roasting process (Demir et al., 2002 and Saklar et al., 2001). Furan is considered as an undesirable compound (Ameur et al, 2006) that is produced due to thermal degradation of carbohydrates, pyrolysis of sugar at high temperature and oxidation of polyunsaturated fatty acids (Crews and Castle, 2007; Boekel, 1998 and Olano and Castro, 1996) and can be linked with color development as furan is considered a useful tool to evaluate browning reaction extension (Fallico et al., 2003).

No information exists about shelf life or the effects of storage temperature, packaging material and storage period on quality degradation of kerisik have been reported. Therefore, it is important to have an independent model that could

accurately predict the progress of chemical reaction occurring in a semi-solid phase during thermal processing and storage (Ahmed et al., 2004) to explain the effects of factors that might play major role in determining the shelf life of kerisik. Lipid oxidation is known as one of the factor that could affects foods shelf life. Hence peroxide value has been selected as the main indicator in this study to further understand about the quality degradation of kerisik. This is because, the mechanical actions such as grinding or pounding could increase kerisik susceptibility to oxidation. Thus, the control of roasting process is significant since both roasting temperature and roasting time could affect the product quality (Lee et al., 2010; Şimşek, 2007; Kahyaoglu and Kaya, 2006 and Özdemir and Devres, 2000).

Objectives of study

The scopes of the study were to establish the optimal processing conditions for kerisik and to determine its shelf life in different packaging materials using an accelerated shelf life testing (ASLT).

The specific objectives of this study were:

- 1. To optimize the roasting time and roasting temperature of kerisik production using response surface methodology (RSM).
- 2. To determine the shelf life of kerisik stored in different packaging materials (nylon and polypropylene films) and temperatures (38 and 48 °C) using accelerated shelf life testing (ASLT) method.

Limitation of study

There were some limitations in this study such as the study was done on specific characteristics of coconut kernels as specified in the introduction part and was purchased from pasar Sri Serdang, Malaysia. The equipments used were from the Faculty of Food Science and Technology, Universiti Putra Malaysia. Other limitations might include the collected data, data interpretations, software used and access to scientific literatures.

REFERENCES/BIBLIOGRAPHY

- Abou-Gharbia, H.A., Shehata, A.A.Y., & Shahidi, F. (2000). Effect of processing on oxidative stability and lipid classes of sesame oil. *Food Research International*, *33*, 331-340.
- Ahmadian-Kouchaksaraei, Z., Varidi, M., Varidi, M.J., & Pourazarang, H. (2014). Influence of processing conditions on the physicochemical and sensory properties of sesame milk: a novel nutritional beverage. *Food Science and Technology*, 57, 299-305.
- Ahmed, J., Shivhare, U. S., & Singh, P. (2004). Colour kinetics and rheology of coriander leaf puree and storage characteristics of the paste. *Food Chemistry*, 84, 605-611.
- Akpinar, K., Karakas, B., Akça, H., & Certel, M. (2011). Determination of HMF in roasted flour/oil mixtures and effect of solvent used in the extraction procedure. *Food Chemistry*, 128, 513-516.
- Alamprese, C., Ratti, S., & Rossi, M. (2009). Effects of roasting conditions on hazelnut characteristics in a two-steps process. *Journal of Food Engineering*, 95, 272-279.
- Altaki, M. S., Santos, F. J., & Galceran, M. T. (2011). Occurence of furan in coffee from spanish market. Contribution of brewing and roasting. *Food Chemistry*, *126*, 1527-1532.
- Ameur, L. A., Trystram, G., & Aragon, B. I. (2006). Accumulation of 5hydroxymethyl-2-furfural in cookies during the baking process. Validation of an extraction method. *Food Chemistry*, 98, 790-796.
- Anese, M., Nicoli, M.C., Verardo, G., Munari, M., Mirolo, G., & Bortolomeazzi, R. (2014). Effect of vacuum roasting on acrylamide formation and reduction in coffee beans. *Food Chemistry*, 145, 168-172.
- AOCS. (2009). *Acid value (Cd 3d-63)*. USA: Official methods and Recommended Practices of the American Oil Chemists' Society.
- AOCS. (2009). *p-Anisidine value (Cd 18-90)*. USA: Official Methods and Recommended Practices of the American Oil Chemists' Society.
- AOCS. (2009). *Peroxide value acetic acid-isooctane method (Cd 8b-90)*. USA: Official Methods and Recommended Practices of the American Oil Chemists' Society.
- Atalar, I., & Dervisoglu, M. (2015). Optimization of spray drying process parameters for kefir powder using response surface methosology. *Food Science and Technology*, 60, 751-757.
- Bachir bey, M., Meziant, L., Benchikh, Y., & Louaileche, H. (2014). Deployment of response surface methodology to optimize recovery of dark fresh fig (*ficus*

carica L., *var. azenjar*) total phenolic compounds and antioxidant activity. *Food Chemistry*, *162*, 277-282.

- Bakkalbaşi, E., Yilmaz, Ö. M., Javidipour, I., & Artik, N. (2012). Effects of packaging materials, storage conditions and variety on oxidative stability of shelled walnuts. *LWT-Food Science and Technology*, *46*, 203-209.
- Beliz, H. D., Grosch, W., & Schieberle, P. (2004). *Food Chemistry* (3rd ed.). Berlin: Springer.
- Beliz, H. D., Grosch, W., & Schieberle, P. (2009). *Food Chemistry* (4th revised ed.). (Extended, Ed.) Verlag berlin Heidelberg: Springer.
- Bhunia, K., Sablani, S.S., Tang, J., & Rasco, B. (2013). Migration of chemical compounds from packaging polymers during microwave, conventional heat treatment, and storage. *Comprehensive Reviews in Food Science and Food Safety*, 12, 523-545.
- Boekel, V. M.A.J.S. Food Chem, 62, (1998) 403.
- Bomfim, M.V.J., Zamith, H.P.S., & Abrantes, S.M.P. (2011). Migration of3caprolactam residues in packaging intended for contact with fatty foods. *Food Control, 22,* 681-684.
- Borompichaichartkul, C., Luengsode, K., Chinprahast, N., & Devahastin, S. (2009). Improving quality of macadamia nut (Macadamia integrifolia) through the use of hybrid drying process. *Journal of Food Engineering*, *93*, 348-353.
- Box, G. E., & Draper, N. (1987). *Empirical model-building and response surfaces*. New York: John Wiley.
- Buckholz, L. L., Daun, H., & Stier, E. (1980). Influence of roasting time on sensory attributes of fresh roasted peanuts. *Journal of Food Science*, 45, 547-554.
- Calligaris, S., Pieve, S. D., Kravina, G., Manzocco, L., & Nicoli, C. M. (2008). Shelf life prediction of bread sticks using oxidation indices: a validation study. *Journal of Food Science*, 51-56.
- Cämmerer, B., & Kroh, L. W. (2009). Shelf life of linseeds and peanuts in relation to roasting. *LWT-Food Science and Technology*, *42*, 545-549.
- Ćepo, D.V., Mornar, A., Nigović, B., Kremer, D., Radanović, D. and Dragojević, I.V. (2014). Optimization of roasting conditions as an useful approach for increasing antioxidant activity of carob powder. *Food Science and Technology*, *58*, 578-586.
- Chan, E., & Elevitch, C. R. (2006, April). Cocos nucifera (coconut) Arecaceae (palm family). *Special Profiles for Pacific Island Agroforestry, ver 2.1*, 1-27. Retrieved Febuary 1, 2014, from www.taditionaltree.org
- Che Man, Y. B., & Wan Hussain, W. R. (1998). Comparison of the frying performance of refined, bleached and deodorized palm olein and coconut oil. *Journal of Food Lipids, 5*, 197-210.

- Chiou, R. Y., Liu, C. D., Liu, C. P., Ferng, S., & Tsai, R. T. (1992). Characterization of peanut kernels as affected by harvest date and drying practices. *J.Agric.Food Chem.*, 40, 1536-1540.
- CODEX ALIMENTARIUS. (2014, February 25). Retrieved from CODEX STANDARD FOR NAMED VEGETABLE OILS: http://www.codexalimentarius.net/web/index en.jsp
- Conte, A., Scrocco, C., Brescia, I., & Nobile, M. A. (2009). Packaging strategies to prolong the shelf life of minimally processed lampascioni. *Journal of Food Enginnering*, 90, 199-206.
- Crews, C. and Castle, C. (2007). A review of the occurrence, formation and analysis of furan in heat processed foods. *Trends in Food Science and Technology*, 18, 365-372.
- Dayrit, F.M., Bunafe, O.E.M., Chainani, E.T., Vera, I.M.S.D., Dimzon, I.K.D., Gonzales, E.G., & Santos, J.E.R. (2007). Standards for essential composition and quality factors of commercial virgin coconut oil and its differentiation from RBD coconut oil and copra oil. *Philippine Journal of Science*, 136 (2), 119-129.
- Demir, A. D., Celayeta, J. M., Cronin, K., & Abodayeh, K. (2002). Modeling of the kinetics of colour change in hazelnuts during air roasting. *Journal of Food Engineering*, 55, 283-292.
- Durmaz, G., & Gökmen, V. (2010). Determination of 5-hydroxymethyl-2-furfural and 2-furfural in oils as indicator of heat pre-treatment. *Food Chemistry*, 123, 912-916.
- Durmaz, G., & Gökmen, V. (2011). Changes in oxidative stability, antioxidant capacity and phytochemical composition of Pistachia terebinthus oil with roasting. *Food Chemistry*, 128, 410-414.
- Eichner, K. (1981). Antioxidative effects of maillard reaction intermediates. *Progress in Food&Nutrition Science*, 5, 441-451.
- Fallico, B., Arena, E., & Zappalá, M. (2003). Roasting of hazelnuts; role of oil in colour development and hydroxymethylfurfural formation. *Food Chemistry*, *81*, 569-573.
- Fan, G., Han, Y., Gu, Z., & Chen, D. (2008). Optimizing conditions for anthocyanins extraction from purple sweet potato using response surface methodology (RSM). *LWT*, 41, 155-160.
- Farhoosh, R., Zohreh, S., & Yazdi, H. (2013). Shelf life prediction of olive oils using empirical model developed at low and high temperatures. *Food Chemistry*, *141*, 557-565.
- Federal Agricultural and Marketing Authority. (2011). Retrieved 9 January 2015 from <u>http://www.fama.gov.my/documents/10157/92f366fe-fdb2-49co-bdb2-f62d975c012a</u>

- Fouskake, M., Karametsi, K., & Chaniotakis, N. A. (2009). Method for the determination of water content in sultana raisins using a water activity probe. *Food Chemistry*, 82, 133-137.
- Galić, K., Ćurić, D., & Gabrić, D. (2009). Shelf life of packaged bakery goods-a review. *Critical Reviews in Food Science and Nutrition*, 49(5), 405-426.
- Gallina, A., Stocco, N., & Mutirelli, F. (2010). Karl Fischer titration to determine moisture in honey: A new simplified approach. *Food Control, 21*, 942-944.
- Gamlı, O. F., & Hayoglu, I. (2007). The effect of the different packaging and storage conditions on the quality of pistachio nut paste. *Journal of Food Engineering*, 78, 443-448.
- Gan, H.L., Tan, C.P., Che Man, Y.B., NorAini, I., & Nazimah, S. (2005). Monitoring the storage stability of RBD palm olein using the electronic nose. *Food Chemistry*, 89, 271-282.
- García, P. G., López, A. L., & Fernández, A. G. (2008). Study of the shelf life of ripe olives using an accelerated test approach. *Journal of Food Engineering*, *84*, 569-575.
- Gharby, S., Harhar, H., Aume, D. G., Haddad, A., Matthäus, B., & Charrouf, Z. (2011). Oxidative stability of edible argan oil: A two-year study. *LWT-Food Science and Technology*, 44, 1-8.
- Ghazali, H. M., Tan, A., Abdulkarim, S. M., & Dzulkifly, M. H. (2009). Oxidative stability of virgin coconut oil compared with RBD palm olein in deep-fat frying of fish crackers. *Journal of Food, Agriculture & Environment, 7*(3&4), 23-27.
- Gimenez, B., Guillen, M.C.G., Mateos, M.P., Montero, P., & Ruiz, G.M. (2011). Evaluation of lipid oxidation in horse mackerel patties covered with boragecontaining film during frozen storage. *Food Chemistry*, 124, 1393-1403.
- Gonșalves, E. M., Abreu, M., Brandaio, T. R., & Silva, C. C. (2011). Degradation kinetics of colour, vitamin c and drip loss in frozen brocolli (*Brassica oleracea L. sp*) during storage of isothermal and non-isothermal conditions. *International Journal of Refrigeration, 34*, 2136-2144.
- Granato, D., Ribeiro, J. C., Castro, I. A., & Masson, M. L. (2010). Sensory evaluation and physicohemical optimisation of soy-based desserts using response surface methodology. *Food Chemistry*, 121, 899-906.
- Guarte R.C, M. W., & M, K. (1996). Drying characteristics of copra and coconut oil. *Postharvest Biology and Technology*, *9*, 361-372.
- Gujral, H.S., Sharma, P., & Rachna, S. (2011). Effect of sand roasting on beta glucan extractability, physicochemical and antioxidant properties of oats. *Food Science and Technology*, 44, 2223-2230.

- Hafiza, Y., Linforth, R.S.T., & Cook, D.J. (2014). Flavour generation during commercial barley and malt roasting operations: A time course study. *Food Chemistry*, 145, 378-387.
- Hansen, B. Ethnic cook coconut rice for breakfast. *Los Angeles Times*, February 11, 1993, p.34.
- Ho, C.W., Wan Aida, W.M., Maskat, M.Y., & Osman, H. (2007). Changes in volatile compounds of palm sap (Arenga pinnata) during the heating process for production of palm sugar. *Food Chemistry*, *102*, 1156-1162.
- Holse, M., Skov, T., & Hansen, A. (2012). Oxidative storage stability of roasted marama beans (Tylosema esculentum). *Food Research International, 47,* 385-391.
- Homayoonfal, M., Khodaiyan, F., & Mousavi, M. (2015). Modelling and optimising of physicochemicals features of walnut-oil beverage emulsions by implementation of response surface methodology: effect of preparation conditions on emusion stability. *Food Chemistry*, 174, 649-659.
- Hussain, S. R., Terao, J., & Mathuushita, S. (1986). Effects of browning products of phospholipids on autoxidation of methyl lionete. *Journal of American Oil Chemist Society*, 60, 1528-1533.
- Ilaiyaraja, N., Likhith, K.R., Babu, G.R.S., & Khanum, F. (2015). Optimisation of extraction of bioactive compounds from *feronia limonia* (wood apple) fruit using response surface methodology (RSM). *Food Chemistry*, 173,348-354.
- Institut Penyelidikan dan Kemajuan Pertanian Malaysia. (1991). Perusahaan memproses kerisik (48 siri panduan untuk usahawan) ISSSN 0227-4767. Kuala Lumpur, Malaysia.
- Institut Penyelidikan dan Kemajuan Pertanian Malaysia. (1993). Reka bentuk dan penilaian alat pengering kelapa parut bagi pengeluaran kerisik (Laporan MARDI No. 169). Kuala Lumpur, Malaysia.
- Jayalekshmy, A., & Mathew, A. G. (1990). Changes in carbohydrates and proteins of coconut during roasting. *Food Chemistry*, *37*, 123-134.
- Jayalekshmy, A., Narayanan, C.S., & Mathew, A.G. (1991). Identification of volatile flavor compounds in roasted coconut. *JAOCS*, 68, 873-880.
- Jena, S., & Das, H. (2012). Shelf life prediction of aluminium foil laminated polyethylene packed vacuum dried coconut milk powder. *Journal of Food Engineering*, 108, 135-142.
- Johan, G.S.M.C., Russly, A.R., Bakar, J., Yaakob, B.C.M. and Rusul, G. (2007). Pasteurization, development and storage of *sous vide rendang* (spicy beef stew). *Journal of Food Service*, 18, 251-263.
- Kahyaoglu, T. (2008). Optimization of the pistachio nuts roasting process using response surface methodology and gene expression programming. *LWT*, 41, 21-33.

- Kahyaoglu, T., & Kaya, S. (2006). Modeling of moisture, color and texture changes in sesame seeds during the conventional roasting. *Journal of Food Engineering*, 75, 167-177.
- Kim, H.G., Kim, G.W., O. H., Yoo, S.Y., Kim, Y.O., & O, M.S. (2011). Influence of roasting on the antioxidant activity of small black soybean (*glycine max* L. Merrill). *Food Science and Technology*, 44, 992-998.
- Kinderlerer, J. L. (1994). Degradation of the lauric acid oils. *International Biodeterioration&Biodegradation*, 1994, 345-354.
- Kong, K. W., Ismail, A., Tan, C. P., & Rajab, N. F. (2010). Optimization of oven drying conditions for lycopene content and lipophilic antioxidant capacity in a by-product of the pink guava puree industry using response surface methodology. *LWT-Food Science and Technology*, 43, 729-735.
- Kothe, L., Zimmermann, B.F., & Galensa, R. (2013). Temperature influences epimerization and composition of flavanol monomers, dimers and trimers during cocoa bean roasting. *Food Chemistry*, 141, 3656-3663.
- Kumar, S.N. & Balakrishna, A. (2009). Seasonal variations in fatty acid composition of oil in developing coconut. *Journal of Food Quality*, 32, 158-176.
- Lasekan, O., & Abbas, K. (2011). Investigation of the roasting conditions with minimal acrylamide generation in tropical almond (Terminalia catappa) nuts by response surface methodology. *Food Chemistry*, 125, 713-718.
- Lee, S. W., Jeung, M. K., Park, M. H., Lee, S. Y., & Lee, J. H. (2010). Effects of roasting conditions of sesame seeds on the oxidative stability of pressed oil during thermal oxidation. *Food Chemistry*, 118, 681-685.
- Lee, S. Y., & Krochta, J. M. (2002). Accelerated shelf life testing of whey-proteincoated peanuts analyzed by static headspace gas chromatography. J. Agric. Food Chem., 50, 2022-2028.
- Limbo, S., Torri, L., Sinelli, N., Franzetti, L., & Casiraghi, E. (2010). Evaluation and predictive modelling of shelf life of minced beef stored in high oxygen modified atmosphere packaging at different temperatures. *Meat Science*, *84*, 129-136.
- Liu, Y., & Kitts, D.D. (2011). Confirmation that maillard reaction is the principal contributor to the antioxidant capacity of coffee brews. *Food Research International*, 44, 2418-2424.
- Lopez, A., & Pique, M. T. (1997). Influence of the drying conditions on the hazelnut quality. III Browning. *Drying Technology*, 15, 989-1002.
- Ludwig, I.A., Bravo, J., Pena, M.P.D., & Cid, C. (2013). Effect of sugar addition (torrefacto) during roasting process on antioxidant capacity and phenolics of coffee. *Food Science and Technology*, *51*, 553-559.

- Mandal, M. D., & Mandal, S. (2011). Coconut (*Cocos nucifera L.: Arecaceae*):In health promotion and disease prevention. *Asian Pacific Journal of Tropical Medicine*, 241-247.
- Mannekote, J. K., & Kailas, S. V. (2012). The effect of oxidation on the tribological performance of few vegetable oils. *Journal of Material Research and Technology*, 1(2), 91-95.
- Manzocco, L., Pannozo, A., & Calligaris, S. (2012). Accelerated shelf life testing (ASLT) of oils by light and temperature exploitations. *J. Am. Oil Chem. Soc.*, *89*, 577-583.
- Martins, A.C., Bukman, L., Vargas, A.M.M., Barizão, É.O., Moraes, J.C.G. Visentainer, J.V., & Almeida, V.C. (2013). The antioxidant activity of teas measured by the FRAP method adapted to the FIA system: Optimising the conditions using the response surface methodology. *Food Chemistry*, 138, 574-580.
- Mayer, K. P. (1985). Infra-red roasting of nuts, particularly hazelnuts. *Confectionary Production, 51*, 313-316.
- Mensitieri, G., Maio, E.D., Buonocore, G.G., Nedi, I., Oliviero, M., Sansone, L., & Iannace, S. (2011). Processing and shelf life issues of selected food packaging materials and structures from renewable resources. *Trends in Food Science&Technology, 22,* 72-80.
- Mensitieri^b, G., Scherillo, G., & Iannace, S. (2013). Flexible packaging structures for high pressure treatments. *Innovative Food Science and Emerging Technologies*, 17, 12-21.
- Mexis, S. T., Badeka, A. V., & Kontominas, M. G. (2009). Quality evaluation of raw ground almond kernels (Prunus dulas): Effect of active and modified atmosphere packaging, container oxygen barrier and storage conditions. *Innovative Food Science and Emerging Technologies, 10*, 580-589.
- Mohammad, L. (2010, May 3). Projek Kerisik. Retrieved 17 August 2014 from http:// http://projekkerisik.blogspot.com/
- Mohd, S.M.S., Mohd, S.M.Z., Norazmir, M.N., & Rosmaliza, M. (2013). Factors that restrict young generation to practice malay traditional festive foods. In Proceedia-Social and Behavioral Sciences 101, Proceedings of the AMER International Conference on Quality of Life, Langkawi, Malaysia, April 6-8, 2013.
- Moss, J. R., & Otten, L. (1989). A relationship between color development and moisture content during roasting of peanut. *Canadian Institute of Food Science and Technology*, 22, 34-39.
- Nair, K.G.P., Rajamohan, T., & Kurup, P.A. (1998). Coconut kernel protein modifies the effect of coconut oil on serum lipids. *Plant Foods for Human Nutrition*, 53, 133-144.

- Nazrul, A.S. (2010, May 31). Mahu eksport kerisik ke pasaran dunia Retrieved 17 August 2014 from <u>http://www.utusan.com.my/utusan/info.asp?y=2010&dt=0531&pub=Utusan</u> <u>Malaysia&sec=Agrobiz&pg=ag_02.htm</u>
- Nie, S., Huang, J., Hu, J., Zhang, Y., Wang, S., Li, C., Marcone, M., & Xie, M. (2013). Effect of pH, temperature and heating time on the formation of furan in sugar–glycine model systems. *Food Science and Human Wellness*, 2, 87-92.
- Ng, S. C., Anderson, A., Coker, J., & Ondrus, M. (2007). Characterization of lipid oxidation products in quinoa (Chenopodium quinoa). *Food Chemistry*, 101, 185-192.
- Nobile, M.A.D., Buonocore, G.G., Palmieri, L., Aldi, A., & Acierno, D. (2002). Moisture transport properties of polyamides copolymers intended for food packaging applications. *Journal of Food Engineering*, *53*, 287-293.
- Oracz, J., & Nebesny, E. (2014). Influence of roasting conditions on the biogenic amine content in cocoa beans of different *theobroma cacao* cultivars. *Food Research International*, 55, 1-10.
- Olano, A. and Castro, M. In: L.M.L. nollet (Ed.). Handbook of Food Analysis, Marcel Dekker, New York, 1996, p.1683.
- Owczarek-Fendor, A., Meulenaer, B.D., Scholl, G., Adams, A., Lancker, F.V., Eppe, G., Pauw, E.D., Scippo, M.L., & Kimpe, N.D. (2012). Furan formation in starch-based model systems containing carbohydrates in combination with proteins, ascorbic acid and lipids. *Food Chemistry*, 133, 816-821.
- Özdemir, M., & Devres, O. (2000). Analysis of color development during roasting of hazelnuts using response surface methodology. *Journal of Food Engineering*, 45, 17-24.
- Özdemir, M., Ackurt, F., Yildiz, M., Biringen, G., Gürcan, T., & Löker, M. (2001). Effect of roasting on some nutrients of hazelnuts (corylus avellena l.). *Food Chemistry*, 73, 185-190.
- Ozel, M.Z., Yanik, D.K., Gogus, F., Hamilton, J.F., & Lewis, A.C. (2014). Effect of roasting method and oil reduction on volatiles of roasted *pistachia terebinthus* using direct thermal desorption-GCxGC-TOF/MS. *Food Science and Technology*, *59*, 283-288.
- Özkanlı, O., & Kaya, A. (2007). Storage stability of butter oils produced from sheep's non-pasteurized and pasteurized milk. *Food Chemistry*, 100, 1026-1031.
- Palazón, M.A., Conesa, D.P., Abellán, P., Ros, G., Romero, F., & Vidal, M.F. (2009). Determination of shelf-life of homogenized apple-based beikost storage at different temperatures using Weibull hazard model. *Food Science and Technology*, 42, 319-326.

- Pua, C.K., Hamid, N.S.A., Tan, C.P., Mirhosseini, H., Rahman, R.A., & Rusul, G. (2008). Storage stability of jackfruit (Artocarpus heterophyllus) powder packaged in aluminium laminated polyethylene and metallized co-extruded biaxially oriented polypropylene during storage. *Journal of Food Engineering*, 89, 419-428.
- Pua^b, C. K., N. S., Tan, C. P., Mirhosseini, H., R. B., & Rusul, G. (2010). Optimization of drum drying processing parameters for production of jackfruit (*Artocarpus heterophyllus*) powder using response surface methodology. *LWT-Food Science and Technology*, 43, 343-349.
- Pumilia, G., Cichon, M.J., Cooperstone, J.L., Giuffrida, D., Dugo, G., & Schwartz, S.J. (2014). Changes in chlorophylls, chlorophyll degradation products and lutein in pistachio kernels (*pistachia vera* L.) during roasting. *Food Research International*, 65, 193-198.
- Puri, M., Sharma, D., Barrow, C.J., & Tiwary, A.K. (2012). Optimisation of novel method for the extraction of steviosides from *stevia rebaudiana* leaves. *Food Chemistry*, 132, 1113-1120.
- Raghavendra, S. N., & Raghavarao, K. S. (2010). Effect of different treatments for the destabilization of coconut milk emulsion. *Journal of Food Engineering*, *97*, 341-347.
- Ragnarsson, J. O., & Labuza, T. P. (1976). Accelerated shelf life testing for oxidation rancidity in foods. A review. England: Applied Science Publisher Ltd.
- Rattanathanalerk, M., Chiewchan, N., & Srichumpoung, W. (2005). Effect of thermal processing on the quality loss of pineapple juice. *Journal of Food Engineering*, 66, 259-265.
- Rendón, M.Y., Salva, T.D.J.G., & Bragagnolo, N. (2014). Impact of chemical changes on the sensory characteristics of coffee beans during storage. *Food Chemistry*, 147, 279-286.
- Risch, S.J. (2009). Food Packaging History and Innovations. J. Agric. Food Chem., 57, 8089-8092.
- Robertson, G. L. (1993). Shelf life of foods. In Food Packaging -Principles and Practice. In *Shelf life of foods* (pp. 338-380). New York: Marcel Dekker Inc.
- Saad, B., Wai, W.T., Lim, B.P., & Muhammad, I.S. (2007). Flow injection determination of anisidine value in palm oil samples using a triiodide potentiometric detector. *Analytica Chimica Acta*, *591*, 248-254.
- Saikia, S., Manhot, N.K., & Mahanta, C.L. (2015). Optimisation of phenolic extraction fromAverrhoa carambola pomace by response surface methodology and its microencapsulation by spray and freeze drying. *Food Chemistry*, 171, 144-152.

- Saklar, S., Katnas, S., & Ungan, S. (2001). Determination of optimum hazelnut roasting conditions. *International Journal of Food Science and Technology*, 36, 271-281.
- Shafisoltani, M., Salehifar, M., & Hashemi, M. (2014). Effects of enzymatic treatment using Response Surface Methodology on the quality of bread flour. *Food Chemistry*, 148, 176-183.
- Shakerardekani, A., & Roselina. K. (2013). Effect of different types of plastic packagings films on the moisture and aflatoxin contents of pistachio nuts during storage. J. Food Sci. Technol., 50(2), 409-411.
- Şimşek, A. (2007). The use of 3D-nonlinear regression analysis in mathematics modeling of colour change in roasted hazelnuts. *Journal of Food Engineering*, 78, 1361-1370.
- Singh, B. P., Jha, A., Sharma, N., & Rasane, P. (2013). Optimization of a process and development of a shelf life prediction model for instant multigrain dalia mix. *Journal of Food Process Engineering*, *36*, 811-823.
- Siriphanich, J., Saradhuldhat, P., Romphophak, T., Krisanapook, K., Pathaveerat, S. and Tongchitpakdee, S. (2011). Coconut (*cocos nucifera* L.). In *Postharvest Biology and Technology of Tropical and Subtropical Fruits*, ed. E. Yahia, pp. 8-35. Thailand: Woodhead Publishing Limited.
- Sirisompong, W., Jirapakkul, W., & Klinkesorn, U. (2011). Response surface optimization and characteristics of rambutan (*nephelium lappaceum* L.) kernel fat by hexane extraction. *Food Science and Technology*, 44, 1946-1951.
- Tananuwong, K., & Malila, Y. (2011). Changes in physicochemical properties of organic hulled rice during storage under different conditions. *Food Chemistry*, 125, 179-185.
- Tan, C. Coconut products. *The Strait Times*, March, 24, 2013.
- Tazi, S., Plantevin, F., Falco, C. D., Pulgserver, A., & Ajandouz, E. H. (2009). Effects of light, temperature and water activity on the kinetics of lipoxidation in almond-based products. *Food Chemistry*, *115*, 958-964.
- Toci, A.T., Neto, V.J.M.F., Torres, A.G., & Farah, A. (2013). Changes in triacylglycerols and free fatty acids composition during storage of roasted coffee. *Food Science and Technology*, *50*, 581-590.
- Top exports-Coconut (copra) oil-2011.n.d. Retrieved 28 February 2014 from FAOSTAT: http://faostat.fao.org/site/342/default.aspx
- Top exports-coconut-2011.n.d. Retrieved 19 February 2014 from FAOSTAT: http://faostat.fao.org/site/342/default.aspx
- Top productions-coconuts-2011.n.d. Retrieved 19 February 2014 from FAOSTAT: <u>http://faostat.fao.org/site/339/default.aspx</u>

- Verardo, V., Ferioli, F., Riciputi, Y., Iafelice, G., Marconi, E., & Florenza, M. (2009). Evaluation of lipid oxidation in spaghetti pasta enriched with long chain n-3 polyunsaturated fatty acids under different storage conditions. *Food Chemistry*, 114, 472-477.
- Vaidya, B., & Jong, B. E. (2013). Effect of temperature on oxidation kinetics of walnut and grapeseed oil. *Food Sci. Biotechnology*, 22, 273-279.
- Vignoli, J.A., Viegas, M.C., Bassoli, D.G., & Benassi, MDT. (2014). Roasting process affects differently the bioactive compounds and the antioxidant activity of arabica and robusta coffees. *Food Research International*,61, 279-285.
- Wadikar, D.D., Majumdar, T.K., Nanjappa, C., Premavalli, K.S., & Bawa, A.S. (2008). Development of shelf stable pepper based appetizers by response surface methodology (RSM). *LWT*, 41, 1400-1411.
- Waisundara, V.Y., Perera, C.O., & Barlow, P.J. (2007). Effects of different pretreatments of fresh coconut kernels on some of the quality attributes of the coconut milk extracted. *Food Chemistry*, 101, 771-777.
- Wegener, J. W., & Sánchez, P. L. (2010). Furan levels in fruit and vegetable juices, nutrition drinks and bakery products. *Analytica Chimica Acta*, 672, 55-60.
- Xu, Q., Shen, Y., Wang, H., Zhang, N., Xu, S., & Zhang, L. (2013). pplication of response surface methodology to optimise extraction of flavonoids fromfructus sophorae. *Food Chemistry*, 138, 2122-2129.
- Yang, J., Pan, Z., Takeoka, G., Mackey, B., Bingol, G., Brandl, M.T., Garcin, K., McHugh, T.H., & Wang, Hua. (2013). Shelf-life of infrared dry-roasted almonds. *Food Chemistry*, 138,671-678.
- Yang^b, J., Pan, Z., Takeoka, G., Mackey, B., Bingol, G., Brandl, M.T., Garcin, K., McHugh, T.H., & Wang, H. (2012). Shelf-life of infrared dry-roasted almonds. *Food Chemistry*, 138, 671-678.
- Yeretzian, C., Pascual, E.C., & Goodman, B.A. (2012). Effects of roasting conditions and grinding on free radical contents of coffee beans stored in air. *Food Chemistry 131*, 811-816.
- Yin, H. F., Fan, G. J., & Gu, Z. X. (2010). Optimization of culture parameters of selenium enrinched yeast *(saccharomyces cerevisiae)* by response surface methodology (RSM). *LWT-Food Science and Technology, 43*, 666-669.
- Youn, K. S., & Chung, H. S. (2012). Optimization of the roasting temperature and time for preparation of coffee-like maize beverage using the response surface methodology. *LWT-Food Science and Technology*, 46, 305-310.
- Zhou, G., Fu, L., & Li, X. (2015). Optimisation of ultrasound-assisted extraction conditions for maximal recovery of active monacolins and removal of toxic citrinin from red yeast rice by a full factorial design coupled with response surface methodology. *Food Chemistry*, 170, 186-192.

- Zoller, O., Sager, F., & Reinhard, H. (2007). Furan in food: headspace method and product survey. *Food Additives and Contaminants*, 24, 91-107.
- Żyżelewicz, D., Budryn, G., Krysiak, W., Oracz, J., Nebesny, E., & Bojczuk, M. (2014). Influence of roasting conditions on fatty acid composition and oxidative changes of cocoa butter extracted from cocoa bean of Forastero variety cultivated in Togo. *Food Research International, 63,* 328-343.

