



UNIVERSITI PUTRA MALAYSIA

***REACTIVE HYPEREMIA RESPONSE AND ULTRASONOGRAPHIC
FEATURES IN DETERMINING TISSUE RECOVERY IN RATS***

YAPP JONG HENG

FK 2018 70



**REACTIVE HYPEREMIA RESPONSE AND ULTRASONOGRAPHIC
FEATURES IN DETERMINING TISSUE RECOVERY IN RATS**

By

YAPP JONG HENG

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

July 2018

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

REACTIVE HYPEREMIA RESPONSE AND ULTRASONOGRAPHIC FEATURES IN DETERMINING TISSUE RECOVERY IN RATS

By

YAPP JONG HENG

July 2018

Chairman : Associate Professor Ir. Raja Mohd Kamil b. Raja Ahmad, PhD
Faculty : Engineering

Frequent repositioning of patient is important to relief pressure and correct damages on skin areas induced by pressure in order to avoid pressure ulcer (PU) development. Prolonging the recovery time for pressure relief may be beneficial but the amount of recovery time is unknown and worth investigating. In this study, the relationship between different recovery time and weight-bearing tissue conditions subjected to repetitive loading is investigated through histopathology examination in animal model using Sprague-Dawley rats. Furthermore, the effect of different recovery times on non-invasive PU indicators using peak reactive hyperemia (RH) trend and ultrasonographic (US) features are investigated. The hypothesis being tested is that different recovery time that simulates insufficient and sufficient recovery time could affect the RH trend and US feature. This study also compares the peak RH trends and US features with the tissue condition under histopathological examination to determine the potential use of these features in determining the adequate amount of recovery time of weight-bearing tissue in rats non-invasively.

Twenty-one male Sprague-Dawley rats (seven per group), with body weight of 385-485g, were categorised into three groups and subjected to different recovery times, each with three repetitive loading cycles at skin tissues above of right trochanter area. The first, second, and third groups were subjected to short (3 min), moderate (10 min), and prolonged (40 min) recovery, respectively, while applying fixed loading time and pressure (10 min and 50 mmHg, respectively). Peak RH was measured during the three cycles to determine RH trend (increasing, decreasing, and inconsistent). All rat tissues at the trochanter area were evaluated using ultrasound at pre- and post-experiment and rated by two raters to categorise the severity of tissue changes (no, mild, moderate, and severe change). The tissue condition was also evaluated using histopathological examination to distinguish between normal and abnormal tissues.

It was found that as the recovery time increases (3 min vs. 10 min vs. 40 min), the number of samples with normal tissue (43% vs. 43% vs. 100.00%) and inconsistent RH response trend (29% vs. 57% vs. 72%) also increases. In addition, as the recovery time decreases (40 min vs. 10 min vs. 3 min), there is an increased number of samples with abnormal tissue (0% vs. 57% vs. 57%) and increasing RH response trend (14% vs. 29% vs. 57%). Subsequently, most of the samples with increasing RH trend is related to abnormal tissue (71%); while inconsistent RH trends is more related to normal tissue (82%). However, the recovery time has no effect to the tissue condition evaluated under ultrasonographic. Moreover, there is no relationship between the tissue conditions evaluated under ultrasonographic and histopathological examination.

The results suggest that different recovery times affect the tissue condition and RH response trend during repetitive loading. In addition, RH trend over repetitive loading may serve as a new feature for determining tissue condition that leading to pressure ulcer. This animal study may be extended to human studies in the future.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

HYPEREMIA REAKTIF DAN CIRI-CIRI ULTRASONOGRAFI DALAM MENENTUKAN PEMULIHAN TISU TIKUS

Oleh

YAPP JONG HENG

Julai 2018

Pengerusi : Profesor Madya Ir. Raja Mohd Kamil b. Raja Ahmad, PhD
Fakulti : Kejuruteraan

Kekerapan mengangkat dan mengalihkan kedudukan pesakit adalah penting untuk memulihkan kawasan kulit yang cedera daripada tertekan supaya tekanan ulser boleh dicegah. Memanjangkan masa pemulihan untuk kawasan kulit tertekan mungkin berfaedah, manakala masa untuk penuh pemulihan adalah kurang dipahami dan patut dikaji. Oleh sebab itu, hubungan antara masa pemulihan yang berbeza terhadap keadaan kulit tertekan semasa tekanan berulang telah dikaji dalam penyelidikan ini dengan menggunakan pemeriksaan histopatologi terhadap jenis tikus Sprague-Dawley. Di samping itu, kesan masa pemulihan yang berbeza semasa tekanan berulang terhadap ujian penunjuk tekanan ulser bukan jenis invasif seperti trend hyperemia reaktif (RH) dan ciri-ciri ultrasonografi (US) juga dikaji dalam penyelidikan ini. Hipotesis yang diuji adalah masa pemulihan yang berbeza untuk mensimulasikan masa pemulihan yang tidak mencukupi dan mencukupi boleh memberi kesan terhadap trend RH dan ciri-ciri US. Tambahan pula, puncak trend RH dan ciri-ciri US juga dibandingkan dengan keadaan tisu yang diperiksa melalui ujian histopathology untuk menentukan potensi ujian-ujian penunjuk tersebut dalam menentukan masa pemulihan yang mencukupi untuk kawasan kulit tertekan daripada tikus secara tidak invasif.

Sebanyak dua puluh satu tikus jantan jenis Sprague-Dawley (tujuh tikus bagi setiap kumpulan) dengan berat badan 385-485g telah dikategorikan kepada tiga kumpulan. Setiap kumpulan diberi tiga tekanan berulang terhadap kawasan kulit *trochanter* kanan dengan masa pemulihan yang berbeza. Masa pemulihan untuk kumpulan-kumpulan tersebut adalah masa pendek (3 min) untuk kumpulan pertama, masa sederhana (10 min) untuk kumpulan kedua, dan masa panjang (40 min) untuk kumpulan ketiga. Manakala, masa (10 min) dan magnitud tekanan (50 mmHg) adalah tetap bagi semua kumpulan. Puncak RH semasa tiga tekanan berulang telah direkod untuk menentukan trend RH (meningkat, menurun, dan tidak konsisten). Semua

kawasan tisu di *trochanter* kanan bagi pra- dan pasca-eksperimen diimbas dengan diagnostik ultrasound dan dinilai oleh dua ahli radiologi untuk mengkategorikan keterukan perubahan tisu (tiada, ringan, sederhana, atau perubahan teruk). Di samping itu, kawasan tisu tertekan juga dinilai dengan analisis gambaran histopatologi untuk membezakan sama ada keadaan tisu sihat atau cedera.

Keputusan kajian menunjukkan bahawa memanjangkan masa pemulihan (3 min vs. 10 min vs. 40 min) menambah bilangan subjek dengan keadaan tisu sihat (43% vs. 43% vs. 100%) dan trend RH tidak konsisten (29% vs. 57% vs. 72%). Di samping itu, memendekkan masa pemulihan (40 min vs. 10 min vs. 3 min) menambah bilangan subjek dengan keadaan tisu cedera (0% vs. 57% vs. 57%) dan trend RH meningkat (14% vs. 29% vs. 57%). Di samping itu, subjek dengan trend RH meningkat adalah lebih berkaitan dengan tisu cedera (71%), manakala trend RH tidak konsisten adalah lebih berkaitan dengan tisu sihat (82%). Manakala, masa pemulihan yang berbeza tidak memberi keberkesanan terhadap tisu diimbas dengan diagnostik ultrasound. Selain itu, tiada perbezaan antara kategori perubahan tisu hasil imbasan ultrasound terhadap keadaan tisu hasil pemeriksaan histopathologi.

Kajian menunjukkan masa pemulihan yang berbeza memberi kesan terhadap keadaan tisu dan trend RH semasa tekanan berulang. Selain itu, trend RH dalam tekanan berulang berpotensi untuk berfungsi sebagai penunjuk baru untuk menentukan keadaan kulit tekanan yang berisiko tekanan ulser. Kajian haiwan ini boleh dilanjutkan kepada model manusia pada masa hadapan.

ACKNOWLEDGEMENTS

First and foremost, I would like to express my deepest gratitude to my advisor Assoc. Prof. Ir. Dr. Raja Mohd Kamil bin Raja Ahmad for the continuous support of my PhD study. I thank him for his kindness, motivation and patient guidance to my research in all the time and writing of this thesis. I appreciate all his contributions of time and ideas to teach me how good experimental and administrative are done.

I would also like to thank to the rest of my thesis committee: Prof. Datin Dr. Rozi binti Mahmud, Prof. Dr. Md. Zuki bin Abu Bakar, Assoc. Prof. Dr. Siti Anom binti Ahmad, Assoc. Prof. Dr. Norhafizah binti Mohtarrudin, Dr. Loqman Mohamad Yusof, and Dr. Ezamin bin Abdul Rahim, for their insightful comments and encouragement. I thank all of them for their profession in helping my research that has multiple aspects related with electrical and electronics, animal use, US assessment, and histopathology examination. I believe that I am tremendously fortunate to have worked with such a professional and great team.

Last but not the least, a special thanks to my family for supporting me spiritually and financially throughout my PhD life and writing this thesis.

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Raja Mohd Kamil Raja Ahmad, PhD

Associate Professor Ir.
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Siti Anom Ahmad, PhD

Associate Professor Ir.
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Rozi Mahmud, MBBS UM, MMedRad UKM

Professor Medical
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Ezamin Abdul Rahim, MD

Medical Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Norhafizah Mohtarrudin, MBBS, MPath

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Md Zuki Abu Bakar @ Zakaria, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____

Date: _____

Name and Matric No.: Yapp Jong Heng, GS35077

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xx
 CHAPTER	
1 INTRODUCTION	1
1.1 Problem statements	2
1.2 Hypothesis	3
1.3 Aim and objectives	3
1.4 Project scope	4
1.5 Research significance	5
1.6 Thesis outline	5
 2 LITERATURE REVIEW	 7
2.1 Pressure ulcer definition, prevalence, and effects	7
2.2 Etiology of pressure ulcer	7
2.2.1 Pressure	8
2.2.1.1 Cell deformation	8
2.2.1.2 Tissue ischemia	9
2.2.1.3 Reperfusion injury	10
2.2.1.4 Impaired lymphatic drainage	11
2.2.2 Prolonged pressure	12
2.2.3 Increased skin temperature	13
2.3 Risk factors of pressure ulcer	14
2.4 Prevention of pressure ulcer	15
2.4.1 Ineffectiveness of turning and using special support surface	15
2.4.2 Benefits of optimal repositioning	16
2.5 Variation of PU development among individuals	16
2.6 Pressure application for pressure ulcer study	16
2.6.1 Rig structure of pressure control system	17
2.6.2 Pressure electronic sensor	20
2.6.3 Linear movement technique	21
2.6.4 Control algorithm	21
2.6.5 Embedded system platform	22
2.6.6 System software and driver	22
2.7 Measurements of weight-bearing tissue response	23

2.7.1	Reactive hyperemia response	23
2.7.1.1	Parameters of reactive hyperemia	24
2.7.1.2	Laser Doppler flowmetry system	26
2.7.2	Ultrasonography evaluation	28
2.7.2.1	High frequency ultrasound system	30
2.7.2.2	Ultrasonographic features for the evaluation of tissue changes	31
2.7.2.3	Image enhancement technique	32
2.7.3	Histopathology examination	44
3	METHODOLOGY	45
3.1	Research design	46
3.2	Ethical approval	47
3.3	Subjects	47
3.4	Sample size	48
3.5	Instrumentation	48
3.5.1	Pressure control system	49
3.5.1.1	Hardware design	49
3.5.1.2	Software design for system requirement	50
3.5.2	Temperature monitoring system	65
3.5.3	Laser Doppler flowmetry system	66
3.5.4	High frequency ultrasound system	66
3.5.5	Light microscope	67
3.6	Data collection protocols	68
3.6.1	Preparation procedure	69
3.6.2	Screening procedure	70
3.6.3	Experimental procedure	70
3.6.4	Data post-processing and analysis	71
3.6.4.1	Harvested tissues	72
3.6.4.2	Recovery time	72
3.6.4.3	Post-processing of skin blood flow signal	73
3.6.4.4	Post-processing of ultrasound images	75
3.7	Statistical analysis	87
3.8	Summary	90
4	RESULTS	91
4.1	Biometric data	91
4.2	Pressure applied on skin	92
4.3	Skin blood flow response	95
4.4	Reactive hyperemia response trend	96
4.5	Ultrasonographic evaluation and inter-raters reliability	98
4.5.1	Percentage agreement	102
4.5.2	Kappa coefficient	103
4.6	Histopathology examination	106
4.7	Effects of recovery time on tissues	108
4.7.1	Recovery time vs. Histopathological examination	109
4.7.2	Recovery time vs. Reactive hyperemia trend	110
4.7.3	Recovery time vs. Ultrasonographic evaluation	111

4.8	Comparison between reactive hyperemia trend and ultrasonographic evaluation in determining tissue condition	113
4.8.1	Reactive hyperemia trend vs. Histopathological examination	113
4.8.2	Ultrasonographic evaluation vs. Histopathology examination	116
5	DISCUSSION	122
5.1	Summary of major contributions	122
5.2	Effect of recovery time on weight-bearing tissues	123
5.2.1	Recovery time vs. histopathological examination	124
5.2.2	Recovery time vs. Reactive hyperemia trend	124
5.2.2.1	Increasing RH trend	125
5.2.2.2	Inconsistent RH trend	126
5.2.2.3	Decreasing RH trend	126
5.2.3	Recovery time vs. Ultrasonographic features	127
5.2.4	Variation of tissue responses to identical pressure induced damage	128
5.3	Determining tissue conditions with reactive hyperemia and ultrasound features	128
5.3.1	Reactive hyperemia trend vs. Histopathology examination	129
5.3.2	Ultrasonographic features vs. Histopathological examination	130
6	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	132
6.1	Conclusion	132
6.2	Recommendations for future research	133
6.3	Contributions	133
	REFERENCES	135
	APPENDICES	157
	BIODATA OF STUDENT	255
	PUBLICATION	256

LIST OF TABLES

Table	Page
1 PU studies related with varied strain magnitude	9
2 PU studies related to skin blood flow response during loading	10
3 PU studies related to reperfusion injury	11
4 PU studies related to impaired lymphatic drainage	12
5 PU studies related to varied loading duration	13
6 PU studies related with varied skin temperature	14
7 PU studies related with RH response	24
8 Skin blood flow parameters to characterise the RH response	26
9 PU studies related with US	29
10 Specifications of ultrasound probes from Cortex Technology	31
11 Image enhancement techniques for medical images	34
12 Purpose and disadvantage of medical image enhancement techniques	41
13 PU studies using histopathology examination	44
14 Relationship of input force and output voltage for force sensor	56
15 Effects of PID gain on controlling pressure application	60
16 Intensity level conversion with histogram equalization	84
17 Detailed biometrical information of all subjects	92
18 Biometrical data of rat between groups given in mean (SD) and significant tested using ANOVA	92
19 Comparisons of the applied pressures (mean $\pm$ SD) between groups with different recovery time	95
20 Comparisons of the skin perfusion (mean $\pm$ SD) between groups with different recovery time	96

21	Peak SBF of each subject with 3 minutes recovery time during three repetitive loading	97
22	Peak SBF of each subject with 10 minutes recovery time during three repetitive loading	97
23	Peak SBF of each subject with 40 minutes recovery time during three repetitive loading	98
24	Severity of tissue changes and percent agreement for both raters	102
25	Counts of subjects with severity of tissue changes (in number)	103
26	Counts of subjects with severity of tissue changes (in percentage)	104
27	Histopathological examination results	108
28	Relationship between tissue condition and recovery time in terms of number of samples	109
29	Significance of the relationship between recovery times in terms of histopathology results	109
30	Number of samples with RH trends	110
31	Significance of the relationship between recovery times in terms of RH trends	111
32	Number of samples with severity of tissue change	112
33	Significance of the relationship between recovery times in terms of US categories	112
34	RH response trend and histopathological examination for each subject	114
35	Number of samples with tissue conditions response to RH trend	114
36	Contingency table (2 x 2) for paired RH trends (Increasing vs. Decreasing) and histopathological assessment	115
37	Contingency table (2 x 2) for paired RH trends (Increasing vs. Inconsistent) and histopathological assessment	115
38	Contingency table (2 x 2) for paired RH trends (Decreasing vs. Inconsistent) and histopathological assessment	115
39	Significance of the comparisons of the RH trends in terms of histopathology results	115

40	US evaluation and histopathological examination for each subject	117
41	Number of samples with tissue conditions examined under histopathology and ultrasonic evaluation	117
42	Contingency table (2 x 2) for paired US evaluations (No change vs. Mild change) and histopathological assessment, from rater 1 data	118
43	Contingency table (2 x 2) for paired US evaluations (No change vs. Moderate change) and histopathological assessment, from rater 1 data	118
44	Contingency table (2 x 2) for paired US evaluations (No change vs. Severe change) and histopathological assessment, from rater 1 data	118
45	Contingency table (2 x 2) for paired US evaluations (Mild change vs. Moderate change) and histopathological assessment, from rater 1 data	119
46	Contingency table (2 x 2) for paired US evaluations (Mild change vs. Severe change) and histopathological assessment, from rater 1 data	119
47	Contingency table (2 x 2) for paired US evaluations (Moderate change vs. Severe change) and histopathological assessment, from rater 1 data	119
48	Contingency table (2 x 2) for paired US evaluations (No change vs. Mild change) and histopathological assessment, from rater 2 data	119
49	Contingency table (2 x 2) for paired US evaluations (No change vs. Moderate change) and histopathological assessment, from rater 2 data	120
50	Contingency table (2 x 2) for paired US evaluations (No change vs. Severe change) and histopathological assessment, from rater 2 data	120
51	Contingency table (2 x 2) for paired US evaluations (Mild change vs. Moderate change) and histopathological assessment, from rater 2 data	120
52	Contingency table (2 x 2) for paired US evaluations (Mild change vs. Severe change) and histopathological assessment, from rater 2 data	120
53	Contingency table (2 x 2) for paired US evaluations (Moderate change vs. Severe change) and histopathological assessment, from rater 2 data	120
54	Significance of the comparisons for the severity of tissue changes in terms of histopathology results (rater 1)	121
55	Significance of the comparisons for the severity of tissue changes in terms of histopathology results (rater 2)	121

LIST OF FIGURES

Figure	Page
1 Blood vessels lined with endothelium	xxii
2 Pressure ulcer at sacrum (buttock) area	7
3 Lying positions	15
4 Computer-controlled indenter system consisting (A) lever, (B) motor, (C) stage, (D) load cell, and (E) cooling system	17
5 Pressure application device	18
6 Computer-controlled pressure system	19
7 Pressure application on trochanter area of rat	19
8 RH time domain parameters at post-loading	25
9 Overview of the theory behind laser Doppler flowmetry	27
10 LDF probe and probe holder	28
11 Examples of (a) pre- and (b) post- experimental ultrasound images (60 x 260 μ m resolution) with underlying skin tissues above of trochanter area of a Sprague Dawley rat (third subject of group with 10 min recovery time during repetitive loading in this study). The presence of oedema (circle), distorted tissue layers (arrows) and increased layer thickness on post-experimental image	32
12 Ultrasound images from previous pressure ulcer studies show presence of oedema at skin layer (dark area circled in Image A and hypo-echogenic layer in Image D) and discontinuous fascia layer (dashed circle in Image B and arrow in Image C)	33
13 Images of original, enhanced with HE, and enhanced with BBHE. Image enhanced with BBHE has mean brightness preserved as in original image	35
14 CLAHE clipping a histogram and uniform redistribution of the pixel values for image enhancement. Some pixels are exceeding the clip limit after the redistribution	36
15 Original and negative image of brain tumor image	37

16	Original image and its enhancement using thresholding technique	38
17	Original image (left) and enhanced image with log transformation (right)	38
18	Plot of power law transformation	40
19	Input image and its histogram and cumulative distribution	43
20	Flow of methodology	46
21	Time schedule of measurements for ultrasonic evaluation (US), pressure, SBF, skin temperature, and histological examination (HS). Ultrasonic evaluation was performed at pre- (white rectangle) and post-application of repetitive loading (dark grey rectangle) and rats were euthanized at the end for histopathological examination	47
22	Pressure control system, LDF system, and computer controlled system. (A) Rig structure, (B) linear actuator, (C) Perspex indenter, (D) flexible force sensor, (E) LDF probe, (F) temperature sensor, (G) LDF system, and (H) computer with LabVIEW graphical user interface	49
23	System block diagram for pressure monitoring, skin blood flow monitoring, temperature monitoring and pressure adjusting	50
24	Graphical user interface created with LabVIEW: (a) Repetitive loading setting panel for pressure magnitude, loading duration, and recovery duration. (b) Interface for monitoring pressure (blue signal), skin blood flow (red signal), and skin temperature (yellow signal)	51
25	Pressure control system block diagram	52
26	Closed loop flow chart for controller	53
27	Calibration apparatus set up	55
28	Force-to voltage schematic circuit of Flexiforce® sensor A401	55
29	Graph of input force against output voltage	57
30	PID Controller in LabVIEW	59
31	Duty cycle (percentage) to time (millisecond) VI	59
32	PID Controller in LabVIEW with error descriptions	60
33	System response with tuned PID controller	62

34	Simulation response with multiple set points	62
35	Hardware testing using rat model	63
36	Hardware testing response with multiple set points	64
37	Performance test of pressure control system and LDF system on human's hand	64
38	Pressure applications and skin blood flow response	65
39	DermaScan C USB High Resolution Ultrasound system components: A) power supply and cable, B) main unit to laptop power cable, C) main unit, D) 2D probe, E) ultrasound gel, and F) probe holder and accessories	67
40	Nikon Eclipse 80i microscope (A) and QImaging Micropublisher 3.3 RTV camera (B) for histopathology examination	68
41	The subject lay prone on the heating mat at base of pressure control system, and the tests were performed on the skin above of right trochanter area with following measurement devices: (A) temperature sensor for monitoring the skin temperature adjacent to trochanter area, (B) laser Doppler flowmetry probe inserted in the middle of 2 cm-diameter Perspex indenter for monitoring changes in skin perfusion, (C) FlexiForce sensor placed at the end of the indenter for recording and controlling the applied pressure, and (D) linear actuator for pressure application on the skin	69
42	Schematic of pressure application during baseline and three repetitive loadings for groups with (a) short, (b) moderate, and (c) prolonged recovery times	71
43	Preparation of tissue samples for histopathology: (a) dissection, (b) fixation, (c) embedment in paraffin, and (d) stained tissue on slide	72
44	Steps of skin blood flow signal processing: (a) Raw SBF signal, (b) low-pass filtered SBF signal, (c) peak RH extraction from filtered SBF signal, and (d) RH trend formation	74
45	Tissue layers characterized by colours with varied intensities	75
46	RGB ultrasound image of a Sprague Dawley Rat (forth subject of Group C with 40 min recovery time during repetitive loading in this study)	76
47	Greyscale image and its histogram and cumulative distribution	77

48	Entropy curve with different clip limit values: (Left) before fitting, (right) after fitting. The maximum curvature point is used to determine the suitable clip limit value for CLAHE application	80
49	Entropy curve with different block sizes	81
50	Image after CLAHE and its histogram and cumulative distribution. Muscle layer and trochanter bone (white arrow) were observed after the image enhancement	82
51	Image after histogram equalization and its histogram and cumulative distribution. Muscle layer and trochanter bone (black arrow) were observed after the image enhancement	85
52	Original greyscale image and enhanced image using histogram equalization	86
53	2 x 2 contingency table for Fisher's exact test	88
54	2 x 2 contingency table for odd ratios (OR)	89
55	Average applied pressure on seven subjects with 3 min recovery time during three repetitive loading	94
56	Average applied pressure on seven subjects with 10 min recovery time during three repetitive loading	94
57	Average applied pressure on seven subjects with 40 min recovery time during three repetitive loading	94
58	Examples of filtered skin perfusion signals obtained from three rats subjected to 10 minute recovery: A. increasing, B. decreasing, and C. inconsistent RH trend (green line) from three repetitive loadings	98
59	Examples of post-processed US images using "adaphisteq" function in Matlab to observe tissue changes at skin layer. The pre- and post-experiment images were obtained from rat A2 that subjected to 3 min recovery and rated as "obvious changes" by both raters due to the obvious oedema (red circles), distorted layer (yellow arrows), and increased thickness of dermal and hypodermal at post-images compared to pre-images	100
60	Examples of post-processed US images using "histeq" function in Matlab to observe tissue changes at muscle layer. The pre- and post-experiment images were obtained from rat A2 that subjected to 3 min recovery and rated as "obvious changes" by both raters due to the obvious oedema (red circles) at post-images compared to pre-images	101

- 61 Histopathology examination from histological slices of stained tissues above trochanter: (A) Occasional neutrophil (from subject A5), (B) mild neutrophils (from subject A7), (C) extravasated red blood cells into interstitial tissue (from subject A2), (D) longitudinal tears (from subject A4), and (E) pinkish area at interstitial tissues that represent oedema (from subject B3). Bar represents 50 μ m

107



LIST OF ABBREVIATIONS

PU	Pressure ulcer
RH	Reactive hyperemia
US	Ultrasonographic
HE	Histopathological examination
NI	National Instrument
NPUAP	National PU Advisory Panel
EPUAP	European PU Advisory Panel
LDF	Laser Doppler Flowmetry
PID	Proportional-integral-derivative

GLOSSARY OF TERMS

Endothelium An exceedingly thin single sheet of endothelial cells lining the inner wall of arteries and veins, while capillaries have walls that are only one endothelial cell in thickness (Alberts et al., 2014; Alberts et al., 2002; Chabner, 2013), as shown in Figure 1




Figure 1 : Blood vessels lined with endothelium
(Source: Chabner, 2013)

Vasodilator An agent that causes vasodilation, such as a nerve or hormone, that widens the blood vessels, which in turn decreases resistance to blood flow and lowers blood pressure (Fundukian, 2011; Soto, 2015)

CHAPTER 1

INTRODUCTION

Pressure ulcer (PU) is an area of localised tissue breakdown due to pressure and/or shear force sustained (Porter-Armstrong et al., 2018; Yarkony, 1994), causing prolonged tissue ischemia (lack of blood supply) (Jan et al., 2010; Liao, Burns, et al., 2013) which results in ulceration. The development of PU can also be accelerated with increasing temperature (Kokate et al., 1995; Romanelli et al., 2018) or moisture on skin (Kanj et al., 1998; Niezgoda et al., 2006; Romanelli et al., 2018; Thomas, 2006). PU can occur when subjects are bedridden, or wearing a poorly fitted medical devices, prosthesis or orthosis (Bouten et al., 2003; Boyko et al., 2018). They are more commonly developed once spinal-cord patients become wheelchair-bound and the risk remains high throughout their lifetime (Bhattacharya et al., 2015; Boyko et al., 2018; Krause et al., 2004; Raghavan et al., 2003; Salzberg et al., 1998; Sonenblum et al., 2011; Vidal et al., 1991). PU is a life-threatening complication due to breakdown tissue interacting with the external environment resulting in fatal septic infection. Therefore, the mortality rate of patients is increased compare with other conditions in hospital (9.1 - 11.6% death with PUs vs. 1.8 - 2.6% death for other conditions) (Bauer et al., 2016; Russo et al., 2008). It is also reported that the presence of PU in the patient has nearly three-fold increase in in-hospital death risk (Bo et al., 2003; Clough, 1994; Raju et al., 2015; Thomas et al., 1996). The ulcers could results in longer hospital stay (Chetter et al., 2016) with an additional 8.2 to 42 days (New et al., 2004; Whitty et al., 2017) or approximately three times longer (Gill, 2015; Russo et al., 2008) than those patients without PU. Additional care cost is needed to prevent further deterioration (costs US health care system \$9.1 billion to \$11.6 billion per year (Bauer et al., 2016), and costs National Health Service (NHS) in UK at between £1.4 billion and £2.1 billion, annually (Baillie, 2014; Dealey et al., 2012)). PUs can be developed at any parts of the body like the sacrum, heel, and buttock (Clark et al., 2017; Reger et al., 2009).

An effective protocol for PUs prevention is therefore needed to reduce the heavy burden on society caused by PUs. Repositioning the patients for at least every two hours is the gold standard (Gill, 2015; Lyder et al., 2008) to relieve prolonged pressure exerted on skin tissues repetitively for PUs prevention. In addition, special support surface is used to prevent skin breakdown by changing or redistributing pressure exerted on skin to restore and/or promote blood perfusion at underlying skin. However, from a survey, only 57 percent of respondents agreed on the effectiveness of the 2 hours repositioning in their intensive care units and remain at high risk of PU development (Latimer et al., 2015; Lyder et al., 2008), and using a different support surface with different repositioning frequencies does not influence the incidence of early PU (stage I) (Defloor et al., 2005; Yap et al., 2018).

Personalizing the repositioning interval has also been suggested to relieve the pressure on the skin for a sufficient amount of time and may alleviate the development of PU (Barnett et al., 1995; Pickham et al., 2016). In estimating an optimal body repositioning interval, the full tissue recovery time must be investigated. In order to know the full tissue recovery time, the tissue condition of an individual with applied recovery time must be evaluated in-vivo and non-invasive evaluation method is the only options for patient.

Reactive hyperemia (RH) response and ultrasonographic (US) features have been suggested as suitable features to identify people at risk of PU non-invasively (Liao, Burns, et al., 2013; Lucas et al., 2014). Previous skin blood flow and PU studies found that the degree of RH is related to the degree of ischemic event (a restriction in blood supply to tissues), and proportional to the pressure magnitude (Lachenbruch et al., 2015; Manorama et al., 2010), loading duration (Liao, Burns, et al., 2013; Matsubara et al., 1998; Thorfinn et al., 2007; Thorfinn et al., 2006), and increased temperature (Lachenbruch et al., 2015; Tzen et al., 2010). Studies related to early detection of PU using US features shows that the patients with at risk of PUs may be presented with oedema (Schäfer et al., 2015), distorted layered structure (Schäfer et al., 2015) and increased dermal and hypodermal thickness (Grap et al., 2015) that are considered as features of tissue changes induced by pressure (Schäfer et al., 2015) associated with inflammation (Grap et al., 2015).

1.1 Problem statements

The incidences of PU still exist even with two hours body repositioning and/or applying special support surface to relieve the pressure (Defloor et al., 2005; Latimer et al., 2015; Lyder et al., 2008; Salisbury, 1985). Reactive hyperemia response occurred after pressure relief could allow tissue prevented and recovered from pressure-induced ischemia (Jan, Liao, Rice, et al., 2013; Thorfinn et al., 2007). Therefore, pressure relief with sufficient recovery time can probably prevent pressure ulcer by fully reversing the tissue damages induced on weight-bearing tissues. The problem is how much recovery time is sufficient? This problem is investigated here.

Furthermore, the existing prevention approaches may not be sufficient for all individuals due to individual characteristics and tolerance for pressure (Braden et al., 2000; Coleman et al., 2014). Both individual susceptibility and local tolerance for pressure are affected by the conditions of individual such as health status and exposure to risk factors (Coleman et al., 2014). This results in varying development time of pressure ulcers between subjects. Therefore, individualised PU prevention method (Barnett et al., 1995; Källman, 2015; Kwong et al., 2016) is essential to alleviate the PU development. An effective protocol for determining tissue condition is therefore needed to indicate the effectiveness of prevention strategy that applied on the individual. The problem is how to determine the weight-bearing tissues condition accurately and non-invasively. The RH trend and US features that can be potentially used in determining weight bearing tissue condition non-invasively need to be

validated with histopathological examination and is addressed in this prospective study.

1.2 Hypothesis

The following hypotheses are tested:

- I. Different recovery time during repetitive loading can affects the tissue condition and response.
 - a. Short recovery time gives insufficient tissue recovery during repetitive loading and is presented as abnormal tissue under histopathological examination; while prolonged recovery time can promote tissue recovery and presented as normal tissue.
 - b. Peak RH increases from cycle to cycle of repetitive loading with short recovery time, and the peak does not increase with prolonged recovery time.
 - c. US assessment detects severe tissue changes induced by repetitive loading with short recovery time; while no/mild tissue changes with prolonged recovery time.
- II. Peak RH trend and US features can be used to indicate tissue condition non-invasively.
 - a. Peak RH increases from cycle to cycle of repetitive loading indicates accumulation of ischemic event at weight-bearing tissue and the tissue is in abnormal condition under histopathological examination; and the peak RH with non-increases from cycle to cycle is in normal condition under histopathological examination.
 - b. No tissue changes under US assessment has normal tissue condition under histopathological examination; while mild to severe tissue changes under US assessment has abnormal tissue condition under histopathological examination.

1.3 Aim and objectives

The aim of this study is to investigate the possibility of using certain features (RH trends and US features) measured using non-invasive modalities in determining sufficient recovery time and the condition of weight-bearing tissue of rat during repetitive loading. Firstly, the tissue condition under histopathology (for validating the effect of different recovery time), RH response trend and US feature changes were

investigated with different recovery time that simulate sufficient and insufficient recovery time during repetitive loading. Secondly, the tissue condition with RH response trends and US feature changes were examined using histopathology study.

The objectives of this study are listed below.

1. To establish a relationship between tissue recovery time and tissue condition examined under histopathology.
2. To establish a relationship between tissue recovery time and RH trend.
3. To establish a relationship between tissue recovery time and US feature changes.
4. To establish a relationship between RH trends and tissue condition examined under histopathology.
5. To establish a relationship between US features and tissue condition examined under histopathology.

1.4 Project scope

In this project a pressure control system that applies repetitive loading on the skin of anaesthetised rats is developed. The repetitive loading is able to restrict and restore blood perfusion at underlying skin to induce ischemic-reperfusion event that simulates the loading and recovery condition on the skin of bedridden patients with interval repositioning. This prospective study employed 21 rats (seven rats per group); as such, the findings are derived from a limited population size to explore the effects of recovery time on tissue condition, RH trend, and US features during repetitive loading. Nevertheless, the number of samples in this study ($n = 21$ rats divided into three groups) is bigger than previous related PU studies (Anke Stekelenburg et al., 2007: 10 Brown-Norway rats divided into three groups; Herrman *et al.*, 1999: 7 fuzzy rats; Sari *et al.*, 2008: 8 rats divided into three groups). Inducing pressure on human subject with unknown effect and histopathology examination could lead to tissue damage and will violate ethics consideration. This strengthen the reason of using animal model to test the hypothesis stated in this study. Once the hypothesis is proven, transitioning to human study is possible to validate the findings in this study on human subject.

Three recovery times (prolonged-40 min, moderate-10 min, and short-3 min) during repetitive loading was applied on the skin of 21 male rats to simulate sufficient and insufficient recovery time for the weight-bearing tissues in reversing the damages induced by repetitive loading. The scope of this implementation is related to preventative interventions for early PU from the aspect of pressure relief during recovery time, with the use of RH trends (peak RH) and US featured to detect changes (presence of oedema, distorted layered structure, and increased dermal and hypodermal thickness) measured using laser Doppler perfusion and ultrasound, respectively. The project investigates the effects of different recovery time on tissue condition, RH trends, and US feature changes. The investigations were performed to

determine the relationship of different recovery time in term of tissue condition and responses. Besides, the potential features used for estimating optimal tissue recovery time non-invasively can be determined.

Furthermore, the potential features were also related with tissue condition examined using histopathology study to determine the use of the features in determining tissue condition non-invasively.

1.5 Research significance

This study enhanced the understanding of PU by establishing the relationship between weight-bearing tissue and different recovery time during repetitive loading. Findings from this study supports the use of recovery time in preventing weight-bearing tissue approaches to early stage of PU. In addition, this study investigates the prospect of using RH response trends and/or US features for determining condition of weight-bearing tissues of an individual during repetitive loading. The RH response trend and/or US features that are associated with the tissue condition examined under histopathology should be further included into clinical protocol as a diagnostic strategy in determining tissue condition non-invasively. Furthermore, this study may contribute towards the development of PU preventive strategy in the aspect of achieving optimal recovery time during repetitive loading. Findings from this study may assist in future design of clinical protocol, e.g. achieve optimal body repositioning time interval or optimal time of alternating pressure on air mattress, using the features (RH trend and/or US features) to prevent the development of PU.

1.6 Thesis outline

This thesis is divided into five chapters consisting of thorough details of the study.

Chapter 1 provides the research background, existing problems to carry out this thesis, aims and objectives to accomplish this work, project scopes and research significance.

Chapter 2 covers a comprehensive review on PU, and the detection and prevention methods of PU development. In recent research, the RH and US features were introduced to have the benefits go further in preventing PU by determining the optimal recovery time and tissue condition of an individual.

Chapter 3 describes the methodology of the research. In this chapter the number and type of subject, measurement equipment, measurement considerations, and data collection protocol are detailed.

Chapter 4 shows the results of RH trend, US feature, and histopathology examination that induced by the three different recovery times. The RH trend and US feature are compared with the tissue condition based histopathology examination to determine the potential use of these features in determining weight-bearing tissues condition.

Chapter 5 discusses the results. The potential and limitation of the features in responding recovery time and determining tissue condition were presented to determine their potential use in preventing PU.

Chapter 6 presents the major findings of the study. Future work and recommendations are also highlighted in this chapter.



REFERENCES

- Adyanthaya, S., and Jose, M. (2013). Quality and safety aspects in histopathology laboratory. *Journal of oral and maxillofacial pathology: JOMFP*, 17(3): 402.
- Agrawal, P., Chourasia, V., Kapoor, R., and Agrawal, S. (2014). A Comprehensive Study of the Image Enhancement Techniques. *International Journal of Advance Foundation and Research in Computer (IJAFRC)*, 1: 85-89.
- Alberts, B., Johnson, A., Lewis, J., Morgan, D., Raff, M., Roberts, K., and Walter, P. (2014). *Molecular Biology of the Cell, Sixth Edition*: Taylor & Francis Group.
- Alberts, B., Johnson, A., Lewis, J., Walter, P., Raff, M., and Roberts, K. (2002). *Molecular Biology of the Cell 4th Edition*: International Student Edition: Routledge.
- Aljanabi, M. A., Hussain, Z. M., and Lu, S. F. (2018). An Entropy-Histogram Approach for Image Similarity and Face Recognition. *Mathematical Problems in Engineering*, 2018.
- Allen, J., and Howell, K. (2014). Microvascular imaging: techniques and opportunities for clinical physiological measurements. *Physiological measurement*, 35(7): R91.
- Amlung, S. R., Miller, W. L., and Bosley, L. M. (2001). The 1999 national pressure ulcer prevalence survey: a benchmarking approach. *Advances in skin & wound care*, 14(6): 297-301.
- Andersen, E. S., and Karlsmark, T. (2008). Evaluation of four non-invasive methods for examination and characterization of pressure ulcers. *Skin Research and Technology*, 14(3): 270-276.
- Aoi, N., Yoshimura, K., Kadono, T., Nakagami, G., Iizuka, S., Higashino, T., . . . Sanada, H. (2009). Ultrasound assessment of deep tissue injury in pressure ulcers: possible prediction of pressure ulcer progression. *Plastic and reconstructive surgery*, 124(2): 540-550.
- Assous, S., Humeau, A., Tartas, M., Abraham, P., and L'Huillier, J.-P. (2006). S-transform applied to laser Doppler flowmetry reactive hyperemia signals. *IEEE Transactions on Biomedical Engineering*, 53(6): 1032-1037.
- Aström, K. J., and Murray, R. M. (2010). *Feedback systems: an introduction for scientists and engineers*: Princeton university press.
- Auer, S. K., Salin, K., Anderson, G. J., and Metcalfe, N. B. (2018). Individuals exhibit consistent differences in their metabolic rates across changing thermal conditions. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 217: 1-6.
- Bader, D. (1990). The recovery characteristics of soft tissues following repeated loading. *Journal of rehabilitation research and development*, 27(2): 141.

- Bader, D., Barnhill, R., and Ryan, T. (1986). Effect of externally applied skin surface forces on tissue vasculature. *Archives of physical medicine and rehabilitation*, 67(11): 807-811.
- Baillie, L. (2014). *Developing Practical Nursing Skills, Fourth Edition*: Taylor & Francis.
- Baird, M. S. (2015). *Manual of Critical Care Nursing-E-Book: Nursing Interventions and Collaborative Management*: Elsevier Health Sciences.
- Bajrovic, F., Čenčur, M., Hožič, M., Ribarič, S., and Stefanovska, A. (2000). The contribution of lumbar sympathetic neurones activity to rat's skin blood flow oscillations. *Pflügers Archiv*, 439(1): r158-r160.
- Ballabeni, A., Apollonio, F., Gaiani, M., and Remondino, F. (2015). ADVANCES IN IMAGE PRE-PROCESSING TO IMPROVE AUTOMATED 3D RECONSTRUCTION. *International Archives of the Photogrammetry, Remote Sensing & Spatial Information Sciences*.
- Ballard, G., and Association, W. D. K. (1993). PID versus on-off control, why use a valve positioner?
- Barbenel, J., Jordan, M., Nicol, S., and Clark, M. (1977). Incidence of pressure-sores in the Greater Glasgow Health Board area. *The Lancet*, 310(8037): 548-550.
- Barnett, R. I., and Ablarde, J. A. (1995). Skin vascular reaction to short durations of normal seating. *Archives of physical medicine and rehabilitation*, 76(6): 533-540.
- Bauer, K., Rock, K., Nazzal, M., Jones, O., and Qu, W. (2016). Pressure Ulcers in the United States' Inpatient Population From 2008 to 2012: Results of a Retrospective Nationwide Study. *Ostomy/wound management*, 62(11): 30-38.
- Benes, R., and Riha, K. (2011). Noise reduction in medical ultrasound images. *Electron Revenue*, 2: 1-8.
- Bergstrand, S. (2014). *Preventing pressure ulcers by assessment of the microcirculation in tissue exposed to pressure*. Linköping University Electronic Press.
- Bergstrom, N. (2005). Patients at risk for pressure ulcers and evidence-based care for pressure ulcer prevention (pp. 35-50): Springer.
- Bergstrom, N., Allman, R., Carlson, C., Eaglestein, W., Frantz, R., and Garber, S. (1992). Panel for the Prediction and Prevention of Pressure Ulcers in Adults. Pressure ulcers in adults: Prediction and prevention. Clinical Practice Guideline, No. 3, AHCPR Publication No. 92-0047. Rockville, MD: Agency for Health Care Policy and Research. *Public Health Service, US Department of Health and Human Services*.
- Bhattacharya, S., and Mishra, R. (2015). Pressure ulcers: current understanding and newer modalities of treatment. *Indian Journal of Plastic Surgery: Official Publication of the Association of Plastic Surgeons of India*, 48(1): 4.

- Bliss, M. R. Hyperaemia. *Journal of tissue viability*, 8(4): 4-13. doi: 10.1016/S0965-206X(98)80028-4
- Bo, M., Massaia, M., Raspo, S., Bosco, F., Cena, P., Molaschi, M., and Fabris, F. (2003). Predictive Factors of In-Hospital Mortality in Older Patients Admitted to a Medical Intensive Care Unit. *Journal of the American Geriatrics Society*, 51(4): 529-533.
- Boussouar, A., Meziane, F., and Crofts, G. (2017). Plantar fascia segmentation and thickness estimation in ultrasound images. *Computerized Medical Imaging and Graphics*, 56: 60-73.
- Bouten, C., Oomens, C., Colin, D., and Bader, D. (2005). The aetiopathology of pressure ulcers: a hierarchical approach *Pressure ulcer research* (pp. 1-9): Springer.
- Bouten, C. V., Oomens, C. W., Baaijens, F. P., and Bader, D. L. (2003). The etiology of pressure ulcers: skin deep or muscle bound? *Archives of physical medicine and rehabilitation*, 84(4): 616-619.
- Bower, K. M. (2003). *When to use Fisher's exact test*. Paper presented at the American Society for Quality, Six Sigma Forum Magazine.
- Boyko, T. V., Longaker, M. T., and Yang, G. P. (2018). Review of the current management of pressure ulcers. *Advances in wound care*, 7(2): 57-67.
- Braden, B., and Bergstrom, N. (2000). A conceptual schema for the study of the etiology of pressure sores. *Rehabilitation Nursing*, 25(3): 105-110.
- Brienza, D., Deppisch, M., Gillespie, C., Goldberg, M., Gruccio, P., Jordan, R., . . . Sylvia, C. (2015). Do Lift Slings Significantly Change the Efficacy of Therapeutic Support Surfaces? A National Pressure Ulcer Advisory Panel White Paper.
- Brienza, D. M., Karg, P. E., Geyer, M. J., Kelsey, S., and Treffler, E. (2001). The relationship between pressure ulcer incidence and buttock-seat cushion interface pressure in at-risk elderly wheelchair users. *Archives of physical medicine and rehabilitation*, 82(4): 529-533.
- BRITO, M. V. H., Yasojima, E. Y., Machado, A. A., SILVEIRA, M. P. P. R., TEIXEIRA, R. K. C., Yamaki, V. N., and COSTA, F. L. d. S. (2017). DIFFERENT PROTOCOLS OF POSTCONDITIONING DOES NOT ATTENUATE MESENTERIC ISCHEMIA-REPERFUSION INJURY AFTER SHORT-TERM REPERFUSION. *ABCD. Arquivos Brasileiros de Cirurgia Digestiva (São Paulo)*, 30(1): 27-29.
- Capp, C., Dorwart, W., Elias, N., Hillman, S., Lancaster, S., Nair, R., . . . Smith, D. (2004). Post pressure hyperemia in the rat. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 137(3): 533-546.
- Carolan-Rees, G., Tweddel, A., Naka, K., and Griffith, T. (2002). Fractal dimensions of laser Doppler flowmetry time series. *Medical engineering & physics*, 24(1): 71-76.

- Ceelen, K. (2003). Aetiology of pressure ulcers A literature review. *Part I of MSc-thesis, Eindhoven university of Technology, Faculty of Biomedical Engineering*.
- Ceelen, K., Stekelenburg, A., Loerakker, S., Strijkers, G., Bader, D., Nicolay, K., . . . Oomens, C. (2008). Compression-induced damage and internal tissue strains are related. *Journal of biomechanics*, 41(16): 3399-3404.
- Chabner, D.-E. (2013). *The Language of Medicine-E-Book*: Elsevier Health Sciences.
- Chang, Y.-C., Chang, C.-M., Lai, L.-C., and Chen, L.-H. (2014). Contrast enhancement with considering visual effects based on gray-level grouping. *Journal of Marine Science and Technology*.
- Charan, J., and Biswas, T. (2013). How to calculate sample size for different study designs in medical research? *Indian journal of psychological medicine*, 35(2): 121.
- Charan, J., and Kantharia, N. (2013). How to calculate sample size in animal studies? *Journal of Pharmacology and Pharmacotherapeutics*, 4(4): 303.
- Chen, Y., DeVivo, M. J., and Jackson, A. B. (2005). Pressure ulcer prevalence in people with spinal cord injury: age-period-duration effects. *Archives of physical medicine and rehabilitation*, 86(6): 1208-1213.
- Chetter, I., Oswald, A., Fletcher, M., Dumville, J., and Cullum, N. (2016). A survey of patients with surgical wounds healing by secondary intention; an assessment of prevalence, aetiology, duration and management. *Journal of tissue viability*.
- Clark, M., Semple, M. J., Ivins, N., Mahoney, K., and Harding, K. (2017). National audit of pressure ulcers and incontinence-associated dermatitis in hospitals across Wales: a cross-sectional study. *BMJ open*, 7(8): e015616.
- Clough, N. (1994). The cost of pressure area management in an intensive care unit. *Journal of Wound Care*, 3(1): 33-35.
- Coleman, S., Nixon, J., Keen, J., Wilson, L., McGinnis, E., Dealey, C., . . . Schols, J. M. (2014). A new pressure ulcer conceptual framework. *Journal of advanced nursing*, 70(10): 2222-2234.
- Cortex, T. (*DermaLab® Series SkinLab Combo Instruction Manual* (pp. 41). Retrieved from <https://www.deernz.org/sites/dinz/files/Instruction%20Manual%20SkinLab%20Z5010108%20UK.pdf>
- Cracowski, J.-L., Minson, C. T., Salvat-Melis, M., and Halliwill, J. R. (2006). Methodological issues in the assessment of skin microvascular endothelial function in humans. *Trends in pharmacological sciences*, 27(9): 503-508.
- Cui, F.-F., Pan, Y.-Y., Xie, H.-H., Wang, X.-H., Shi, H.-X., Xiao, J., . . . Jiang, L.-P. (2016). Pressure combined with ischemia/reperfusion injury induces deep tissue injury via endoplasmic reticulum stress in a rat pressure ulcer model. *International journal of molecular sciences*, 17(3): 284.

- Dahiru, T. (2008). P-value, a true test of statistical significance? A cautionary note. *Annals of Ibadan postgraduate medicine*, 6(1): 21-26.
- Daniel, R. K., Priest, D. L., and Wheatley, D. C. (1981). Etiologic factors in pressure sores: an experimental model. *Archives of physical medicine and rehabilitation*, 62(10): 492-498.
- Davis, K. G., and Kotowski, S. E. (2015). Role of Bed Design and Head-of-Bed Articulation on Patient Migration. *Journal of nursing care quality*, 30(3): E1.
- Dealey, C., Posnett, J., and Walker, A. (2012). The cost of pressure ulcers in the United Kingdom. *Journal of Wound Care*, 21(6).
- Deepa, S., and Krishnaveni, K. (2018). A Review on Image Enhancement Techniques. *International Journal of Engineering Research in Computer Science and Engineering*, 5(3): 295-299.
- Defloor, T., De Bacquer, D., and Grypdonck, M. H. (2005). The effect of various combinations of turning and pressure reducing devices on the incidence of pressure ulcers. *International journal of nursing studies*, 42(1): 37-46.
- Defloor, T., and Grypdonck, M. H. (1999). Sitting posture and prevention of pressure ulcers. *Applied Nursing Research*, 12(3): 136-142.
- Defloor, T., Vanderwee, K., Wilborn, D., and Dassen, T. (2006). Pressure ulcer prevention and repositioning *Science and Practice of Pressure Ulcer Management* (pp. 67-73): Springer.
- Dell, R. B., Holleran, S., and Ramakrishnan, R. (2002). Sample size determination. *Ilar Journal*, 43(4): 207-213.
- du Prel, J.-B., Röhrig, B., Hommel, G., and Blettner, M. (2010). Choosing statistical tests: part 12 of a series on evaluation of scientific publications. *Deutsches Ärzteblatt International*, 107(19): 343.
- Easwaran, S. (2009). *Simplified Teaching and Understanding of Histogram Equalization in Digital Image Processing*. Paper presented at the American Society for Engineering Education.
- Emam, K. E. (1999). Benchmarking Kappa: Interrater agreement in software process assessments. *Empirical Software Engineering*, 4(2): 113-133.
- Ertugrul, N. (2000). Towards virtual laboratories: A survey of LabVIEW-based teaching/learning tools and future trends. *International Journal of Engineering Education*, 16(3): 171-180.
- Exton-Smith, A., and Sherwin, R. (1961). The prevention of pressure sores significance of spontaneous bodily movements. *The Lancet*, 278(7212): 1124-1126.
- Fam, R., and Lepage, Y. (2018). Validating analogically generated Indonesian words using Fisher's exact test.

- Festing, M. F., and Altman, D. G. (2002). Guidelines for the design and statistical analysis of experiments using laboratory animals. *ILAR journal*, 43(4): 244-258.
- Figueiredo Filho, D. B., Paranhos, R., Rocha, E. C. d., Batista, M., Silva Jr, J. A. d., Santos, M. L. W. D., and Marino, J. G. (2013). When is statistical significance not significant? *Brazilian Political Science Review*, 7(1): 31-55.
- Fragala, M., and Pontani-Bailey, L. (2005). Proper positioning of clients: A risk for caregivers. *Workplace Health & Safety*, 53(10): 438.
- Fredriksson, I., Fors, C., and Johansson, J. (2007). Laser Doppler flowmetry-a theoretical framework. *Department of Biomedical Engineering, Linköping University*: 6-7.
- Freeman, J. V., and Campbell, M. J. THE ANALYSIS OF CATEGORICAL DATA: FISHER'S EXACT TEST.
- Fundukian, L. J. (2011). *The Gale Encyclopedia of Medicine, ; 6 Volume Set*: Gale.
- Gai, S., Zhang, B., Yang, C., and Yu, L. (2018). Speckle noise reduction in medical ultrasound image using monogenic wavelet and Laplace mixture distribution. *Digital Signal Processing*, 72: 192-207.
- Garber, S. L., and Rintala, D. H. (2003). Pressure ulcers in veterans with spinal cord injury: a retrospective study. *Journal of rehabilitation research and development*, 40(5): 433.
- Gawlitta, D., Oomens, C. W., Bader, D. L., Baaijens, F. P., and Bouten, C. V. (2007). Temporal differences in the influence of ischemic factors and deformation on the metabolism of engineered skeletal muscle. *Journal of Applied Physiology*, 103(2): 464-473.
- Gebhardt, K., and Bliss, M. R. (1994). Preventing pressure sores in orthopaedic patients-is prolonged chair nursing detrimental? *Journal of tissue viability*, 4(2): 51-54.
- Gedamu, H., Hailu, M., and Amano, A. (2014). Prevalence and associated factors of pressure ulcer among hospitalized patients at Felegehiwot referral hospital, Bahir Dar, Ethiopia. *Advances in Nursing*, 2014.
- Gefen, A. (2009). Reswick and Rogers pressure-time curve for pressure ulcer risk. Part 2. *Nursing Standard*, 23(46): 40-44.
- Gefen, A., Farid, K. J., and Shaywitz, I. (2013). A review of deep tissue injury development, detection, and prevention: shear savvy. *Ostomy/wound management*, 59(2): 26-35.
- Gefen, A., van Nierop, B., Bader, D. L., and Oomens, C. W. (2008). Strain-time cell-death threshold for skeletal muscle in a tissue-engineered model system for deep tissue injury. *Journal of biomechanics*, 41(9): 2003-2012.

- Ghazanfari, M., Vogt, L., Banzer, W., and Rhodius, U. (2002). Reproducibility of non-invasive blood flow measurements using laser Doppler spectroscopy. *Physikalische Medizin Rehabilitationsmedizin Kurortmedizin*, 12(6): 330-336.
- Ghosh, S., Derle, A., Ahire, M., More, P., Jagtap, S., Phadatare, S. D., . . . Shinde, V. S. (2013). Phytochemical analysis and free radical scavenging activity of medicinal plants *Gnidia glauca* and *Dioscorea bulbifera*. *PloS one*, 8(12): e82529.
- Gidlöf, A., Lewis, D., and Hammersen, F. (1988). The effect of prolonged total ischemia on the ultrastructure of human skeletal muscle capillaries. A morphometric analysis. *International journal of microcirculation, clinical and experimental/sponsored by the European Society for Microcirculation*, 7(1): 67-86.
- Gill, E. C. (2015). Reducing hospital acquired pressure ulcers in intensive care. *BMJ Open Quality*, 4(1): u205599. w203015.
- Gräbner, D., Tintelott, M., Dumstorff, G., and Lang, W. (2017). *Low-Cost Thin and Flexible Screen-Printed Pressure Sensor*. Paper presented at the Multidisciplinary Digital Publishing Institute Proceedings.
- Grap, M. J., Burk, R. S., Lucas, V., Munro, C. L., Wetzel, P. A., and Schubert, C. M. (2015). Use of high frequency ultrasound to detect changes in skin integrity: An image evaluation validation procedure. *Intensive and Critical Care Nursing*, 31(3): 141-147.
- Gray, R. J., Voegeli, D., and Bader, D. L. (2016). Features of lymphatic dysfunction in compressed skin tissues—Implications in pressure ulcer aetiology. *Journal of tissue viability*, 25(1): 26-31.
- Guan, Y., Cai, B., Wu, X., Peng, S., Gan, L., Huang, D., . . . Liu, J. (2017). MicroRNA-352 regulates collateral vessel growth induced by elevated fluid shear stress in the rat hind limb. *Scientific reports*, 7(1): 6643.
- Guihan, M., Bates-Jenson, B. M., Chun, S., Parachuri, R., Chin, A. S., and McCreath, H. (2012). Assessing the feasibility of subepidermal moisture to predict erythema and stage 1 pressure ulcers in persons with spinal cord injury: a pilot study. *The journal of spinal cord medicine*, 35(1): 46-52.
- Gust, M. J., Hong, S. J., Fang, R. C., Lanier, S. T., Buck, D. W., Nuñez, J. M., . . . Mustoe, T. A. (2017). Adipose Tissue Drives Response to Ischemia-Reperfusion Injury in a Murine Pressure Sore Model. *Plastic and reconstructive surgery*, 139(5): 1128e-1138e.
- Gwet, K. L. (2014). *Handbook of inter-rater reliability: The definitive guide to measuring the extent of agreement among raters*: Advanced Analytics, LLC.
- Hagisawa, S., Ferguson-Pell, M., Cardi, M., and Miller, S. D. (1994). Assessment of skin blood content and oxygenation in spinal cord injured subjects during reactive hyperemia. *Journal of rehabilitation research and development*, 31: 1-1.

- Haisma, J. A., Van Der Woude, L. H., Stam, H. J., Bergen, M. P., Sluis, T. A., Post, M. W., and Bussmann, J. B. (2007). Complications following spinal cord injury: occurrence and risk factors in a longitudinal study during and after inpatient rehabilitation. *Journal of rehabilitation medicine*, 39(5): 393-398.
- Hallgren, K. A. (2012). Computing inter-rater reliability for observational data: an overview and tutorial. *Tutorials in quantitative methods for psychology*, 8(1): 23.
- Hargreaves, K. M., and Berman, L. H. (2015). *Cohen's Pathways of the Pulp*: Elsevier.
- Haugen, F. (2010). The Good Gain method for PI (D) controller tuning. *Tech Teach*: 1-7.
- Heerfordt Sjökvist, J. (2016). Enhanced Display of Mitoses in Hematoxylin-Eosin Digital Pathology Images.
- Helvig, E. I., and Nichols, L. W. (2012). Use of high-frequency ultrasound to detect heel pressure injury in elders. *Journal of Wound Ostomy & Continence Nursing*, 39(5): 500-508.
- Herrman, E. C., Knapp, C. F., Donofrio, J. C., and Salcido, R. (1999). Skin perfusion responses to surface pressure-induced ischemia: implication for the developing pressure ulcer. *Journal of rehabilitation research and development*, 36(2): 109.
- Hickson, M. (2006). Malnutrition and ageing. *Postgraduate Medical Journal*, 82(963): 2-8.
- Higashino, T., Nakagami, G., Kadono, T., Ogawa, Y., Iizaka, S., Koyanagi, H., . . . Sanada, H. (2014). Combination of thermographic and ultrasonographic assessments for early detection of deep tissue injury. *International wound journal*, 11(5): 509-516.
- Hirata, Y., Nagata, D., Suzuki, E., Nishimatsu, H., Suzuki, J.-i., and Nagai, R. (2010). Diagnosis and treatment of endothelial dysfunction in cardiovascular disease. *International heart journal*, 51(1): 1-6.
- Hölzle, F., Loeffelbein, D. J., Nolte, D., and Wolff, K.-D. (2006). Free flap monitoring using simultaneous non-invasive laser Doppler flowmetry and tissue spectrophotometry. *Journal of cranio-maxillofacial surgery*, 34(1): 25-33.
- Humeau, A., Koïtka, A., Abraham, P., Saumet, J.-L., and L'Huillier, J.-P. (2004). Time–frequency analysis of laser Doppler flowmetry signals recorded in response to a progressive pressure applied locally on anaesthetized healthy rats. *Physics in medicine and biology*, 49(5): 843.
- Jagatheeswari, P., Suresh Kumar, S., and Rajaram, M. (2009). Contrast enhancement for medical images based on histogram equalization followed by median filter.
- Jan, Y.-K., Jones, M. A., Rabadi, M. H., Foreman, R. D., and Thiessen, A. (2010). Effect of wheelchair tilt-in-space and recline angles on skin perfusion over the ischial tuberosity in people with spinal cord injury. *Archives of physical medicine and rehabilitation*, 91(11): 1758-1764.

- Jan, Y.-K., Lee, B., Liao, F., and Foreman, R. D. (2012). Local cooling reduces skin ischemia under surface pressure in rats: an assessment by wavelet analysis of laser Doppler blood flow oscillations. *Physiological measurement*, 33(10): 1733.
- Jan, Y.-K., Liao, F., Jones, M. A., Rice, L. A., and Tisdell, T. (2013). Effect of durations of wheelchair tilt-in-space and recline on skin perfusion over the ischial tuberosity in people with spinal cord injury. *Archives of physical medicine and rehabilitation*, 94(4): 667-672.
- Jan, Y.-K., Liao, F., Rice, L. A., and Woods, J. A. (2013). Using reactive hyperemia to assess the efficacy of local cooling on reducing sacral skin ischemia under surface pressure in people with spinal cord injury: a preliminary report. *Archives of physical medicine and rehabilitation*, 94(10): 1982-1989.
- Jasperse, J. L., Shoemaker, J. K., Gray, E. J., and Clifford, P. S. (2015). Positional differences in reactive hyperemia provide insight into initial phase of exercise hyperemia. *Journal of Applied Physiology*, 119(5): 569-575.
- Jiang, L.-P., Tu, Q., Wang, Y., and Zhang, E. (2011). Ischemia-reperfusion injury-induced histological changes affecting early stage pressure ulcer development in a rat model. *Ostomy/wound management*, 57(2): 55-60.
- Jyothi, G., Sushma, C., and Veeresh, D. (2015). Luminance based conversion of gray scale image to RGB image. *International Journal of Computer Science and Information Technology Research*, 3(3): 279-283.
- Källman, U. (2015). *Evaluation of Repositioning in Pressure Ulcer Prevention*. Linköping University Electronic Press.
- Kanan, C., and Cottrell, G. W. (2012). Color-to-grayscale: does the method matter in image recognition? *PloS one*, 7(1): e29740.
- Kanj, L. F., Wilking, S. V. B., and Phillips, T. J. (1998). Continuing medical education: Pressure ulcers. *Journal of the American Academy of Dermatology*, 38(4): 517-538.
- Kanno, N., Nakamura, T., Yamanaka, M., Kouda, K., and Tajima, F. (2009). Low-echoic lesions underneath the skin in subjects with spinal-cord injury. *Spinal Cord*, 47(3): 225-229.
- Kasuya, A., Sakabe, J.-i., and Tokura, Y. (2014). Potential application of in vivo imaging of impaired lymphatic duct to evaluate the severity of pressure ulcer in mouse model. *Scientific reports*, 4: 4173.
- Kaur, R., and Taqdir. (2016). Image Enhancement Techniques - A Review. *International Research Journal of Engineering and Technology*, 3(3): 1308-1315.
- Kaur, S., and Kaur, P. (2016). A Novel Brightness Preserving Histogram Equalization Technique for Image Contrast Enhancement. *International Journal of Science and Research*, 5(9): 1305-1309.

- Kim, H.-Y. (2014). Analysis of variance (ANOVA) comparing means of more than two groups. *Restorative dentistry & endodontics*, 39(1): 74-77.
- Kim, H.-Y. (2017). Statistical notes for clinical researchers: chi-squared test and Fisher's exact test. *Restorative dentistry & endodontics*, 42(2): 152-155.
- Kim, J. (2015). How to Choose the Level of Significance: A Pedagogical Note.
- Kim, S. J., Min, B. S., Lim, D. K., and Lee, J. H. (2013). *Determining parameters in contrast limited adaptive histogram equalization*. Paper presented at the Proceedings of the 7th International Conference on Information Security and Assurance.
- Kind, G. M., Buntic, R. F., Buncke, G. M., Cooper, T. M., Siko, P. P., and Buncke Jr, H. J. (1998). The effect of an implantable Doppler probe on the salvage of microvascular tissue transplants. *Plastic and reconstructive surgery*, 101(5): 1268-1273.
- Kitamura, A., Nakagami, G., Yoshida, M., Noguchi, H., Nishijima, Y., Minematsu, T., . . . Takahashi, K. (2014). Visualization of Tumor Necrosis Factor- α Distributions Within Pressure Ulcer Tissue Using the Wound Blotting Method: A Case Report and Discussion. *Wounds: a compendium of clinical research and practice*, 26(11): 323-329.
- Koivunen, M., Hjerpe, A., Luotola, E., Kauko, T., and Asikainen, P. (2018). Risks and prevalence of pressure ulcers among patients in an acute hospital in Finland. *Journal of Wound Care*, 27(Sup2): S4-S10.
- Kokate, J. Y., Leland, K. J., Held, A. M., Hansen, G. L., Kveen, G. L., Johnson, B. A., . . . Iaizzo, P. A. (1995). Temperature-modulated pressure ulcers: a porcine model. *Archives of physical medicine and rehabilitation*, 76(7): 666-673.
- Köksoy, C., Akkoca, M., Tokgöz, S., Çetinkaya, A., Sevim, Y., and Demirel-Yılmaz, E. (2018). Venous stent placement ameliorates cutaneous microvascular function in ilio caval venous obstruction. *Journal of Vascular Surgery: Venous and Lymphatic Disorders*, 6(1): 57-65.
- Kolaczowska, E., and Kubes, P. (2013). Neutrophil recruitment and function in health and inflammation. *Nature Reviews Immunology*, 13(3): 159-175.
- Koller, A., and Bagi, Z. (2002). On the role of mechanosensitive mechanisms eliciting reactive hyperemia. *American Journal of Physiology-Heart and Circulatory Physiology*, 283(6): H2250-H2259.
- Kosiak, M. (1961). Etiology of decubitus ulcers. *Arch Phys Med Rehabil*, 42: 19-29.
- Krause, J. S., and Broderick, L. (2004). Patterns of recurrent pressure ulcers after spinal cord injury: identification of risk and protective factors 5 or more years after onset. *Archives of physical medicine and rehabilitation*, 85(8): 1257-1264.
- Kravtcova, A. (2017). Blood vessel detection in medical procedures using laser Doppler flowmetry.

- Kumar, M., and Rana, A. (2016). Image Enhancement using Contrast Limited Adaptive Histogram Equalization and Wiener filter. *International Journal Of Engineering And Computer Science*, 5(6).
- Kumar, S., and Verma, P. (2012). Comparison of Different Enhanced Image Denoising with Multiple Histogram Techniques. *International Journal of Soft Computing and Engineering (IJSCE)* ISSN: 2231-2307.
- Kuo, P. J., Chew, K. Y., Kuo, Y. R., and Lin, P. Y. (2014). Comparison of outcomes of pressure sore reconstructions among perforator flaps, perforator-based rotation fasciocutaneous flaps, and musculocutaneous flaps. *Microsurgery*, 34(7): 547-553.
- Kwong, E. W., Hung, M. S., and Woo, K. (2016). Improvement of pressure ulcer prevention care in private for-profit residential care homes: an action research study. *BMC geriatrics*, 16(1): 192.
- Lachenbruch, C., Tzen, Y.-T., Brienza, D., Karg, P. E., and Lachenbruch, P. A. (2015). Relative contributions of interface pressure, shear stress, and temperature on ischemic-induced, skin-reactive hyperemia in healthy volunteers: a repeated measures laboratory study. *Ostomy/wound management*, 61(2): 16-25.
- Lahmann, N. A., and Kottner, J. (2011). Relation between pressure, friction and pressure ulcer categories: a secondary data analysis of hospital patients using CHAID methods. *International journal of nursing studies*, 48(12): 1487-1494.
- Landis, J. R., and Koch, G. G. (1977). The measurement of observer agreement for categorical data. *biometrics*: 159-174.
- Länne Rosenlund, H. (2016). Redesign of a generic human limb pressure device—towards early diagnosis of pressure ulcer risk patients.
- Latimer, S., Chaboyer, W., and Gillespie, B. M. (2015). The repositioning of hospitalized patients with reduced mobility: a prospective study. *Nursing open*, 2(2): 85-93.
- Laude, K., Thuillez, C., and Richard, V. (2001). Coronary endothelial dysfunction after ischemia and reperfusion: a new therapeutic target? *Brazilian Journal of Medical and Biological Research*, 34(1): 1-7.
- Lavana, K. K., and Kumar, R. (2011). A comparative study of image enhancement using histogram approach. *Int. J. Comput. Appl*, 32(5): 1-6.
- Lehoux, S., Castier, Y., and Tedgui, A. (2006). Molecular mechanisms of the vascular responses to haemodynamic forces. *Journal of internal medicine*, 259(4): 381-392.
- Lehoux, S., and Lévy, B. I. (2006). Collateral Artery Growth Making the Most of What You Have. *Circulation research*, 99(6): 567-569.
- Li, G., Kiel, J. W., Cardenas, D. P., Bryan, H., and Duong, T. Q. (2013). Postocclusive reactive hyperemia occurs in the rat retinal circulation but not in the choroid. *Investigative ophthalmology & visual science*, 54(7): 5123-5131.

- Liao, F., Burns, S., and Jan, Y.-K. (2013). Skin blood flow dynamics and its role in pressure ulcers. *Journal of tissue viability*, 22(2): 25-36.
- Liao, F., O'Brien Jr, W. D., and Jan, Y.-K. (2013). Assessing complexity of skin blood flow oscillations in response to locally applied heating and pressure in rats: Implications for pressure ulcer risk. *Physica A: Statistical Mechanics and its Applications*, 392(20): 4905-4915.
- Liberman, A. M. (2005). How much more likely? The implications of odds ratios for probabilities. *American Journal of Evaluation*, 26(2): 253-266.
- Lin, L., Chu, H.-C., and Lu, Y.-W. (1999). A simulation program for the sensitivity and linearity of piezoresistive pressure sensors. *J. Microelectromech. Syst*, 8(4): 514-522.
- Linder-Ganz, E., and Gefen, A. (2004). Mechanical compression-induced pressure sores in rat hindlimb: muscle stiffness, histology, and computational models. *Journal of Applied Physiology*, 96(6): 2034-2049.
- Lindgren, M., Unosson, M., Fredrikson, M., and Ek, A. C. (2004). Immobility—a major risk factor for development of pressure ulcers among adult hospitalized patients: a prospective study. *Scandinavian journal of caring sciences*, 18(1): 57-64.
- Lucas, V. S., Burk, R. S., Creehan, S., and Grap, M. J. (2014). Utility of high-frequency ultrasound: moving beyond the surface to detect changes in skin integrity. *Plastic surgical nursing: official journal of the American Society of Plastic and Reconstructive Surgical Nurses*, 34(1): 34.
- Lyder, C. H., and Ayello, E. A. (2008). Pressure ulcers: a patient safety issue.
- Ma, J., Fan, X., Yang, S. X., Zhang, X., and Zhu, X. (2017). Contrast Limited Adaptive Histogram Equalization Based Fusion for Underwater Image Enhancement: Preprints.
- Magudeeswaran, V., and Ravichandran, C. (2013). Fuzzy logic-based histogram equalization for image contrast enhancement. *Mathematical Problems in Engineering*, 2013.
- Maini, R., and Aggarwal, H. (2010). A comprehensive review of image enhancement techniques. *arXiv preprint arXiv:1003.4053*.
- Manikpuri, U., and Yadav, Y. (2014). Image Enhancement Through Logarithmic Transformation. *Image*, 1(8).
- Manorama, A. A., Baek, S., Vorro, J., Sikorskii, A., and Bush, T. R. (2010). Blood perfusion and transcutaneous oxygen level characterizations in human skin with changes in normal and shear loads—Implications for pressure ulcer formation. *Clinical biomechanics*, 25(8): 823-828.
- Martinelli, L., Hurschler, C., and Rosenbaum, D. (2006). Comparison of capacitive versus resistive joint contact stress sensors. *Clinical Orthopaedics and Related Research®*, 447: 214-220.

- Matsubara, J., Narumi, J., Nagasue, M., Sakamoto, S., Yuasa, K., and Shimizu, T. (1998). Postocclusive reactive hyperemia during vascular reconstruction. *International Journal of Angiology*, 7(3): 222-227.
- Matsunaga, T., Warltier, D. C., Weihrauch, D. W., Moniz, M., Tessmer, J., and Chilian, W. M. (2000). Ischemia-induced coronary collateral growth is dependent on vascular endothelial growth factor and nitric oxide. *Circulation*, 102(25): 3098-3103.
- Mattoon, J. S., and Nyland, T. G. (2014). *Small Animal Diagnostic Ultrasound - E-Book*: Elsevier Health Sciences.
- McCord, J. M. (1985). Oxygen-derived free radicals in postischemic tissue injury. *New England Journal of Medicine*, 312(3): 159-163.
- McGarr, G. W., Hodges, G. J., Mallette, M. M., and Cheung, S. S. (2018). Ischemia-reperfusion injury alters skin microvascular responses to local heating of the index finger. *Microvascular research*, 118: 12-19.
- McGuinness Abdollahi, Z. (2015). *Intra-operative optical monitoring of bowel tissue viability based on photoplethysmography and laser doppler flowmetry*. City University London.
- McHugh, M. L. (2012). Interrater reliability: the kappa statistic. *Biochemia medica*, 22(3): 276-282.
- McKinley, W. O., Jackson, A. B., Cardenas, D. D., and Michael, J. (1999). Long-term medical complications after traumatic spinal cord injury: a regional model systems analysis. *Archives of physical medicine and rehabilitation*, 80(11): 1402-1410.
- Menezes, I. A. C., Cunha, C. L. P. d., Carraro Júnior, H., and Luy, A. M. (2018). Perfusion index for assessing microvascular reactivity in septic shock after fluid resuscitation. *Revista Brasileira de terapia intensiva*, 30(2): 135-143.
- Min, B. S., Lim, D. K., Kim, S. J., and Lee, J. H. (2013). A novel method of determining parameters of CLAHE based on image entropy. *International Journal of Software Engineering and Its Applications*, 7(5): 113-120.
- Monfre, J. M. (2016). *Pressure ulcers Wound Healing-New insights into Ancient Challenges*: InTech.
- Musu, D., Rossi-Fedele, G., Campisi, G., and Cotti, E. (2016). Ultrasonography in the diagnosis of bone lesions of the jaws: a systematic review. *Oral surgery, oral medicine, oral pathology and oral radiology*, 122(1): e19-e29.
- Nagy, J. A., Benjamin, L., Zeng, H., Dvorak, A. M., and Dvorak, H. F. (2008). Vascular permeability, vascular hyperpermeability and angiogenesis. *Angiogenesis*, 11(2): 109-119.
- National_Instruments. *NI myRIO-1900 User Guide and Specifications*.

- New, P. W., Rawicki, H. B., and Bailey, M. J. (2004). Nontraumatic spinal cord injury rehabilitation: pressure ulcer patterns, prediction, and impact. *Archives of physical medicine and rehabilitation*, 85(1): 87-93.
- Newson, T., and Rolfe, P. (1982). Skin surface PO₂ and blood flow measurements over the ischial tuberosity. *Archives of physical medicine and rehabilitation*, 63(11): 553-556.
- Ng, L. G., Qin, J. S., Roediger, B., Wang, Y., Jain, R., Cavanagh, L. L., . . . Grimbaldston, M. A. (2011). Visualizing the neutrophil response to sterile tissue injury in mouse dermis reveals a three-phase cascade of events. *Journal of Investigative Dermatology*, 131(10): 2058-2068.
- Nicholson, P., Leeman, A., O'NEILL, C., Dobbs, S., Deshmukh, A., Denham, M., . . . Dobbs, R. (1988). Pressure sores: effect of Parkinson's disease and cognitive function on spontaneous movement in bed. *Age and ageing*, 17(2): 111-115.
- Niezgoda, J. A., and Mendez-Eastman, S. (2006). The effective management of pressure ulcers. *Advances in skin & wound care*, 19(1): 3-15.
- Nithya, K., Karthik, M. K., and Ramasamy, K. (2014). Comparison of Contrast Enhancement Technique with Partitioned Iterated Function System.
- Olsen Engineering, L. What are the benefits of Electric Linear Actuators compared with Pneumatic Systems? <http://www.consultolsen.com/downloads/benefits3.pdf>. Retrieved 28 August 2018.
- Olsson, R. A. (1975). Myocardial reactive hyperemia. *Circulation research*, 37(3): 263-270.
- Omair, A. (2014). Understanding the process of statistical methods for effective data analysis. *Journal of Health Specialties*, 2(3): 100.
- Oomens, C., Loerakker, S., and Bader, D. (2010). The importance of internal strain as opposed to interface pressure in the prevention of pressure related deep tissue injury. *Journal of tissue viability*, 19(2): 35-42.
- Oomens, C. W. J., Bressers, O., Bosboom, E., Bouten, C., and Bader, D. (2003). Can loaded interface characteristics influence strain distributions in muscle adjacent to bony prominences? *Computer methods in biomechanics and biomedical engineering*, 6(3): 171-180.
- Oremus, M., Oremus, C., Hall, G. B., McKinnon, M. C., ECT, and Team, C. S. R. (2012). Inter-rater and test-retest reliability of quality assessments by novice student raters using the Jadad and Newcastle–Ottawa Scales. *BMJ open*, 2(4): e001368.
- Osuala, E. O. (2014). Innovation in prevention and treatment of pressure ulcer: Nursing implication. *Tropical Journal of Medical Research*, 17(2): 61.

- Panel, N. P. U. A. (2014). European Pressure Ulcer Advisory Panel and Pan Pacific Pressure Injury Alliance. *Prevention and Treatment of Pressure Ulcers: Quick Reference Guide*.
- Parr, E., Ferdinand, P., and Roffe, C. (2017). Management of Acute Stroke in the Older Person. *Geriatrics*, 2(3): 27.
- Patel, B., and Patel, R. (2014). An analytical study on comparison of different image compression formats. *International Journal for Innovative Research in Science & Technology (IJRST)*, 1(7): 24-31.
- Patel, D. J., and Burton, A. C. (1956). Reactive hyperemia in the human finger. *Circulation research*, 4(6): 710-712.
- Patel, O., Maravi, Y. P., and Sharma, S. (2013). A comparative study of histogram equalization based image enhancement techniques for brightness preservation and contrast enhancement. *arXiv preprint arXiv:1311.4033*.
- Patel, S., Knapp, C. F., Donofrio, J. C., and Salcido, R. (1999). Temperature effects on surface pressure-induced changes in rat skin perfusion: implications in pressure ulcer development. *Journal of rehabilitation research and development*, 36(3): 189.
- Paul, D. (2014). Comparative Study of Different Image Enhancement Methods. *International Journal of Engineering Research & Technology*, 3(1): 1835-1847.
- Peirce, S. M., Skalak, T. C., and Rodeheaver, G. T. (2000). Ischemia-reperfusion injury in chronic pressure ulcer formation: A skin model in the rat. *Wound repair and regeneration*, 8(1): 68-76.
- Perimed_Instrument. Laser Doppler Probe. http://www.perimed-instruments.com/brochures/Laser_Doppler_Probes_Perimed.pdf. Retrieved 5 Dec 2013.
- Pickham, D., Ballew, B., Ebong, K., Shinn, J., Lough, M. E., and Mayer, B. (2016). Evaluating optimal patient-turning procedures for reducing hospital-acquired pressure ulcers (LS-HAPU): study protocol for a randomized controlled trial. *Trials*, 17(1): 190.
- Pizer, S. M., Johnston, R. E., Ericksen, J. P., Yankaskas, B. C., and Muller, K. E. (1990). *Contrast-limited adaptive histogram equalization: speed and effectiveness*. Paper presented at the Visualization in Biomedical Computing, 1990., Proceedings of the First Conference on.
- Pleiner, J., Schaller, G., Mittermayer, F., Marsik, C., MacAllister, R. J., Kapiotis, S., . . . Wolzt, M. (2008). Intra-arterial vitamin C prevents endothelial dysfunction caused by ischemia-reperfusion. *Atherosclerosis*, 197(1): 383-391.
- Ploquin, M., Basarab, A., and Kouamé, D. (2015). Resolution enhancement in medical ultrasound imaging. *Journal of Medical Imaging*, 2(1): 017001.

- Pompeii, L. A., Lipscomb, H. J., Schoenfisch, A. L., and Dement, J. M. (2009). Musculoskeletal injuries resulting from patient handling tasks among hospital workers. *American journal of industrial medicine*, 52(7): 571-578.
- Poredos, P., and Jezovnik, M. K. (2013). Testing endothelial function and its clinical relevance. *Journal of atherosclerosis and thrombosis*, 20(1): 1-8.
- Porter-Armstrong, A. P., Adams, C., Moorhead, A. S., Donnelly, J., Nixon, J., Bader, D. L., . . . Stinson, M. D. (2013). Do high frequency ultrasound images support clinical skin assessment? *ISRN nursing*, 2013.
- Porter-Armstrong, A. P., Moore, Z. E., Bradbury, I., and McDonough, S. (2018). Education of healthcare professionals for preventing pressure ulcers. *Cochrane Database of Systematic Reviews*(5).
- Pretto, E. (1991). *Reperfusion injury of the liver*. Paper presented at the Transplantation proceedings.
- Quintavalle, P. R., Lyder, C. H., Mertz, P. J., Phillips-Jones, C., and Dyson, M. (2006). Use of high-resolution, high-frequency diagnostic ultrasound to investigate the pathogenesis of pressure ulcer development. *Advances in skin & wound care*, 19(9): 498-505.
- Raghavan, P., Raza, W., Ahmed, Y., and Chamberlain, M. (2003). Prevalence of pressure sores in a community sample of spinal injury patients. *Clinical Rehabilitation*, 17(8): 879-884.
- Rahman, S. A., Soin, N., and Ibrahim, F. (2014). *Analysis of MEMS diaphragm of piezoresistive intracranial pressure sensor*. Paper presented at the Biomedical Engineering and Sciences (IECBES), 2014 IEEE Conference on.
- Rajput, S., and Suralkar, S. (2013). Comparative study of image enhancement techniques. *International Journal of Computer Science and Mobile Computing-A Monthly Journal of Computer Science and Information Technology*, 2(1): 11-21.
- Raju, D., Su, X., Patrician, P. A., Loan, L. A., and McCarthy, M. S. (2015). Exploring factors associated with pressure ulcers: A data mining approach. *International journal of nursing studies*, 52(1): 102-111.
- Rana, S., and Rana, S. (2015). A Review of Medical Image Enhancement Techniques for Image Processing. *International Journal of Current Engineering and Technology*, 5(2): 1282-1286.
- Reger, S. I., and Ranganathan, V. K. (2009). The importance of the microenvironment of support surfaces in the prevalence of pressure ulcers *Bioengineering research of chronic wounds* (pp. 85-100): Springer.
- Reswick, J., and Rogers, J. (1976). Experience at Rancho Los Amigos Hospital with devices and techniques to prevent pressure sores *Bed sore biomechanics* (pp. 301-310): Springer.

- Revie, W. C., and Green, P. (2016). Current problems and perspectives on colour in medical imaging. *Electronic Imaging*, 2016(20): 1-5.
- Rich, S. E., Margolis, D., Shardell, M., Hawkes, W. G., Miller, R. R., Amr, S., and Baumgarten, M. (2011). Frequent manual repositioning and incidence of pressure ulcers among bed-bound elderly hip fracture patients. *Wound repair and regeneration*, 19(1): 10-18.
- Robotics, T. "FlexiForce® Sensor". Retrieved 18 Nov, 2017, from www.trossenrobotics.com/flexiforce-1lb-resistive-force-sensor.aspx.
- Romanelli, M., Clark, M., Gefen, A., and Ciprandi, G. (2018). *Science and Practice of Pressure Ulcer Management*: Springer London.
- Ruch, R., Patton, H., Ruch, R., and Patton, H. (1965). Energy metabolism. *Physiology and biophysics*, 19th ed. Philadelphia, PA: Saunders Pr, 1064.
- Russo, C. A., and Elixhauser, A. (2006). Hospitalizations related to pressure sores, 2003.
- Russo, C. A., Steiner, C., and Spector, W. (2008). Hospitalizations related to pressure ulcers among adults 18 years and older, 2006.
- Ruxton, G. D., and Neuhäuser, M. (2013). Review of alternative approaches to calculation of a confidence interval for the odds ratio of a 2× 2 contingency table. *Methods in Ecology and Evolution*, 4(1): 9-13.
- Saba, L., Rozen, W. M., Alonso-Burgos, A., and Ribuffo, D. (2014). *Imaging for Plastic Surgery*: CRC Press.
- Salisbury, R. E. (1985). Transcutaneous PO₂ monitoring in bedridden burn patients: a physiological analysis of four methods to prevent pressure sores. *Chronic ulcers of the skin*. New York: McGraw-Hill: 189-195.
- Salzberg, A. C., Byrne, D. W., Cayten, G. C., Kabir, R., van Niewerburgh, P., Viehbeck, M., . . . Jones, E. C. (1998). Predicting and preventing pressure ulcers in adults with paralysis. *Advances in skin & wound care*, 11(5): 237-246.
- Sari, Y., Nakagami, G., Kinoshita, A., Huang, L., Ueda, K., Iizaka, S., . . . Sugama, J. (2008). Changes in serum and exudate creatine phosphokinase concentrations as an indicator of deep tissue injury: a pilot study. *International wound journal*, 5(5): 674-680.
- Schäfer, G., Dobos, G., Lünemann, L., Blume-Peytavi, U., Fischer, T., and Kottner, J. (2015). Using ultrasound elastography to monitor human soft tissue behaviour during prolonged loading: A clinical explorative study. *Journal of tissue viability*, 24(4): 165-172.
- Schaper, W., and Schaper, J. (2012). *Collateral Circulation: Heart, Brain, Kidney, Limbs*: Springer US.

- Schubert, V., and Fagrell, B. (1990). Postocclusive reactive hyperemia and thermal response in the skin microcirculation of subjects with spinal cord injury. *Scandinavian journal of rehabilitation medicine*, 23(1): 33-40.
- Seida, M., Wagner, H., Vass, K., and Klatzo, I. (1988). Effect of aminophylline on postischemic edema and brain damage in cats. *Stroke*, 19(10): 1275-1282.
- Shahbaz, R. (2015). Evaluation of Endothelial Dysfunction Using Laser Doppler Flowmetry in Diabetes Mellitus and Peripheral Arterial Occlusive Disease Patients. *Journal of Vascular Surgery*, 61(6): 92S. doi: 10.1016/j.jvs.2015.04.179
- Shahrokhi, M., and Zomorodi, A. (2013). Comparison of PID controller tuning methods. *Department of Chemical & Petroleum Engineering Sharif University of Technology*.
- Shaked, E., and Gefen, A. (2013). Modeling the effects of moisture-related skin-support friction on the risk for superficial pressure ulcers during patient repositioning in bed. *Frontiers in bioengineering and biotechnology*, 1: 9.
- Shweta, and Viswanath, K. (2017). A Review of Enhancement Techniques on Medical Images. *International Journal of Advanced Computing and Electronics Technology*, 4(3): 26-30.
- Sihvola, H., and Savolainen, J. (2016). Pressure ulcer prevention and early detection: an introductory guide in English.
- Silva, C. F. R., Santana, R. F., Oliveira, B. G. R. B., and Carmo, T. G. (2017). High prevalence of skin and wound care of hospitalized elderly in Brazil: a prospective observational study. *BMC research notes*, 10(1): 81.
- Singh, B. B., and Patel, S. (2017). Efficient Medical Image Enhancement using CLAHE Enhancement and Wavelet Fusion. *International Journal of Computer Applications*, 167(5).
- Singh, H., and Singh, A. P. V. (2017). A Comparative Analysis on Histogram Equalization Techniques for Medical Image Enhancement. *International Journals of Advanced Research in Computer Science and Software Engineering*, 7(6): 364-370.
- Singh, T. R., Singh, O. I., Singh, K. M., Sinam, T., and Singh, T. R. (2010). Image enhancement by adaptive power-law transformations. *Bahria University Journal of Information & Communication Technology*, 3(1): 29-37.
- Sonenblum, S. E., and Sprigle, S. H. (2011). The impact of tilting on blood flow and localized tissue loading. *Journal of tissue viability*, 20(1): 3-13.
- Sonker, D., and Parsai, M. (2013). Comparison of Histogram Equalization Techniques for Image Enhancement of Grayscale images of Dawn and Dusk. *International Journal of Modern Engineering Research (IJMER)*, 3(4).
- Soto, C. (2015). *ECG: Essentials of Electrocardiography*: Cengage Learning.
- Sperelakis, N. (2012). *Physiology and Pathophysiology of the Heart*: Springer US.

- Spriestersbach, A., Röhrig, B., du Prel, J.-B., Gerhold-Ay, A., and Blettner, M. (2009). Deskriptive Statistik. *Dtsch Arztebl Int*, 106(36): 578-583.
- Sprigle, S., Linden, M., and Riordan, B. (2002). Characterizing reactive hyperemia via tissue reflectance spectroscopy in response to an ischemic load across gender, age, skin pigmentation and diabetes. *Medical engineering & physics*, 24(10): 651-661.
- Sripriya, T., and Jeyalakshmi, V. (2008). Survey on Pressure Sensors in the Previous Decades.
- Stekelenburg, A., Oomens, C., Strijkers, G., Nicolay, K., and Bader, D. (2006). Compression-induced deep tissue injury examined with magnetic resonance imaging and histology. *Journal of Applied Physiology*, 100(6): 1946-1954.
- Stekelenburg, A., Strijkers, G. J., Parusel, H., Bader, D. L., Nicolay, K., and Oomens, C. W. (2007). Role of ischemia and deformation in the onset of compression-induced deep tissue injury: MRI-based studies in a rat model. *Journal of Applied Physiology*, 102(5): 2002-2011.
- Stephens, M., and Bartley, C. (2018). Understanding the association between pressure ulcers and sitting in adults what does it mean for me and my carers? Seating guidelines for people, carers and health & social care professionals. *Journal of tissue viability*, 27(1): 59-73.
- Stoller, M., and Seiler, C. (2014). Pathophysiology of coronary collaterals. *Current cardiology reviews*, 10(1): 38-56.
- Stotts, N. A., Deosaransingh, K., Roll, F. J., and Newman, J. (1998). Underutilization of pressure ulcer risk assessment in hip fracture patients. *Advances in skin & wound care*, 11(1): 32-39.
- Suresh, K., and Chandrashekara, S. (2012). Sample size estimation and power analysis for clinical research studies. *Journal of human reproductive sciences*, 5(1): 7.
- Szumilas, M. (2010). Explaining odds ratios. *Journal of the Canadian academy of child and adolescent psychiatry*, 19(3): 227.
- Tan, L. K. (2006). Image file formats. *Biomed Imaging Interv J*, 2(1): e6.
- Tayyib, N., Coyer, F., and Lewis, P. (2013). Pressure ulcers in the adult intensive care unit: a literature review of patient risk factors and risk assessment scales. *Journal of Nursing Education and Practice*, 3(11): p28.
- Tehrani, K. A., and Mpanda, A. (2012). PID Control Theory *Introduction to PID Controllers-Theory, Tuning and Application to Frontier Areas*: InTech.
- Teixeira, B. C., Lopes, A. L., Macedo, R. C. O., Correa, C. S., Ramis, T. R., Ribeiro, J. L., and Reischak-Oliveira, A. (2014). Inflammatory markers, endothelial function and cardiovascular risk. *Jornal Vascular Brasileiro*, 13(2): 108-115.
- Tekscan. *Flexiforce® Sensor User's Manual*: South Boston.

- Thamizharasi, A., and Jayasudha, J. (2016). AN ILLUMINATION INVARIANT FACE RECOGNITION BY ENHANCED CONTRAST LIMITED ADAPTIVE HISTOGRAM EQUALIZATION. *ICTACT Journal on Image & Video Processing*, 6(4).
- Thamman, P., and Verma, A. (2014). Contrast Enhancement of Medical Images-A Review. *International Journal of Innovations & Advancement in Computer Science*, 3(4): 124-128.
- Thomas, D. R. (2006). Prevention and treatment of pressure ulcers. *Journal of the American Medical Directors Association*, 7(1): 46-59.
- Thomas, D. R., Goode, P. S., Tarquine, P. H., and Allman, R. M. (1996). Hospital-Acquired Pressure Ulcers and Risk of Death. *Journal of the American Geriatrics Society*, 44(12): 1435-1440.
- Thorfinn, J., Sjöberg, F., and Lidman, D. (2007). Perfusion of buttock skin in healthy volunteers after long and short repetitive loading evaluated by laser Doppler perfusion imager. *Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery*, 41(6): 297-302.
- Thorfinn, J., Sjöberg, F., Sjöstrand, L., and Lidman, D. (2006). Perfusion of the skin of the buttocks in paraplegic and tetraplegic patients, and in healthy subjects after a short and long load. *Scandinavian journal of plastic and reconstructive surgery and hand surgery*, 40(3): 153-160.
- Togawa, T., Tamura, T., and Oberg, P. A. (1997). *Biomedical transducers and instruments*: CRC press.
- Tsai, D.-Y., Lee, Y., and Matsuyama, E. (2008). Information entropy measure for evaluation of image quality. *Journal of digital imaging*, 21(3): 338-347.
- Tsuji, S., Ichioka, S., Sekiya, N., and Nakatsuka, T. (2005). Analysis of ischemia-reperfusion injury in a microcirculatory model of pressure ulcers. *Wound repair and regeneration*, 13(2): 209-215.
- Turner, P. V., Brabb, T., Pekow, C., and Vasbinder, M. A. (2011). Administration of substances to laboratory animals: routes of administration and factors to consider. *Journal of the American Association for Laboratory Animal Science*, 50(5): 600-613.
- Tzen, Y.-T., Brienza, D. M., Karg, P., and Loughlin, P. (2010). Effects of local cooling on sacral skin perfusion response to pressure: implications for pressure ulcer prevention. *Journal of tissue viability*, 19(3): 86-97.
- Tzen, Y.-T., Brienza, D. M., Karg, P. E., and Loughlin, P. J. (2013). Effectiveness of local cooling for enhancing tissue ischemia tolerance in people with spinal cord injury. *The journal of spinal cord medicine*, 36(4): 357-364.
- Ünal, S., Özmen, S., Demİr, Y., Yavuzer, R., Latİfogu, O., and Atabay, K. (2001). The effect of gradually increased blood flow on ischemia-reperfusion injury. *Annals of plastic surgery*, 47(4): 412-416.

- Van der Wielen, H., Post, M., Lay, V., Gläsche, K., and Scheel-Sailer, A. (2016). Hospital-acquired pressure ulcers in spinal cord injured patients: time to occur, time until closure and risk factors. *Spinal Cord*, 54(9): 726.
- VanGilder, C., MacFarlane, G. D., and Meyer, S. (2008). Results of nine international pressure ulcer prevalence surveys: 1989 to 2005. *Ostomy/wound management*, 54(2): 40-54.
- Vidal, J., and Sarrias, M. (1991). An analysis of the diverse factors concerned with the development of pressure sores in spinal cord injured patients. *Paraplegia*, 29(4): 261-267.
- Wang, Q., Kong, L., Sprigle, S., and Hayward, V. (2006). *Portable gage for pressure ulcer detection*. Paper presented at the Engineering in Medicine and Biology Society, 2006. EMBS'06. 28th Annual International Conference of the IEEE.
- Weststrate, J. T. M. (2005). *The value of pressure ulcer risk assessment and interface pressure measurements in patients: A nursing perspective*.
- Whitty, J. A., McInnes, E., Bucknall, T., Webster, J., Gillespie, B. M., Banks, M., . . . Roberts, S. (2017). The cost-effectiveness of a patient centred pressure ulcer prevention care bundle: Findings from the INTACT cluster randomised trial. *International journal of nursing studies*, 75: 35-42.
- Wilson, D. (2010). *Clinical Veterinary Advisor-E-Book: The Horse*: Elsevier Health Sciences.
- Wong, H., Kaufman, J., Baylis, B., Conly, J. M., Hogan, D. B., Stelfox, H. T., . . . Ho, C. H. (2015). Efficacy of a pressure-sensing mattress cover system for reducing interface pressure: study protocol for a randomized controlled trial. *Trials*, 16(1): 434.
- Xiong, S., Goonetilleke, R. S., Witana, C. P., and Rodrigo, W. (2010). An indentation apparatus for evaluating discomfort and pain thresholds in conjunction with mechanical properties of foot tissue in vivo. *Journal of rehabilitation research and development*, 47(7): 629-641.
- Yamamoto, H., Tomoike, H., Shimokawa, H., Nabeyama, S., and Nakamura, M. (1984). Development of collateral function with repetitive coronary occlusion in a canine model reduces myocardial reactive hyperemia in the absence of significant coronary stenosis. *Circulation research*, 55(5): 623-632.
- Yap, T. L., Kennerly, S. M., Horn, S. D., Bergstrom, N., Datta, S., and Colon-Emeric, C. (2018). TEAM-UP for quality: a cluster randomized controlled trial protocol focused on preventing pressure ulcers through repositioning frequency and precipitating factors. *BMC geriatrics*, 18(1): 54.
- Yarkony, G. M. (1994). Pressure ulcers: a review. *Archives of physical medicine and rehabilitation*, 75(8):908-17.
- Youney, J. (2007). A comparison and evaluation of common pid tuning methods.

- Yu, X., Xu, J., Huang, G., Zhang, K., Qing, L., Liu, W., and Xu, W. (2017). Bubble-induced endothelial microparticles promote endothelial dysfunction. *PloS one*, 12(1): e0168881.
- Yuen, J. C., and Feng, Z. (2000). Monitoring free flaps using the laser Doppler flowmeter: five-year experience. *Plastic and reconstructive surgery*, 105(1): 55-61.
- Yvonne-Tee, G. B., Rasool, A. H. G., Halim, A. S., and Rahman, A. R. A. (2005). Reproducibility of different laser Doppler fluximetry parameters of postocclusive reactive hyperemia in human forearm skin. *Journal of pharmacological and toxicological methods*, 52(2): 286-292.
- Zhang, Y.-Z., Xu, T., Gong, H.-Y., Li, C.-Y., Ye, X.-H., Lin, H.-J., . . . Wu, X.-H. (2016). Application of high-resolution ultrasound, real-time elastography, and contrast-enhanced ultrasound in differentiating solid thyroid nodules. *Medicine*, 95(45).
- Zuo, X.-L., and Meng, F.-J. (2015). A care bundle for pressure ulcer treatment in intensive care units. *International Journal of Nursing Sciences*, 2(4): 340-347.