UNIVERSITI PUTRA MALAYSIA

IDENTIFICATION OF BASAL STEM ROT DISEASE IN OIL PALM TREE USING THERMAL IMAGING TECHNIQUE

GHAIBULNA BINTI ABDOL LAJIS

FK 2018 149

IDENTIFICATION OF BASAL STEM ROT DISEASE IN OIL PALM TREE USING THERMAL IMAGING TECHNIQUE

By

GHAIBULNA BINTI ABDOL LAJIS

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science

August 2018

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

IDENTIFICATION OF BASAL STEM ROT DISEASE IN OIL PALM TREE USING THERMAL IMAGING TECHNIQUE

By

GHAIBULNA BINTI ABDOL LAJIS

August 2018

Chairman : Siti Khairunniza Bejo, PhD
Faculty : Engineering

Basal stem rot (BSR), caused by *Ganoderma boninense* is known as one of the deadliest diseases in the oil palm plantations in Southeast Asia. *Ganoderma* could reduce the productivity of oil palm plantations and possibly reduce the market value of palm oil in Malaysia. The available technique of BSR detection is time-consuming and human dependence. This study focuses on detecting the oil palm tree infected by BSR using thermal imaging technique. In order to find a suitable time to capture the thermal images, thermal images of canopy and trunk sections of the oil palm trees from healthy and BSR-infected trees were captured in the morning (9 to 12 pm) and afternoon (12 to 3 pm) session. The images were pre-processed using FLIR QuickReport 1.2 (FLIR Systems, Inc., Oregon, United States). The images were then processed using MATLAB software (Version R2016b, The MathWorks Inc., Massachusetts, United States) to extract pixel value representing thermal properties of the trees. After that, statistical analysis was done using these pixel values. The result from T-test has shown that thermal images taken at canopy section during the afternoon session have a significant difference ($\alpha<0.05$) between healthy and BSR-infected trees. There were four features extracted from the images of canopy section namely minimum, maximum, mean and standard deviation value. Based on the statistical analysis, only mean of the pixel value gave a significant difference with a P value of 0.0052. For the maximum feature, all the data has the same value regardless of the healthiness condition, hence this feature will not be used for further analysis. Four different types of classifier namely, Linear Discriminant Analysis (LDA), Quadratic Discriminant Analysis (QDA), Support Vector Machine (SVM) and k-nearest neighbour (kNN) were used and compared. Input parameters were taken from different combinations of the features to classify the healthy and BSR-infected trees. The result showed that quadratic SVM with the input parameter

using a combination of minimum and mean gave the highest percentage of accuracy with 67.0%. In order to improve the accuracy, new indices called; healthy variance (Y_H) BSR-infected variance (Y_{UH}) and all variance (Y_{ALL}) were developed based on the squared value of the difference between the mean intensity value of an oil palm tree and the averaged mean intensity value of healthy, BSR-infected and all samples accordingly. However, it only gave the best accuracy at 62.3% from the combination of minimum, mean, standard deviation and Y_{UH} using linear SVM classifier. As a result, the Principal Component Analysis (PCA) was introduced to extract the most suitable features among six features available. The score plot of PC1 versus PC3 has shown that there were two distinguishable trendlines where the BSR-infected tree is located outside the trendline of the healthy trees. Values of PC1 and PC3 were later used for classification using all fourteen different types of classification model. Based on the results, the quadratic SVM model gave the best classification with the highest accuracy of 89.2% for the training set and 84.4% for the test set. Based on this study, it can be concluded that thermal imaging has the potential for BSR detection in oil palm trees.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

MENGENALPASTI PENYAKIT REPUT PANGKAL BATANG PADA POKOK KELAPA SAWIT MENGGUNAKAN TEKNIK PENGIMEJAN TERMA

Oleh

GHAIBULNA BINTI ABDOL LAJIS

Ogos 2018

Pengerusi : Siti Khairunniza Bejo, PhD
Fakulti : Kejuruteraan

Reput pangkal batang (RPB) adalah sejenis penyakit yang disebabkan oleh sejenis fungus, *Ganoderma boninense*. Ia dikenali sebagai penyakit paling berbahaya di ladang kelapa sawit sekitar Asia Tenggara. *Ganoderma* boleh mengurangkan kadar produktiviti ladang kelapa sawit dan juga berpotensi mengurangkan nilai pasaran minyak sawit di Malaysia. Teknik pengesanan RPB sedia ada memerlukan masa yang lama dan terlalu bergantung dengan tenaga kerja manusia. Kajian ini menumpukan pengenalpastian sifat terma pokok yang sihat dan yang dijangkiti penyakit RPB menggunakan teknik pengimejan terma. Bagi mendapatkan masa yang sesuai untuk mengambil imej terma, imej terma pada bahagian kanopi dan batang pokok kelapa sawit telah diambil daripada pokok yang sihat dan pokok yang dijangkiti RPB ketika sesi pagi (9 hingga 12 pm) dan sesi petang (12 hingga 3 pm). Imej terma tersebut telah dipra-proses menggunakan perisian FLIR QuickReport 1.2 (FLIR Systems, Inc., Oregon, United States) dan kemudiannya diproses menggunakan perisian MATLAB (Versi R2016b, The MathWorks Inc., Massachusetts, United States) untuk mengestrak nilai piksel yang mewakili sifat terma pokok. Selepas itu, analisis statistik dilakukan menggunakan nilai tersebut. Hasil daripada ujian-T menunjukkan imej terma yang diambil pada bahagian kanopi ketika sesi petang mempunyai perbezaan yang signifikan ($\alpha<0.05$) antara pokok yang sihat dan dijangkiti RPB. Terdapat empat sifat yang disari daripada imej pada bahagian kanopi iaitu nilai minimum, maksimum, purata dan sisihan piawai. Hasil daripada analisis statistik, hanya purata nilai piksel yang memberikan perbezaan signifikan dengan nilai P sebanyak 0.0052. Bagi nilai maksimum pula, kesemua data mempunyai nilai yang sama walaupun berbeza keadaan kesihatan. Oleh itu, sifat ini tidak akan digunakan untuk analisis seterusnya. Empat belas jenis model pengelasan yang berbeza iaitu Analisis Pembeza Layan Linear (LDA),

Analisis Pembeza Layan Kuadratik (QDA), Mesin Vektor Sokongan (SVM) dan Jiran Terhampir-k (kNN) digunakan dan dibandingkan. Parameter input daripada gabungan sifat yang berbeza telah digunakan untuk mengelaskan pokok yang sihat dan dijangkiti RPB. Keputusan menunjukkan SVM kuadratik dengan gabungan sifat minimum dan purata telah menghasilkan peratus ketepatan tertinggi dengan nilai 67.0%. Bagi mempertingkatkan ketepatan pengelasan, indeks baru yang dikenali sebagai variasi sihat (Y_H), variasi jangkitan RPB (Y_{UH}) dan variasi kesemuaya (Y_{ALL}) telah dihasilkan berdasarkan nilai kuasa dua perbezaan purata pokok dengan purata kesemua pokok yang dikira berdasarkan sampel pokok yang diambil daripada pokok yang sihat, dijangkiti RPB dan gabungan kedua-duanya. Walau bagaimanapun, ia hanya memberikan peratus ketepatan tertinggi pada 62.3% sahaja daripada gabungan sifat minimum, purata, sisihan piawai dan indeks Y_{UH} melalui pengelas SVM linear. Hasilnya, AKP telah digunakan untuk mengekstrak sifat yang paling sesuai di antara enam sifat yang ada. Plot skor PC1 versus PC3 telah menunjukkan bahawa ada dua garis arah aliran yang boleh dibezakan di mana pokok yang dijangkiti RPB terletak di luar garis arah aliran pokok yang sihat. Nilai PC1 dan PC3 kemudiannya digunakan untuk pengelasan menggunakan empat pengelas berbeza. Hasil keputusan menunjukkan pengelas SVM kuadratik memberi peratus ketepatan tertinggi dengan nilai 89.2% untuk set latihan dan 84.4% untuk set ujian. Berdasarkan kajian ini, dapat disimpulkan bahawa teknik pengimejan terma mempunyai potensi untuk mengenalpasti jangkitan RPB pada pokok kelapa sawit.

ACKNOWLEDGEMENTS

First and foremost, I'm grateful to Allah, the Almighty for providing me with this opportunity and for granting me the capability to proceed successfully.

I would like to express my sincere gratitude to my supervisor, Associate Professor Dr. Siti Khairunniza Bejo for her continuous guidance, advice, comments and encouragement. Her efforts ensured that this study was completed is highly appreciated. I would also like to thank my research supervisory committees: Associate Professor Dr. Samsuzana Abd Aziz, Associate Professor Dr. Tofael Ahamed and Dr. Hj Idris Abu Seman, for their encouragement and insightful comments. Special thanks are also due to staff members at Department of Biological and Agricultural Engineering, Faculty of Engineering, Universiti Putra Malaysia (UPM) and Ganoderma and Diseases Research for Oil Palm Unit, Malaysian Palm Oil Board (MPOB) for their kindness, help and technical support throughout the journey of this thesis. I also appreciate the support given by the member of Faculty of Life and Environmental Sciences, University of Tsukuba for their assistance in conducting the research.

I am grateful to my fellow friends in UPM: Rasydan, Zul, Nik and Dini, for the stimulating discussions, for the sleepless nights we were working together before deadlines, and for all the fun we have had in the last three years. Also, I thank my friends in University of Tsukuba: Nazia-san and others for always making me feel at home even when I was thousands of miles away.

My sincere gratitude to my family: my parents Abdol Lajis and Hamizah and my brother, Ghaibullah, for their patience and support throughout my life. Million thanks to my long-time friends, Asmi, Ariff, Am, Aziz, Ayun, Asmah and Shiro for always being there and believing in me.

I would also like to express my gratitude to the Ministry of Higher Education (MOHE) and Universiti Putra Malaysia (UPM) for sponsoring this research under Putra IPB Grant.

Finally, thank you to those who were directly or indirectly related to this study.

I certify that a Thesis Examination Committee has met on 27 August 2018 to conduct the final examination of Ghaibulna binti Abdol Lajis on her thesis entitled "Identification of Basal Stem Rot Disease in Oil Palm Tree Using Thermal Imaging Technique" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Hasfalina binti Che Man, PhD
Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Norhashila binti Hashim, PhD
Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Chong Khim Phin, PhD
Associate Professor
Universiti Malaysia Sabah
Malaysia
(External Examiner)

RUSLI HAJI ABDULLAH, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 27 September 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Siti Khairunniza Bejo, PhD
Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Samsuzana Abd Aziz, PhD
Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Hj Idris Abu Seman, PhD
Head
Ganoderma and Diseases Research for Oil Palm Unit
Malaysian Palm Oil Board
(Member)

Tofael Ahamed, PhD
Associate Professor
Faculty of Life and Environmental Sciences
University of Tsukuba
(Member)

ROBIAH BINTI YUNUS, PhD
Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by Graduate Student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Ghaibulna binti Abdol Lajis (GS43756)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature : _____
Name of Chairman of
Supervisory Committee : Associate Professor Dr. Siti Khairunniza Bejo

Signature : _____
Name of Member of
Supervisory Committee : Associate Professor Dr. Samsuzana Abd Aziz

Signature : _____
Name of Member of
Supervisory Committee : Dr. Hj Idris Abu Seman

Signature : _____
Name of Member of
Supervisory Committee : Associate Professor Dr. Tofael Ahamed

TABLE OF CONTENTS

LIST OF TABLES

LIST OF FIGURES

CHAPTER 1

INTRODUCTION

1.1 General Overview

Oil palm is one of the most important crops in Malaysia. In 2016, the agriculture sector stood at 8.1% or RM89.5 billion to the Gross Domestic Product (GDP). Oil palm was a major contributor to the GDP of agriculture sector at 43.1% followed by other agriculture (19.5%), livestock (11.6%), fishing (11.5%), forestry & logging (7.2%) and rubber (7.1%) (Department of Statistics Malaysia, 2017). Malaysia is one of the top producers and exporters of the palm oil globally, producing 103.96 million tonnes of oil palm fresh fruit bunch in 2017 (Economics and Industry Development Division, 2018).

Oil palms have an extensive range of products and applications in both food and non-food sector. Its multi-purpose property makes oil palm industry high in demand. The potential productivity of oil palm is several times greater than the other oil-producing crops, given that the crop is managed appropriately, much less land is required to produce the same amount of vegetable oil compared with other vegetable oil crops (Donough et al., 2009).

However, the oil palm tree in Malaysia is no exception to disease. Among other diseases, a fungal disease caused by *Ganoderma boninense*, known as basal stem rot (BSR) leads to great losses in oil palm production. BSR is notorious as one of the most destructive diseases at oil palm tree in plantations around Southeast Asia, particularly in Malaysia and North Sumatra (Flood et al., 2000). It attacks the basal stem of oil palm trees making them rot and slowly affecting xylem of the trees and causing water transportation to the upper part of oil palm become disturbed, thus the leaves at the oil palm's frond turn yellow. It is a major threat to the palm oil industry as it can reduce the yield of oil palm, hence causing economic losses to oil palm plantations (Kamu et al., 2016).

Different treatment methods were invented to treat BSR infection. Among the common methods used in controlling BSR infection are fungicide treatment (George et al., 1996) and biological control (Sapak et al., 2006). Another common method used in treating BSR-infected oil palm trees is by removing the infected oil palms and mounding the soil around the tree or a mixture of these two methods (Darus & Abu Seman, 2002). However, none of the methods mentioned is completely adequate in stopping and minimizing the effects of the

disease on the yield production as well as the health of the trees (Singh, 1991). Some plantations purposely leave the trees without any treatment to reduce the treatment cost, causing the yield production to reduce over time.

It was found that the existing treatments for BSR infection were not completely efficient in treating the oil palm trees in the later stages of infection. Therefore, detection of BSR infection is needed in order to treat the trees as soon as possible before the infection in oil palm plantation becomes worse. Currently, there are three approaches to detect BSR; manual, laboratory-based and remote sensing method. The manual observation is the most common technique to detect BSR. It was done by detecting the *Ganoderma* specific foliar symptoms and fungus fruiting bodies which grow as a parasite on the trunks. However, it requires a labour force with the knowledge to identify BSR-infected trees and differentiate other unhealthy trees due to different causes. It is also time-consuming as the oil palm trees are needed to be observed carefully and closely to check for the symptom one by one.

For laboratory-based approaches, the test sample is extracted by drilling the trunk of the oil palm tree. Ganoderma Selective Medium (GSM) test and Polymerase chain reaction (PCR) analysis are done using the extracted samples for the isolation, growth and identification of the fungus in the samples (Lim & Fong, 2005). The diagnostics method used are usually difficult to do due to its complex procedures. Some of the methods are also considered as expensive because of the cost needed to provide all necessary components for the analysis (Lelong et al., 2010; Wahab et al., 2011).

Non-invasive remote sensing based approaches have also been explored in order to identify and map the BSR-infected trees using different techniques such as ground-based, airborne and space-borne remote sensing. Hyperspectral and multispectral remote sensing approaches had shown the capability of identifying healthy and BSR-infected trees from recent studies (Khairunniza-Bejo et al., 2015; Mohamad Anuar et al., 2015). Terrestrial Laser Scanning (TLS) (Khairunniza-Bejo & Vong, 2014), Intelligent Electronic Nose (E-Nose) system (Abdullah et al., 2012; Azahar et al., 2011), tomographic sensor (Hamidon & Mukhlisin, 2014) and Microfocus X-ray Fluorescence (μXRF) (Meor Yusoff et al., 2009) also showed positive results in detecting BSR-infected tree. These reports showed that the techniques used are capable of early detection of BSR and differentiating the healthy and BSR-infected trees. However, some of the approaches were still limited to further classify the severity level of BSR infection (Khosrokhani et al., 2016).

Thermal imaging is a method of detecting infrared radiation of an object represented by an image. All objects with a temperature greater than absolute

zero (-273°C) emit infrared radiation, however, human's vision is limited to visible spectrum electromagnetic radiation. Thermal imaging extends human's limited vision beyond the boundary to view the infrared radiation. Thermal imaging has been used in various applications in different industries such as detecting moisture and building failure, including leaks. It is also used in medical field to detect peripheral vascular disorders (Bagavathiappan et al., 2009), to aid in detecting breast cancer (Arora et al., 2008) and to assist in detecting bone fracture injury (Cook et al., 2005). For agricultural industry, it is used to determine the plant physiological state and irrigation scheduling (Jones, 2004) and yield forecasting (Smith et al., 1985; Stajnko et al., 2004). Water stress in sunflower leaves can also be analysed from thermal image processing system (Hashimoto et al., 1984). It was also employed in detecting infestation by *Cryptolestes ferrugineus* inside wheat kernels (Manickavasagan et al., 2008) and to evaluate the damage to fruits and vegetables due to microbial activities (Hellebrand et al., 2002). Sankaran et al. (2013) used a combination of visible, near infrared and thermal imaging techniques to detect citrus greening disease in citrus trees. Based on the literature, it can be concluded that thermal imaging was able to analyse the water stress of plants as well as detecting disease. As BSR infection affected the water transportation of oil palm, the water stress of the BSR infected tree will vary from the healthy tree. Therefore, there is a potential use of thermal imaging in detecting BSR infection.

1.2 Problem Statement

A variety of study and approach has been explored for the detection of BSR disease in oil palm trees including manual visual inspection, laboratory analysis and remote sensing. The capability of these techniques to detect BSR in oil palm tree is still limited in term of the labour force, cost and time. It is also noted that some approaches still have a limitation in detecting BSR infection at the early stage.

Thermal imaging has been used in agricultural applications for disease detection and water stress analysis. Previous studies showed a good potential of using thermal imaging in detecting oil palm tree infected by BSR as BSR infection also affected the water stress of the oil palm tree. However, up to recently, the study on detecting the infected BSR tree in palm oil plantation using thermal imaging is still a largely unexplored research area to date. This research will explore and evaluate the potential of thermal imaging in detecting BSR at oil palm trees.

1.3 Objectives

The goal of this research is to explore the potential of using thermal imaging to identify oil palm trees infected with Basal Stem Rot (BSR) disease. Specific objectives are:

i. To identify the suitable time and tree section for thermal image acquisition.
ii. To identify the suitable features to differentiate healthy and BSR-infected trees extracted from thermal images.
iii. To classify BSR-infected trees using machine learning techniques.

1.4 The Scope of the Study

This study explored the potential of using thermal images to detect BSR disease at matured oil palm trees. The healthiness status of the trees was classified by the expert from Malaysian Palm Oil Board (MPOB). Therefore, the differences of the tree condition were assumed due to *Ganoderma boninense* infection, not by other factors. The emissivity of the thermal camera was fixed to 0.98 to ensure all the images are in the same condition. This study was focused on analysing the variation of thermal in intensity, not in detecting the temperature of the trees.

REFERENCES

Abdullah, A. H., Adom, A., Md Shakaff, A. Y., Ahmad, M. N., Zakaria, A., Ahmad Saad, F. S., … Kamarudin, L. M. (2012). Hand-Held Electronic Nose Sensor Selection System for Basal Stamp Rot (BSR) Disease Detection. In *2012 Third International Conference on Intelligent Systems Modelling and Simulation* (pp. 737–742). https://doi.org/10.1109/ISMS.2012.139

Abdullah, A. H., Adorn, A. H., Md Shakaff, A. Y., Ahmad, M. N., Saad, M. A., Tan, E., … Zakaria, A. (2011). *Electronic nose system for ganoderma detection*. *Sensor Letters* (Vol. 9).

Abdullah, F., Ilias, G. N. M., Nelson, M., Mohd Zainudin, N. A. I., & Umi Kalsom, Y. (2003). Disease assessment and the efficacy of Trichoderma as a biocontrol agent of basal stem rot of oil palm. *Research Bulletin Science Putra*, *11*, 31–33.

Abu Seman, I., Kushairi, A., Darus, A., & Wahid, M. B. (2006). Technique for Inoculation of Oil Palm. *MPOB Information Series*, *321*(314), 1–4.

Abu Seman, I., Mohd Su'ud, M., Loonis, P., & Wahid, M. B. (2010). GanoSken for early detection of ganoderma infection in oil palm. *MPOB Information Series*, *June 2010*(MPOB TT No. 442), 3–6.

Abu Seman, I., & Rafidah, A. R. (2008). Polyclonal antibody for detection of Ganoderma. *MPOB Information Series*, *June 2008*(MPOB TT No. 405).

Abu Seman, I., S., I., & Darus, A. (2004). Innovative Technique of Sanitation for Controlling Ganoderma at Replanting. *MPOB Information Series*, *June 2004*(MPOB TT No 213), 3–6.

Abu Seman, I., Yamaoka, M., Hayakawa, S., Wahid, M. B., Noorhasimah, I., & Darus, A. (2003). PCR Technique for Detection Ganoderma. *MPOB Information Series*, *June 2003*(MPOB TT No. 188), 1–4.

Adam, H., Collin, M., Richaud, F., Beulé, T., Cros, D., Omoré, A., … Tregear, J. W. (2011). Environmental regulation of sex determination in oil palm:

current knowledge and insights from other species. *Annals of Botany*, *108*(8), 1529–1537. https://doi.org/10.1093/aob/mcr151

Ahmadi, P., Muharam, F. M., Ahmad, K., Mansor, S., & Abu Seman, I. (2017). Early Detection of Ganoderma Basal Stem Rot of Oil Palms Using Artificial Neural Network Spectral Analysis. *Plant Disease*, *101*(6), 1009–1016. https://doi.org/10.1094/PDIS-12-16-1699-RE

Ajanki, A. (2007). Example of k-nearest neighbour classification. Retrieved from https://en.wikipedia.org/wiki/K-nearest_neighbors_algorithm#/media/File:KnnClassification.svg

Alpaydin, E. (2010). *Introduction to Machine Learning*. (T. Dietterich, C. Bishop, D. Heckerman, M. Jordan, & M. Kearns, Eds.) (Second). London: The MIT Press. https://doi.org/10.1016/j.neuroimage.2010.11.004

Amon, F., Hamins, A., Bryner, N., & Rowe, J. (2008). *Meaningful performance evaluation conditions for fire service thermal imaging cameras*. *Fire Safety Journal* (Vol. 43). https://doi.org/10.1016/j.firesaf.2007.12.006

Arango, M., Martínez, G., & Torres, G. (2016). Advances in the Interpretation of Tomographic Images as an Early Detection Method of Oil Palm Affected by Basal Stem Rot in Colombia. *Plant Disease*, *100*(8), 1559–1563. https://doi.org/10.1094/PDIS-12-15-1473-RE

Arora, N., Martins, D., Ruggerio, D., Tousimis, E., Swistel, A. J., Osborne, M. P., & Simmons, R. M. (2008). Effectiveness of a noninvasive digital infrared thermal imaging system in the detection of breast cancer. *The American Journal of Surgery*, *196*(4), 523–526. https://doi.org/10.1016/j.amjsurg.2008.06.015

Azahar, T. M., Che Mustapha, J., Mazliham, S., & Boursier, P. (2011). Temporal Analysis of Basal Stem Rot Disease in Oil Palm Plantations: An Analysis on Peat Soil. *International Journal of Engineering & Technology IJET-IJENS*, *11*(3), 96–101.

Bagavathiappan, S., Thangavelu, S., John, P., Jayakumar, T., Raj, B., Karunanithi, R., … Jagadeesan, K. (2009). Infrared thermal imaging for detection of peripheral vascular disorder. *Journal of Medical Physics*, *34*(1),

43–47. https://doi.org/10.4103/0971-6203.48720

Ballester, C., Jiménez-Bello, M. A., Castel, J. R., & Intrigliolo, D. S. (2013). Usefulness of thermography for plant water stress detection in citrus and persimmon trees. *Agricultural and Forest Meteorology*, *168*, 120–129. https://doi.org/10.1016/J.AGRFORMET.2012.08.005

Banerjee, K., Krishnan, P., & Mridha, N. (2018). Application of thermal imaging of wheat crop canopy to estimate leaf area index under different moisture stress conditions. *Biosystems Engineering*, *166*, 13–27. https://doi.org/10.1016/J.BIOSYSTEMSENG.2017.10.012

Battalwar, P., Gokhale, J., & Bansod, U. (2015). Infrared Thermography and IR Camera. *International Journal of Research in Science & Engineering*, *1*(3), 9–14.

Benjamin, M., & Chee, K. H. (1995). Basal stem rot of oil palm - a serious problem on inland soils. *MAPPS Newsletter*, *19*(1), 3.

Berdugo, C. A., Zito, R., Paulus, S., & Mahlein, A. K. (2014). Fusion of sensor data for the detection and differentiation of plant diseases in cucumber. *Plant Pathology*, *63*(6), 1344–1356. https://doi.org/10.1111/ppa.12219

Bhange, M., & Hingoliwala, H. A. (2015). Smart Farming: Pomegranate Disease Detection Using Image Processing. *Procedia Computer Science*, *58*, 280–288. https://doi.org/10.1016/j.procs.2015.08.022

Campbell, J. B. (2002). *Introduction to Remote Sensing*. CRC Press. Retrieved from https://books.google.com.my/books?id=1KfQxsN0vp8C

Chang, A., Jung, J., Maeda, M., & Landivar, J. (2017). *Crop height monitoring with digital imagery from Unmanned Aerial System (UAS)*. *Computers and Electronics in Agriculture* (Vol. 141). https://doi.org/10.1016/j.compag.2017.07.008

Chavarria, G., & dos Santos, H. P. (2012). Plant Water Relations: Absorption, Transport and Control Mechanisms. *Advances in Selected Plant Physiology Aspects*. https://doi.org/10.5772/33478

Chemat, S. (2017). *Edible Oils: Extraction, Processing, and Applications*. (S. Chemat, Ed.). CRC Press. Retrieved from https://books.google.com.my/books?id=klAsDwAAQBAJ

Chen, Y., Chen, D., Li, Y., & Chen, L. (2010). Otsu's thresholding method based on gray level-gradient two-dimensional histogram. In *2010 2nd International Asia Conference on Informatics in Control, Automation and Robotics (CAR 2010)* (Vol. 3, pp. 282–285). https://doi.org/10.1109/CAR.2010.5456687

Chung, G. F. (2015). Effect of Pests and Diseases on Oil Palm Yield. In O. M. Lai, C. P. Tan, & C. C. Akoh (Eds.), *Palm Oil: Production, Processing, Characterization, and Uses*. Elsevier Science. Retrieved from https://books.google.com.my/books?id=6uRxCgAAQBAJ

Cook, R. J., Thakore, S., & Nichol, N. M. (2005). Thermal imaging: A hotspot for the future? *Injury Extra*, *36*, 395–397.

Corley, R. H. V., & Tinker, P. B. H. (2008). *The Oil Palm* (4th ed.). John Wiley and Sons. Retrieved from https://books.google.com.my/books?id=NtCo1TdXuQkC

Cortes, C., & Vapnik, V. (1995). Support-Vector Networks. *Machine Learning*, *20*, 273–297. https://doi.org/10.1023/A:1022627411411

Danielson, M. (2016). Why palm oil is so attractive to food manufacturers. Retrieved April 30, 2018, from https://www.palmoilhealth.org/faq/palm-oil-attractive-food-manufacturers/

Danno, A. K., Miyazato, M., & Ishiguro, E. (1978). Quality evaluation of agricultural products by infrared imaging method, 1: Grading of fruits for bruise and other surface defects. *Memoirs of the Faculty of Agriculture, Kagoshima University1*.

Danno, A. K., Miyazato, M., & Ishiguro, E. (1980). Quality evaluation of agricultural products by infrared imaging method. III. Maturity evaluation of fruits and vegetables. *Memoirs of the Faculty of Agriculture, Kagoshima University*.

Darus, A., & Abu Seman, I. (1991). A selective medium for the isolation of Ganoderma from diseased tissues. In *Proceeding of the 1991 International Palm Oil Conference: Progress, Prospects and Challenge Towards the 21st Century - Agriculture Conference* (pp. 517–519).

Darus, A., & Abu Seman, I. (2002). Progress and Research on Ganoderma Basal Stem Rot of Oil Palm. In *Seminar Recent Progress in the Management of Peat and Ganoderma* (p. 50). Bangi: Malaysian Palm Oil Board.

Darus, A., Abu Seman, I., & Hashim, K. (1993). Confirmation of Ganoderma infected palm by drilling technique. In S. Jalani, A. Darus, N. Rajanaidu, D. Tayeb, K. Paranjothy, M. B. Wahid, … K. C. Chang (Eds.), *Proceeding of 1993 PORIM International Palm Oil Congress: Update and Vision (Agriculture)* (pp. 735–738).

Darus, A., Abu Seman, I., & Hassan, A. H. (1989). Significance of the black line within oil palm tissue decayed by Ganoderma boninense. *Elaieis*, *1*(1), 11–16.

Darus, A., Abu Seman, I., & Singh, G. (2000). Status of Ganoderma in oil palm. In J. Flood, P. D. Bridge, & M. Holderness (Eds.), *Ganoderma. Diseases of Perennial Crops.* (pp. 49–68). Wallingford: CABI.

Department of Environment. (1999). *Industrial Processes & The Environment (Handbook 3): Crude Palm Oil Industry*. (M. I. Thani, R. Hussin, W. R. Wan Ibrahim, & M. S. Sulaiman, Eds.).

Department of Statistics Malaysia. (2017, December 22). Selected Agricultural Indicators, Malaysia, 2017, pp. 1–4.

Dey, A. K., Sharma, M., & Meshram, M. R. (2016). Image Processing Based Leaf Rot Disease, Detection of Betel Vine (Piper betle L.). *Procedia Computer Science*, *85*, 748–754. https://doi.org/10.1016/j.procs.2016.05.262

Donough, C. R., Witt, C. W., & Fairhurst, T. H. (2009). Yield Intensification in Oil Palm Plantations through Best Management Practice. *Better Crops*, *93*(1), 12–14.

Economics and Industry Development Division. (2018). *Overview of the Malaysian Oil Palm Industry 2017*. *Malaysian Palm Oil Board*. Retrieved from http://bepi.mpob.gov.my

Eslamian, S. (2014). *Handbook of Engineering Hydrology: Fundamentals and Applications*. CRC Press. Retrieved from https://books.google.com.my/books?id=SH3NBQAAQBAJ

Flood, J., Hasan, Y., Turner, P. D., & O'Grady, E. B. (2000). The spread of Ganoderma from infective sources in the field and its implications for management of the disease in oil palm. In J. Flood, P. D. Bridge, & M. Holderness (Eds.), *Ganoderma. Diseases of Perennial Crops.* (pp. 101–112). Wallingford: CABI. https://doi.org/10.1079/9780851993881.0101

Frater, M. R., & Arnold, J. F. (2000). Coding of 12-bit video from thermal imaging systems. *Signal Processing: Image Communication*, *15*(10), 907–916. https://doi.org/10.1016/S0923-5965(99)00038-7

George, S. T., Chung, G. F., & Zakaria, K. (1996). Updated results (1990-1995) on trunk injection of fungicides for the control of Ganoderma basal stem rot. In A. Darus, M. B. Wahid, R. N, M. T. D, P. K, S. C. Cheah, … R. S (Eds.), *Proceedings of the 1996 PORIM International Palm Oil Congress - Agriculture Conference* (pp. 508–515). Kuala Lumpur: PORIM.

George, S. T., Chung, G. F., & Zakaria, K. (2000). Benefits of Soil Mounding Tall Palms in a High Ganoderma Incidence Area in Lower Perak. In *International Planters Conference 2000* (pp. 565–576). Kuala Lumpur.

Giorleo, G., & Meola, C. (2002). Comparison between pulsed and modulated thermography in glass–epoxy laminates. *NDT & E International*, *35*(5), 287–292. https://doi.org/10.1016/S0963-8695(01)00062-7

Gong, A., Yu, J., He, Y., & Qiu, Z. (2013). Citrus yield estimation based on images processed by an Android mobile phone. *Biosystems Engineering*, *115*, 162–170. https://doi.org/10.1016/j.biosystemseng.2013.03.009

Gonzalez, R. C., Woods, R. E., & Eddins, S. L. (2009). Introduction. In *Digital Imaging Processing Using MATLAB* (2nd ed.). Gatesmark Publishing. https://doi.org/10.1109/IEMBS.2011.6091204

Grover, S., & Gusain, V. (2014). Digital image processing. *International Journal of Innovative Research in Technology*, *1*(6), 70–73. Retrieved from http://dl.acm.org/citation.cfm?id=518399

Habi Mat Dian, N. L., Abd Hamid, R., Kanagaratnam, S., Awg Isa, W. R., Mohd Hassim, N. A., Ismail, N. H., … Mat Sahri, M. (2017). Palm Oil and Palm Kernel Oil: Versatile Ingredients for Food Applications. *Journal of Oil Palm Research*, *29*(4), 487–511. https://doi.org/10.21894/jopr.2017.00014

Hamidon, N. A., & Mukhlisin, M. (2014). A Review of Application of Computed Tomography on Early Detection of Basal Stem Rot Disease. *Jurnal Teknologi (Sciences & Engineering)*, *70*(3).

Harun, M. H., & Md Noor, M. R. (2006). Canopy temperature difference (CTD) for detecting stress in oil palm. *MPOB Information Series*, *June 2006*(MPOB TT No. 309).

Harun, M. H., S., I., & Abu Seman, I. (2005). Gas exchange responses of oil palm to Ganoderma boninense infection. *Asian Journal of Plant Sciences*. https://doi.org/10.3923/ajps.2005.438.444

Hasan, Y., & Turner, P. D. (1998). The comparative importance of different oil palm tissues as infection sources for basal stem rot in replantings. *The Planter*, *74*(864), 119–135.

Hashim, K. (1990). *Basal Stem Rot of Oil Palm : Incidence, Etiology and Control*. Universiti Pertanian Malaysia.

Hashim, K. (1995). Basal stem rot of oil palm caused by Ganoderma boninense: an update. In B. S. Jalani (Ed.), *Proceeding 1993 PORIM International Palm Oil Congress - Agriculture* (pp. 739–749). Kuala Lumpur: Palm Oil Research Institute of Malaysia.

Hashimoto, Y., Ino, T., Kramer, P. J., Naylor, A. W., & Strain, B. R. (1984). Dynamic Analysis of Water Stress of Sunflower Leaves by Means of a Thermal Image Processing System. *Plant Physiology*, *76*, 266–269. https://doi.org/10.1104/pp.76.1.266

Hassan Abdalla, S., & Fattoh Osman, S. E. (2016). Digital Image Processing Technology based on MATLAB. *International Journal of Advanced Research in Computer Science*, *7*(3).

Heldman, D. R., & Moraru, C. I. (2014). *Encyclopedia of Agricultural, Food, and Biological Engineering, Second Edition* - (2nd ed.). CRC Press. Retrieved from https://books.google.com.my/books?id=MU5ZDwAAQBAJ

Hellebrand, H. J., Beuche, H., & Linke, M. (2002). Thermal Imaging. In J. Blahovec & M. Kutílek (Eds.), *Physical Methods in Agriculture: Approach to Precision and Quality* (pp. 411–427). Boston, MA: Springer US. https://doi.org/10.1007/978-1-4615-0085-8_22

Ho, C. T., & Hashim, K. (1997). Usefulness of soil mounding treatments in prolonging productivity of prime-aged Ganoderma infected palms. *The Planter*, *73*, 239–244.

Ho, Y. W., & Nawawi, A. (1985). Ganoderma boninense Pat. from based stem rot of oil palm [Elaeis guineensis] in Peninsular Malaysia. *Pertanika*, *8*(3), 425–428.

Husin, Z., Md Shakaff, A. Y., Abdul Aziz, A. H., & S Mohamed Farook, R. (2012). Feasibility Study on Plant Chili Disease Detection Using Image Processing Techniques. *2012 Third International Conference on Intelligent Systems Modelling and Simulation*, 291–296. https://doi.org/10.1109/ISMS.2012.33

Imran, S. N. M., Baharudin, F., Ali, M. F., & Rahiman, M. H. F. (2018). Chemical Relationship on Detection of Ganoderma Disease on Oil Palm Tree System. *IOP Conference Series: Earth and Environmental Science*, *140*(1). https://doi.org/10.1088/1755-1315/140/1/012048

Ishaq, I., Alias, M. S., Kadir, J., & Kasawani, I. (2014). *Detection of basal stem rot disease at oil palm plantations using sonic tomography*. *Journal of Sustainability Science and Management* (Vol. 9).

Jafari, M., Minaei, S., & Safaie, N. (2017). Detection of pre-symptomatic rose powdery-mildew and gray-mold diseases based on thermal vision. *Infrared Physics and Technology*, *85*, 170–183. https://doi.org/10.1016/j.infrared.2017.04.023

Jones, H. G. (1999). Use of infrared thermometry for estimation of stomatal conductance as a possible aid to irrigation scheduling. *Agricultural and Forest Meteorology*, *95*(3), 139–149. https://doi.org/10.1016/S0168-1923(99)00030-1

Jones, H. G. (2004). Irrigation scheduling: Advantages and pitfalls of plant-based methods. *Journal of Experimental Botany*, *55*(407), 2427–2436. https://doi.org/10.1093/jxb/erh213

Jones, H. G., Stoll, M., Santos, T., de Sousa, C. I. G., Chaves, M. M., & Grant, O. M. (2002). Use of infrared thermography for monitoring stomatal closure in the field: application to grapevine. *Journal of Experimental Botany*, *53 378*, 2249–2260.

Kamu, A., Chong, K. P., Abu Seman, I., & Ho, C. M. (2016). Economic Loss due to Ganoderma Disease in Oil Palm. *International Journal of Economics and Management Engineering*, *10*(2), 604–608. Retrieved from https://waset.org/Publication/economic-loss-due-to-ganoderma-disease-in-oil-palm/10003869

Khairunniza-Bejo, S., & Vong, C. N. (2014). Detection of Basal Stem Rot (BSR) Infected Oil Palm Tree Using Laser Scanning Data. *Agriculture and Agricultural Science Procedia*, *2*, 156–164. https://doi.org/10.1016/j.aaspro.2014.11.023

Khairunniza-Bejo, S., Yusoff, Y., Nik Yusoff, N. S., Abu Seman, I., & Mohamad Anuar, I. (2015). Identification of Healthy and BSR-Infected Oil Palm Trees Using Color Indices. *International Journal of Biological, Biomolecular, Agricultural, Food and Biotechnological Engineering*, *9*(8), 876–879. Retrieved from https://waset.org/Publication/identification-of-healthy-and-bsr-infected-oil-palm-trees-using-color-indices/10001908

Khosrokhani, M., Khairunniza-Bejo, S., & Pradhan, B. (2016). Geospatial technologies for detection and monitoring of Ganoderma basal stem rot infection in oil palm plantations: a review on sensors and techniques. *Geocarto International*. https://doi.org/10.1080/10106049.2016.1243410

Kumar, G., & Bhatia, P. K. (2014). A Detailed Review of Feature Extraction in Image Processing Systems. In *2014 Fourth International Conference on Advanced Computing & Communication Technologies* (pp. 5–12).

https://doi.org/10.1109/ACCT.2014.74

Lee, D.-J., Schoenberger, R., Archibald, J., & McCollum, S. (2008). Development of a machine vision system for automatic date grading using digital reflective near-infrared imaging. *Journal of Food Engineering*, *86*(3), 388–398. https://doi.org/10.1016/J.JFOODENG.2007.10.021

Lelong, C. C. D., Roger, J.-M., Brégand, S., Dubertret, F., Lanore, M., A Sitorus, N., … Caliman, J.-P. (2010). Evaluation of Oil-Palm Fungal Disease Infestation with Canopy Hyperspectral Reflectance Data. *Sensors (Basel, Switzerland)*, *10*(1), 734–747. https://doi.org/10.3390/s100100734

Liaghat, S., Mansor, S., Ehsani, R., Shafri, H. Z. M., Meon, S., & Sankaran, S. (2014). Mid-infrared spectroscopy for early detection of basal stem rot disease in oil palm. *Computers and Electronics in Agriculture*, *101*, 48–54. https://doi.org/10.1016/j.compag.2013.12.012

Lillesand, T., Kiefer, R. W., & Chipman, J. (2014). *Remote Sensing and Image Interpretation*. John Wiley and Sons. Retrieved from https://books.google.com.my/books?id=AFHDCAAAQBAJ

Lim, H. P., & Fong, Y. K. (2005). Research on basal stem rot (BSR) of ornamental palms caused by basidiospores from Ganoderma boninense. *Mycopathologia*, *159*(1), 171–179. https://doi.org/10.1007/s11046-004-4440-6

Lim, K. H., Chuah, J. H., & Ho, C. H. (1993). Effects of soil heaping on Ganoderma infected oil palms. In *Proceedings of the 1993 PORIM International Palm Oil Congress "Update and Vision" (Agriculture)* (pp. 735–738). Bangi: Palm Oil Research Institute of Malaysia.

Lim, K. H., & Udin, W. (2010). Management of Ganoderma in peat soil in Indonesi. In *2010 Second International Seminar Oil Palm Diseases: Advances in Ganoderma Research and Management* (p. 11). Yogyakarta.

Lisnawita, Hanum, H., & Tantawi, A. R. (2016). Survey of Basal Stem Rot Disease on Oil Palms (Elaeis guineensis Jacq.) in Kebun Bukit Kijang,North Sumatera, Indonesia. *IOP Conference Series: Earth and Environmental Science*, *41*(1). https://doi.org/10.1088/1755-1315/41/1/012007

Malaysian Palm Oil Council. (2012). The Oil Palm Tree. Retrieved April 30, 2018, from http://www.mpoc.org.my/The_Oil_Palm_Tree.aspx

Manickavasagan, A., Jayas, D. S., & White, N. D. G. (2008). Thermal imaging to detect infestation by Cryptolestes ferrugineus inside wheat kernels. *Journal of Stored Products Research*, *44*, 186–192. https://doi.org/10.1016/j.jspr.2007.10.006

Markom, M. A., Shakaff, A. Y. M., Adom, A. H., Ahmad, M. N., Hidayat, W., Abdullah, A. H., & Fikri, N. A. (2009). Intelligent electronic nose system for basal stem rot disease detection. *Computers and Electronics in Agriculture*, *66*(2), 140–146. https://doi.org/10.1016/J.COMPAG.2009.01.006

Meor Yusoff, M. S., Khalid, M. A., & Abu Seman, I. (2009). Identification of Basal Stem Rot Disease in Local Palm Oil By Microfocus Xrf. *Journal of Nuclear and Related Technologies*, *6*(1), 282–287.

Merlot, S., Mustilli, A.-C., Genty, B., North, H., Lefebvre, V., Sotta, B., … Giraudat, J. (2002). Use of infrared thermal imaging to isolate Arabidopsis mutants defective in stomatal regulation. *The Plant Journal*, *30*(5), 601–609. https://doi.org/10.1046/j.1365-313X.2002.01322.x

Meunkaewjinda, A., Kumsawat, P., Attakitmongcol, K., & Srikaew, A. (2008). Grape leaf disease detection from color imagery using hybrid intelligent system. *2008 5th International Conference on Electrical Engineering/Electronics, Computer, Telecommunications and Information Technology*, 513–516. https://doi.org/10.1109/ECTICON.2008.4600483

Mohamad Anuar, I., Abu Seman, I., Nor Maris, N., & Bahrom, E. (2015). Spectral based Analysis of Airborne Hyperspectral Remote Sensing Image for Detection of Ganoderma Disease in Oil Palm. In *Int'l Conference on Biological and Environmental Science (BIOES 2015)* (pp. 13–20). Phuket.

Mohamad Anuar, I., Abu Seman, I., Wahid, M. B., Mohd Noor, N., & Mohd Shafri, H. Z. (2012). Field spectroscopy for detection of Ganoderma disease in oil palm. *MPOB Information Series*, (MPOB TT No 532).

Ng, E. Y. K. (2009). A review of thermography as promising non-invasive detection modality for breast tumor. *International Journal of Thermal*

Sciences, *48*(5), 849–859. https://doi.org/10.1016/J.IJTHERMALSCI.2008.06.015

Othman, N., Manan, Z. A., Wan Alwi, S. R., & Sarmidi, M. R. (2010). A Review of Extraction Technology for Carotenoids and Vitamin E Recovery from Palm Oil. *Journal of Applied Science*, *10*(12), 1187–1191. https://doi.org/10.3923/jas.2010.1187.1191

Otsu, N. (1979). A Threshold Selection Method from Gray-Level Histograms. *IEEE Transactions on Systems, Man, and Cybernetics*, *9*(1), 62–66. Retrieved from http://ieeexplore.ieee.org/document/4310076/

Owoyele, B. V., & Owolabi, G. O. (2014). Traditional oil palm (Elaeis guineensis jacq.) and its medicinal uses: A review. *Tang [Humanitas Medicine]*, *4*(3), 16.1-16.8. https://doi.org/10.5667/tang.2014.0004

Paterson, R. R. M., Holderness, M., Kelly, J., Miller, R., & O'Grady, E. B. (2000). In vitro diodegradation of oil-palm stem using macroscopic fungi from South East Asia: a preliminary investigation. In J. Flood, P. D. Bridge, & M. Holderness (Eds.), *Ganoderma. Diseases of Perennial Crops.* (pp. 129–138). Wallingford.

Peng, D., Huang, J., Li, C., Liu, L., Huang, W., Wang, F., & Yang, X. (2014). *Modelling paddy rice yield using MODIS data*. *Agricultural and Forest Meteorology* (Vol. 184). https://doi.org/10.1016/j.agrformet.2013.09.006

Phadikar, S., & Sil, J. (2008). Rice disease identification using pattern recognition techniques. In *Proceedings of 11th International Conference on Computer and Information Technology (ICCIT 2008)* (pp. 420–423). https://doi.org/10.1109/ICCITECHN.2008.4803079

Rahkonen, J., & Jokela, H. (2003). Infrared Radiometry for Measuring Plant Leaf Temperature during Thermal Weed Control Treatment. *Biosystems Engineering*, *86*(3), 257–266. https://doi.org/10.1016/S1537-5110(03)00138-7

Rao, A. K. (1990). Basal stem rot "Ganoderma" problem in oil palm smallholdings - the West Johor experience. In *Proceedings of Ganoderma workshop* (pp. 36–48). Kuala Lumpur: Palm Oil Research Institute of Malaysia.

Rees, R. W., Flood, J., Hasan, Y., Potter, U., & Cooper, R. M. (2009). Basal stem rot of oil palm (Elaeis guineensis); mode of root infection and lower stem invasion by Ganoderma boninense. *Plant Pathology*, *58*(5), 982–989. https://doi.org/10.1111/j.1365-3059.2009.02100.x

Riazi, M. R., & Chiaramonti, D. (2017). Introduction: Biofuel Production and Processing Technology. In M. R. Riazi & D. Chiaramonti (Eds.), *Biofuels Production and Processing Technology* (pp. 1–9).

Roughgarden, T., & Valiant, G. (2016). CS168: The Modern Algorithmic Toolbox Lecture #8: PCA and the Power Iteration Method.

Rubin, G. D. (2014). Computed Tomography: Revolutionizing the Practice of Medicine for 40 Years. *Radiology*, *273*(2S), S45–S74. https://doi.org/10.1148/radiol.14141356

Sambanthamurthi, R., Singh, R., Ghulam Kadir, A. P., Ong-Abdullah, M., & Kushairi, A. (2009). Opportunities for the Oil Palm via Breeding and Biotechnology. In *Breeding Plantation Tree Crops: Tropical Species* (pp. 377–421). https://doi.org/10.1007/978-0-387-71201-7_11

Sankaran, S., Maja, J., Buchanon, S., & Ehsani, R. (2013). Huanglongbing (Citrus Greening) Detection Using Visible, Near Infrared and Thermal Imaging Techniques. *Sensors*, *13*, 2117–2130. https://doi.org/10.3390/s130202117

Santoso, H., Gunawan, T., Jatmiko, R. H., Darmosarkoro, W., & Minasny, B. (2010). Mapping and identifying basal stem rot disease in oil palms in North Sumatra with QuickBird imagery. *Precision Agriculture*, *12*, 233–248.

Sapak, Z., Meon, S., & Zainal Abidin, M. A. (2006). Isolation and characterization of microbial endophytes from oil palm roots: Implication as biocontrol agents against Ganoderma. *Planter*, *82*, 587–597.

Sarawak Land Consolidation And Rehabilitation Authority. (2012). Pests and Diseases Control. Retrieved April 30, 2018, from http://www.salcra.gov.my/en/sustainable-plantation/pest-diseases-control.html

Sauki, A., Zazarli Shah, M. S., & Wan Bakar, W. Z. (2015). Application of Ester based Drilling Fluid for Shale Gas Drilling. *IOP Conference Series: Materials Science and Engineering*, *83*(1). https://doi.org/10.1088/1757-899X/83/1/012012

Sethupathi, S., Chai, S.-P., & Mohamed, A. R. (2008). Utilization of oil palm as a source of renewable energy in Malaysia. *Renewable and Sustainable Energy Reviews*, *12*(9), 2404–2421. https://doi.org/10.1016/j.rser.2007.06.006

Shuit, S. H., Tan, K. T., Lee, K. T., & Kamaruddin, A. H. (2009). Oil palm biomass as a sustainable energy source: A Malaysian case study. *Energy*, *34*(9), 1225–1235. https://doi.org/10.1016/J.ENERGY.2009.05.008

Singh, G. (1991). Ganoderma - the scourge of oil palms [Elaeis guineensis] in the coastal areas [Peninsular Malaysia]. *Planter*, *67*(786), 421–444.

Singh, V., & Misra, A. K. (2017). Detection of plant leaf diseases using image segmentation and soft computing techniques. *Information Processing in Agriculture*, *4*, 41–49. https://doi.org/10.1016/j.inpa.2016.10.005

Smith, R. C. G., Barrs, H. D., Steiner, J. L., & Stapper, M. (1985). Relationship between wheat yield and foliage temperature: theory and its application to infrared measurements. *Agricultural and Forest Meteorology*, *36*(2), 129–143. https://doi.org/10.1016/0168-1923(85)90005-X

Spruyt, V. (2014). The Curse of Dimensionality in Classification. Retrieved from http://www.visiondummy.com/2014/04/curse-dimensionality-affect-classification/

Stajnko, D., Lakota, M., & Hočevar, M. (2004). Estimation of number and diameter of apple fruits in an orchard during the growing season by thermal imaging. *Computers and Electronics in Agriculture*, *42*(1), 31–42. https://doi.org/10.1016/S0168-1699(03)00086-3

Susanto, A., Sudharto, P. S., & Purba, R. Y. (2005). Enhancing biological control of basal stem rot disease (Ganoderma boninense) in oil palm plantations. *Mycopathologia*, *159*, 153–157. https://doi.org/10.1007/s11046-004-4438-0

Teoh, C. H. (2002). The palm oil industry in Malaysia. From seed to frying pan., *14*(2), 24–33.

Turner, P. D. (1981). *Oil palm diseases and disorders.* Kuala Lumpur, Malaysia: Oxford Univ. Press.

Turner, P. D., & Gillbanks, R. A. (1974). *Oil Palm Cultivation and Management*. Kuala Lumpur: Yau Seng Press.

Varith, J., Hyde, G. M., Baritelle, A. L., Fellman, J. K., & Sattabongkot, T. (2003). Non-contact bruise detection in apples by thermal imaging. *Innovative Food Science and Emerging Technologies*, *4*, 211–218. https://doi.org/10.1016/S1466-8564(03)00021-3

Verheye, W. H. (2010a). Growth and Production of Oil Palm. In W. H. Verheye (Ed.), *Land Use, Land Cover and Soil Sciences*. UNESCO-EOLSS Publishers. Retrieved from https://biblio.ugent.be/publication/1009126/file/1009127.pdf

Verheye, W. H. (2010b). *Soils, Plant Growth and Crop Production - Volume II*. (W. H. Verheye, Ed.). EOLSS Publications. Retrieved from https://books.google.com.my/books?id=wwswCwAAQBAJ

Verleysen, M., & François, D. (2005). The Curse of Dimensionality in Data Mining and Time Series Prediction. In J. Cabestany, A. Prieto, & F. Sandoval (Eds.) (pp. 758–770). Berlin, Heidelberg: Springer Berlin Heidelberg.

Wahab, A., Meon, S., Paterson, R., Zainal Abidin, M. A., & Lima, N. (2011). Ergosterol analyses of oil palm seedlings and plants infected with Ganoderma. *Crop Protection*, *30*(11), 1438–1442. https://doi.org/10.1016/j.cropro.2011.07.004

Wong, L. C., Bong, C. F. J., & Abu Seman, I. (2012). Ganoderma Species Associated with Basal Stem Rot Disease of Oil Palm. *American Journal of Applied Sciences*, *9*(6), 879–885. https://doi.org/10.3844/ajassp.2012.879.885

Xu, L., & Zhao, Y. (2010). *Automated strawberry grading system based on image*

processing. *Computers and Electronics in Agriculture* (Vol. 71). https://doi.org/10.1016/j.compag.2009.09.013

Yao, Q., Chen, G., Wang, Z., Zang, C., Yang, B., & Tang, J. (2017). Automated detection and identification of white-backed planthoppers in paddy fields using image processing. *Journal of Integrative Agriculture*, *16*(7), 1547–1557. https://doi.org/10.1016/S2095-3119(16)61497-1

Yao, Q., Guan, Z., Zhou, Y., Tang, J., Hu, Y., & Yang, B. (2009). *Application of Support Vector Machine for Detecting Rice Diseases Using Shape and Color Texture Features*. *Engineering Computation, International Conference on* (Vol. 0). https://doi.org/10.1109/ICEC.2009.73

Zhang, S., Wu, X., You, Z., & Zhang, L. (2017). Leaf image based cucumber disease recognition using sparse representation classification. *Computers and Electronics in Agriculture*, *134*, 135–141. https://doi.org/10.1016/j.compag.2017.01.014