



UNIVERSITI PUTRA MALAYSIA

***MODELING SOLUTE TRANSPORT FOR IMPROVED FERTILISER
USE IN RICE PRODUCTION SYSTEM***

ABDIKANI ABDULLAHI MO'ALLIM

FK 2018 169



**MODELING SOLUTE TRANSPORT FOR IMPROVED FERTILISER
USE IN RICE PRODUCTION SYSTEM**

By

ABDIKANI ABDULLAHI MO'ALLIM

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

June 2018

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This thesis is dedicated to;

My parents for their endless love and wishes for their son to achieve this higher dream,

My two elder sisters, Naima and Hamdi Abdullahi, who has been supportive during my study,

My sibilings, Decca, Abdikarim, and Abdiaziz Abdullahi, who have been a source of inspiration to me throughout my study.



© COPYRIGHT

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**MODELING SOLUTE TRANSPORT FOR IMPROVED FERTILISER
USE IN RICE PRODUCTION SYSTEM**

By

ABDIKANI ABDULLAHI MO'ALLIM

June 2018

Chairman : Md Rowshon Kamal, PhD
Faculty : Engineering

Quantification of water and nutrients and their interactions of a paddy field environment are crucial for the improved utilization of fertilizers for the sustainable rice production. Solutes runoff and leaching are two direct pathways of nutrient pollution from paddy fields to water resources systems. Due to the dynamic nature of paddy fields, solute transport and transformation process are complex and difficult to understand. The past investigation on the water balance components using multifarious parameters did not reflect the true condition of paddy field environments. Quantification of agrochemical losses from paddy fields are generally related to the amount of inflow and outflow water in the paddy field environment which yet to be measured accurately. In order to overcome the challenges, the modern monitoring devices together with sensors and data logging system were installed for intensive field observations in a paddy and developed empirical models to quantify the solute losses through the surface and sub-surface water leaving from a paddy field system for the better utilization of fertilizers (N, P, K). The intensive field investigation was carried out in a paddy plot at Sawah Sempadan compartment of the Tanjung Karang Rice Irrigation Scheme (TAKRIS) for two rice growing seasons (January-April and July-October) in 2017.

Firstly, the water balance components in a paddy plot was analysed from the intensive field observations with 1-10 minutes interval of a paddy field. Water balance analysis results revealed that irrigation water accounted 59.6 % of the total water input (irrigation and rainfall) during the January to April (Off Season). However, about 76.2% of total water input during the July to October (Main season). The amount of rainfall contributed to 23.8% and 40.4% of total water input in the main season and off-season, respectively. Drainage flow accounted 37.3% and 43.7% of the total water input during off season and main season, respectively. The

daily evapotranspiration accounted 41.7% and 61% of total water input during off-season and main season, respectively. Observed seepage and percolation of 17.1% to 19.2% of total water input accounted during both seasons respectively. The yield of the experimental plot was obtained 2.5 t/ha and 2.7 t/ha for the off season and main season, respectively. Finally, the water productivity index was analyzed 0.72 kg m⁻³ during off-season and 0.78 kg m⁻³ during main season, respectively.

Based on solute transport analysis, the accumulated total nitrogen (T-N) of 50.3% to 49.7% estimated in the top 40 cm soil layer while 49.7 % to 53 % T-N as leachate obtained below 40 cm soil layer (40-100 cm) during off season and main season, respectively. About 85% of N leaching losses were in the form of NO₃⁻, however there was still a large quantity of NO₃⁻ remained below root zone that contributes the groundwater. The total leaching loss of T-N was 34.9 and 27.9 kg/ha during off and main seasons respectively. The estimated loss of total phosphorous during the two rice growing seasons were 3 and 1.7 kg/ha, respectively. The total amount of T-N, T-P and K loss through drainage were 27.7 and 18.5, 2.2 and 1.1, 5.9 and 3.5 kg/ha during off-season and main season, respectively.

The Hydrus-1D was applied to simulate water and solute movement under different soil depths of 20, 40, 60, 80 and 100 cm in real paddy environment experiments. The simulated and observed water flow and nutrient leaching were in good agreement ($R^2 = 0.98$, RMSE = 0.24). Hydrus-1D simulation showed the similar patterns of the water and solute movement under different soil depths during the study period. The observed and simulated N, P and K concentration in paddy was high due to fertilization and other climatic factors. Therefore, reduction of excessive fertilizer rate especially during early rice growing period and adaptation of water saving techniques can reduce the pollutant risks from paddy soil.

Regression analyses were performed for the development of the improved fertilizer use models. Multiple linear regression analysis was performed to know the relationships between EC versus solutes (N, P and K) during the both seasons. The polynomial regression analysis was fitted to evaluate whether EC changes has an impact on N, P and K concentrations in paddy field. Finally, empirical models were established to estimate the concentrations of N, P and K using two rice growing season data. MS Excel solver program were used to develop the empirical models. The results obtained a strong agreement between observed and predicted N, P, and K with the determination coefficients (R^2) of 0.91 and 0.95 during the both seasons. Therefore, the models could be useful in predicting the solute concentration changes within root zone and below root zone during entire rice growing season for better utilization of fertilizers.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PERMODELAN PENGANGKUTAN BAHAN LARUT BAGI
PENGUNAAN BAJA YANG LEBIH BAIK DALAM SISTEM
PENGELUARAN BERAS**

Oleh

ABDIKANI ABDULLAHI MO'ALLIM

Jun 2018

Pengerusi : Md Rowshon Kamal, PhD
Fakulti : Kejuruteraan

Kuantifikasi air dan nutrien serta interaksi mereka di dalam persekitaran padi adalah penting untuk penggunaan baja yang lebih baik bagi pengeluaran beras yang berkekalan. Bahan larut larian dan pengusaran adalah dua laluan langsung pencemaran nutrien dari sawah padi kepada sistem sumber air. Oleh kerana sifat padi yang dinamik, proses pengangkutan dan transformasi bahan larut menjadi kompleks dan sukar difahami. Penyiasatan lepas berkenaan komponen-komponen keseimbangan air menggunakan parameter yang pelbagai tidak mencerminkan keadaan sebenar persekitaran sawah padi. Kuantifikasi kerugian agrokimia dari sawah secara umumnya berkait dengan jumlah aliran masuk dan keluar air dalam persekitaran sawah padi yang belum diukur dengan tepat. Untuk mengatasi cabaran-cabaran ini, alat pemantauan moden bersama-sama dengan sensor dan sistem pengelogan data dipasang untuk pemerhatian lapangan yang intensif di dalam padi dan model empirikal yang dibangunkan bagi mengira kerugian bahan larut melalui permukaan dan sub-permukaan air yang meninggalkan sistem sawah padi untuk penggunaan baja yang lebih baik (N, P, K). Penyiasatan lapangan intensif dilakukan di plot padi di kompartmen Sawah Sempadan, Skim Pengairan Padi Tanjung Karang (TAKRIS) untuk dua musim penanaman padi (Januari-April dan Julai-Oktober) pada tahun 2017.

Pertama, komponen-komponen keseimbangan air dalam plot padi dianalisis dari pemerhatian lapangan intensif dengan jarak masa 1-10 minit pada sawah padi. Hasil analisa keseimbangan air menunjukkan bahawa air pengairan menyumbang 59.6% daripada jumlah input air (pengairan dan hujan) dari Januari hingga April (luar musim). Walau bagaimanapun, kira-kira 76.2% daripada jumlah input air diperhatikan dari bulan Julai hingga Oktober (musim utama). Jumlah air hujan menyumbang kepada 23.8% dan 40.4% daripada jumlah input air pada musim utama

dan luar musim. Manakala air saliran menyumbang 37.3% dan 43.7% daripada jumlah input air semasa luar musim dan musim utama. Evapotranspirasi harian menyumbang 41.7% dan 61% daripada jumlah input air pada luar musim dan musim utama. Rembesan dan perkolasi sebanyak 17.1% kepada 19.2% diperhatikan daripada jumlah input air yang diambil kira dalam kedua-dua musim. Hasil dari plot eksperimentasi menunjukkan 2.5 t / ha dan 2.7 t / ha bagi luar musim dan musim utama. Akhirnya, indeks produktiviti air dianalisis sebanyak 0.72 kg m⁻³ semasa luar musim dan 0.78 kgm⁻³ semasa musim utama.

Berdasarkan analisis pengangkutan bahan larut, jumlah nitrogen (TN) yang terkumpul sebanyak 50.3% hingga 49.7% dianggarkan dalam lapisan tanah sedalam 40 cm manakala 49.7% hingga 53% TN bahan larut lesap diperoleh di bawah lapisan tanah sedalam 40 cm (40-100 cm) semasa luar musim dan musim utama. Kira-kira 85% daripada kehilangan pengurusan N adalah dalam bentuk NO₃⁻; namun, masih ada sejumlah besar NO₃⁻ yang kekal di bawah zon akar yang menyumbang air bawah tanah. Jumlah kehilangan pengurusan T-N adalah 34.9 dan 27.9 kg / ha masing-masing pada luar musim dan musim utama. Anggaran kehilangan jumlah fosforus selama dua musim penanaman padi masing-masing adalah 3 dan 1.7 kg / ha. Jumlah kerugian T-N, T-P dan K melalui saliran ialah 27.7 dan 18.5, 2.2 dan 1.1, 5.9 dan 3.5 kg / ha semasa luar musim dan musim utama.

Hidrus-1D digunakan untuk mensimulasikan pergerakan air dan bahan larut di bawah kedalaman tanah yang berbeza dari 20, 40, 60, 80 dan 100 cm dalam eksperimen persekitaran padi sebenar. Aliran air yang disimulasi dan diperhatikan dan pengurusan nutrien adalah sepadan ($R^2 = 0.98$, $RMSE = 0.24$). Simulasi Hydrus-1D menunjukkan corak pergerakan air dan bahan larut yang sama di bawah kedalaman tanah yang berbeza semasa tempoh kajian. Kepekatan N, P dan K yang diperhatikan dan disimulasikan dalam padi adalah tinggi disebabkan faktor persenyawaan dan iklim lain. Oleh itu, pengurangan kadar baja yang berlebihan terutamanya semasa tempoh awal penanaman padi dan penyesuaian teknik penjimatan air dapat mengurangkan risiko pencemaran sawah padi.

Analisis regresi dilakukan bagi pembangunan model penggunaan baja yang lebih baik. Analisis regresi linier berganda dilakukan untuk mengetahui hubungan antara EC dan bahan larut (N, P dan K) semasa kedua-dua musim. Analisis regresi polinomial dipasang untuk menilai sama ada atau tidak perubahan EC mempunyai kesan terhadap kepekatan N, P dan K dalam sawah padi. Akhirnya, model empirikal ditubuhkan bagi menganggar kepekatan N, P dan K menggunakan dua data musim penanaman padi. Program penyelesaian MS Excel digunakan untuk membangunkan model empirikal. Hasil kajian memperoleh persefahaman yang kuat antara N, P, dan K yang diperhatikan dan yang dianggarkan dengan koefisien penentuan (R^2) 0.91 dan 0.95 semasa kedua-dua musim. Oleh itu, model-model ini boleh dikatakan berguna dalam meramal perubahan konsentrasi bahan larut dalam zon akar dan di bawah zon akar sepanjang musim penanaman padi untuk penggunaan baja yang lebih baik.

ACKNOWLEDGEMENTS

In the name of Allah, Most Gracious, Most Merciful, all praise and thanks are due to Allah, and Peace and blessings are upon his messenger. Although I had undergone many obstacles and queries throughout the completion, I am so glad that my Ph.D. project was carried out and conducted successfully by the help of Allah the Almighty.. I would like to thank my committee members who were more than generous with their expertise and precious time. A special thank to Dr. Md. Rowshon Kamal, my committee chairman for his endless support, invaluable guidance, constant encouragement, and supervision throughout completion of the project. I should consider myself fortunate to have him as my guide. Thank you Dr. Aimrun Wayakok and Assoc. Prof. Dr. Hasfalina Bin Che Man for agreeing to serve on my committee, without whose extraordinary help, I would not have been able to accomplish my task. Equally worthy of mention is the encouragement received from Dr. Amimul Ahsan who was initially part of my supervisory committee but left from the University during the study period.

I would like to acknowledge and thank the Government of Malaysia for financial support through the Special Graduate Research Allowance (S-GRA) program, in the Ministry of Science, Technology and Innovation (MOSTI). Sincere appreciation is due to the owner of the experimental plot Mr. Zulkifly Bin Sulaiman and his family for allowing us to conduct our experiment smoothly and providing the necessary help. Heartfelt thanks to the staff of the Department of Biological and Agricultural Engineering for providing the numerous equipment, instruments and moral support for this research work. I would also like to thank Malaysia Meteorology Department for providing us weather data. Also thanks to Engr. Hafiz and Engr. Sabri and the lab staff for their kind assistance during soil and water sample tests. I would like to acknowledge my cousin Engr. Abdirasak Ali, who suffered a serious head injury while he was helping me, during my fieldwork. We wish you healthy and full recovery very soon. I am also grateful to my colleagues and friends who supported me during my data collection.

Without my parents and siblings, it would not have been easy and possible for me to achieve this great work. Thank you my parents and two elder sisters Naima Abdullahi and Hamdi Abdullahi for financial support, unconditional love and care. I love them so much, and I would not have made it this far without them. Sincere gratitude goes to my parents, siblings and friends for their support, love, affection and encouragement.

I certify that a Thesis Examination Committee has met on 27 June 2018 to conduct the final examination of Abdikani Abdullahi Mo'allim on his thesis entitled "Modeling Solute Transport for Improved Fertiliser Use in Rice Production System" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Azmi bin Dato Haji Yahya, PhD

Professor Ir.
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Abdul Rashid bin Mohamed Sharif, PhD

Professor Sr.Gs.
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Desa bin Ahmad, PhD

Professor Ir.
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Md. Abdul Mojid, PhD

Professor
Bangladesh Agricultural University
Bangladesh
(External Examiner)



RUSLI HAJI ABDULLAH, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 30 August 2018

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Md Rowshon Kamal, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Aimrun Wayayok, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Hasfalina Bt. Che Man, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No.: Abdikani Abdullahi Mo'allim, GS41789

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Dr. Md Rowshon Kamal

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Aimrun Wayayok

Signature: _____
Name of Member
of Supervisory
Committee: Associate Professor
Dr. Hasfalina Bt. Che Man

TABLE OF CONTENTS

		Page
ABSTRACT		i
ABSTRAK		iii
ACKNOWLEDGEMENTS		v
APPROVAL		vi
DECLARATION		viii
LIST OF TABLES		xiv
LIST OF FIGURES		xvii
LIST OF ABBREVIATIONS		xxix
LIST OF SYMBOLS		xxx
CHAPTER		
1	INTRODUCTION	1
1.1	Background of the Research	1
1.2	Problem Statements	5
1.3	Aim and Objectives	6
1.4	Scope of the Study	6
1.5	Limitations on the Scope of the Study	6
1.6	Organization of the Thesis	7
2	LITERATURE REVIEW	8
2.1	Challenges of Rice Production in Malaysia	8
2.1.1	History of Rice Production	8
2.1.2	Agricultural Practices	9
2.1.3	Fertilizer Uses in Rice Production System	10
2.1.4	Environmental Impacts	11
2.1.5	Irrigation Practices	12
2.2	Estimation of Water Balance Components from Paddy Fields	13
2.2.1	Water Balance Model	13
2.2.1.1	Application of Water Balance Model	14
2.2.2	Crop Evapotranspiration (ET)	15
2.2.2.1	Computation of Reference Crop Evapotranspiration (ET ₀)	16
2.2.2.2	Application of FAO Penman-Monteith Method	16
2.2.3	Seepage and Deep Percolation (SP and DP)	17
2.2.4	Water Requirements	17
2.3	Solute Transport Phenomena in Paddy Fields	18
2.3.1	Nitrogen Transport	19
2.3.2	Phosphorus Transport	20
2.3.3	Pesticide Concentrations	22
2.4	Some Water Quality Parameters	23
2.4.1	pH and Temperature	23

2.4.2	Electrical Conductivity (EC) and Total Suspended Solids (TDS)	23
2.4.3	Dissolved Oxygen (DO)	24
2.5	Water Pollution	25
2.5.1	Sources of Water Pollution	26
2.5.2	Point Source Pollution	26
2.5.3	Non-point source pollution	27
2.6	Methods of Solutes Reduction from Agricultural Lands	27
2.6.1	Water Saving Irrigation (WSI)	27
2.6.2	Drainage Control	29
2.6.3	Recycling irrigation system	30
2.7	Applications of Numerical Models for Solute Transport Studies	32
2.7.1	LEACHM Model	32
2.7.2	SWAP Model	33
2.7.3	PCPF-1 model	34
2.7.4	VS2DT model	34
2.7.5	HYDRUS-1D model	35
2.8	Summary	37
3	MATERIALS AND METHODS	38
3.1	Introduction	38
3.2	Location of the Study Area	39
3.2.1	History and General Information of the Scheme	40
3.2.2	Climate Patterns	42
3.2.3	Other Climatic Features of the Study Area	42
3.2.4	Reference Crop Evapotranspiration (ET _o)	46
3.2.5	Soils	47
3.2.6	Water Resources	47
3.2.7	Existing Development and Current Practices	48
3.2.7.1	Main Conveyance System	48
3.2.7.2	Tertiary Distribution and Drainage Network	51
3.2.7.3	Current Cropping Practice and Irrigation Water Management	51
3.2.7.4	Current Water Management Issues	52
3.2.8	Site Visits and Meetings with Scheme Management	53
3.3	Layout of Equipment Installed in a Paddy Field	55
3.4	Monitoring Water Balance Components in Paddy Field	57
3.4.1	Water Balance Model in a Paddy Field	57
3.4.1.1	Irrigation Supply (IR)	60
3.4.1.2	Rainfall (RF)	61
3.4.1.3	Evapotranspiration (ET)	62
3.4.1.4	Seepage-Percolation (SP)	63
3.4.1.5	Drainage Flow	64
3.4.1.6	Ponding Water Depth	65
3.4.1.7	Calculation of Irrigation Performance Indices	66
3.4.2	Downloading of Data from Datalogger	67

	3.4.2.1	Irrigation Supply	67
	3.4.2.2	Field Drainage	68
	3.4.2.3	Fluctuation Water Level in a Paddy Plot	69
	3.4.2.4	Rainfall	69
3.5		Monitoring Solute Transport (N, P, K) in Rice Production	70
	3.5.1	Collection of Leachate from Paddy Field	70
	3.5.2	Analysis of Paddy Field's Leachate in Laboratory	72
	3.5.3	Analysis of Nutrients Loss (N, P and K)	73
3.6		Dynamics of Solutes (N, P and K) Transport using 1-D Hydrus Numerical Model	74
	3.6.1	Hydrus-1D model	74
	3.6.2	Water Flow	74
	3.6.3	Nitrogen Transport and Transformations	75
	3.6.4	Model Parameters	76
	3.6.5	Estimation of Soil Hydraulic Parameters	76
	3.6.6	Root Water Uptake	76
	3.6.7	Solute Transport and Reaction/Transformation Parameters	76
	3.6.8	Initial and boundary conditions	77
	3.6.9	Model evaluation criteria	77
3.7		Development of Empirical Models for Improved Fertilizers Use	78
4		RESULTS AND DISCUSSION	79
4.1		Soil and Climate in the Study Plot	79
	4.1.1	Soil Conditions	79
	4.1.2	Climate Conditions	80
4.2		Analysis of Water Balance Components	81
	4.2.1	Evapotranspiration (ET)	82
	4.2.2	Rainfall (RF) and Effective Rainfall (ER)	90
	4.2.3	Irrigation Supply (IR)	93
	4.2.4	Drainage Requirements (DR)	96
	4.2.5	Seepage and Percolation (SP)	98
	4.2.6	Fluctuation of Standing Water Depth (SW)	100
	4.2.7	Crop and irrigation water requirements during two consecutive rice growing season	102
	4.2.8	Water productivity index (WPI)	103
	4.2.9	Summary of Water balance component analysis	104
4.3		Solutes (N, P and K) Losses from Paddy Field	108
	4.3.1	Solutes Concentration in Flooding water	108
	4.3.2	Solutes Concentration in Irrigation Water	111
	4.3.3	Solutes (N, P and K) Concentration in Drainage Water	114
	4.3.4	Solutes (N, P and K) Concentration in Subsurface Water	117
	4.3.5	Leaching Losses of solutes (NPK) from Paddy Field during two rice growing season	129
	4.3.6	Nutrient losses via drainage during off and main seasons	147

4.3.7	Summary	148
4.4	Simulation of water and solute movement using Hydrus-1D model	149
4.4.1	Water movement simulation analysis	149
4.4.1.1	Model assessment	149
4.4.1.2	Pressure head	150
4.4.1.3	Water content	152
4.4.1.4	Root water uptake	154
4.4.1.5	Water flux	156
4.4.2	Solute transport simulation analysis	159
4.4.2.1	Model assessment	159
4.4.2.2	Spatio-temporal dynamics of solutes	160
4.4.2.3	Simulation of solute fluxes at the bottom of the soil layer	169
4.5	Model development	171
4.5.1	Development of EC models	171
4.5.2	Development of N, P and K models	178
4.5.2.1	Based on EC data	178
4.5.2.2	Nutrients Availability in Field with Time	185
4.5.3	Guidelines for developing of Decision support system	196
4.5.4	Summary	198
5	CONCLUSIONS AND RECOMMENDATIONS	202
5.1	Conclusion	202
5.2	Recommendations for Future Research Works	203
	REFERENCES	205
	APPENDICES	232
	BIODATA OF STUDENT	234
	LIST OF PUBLICATIONS	235

LIST OF TABLES

Table	Page
2.1 Fertilizer use practices in the unadjusted farmer sustainability index (FSI) based on rice check 2014	11
3.1 General Features of Tanjung Karang Rice Irrigation Scheme	41
3.2 Irrigation Command Areas by Compartments of the Scheme	52
3.3 Crop coefficient Kc values for rice	63
3.4 Solutes reaction parameters	77
4.1 Physical properties of the soil in the experimental field during off-season (January – April 2017)	79
4.2 Physical properties of the soil in the experimental field during Main season (July – October 2017)	79
4.3 Chemical properties of the soil in the experimental field during off-season (January – April 2017)	80
4.4 Chemical properties of the soil in the experimental field during Main season (July – October 2017)	80
4.5 Mean monthly weather parameters during two rice growing seasons	81
4.6 Summary of water productivity index, irrigation water requirement and crop water requirement during both seasons	104
4.7 Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP and K in flooding water depth in a paddy field during the off-season	109
4.8 Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP and K in flooding water depth in a paddy field during the main season	110
4.9 Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP, and K in irrigation water during off-season	112
4.10 Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP, and K in irrigation water during main season	113
4.11 Concentration of T-N, NO ₃ ⁻ , NH ₄ ⁺ , T-P, and K in drainage water during off-season	115
4.12 Concentration of T-N, NO ₃ ⁻ , NH ₄ ⁺ , T-P, and K in drainage water during main season	115

4.13	Concentration of Total Nitrogen in Sub-surface water during off-season	119
4.14	Concentration of Total Nitrogen in Sub-surface water during main season	120
4.15	Concentration of Nitrate Nitrogen in Sub-surface water during off-season	121
4.16	Concentration of Nitrate-N in subsurface water during main season	122
4.17	Concentration of Ammonium Nitrogen in subsurface water during off-season	123
4.18	Concentration of Ammonium Nitrogen in subsurface water during main season	124
4.19	Concentration of Total Phosphorous in subsurface water during off-season	125
4.20	Concentration of Total Phosphorous in subsurface water during main season	126
4.21	Concentration of Potassium in subsurface water during off-season	127
4.22	Concentration of Potassium in subsurface water during main season	128
4.23	Cumulative of Total Nitrogen under different soil depths during off-season	131
4.24	Cumulative of Total Nitrogen under different soil depths during main season	132
4.25	Cumulative of Nitrate-N under different soil depths during off-season	132
4.26	Cumulative of Nitrate-N under different soil depths during main season	133
4.27	Cumulative of Ammonium-N under different soil depths during off-season	133
4.28	Cumulative of Ammonium-N under different soil depths during main season	134
4.29	Cumulative of Total Phosphorous under different soil depths during off-season	134
4.30	Cumulative of Total Phosphorous under different soil depths during main season	135

4.31	Cumulative of Total Potassium under different soil depths during off-season	135
4.32	Cumulative of Total Potassium under different soil depths during main season	136
4.33	Optimized values of soil hydraulic parameters	150
4.34	Statistical parameters of the model evaluation	160
4.35	Developed multi-linear equations of EC in surface, drainage and subsurface water during main season	199
4.36	The performance of developed equations of T-N, T-P, and K in flooding water based on EC	199
4.37	The performance of developed equations of T-N, T-P, and K in drainage water based on EC	199
4.38	The performance of developed equations of T-N, T-P, and K in subsurface water based on EC	200
4.39	The performance of developed equations of T-N, T-P, and K in flooding water based on fertilizer application date	200
4.40	The performance of developed equations of T-N, T-P, and K in subsurface water based on fertilizer application date	201

LIST OF FIGURES

Figure		Page
2.1	Global fertilizer (N+P ₂ O ₅ +K ₂ O) consumption	11
2.2	Suggested Crop Coefficient (k _c) Values for Rice (MR84 variety)	15
3.1	Flowchart for the Development of the improved fertilizer use for rice production using intensive field investigation and laboratory analysis	38
3.2	Location of the Study Area at Integrated Agriculture Authority, Kula Selangor	39
3.3	Location of the experimental plot at Tanjung Karang rice irrigation scheme (TAKRIS) Malaysia	40
3.4	Mean Monthly Variations of Minimum, Average and Maximum Rainfall (1991-2010) in the Project Area	42
3.5	General Mean Monthly Climatic Conditions at MARDI Tg. Karang Station	43
3.6	Recorded Mean Monthly Pan Evaporation in the Project Area (1991-2010)	44
3.7	Recorded Mean Monthly Temperature (°C) in the Project Area (1991-2010)	44
3.8	Recorded Mean Monthly Relative Humidity (RH) in the Project Area (1991-2010)	45
3.9	Recorded Mean Monthly Observed Sunshine Duration in the Project Area (1991-2010)	45
3.10	Recorded Mean Monthly Sunshine Duration in the Project Area (1991-2010)	46
3.11	Daily Reference Crop Evapotranspiration (ET _o) using FAO Penman-Monteith Method	47
3.12	Recorded Monthly Discharges at Bernam River Headwork (1991-2002)	48
3.13	Overall Water Resources Configuration of the Tanjung Karang Irrigation Scheme	49
3.14	Diversion for irrigation supply to feeder canal at Bernam River Headworks (BRH)	50

3.15	Irrigation control structure in the main canal at Sg. Burong	50
3.16	Tertiary canal with adjustable off-take for distributing water from the Main Canal to the fields	51
3.17	Photograph with DID and DOA Staff at the IADA Irrigation Scheme, Kuala Selangor	54
3.18	Layout of the equipment installation for the field investigation of water balance components and solute transport in a Paddy Field	55
3.19	Installed equipment for monitoring solute transport and components of water balance model in a paddy field	56
3.20	Schematic diagram of water balance component in paddy fields	58
3.21	Typical view of the paddy field after installation of the equipment	59
3.22	Monitoring irrigation water supply at 1-minute interval by RBC flume with flow sensor and datalogger in the experimental plot	60
3.23	Monitoring rainfall at 10-minutes interval by RBC flume with sensor and datalogger in the experimental plot	61
3.24	Monitoring daily crop evapotranspiration using ET gage and field lysimeters with Marriott tube in the experimental plot	62
3.25	Monitoring daily seepage-percolation using field lysimeters and Marriott tube in the experimental plot	64
3.26	Monitoring drainage flow at 1-minute interval from field using Flow Sensor and Data Logger during rice growing season at the experimental plot	65
3.27	Monitoring of flooding water depth at 10-minutes interval in the experimental plot during rice growing season	66
3.28	Typical view of downloading recorded data from a field device at the experimental plot	67
3.29	Typical view of irrigation data downloading process	68
3.30	Typical view of drainage data downloading process	68
3.31	Typical view of water level data downloading process	69
3.32	Typical view of rainfall data downloading process	69
3.33	Porous cup Soil water samplers installed at 20 cm, 40 cm, 60 cm, 80 cm and 100 cm in a paddy field	70

3.34	Typical field view of five porous cups soil water samplers installed at 20 cm, 40 cm, 60 cm, 80 cm and 100 cm in a paddy field	70
3.35	Typical field view of five porous cups soil water samplers installed at 20 cm, 40 cm, 60 cm, 80 cm and 100 cm in a paddy field	71
3.36	Water samples collected from the paddy field and its subsurface at 20 cm incremental depths	72
3.37	DR/890 colorimeter for NPK measurements during the experimental period	73
4.1	Monthly used water balance components in a paddy field during two consecutive rice growing seasons in 2017	81
4.2	Daily measured crop evapotranspiration in a paddy plot during the off-season (January to April 2017)	82
4.3	Daily measured crop evapotranspiration in a paddy plot during the main season (July to October 2017)	83
4.4	Daily crop evapotranspiration in a paddy plot during the off-season using CropWat 8.0 (January to April 2017)	84
4.5	Daily crop evapotranspiration in a paddy plot during the main season using CropWat 8.0 (July to October 2017)	84
4.6	Daily computed reference evapotranspiration using CropWat 8.0 during the off-season (January to April 2017)	85
4.7	Daily computed reference evapotranspiration using CropWat 8.0 during the main season (July to October 2017)	86
4.8	Comparison of daily measured crop evapotranspiration in a paddy plot during two rice growing seasons (January to October 2017)	87
4.9	Comparison of weekly measured crop evapotranspiration in a paddy plot during two rice growing seasons (January to October 2017)	87
4.10	Comparison of daily crop evapotranspiration in a paddy plot during two rice growing seasons using CropWat 8.0 (January to October 2017)	88
4.11	Comparison of weekly crop evapotranspiration in a paddy plot during two rice growing seasons using CropWat 8.0 (January to October 2017)	88
4.12	Comparison of daily computed reference evapotranspiration using CropWat 8.0 during two rice growing season (January to October 2017)	89

4.13	Comparison of weekly computed reference evapotranspiration using CropWat 8.0 during two rice growing season (January to October 2017)	89
4.14	Daily observed rainfall using data-logger during the off-season (January to April 2017)	90
4.15	Daily observed rainfall using data-logger during the main season (July to October 2017)	91
4.16	Comparison of daily variation of rainfall during two rice growing season (January to October 2017)	91
4.17	Comparison of weekly variation of rainfall during two rice growing season (January to October 2017)	92
4.18	Monthly effective rainfall during two rice irrigation seasons in 2017	92
4.19	Observed daily irrigation water supply during the off-season	94
4.20	Observed daily irrigation water supply during the main season	94
4.21	Estimated irrigation water volume at weekly interval during rice growing periods in both seasons	95
4.22	Estimated irrigation water volume at 10-days interval during rice growing periods in both seasons	95
4.23	Recorded daily drainage water from a paddy field during the off-season	96
4.24	Recorded daily drainage water from a paddy field during the main season	97
4.25	Estimated the volume of drainage flow at weekly interval during rice growing period for both seasons	97
4.26	Estimated the volume of drainage flow at 10-days interval during rice growing period for both seasons	98
4.27	Measured daily seepage and percolation water from a paddy field during the off-season	99
4.28	Measured daily seepage and percolation water from a paddy field during the main season	99
4.29	Comparison of weekly deep percolation water during two rice growing season (January to October 2017)	100

4.30	Observed daily fluctuation of standing water depth during the off-season	101
4.31	Observed daily fluctuation of standing water depth during the main season	101
4.32	Estimated crop water requirement for a paddy plot during both rice growing seasons in 2017	103
4.33	Observed weekly irrigation, drainage and rainfall water during the off-season (January to April 2017)	105
4.34	Observed weekly irrigation, drainage and rainfall water during the main season (July to October 2017)	105
4.35	Observed monthly irrigation, drainage and rainfall water during the off-season (January to April 2017)	106
4.36	Observed monthly irrigation, drainage and rainfall water during the main season (July to October 2017)	106
4.37	Observed weekly irrigation, drainage, evapotranspiration, seepage percolation and rainfall water during the off-season (January to April 2017)	107
4.38	Observed weekly irrigation, drainage, evapotranspiration, seepage percolation and rainfall water during the main season (July to October 2017)	107
4.39	Observed monthly water balance components during the two rice growing seasons (January to October 2017)	108
4.40	Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP, and K in flooding water depth during off-season	111
4.41	Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP, and K in flooding water depth during main season	111
4.42	Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP, and K in irrigation water during off-season	113
4.43	Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP, and K in irrigation water during main season	114
4.44	Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP and K in drainage water during the off-season	116
4.45	Concentration of TN, NO ₃ ⁻ , NH ₄ ⁺ , TP and K in drainage water during the main season	116

4.46	Dynamic characteristics of Total-N in subsurface water during the off-season	119
4.47	Dynamic characteristics of T-N in subsurface water during the main season	120
4.48	Dynamic characteristics of Nitrate-N in subsurface water during the off-season	121
4.49	Dynamic characteristics of Nitrate-N in subsurface water during the main season	122
4.50	Dynamic characteristics of Ammonium-N in subsurface water during the off-season	123
4.51	Dynamic characteristics of Ammonium-N in subsurface water during the main season	124
4.52	Dynamic characteristics of Total-P in subsurface water during the off-season	125
4.53	Dynamic characteristics of Total-P in subsurface water during the main season	126
4.54	Dynamic characteristics of K in subsurface water during the off-season	127
4.55	Dynamic characteristics of K in subsurface water during the main season	128
4.56	Cumulative of Total-N within root zone (0-40 cm) during the off-season	136
4.57	Cumulative of Total-N within root zone (0-40 cm) during the main season	137
4.58	Cumulative of Total-N below root zone (60-100 cm) during the off-season	137
4.59	Cumulative of Total-N below root zone (60-100 cm) during the main season	138
4.60	Cumulative of Nitrate-N within root zone (0-40 cm) during the off-season	138
4.61	Cumulative of Nitrate-N within root zone (0-40 cm) during the main season.	139
4.62	Cumulative of Nitrate-N below root zone (60-100 cm) during the off-season.	139

4.63	Cumulative of Nitrate-N below root zone (60-100 cm) during the main season	140
4.64	Cumulative of Ammonium-N within root zone (0-40 cm) during the off-season	140
4.65	Cumulative of Ammonium-N within root zone (0-40 cm) during the main season	141
4.66	Cumulative of Ammonium-N below root zone (60-100 cm) during the off-season	141
4.67	Cumulative of Ammonium-N below root zone (60-100 cm) during the main season	142
4.68	Cumulative of Total-P within root zone (0-40 cm) during the off-season	142
4.69	Cumulative of Total-P within root zone (0-40 cm) during the main season	143
4.70	Cumulative of Total-P below root zone (60-100 cm) during the off-season	143
4.71	Cumulative of Total-P below root zone (60-100 cm) during the main season	144
4.72	Cumulative of Potassium within root zone (0-40 cm) during the off-season	144
4.73	Cumulative of Potassium within root zone (0-40 cm) during the main season	145
4.74	Cumulative of Potassium below root zone (60-100 cm) during the off-season	145
4.75	Cumulative of Potassium below root zone (60-100 cm) during the main season	146
4.76	Leaching losses of T-N, NO ₃ ⁻ , NH ₄ ⁺ , T-P, and K during two rice growing seasons	146
4.77	Drainage losses of T-N, NO ₃ ⁻ , NH ₄ ⁺ , T-P, and K during two rice growing seasons	148
4.78	Simulated pressure heads at different soil depths 20, 40, 60, 80 and 100 cm during the off-season	151
4.79	Simulated pressure heads at different soil depths 20, 40, 60, 80 and 100 cm during the main season	151

4.80	Simulated and observed soil water content within root zone (0-40 cm) during the off-season	152
4.81	Simulated and observed soil water content below root zone (60-100 cm) during the off-season	153
4.82	Simulated and observed soil water content within root zone (0-40 cm) during the main season	153
4.83	Simulated and observed soil water content below root zone (60-100 cm) during the main season	154
4.84	Simulated actual daily water uptake and simulated cumulative water uptake during the off-season	155
4.85	Simulated actual daily root water uptake and simulated cumulative water uptake during the main season	155
4.86	Simulated water fluxes within root zone (0-40 cm) during the off-season	157
4.87	Simulated water fluxes below root zone (60-100 cm) during the off-season	157
4.88	Simulated water fluxes within root zone (0-40 cm) during the main season	158
4.89	Simulated water fluxes below root zone (60-100 cm) during the main season	158
4.90	Comparison between simulated and observed NO_3^- concentration within root zone (0-40 cm) during the off-season	161
4.91	Comparison between simulated and observed NO_3^- concentration below root zone (60-100 cm) during the off-season	162
4.92	Comparison between simulated and observed NH_4^+ concentration within root zone (0-40 cm) during the off-season	162
4.93	Comparison between simulated and observed NH_4^+ concentration below root zone (60-100 cm) during the off-season	163
4.94	Comparison between simulated and observed T-P concentration within root zone (0-40 cm) during the off-season	163
4.95	Comparison between simulated and observed T-P concentration within below zone (60-100 cm) during the off-season	164
4.96	Comparison between simulated and observed K concentration within root zone (0-40 cm) during the off-season	164

4.97	Comparison between simulated and observed K concentration below root zone (60-100 cm) during the off-season	165
4.98	Comparison between simulated and observed NO ₃ ⁻ concentration within root zone (0-40 cm) during the main season	165
4.99	Comparison between simulated and observed NO ₃ ⁻ concentration below root zone (60-100 cm) during the main season	166
4.100	Comparison between simulated and observed NH ₄ ⁺ concentration within root zone (0-40 cm) during the main season	166
4.101	Comparison between simulated and observed NH ₄ ⁺ concentration below root zone (60-100 cm) during the main season	167
4.102	Comparison between simulated and observed T-P concentration within root zone (0-40 cm) during the main season	167
4.103	Comparison between simulated and observed T-P concentration below root zone (60-100 cm) during the main season	168
4.104	Comparison between simulated and observed K concentration within root zone (0-40 cm) during the main season	168
4.105	Comparison between simulated and observed K concentration below root zone (60-100 cm) during the main season	169
4.106	Simulated cumulative leaching fluxes of NO ₃ ⁻ , NH ₄ ⁺ , T-P and K at the bottom of the soil layer (100 cm) during the off-season	170
4.107	Simulated cumulative leaching fluxes of NO ₃ ⁻ , NH ₄ ⁺ , T-P and K at the bottom of the soil layer (100 cm) during the main season	170
4.108	Relationship between EC and T-N in flooding water during the rice growing season	172
4.109	Relationship between EC and T-P in flooding water during the rice growing season	172
4.110	Relationship between EC and K in flooding water during the rice growing season	173
4.111	Relationship between EC and T-N in drainage water during the rice growing season	173
4.112	Relationship between EC and T-P in drainage water during the rice growing season	174
4.113	Relationship between EC and K in drainage water during the rice growing season	174

4.114	Relationship between EC and T-N within root zone during the rice growing season	175
4.115	Relationship between EC and T-P within root zone during the rice growing season	175
4.116	Relationship between EC and K within root zone during the rice growing season	176
4.117	Relationship between EC and T-N below root zone during the rice growing season	176
4.118	Relationship between EC and T-P below root zone during the rice growing season	177
4.119	Relationship between EC and T-P below root zone during the rice growing season	177
4.12	Relationship between T-N and EC in flooding water during the rice growing season	178
4.121	Relationship between T-P and EC in flooding water during the rice growing season	179
4.122	Relationship between K and EC in flooding water during the rice growing season	179
4.123	Relationship between T-N and EC in drainage water during the rice growing season	180
4.124	Relationship between T-P and EC in drainage water during the rice growing season	181
4.125	Relationship between K and EC in drainage water during the rice growing season	181
4.126	Relationship between T-N and EC within root zone during the rice growing season	182
4.127	Relationship between T-P and EC within root zone during the rice growing season	183
4.128	Relationship between K and EC within root zone during the rice growing season	183
4.129	Relationship between T-N and EC below root zone during the rice growing season	184
4.130	Relationship between T-P and EC below root zone during the rice growing season	184

4.131	Relationship between K and EC below root zone during the rice growing season	185
4.132	Observed and simulated T-N under different soil depths during the off-season	186
4.133	Observed and simulated T-P under different soil depths during the off-season	186
4.134	Observed and simulated K under different soil depths during the off-season	187
4.135	Observed and simulated of N, P and K in flooding water during the off-season	187
4.136	Observed and simulated T-N under different soil depths during the main season	188
4.137	Observed and simulated T-P under different soil depths during the main season	188
4.138	Observed and simulated K under different soil depths during the main season	189
4.139	Observed and simulated of N, P and K in flooding water during the main season	189
4.140	Observed and simulated T-N within root zone during the off-season	190
4.141	Observed and simulated T-P within root zone during the off-season	191
4.142	Observed and simulated K within root zone during the off-season	191
4.143	Observed and simulated T-N below root zone during the off-season	192
4.144	Observed and simulated T-P below root zone during the off-season	192
4.145	Observed and simulated K below root zone during the off-season	193
4.146	Observed and simulated T-N within root zone during main rice growing season	193
4.147	Observed and simulated T-P within root zone during main rice growing season	194
4.148	Observed and simulated K within root zone during main rice growing season	194
4.149	Observed and simulated T-N below root zone during main rice growing season	195

4.150	Observed and simulated T-P below root zone during main rice growing season	195
4.151	Observed and simulated K below root zone during main rice growing season	196
4.152	Architecture of Web Paddy GIS	197
4.153	Flow chart of DSS in Web Paddy GIS	198



LIST OF ABBREVIATIONS

MOA	Ministry of Agriculture
IPCC	Intergovernmental Panel on Climate Change
DOE	Department of Environment
SSL	self-sufficiency level
FAO	Food and Agriculture Organization
ICID	International Commission of Irrigation and Drainage
IRRI	International Rice Research Institute
ET	Evapotranspiration
ET _o	Reference Evapotranspiration
RF	Rainfall
DR	Drainage
WL	Water Level
IR	Irrigation
K _c	Crop Coefficient
Mg/L	Milligram per litter
Kg/ha	Kilogram per hectare
mm	Millimetre
cm	centimetre
mm day ⁻¹	Millimetre per day
cm day ⁻¹	Cent meter per day

LIST OF SYMBOLS

ρ	Density
Δ	slope of the saturation vapour pressure temperature relationship
γ	psychrometric constant
α	albedo or canopy reflection coefficient
ω	sunset hour angle
ϕ	latitude
δ	solar declination
σ	Stefan-Boltzman constant
$e^{\circ}(T_{\min})$	saturation vapour pressure at daily minimum temperature
$e^{\circ}(T_{\max})$	saturation vapour pressure at daily maximum temperature
Δt	length of time interval
Δz	effective soil depth

CHAPTER 1

INTRODUCTION

1.1 Background of the Research

Rice is the main food crop in Malaysia. Rice production of the country has achieved 72% self-sufficiency level (SSL) with an annual production of 3.5 million tonnes a year (MOA, 2017). The Agriculture and Agro-based Industry Ministry targets the country to achieve a 100 percent self-sufficiency level (SSL) in paddy production by 2020. Irrigation is crucial to the world's food grain production because 40% of all crops and close to 60% of cereal production comes from irrigated agriculture (Phogat et al. 2010), even though irrigated lands comprise only 20% of the arable land (FAO, 2004). In Asia, irrigated agriculture uses 90% of the total freshwater, and more than half of this irrigates rice. About 75% of the global rice volume is produced in the irrigated low lands (Cantrell 2004). There are an estimated 150 million hectares of rice lands worldwide, 50% of which are irrigated, usually with continuous flooding for most of the crop season (Ghosh and Bhat 1998). In many irrigated areas, rice is grown as a monoculture with two rice crops every year. Global water and food security are one of the most important challenges in the 21st century to supply sufficient food for the increasing population while sustaining the stressed environment threatened by climate change. One of the main causes of land degradation is intensifying chemical fertilizer and pesticide usage (Bagheri et al. 2008; Bala et al. 2014 and Jean et al. 2015). In addition, a policy implemented by the government in term of subsidies is one of the encouragement factors for paddy farmers to adopt chemical substance in paddy cultivation techniques.

Rice production is one of the major non-point sources of fertilizers and pesticide pollution in Malaysia. Monitoring of these fertilizers and pesticide concentrations in river systems detected a number of chemicals commonly used in paddy fields, and these concentrations may appear to have adverse effects on the aquatic ecosystem. Most of the paddy fields are treated with fertilizers, herbicides, fungicides and insecticides which are applied during the crop season accordingly. Therefore, rice production is one of the major nonpoint sources of pollution. The typical paddy field in Malaysia is susceptible to fertilizers, pesticides and herbicides runoff since the chemicals are applied directly onto paddy water. Due to excessive use of chemicals in rice production in Malaysia compared to other Asian countries, it may presume that a significant amount of runoff along with fertilizers losses is occurring from paddy fields. Pesticides runoff losses from paddy fields range from a few percent to more than 50% of the applied amount depending on the water management (Maru, 1991 and Sudo et al., 2002). Inao et. al. (2008) highlighted that it is important to develop and validate mathematical models adapted to paddies in the Asian region in order to establish a realistic assessment and management procedure for environmentally-friendly rice production. Therefore, water quantity and distribution

during the growing season is one of the key factors controlling crop growth and nutrients uptake in rice production.

There is increasing pressure on primary producers to reduce the environmental impacts of agricultural production. However, agricultural crop production systems are inherently sprinkling with respect to nutrients. Therefore, the challenge to producers is to manage their crop production systems in order to minimize environmental losses of nutrients, while achieving crop yield and quality targets. Many strategies have been developed in recent years to meet this challenge (Zebarth et al., 2009). These include: development of new tools to measure crop N status in order to refine in-season fertilizer N management, development of new soil N tests to improve prediction of soil N supply, development of new fertilizer N products with release patterns more closely matched to crop N uptake patterns, and development of site-specific N management strategies. A number of studies highlighted that water contaminated by nitrate is not potable and at high concentrations can be a serious risk for human health (Al-Redhaiman 2000; Anjana and Iqbal, 2007). Moreover, the water industry must bear additional costs to remove nitrates from groundwater sources (Harris et al. 1992; Cameron and Schipper 2010). The detrimental impacts of nitrate loss from the soil have toxicological implications for animals and humans (Camargo and Alonso, 2006), and also on the environment leading to the eutrophication of freshwater (London, 2005), and marine ecosystems (Beman et al. 2005). In rice production, the mineral commercial fertilizers anhydrous ammonia, urea, ammonium sulfate and ammonium nitrate are commonly applied. They are particularly soluble for easy assimilation by crops. Both urea and ammonia are converted to nitrate at different rates depending on the nature of the soil and of the climatic conditions, thus leading to various loss mechanisms either by volatilization for ammonia or runoff for nitrate or urea after heavy rainfall and leaching into groundwater (Vitosh et al. 1995; Jarvis et al. 2011). The most effective ways of improving the efficiency of fertilizers (N, P and K) use in agricultural crop production is the matching the supply of fertilizers to the crop demand in both space and time. This can be achieved by supplementing the supply of fertilizers from the soil with the appropriate rate and form of fertilizers, at the right time, and at the right location. While the concept is simple, this is difficult to achieve in practice due to substantial variation in both crop demand and in soil supply of fertilizers across years and among and within fields.

The increase in rice production can be achieved by efficient and good agricultural management practices, water and nutrient inputs. However, the use of conventional practice with poor management of plant nutrients and excessive fertilizer application will have a major impact in efforts to increase rice production in Malaysia. The importance of fertilizers for achieving increase crop production must be emphasized. Many rice varieties, particularly the high-yielding improved varieties currently being introduced into many traditional farming systems throughout the world, respond markedly to fertilization. When used in conjunction with good management practices (thorough land preparation, controlled irrigation, timely weeding) fertilizers can increase yields many times over. On the other hand, if used improperly fertilizers can

damage crops, waste money, or possibly lead to a dependence on scarce chemical inputs.

Fertilizers are usually applied in Malaysia by conventional means. In paddy fields, the applied fertilizers, after dissolution, not only transport over and infiltrate into the soil, but also diffuse out and channel in all possible directions due to the transverse variation of water velocity and depth (Strelkoff et al. 2003). Deterioration of water quality in streams and lakes continues to be a significant issue in many counties (ICID, 2013). Causative factors include not only pollutants from various point sources but also those from non-point sources. The intensive use of agrochemicals for rice cultivation has been responsible for making paddy fields as a significant contributor of non-point source pollution. One of the most important factors contributing to this problem is the large amount of nutrients from non-point sources, especially agricultural paddy fields in which excessive chemical fertilizer application increases the concentration chemical loads (e.g. Nitrate nitrogen) in groundwater, which in turn results in the eutrophication of public water areas (Feng et al. 2004). Our understanding of nutrients/fertilizers transformation in paddy fields is limited due to complex interactions between soil, water, and biomass (Nakasone et al. 2004) and its behavior in relation to plant growth in paddy soils has been extensively studied (De Datta, 1986). Therefore, forecasting hydrological pathways and pollutants (nutrient and pesticides) behavior in paddy soil appear to be crucial in order to define the specific management practices controlling non-point source pollution and preserving water resources (Tournebize et al. 2006).

Water management is a difficult task for a large rice irrigation system. Different sub-systems, such as soil, water, climate, nutrients, plant, management systems and their complex dynamics work in the paddy field environment. Furthermore, an individual irrigation scheme has its physical and unique characteristics. The effects of climate change are significant on water demand for irrigation that is continuously being aggravated by unsustainable practices like over-use of chemical fertilizers and poor water management. Excessive irrigation deliveries generate a huge amount of return flows containing fertilizers, insecticides, and pesticides from paddy fields in Malaysia. Eventually, drainage water from paddy fields loses the essential agrochemicals and pollutes the surface water resources. Poor and uneven water distributions were often criticized as the major bottleneck in attaining efficient water use in rice irrigation systems in Malaysia (JICA and DID, 1998; Rowshon et al. 2009). Ramli et al. (2012) reported that the removal of fertilizers' subsidy reduced rice production and the self-sufficient level (SSL) in Malaysia and overuse of agrochemicals results in severe environmental problems. Therefore, optimum application of fertilizers is crucial to minimize their loss and reduce environmental pollution. In this regard, reuse of drainage water has a potential to play a vital role in profitable rice production.

Many endeavors have been made to define nutrient rates as ideas of economic advantages, including mass-balance (Scarf et al. 2006), economically ideal fertilizer rate (Neeteson and Wadman, 1987; Sawyer et al. 2006), maximum return to N

(Sawyer et al. 206) and information based ideal N compost (Zhu and Chen, 2002). Regardless, the vast majority of these suggestions addresses just site-specific ideal N rate. Such measures would be hard to reach out to a vast zone having distinctive rice assortments, environments, and cropping strategies. In addition, less consideration has been given as the measure of naturally ideal N rates. Our utilization of N manure has ended up far-reaching, bringing about serve natural issues (Chen et al. 2011). Hafeez et al. (2007) reported that rice production remained profitable despite high pumping costs to extract water from shallow tube wells for supplementary irrigation during dry seasons. In addition, small irrigation pumps owned by farmers can play an important role in capturing excess water that is drained from paddy fields to be reused in irrigation.

Excessive water and fertilizer inputs have led to a series of environmental problems in agricultural production areas around the world. Identifying the fates of water and nutrients is crucial to develop best management strategies in intensive agricultural production systems (Liang et al. 2018). Leon and Kohyama (2017) suggested that controlling and predicting nutrient losses into subsurface and surface water is vital for evaluating the environmental impacts of rice cultivation. Nutrient loss mechanism is complicated and shows remarkably regional differences due to spatial heterogeneities of underlying surface conditions, climate and agricultural practices (Zhang et al. 2016a). The regional heterogeneities of climate, underlying surface conditions and agricultural management practices also result in the remarkable regional differences of nutrient losses. A large number of applications also existed in the studies of hydrology and environmental observations or simulations, such as univariate analyses of extreme events and dam regulations (Zhang et al. 2016b).

Quantification of the amount of water used is very crucial for understanding and finding water use efficiency to an irrigation system level. Irrigation return flow consists of surface and subsurface flows. Water balance models, considering both components, can predict the return flow for re-use in paddy fields (Chowdary et al. 2004). Several mathematical models are available to describe water balance and behaviors of nutrients and pesticides in flooded rice fields. Some models describing the fate of nitrogen in rice fields focus on various processes taking place in flooded water: PADDY (Pesticide Paddy Field Model) (Inao and Kitamura 1999), PCPF-1 (Pesticide Concentration in Paddy Field, v.1) (Watanabe et al. 2007), RICEWQ (Rice Water Quality) (Williams et al. 1999; Karpouzas and Capri, 2006), and PADDIMOD (Jeon et al. 2005), while others describe mass transport in flooded water and the soil underneath. Chung et al. (2003) developed GLEAMS-PADDY model to describe nutrient loading in surface water and groundwater bodies. Chowdary et al. (2004), developed and applied a simple model for assessing the concentration of nitrates in water percolating out of the flooded rice fields. Tournebize et al. (2006), developed a coupled model (PCPF-SWMS) for simulating the fate and behavior of pollutants in water and soil of paddy fields. In GLEAMS-PADDY model, (Chung et al. 2003), N balance is separated into NH₄-N and NO₃-N balance and applied to ponding water and underlying soil. PADDIMOD (Jeon et al.

2005), describes N balance as the total inorganic N without focusing on NH₄-N and NO₃-N balance separately.

Through the literature, no study has been done yet in Malaysia for modeling solute transport for better fertilizer utilization towards improving water productivity and surface water pollution control in paddy fields. After application of fertilizers to the field, it dissolves, transports over the land surface and infiltrates into the soil by irrigation water. In deep soil layer, the HYDRUS-1D model can demonstrate appropriately the distribution of fertilizers through the gravitational one-dimensional flow (Taftah and Sepaskhah, 2012). Numerical modeling has an important significance for improving fertilization methods (Abbasi et al. 2003) and reducing agricultural non-point source pollution attributable to improper fertilization (Bradford and Katopodes, 1998). Thus, it's very important to establish an improved fertilizer use to the governmental authorities and farmers.

1.2 Problem Statements

Food security is key global challenges. By 2050 the world will need to increase crop production to feed its projected 9 billion people (FAO, 2009). Climate change brings a serious impact on water resources, which affects the food security. Therefore, the rice production system must enable to respond to challenges posed by the effects of climate change on precious land and water resources.

It is well understood that over fertilization is a major problem in intensive agricultural production areas, resulting in the enrichment of air, soil, and water with reactive nitrogen leading to the impairment of ecosystem functions. The over-fertilization entails unnecessary economic expenditure for farmers. Nitrogen loss from agricultural fields is the main cause of eutrophication. Few studies have determined the efficiency of N use and loss of paddy fields (Yang et al. 2014) and fewer studies have focused under flooded condition (Kiran et al. 2010). The excessive N and P fertilizer use with decreasing fertilizer use efficiencies in agriculture has resulted in large amounts of N and P elements entering ambient water bodies and the atmosphere through various means (Xing and Zhu 2000; Yoshinaga et al. 2007; Ni et al. 2007; Li et al. 2008). Meanwhile, the transport of agrochemicals from paddy fields pollutes the lakes and streams which lead to harm human health.

Due to the dynamic nature of paddy fields, N transport and transformation process are complex and difficult to understand. Quantification of water flow and nutrient losses in rice field becomes a challenge as rice is highly water and nitrogen demanding crop, thus, play key role for contributing pollutants in both surface and subsurface waters. In addition, rice field can lead a considerable nutrient loss by leaching under irrigated conditions and excess use of fertilizers. Indeed, no study is reported yet on this important aspect in Malaysia. Thus, this study focused to

develop the better fertilizer utilization for improving water productivity and surface water pollutant control.

1.3 Aim and Objectives

The main objective of this study is to develop the improved fertilizer use model for rice production through the investigation of intensive water and solutes balance model integrated with 1-D solute transport numerical model. The specific objectives are:

1. To analyze water balance components in modeling solute transports in growing paddy field.
2. To estimate nutrients (N, P, and K) loads and water quality parameters from paddy fields to surface and subsurface water systems.
3. To characterize the spatio-temporal dynamics of nutrients using the HYDRUS-1D numerical model in paddy fields, and
4. To develop empirical models for the better utilization of nutrients in a rice-production system using EC and fertilizer application date.

1.4 Scope of the Study

This study focused on the modeling and development of improved fertilizer use for rice production under real paddy environment at Tanjung Karang Rice Irrigation Scheme (TAKRIS). An intensive investigation of water and solutes losses from surface and subsurface water throughout two consecutive rice-growing seasons was conducted. It was also within the scope of the study to utilize Hydrus-1D model, calibrate and validate it with the aim of using the model to predict water and solutes movement under different soil depths in the experimental plot. Multiple and polynomial regression analysis were performed, and different empirical equations were developed to estimate EC, T-N, T-P and K in the experimental plot. The main focus of the field experiments was the development of empirical equations to predict nutrient (N, P and K) concentrations at the surface, subsurface and drainage water for the better use of fertilizers.

1.5 Limitations on the Scope of the Study

Although the study has achieved its objectives as set out, the following limitations on the scope of the study are highlighted. First, the major limitation of the study is that the field investigation has been carried out in a specific paddy plot for two seasons in a year only. Secondly, the validation was not possible due to the climatic variations between two rice growing seasons. Thirdly, the maintaining the standing water depth was not possible in a single plot. Finally, only one set porous cup with

tube installed at different depths in a paddy field. It could be better if they would be installed at three different locations in the paddy field.

1.6 Organization of the Thesis

Chapter one focuses on the general introduction to the work, gives the statement of the problem, the objectives of the study, scope and limitation of the study and the thesis outlined. Chapter two contains literature review which discusses challenges of rice production in Malaysia, estimation of water balance component analysis from paddy field, solute transport from agricultural fields, methods of reducing fate and solute transport from agricultural fields, in particular, paddy fields. It also contains different surface and subsurface models such as LEACHM, SWAP, PCPF, VS2DT and HYDRUS-1D to simulate water flow and agrochemical (N, P and K). Chapter three discusses the location, soil and climate conditions within the study area. It further describes the description of the experimental setup, sample collection, lab tests, estimation of water balance components, solute transport analysis, development of statistical models, and utilization of Hydrus-1D procedures. Chapter four comprises of the results obtained and their discussions. Chapter five contains the conclusions drawn from the entire work, the major research findings and the recommendations of the further studies. At the end, a short bio-data of the candidate and a list of publications from the study are presented.

REFERENCES

- Abbasi F., Feyen, J., Roth, R. L., Sheedy, M., and van Genuchten, M. T. (2003). Water flow and solute transport in furrow-irrigated fields. *Irrigation Science*, 22(2), 57-65.
- Abbasi, M. R., and Sepaskhah, A. R. (2011). Effects of water-saving irrigations on different rice cultivars (*Oryza sativa* L.) in field conditions. *International Journal of Plant Production*, 5(2), 147-153.
- Abdullahi, A. S., Mohammad Soom, M. A., Ahmad, D., and Mohamed Shariff, A. R. (2013). Characterization of rice ('*Oryza sativa*') evapotranspiration using micro paddy lysimeter and class. *Australian Journal of Crop Science*, 7(5), 650.
- Alaerts, G. J., Mahbubar, R., and Kelderman, P. (1996). Performance analysis of a full-scale duckweed-covered sewage lagoon. *Water Research*, 30(4), 843-852.
- Alam, M. M., Siwar, C., bin Toriman, M. E., Molla, R. I., and Talib, B. (2012). Climate change induced adaptation by paddy farmers in Malaysia. *Mitigation and Adaptation Strategies for Global Change*, 17(2), 173-186.
- Ali, H. (2011). *Practices of irrigation & on-farm water management* (Vol. 2). Springer Science and Business Media.
- Allen, R. G., and Pruitt, W. O. (1991). FAO-24 reference evapotranspiration factors. *Journal of irrigation and drainage engineering*, 117(5), 758-773.
- Allen, R. G., Pereira, L. S., Raes, D., and Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9), D05109.
- Almasri, M. N., and Kaluarachchi, J. J. (2005). Modular neural networks to predict the nitrate distribution in ground water using the on-ground nitrogen loading and recharge data. *Environmental Modelling & Software*, 20(7), 851-871.
- Al-Redhaiman, K. N. (2000). Nitrate accumulation in plants and hazards to man and livestock: A review. *J. King Saud Univ*, 12, 143-156.
- Anbumozhi, V., Yamaji, E., and Tabuchi, T. (1998). Rice crop growth and yield as influenced by changes in ponding water depth, water regime and fertigation level. *Agricultural Water Management*, 37(3), 241-253.
- Anbumozhi, V., Yamaji, E., and Tabuchi, T. (1998). Rice crop growth and yield as influenced by changes in ponding water depth, water regime and fertigation level. *Agricultural Water Management*, 37(3), 241-253.

- Angus, J.F. (1991). The evolution of methods for quantifying risk in water limited environments. In: Muchow RC, Bellamy JA (eds) Climatic risk in crop production: models and management for the semiarid tropics and subtropics. CAB International, Wallingford, pp 39-53.
- Anjana, S. U., and Iqbal, M. (2007). Nitrate accumulation in plants, factors affecting the process, and human health implications. A review. *Agronomy for Sustainable Development*, 27(1), 45-57.
- Antonopoulos, V. Z. (2008). Modeling of water and nitrogen balance in the ponded water of rice fields. *Paddy and Water Environment*, 6(4), 387-395.
- Antonopoulos, V. Z. (2010). Modelling of water and nitrogen balances in the ponded water and soil profile of rice fields in Northern Greece. *Agricultural water management*, 98(2), 321-330.
- Anyusheva, M., Lamers, M., La, N., Nguyen, V. V., and Streck, T. (2012). Fate of pesticides in combined paddy rice–fish pond farming systems in Northern Vietnam. *Journal of environmental quality*, 41(2), 515-525.
- Asada, K., Eguchi, S., Tsunekawa, A., Tsuji, M., Itahashi, S., and Katou, H. (2015). Predicting nitrogen leaching with the modified LEACHM model: validation in soils receiving long-term application of animal manure composts. *Nutrient cycling in agroecosystems*, 102(2), 209-225.
- Avvannavar, S. M., and Shrihari, S. (2007). Evaluation of water quality index for drinking purposes for river Netravathi, Mangalore, South India. *Environmental monitoring and assessment*, 143(1-3), 279-290.
- Ayars, J. E., Christen, E. W., and Hornbuckle, J. W. (2006). Controlled drainage for improved water management in arid regions irrigated agriculture. *agricultural water management*, 86(1-2), 128-139.
- Ayob, A. H., Zulkefli, M., Mohammad, C. H., Sharma, M. L., Mohd Zain, M., Aminuddin, B. Y., and Ahmad, A. R. (1996, December). Water and nutrient balance study in the KADA paddy irrigation scheme. In *Aciar Proceedings* (pp. 33-44). Australian Centre for International Agricultural Research.
- Azevedo, A. S., Singh, P., Kanwar, R. S., and Ahuja, L. R. (1997). Simulating nitrogen management effects of subsurface drainage water quality. *Agricultural Systems*, 55(4), 481.
- Bagheri, A., Shabanali Fami, H., Rezvanfar, A., Asadi, A. and Yazdani, S. (2008). Perception of Paddy Farmers towards Sustainable Agricultural Technologies: Case of Haraz Catchments Area in Mazandaran Province of Iran. *American Journal of Applied Sciences*, 5(10): 1384-1391, 2008.

- Bakar, K. B., and Azman, S. (2012). *Garbage Enzyme As An Alternative Method in Treatment of Sullage: Suitable for household user*. LAP LAMBERT Academic Publishing.
- Bala, B. K., Alias, E. F., Fatimah, M. A., Noh, K. M. and Hadi, A. H. A. (2014). *Modelling of Food Security in Malaysia. Simulation Modelling Practice and Theory*, 47 (2014) 152-164.
- Becker, M. W., and Jiang, Z. (2007). Flux-based contaminant transport in a GIS environment. *Journal of hydrology*, 343(3), 203-210.
- Belder, P., Bouman, B. A. M., Cabangon, R., Guoan, L., Quilang, E. J. P., Yuanhua, L., and Tuong, T. P. (2004). Effect of water-saving irrigation on rice yield and water use in typical lowland conditions in Asia. *Agricultural water management*, 65(3), 193-210.
- Beman, J. M., Arrigo, K. R., and Matson, P. A. (2005). Agricultural runoff fuels large phytoplankton blooms in vulnerable areas of the ocean. *Nature*, 434(7030), 211.
- Bhuiyan, S. I. (1982). *Irrigation system management research and selected methodological issues*. IRRI research paper series-International Rice Research Institute.
- Blackmer, A. M. (1987). Losses and transport of nitrogen from soils.
- Bleken, M. A., and Bakken, L. R. (1997). The nitrogen cost of food production: Norwegian society. *Ambio*, 134-142.
- Boesten, J. J. T. I., and Van der Linden, A. M. A. (1991). Modeling the influence of sorption and transformation on pesticide leaching and persistence. *Journal of Environmental Quality*, 20(2), 425-435.
- Bolton, F. R., and Zandstra, H. G. (1981). A soil moisture-based yield model of wetland rainfed rice.
- Boulangé, J., Watanabe, H., Inao, K., Iwafune, T., Zhang, M., Luo, Y., and Arnold, J. (2014). Development and validation of a basin scale model PCPF-1@SWAT for simulating fate and transport of rice pesticides. *Journal of hydrology*, 517, 146-156.
- Bouman, B. A. M., and Tuong, T. P. (2001). Field water management to save water and increase its productivity in irrigated lowland rice. *Agricultural water management*, 49(1), 11-30.
- Bouman, B. A. M., Humphreys, E., Tuong, T. P., and Barker, R. (2007). Rice and water. *Advances in agronomy*, 92, 187-237.

- Bouman, B. A. M., Peng, S., Castaneda, A. R., and Visperas, R. M. (2005). Yield and water use of irrigated tropical aerobic rice systems. *Agricultural Water Management*, 74(2), 87-105.
- Bradford, S. F., and Katopodes, N. D. (1998). Nonhydrostatic model for surface irrigation. *Journal of Irrigation and Drainage Engineering*, 124(4), 200-212.
- Brouwer, F. M., Godeschalk, F. E., Hellegers, P. J. G. J., and Kelholt, H. J. (1995). Mineral balances at farm level in the European Union (No. 137). LEI-DLO.
- Bünemann, E. K., Steinebrunner, F., Smithson, P. C., Frossard, E., and Oberson, A. (2004). Phosphorus dynamics in a highly weathered soil as revealed by isotopic labeling techniques. *Soil Science Society of America Journal*, 68(5), 1645-1655.
- Buresh, R. J., and Witt, C. (2007). Site-specific nutrient management. Fertilizer best management practices, 47.
- Cai, X., McKinney, D. C., and Rosegrant, M. W. (2003). Sustainability analysis for irrigation water management in the Aral Sea region. *Agricultural Systems*, 76(3), 1043-1066.
- Camargo, J. A., and Alonso, Á. (2006). Ecological and toxicological effects of inorganic nitrogen pollution in aquatic ecosystems: a global assessment. *Environment international*, 32(6), 831-849.
- Cameron, S. G., and Schipper, L. A. (2010). Nitrate removal and hydraulic performance of organic carbon for use in denitrification beds. *Ecological Engineering*, 36(11), 1588-1595.
- Cannavo, P., Recous, S., Parnaudeau, V., and Reau, R. (2008). Modeling N dynamics to assess environmental impacts of cropped soils. *Advances in Agronomy*, 97, 131-174.
- Cantrell, R. P., and Hettel, G. P. (2004, February). New challenges and technological opportunities for rice-based production systems for food security and poverty alleviation in Asia and the Pacific. In presented at the FAO Rice Conference, FAO, Rome, Italy.
- Carsel, R. F., Imhoff, J. C., Hummel, P. R., Cheplick, J. M., and Donigian, A. J. A (1998). Model for predicting pesticide and nitrogen fate in the crop root and unsaturated soil zones. User's manual (US Environmental Protection Agency, Athens, GA, 1999).
- Cerejeira, M. J., Viana, P., Batista, S., Pereira, T., Silva, E., Valério, M. J., and Silva-Fernandes, A. M. (2003). Pesticides in Portuguese surface and ground waters. *Water research*, 37(5), 1055-1063.

- Chan, C. S., and Cheong, A. W. (2001). Seasonal weather effects on crop evapotranspiration and rice yield. *Journal of Tropical Agriculture and Food Science*, 29, 77-92.
- Chan, K. Y., Oates, A., Swan, A. D., Hayes, R. C., Dear, B. S., and Peoples, M. B. (2006). Agronomic consequences of tractor wheel compaction on a clay soil. *Soil and Tillage Research*, 89(1), 13-21.
- Chandna, P., Khurana, M. L., Ladha, J. K., Punia, M., Mehla, R. S., and Gupta, R. (2011). Spatial and seasonal distribution of nitrate-N in groundwater beneath the rice-wheat cropping system of India: a geospatial analysis. *Environmental monitoring and assessment*, 178(1-4), 545-562.
- Chen, M., Tang, H., Ma, H., Holland, T. C., Ng, K. S., and Salley, S. O. (2011). Effect of nutrients on growth and lipid accumulation in the green algae *Dunaliella tertiolecta*. *Bioresource technology*, 102(2), 1649-1655.
- Chen, S. K., and Liu, C. W. (2002). Analysis of water movement in paddy rice fields (I) experimental studies. *Journal of hydrology*, 260(1), 206-215.
- Cho, J. Y., and Han, K. W. (2002). Nutrient losses from a paddy field plot in central Korea. *Water, Air, and Soil Pollution*, 134(1-4), 215-228.
- Cho, J. Y., Han, K. W., Choi, J. K., Kim, Y. J., and Yoon, K. S. (2002) N and P losses from a paddy field plot in central Korea. *Soil Science and Plant Nutrition*, 48 (3):301-306.
- Choi, W. J., Han, G. H., Ro, H. M., Yoo, S. H., and Lee, S. M. (2002). Evaluation of nitrate contamination sources of unconfined groundwater in the North Han River basin of Korea using nitrogen isotope ratios. *Geosciences Journal*, 6(1), 47-55.
- Chowdary, V. M., Rao, N. H., and Sarma, P. B. S. (2004). A coupled soil water and nitrogen balance model for flooded rice fields in India. *Agriculture, Ecosystems & Environment*, 103(3), 425-441.
- Chung, S. O., Kim, H. S., and Kim, J. S. (2003). Model development for nutrient loading from paddy rice fields. *Agricultural Water Management*, 62(1), 1-17.
- Cichota, R., and Snow, V. O. (2009). Estimating nutrient loss to waterways—an overview of models of relevance to New Zealand pastoral farms. *New Zealand Journal of Agricultural Research*, 52(3), 239-260.
- Correll, D. L. (1998). The role of phosphorus in the eutrophication of receiving waters: A review. *Journal of Environmental Quality*, 27(2), 261-266.
- Daniel, T. C., Sharpley, A. N., and Lemunyon, J. L. (1998). Agricultural phosphorus and eutrophication: A symposium overview. *Journal of Environmental Quality*, 27(2), 251-257.

- Dash, C. J., Sarangi, A., Adhikary, P. P., and Singh, D. K. (2015). Simulation of Nitrate Leaching under Maize–Wheat Cropping System in a Semiarid Irrigated Area of the Indo-Gangetic Plain, India. *Journal of Irrigation and Drainage Engineering*, 04015053.
- Davis, A. P., and McCuen, R. H. (2005). Stormwater management for smart growth. Springer Science & Business Media.
- De Data, S. K. (1981). Principles and practices of rice production (No. 633.18 D2622p Ej. 2 019781). John Wiley & Sons,.
- De Datta, S. K. (1986). 10. Improving nitrogen fertilizer efficiency in lowland rice in tropical Asia. *Fertilizer research*, 9(1-2), 171-186.
- De Datta, S. K., Abilay, W. P., and Kalwar, G. N. (1973). Water stress effects in flooded tropical rice. *Water Management in Philippine Irrigation Systems: Research and Operations*, 19-36.
- Decrem, M., Spiess, E., Richner, W., and Herzog, F. (2007). Impact of Swiss agricultural policies on nitrate leaching from arable land. *Agronomy for sustainable development*, 27(3), 243-253.
- Deng, Y. J., Song, G. C., and Wen, Z. S. (2008). Rice controlled irrigation techniques further studies. *Heilongjiang Sci Technol Water Conserv*, 36(05), 10-11.
- Department of Irrigation and Drainage (DID). (2009). Malaysian Environmental Quality Report 2009. http://h2o.water.gov.my/man_water/WRP21.pdf.
- Döll, P. (2002). Impact of climate change and variability on irrigation requirements: a global perspective. *Climatic change*, 54(3), 269-293.
- Doorenbos, J., and Kassam, A. H. (1979). Yield response to water. Irrigation and Drainage Paper 33. FAO, Rome. *Yield response to water. Irrigation and Drainage Paper 33. FAO, Rome*.
- Doorenbos, J., and Pruitt, W. O. (1977). Crop water requirements. Irrigation and drainage paper no. 24. FAO, Rome.
- Ebrahimian, H., Liaghat, A., Parsinejad, M., Abbasi, F., and Navabian, M. (2012). Comparison of one-and two-dimensional models to simulate alternate and conventional furrow fertigation. *Journal of Irrigation and Drainage Engineering*, 138(10), 929-938.
- Enright, P., and Madramootoo, C. A. (2004). Phosphorus losses in surface runoff and subsurface drainage waters on two agricultural fields in Quebec. In *Drainage VIII, 21-24 March 2004* (p. 1). American Society of Agricultural and Biological Engineers.

- Eom, K. C. (2001). Environmentally beneficial function of rice culture and paddy soil. Rice culture in Asia. International Commission on Irrigation and Drainage, and Korean National Committee on Irrigation and Drainage, Korea, 28-35.
- Fang, W. A. N. G., Yong-Rong, B. I. A. N., Jiang, X., Hong-Jian, G. A. O., Gui-Fen, Y. U., and Jian-Cai, D. E. N. G. (2006). Residual Characteristics of Organochlorine Pesticides in Lou Soils with Different Fertilization Modes. *1. Pedosphere*, 16(2), 161-168.
- FAO (2004) FAOSTAT Database Agricultural Production; Food and Agriculture Organization of the United Nations: Rome, Italy. Available online: <http://www.apps.fao.org> (accessed 26 November 2004).
- FAO (2005). AQUASTAT on-line database, Food and Agriculture Organization, Rome, <http://faostat.fao.org> last access: 10 October 2005.
- FAO, U. (2009, December). How to feed the world in 2050. In Rome: High-Level Expert Forum.
- FAO. (2015). World fertilizer trends and outlook to 2018. Food and Agriculture organization of the United Nations (FAO), Rome, ISBN 978-92-5-108692-6, Annual Report 14.
- Feddes, R. A., Kowalik, P. J., and Zaradny, H. (1978). Simulation of field water use and crop yield. John Wiley & Sons, New York. Simulation of field water use and crop yield. John Wiley & Sons, New York.
- Feng, Y. W., Yoshinaga, I., Shiratani, E., Hitomi, T., and Hasebe, H. (2004). Characteristics and behavior of nutrients in a paddy field area equipped with a recycling irrigation system. *Agricultural water management*, 68(1), 47-60.
- Fernald, A. G., and Guldán, S. J. (2006). Surface water-groundwater interactions between irrigation ditches, alluvial aquifers, and streams. *Reviews in Fisheries Science*, 14(1-2), 79-89.
- Fisher, M. J., Fausey, N. R., Subler, S. E., Brown, L. C., and Bierman, P. M. (1999). Water table management, nitrogen dynamics, and yields of corn and soybean. *Soil Science Society of America Journal*, 63(6), 1786-1795.
- Fukuda H, and Tsutsui H., (1973). Rice Irrigation in Japan. Overseas Technical Cooperation Agency, Tokyo, Japan.
- Füleký, G. (2014). Accumulation and depletion of fertilizer originated Nitrate-N and Ammonium-N in deeper soil layers. *COLUMELLA: JOURNAL OF AGRICULTURAL AND ENVIRONMENTAL SCIENCES*, 1(2), 7-16.

- Gao, C., Zhu, J. G., Zhu, J. Y., Gao, X., Dou, Y. J., and Hosen, Y. (2004). Nitrogen export from an agriculture watershed in the Taihu Lake area, China. *Environmental Geochemistry and Health*, 26(2), 199-207.
- Gaydon, D. S., Probert, M. E., Buresh, R. J., Meinke, H., Suriadi, A., Dobermann, A., and Timsina, J. (2012). Rice in cropping systems—Modelling transitions between flooded and non-flooded soil environments. *European journal of agronomy*, 39, 9-24.
- Gentry, L. E., David, M. B., Below, F. E., Royer, T. V., and McIsaac, G. F. (2009). Nitrogen mass balance of a tile-drained agricultural watershed in east-central Illinois. *Journal of environmental quality*, 38(5), 1841-1847.
- Gervais, J. A., Luukinen, B., Buhl, K., and Stone, D. (2010). Imidacloprid technical fact sheet. National Pesticide Information Center, Oregon State University Extension Services.
- Ghani, M. A. (1978). Studies on seepage and percolation in selected field conditions in Bangladesh. Unpublished Master of Engineering Thesis. Asian Institute of Technology, Bangkok, Thailand.
- Gheysari, M., Mirlatifi, S. M., Homae, M., Asadi, M. E., and Hoogenboom, G. (2009). Nitrate leaching in a silage maize field under different irrigation and nitrogen fertilizer rates. *Agricultural water management*, 96(6), 946-954.
- Ghosh, B. C., and Bhat, R. (1998). Environmental hazards of nitrogen loading in wetland rice fields. In *Nitrogen, the Confer-Ns* (pp. 123-126).
- Guo, D. L., Mitchell, R. J., and Hendricks, J. J. (2004). Fine root branch orders respond differentially to carbon source-sink manipulations in a longleaf pine forest. *Oecologia*, 140(3), 450-457.
- Hafeez, M. M., Bouman, B. A. M., Van de Giesen, N., and Vlek, P. (2007). Scale effects on water use and water productivity in a rice-based irrigation system (UPRIIS) in the Philippines. *Agricultural water management*, 92(1-2), 81-89.
- Hama, T., Aoki, T., Osuga, K., Sugiyama, S., and Iwasaki, D. (2012). Nitrogen and phosphorus effluent loads from a paddy-field district adopting collective crop rotation. *Water Science and Technology*, 66(5), 1074-1080.
- Hama, T., Nakamura, K., Kawashima, S., Kaneki, R., and Mitsuno, T. (2011). Effects of cyclic irrigation on water and nitrogen mass balances in a paddy field. *Ecological Engineering*, 37(10), 1563-1566.
- Hanson, B. R., Šimůnek, J., and Hopmans, J. W. (2006). Evaluation of urea–ammonium–nitrate fertigation with drip irrigation using numerical modeling. *Agricultural water management*, 86(1-2), 102-113.

- Harris, R. C., and Skinner, A. C. (1992). Controlling diffuse pollution of groundwater from agriculture and industry. *Water and Environment Journal*, 6(6), 569-574.
- Haygarth, P. M., Hepworth, L., and Jarvis, S. C. (1998). Forms of phosphorus transfer in hydrological pathways from soil under grazed grassland. *European Journal of Soil Science*, 49(1), 65-72.
- Healy, R. W. (1990). Simulation of solute transport in variably saturated porous media with supplemental information on modifications to the US Geological Survey's computer program VS2D. Department of the Interior, US Geological Survey.
- Hillel, D. (1998). *Environmental soil physics: Fundamentals, applications, and environmental considerations*. Elsevier, New York, USA.
- Hou, H., Zhou, S., Hosomi, M., Toyota, K., Yosimura, K., Mutou, Y., and Motobayashi, T. (2007). Ammonia emissions from anaerobically-digested slurry and chemical fertilizer applied to flooded forage rice. *Water, air, and soil pollution*, 183(1-4), 37-48.
- Huang, J., Reneau, R. B., and Hagedorn, C. (2000). Nitrogen removal in constructed wetlands employed to treat domestic wastewater. *Water Research*, 34(9), 2582-2588.
- Hukkeri, S. B., and Sharma, A. K. (1980). Water use efficiency of transplanted and direct-sown rice under different water-management practices [India]. *Indian Journal of Agricultural Sciences (India)*.
- Hutson, J. L., and Wagenet, R. J. (1992). LEACHM, Leaching Estimation and CHemistry Model, Version 3.0, Dept (No. 92-3). SCAS Research Report.
- ICID. 2013. Sprinkler and micro irrigated area. Data provided by National Committees, International Commission on Irrigation and Drainage. Available at http://www.icid.org/sprin_micro_11.pdf, accessed in May, 2013.
- IFA (2006) International fertilizer industry association. Statistics nitrogen fertilizer nutrient consumption. Updated June 2006 <http://www.fertilizer.org/ifa/statistics/indicators/tablen.asp>.
- Inao, K., and Kitamura, Y. (1999). Pesticide paddy field model (PADDY) for predicting pesticide concentrations in water and soil in paddy fields. *Pesticide science*, 55(1), 38-46.
- Inao, K., and Kitamura, Y. (1999). Pesticide paddy field model (PADDY) for predicting pesticide concentrations in water and soil in paddy fields. *Pesticide Science*, 55(1), 38-46.

- Inao, K., Watanabe, H., Karpouzas, D. G., and Capri, E. (2008). Simulation models of pesticide fate and transport in paddy environment for ecological risk assessment and management. *Japan Agricultural Research Quarterly: JARQ*, 42(1), 13-21.
- IPCC. 2007. *Climate change 2007: The physical science basis. Contribution of Working Group I to the fourth assessment report of the Intergovernmental Panel on Climate Change*. SD Solomon (Ed.).
- Iqbal, M. T. (2011). Nitrogen leaching from paddy field under different fertilization rates. *Malaysian Journal of Soil Science*, 15, 101-114.
- Ishii, S., Ikeda, S., Minamisawa, K., and Senoo, K. (2011). Nitrogen cycling in rice paddy environments: past achievements and future challenges. *Microbes and Environments*, 26(4), 282-292.
- Iwasaki, N., Inao, K., Iwafune, T., Horio, T., and Obara, H. (2012). Coupling of the PADDY-Large model with geospatial information for predicting paddy pesticide behavior in river basins. *Limnology*, 13(2), 221-235.
- Iwata, S., Tabuchi, T., and Warkentin, B. P. (1995). *Soil-water interactions: mechanisms and applications* (No. Ed. 2). Marcel Dekker, Inc.
- Jamieson, A., Madramootoo, C. A., and Enright, P. (2003). Phosphorus losses in surface and subsurface runoff from a snowmelt event on an agricultural field in Quebec. *Canadian Biosystems Engineering*, 45, 1-1.
- Jarvis, S., Hutchings, N. J., Brentrup, F., Olesen, J. E., and van der Hoek, K. W. (2011). Nitrogen flows in farming systems across Europe. In *European Nitrogen Assessment* (pp. 211-228). Cambridge University Press.
- Jean, C. B., Paul, B. M., Jan, D. G and Alfons, O. L. (2015). Potential Impact of Alternative Agricultural Technologies to Ensure Food Security and Raise Income of Farm Households in Rwanda. *Forum for Development Studies*. Vol. 42, No. 1, 133- 157.
- Jensen, M. E., and Allen, R. G. (Eds.). (2016, April). *Evaporation, evapotranspiration, and irrigation water requirements*. American Society of Civil Engineers.
- Jeon, J. H., Yoon, C. G., Ham, J. H., and Jung, K. W. (2005). Model development for surface drainage loading estimates from paddy rice fields. *Paddy and Water Environment*, 3(2), 93-101.
- Jeon, J. H., Yoon, C. G., Ham, J. H., and Jung, K. W. (2005). Model development for surface drainage loading estimates from paddy rice fields. *Paddy and Water Environment*, 3(2), 93-101.

- Jeong, C.Y., Weindorf, D. C., DeRamus, A., and Goodeaux, L.L. (2011). Surface and subsurface phosphorus losses from sugarcane fields with different management practices. *Water, Air, & Soil Pollution*, 217(1-4), 649-661.
- Ji, X. H., Zheng, S. X., Shi, L. H., and Liao, Y. L. (2008). Effect of fertilization on nutrient leaching loss from different paddy soils in Dongting Lake Area. *Acta Pedofil Sin*, 45(4), 663-671 (English Abstract).
- Jia, Z., Luo, W., Fang, S., Wang, N., and Wang, L. (2006). Evaluating current drainage practices and feasibility of controlled drainage in the YinNan Irrigation District, China. *Agricultural water management*, 84(1-2), 20-26.
- JICA and DID (1998). The study on modernization of irrigation water management system in the Granary area of Peninsular Malaysia. Final Report, March 1998, Japan International Cooperation Agency (JICA).
- Juahir, H., Zain, S. M., Yusoff, M. K., Hanidza, T. T., Armi, A. M., Toriman, M. E., and Mokhtar, M. (2011). Spatial water quality assessment of Langat River Basin (Malaysia) using environmetric techniques. *Environmental monitoring and assessment*, 173(1-4), 625-641.
- Kampen, J., (1970). Water losses and water balance studies in lowland rice irrigation. Ph.D. thesis, Cornell University, USA, 416 pp.
- Kaneki, R. (1991). Water quality of return flow and the purification by using paddy fields. *Journal of the Japanese Society of Irrigation, Drainage and Reclamation Engineering*.
- Karpouzias, D.G., and Capri, E. (2006). Risk analysis of pesticides applied to rice paddies using RICEWQ 1.6. 2v and RIVWQ 2.02. *Paddy and Water Environment*, 4(1), 29-38.
- Karul, C., Soyupak, S., Çilesiz, A.F., Akbay, N., and Germen, E. (2000). Case studies on the use of neural networks in eutrophication modeling. *Ecological modelling*, 134(2), 145-152.
- Khakural, B.R., and Robert, P. C. (1993). Soil nitrate leaching potential indices: using a simulation model as a screening system. *Journal of environmental quality*, 22(4), 839-845.
- Khan, S., Xevi, E., and Meyer, W.S. (2003). Salt, water, and groundwater management models to determine sustainable cropping patterns in shallow saline groundwater regions of Australia. *Journal of crop production*, 7(1-2), 325-340.
- Kibe, K., Takahashi, M., Kameya, T., and Urano, K. (2000). Adsorption equilibriums of principal herbicides on paddy soils in Japan. *Science of the total environment*, 263(1-3), 115-125.

- Kidmose, J., Engesgaard, P., Nilsson, B., Laier, T., and Looms, M. C. (2011). Spatial Distribution of Seepage at a Flow-Through Lake: Lake Hampen, Western Denmark. *Vadose Zone Journal*, 10(1), 110-124.
- Kijne, J., Barron, J., Hoff, H., Rockström, J., Karlberg, L., Gowing, J., and Wichelns, D. (2009). Opportunities to increase water productivity in agriculture with special reference to Africa and South Asia. Stockholm Environment Institute, Project Report.
- Kim, M. Y., Jee, H. K., Lee, S. T., and Kim, M. K. (2006). Prediction of nitrogen and phosphorus transport in surface runoff from agricultural watersheds. *KSCE Journal of Civil Engineering*, 10(1), 53-58.
- Kim, M. Y., Jee, H. K., Lee, S. T., and Kim, M. K. (2006). Prediction of nitrogen and phosphorus transport in surface runoff from agricultural watersheds. *KSCE Journal of Civil Engineering*, 10(1), 53-58.
- Kiran, J. K., Khanif, Y. M., Amminuddin, H., and Anuar, A. R. (2010). Effects of controlled release urea on the yield and nitrogen nutrition of flooded rice. *Communications in soil science and plant analysis*, 41(7), 811-819.
- Koshino, M. (1976). Incoming and outgoing of fertilizer nutrients in cropped lands. In *Science for Better Environment; Proceedings of the International Congress on the Human Environment*.
- Kroes, J. G., Wesseling, J. G., and Van Dam, J. C. (2000). Integrated modelling of the soil–water–atmosphere–plant system using the model SWAP 2.0: an overview of theory and an application. *Hydrological processes*, 14(11-12), 1993-2002.
- Kroeze, C., Aerts, R., van Breemen, N., van Dam, D., Hofschreuder, P., Hoosbeek, M., & Oenema, O. (2003). Uncertainties in the fate of nitrogen I: An overview of sources of uncertainty illustrated with a Dutch case study. *Nutrient Cycling in Agroecosystems*, 66(1), 43-69.
- Kubota, H. (1979). Water and material (N, P) balance in the paddy fields along Lake Kasumigaura. *Trans Jpn Soc Irrig Drain Reclam Eng*, 84, 22-28.
- Kudo, A. (1995). Characteristics of water management and outflow load from a paddy field in a return flow irrigation area. *J. Japanese Society of Irrigation, Drainage and Reclamation Engineering*, 63, 179-184.
- Kumar Reddy, D. H., and Lee, S. M. (2012). Water pollution and treatment technologies. *J Environ Anal Toxicol*, 2, e103.
- Lage, M., Bamouh, A., Karrou, M., and El Mourid, M. (2003). Estimation of rice evapotranspiration using a microlysimeter technique and comparison with FAO Penman-Monteith and Pan evaporation methods under Moroccan conditions. *Agronomie*, 23(7), 625-631.

- Lalonde, V., Madramootoo, C. A., Trenholm, L., and Broughton, R. S. (1996). Effects of controlled drainage on nitrate concentrations in subsurface drain discharge. *Agricultural Water Management*, 29(2), 187-199.
- Lamers, M., Anyusheva, M., La, N., Nguyen, V. V., and Streck, T. (2011). Pesticide pollution in surface-and groundwater by paddy rice cultivation: a case study from Northern Vietnam. *Clean–Soil, Air, Water*, 39(4), 356-361.
- Lappala, E. G., Healy, R. W., and Weeks, E. P. (1987). Documentation of computer program VS2D to solve the equations of fluid flow in variably saturated porous media (pp. 83-4099). Department of the Interior, US Geological Survey.
- Lee, H., Masuda, T., Yasuda, H., and Hosoi, Y. (2014). The pollutant loads from a paddy field watershed due to agricultural activity. *Paddy and Water Environment*, 12(4), 439-448.
- Lee, K. H., and Jose, S. (2005). Nitrate leaching in cottonwood and loblolly pine biomass plantations along a nitrogen fertilization gradient. *Agriculture, ecosystems & environment*, 105(4), 615-623.
- Lee, K. H., and Jose, S. (2005). Nitrate leaching in cottonwood and loblolly pine biomass plantations along a nitrogen fertilization gradient. *Agriculture, ecosystems & environment*, 105(4), 615-623.
- Lee, S. T., Haque, M. A., and Feng, H. Y. (2006). Modeling Water Balance Components in Rice Field Irrigation. *The Institution of Engineers: Kuala Lumpur, Malaysia*, 67, 22-25.
- Lee, T. S., Haque, M. A., and Najim, M. M. M. (2005). Scheduling the cropping calendar in wet-seeded rice schemes in Malaysia. *Agricultural water management*, 71(1), 71-84.
- Leon, A., and Kohyama, K. (2017). Estimating nitrogen and phosphorus losses from lowland paddy rice fields during cropping seasons and its application for life cycle assessment. *Journal of Cleaner Production*, 164, 963-979.
- Li, H., Liang, X., Chen, Y., Tian, G., and Zhang, Z. (2008). Ammonia volatilization from urea in rice fields with zero-drainage water management. *Agricultural Water Management*, 95(8), 887-894.
- Li, W., Xia, Y., Ti, C., and Yan, X. (2011). Evaluation of biological and chemical nitrogen indices for predicting nitrogen-supplying capacity of paddy soils in the Taihu Lake region, China. *Biology and Fertility of Soils*, 47(6), 669-678.
- Li, X., Hu, C., Delgado, J. A., Zhang, Y., and Ouyang, Z. (2007). Increased nitrogen use efficiencies as a key mitigation alternative to reduce nitrate leaching in north china plain. *Agricultural Water Management*, 89(1), 137-147.

- Li, Y. H. (2001). Research and practice of water-saving irrigation for rice in China. In *Water saving irrigation for rice: proceedings of an international workshop, Wuhan, China*. International Water Management Institute (pp. 1-9).
- Li, Y., Šimůnek, J., Wang, S., Yuan, J., and Zhang, W. (2017). Modeling of soil water regime and water balance in a transplanted rice field experiment with reduced irrigation. *Water*, 9(4), 248.
- Li, Y., Šimůnek, J., Zhang, Z., Huang, M., Ni, L., Zhu, L., and Chen, Y. (2015). Water flow and nitrate transport through a lakeshore with different revetment materials. *Journal of Hydrology*, 520, 123-133.
- Li, Y., Šimůnek, J., Zhang, Z., Huang, M., Ni, L., Zhu, L., and Chen, Y. (2015a). Water flow and nitrate transport through a lakeshore with different revetment materials. *Journal of Hydrology*, 520, 123-133.
- Li, Y., Šimůnek, J., Zhang, Z., Huang, M., Ni, L., Zhu, L., and Chen, Y. (2015). Water flow and nitrate transport through a lakeshore with different revetment materials. *Journal of Hydrology*, 520, 123-133.
- Li, Y., Šimůnek, J., Zhang, Z., Jing, L., and Ni, L. (2015b). Evaluation of nitrogen balance in a direct-seeded-rice field experiment using Hydrus-1D. *Agricultural Water Management*, 148, 213-222.
- Liang, H., Hu, K., Batchelor, W. D., Qin, W., and Li, B. (2018). Developing a water and nitrogen management model for greenhouse vegetable production in China: Sensitivity analysis and evaluation. *Ecological Modelling*, 367, 24-33.
- Liang, X. Q., Chen, Y. X., Li, H., Tian, G. M., Ni, W. Z., He, M. M., and Zhang, Z. J. (2007). Modeling transport and fate of nitrogen from urea applied to a near-trench paddy field. *Environmental pollution*, 150(3), 313-320.
- Liu, F. L., Peng, S. Z., Xu, J. Z., and Ding, J. L. (2004). Control irrigation schedules on rice population structure test [J]. *J Shenyang Agric Univ*, 35(5), 417-419.
- London, J.G. Nitrogen study fertilizes fears of pollution. *Nature* 2005, 433, 791.
- Lotse, E. G., Jabro, J. D., Simmons, K. E., and Baker, D. E. (1992). Simulation of nitrogen dynamics and leaching from arable soils. *Journal of contaminant hydrology*, 10(3), 183-196.
- Ma, Y., Feng, S., Huo, Z., and Song, X. (2011). Application of the SWAP model to simulate the field water cycle under deficit irrigation in Beijing, China. *Mathematical and Computer Modelling*, 54(3-4), 1044-1052.
- Madramootoo, C. A., Dodds, G. T., and Papadopoulos, A. (1993). Agronomic and environmental benefits of water-table management. *Journal of irrigation and drainage engineering*, 119(6), 1052-1065.

- Madramootoo, C. A., Wiyo, K. A., and Enright, P. (1992). Nutrient losses through tile drains from two potato fields. *Applied engineering in agriculture*, 8(5), 639-646.
- Maina, M. M., Amin, M. S. M., Rowshon, M. K., Aimrun, W., Samsuzana, A. A., and Yazid, M. A. (2014). Effects of crop evapotranspiration estimation techniques and weather parameters on rice crop water requirement. *Australian Journal of Crop Science*, 8(4), 495.
- Maru, S. (1991). Study on the behavior and fate of pesticides in aquatic environment. *Spec Bull Chiba Agric Exp Stn*, 18, 1-62.
- Miao, Z., Padovani, L., Riparbelli, C., Ritter, A. M., Trevisan, M., and Capri, E. (2003). Prediction of the environmental concentration of pesticide in paddy field and surrounding surface water bodies. *Paddy and Water Environment*, 1(3), 121-132.
- Minh, N. H., Minh, T. B., Kajiwara, N., Kunisue, T., Iwata, H., Viet, P. H., and Tanabe, S. (2007). Pollution sources and occurrences of selected persistent organic pollutants (POPs) in sediments of the Mekong River delta, South Vietnam. *Chemosphere*, 67(9), 1794-1801.
- Ministry of Agriculture and Agro-Based Industry Malaysia (MOA) (2017). *Perangkaan Agro Makanan*; Ministry of Agriculture and Agro-Based Industry Malaysia (MOA): Putrajaya, Malaysia, 2017.
- Molden, D., Oweis, T. Y., Pasquale, S., Kijne, J. W., Hanjra, M. A., Bindraban, P. S., and Hachum, A. (2007). Pathways for increasing agricultural water productivity.
- Mood, N. S., Parsinejad, M., and Mirzaei, F. (2009). Controlled drainage effects on crop yield and water use efficiency under semi-arid condition of Iran. In *World Environmental and Water Resources Congress 2009: Great Rivers* (pp. 1-8).
- Munster, C. L., Skaggs, R. W., Parsons, J. E., Evans, R. O., Gilliam, J. W., and Breve, M. A. (1994). Simulating aldicarb transport in a drained field. *Transactions of the ASAE (USA)*.
- Naito, Y. (1969). Studies on the transpiration and evapotranspiration amount of various crops. *Bull. Tokai-Kinki Natl. Agric. Exp. Stn.*, 18, 49-151.
- Nakano, H., Nakajima, E., Fujii, Y., Shigemori, H., and Hasegawa, K. (2004). Structure-activity relationships of alkaloids from mesquite (*Prosopis juliflora* (Sw.) DC.). *Plant growth regulation*, 44(3), 207-210.
- Nakasone, H., Abbas, M. A., and Kuroda, H. (2004). Nitrogen transport and transformation in packed soil columns from paddy fields. *Paddy and Water Environment*, 2(3), 115-124.

- Naseri, A. A., Hoseini, Y., Moazed, H., Abbasi, F., Samani, H. M. V., and Sakebi, S. A. (2011). Phosphorus transport through a saturated soil column: Comparison between physical modeling and HYDRUS-3D outputs. *Journal of Applied Sciences*, 11(5), 815-823.
- Neeteson, J. J., and Wadman, W. P. (1987). Assessment of economically optimum application rates of fertilizer N on the basis of response curves. *Fertilizer research*, 12(1), 37-52.
- Ng, H. Y. F., Tan, C. S., Drury, C. F., and Gaynor, J. D. (2002). Controlled drainage and subirrigation influences tile nitrate loss and corn yields in a sandy loam soil in Southwestern Ontario. *Agriculture, ecosystems & environment*, 90(1), 81-88.
- NI, Wuzhong., Jianping, L., and Zhaoliang, Z. (2007). Occurrence of nitrification-denitrification and gaseous nitrogen loss process in flooded rice soil. *Progress in Natural Science*, 17(1), 6-10.
- Nolan, B. T., Bayless, E. R., Green, C. T., Garg, S., Voss, F. D., Lampe, D. C., and Bekins, B. A. (2005). Evaluation of unsaturated-zone solute-transport models for studies of agricultural chemicals (No. USGS-OFR-2005-1197). GEOLOGICAL SURVEY RESTON VA.
- Ochoa, C. G., Fernald, A. G., Guldan, S. J., and Shukla, M. K. (2007). Deep percolation and its effects on shallow groundwater level rise following flood irrigation. *Transactions of the ASABE*, 50(1), 73-81.
- Odhiambo, L. O., and Murty, V. V. N. (1996). Modeling water balance components in relation to field layout in lowland paddy fields. I. Model development. *Agricultural Water Management*, 30(2), 185-199.
- Onweremadu, E. U. (2007). Predicting phosphorus sorption characteristics in highly weathered soils of South-Eastern Nigeria. *Res. J. Environ. Sci*, 1(2), 47-55.
- Padovani, L., Capri, E., Padovani, C., Puglisi, E., and Trevisan, M. (2006). Monitoring tricyclazole residues in rice paddy watersheds. *Chemosphere*, 62(2), 303-314.
- Pande, H. K., and Adak, N. K. (1971). Leaching loss of nitrogen in submerged rice cultivation. *Experimental Agriculture*, 7(04), 329-336.
- Papadopoulou-Mourkidou, E., Karpouzas, D. G., Patsias, J., Kotopoulou, A., Milothridou, A., Kintzikoglou, K., and Vlachou, P. (2004). The potential of pesticides to contaminate the groundwater resources of the Axios river basin in Macedonia, Northern Greece. Part I. Monitoring study in the north part of the basin. *Science of the Total Environment*, 321(1-3), 127-146.

- Pathak, B. K., Kazama, F., and Toshiaki, I. (2004). Monitoring of nitrogen leaching from a tropical paddy in Thailand. *Agricultural Engineering International: CIGR Journal*.
- Peng, S. Z., Yang, S. H., Xu, J. Z., Luo, Y. F., and Hou, H. J. (2011). Nitrogen and phosphorus leaching losses from paddy fields with different water and nitrogen managements. *Paddy and Water Environment*, 9(3), 333-342.
- Peng, S. Z., Yang, S. H., Xu, J. Z., Luo, Y. F., and Hou, H. J. (2011). Nitrogen and phosphorus leaching losses from paddy fields with different water and nitrogen managements. *Paddy and Water Environment*, 9(3), 333-342.
- Peng, S., Xu, J., Huang, Q., and Liu, F. (2004). Controlled irrigation of paddy rice and environmental multifunctionality. *Journal of Shenyang Agricultural University*, 35(5-6), 443-445.
- Penman, H. L. (1956). Estimating evaporation. *Eos, Transactions American Geophysical Union*, 37(1), 43-50.
- Phillips, I. R. (2006). Modelling water and chemical transport in large undisturbed soil cores using HYDRUS-2D. *Soil Research*, 44(1), 27-34.
- Phogat, V., Yadav, A. K., Malik, R. S., Kumar, S., and Cox, J. (2010). Simulation of salt and water movement and estimation of water productivity of rice crop irrigated with saline water. *Paddy and Water Environment*, 8(4), 333-346.
- Phong, T. K., Hiramatsu, K., Vu, S. H., Ishihara, S., and Watanabe, H. (2009). Simulation of pesticide behavior in a paddy block by a pesticide fate and transport model. *J. Fac. Agric. Kyushu Univ*, 54, 505-512.
- Pimentel, D., and Levitan, L. (1986). Pesticides: amounts applied and amounts reaching pests. *Bioscience*, 36(2), 86-91.
- Pirmoradian, N., Sepaskhah, A. R., and Maftoun, M. (2004). Effects of water-saving irrigation and nitrogen fertilization on yield and yield components of rice (*Oryza sativa* L.). *Plant production science*, 7(3), 337-346.
- Qiu, W. G., Tang, H., and Wang, C. (2004). Rule of loss of nitrogen in the surface water of rice fields and the control technology. *Journal of Agro-Environment Science*, 23(4), 740-744.
- Ramli, N. N., Shamsudin, M. N., Mohamed, Z., and Radam, A. (2012). The impact of fertilizer subsidy on Malaysia paddy/rice industry using a system dynamics approach. *International Journal of Social Science and Humanity*, 2(3), 213.
- Rashid, M. A., Kabir, W., Khan, L. R., Saleh, A. F. M., and Khair, M. A. (2009). Estimation of water Loss from irrigated rice fields. *SAARC Agri*, 7(1), 29-42.

- Rath, A. K., Swain, B., Ramakrishnan, B., Panda, D., Adhya, T. K., Rao, V. R., and Sethunathan, N. (1999). Influence of fertilizer management and water regime on methane emission from rice fields. *Agriculture, ecosystems & environment*, 76(2), 99-107.
- Rijtema P E (1965) An analysis of actual evapotranspiration. Agric Res Rep 659. Pudoc, Wageningen, 107 p.
- Rosenzweig, C., and Parry, M. L. (1994). Potential impact of climate change on world food supply. *Nature*, 367(6459), 133-138.
- Rowshon, M. K., Amin, M. S. M., Lee, T. S., and Shariff, A. R. M. (2009). GIS-integrated rice irrigation management information system for a river-fed scheme. *Water resources management*, 23(14), 2841-2866.
- Rowshon, M. K., Amin, M. S. M., Mojid, M. A., and Yaji, M. (2014). Estimated evapotranspiration of rice based on pan evaporation as a surrogate to lysimeter measurement. *Paddy and water environment*, 12(1), 35-41.
- Salehi-Lisar, S. Y., and Bakhshayeshan-Agdam, H. (2016). Drought stress in plants: causes, consequences, and tolerance. In *Drought Stress Tolerance in Plants, Vol 1* (pp. 1-16). Springer, Cham.
- Sasakawa, H., and Yamamoto, Y. (1978). Comparison of the uptake of nitrate and ammonium by rice seedlings: influences of light, temperature, oxygen concentration, exogenous sucrose, and metabolic inhibitors. *Plant Physiology*, 62(4), 665-669.
- Saunders, D. L., and Kalff, J. (2001). Denitrification rates in the sediments of Lake Memphremagog, Canada–USA. *Water Research*, 35(8), 1897-1904.
- Sawyer, J., Nafziger, E., Randall, G., Bundy, L., Rehm, G., and Joern, B. (2006). Concepts and rationale for regional nitrogen rate guidelines for corn.
- Scarff, M., Arnold, S. A., Harvey, L. M., and McNeil, B. (2006). Near infrared spectroscopy for bioprocess monitoring and control: current status and future trends. *Critical Reviews in Biotechnology*, 26(1), 17-39.
- Schaap, M. G., Leij, F. J., and Van Genuchten, M. T. (2001). Rosetta: A computer program for estimating soil hydraulic parameters with hierarchical pedotransfer functions. *Journal of hydrology*, 251(3-4), 163-176.
- Schmidt, K. D., and Sherman, I. (1987). Effect of irrigation on groundwater quality in California. *Journal of irrigation and drainage engineering*, 113(1), 16-29.
- Schwarzenbach, R.P., Escher, B.I., Fenner, K., Hofstetter, T.B., Johnson, C.A., von Gunten, U., and Wehrli, B. (2006). The challenge of micropollutants in aquatic systems. *Science* 313, 1072-1077.

- Seol, S.I., Lee, S.M., Han, G.H., Choi, W.J., and Yoo, S.H. (2000) Urea transformation and nitrogen loss in waterlogged soil column. *Agricultural Chemistry and Biotechnology*, 43 (2): 86–93.
- Sharpley, A. N., Kleinman, P. J., Heathwaite, A. L., Gburek, W. J., Folmar, G. J., and Schmidt, J. P. (2008). Phosphorus loss from an agricultural watershed as a function of storm size. *Journal of Environmental Quality*, 37(2), 362-368.
- Shaw, J. N., Bosch, D. D., West, L. T., Truman, C. C., and Radcliffe, D. E. (2001). Lateral flow in loamy to sandy Kandicudults of the Upper Coastal Plain of Georgia (USA). *Geoderma*, 99(1-2), 1-25.
- Shekofteh, H., Afyuni, M., Hajabbasi, M. A., Iversen, B. V., Nezamabadi-Pour, H., Abassi, F., and Shirani, H. (2013). Modeling of Nitrate Leaching from a Potato Field using HYDRUS-2D. *Communications in soil science and plant analysis*, 44(20), 2917-2931.
- Shi, Y., Gao, X., Zhang, D., and Giorgi, F. (2011). Climate change over the Yarlung Zangbo–Brahmaputra River Basin in the 21st century as simulated by a high resolution regional climate model. *Quaternary international*, 244(2), 159-168.
- Shiratani, E., Yoshinaga, I., Feng, Y., and Hasebe, H. (2004). Scenario analysis for reduction of effluent load from an agricultural area by recycling the run-off water. *Water Science and Technology*, 49(3), 55-62.
- Sik Yoon, K., Kyu Choi, J., Gwon Son, J., and Young Cho, J. (2006). Concentration profile of nitrogen and phosphorus in leachate of a paddy plot during the rice cultivation period in southern Korea. *Communications in soil science and plant analysis*, 37(13-14), 1957-1972.
- Simard, R. R., Cluis, D., Gangbazo, G., and Beauchemin, S. (1995). Phosphorus status of forest and agricultural soils from a watershed of high animal density. *Journal of Environmental Quality*, 24(5), 1010-1017.
- Šimůnek, J., van Genuchten, M. T., and Šejna, M. (2008). Development and applications of the HYDRUS and STANMOD software packages and related codes. *Vadose Zone Journal*, 7(2), 587-600.
- Singh, R., Van Dam, J. C., and Feddes, R. A. (2006). Water productivity analysis of irrigated crops in Sirsa district, India. *Agricultural Water Management*, 82(3), 253-278.
- Skaggs, T. H., Trout, T. J., Šimůnek, J., and Shouse, P. J. (2004). Comparison of HYDRUS-2D simulations of drip irrigation with experimental observations. *Journal of irrigation and drainage engineering*, 130(4), 304-310.

- Sogbedji, J. M., Van Es, H. M., Melkonian, J. J., and Schindelbeck, R. R. (2006). Evaluation of the PNM model for simulating drain flow nitrate-N concentration under manure-fertilized maize. *Plant and soil*, 282(1-2), 343-360.
- Soil Survey Staff (1975). *Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys*; USDA: Washington, DC, USA.
- Strebel, O. W. H. M., Duynisveld, W. H. M., and Böttcher, J. (1989). Nitrate pollution of groundwater in western Europe. *Agriculture, ecosystems & environment*, 26(3-4), 189-214.
- Strelkoff, T. S., Tamimi, A. H., and Clemmens, A. J. (2003). Two-dimensional basin flow with irregular bottom configuration. *Journal of irrigation and drainage engineering*, 129(6), 391-401.
- Sudo, M., Kunimatsu, T., and Okubo, T. (2002). Concentration and loading of pesticide residues in Lake Biwa basin (Japan). *Water Research*, 36(1), 315-329.
- Sudo, M., Okubo, T., and Kaneki, R. (2005). Paddy herbicide inputs in the entire river inflow reaching Lake Biwa, Japan. *Limnology*, 6(2), 91-99.
- Sugimoto, K. (1976). Relationship between evapotranspiration and dry matter production of Indica rice. *Tropical agriculture research series*.
- Sutanto, S. J., Wenninger, J., Coenders-Gerrits, A. M. J., and Uhlenbrook, S. (2012). Partitioning of evaporation into transpiration, soil evaporation and interception: a comparison between isotope measurements and a HYDRUS-1D model. *Hydrology and Earth System Sciences*, 16(8), 2605-2616.
- Tabbal, D. F., Bouman, B. A. M., Bhuiyan, S. I., Sibayan, E. B., and Sattar, M. A. (2002). On-farm strategies for reducing water input in irrigated rice; case studies in the Philippines. *Agricultural Water Management*, 56(2), 93-112.
- Tabuchi, T., and Takamura, Y. (1985). Nitrogen and phosphorus outflow from catchment area.
- Tabuchi, T., Takamura, Y., Kubota, H., and Suzuki, S. (1975). On the water pollution of rivers into Kasumigaura Lake basin [Japan], 3: The water quality and load of rivers during manuring period. *Transactions of the Japanese Society of Irrigation, Drainage and Reclamation Engineering (English Abstract)*.
- Tafteh, A., and Sepaskhah, A. R. (2012). Application of HYDRUS-1D model for simulating water and nitrate leaching from continuous and alternate furrow irrigated rapeseed and maize fields. *Agricultural water management*, 113, 19-29.

- Takagi, K., Fajardo, F. F., Inao, K., and Kitamura, Y. (1998). Predicting pesticide behavior in a lowland environment using computer simulation. *Reviews in Toxicology*, 2(1), 269-286.
- Takeda, I., and Fukushima, A. (2004). Phosphorus purification in a paddy field watershed using a circular irrigation system and the role of iron compounds. *Water research*, 38(19), 4065-4074.
- Takeda, I., Fukushima, A., and Tanaka, R. (1997). Non-point pollutant reduction in a paddy-field watershed using a circular irrigation system. *Water Research*, 31(11), 2685-2692.
- Tan X., Shao, D., Gu, W., and Liu, H. (2015). Field analysis of water and nitrogen fate in lowland paddy fields under different water managements using HYDRUS-1D. *Agricultural Water Management*, 150, 67-80.
- Tan, X., Shao, D., Gu, W., and Liu, H. (2015). Field analysis of water and nitrogen fate in lowland paddy fields under different water managements using HYDRUS-1D. *Agricultural Water Management*, 150, 67-80.
- Tan, X., Shao, D., Liu, H., Yang, F., Xiao, C., and Yang, H. (2013). Effects of alternate wetting and drying irrigation on percolation and nitrogen leaching in paddy fields. *Paddy and Water Environment*, 11(1-4), 381-395.
- Tan, X., Shao, D., Liu, H., Yang, F., Xiao, C., and Yang, H. (2013). Effects of alternate wetting and drying irrigation on percolation and nitrogen leaching in paddy fields. *Paddy and Water Environment*, 11(1-4), 381-395.
- Tangang, F. T., and Juneng, L. (2011). Climate projection downscaling for Peninsular Malaysia and Sabah-Sarawak using Hadley Centre PRECIS model. Technical consultation report for National Hydraulic Research Institute of Malaysia (NAHRIM).
- Tavakkoli, A. R., and Oweis, T. Y. (2004). The role of supplemental irrigation and nitrogen in producing bread wheat in the highlands of Iran. *Agricultural Water Management*, 65(3), 225-236.
- Terano, R., Mohamed, Z., Shamsudin, M. N., and Latif, I. A. (2015). Factors influencing intention to adopt sustainable agriculture practices among paddy farmers in Kada, Malaysia. *Asian Journal of Agricultural Research*, 9(5), 268-275.
- Thakur, A. K., Mohanty, R. K., Patil, D. U., and Kumar, A. (2014). Impact of water management on yield and water productivity with system of rice intensification (SRI) and conventional transplanting system in rice. *Paddy and water environment*, 12(4), 413-424.

- Thomas, D. L., Hunt, P. G., and Gilliam, J. W. (1992). Water table management for water quality improvement. *Journal of soil and water conservation*, 47(1), 65-70.
- Tian, H., Chen, G., Liu, M., Zhang, C., Sun, G., Lu, C., and Chappelka, A. (2010). Model estimates of net primary productivity, evapotranspiration, and water use efficiency in the terrestrial ecosystems of the southern United States during 1895–2007. *Forest ecology and management*, 259(7), 1311-1327.
- Tomar, V. S., and O'Toole, J. C. (1980). Design and Testing of a Microlysimeter for Wetland Rice 1. *Agronomy Journal*, 72(4), 689-692.
- Tournebize, J., Watanabe, H., Takagi, K., and Nishimura, T. (2006). The development of a coupled model (PCPF-SWMS) to simulate water flow and pollutant transport in Japanese paddy fields. *Paddy and Water Environment*, 4(1), 39-51.
- Undan, R. C., Phillips, A. L., and Hill, R. W. (1978). Drainage Requirements for Lowland Rice. Paper, (78-2041), 10.
- United States Environmental Protection Agency (USEPA) (2004). Office of Environmental information, Water chemistry laboratory Manual, Washington DC. http://www.epa.gov/owow/monitoring/wsa/WRS_lab_manual.pdf.
- Van Dam, J. C., Huygen, J., Wesseling, J. G., Feddes, R. A., Kabat, P., Van Walsum, P. E. V., and Van Diepen, C. A. (1997). Theory of SWAP version 2.0; Simulation of water flow, solute transport and plant growth in the soil-water-atmosphere-plant environment (No. 71). DLO Winand Staring Centre.
- Van Nguyen, N., and Ferrero, A. (2006). Meeting the challenges of global rice production, 1-9.
- Verma, A. K., Gupta, S. K., and Isaac, R. K. (2014). Calibration and validation of SWAP to simulate conjunctive use of fresh and saline irrigation waters in semi-arid regions. *Environmental Modeling & Assessment*, 19(1), 45-55.
- Vitosh, M. L., Johnson, J. W., and Mengel, D. B. (1995). TH-state Fertilizer Recommendations for Corn, Soybeans, Wheat and Alfalfa. *Archive. lib. msu.edu*.
- Vörösmarty, C. J., Green, P., Salisbury, J., and Lammers, R. B. (2000). Global water resources: vulnerability from climate change and population growth. *science*, 289(5477), 284-288.
- Vörösmarty, C. J., Léveque, C., Revenga, C., Bos, R., Caudill, C., Chilton, J., and Barker, S. (2005). Fresh water. *Millennium ecosystem assessment*, 1, 165-207.

- Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., and Davies, P. M. (2010). Global threats to human water security and river biodiversity. *Nature*, 467(7315), 555.
- Vryzas, Z., Alexoudis, C., Vassiliou, G., Galanis, K., and Papadopoulou-Mourkidou, E. (2011). Determination and aquatic risk assessment of pesticide residues in riparian drainage canals in northeastern Greece. *Ecotoxicology and Environmental Safety*, 74(2), 174-181.
- Vu, S. H., Ishihara, S., and Watanabe, H. (2006). Exposure risk assessment and evaluation of the best management practice for controlling pesticide runoff from paddy fields. Part 1: Paddy watershed monitoring. *Pest Management Science: formerly Pesticide Science*, 62(12), 1193-1206.
- Wah, C. A. (1998). Direct seeded rice in Malaysia: a success story. Asia-Pacific Association of Agricultural Research Institutions.
- Walker, S. H. (1994). Water loss from permanently banded ricefields investigated by field measurement and numerical modelling (Doctoral dissertation, University of Birmingham).
- Walker, S. H., and Rushton, K. R. (1984). Verification of lateral percolation losses from irrigated rice fields by a numerical model. *Journal of hydrology*, 71(3-4), 335-351.
- Wang, D. J., Liu, Q., Lin, J. H., and Sun, R. J. (2004). Optimum nitrogen use and reduced nitrogen loss for production of rice and wheat in the Yangtse Delta region. *Environmental geochemistry and health*, 26(2), 221-227.
- Wang, H., Ju, X., Wei, Y., Li, B., Zhao, L., and Hu, K. (2010). Simulation of bromide and nitrate leaching under heavy rainfall and high-intensity irrigation rates in North China Plain. *Agricultural Water Management*, 97(10), 1646-1654.
- Wang, H., Liu, C., and Zhang, L. (2002). Water-saving agriculture in China: an overview. *Advances in Agronomy*, 75, 135-171.
- Wang, J., Huang, G., Zhan, H., Mohanty, B. P., Zheng, J., Huang, Q., and Xu, X. (2014). Evaluation of soil water dynamics and crop yield under furrow irrigation with a two-dimensional flow and crop growth coupled model. *Agricultural Water Management*, 141, 10-22.
- Wang, L., Liang, T., Chong, Z., and Zhang, C. (2011). Effects of soil type on leaching and runoff transport of rare earth elements and phosphorous in laboratory experiments. *Environmental Science and Pollution Research*, 18(1), 38-45.

- Washington State Department of Ecology (WSDE) (2002). Introduction to water quality index. Retrieved on January 20, 2008, from <http://www.fotsch.org/WQI.htm>.
- Watanabe, H., and Takagi, K. (2000). A simulation model for predicting pesticide concentrations in paddy water and surface soil. I. Model development. *Environmental Technology*, 21(12), 1379-1391.
- Watanabe, H., Inao, K., Vu, S. H., Phong, T. K., Ishihara, S., Takagi, K., and Tournebize, J. (2007). Pesticide exposure assessment in rice paddy areas: a Japanese perspective. *Pesticide Risk Assessment in Rice Paddies: Theory and Practice*, 167-214.
- Watanabe, H., Kakegawa, Y., and Vu, S. H. (2006). Evaluation of the management practice for controlling herbicide runoff from paddy fields using intermittent and spillover-irrigation schemes. *Paddy and Water Environment*, 4(1), 21-28.
- Watanabe, H., Nguyen, M. H. T., Souphasay, K., Vu, S. H., Phong, T. K., Tournebize, J., and Ishihara, S. (2007). Effect of water management practice on pesticide behavior in paddy water. *Agricultural water management*, 88(1), 132-140.
- Webb, T. H., and Lilburne, L. R. (1999). Use of the LEACHM model and the DRASTIC index to map relative risk of groundwater contamination by pesticide leaching. *Journal of Hydrology (New Zealand)*, 271-288.
- West, L. (2006). World water day: a billion people worldwide lack safe drinking water. *About.com*, 03-26.
- WHO (1997) Guidelines for drinking-water quality, V.1, recommendations. World Health Organisation, Geneva.
- Wickham, T. (1972). Predicting yield benefits in lowland rice through a water balance model. *Water managment in Philippine irrigation systems*, 155-181.
- Wickham, T. H. (1978). Water management for lowland rice: water requirements and yield response. *Soils and rice.*, 649-669.
- Williams, M. W., Ritter, A. M., Cheplick, M. J., and Zdinak, C. E. (1999). RICEWQ: Pesticide runoff model for rice crops. Users manual and program documentation. Version, 1(1).
- Willis, T. M., and Black, A. S. (1996). Irrigation increases groundwater recharge in the Macquarie Valley. *Soil Research*, 34(6), 837-847.
- Wopereis, M. C. S., Bouman, B. A. M., Kropff, M. T., Ten Berge, H. F. M., and Maligaya, A. R. (1994). Water use efficiency of flooded rice fields I. Validation of the soil-water balance model SAWAH. *Agricultural Water Management*, 26(4), 277-289.

- Wuzhong, N., Jianping, L., and Zhaoliang, Z. (2007). Occurrence of nitrification-denitrification and gaseous nitrogen loss process in flooded rice soil. *Progress in Natural Science*, 17(1), 6-10.
- Xiao, M., and Chu, L. (2014). The paddy water environmental effect on nitrogen and phosphorus loss on the condition of controlled drainage. In 2014 Montreal, Quebec Canada July 13–July 16, 2014 (p. 1). American Society of Agricultural and Biological Engineers.
- Xie, X. J., Ran, W., Shen, Q. R., Yang, C. Y., Yang, J. J., and Cao, Z. H. (2004). Field studies on ³²P movement and P leaching from flooded paddy soils in the region of Taihu Lake, China. *Environmental geochemistry and health*, 26(2), 237-243.
- Xing, G. X., and Zhu, Z. L. (2000). An assessment of N loss from agricultural fields to the environment in China. *Nutrient Cycling in Agroecosystems*, 57(1), 67-73.
- Xing, G. X., and Zhu, Z. L. (2000). An assessment of N loss from agricultural fields to the environment in China. *Nutrient Cycling in Agroecosystems*, 57(1), 67-73.
- Xue, X., and Hao, M. (2011). Nitrate leaching on loess soils in north-west China: Appropriate fertilizer rates for winter wheat. *Acta Agriculturae Scandinavica, Section B-Soil & Plant Science*, 61(3), 253-263.
- Yamazaki, (1988). Paddy Field Engineering, Asian Institute of Technology, Bangkok, Thailand (, p. 174).
- Yang, C., Huang, G., Chai, Q., and Luo, Z. (2011). Water use and yield of wheat/maize intercropping under alternate irrigation in the oasis field of northwest China. *Field Crops Research*, 124(3), 426-432.
- Yang, S., Peng, S., Hou, H., and , J. (2014). Controlled irrigation and drainage of a rice paddy field reduced global warming potential of its gas emissions. *Archives of Agronomy and Soil Science*, 60(2), 151-161.
- Yang, S., Peng, S., Hou, H., and Xu, J. (2014). Controlled irrigation and drainage of a rice paddy field reduced global warming potential of its gas emissions. *Archives of Agronomy and Soil Science*, 60(2), 151-161.
- Yang, S., Peng, S., Xu, J., He, Y., and Wang, Y. (2015). Effects of water saving irrigation and controlled release nitrogen fertilizer managements on nitrogen losses from paddy fields. *Paddy and water environment*, 13(1), 71-80.
- Yang, S., Peng, S., Xu, J., He, Y., and Wang, Y. (2015). Effects of water saving irrigation and controlled release nitrogen fertilizer managements on nitrogen losses from paddy fields. *Paddy and water environment*, 13(1), 71-80.

- Yashima, S. (1984). Studies on rice double cropping in the Muda area-development of field infrastructure, Tokyo, Japan.
- Yatazawa, M. (1977). Cycling of mineral nutrients in agricultural ecosystems. *Agro-ecosyst*, 4, 177-179.
- Yoshinaga, I., Miura, A., Hitomi, T., Hamada, K., and Shiratani, E. (2007). Runoff nitrogen from a large sized paddy field during a crop period. *Agricultural water management*, 87(2), 217-222.
- Yu, S. E., and Zhang, Z. Y. (2002). Technical system of water-saving irrigation for rice planting in Jiangsu province. *Journal of Hohai University: Natural Sciences*, 30(5), 30-34.
- Yu-Hua, T. I. A. N., Bin, Y. I. N., Lin-Zhang, Y. A. N. G., Shi-Xue, Y. I. N., and Zhao-Liang, Z. H. U. (2007). Nitrogen runoff and leaching losses during rice-wheat rotations in Taihu Lake region, China. *Pedosphere*, 17(4), 445-456.
- Zawawi, M. A. M., Mustapha, S., and Puasa, Z. (2010). Determination of water requirement in a paddy field at Seberang Perak rice cultivation area. *Jl Eng*, 71(4), 32-41.
- Zebarth, B. J., Drury, C. F., Tremblay, N., and Cambouris, A. N. (2009). Opportunities for improved fertilizer nitrogen management in production of arable crops in eastern Canada: A review. *Canadian Journal of Soil Science*, 89(2), 113-132.
- Zerihun, D., Furman, A., Warrick, A. W., and Sanchez, C. A. (2005). Coupled surface-subsurface flow model for improved basin irrigation management. *Journal of irrigation and drainage engineering*, 131(2), 111-128.
- Zhang, H. C., Cao, Z. H., Shen, Q. R., and Wong, M. H. (2003). Effect of phosphate fertilizer application on phosphorus (P) losses from paddy soils in Taihu Lake Region: I. Effect of phosphate fertilizer rate on P losses from paddy soil. *Chemosphere*, 50(6), 695-701.
- ZHANG, J. H., Jian-Li, L. I. U., ZHANG, J. B., CHENG, Y. N., and Wei-Peng, W. A. N. G. (2013). Nitrate-nitrogen dynamics and nitrogen budgets in rice-wheat rotations in Taihu Lake region, China. *Pedosphere*, 23(1), 59-69.
- Zhang, Y. Y., Shao, Q. X., Ye, A. Z., Xing, H. T., and Xia, J. (2016b). Integrated water system simulation by considering hydrological and biogeochemical processes: model development, with parameter sensitivity and autocalibration. *Hydrology and Earth System Sciences*, 20(1), 529.

- Zhang, Y., Zhou, Y., Shao, Q., Liu, H., Lei, Q., Zhai, X., and Wang, X. (2016). Diffuse nutrient losses and the impact factors determining their regional differences in four catchments from North to South China. *Journal of Hydrology*, 543, 577-594.
- Zhang, Z. L., and Xue, J. L. (1996). Study on the intermittent irrigation experiment of rice. *Water Sav Irrig*, 15(6), 23-29.
- Zhao, T. J., and Qiao, X. D. (2010). Suggestions of rice controlled irrigation technology. *Agric Jilin*, 4(05), 122.
- Zhao, X., Xie, Y. X., Xiong, Z. Q., Yan, X. Y., Xing, G. X., and Zhu, Z. L. (2009). Nitrogen fate and environmental consequence in paddy soil under rice-wheat rotation in the Taihu lake region, China. *Plant and soil*, 319(1-2), 225-234.
- Zhao, X., Xie, Y. X., Xiong, Z. Q., Yan, X. Y., Xing, G. X., and Zhu, Z. L. (2009). Nitrogen fate and environmental consequence in paddy soil under rice-wheat rotation in the Taihu lake region, China. *Plant and soil*, 319(1-2), 225-234.
- Zhao, X., Zhou, Y., Min, J., Wang, S., Shi, W., and Xing, G. (2012). Nitrogen runoff dominates water nitrogen pollution from rice-wheat rotation in the Taihu Lake region of China. *Agriculture, ecosystems & environment*, 156, 1-11.
- Zhou, S., Nishiyama, K., Watanabe, Y., and Hosomi, M. (2009). Nitrogen budget and ammonia volatilization in paddy fields fertilized with liquid cattle waste. *Water, air, and soil pollution*, 201(1-4), 135-147.
- Zhou, S., Nishiyama, K., Watanabe, Y., and Hosomi, M. (2009). Nitrogen budget and ammonia volatilization in paddy fields fertilized with liquid cattle waste. *Water, air, and soil pollution*, 201(1-4), 135-147.
- Zhu, T. Y. (1985). Water saving effect and its theory under rice shallow-wet irrigation: North China. *J hydr Eng*, 11, 44-53.
- Zhu, Z., ZHAO-LIANG, X., QiXiao, W., and Freney, J. R. (1997). Mineralization of soil nitrogen. *Nitrogen in soils of China*, 43-66.
- Zhu, Z. L., and Chen, D. L. (2002). Nitrogen fertilizer use in China—Contributions to food production, impacts on the environment and best management strategies. *Nutrient Cycling in Agroecosystems*, 63(2-3), 117-127.
- Zulu, G., Toyota, M., and Misawa, S. I. (1996). Characteristics of water reuse and its effects on paddy irrigation system water balance and the riceland ecosystem. *Agricultural Water Management*, 31(3), 269-283.