



**UNIVERSITI PUTRA MALAYSIA**

***PROPAGATION, GROWTH PERFORMANCE AND YIELD OF  
Piper betle L. AT DIFFERENT NITROGEN AND SHADE LEVELS***

**QUSAY ABDULHAMZA MUTTALEB**

**FP 2018 68**



**PROPAGATION, GROWTH PERFORMANCE AND YIELD OF  
*Piper betle* L. AT DIFFERENT NITROGEN AND SHADE LEVELS**

By

**QUSAY ABDULHAMZA MUTTALEB**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfillment of the Requirements for the Degree of Master of Science**

**February 2018**

## **COPYRIGHT**

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



## DEDICATION

*To my lovely country Iraq*

*To my parents, may God have mercy on them*

*To my wife, brothers, sisters and daughters whose encourage and gave me the power  
to achieve my goal*

*and*

*To my friends who boost me all those past years*



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

**PROPAGATION, GROWTH PERFORMANCE AND YIELD OF  
*Piper betle* L. AT DIFFERENT NITROGEN AND SHADE LEVELS**

By

**QUSAY ABDULHAMZA MUTTALEB**

**February 2018**

**Chairman : Associate Professor Thohirah Lee Abdullah @ Lee Chin Chin, PhD**  
**Faculty : Agriculture**

*Piper betle* L., an evergreen perennial climber from the family *Piperaceae*, is an important medicinal plant in Malaysia. Despite its importance, knowledge on its propagation is still lacking. The present study was conducted to explore the feasibility of *in vitro* and *in vivo* systems of propagation. Sterilization of explants prior to *in vitro* procedures was carried out in 20% Clorox for 20 minutes. *In vitro* culture medium with 1.0 mg/L BAP was the best medium for shoot production. The study also examined *in vivo* rooting of cuttings and growth performance as affected by different indole 3 butyric acid (IBA) treatments and different nitrogen rates (0, 50 100, 150 kg/ha) and shade levels (0%, 30% and 50 %). Morphological parameters were recorded at 30, 60 and 90 days after planting. Semi-hardwood cuttings treated with 2000 mg/L IBA showed the best rooting performance (100%). Plants were significantly taller in treatment at 50 % shade and 100 kg N/ha. At first (H1), second (H2) and third (last) (H3) harvests, maximum height for each harvest was recorded at 70, 105.67 and 184.67 cm respectively. The number of branches was highest at 26.00 from H3, followed by H2 (10.33) and H1 (6.00). The number of leaves from H1, H2 and H3 harvests were 18.00, 46.33 and 171.00 respectively. The total leaf area recorded from H1, H2 and H3 were 653.17, 4108.28 and 10401.17 cm<sup>2</sup> respectively. The fresh weight of leaves were recorded at 18.12 g (H1), 91.07 g (H2) and 185.15 g (H3), while the dry weight of leaves were recorded at 21.55 g (H3) 10.91 g (H2) and 2.40 g (H1), significantly higher at 30 % shade with 100 kg N/ha. With respect to physiological parameters, photosynthesis (PR) and transpiration (TR) rates and stomatal conductance (SC) increased with increasing light intensity with treatment at 30 % shade and 100 kg N/ha giving the highest PR of 10.13  $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ , SC 0.13  $\text{mol}/\text{m}^2/\text{s}$  and TR 1.60  $\mu\text{mol}/\text{m}^2/\text{s}$ . Chlorophyll content (a and ab) did not show any significant difference in the 30% and 50 % shade levels (1.55 and 1.49 mg/cm<sup>2</sup>) but was highest at 100 kg N/ha (1.76 mg/cm<sup>2</sup>). Chlorophyll b was highest at 50% shade which suggests that chlorophyll-a biosynthesis in relation to chlorophyll-b was

significant in low light intensity. Total phenolic content (TPC) (highest at 1.19 mg (GAE)/g), total flavonoid content (TFC) (highest at 128.13 mg Quercetin/g) and antioxidant activities (DPPH) (highest at 85.15%) and FRAP 7.87  $\mu\text{mol Fe(II)}/\text{g dry}$ ) were highly significant in the control treatment.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PEMBIAKAN, PRESTASI PERTUMBUHAN DAN HASIL *Piper betle* L.  
PADA PARAS NITROGEN DAN KETEDUHAN YANG BERBEZA**

Oleh

**QUSAY ABDULHAMZA MUTTALEB**

**Februari 2018**

**Pengerusi : Profesor Madya Thohirah Lee Abdullah @ Lee Chin Chin, PhD**  
**Fakulti : Pertanian**

*Piper betle* L., satu spesies malar hijau pemanjat daripada keluarga Piperaceae, adalah satu spesies perubatan yang penting di Malaysia. Walaupun demikian, pengetahuan mengenai pembiakannya masih berkurangan. Kajian ini dijalankan untuk meneroka kebolehlaksanaan sistem pembiakan in vitro dan in vivo. Pengsterilan eksplan sebelum prosidur in vitro telah dilakukan dengan menggunakan 20% Clorox selama 20 minit. Medium kultur yang mengandungi 1.0 mg/L BAP didapati medium yang terbaik dari segi pengeluaran tunas. Kajian juga meneliti daya pengakaran keratan secara in vivo dan prestasi pertumbuhan ekoran rawatan dengan asid indol 3 butyric (IBA) serta kadar nitrogen (0, 50 100, 150 kg/ha) and paras keteduhan (0%, 30% and 50 %). Parameter morfologi telah dicatat pada 30, 60 and 90 hari selepas penanaman. Keratan separa-matang yang dirawat dengan 2000 mg/L IBA menunjukkan prestasi pengakaran yang terbaik (100%). Tanaman didapati lebih tinggi dengan ketara daripada rawatan 50% keteduhan dan 100 kg N/hektar. Pada tuaian pertama (H1), kedua (H2) dan ketiga (terakhir) (H3), ketinggian maksimum bagi setiap tuaian adalah masing-masing pada 70, 105.67 dan 184.67 cm. Bilangan dahan didapati tertinggi pada H3 (26.00) diikuti dengan H2 (10.33) dan H1 (6.00). Bilangan daun daripada H1, H2 dan H3 adalah masing-masing 18.00, 46.33 dan 171.00. Jumlah keluasan daun daripada H1, H2 and H3 adalah masing-masing 653.17, 4108.28 dan 10401.17 cm<sup>2</sup>. Berat basah daun segar telah direkodkan pada 18.12 (H1), 91.07 (H2) dan 185.15 g (H3), sementara berat kering daun adalah pada 21.55 (H3), 10.91 (H2) dan 2.40 g (H1), iaitu lebih tinggi dan ketara pada 30% keteduhan dengan 100 kg N/ha. Dalam kajian yang sama, keluasan daun spesifik (SLA) telah mencatatkan 272.44 (H1), 376.57 (H2) dan 483.82 cm<sup>2</sup>/g. Kadar pertumbuhan tanaman (CGR) telah merekodkan 0.362 g daripada tuaian 30-60 hari diikuti oleh 1.794 daripada tuaian 60-90 hari dengan rawatan 100 N kg/hektar. Nisbah akar-pucuk (RSR) daripada H1 dan H2 adalah 0.79 g diikuti oleh H3 pada 0.43 g dengan ketinggian yang ketara pada keteduhan 30% dan 100 kg N/hektar. Bagi parameter fisiologi, kadar fotosintesis (PR)

dan transpirasi (TR), serta konduktans stomata (SC) meningkat dengan peningkatan keamatan cahaya dengan rawatan keteduhan 30% dan 100 kg N/hektar dengan mencatatkan nilai tertinggi PR pada  $10.13 \mu\text{mol CO}_2 / \text{m}^2/\text{s}$ , SC  $0.13 \text{ mol}/\text{m}^2/\text{s}$  dan TR  $1.60 \mu\text{mol}/\text{m}^2/\text{s}$ . Kandungan klorofil (a dan b) tidak menunjukkan perbezaan yang ketara diparas keteduhan 30% dan 50% ( $1.55$  dan  $1.49 \text{ mg}/\text{cm}^2$ ), tetapi memuncak tinggi pada  $100 \text{ kg N/hektar}$  ( $1.76 \text{ mg}/\text{cm}^2$ ). Klorofil-b mencapai ketinggian maksimum pada keteduhan 50%, dan ini membawa erti bahawa biosintesis klorofil-a yang berkaitan dengan klorofil- b adalah rendah dengan ketara dalam keamatan cahaya yang rendah. Kandungan fenolik total (TPC) (tertinggi pada  $1.19 \text{ mg(GAE)}/\text{g}$ ), kandungan flavonoid total (TFC) (tertinggi pada  $128.13 \text{ mg Quercetin}/\text{g}$ ) dan aktiviti anti-oksidan (DPPH) (tertinggi pada  $85.15\%$ ) dan FRAP pada  $7.87 \mu\text{mol Fe(II)}/\text{g dry}$ ) adalah ketara pada rawatan kawalan.





## ACKNOWLEDGEMENTS

*In the name of-of Allah the Merciful*

Alhamdulillah, all praise God for his intensity and blessings in the performance of this letter.

Special appreciate goes to my supervisor, Dr. Thohirah Lee Abdullah @ Lee Chin Chin and co-work Dr. Siti Aisha Hassan, and Encik Azmi Abd Rashid, for continued supervision and support. Their valuable help from constructive comments and suggestions throughout the experiments and thesis work contributed to the success of this research. Do not forget, I appreciate Staff, Encik Mat Yusof Suki, Encik Azahar Othman, Encik Mohd Yusoff Mohd Yassin for their support and knowledge. I also extend my thanks to all technicians and staff of the Crop Science Department for their assistance. I would like to thank all my friends, especially Omar, Imad, Firas, Dhualfiqar, Mustafa, Chen, Hilme, Tan, Suzanne, Catherine, and others for their kindness, help and moral support during my studies. Thank you for friendship and memories. Last but not least, my deepest gratitude to my beloved uncle haj Kadhom, brothers, sisters and family for their endless love, help, support, prayer and encouragement. As my wife is wonderful and his daughters are very grateful and appreciate their support, love and care. For those who have contributed indirectly to this research, kindness means a lot to me. Thank you so much.

***Qusay Abdulhamza Muttaleb, 2018***

I certify that a Thesis Examination Committee has met on 12 June 2018 to conduct the final examination of Qusay Abdulhamza Muttaleb on his thesis entitled "Propagation, Growth Performance and Yield of *Piper betle* L. at Different Nitrogen and Shade Levels" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

**Adam b Puteh, PhD**

Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Chairman)

**Puteri Edaroyati bt Megat Wahab, PhD**

Senior Lecturer

Faculty of Agriculture

Universiti Putra Malaysia

(Internal Examiner)

**Asgar Ali, PhD**

Professor

University of Nottingham Malaysia

Malaysia

(External Examiner)



**RUSLI HAJI ABDULLAH, PhD**

Professor and Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 31 October 2018

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Thohirah Lee Abdullah @ Lee Chin Chin, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Siti Aisha Hassan, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Azmi Abd. Rashid, M.Phil, PhD**

Senior Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Name and Matric No: Qusay Abdulhamza Muttaleb, GS42149

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: \_\_\_\_\_  
Name of Chairman  
of Supervisory Committee: Associate Professor  
Dr. Thohirah Lee Abdullah @ Lee Chin Chin

Signature: \_\_\_\_\_  
Name of Member  
of Supervisory Committee: Associate Professor  
Dr. Siti Aisha Hassan

Signature: \_\_\_\_\_  
Name of Member  
of Supervisory Committee: Dr. Azmi Abd. Rashid, M.Phil

## TABLE OF CONTENTS

	Page
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiv
<b>LIST OF FIGURES</b>	xv
<b>LIST OF ABBREVIATIONS</b>	xviii
 <b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	1
 <b>2 LITERATURE REVIEW</b>	3
2.1 Taxonomy of Betel	3
2.2 Distribution and Origin of Betel	3
2.3 Significance of Traditional Medicine	4
2.4 Importance of Medicinal Plants	4
2.4.1 Importance of Betel as a Medicinal Plant	5
2.4.2 Uses of Betel	5
2.4.3 Chemical Constituents in Betel	6
2.4.3.1 Phenol	7
2.4.3.2 Flavonoid	9
2.4.3.3 Antioxidant Properties of Betel	9
2.4.3.4 Antioxidant Activity of Betel	9
2.5 Propagation of Betel	10
2.5.1 Propagation of Betel in Field Conditions	11
2.5.2 Factors Affecting on the Propagation of Betel in Field	11
2.5.2.1 Media	11
2.5.2.2 Light Intensity	11
2.5.2.3 Fertilizers	12
2.5.3 Soil and Foliar Analysis	13
2.5.4 Effect of Different Growth Factors on Physiological Parameters of Betel Yield	14
2.5.5 Effects of Different Growth Factors on Antioxidant Activity of Betel	14
2.6 Micropropagation of Betel	14
2.6.1 The Role of Plant Tissue Culture in Micropropagation of Betel	15
2.6.2 Factors Effecting on Micropropagation	15
2.6.2.1 Explant Source	16
2.6.2.2 Sterilization	16
2.6.2.3 Tissue Culture Media	16
2.6.2.4 Culture Browning	17

2.6.2.5	Plant Growth Hormones	17
2.6.2.6	Response to Media	17
2.6.2.7	Response to Plant Growth Regulators	18
2.6.2.8	Effect of Plant Growth Regulators on Shoots Induction and Multiplication	19
2.6.2.9	Effect of Plant Growth Regulators on Roots Induction	19
2.6.2.10	Histology of Propagated Roots and Stems	20
<b>3</b>	<b>EFFECT OF SHADE LEVELS AND NITROGEN RATES ON THE MORPHO-PHYSIOLOGICAL TRAITS AND SECONDARY METABOLITES OF <i>Piper betle</i> L.</b>	<b>21</b>
3.1	Introduction	21
3.2	Materials and Methods	22
3.2.1	Location of the Experiment	22
3.2.2	Shading	22
3.2.3	Seedlings Preparation	23
3.2.4	The Design of Field Experiment	24
3.2.5	Fertilization and Watering	25
3.2.6	Harvesting	25
3.2.7	Morphological Parameters	25
3.2.7.1	Plant Height (cm)	25
3.2.7.2	Number of Branches	26
3.2.7.3	Number of Leaves	26
3.2.7.4	Fresh Leaves Weight (g)	26
3.2.7.5	Dry Leaves Weight (g)	26
3.2.7.6	Total Leaf Area (cm <sup>2</sup> )	27
3.2.7.7	Specific Leaf Area (SLA) cm <sup>2</sup> /g	27
3.2.7.8	Root-shoot Ratio (RSR)	27
3.2.7.9	Biomass Dry Weight (g)	27
3.2.7.10	Crop Growth Rate (CGR) g/period	28
3.2.7.11	Tissue Analysis	28
3.2.7.12	Media Analysis	28
3.2.8	Physiological Parameters	29
3.2.8.1	Photosynthetic Rate, Conductance and Transpiration Rate	29
3.2.8.2	Chlorophyll Content (mg/g fresh weight)	30
3.2.8.3	Secondary Metabolites	30
3.3	Statistical Analysis	32
3.4	Results	32
3.4.1	Morphological Results	32
3.4.1.1	Plant Height	32
3.4.1.2	Number of Branches	34
3.4.1.3	Number of Leaves	36
3.4.1.4	Fresh Leaf Weight	38
3.4.1.5	Dry Leaf Weight	40
3.4.1.6	Total Leaf Area	42
3.4.1.7	Specific Leaf Area	44



3.4.1.8	Root-shoot Ratio	46
3.4.1.9	Biomass Dry Weight (g)	48
3.4.1.10	Crop Growth Rate (g)	50
3.4.1.11	Tissue Analysis	51
3.4.1.12	Media Analysis	53
3.4.1.13	Correlation Between Morphological Parameters of <i>P. betle</i> as Affected by Shades and Nitrogen Rates	55
3.4.2	Physiological Results	61
3.4.2.1	Photosynthetic Rate (PR)	61
3.4.2.2	Stomatal Conductance (SC)	61
3.4.2.3	Transpiration Rate (TR)	62
3.4.2.4	Chlorophyll Content	63
3.4.2.5	Secondary Metabolites	65
3.5	Discussion	71
3.6	Conclusion	75
<b>4</b>	<b>ROOTING OF STEM CUTTINGS WITH DIFFERENT INDOLE 3 BUTYRIC ACID (IBA) TREATMENTS AND DEVELOPMENT OF MICROPROPAGATION PROTOCOL FOR <i>Piper betle</i> L. USING NODE EXPLANT</b>	<b>77</b>
4.1	Introduction	77
4.2	Materials and Methods	78
4.2.1	Rooting of the Stem Cuttings with Different Indole 3 Butyric Acid (IBA) Treatments	78
4.2.2	Adventitious Root Initiation and Development (Root Anatomy and Histology)	79
4.2.3	Development of Micropropagation Protocol of <i>P. betle</i> Using Node Explant	80
4.2.3.1	Reducing Microbial Contamination and Increasing the Explant Survival by Clorox Treatment	80
4.2.3.2	Determination of the Optimum BAP Concentration in Inducing Shoot Multiplication from the Node Culture of <i>P. betle</i>	80
4.3	Result	80
4.3.1	The Results of the Rooting Study	80
4.3.1.1	The Percentage of Rooting (%)	80
4.3.1.2	The Number of the Roots	81
4.3.1.3	The Fresh Weight of the Roots (g)	81
4.3.1.4	The Dry Weight of the Roots (g)	81
4.3.1.5	The Roots Length (cm)	81
4.3.1.6	The Diameter of the Roots (mm)	82
4.3.1.7	The Surface Area of the Roots (cm <sup>2</sup> )	82
4.3.1.8	The Root Volume (cm <sup>3</sup> )	82
4.3.2	The Results of the Histological Study	84
4.3.3	The Results of Study on Development of Micropropagation Protocol of <i>P. betle</i> Using Node Explant	86



4.3.3.1	The Effect of Different Clorox Concentrations with Different Immersion Times on the Node Explants of <i>P. betle</i>	86
4.3.3.2	The effect of Different Concentrations of Benzyl Amino Purine (BAP) on the Explant growth of <i>P. betle</i>	87
4.4	Discussion	89
4.5	Conclusions	91
<b>5</b>	<b>SUMMARY, GENERAL CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH</b>	<b>93</b>
5.1	Summary	93
5.2	General conclusion	95
5.3	Recommendations for Future Studies	95
	<b>REFERENCES</b>	<b>96</b>
	<b>APPENDICES</b>	<b>125</b>
	<b>BIODATA OF STUDENT</b>	<b>145</b>
	<b>LIST OF PUBLICATIONS</b>	<b>146</b>

## LIST OF TABLES

Table	Page
2.1 Chemical composition of fresh betel leaf	6
2.2 Chemical compounds of <i>P. betle</i> and their uses	8
3.1 The rates and fertilization cycle	25
3.3 Effect of different shade levels and nitrogen rates on nutrients content in leaf tissue of <i>P. betle</i> after 90 days from transplanting	52
3.4 Initial states of media chemical properties, before planting	53
3.5 Media chemical properties after three months period of the experiment	54
3.6 Correlation between all morphological components in <i>P. betle</i> as affected by shades and nitrogen rates for first harvest (30 days)	56
3.7 Correlation between all morphology parameters components in <i>P. betle</i> as affected by shades and nitrogen rates for second harvest (60 days)	58
3.8 Correlation between all morphology parameters components in <i>P. betle</i> as affected by shades and nitrogen rates for third harvest (H3 90 days)	60
3.9 Correlation coefficient between measured physiological parameters	70
4.1 Mean comparison of the root percentages, the root number, the dry root weight and the fresh root weight on of <i>P. betle</i> L. as affected by different concentrations of Indole 3 Butyric Acid (IBA) and the types of cuttings (softwood and semi-hardwood)	83
4.2 Mean comparison of the root length, the root diameter, the rooting surface, the root and the root volume of <i>P. betle</i> L. among different concentrations of Indole 3 Butyric Acid (IBA) and the types of cuttings (softwood and semi-hardwood)	83
4.3 Mean comparison among the treatments for the effect of different concentrations of Clorox with various immersion times for the sterilisation explant of <i>P. betle</i> L.	86
4.4 Mean comparison among the treatments for the effect of different concentrations of benzyl aminopurine (BAP) on the explants of <i>P. betle</i>	88

## LIST OF FIGURES

Figure	Page
2.1 South east asian region (green highlighted) shows the significant areas of <i>P. betle</i> consumption in the world	4
3.1 Experiment layout showing nested design (Rep) Replication, (N) Nitrogen, (N1) control, (N2) 50 kg nitrogen / hectare, (N3) 100 kg nitrogen / hectare, (N4) 150 kg nitrogen / hectare, (o) seedling in plot	22
3.2 Measurement light intensity by portable light meter	23
3.3 Seedlings of <i>P. betle</i> 28 days old used in experiment	23
3.4 Experimental site showing polybags having media and transplanted plant	24
3.5 Media containing soil, coco peat and sand at ratio of 3:2:1 in the polybags	24
3.6 Pruning of main tip shoot	26
3.7 Measurement photosynthesis rate, stomata conductance and transpiration rate by Li-Cor 6400 Portable Photosynthesis system	29
3.8 Effect of different shade levels and nitrogen rates on plant height of <i>P. betle</i> , H1 = Harvest 1 (30 days), H2 = Harvest 2 (60 days) and H3 = Harvest 3 (90 days)	33
3.9 Effect of different shade levels and nitrogen rates on number of brunches of <i>P. betle</i> , H1 = Harvest 1 (30 days), H2 = Harvest 2 (60 days) and H3 = Harvest 3 (90 days)	35
3.10 Effect of different shade levels and nitrogen rates on number of leaves of <i>P. betle</i> , H1 = Harvest 1 (30 days), H2 = Harvest 2 (60 days) and H3 = Harvest 3 (90 days)	37
3.11a Effect of different shade levels and nitrogen rates on fresh leaf weight of leaves of <i>P. betle</i> , Harvest 1 (30 days)	38
3.11b Effect of different shade levels and nitrogen rates on fresh leaf weight of leaves of <i>P. betle</i> , Harvest 2 (60 days)	39
3.11c Effect of different shade levels and nitrogen rates on fresh leaf weight of leaves of <i>P. betle</i> , Harvest 3 (90 days)	of 39

3.11d	Effect of different shade levels and nitrogen rates on fresh leaf weight of leaves of <i>P. betle</i> , total of leaf weight for three harvest	40
3.12	Effect of different shade levels and nitrogen rates on day leaf weight of leaves of <i>P. betle</i> , H1 = Harvest 1 (30 days), H2 = Harvest 2 (60 days) and H3 = Harvest 3 (90 days)	41
3.13	Effect of different shade levels and nitrogen rates on total leaf area of <i>P. betle</i> , H1 = Harvest 1 (30 days), H2 = Harvest 2 (60 days) and H3 = Harvest 3 (90 days)	43
3.14	Effect of different shade levels and nitrogen rates on specific leaf area of <i>P. betle</i> , H1 = Harvest 1 (30 days), H2 = Harvest 2 (60 days) and H3 = Harvest 3 (90 days)	45
3.15	Effect of different shade levels and nitrogen rates on root- shoot ratio of <i>P. betle</i> , H1 = Harvest 1 (30 days), H2 = Harvest 2 (60 days) and H3 = Harvest 3 (90 days)	47
3.16	Effect of different shade levels and nitrogen rates on biomass dry weight of <i>P. betle</i> , H1 = Harvest 1 (30 days), H2 = Harvest 2 (60 days) and H3 = Harvest 3 (90 days)	49
3.17	Effect of different shade levels and nitrogen rates on crop growth rate of <i>P. betle</i> , A = the interval time between harvest 1 and harvest, B = the interval time between harvest 2 and harvest 3	50
3.18	Effect of different nitrogen rates and shade levels on photosynthetic rate (PR), of <i>P. betle</i>	61
3.19	Effect of different N rates and shade levels on stomata conductance (SC) of <i>P. betle</i>	62
3.20	Effect of different nitrogen rates and shade levels on transpiration rate (TR) of <i>P. betle</i>	62
3.21	Effect of different nitrogen rates and shade levels on chlorophyll (a) content in <i>P. betle</i>	63
3.22	Effect of different nitrogen rates and shade levels on chlorophyll (b) content in <i>P. betle</i>	64
3.23	Effect of different nitrogen rates and shade levels on chlorophyll (a + b) content in <i>P. betle</i>	64
3.24	Effect of different Nitrogen rates and shade levels on total phenolic accumulation (TPC) of <i>P. betle</i>	65

3.25	Effect of different Nitrogen rates and shade levels on total flavonoids contents (TFC) of <i>P. betle</i>	66
3.26	Effect of different Nitrogen rates and shade levels on 1,1-Diphenyl-2-picryl-hydrazyl (DPPH) of <i>P. betle</i>	67
3.27	Effect of different Nitrogen rates and shade levels on Ferric Reducing Antioxidant Power (FRAP) of <i>P. betle</i>	68
4.1	Rooting media (sand at 9 kg) in trays (size of 25×40 cm) is used for rooting of stem cuttings	78
4.2	The Effect different IBA concentrations on the types of cuttings. A: Softwood cuttings and B: Semi hardwood cuttings after 28 days of treatment	84
4.3	Cross section of the stem of <i>P. betle</i> L. (X4) showing, Epidermis (E) cortex (Cx). Endodermis (Ed), phloem (Ph), vascular cambium (Cb), xylem (X), and pith (Pi). A: Cross section of stem 0 day of softwood. B: Cross section of stem 0 day of semi-hardwood. C: Cross section of the stem 3 days of softwood showing differentiation to root initials occurring near the vascular bundle (arrow). D: Cross section of the stem 3 days of semi-hardwood showing development to root initials occurring near the vascular bundle (arrow). E: Cross section of the stem 6 days of softwood showing some development of root initials organized root primordial (Rp) (arrow). F: Cross section of the stem 6 days of semi-hardwood showing subsequent development of root initials organized root primordial (Rp) (arrow)	85
4.4	The effect of different Clorox concentrations and different immersion times on <i>P. betle</i> explants after five weeks of culture. (A) Fungal contamination, (B) Bacterial contaminations and Browning, (C) Dead explant, and (D) Survived explants without microbial condemnation	87
4.5	The impact of the BAP in combination with shoot regeneration of <i>P. betle</i> . (A & C) Number of the explant with new leaves (arrow), (B) Percentage of the axillary buds (arrow), and (D) Number of leaves (arrow) produced per explant after ten weeks of culture	88
4.6	Rooting explant of <i>P betle</i> in (0.0 MS)	89

## LIST OF ABBREVIATIONS

DPPH	1,1-Diphenyl-2-picrylhydrazyl
TPTZ	2,4, 6-tri-(2-pyridyl)-striozone
AlCl <sub>3</sub>	Aluminum chloride
ANOVA	Analysis of Variance
BAP	benzylaminopurine
BDW	biomass dry weight
Ca	Calcium
CEC	Cation Exchange Capacity
cm mol/kg	centi mol per kilogram
cm	centimeter
cm <sup>3</sup>	centimeter cubic
cm <sup>2</sup>	centimeter square
cm <sup>2</sup> /g	centimeter square per gram
Chl (a+b)	Chlorophyll (a + b)
Chl (a)	Chlorophyll (a)
Chl (b)	Chlorophyll (b)
pH	concentration of H <sup>+</sup>
CGR	Crop Growth Rate
DAT	Days After Transplanting
°C	Degree centigrade
DLW	Dry Leaf Weight
F	Fertilization
FCR	Folin-Ciocalteu reagent
FAF	formalin: acetic acid

FRAP	Free Reducing Antioxidant Potential
FLW	Fresh Leaf Weight
GA	Gallic acid
g	gram
g/kg	gram per kilogram
g/L	gram per liter
H	Harvest
ha	hectare
h	hour
H <sub>2</sub> O <sub>2</sub>	Hydrogen peroxide
IBA	Indole-3-butyric acid
FeCl <sub>2</sub>	Iron (II) chloride
L	liter
Mg	Magnesium
µg	microgram
µg/L	microgram per liter
µml	micromillimeter
µmol/m <sup>2</sup> /s	Micromole per meter square per second
µmol/mol	Micromole per mol
Mg	milligram
ml/L	milligram per liter
mm	millimeter
mm mole	millimole
mmolHCL	millimole Hydrochloric acid
mmol/m <sup>2</sup> /s	Millimole per meter square per second
MS	Murashige and Skoog

N	Nitrogen
NB	Number of Branches
NL	Number of Leaves
%shade	percentage of shade
P	Phosphorus
PR	Photosynthetic Rate
PH	Plant Height
K	Potassium
KCl	Potassium Chloride
RCBD	Randomized Complete Block Design
RSR	Root-Shoot Ratio
NaOH	Sodium hydroxide
SLA	Specific Leaf Area
SD	Standard Deviation
SAS	Statistical Analysis Software
G <sub>s</sub>	Stomata Conductance
TBA	tertiary butyl-alcohol
TFC	Total Flavanones Content
TLA	Total Leaf Area
TPC	Total Phenolics Content
E	Transpiration rate
V	volume



## CHAPTER 1

### INTRODUCTION

Human interest in herbal medicine is increasing day by day. Awareness regarding the potentiality of herbal medicine in curing certain physiological disorders in human has increase day by day after observing the side effect of synthetic drugs (Gopalakrishnan *et al.*, 2012). Herbal medicines are not new for health practices and it has been used since human civilization existed (Alsarhan *et al.*, 2014). It played an imperative role to treat the different types of human ailments (Sharma *et al.*, 2013). According to the report of World Health Organization (2003), 80% of the world population still relies on the traditional/ herbal medicine for their primary health care. These medicinal plants or phytomedicine refers to the use of any plant part such as seeds, leaves, bark or flowers for curing physiological disorders in human. The plants are found in different regions of the world, with a few of these have already been discovered and millions are still needed to be explored.

In Southern Asian region, Malaysia is a forest enrich country possessing a variety of plants in its agroecosystem. These plants can be used as herbal medicine, however, few of these have already play a part in traditional Malaysian health care system due to their therapeutic efficacy (Alsarhan *et al.*, 2014). In this connection, *Piper betle* Linn from the Piperaceae family is one of the most precious medicinal herbs found in central and eastern Malaysia. It is commonly known as “Sirih” in the native language, and is second to tea and coffee based on daily consumption in Southern Asian region (Alsarhan *et al.*, 2014). In spite of its medicinal properties, *P. betle* is commercially cultivated for its leaves. The leaves are used as masticator for chewing purpose in many parts of the world especially in the tropical and sub-tropical countries of the world (Pradhan *et al.*, 2014).

Piperaceae family is already very peculiar as it contains well-known plants such as black pepper and kava. There are about 100 varieties of *P. betle* found throughout the world and out of this 40 are found only in India and 30 in West Bengal (Guha, 1997; Samanta, 1994). However, the most probable place of origin of betel vine is Malaysia (Chattopadhyay and Maiti, 1967). This plant is perennial and dioecious in nature, with evergreen and shade loving creeping characteristics. Every part of the plant has high medicinal value and several attributes such as digestive, carminative, stimulant, antiseptic and antifungal activities have been described. A phenolic compound, hydroxyl-chavicol, with anti-carcinogenic property has also been identified in *P. betle* leaves (Verma *et al.*, 2004).

Stem cuttings either as semi hardwood or softwood cuttings are among the most probable techniques for vegetative propagation in most plants. (Tchoundjeu *et al.*, 2004; Zlesak, 2012). This technique can also play a significant role in the clonal propagation of *P. betle*. The triggering of adventitious root initiation and development

in stem cuttings treated with auxins is well discussed in many plants (Erturk *et al.*, 2010; Hartmann *et al.*, 2010; Bolat, 1995). Nevertheless, exogenous plant growth regulator such as auxin plays an important role in inducing rooting in stem cutting. Among the auxin that can induce rooting in stem cutting include indole 3 butyric acid (IBA) (Hartmann *et al.*, 2010; Yahya *et al.*, 2009). In *P. betle* no report has been made on the use of IBA in stimulating root formation from stem cutting.

On the other hand, in order to supply the active compounds for pharmaceutical and herbal industries, tissue culturing of medicinal plants is widely used. The culturing techniques nowadays have been often observed in conservation of genetic material of many vulnerable medicinal plants (Sidhu, 2011). In that case, micro-propagation is a good technique in order to increase the vegetative growth. Furthermore, this process insures a good regular supply of medicinal plants as multiplication from plants tissues or seeds occur with minimum space and time (Prakash and Van Staden, 2007). Furthermore, the less abundance of sunlight in Malaysia due to cloudy weather throughout the year can also reduce the phenol and flavonoid content of *P. betle* that do not only affect the yield but lower down the plant medicinal properties too. Due to this, efforts should be taken to increase the biomass by improving plant vegetative growth by application of high nutrient especially nitrogen to the soil at the vegetative growth phase of *P. betle*.

Currently, not much research has been carried out to solve the above problem. Therefore, this study was conducted with the objective:

- i. To determine the effect of shade levels and nitrogen rates on the morphological traits, growth and yield of *P. betle*.
- ii. To determine the role of shade and nitrogen on physiological traits and secondary metabolites of *Piper betle* L.
- iii. To determine the types of cuttings and concentration of IBA in rooting of stem cuttings of *P. betle* and to develop micropropagation protocol for *P. betle* using node explant.

## REFERENCES

- Abad, M., Noguera, P., Puchades, R., Maquieira, A., and Noguera, V. (2002). Physico-chemical and chemical properties of some coconut coir dusts for use as a peat substitute for containerised ornamental plants. *Bioresource technology*, 82(3), 241-245.
- Abbasi, B. H., Khan, M. A., Mahmood, T., Ahmad, M., Chaudhary, M. F., and Khan, M. A. (2010). Shoot regeneration and free-radical scavenging activity in *Silybum marianum* L. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 101(3), 371-376.
- Abdullah, T., Misrol, M., Stanslas, J., Abd Aziz, M., and Goh, L. (2014). Effect of shade and morphological characterization of *Janggut Adam* (*Tacca sp.*) as a promising native ornamental and medicinal plant. *Transanction of the Malaysian Society of Plant Physiology*, 22, 13-15.
- Agarwal, T., Singh, R., Shukla, A. D., Waris, I., and Gujrati, A. (2012). Comparative analysis of antibacterial activity of four *Piper betle* varieties. *Adv Appl Sc Res*, 3, 698-705.
- Ahmad, I., Mehmood, Z., and Mohammad, F. (1998). Screening of some Indian medicinal plants for their antimicrobial properties. *Journal of ethnopharmacology*, 62(2), 183-193.
- Ahmad, N., Fazal, H., Abbasi, B. H., Rashid, M., Mahmood, T., and Fatima, N. (2010). Efficient regeneration and antioxidant potential in regenerated tissues of *Piper nigrum* L. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 102(1), 129-134.
- Ahuja, S., and Ahuja, U. (2011). Betel Leaf and Betel Nut in India: History and Uses. *Asian Agri-History*, 15(1).
- Ajith, A., Parani, M., Rao, C., Latha, R., and Balakrishna, P. (1997). Micropropagation and genetic fidelity studies in *Piper longum* L. *Biotechnology of Spices, Medicinal and Aromatic Plants, Indian Society for Spices*, Calicut, India, 94-97.
- Ajithkumar, D., and Seeni, S. (1998). Rapid clonal multiplication through in vitro axillary shoot proliferation of *Aegle marmelos* (L.) Corr., a medicinal tree. *Plant Cell Reports*, 17(5), 422-426.
- Ajithkumar, K., Jayachandran, B., and Ravi, V. (2002). Influence of shade regimes on photosynthetic rate and stomatal characters of ginger (*Zingiber officinale* R.). *Journal of Spices and Aromatic Crops*, 11(1), 26-29.

- Akbar, A., and SHAJAHAN, A. (2014). Somatic embryogenesis through cell suspension culture in Indian commercial cultivars of banana (*Musa Spp.*). *Doctor Thesis, Bharathidasan University, India*
- Ali, S., and Mehta, R. (1968). Preliminary pharmacological and anthelmintic studies of the essential oil of *Piper betle* L.
- Al-Khayri, J. M., and Al-Bahrany, A. M. (2001). In vitro micropropagation of *Citrus aurantifolia* (lime). *Current Science*, 1242-1246.
- Almaliotis, D., Therios, I., and Karatassiou, M. (1996). Effects of nitrogen fertilization on growth, leaf nutrient concentration and photosynthesis in three peach cultivars. *Paper presented at the II International Symposium on Irrigation of Horticultural Crops* 449, 529-534.
- Alsarhan, A., Sultana, N., Al-Khatib, A., and Kadir, M. R. A. (2014). Review on some Malaysian traditional medicinal plants with therapeutic properties. *Journal of Basic & Applied Sciences*, 10, 149.
- Alvira, P., Tomás-Pejó, E., Ballesteros, M., and Negro, M. (2010). Pretreatment technologies for an efficient bioethanol production process based on enzymatic hydrolysis: a review. *Bioresource technology*, 101(13), 4851-4861.
- Amalia, H., Sitompul, R., Hutaaruk, J., Andrianjah, A., and Mun'im, A. (2016). Effectiveness of *Piper betle* leaf infusion as a palpebral skin antiseptic. *Universa Medicina*, 28(2), 83-91.
- Ammirato, P. (1988). Role of ABA in the regulation of somatic embryogenesis. *Hort. Science*, 23(3), 520-520.
- Amonkar, A., Nagabhushan, M., D'souza, A., and Bhide, S. (1986). Hydroxychavicol: a new phenolic antimutagen from betel leaf. *Food and chemical toxicology*, 24(12), 1321-1324.
- Anjaria, J., Parabia, M., and Dwivedi, S. (2002). Ethnovet heritage–Indian ethnoveterinary medicine an overview. *Pathik Enterprise, Ahmedabad, India*.
- Antolovich, M., Prenzler, P. D., Patsalides, E., McDonald, S., and Robards, K. (2002). Methods for testing antioxidant activity. *Analyst*, 127(1), 183-198.
- Arambewela, L., Arawwawala, L., and Ratnasooriya, W. (2005). Antidiabetic activities of aqueous and ethanolic extracts of *Piper betle* leaves in rats. *Journal of Ethnopharmacology*, 102(2), 239-245.
- Arambewela, L., Arawwawala, M., and Rajapaksa, D. (2006). *Piper betle*: a potential natural antioxidant. *International journal of food science & technology*, 41(s1), 10-14.

- Asai, H., Samson, B. K., Stephan, H. M., Songyikhangsuthor, K., Homma, K., Kiyono, Y., et al. (2009). Biochar amendment techniques for upland rice production in Northern Laos: 1. Soil physical properties, leaf SPAD and grain yield. *Field Crops Research*, 111(1), 81-84.
- Ashish, M., Huma, A., and Maheshwari, M. (2009). Antioxidant activity of ethanolic extract of *Piper betel* leaves. *Journal of Pharmacy Research*, 2(3), 491-494.
- Babaei, N., Abdullah, N. A. P., Saleh, G., and Abdullah, T. L. (2013). Control of contamination and explant browning in *Curculigo latifolia* *In vitro* cultures. *Journal of Medicinal Plants Research*, 7(8), 448-454.
- Babu, K. N., Rema, J., Pillai, G. S., Divakaran, M., Ravindran, P., and Pete, K. (1992). Micropropagation of betelvine (*Piper betle* L.). *Journal of Spices and Aromatic Crops*, 1(2), 160-162.
- Babu, K. N., Rema, J., Pillai, S. R. D., Samsudeen, K., and Ravindran, P. (1996). Micropropagation of an endangered species of *Piper barberi* gamble and its conservation. *Indian Journal of Plant Genetic Resources*, 9(1), 179-182.
- Badjatia, N., Satyam, A., Singh, P., Seth, A., and Sharma, A. (2010). Altered antioxidant status and lipid peroxidation in Indian patients with urothelial bladder carcinoma. In *Urologic Oncology: Seminars and Original Investigations*, 360-367.
- Bailey, S., Walters, R. G., Jansson, S., and Horton, P. (2001). Acclimation of *Arabidopsis thaliana* to the light environment: the existence of separate low light and high light responses. *Planta*, 213(5), 794-801.
- Bajji, M., Kinet, J.-M., and Lutts, S. (2002). The use of the electrolyte leakage method for assessing cell membrane stability as a water stress tolerance test in durum wheat. *Plant Growth Regulation*, 36(1), 61-70.
- Baksha, R., Jahan, M. A. A., Khatun, R., and Munshi, J. L. (2005). Micropropagation of *Aloe barbadensis* Mill. through in vitro culture of shoot tip explants. *Plant Tissue Cult Biotechnol*, 15(2), 121-126.
- Balaraju, K., Agastian, P., and Ignacimuthu, S. (2009). Micropropagation of *Swertia chirata* buch.-Hams. ex Wall.: a critically endangered medicinal herb. *Acta physiologiae plantarum*, 31(3), 487-494.
- Balbuena, T. S., Santa-Catarina, C., Silveira, V., Kato, M. J., and Floh, E. I. S. (2009). *In vitro* morphogenesis and cell suspension culture establishment in *Piper solmsianum* C. DC.(Piperaceae). *Acta Botanica Brasilica*, 23(1), 274-281.
- Baligar, V., and Fageria, N. (1999). Plant nutrient efficiency: towards the second paradigm. *Siqueira, jo; moreira, fms; lopes, as; guilherme, lrg*, 183-204.



- Basak, S., and Guha, P. (2015). Modelling the effect of essential oil of betel leaf (*Piper betle* L.) on germination, growth, and apparent lag time of *Penicillium expansum* on semi-synthetic media. *International journal of food microbiology*, 215, 171-178.
- Basak, U., Das, A., and Das, P. (2000). Rooting response in stem cuttings from five species of mangrove trees: effect of auxins and enzyme activities. *Marine Biology*, 136(1), 185-189.
- Basra, A. (2000). *Plant growth regulators in agriculture and horticulture: their role and commercial uses*: CRC Press.
- Basra, S. M., Iqbal, S., and Afzal, I. (2014). Evaluating the response of nitrogen application on growth, development and yield of quinoa genotypes. *International Journal of Agriculture & Biology*, 16(5), 886-892.
- Beadle, C., Coombs, J., Hall, D., LONG, S., and SCURLOCK, J. (1987). Techniques in bioproductivity and photosynthesis. *Techniques in bioproductivity and photosynthesis*. Pergamon Oxford
- Behn, H., Albert, A., Marx, F., Noga, G., and Ulbrich, A. (2010). Ultraviolet-B and photosynthetically active radiation interactively affect yield and pattern of monoterpenes in leaves of peppermint (*Mentha × piperita* L.). *Journal of agricultural and food chemistry*, 58(12), 7361-7367.
- Bekheet, S., Taha, H., and Gabr, A. (2013). Protocol for in vitro morphogenesis and hairy root cultures of milk thistle (*Silybum marianum* L. Gaertn). *Journal of applied sciences research*, 9(1), 860-866.
- Bergquist, S. Å., Gertsson, U. E., Nordmark, L. Y., and Olsson, M. E. (2007). Ascorbic acid, carotenoids, and visual quality of baby spinach as affected by shade netting and postharvest storage. *Journal of agricultural and food chemistry*, 55(21), 8444-8451.
- Bhat, S., Chandel, K., and Malik, S. (1995). Plant regeneration from various explants of cultivated *Piper* species. *Plant Cell Reports*, 14(6), 398-402.
- Bhattacharya, S., Banerjee, D., Bauri, A., Chattopadhyay, S., and Bandyopadhyay, S. (2007). Healing property of the Piper betel phenol, allylpyrocatechol against indomethacin-induced stomach ulceration and mechanism of action. *World Journal of Gastroenterology: WJG*, 13(27), 3705.
- Bhide, S., Zariwala, M., Amonkar, A., and Azuine, M. (1991). Chemopreventive efficacy of a betel leaf extract against benzo [a] pyrene-induced forestomach tumors in mice. *Journal of ethnopharmacology*, 34(2-3), 207-213.
- Bhojwani, S. S., and Razdan, M. K. (1986). *Plant tissue culture: theory and practice* (Vol. 5): Elsevier.

- Bhojwani, S., and Razdan, M. (1996). Haploid production. Plant tissue culture: theory and practice, a revised edition. *Amsterdam: Elsevier Science BV*, 167-213.
- Blevins, D. G. (1994). Uptake, translocation, and function of essential mineral elements in crop plants. *Physiology and Determination of Crop Yield. Madison: ASA, CSSA, SSSA*, 259-275.
- Bojović, B., and Stojanović, J. (2006). Some wheat leaf characteristics in dependence of fertilization. *Kragujevac journal of Science*, 28, 139-146.
- Bolat, I. (1995). The effect of the date of fall budding on bud-take and the quality on shoot development from the inserted bud in apricot. In 2. *Turkish National Horticulture Congress, Adana (Turkey)*, 3-6 Oct 1996. Cukurova University, Faculty of Agriculture.
- Bonfill, M., Cusidó, R. M., Palazón, J., Piñol, M. T., and Morales, C. (2002). Influence of auxins on organogenesis and ginsenoside production in *Panax ginseng* calluses. *Plant cell, tissue and organ culture*, 68(1), 73-78.
- Boro, P., Shorma Deka, A., and Kalita, M. (1998). Clonal propagation of *alternanthera sessilis*: A *biopharmaceutically* potent herbal medicinal plant. *J. Phytol. Res*, 11, 103-106.
- Bosland, P. W., Votava, E. J., and Votava, E. M. (2012). *Peppers: vegetable and spice capsicums* (Vol. 22): *Cabi*.
- Bouza, L., Jacques, M., and Miginiac, E. (1994). *In vitro* propagation of *Paeonia suffruticosa* Andr. cv. 'Mme de Vatry': developmental effects of exogenous hormones during the multiplication phase. *Scientia Horticulturae*, 57(3), 241-251.
- Braca, A., De Tommasi, N., Di Bari, L., Pizza, C., Politi, M., and Morelli, I. (2001). Antioxidant principles from *bauhinia tarapotensis*. *Journal of natural products*, 64(7), 892-895.
- Brand-Williams, W., Cuvelier, M.-E., and Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT-Food science and Technology*, 28(1), 25-30.
- Briskin, D. P., and Gawienowski, M. C. (2001). Differential effects of light and nitrogen on production of hypericins and leaf glands in *Hypericum perforatum*. *Plant physiology and Biochemistry*, 39(12), 1075-1081.
- Casado, J., Navarro, M., Utrilla, M., Martinez, A., and Jimenez, J. (2002). Micropropagation of *Santolina canescens* Lagasca and *In vitro* volatiles production by shoot explants. *Plant cell, tissue and organ culture*, 69(2), 147-153.

- Casey, C. A., Mangan, F. X., Herbert, S. J., Barker, A. V., and Carter, A. K. (2002). The effect of light intensity and nitrogen fertilization on plant growth and leaf quality of ngo gai (*Eryngium foetidum* L.) in Massachusetts. *Paper presented at the XXVI International Horticultural Congress: The Future for Medicinal and Aromatic Plants* 629, 215-229.
- Chakraborty, D., and Shah, B. (2011). Antimicrobial, antioxidative and antihemolytic activity of Piper betel leaf extracts. *International Journal of Pharmacy and Pharmaceutical Sciences*, 3(3), 192-199.
- Chalupa, V. (1977). Organogenesis in Norway spruce and Douglas fir tissue cultures. *Commun Inst For Cech*, 10, 79-87.
- Chan, H. (1978). Soil survey interpretations for improved rubber production in peninsular Malaysia. Soil-resources data for agricultural development. *Hawaii Agricultural Experiment Station, Honolulu, Hawaii, USA*, 41-66.
- Chatterjee, A., and Pakrashi, S. C. (1991). The treatise on Indian medicinal plants: vol. 1. New Delhi: Publications and Information Directorate, CSIR 172p.-illus., col. illus. ISBN 8172360118 *En Icones. Includes authentic Sanskrit slokas in both Devnagri and Roman scripts. Plant records. Geog*, 6.
- Chattopadhyay, S. B., and Maiti, S. (1967). *Diseases of betel vine and spices*: Indian Council of Agricultural Research Krishi Anusandhan Bhavan Pusa; New Delhi.
- Chaturvedi, R., Razdan, M., and Bhojwani, S. (2004). *In vitro* clonal propagation of an adult tree of neem (*Azadirachta indica* A. Juss.) by forced axillary branching. *Plant Science*, 166(2), 501-506.
- Chaveerach, A., Mookkamul, P., Sudmoon, R., and Tanee, T. (2006). Ethnobotany of the genus Piper (Piperaceae) in Thailand. *Ethnobotany Research & Applications*, 4:223-231
- Chawla, H. (2002). Introduction to plant biotechnology: Science Publishers. *International Book Distributing, Lucknow, India*, pp 39–56.
- Chowdhury, M., and Johri, J. (2003). Seasonal variations in chlorophyll content and chlorophyllase activity in Bangla and Mitha varieties of betel vine (*Piper betle* L.) grown in different soil treatments. *Plant Physiol.*, 48:115- 119.
- Chu, I. Y. (1992). Perspective of micropropagation industry. In *Transplant production systems* (pp. 137-150): Springer.
- Croteau, R. (1987). Biosynthesis and catabolism of monoterpenoids. *Chemical Reviews*, 87(5), 929-954.
- CSIR, New Delhi (1969). Council of Scientific and Industrial Research, New Delhi. *The Wealth of India*, 8, 84-94.



- Cushnie, T. T., and Lamb, A. J. (2005). Antimicrobial activity of flavonoids. *International journal of antimicrobial agents*, 26(5), 343-356.
- Dai, Y., Shen, Z., Liu, Y., Wang, L., Hannaway, D., and Lu, H. (2009). Effects of shade treatments on the photosynthetic capacity, chlorophyll fluorescence, and chlorophyll content of *Tetrastigma hemsleyanum* Diels et Gilg. *Environmental and experimental botany*, 65(2), 177-182.
- Daniel, M., and Daniel, M. (1991). Methods in plant chemistry and economic botany: *Kalyani publisher New Delhi, India*. 104
- Das, S., Parida, R., Sandeep, I. S., Nayak, S., and Mohanty, S. (2016). Biotechnological intervention in betelvine (*Piper betle* L.): A review on recent advances and future prospects. *Asian Pacific journal of tropical medicine*, 9(10), 938-946.
- Datta, A., Ghoshdastidar, S., and Singh, M. (2011). Antimicrobial property of Piper betel leaf against clinical isolates of bacteria. *International Journal of Pharma Sciences and Research*, 2(3), 104-109.
- Davies Jr, F., Lazarte, J., and Joiner, J. (1982). Initiation and development of roots in juvenile and mature leaf bud cuttings of *Ficus pumila* L. *American Journal of Botany*, 804-811.
- Davies, F. (1984). Shoot RNA, cambial activity and indolebutyric acid effectivity in seasonal rooting of juvenile and mature *Ficus pumila* cuttings. *Physiologia Plantarum*, 62(4), 571-575.
- Davies, F. T. (1978). A histological and physiological analysis of adventitious root formation in juvenile and mature cuttings of *Ficus pumila* L. *PhD Thesis. University of Florida*.
- Davies, P. J. (2010). The plant hormones: their nature, occurrence, and functions. In *Plant hormones* (pp. 1-15): Springer.
- De Klerk, G.-J. (2002). Rooting of micro cuttings: theory and practice. *In Vitro Cellular & Developmental Biology-Plant*, 38(5), 415-422.
- de Paiva Neto, V. B., and Otoni, W. C. (2003). Carbon sources and their osmotic potential in plant tissue culture: does it matter? *Scientia Horticulture*, 97(3), 193-202.
- Devadas, M., and Chandini, S. (2000). Growth of bush pepper (*Piper nigrum* L.) plants as influenced by light and nutrients. *Journal of Spices and Aromatic Crops*, 9(2), 105-109.
- Dhote, S., Devi, P. U., Pathak, A., and Goswami, R. (2007). Studies on Anti-genotoxic Effect of *Piper betle* Leaves. *Journal of Natural Remedies*, 7(2), 247-251.

- Dwivedi, V., and Tripathi, S. (2014). Review study on potential activity of *Piper betle*. *J Pharmacogn Phytochem*, 3(4), 93-98.
- Epstein, E., and Bloom, A. (2005). Inorganic components of plants. Mineral nutrition of plants: principles and perspectives, 2nd edn. Sinauer Associates, Inc., Massachusetts, 44-45.
- Epstein, E., and Ludwig- Müller, J. (1993). Indole- 3- butyric acid in plants: occurrence, synthesis, metabolism and transport. *Physiologia plantarum*, 88(2), 382-389.
- Erturk, Y., Ercisli, S., Haznedar, A., and Cakmakci, R. (2010). Effects of plant growth promoting rhizobacteria (PGPR) on rooting and root growth of kiwifruit (*Actinidia deliciosa*) stem cuttings. *Biological research*, 43(1), 91-98.
- Etampawala, E., Tennakoon, K., Gunatilleke, C., and Gunatilleke, I. (2002). Studies on propagation, optimal growth conditions and fruit formation of the medicinal plant *Piper longum* L. *Ceylan Journal of Sciences (Biological Sciences)*, 30, 67-77.
- Evans, J., and Poorter, H. (2001). Photosynthetic acclimation of plants to growth irradiance: the relative importance of specific leaf area and nitrogen partitioning in maximizing carbon gain. *Plant, Cell & Environment*, 24(8), 755-767.
- Evans, P. H., Bowers, W. S., and Funk, E. J. (1984). Identification of fungicidal and nematocidal components in the leaves of *Piper betle* (Piperaceae). *Journal of Agricultural and Food chemistry*, 32(6), 1254-1256.
- Fageria, N. (2005). Soil fertility and plant nutrition research under controlled conditions: basic principles and methodology. *Journal of plant nutrition*, 28(11), 1975-1999.
- Fageria, N. (2007). Soil fertility and plant nutrition research under field conditions: Basic principles and methodology. *Journal of plant nutrition*, 30(2), 203-223.
- Fageria, N. K., and Baligar, V. C. (2003). Fertility management of tropical acid soils for sustainable crop production. *Handbook of soil acidity*, 359-385.
- Fageria, N., and Baligar, V. (2005). Enhancing nitrogen use efficiency in crop plants. *Advances in agronomy*, 88, 97-185.
- Fageria, N., and Baligar, V. (2008). Ameliorating soil acidity of tropical Oxisols by liming for sustainable crop production. *Advances in agronomy*, 99, 345-399.
- Fageria, N., Baligar, V., and Bailey, B. (2005). Role of cover crops in improving soil and row crop productivity. *Communications in soil science and plant analysis*, 36(19-20), 2733-2757.

- Fageria, N., Baligar, V., and Clark, R. (2002). Micronutrients in crop production. *Advances in Agronomy*, 77, 185-268.
- Fageria, N., Baligar, V., and Li, Y. (2008). The role of nutrient efficient plants in improving crop yields in the twenty first century. *Journal of plant nutrition*, 31(6), 1121-1157.
- Fageria, N., Filho, M. B., Moreira, A., and Guimarães, C. (2009). Foliar fertilization of crop plants. *Journal of plant nutrition*, 32(6), 1044-1064.
- Falkiner, F. (1990). The criteria for choosing an antibiotic for control of bacteria in plant tissue culture. *International Association for Plant Tissue Culture*, 60: 13-23
- Fauziah, C.I., Jamilah, I. and S. R. Syed Omar. 1997. An Evolution of cation exchange capacity methods for acid tropical soils. *Pertanika J. Trop. Agric. Sci.* 20(2/3): 113-119
- Ferreira, I. C., Baptista, P., Vilas-Boas, M., and Barros, L. (2007). Free-radical scavenging capacity and reducing power of wild edible mushrooms from northeast Portugal: Individual cap and stipe activity. *Food chemistry*, 100(4), 1511-1516.
- Firuzi, O., Lacanna, A., Petrucci, R., Marrosu, G., and Saso, L. (2005). Evaluation of the antioxidant activity of flavonoids by “ferric reducing antioxidant power” assay and cyclic voltammetry. *Biochimica et Biophysica Acta (BBA)-General Subjects*, 1721(1), 174-184.
- Fitchet, M. (1990). Establishment of *P. nigrum* in vitro. *Acta Horti*, 257, 285-291.
- Fraternal, D., Giamperi, L., Ricci, D., and Rocchi, M. (2002). Micropropagation of *Bupleurum fruticosum*: The effect of triacontanol. *Plant cell, tissue and organ culture*, 69(2), 135-140.
- Gagnon, B., and Berrouard, S. (1994). Effects of several organic fertilizers on growth of greenhouse tomato transplants. *Canadian journal of plant science*, 74(1), 167-168.
- Galal, M., Bashir, A., Salih, A., and Adam, S. (1991). Activity of water extracts of *Albizzia anthelmintica* and *A. lebbek* barks against experimental *Hymenolepis diminuta* infection in rats. *Journal of ethnopharmacology*, 31(3), 333-337.
- Gardner, P. T., White, T. A., McPhail, D. B., and Duthie, G. G. (2000). The relative contributions of vitamin C, carotenoids and phenolics to the antioxidant potential of fruit juices. *Food chemistry*, 68(4), 471-474.
- Garro-Monge, G., Gatica-Arias, A. M., and Valdez-Melara, M. (2008). Somatic embryogenesis, plant regeneration and acemannan detection aloe (*Aloe barbadensis* MILL.). *Agronomia Costarricense*, 32(2).

- Gebicka, L., and Banasiak, E. (2009). Flavonoids as reductants of ferryl hemoglobin. *Acta Biochim Pol*, 56(3), 509-513.
- George, E. F., and Sherrington, P. D. (1984). *Plant propagation by tissue culture: Exegetics Ltd. In Handbook and Directory of Commercial Laboratories, Basingstoke England.*, pp. 223- 227
- George, E. F., Hall, M. A., and De Klerk, G.-J. (2007). *Plant propagation by tissue culture: volume 1. the background* (Vol. 1): Springer Science & Business Media.
- Ghasemzadeh, A., and Ghasemzadeh, N. (2011). Effects of shading on synthesis and accumulation of polyphenolic compounds in ginger (*Zingiber officinale* Roscoe) varieties. *Journal of Medicinal Plants Research*, 5(11), 2435-2441.
- Ghasemzadeh, A., Jaafar, H. Z., and Rahmat, A. (2011). Effects of solvent type on phenolics and flavonoids content and antioxidant activities in two varieties of young ginger (*Zingiber officinale* Roscoe) extracts. *Journal of Medicinal Plants Research*, 5(7), 1147-1154.
- Ghasemzadeh, A., Jaafar, H. Z., Rahmat, A., Wahab, P. E. M., and Halim, M. R. A. (2010). Effect of different light intensities on total phenolics and flavonoids synthesis and anti-oxidant activities in young ginger varieties (*Zingiber officinale* Roscoe). *International Journal of Molecular Sciences*, 11(10), 3885-3897.
- Gilani, A. H. (2005). Trends in ethnopharmacology. *Journal of ethnopharmacology*, 100(1), 43-49.
- Girouard, R. M. (1967). Initiation and development of adventitious roots in stem cuttings of *Hedera helix*: anatomical studies of the Juvenile growth phase. *Canadian Journal of Botany*, 45(10), 1877-1881.
- Gogtay, N., Bhatt, H., Dalvi, S., and Kshirsagar, N. (2002). The use and safety of non-allopathic Indian medicines. *Drug safety*, 25(14), 1005-1019.
- Gopalakrishnan, S., Rajameena, R., and Vadivel, E. (2012). Antimicrobial activity of the leaves of *Myxopyrum serratum* AW Hill. *Int. J. Pharm. Sci. Drug Res*, 4(1), 31-34.
- Gopalan, C., Ramasastri, B.V. and Balasubramanian, S.C (1984). Nutritive Value of Indian Foods, *National Institute of Nutrition (ICMR), Hyderabad, India* pp. 108.
- Graham, T. L. (1998). Flavonoid and flavonol glycoside metabolism in *Arabidopsis*. *Plant Physiology and Biochemistry*, 36(1-2), 135-144.

- Gu, X.-D., Sun, M.-Y., Zhang, L., Fu, H.-W., Cui, L., Chen, R.-Z., et al. (2010). UV-B induced changes in the secondary metabolites of *Morus alba* L. leaves. *Molecules*, 15(5), 2980-2993.
- Guha, P. (1997). Paan Theke Kutir Silpa Sambhabana. Exploring Betel Leaves for Cottage Industry”, In: *Krishi, Khadya-O-Gramin Bikash Mela–A Booklet published by the Agricultural and Food Engineering Department, IIT, Kharagpur, India*, 15-19.
- Guha, P. (2006). Betel leaf: the neglected green gold of India. *J Hum Ecol*, 19(2), 87-93.
- Guha, P. and Jain, R.K (1997). Processing and Marketing of Betel Leaf (*Piper betle* L.). *Agricultural and Food Engineering Department, IIT, Kharagpur, India*
- Guha, P., and Jain, R. (1997). Status report on production, processing and marketing of betel leaf (*Piper betle* L.). *Agricultural and Food Engineering Department, IIT, Kharagpur, India*.
- Gundala, S. R., Yang, C., Mukkavilli, R., Paranjpe, R., Brahmabhatt, M., Pannu, V., et al. (2014). Hydroxychavicol, a betel leaf component, inhibits prostate cancer through ROS-driven DNA damage and apoptosis. *Toxicology and applied pharmacology*, 280(1), 86-96.
- Haboudane, D., Miller, J. R., Tremblay, N., Zarco-Tejada, P. J., and Dextraze, L. (2002). Integrated narrow-band vegetation indices for prediction of crop chlorophyll content for application to precision agriculture. *Remote sensing of environment*, 81(2), 416-426.
- Haissig, B. E., and Davis, T. D. (1994). A historical evaluation of adventitious rooting research to 1993. In *Biology of adventitious root formation* (pp. 275-331): Springer.
- Halliwell, B. (1991). Drug antioxidant effects. *Drugs*, 42(4), 569-605.
- Hammerschlag, F. (1982). Factors influencing in vitro multiplication and rooting of the plum rootstock Myrobalan (*Prunus cerasifera* Ehrh.). *Journal American Society for Horticultural Science*.
- Han, H., Zhang, S., and Sun, X. (2009). A review on the molecular mechanism of plants rooting modulated by auxin. *African Journal of Biotechnology*, 8(3).
- Harbage, J. F., and Stimart, D. P. (1996). Effect of pH and 1H-indole-3-butyric acid (IBA) on rooting of apple micro cuttings. *Journal of the American Society for Horticultural Science*, 121(6), 1049-1053.
- Harbage, J. F., Stimart, D. P., and Evert, R. F. (1993). Anatomy of Adventitious Root Formation in Micro cuttings of *Malus domestica* Borkh. Gala'. *Journal of the American Society for Horticultural Science*, 118(5), 680-688.



- Harborne, J. B. (2014). Introduction to ecological biochemistry: *Academic press*. London.
- Harper, J. (1994). Nitrogen metabolism. Physiology and determination of crop yield (*physiology and de*), 285-302.
- Hartmann, H. T., Kester, D. E., and DAVIES, J. (2002). FT; GENEVE, RL Plant propagation: principles and practices: *New Jersey: Prentice-Hall. Englewood Cliffs, NJ*, pp 336-337
- Hartmann, H. T., Kester, D. E., Davies, F. T., and Geneve, R. L. (1997). Plant propagation: principles and practices: *Prentice-Hall Inc. London*, pp. 770
- Hassan, S. A., Mijin, S., Yusoff, U. K., Ding, P., and Wahab, P. E. M. (2012). Nitrate, ascorbic acid, mineral and antioxidant activities of *Cosmos caudatus* in response to organic and mineral-based fertilizer rates. *Molecules*, 17(7), 7843-7853.
- Hegde, N., Patil, S., and Shashidhar, V. (2010). Effect of organic nutrition on the performance of betel vine (*Piper betle* L.) 'Ambadi'. In *XXVIII International Horticultural Congress on Science and Horticulture for People (IHC2010): International Symposium on 933*, 273-278.
- Hemm, M. R., Rider, S. D., Ogas, J., Murry, D. J., and Chapple, C. (2004). Light induces phenylpropanoid metabolism in *Arabidopsis* roots. *The Plant Journal*, 38(5), 765-778.
- Hollman, P. C., van Trijp, J. M., Buysman, M. N., Mengelers, M. J., de Vries, J. H., and Katan, M. B. (1997). Relative bioavailability of the antioxidant flavonoid quercetin from various foods in man. *FEBS letters*, 418(1-2), 152-156.
- Hu, C., and Wang, P. (1983). Meristem, shoot tip and bud cultures. In: *Handbook of plant cell culture (USA)*, Vol 1., p. 91.
- Huang, D., Ou, B., and Prior, R. L. (2005). The chemistry behind antioxidant capacity assays. *Journal of agricultural and food chemistry*, 53(6), 1841-1856.
- Hussain, A., Naz, S., Nazir, H., and Shinwari, Z. K. (2011). Tissue culture of black pepper (*Piper nigrum* L.) in Pakistan. *Pak. J. Bot*, 43(2), 1069-1078.
- Hussain, A., Qarshi, I. A., Nazir, H., and Ullah, I. (2012). Plant tissue culture: current status and opportunities. In *Recent advances in plant in vitro culture*: In Tech, Rijeka, pp 1–28, ISBN:978-953-51-0787-3
- Ibrahim, M. H., and Jaafar, H. Z. (2012). Primary, secondary metabolites, H<sub>2</sub>O<sub>2</sub>, malondialdehyde and photosynthetic responses of *Orthosiphon stimaneus* Benth. to different irradiance levels. *Molecules*, 17(2), 1159-1176.

- Ibrahim, M. H., Jaafar, H. Z., Karimi, E., and Ghasemzadeh, A. (2013). Impact of organic and inorganic fertilizers application on the phytochemical and antioxidant activity of Kacip Fatimah (*Labisia pumila Benth*). *Molecules*, 18(9), 10973-10988.
- Igamberdiev, A. U., Shen, T., and Gardeström, P. (2006). Function of mitochondria during the transition of barley protoplasts from low light to high light. *Planta*, 224(1), 196-204.
- Imam, M., Pariari, A., and SAHU, K. (2012). Effect of integrated nitrogen application on growth and yield of betel vine [*Piper betle* L.]. *Journal of Crop and Weed*, 8(1), 141-144.
- Ingersoll, C. M., Niesenbaum, R. A., Weigle, C. E., and Lehman, J. H. (2010). Total phenolics and individual phenolic acids vary with light environment in *Lindera benzoin*. *Botany*, 88(11), 1007-1010.
- Jaafar, H. Z. E., Mohd Hafiz, I., & Philip, E. (2009). Leaf gas exchange properties of three varieties of *Labisia pumila Benth*. under greenhouse conditions. *J. Trop. Plant Physiol*, 3, 16-24.
- Jagetia, G. C., and Baliga, M. S. (2004). The evaluation of nitric oxide scavenging activity of certain Indian medicinal plants in vitro: a preliminary study. *Journal of Medicinal Food*, 7(3), 343-348.
- Jamil Mohammad, M., and Zuraiqi, S. (2002). Enhancement of Yield and Nitrogen and Water Use Efficiencies by Nitrogen Drip- Fertigation of Garlic. *Journal of plant nutrition*, 26(9), 1749-1766.
- Jana, B. (1995). Betel leaf: A Cash Crop of Villages of Bengal. *Asaboni*, Flat-203, 184.
- Jana, B. (1996). Improved technology for betel leaf cultivation. In *A paper presented in the "Seminar-cum-Workshop on Betel leaf Marketing", held at State cashew nut farm, Directorate of Agricultural Marketing, Digha, Midnapur (WB), India.*
- Jarvis, B., Ali, A., and Shaheed, A. (1983). Auxin and boron in relation to the rooting response and ageing of mung bean cuttings. *New phytologist*, 95(4), 509-518.
- Jaynes, D., Colvin, T., Karlen, D., Cambardella, C., and Meek, D. (2001). Nitrate loss in subsurface drainage as affected by nitrogen fertilizer rate. *Journal of Environmental Quality*, 30(4), 1305-1314.
- Jeng, J.-H., Chen, S.-Y., Liao, C.-H., Tung, Y.-Y., Lin, B.-R., Hahn, L.-J., et al. (2002). Modulation of platelet aggregation by areca nut and betel leaf ingredients: roles of reactive oxygen species and cyclooxygenase. *Free Radical Biology and Medicine*, 32(9), 860-871.

- Jensen, W. A. W. A. (1962). *Botanical histochemistry: principles and practice* (No. QK 861. J46).
- Jeong, K. Y., Pasian, C. C., and Tay, D. (2006). Response of six Begonia species to different shading levels. In *A XXVII International Horticultural Congress-IHC2006: International Symposium on Advances in Environmental Control, Automation* 761, 215-220.
- Jeong, K. Y., Pasian, C. C., McMahon, M., and Tay, D. (2009). Growth of six Begonia species under shading. *The Open Horticulture Journal*, 2, 22-28.
- Johansen, D. A. (1940). *Plant microtechnique*: McGraw-Hill Book Company, Inc: London; p. 530.
- Johkan, M., Shoji, K., Goto, F., Hashida, S.-n., and Yoshihara, T. (2010). Blue light-emitting diode light irradiation of seedlings improves seedling quality and growth after transplanting in red leaf lettuce. *HortScience*, 45(12), 1809-1814.
- Jones Jr, J. B. (2001a). *Laboratory guide for conducting soil tests and plant analysis*. CRC press; p 212 - 213.
- Jones Jr, J. B. (2001c). Soil Analysis (Testing). *Laboratory guide for conducting soil tests and plant analysis*. CRC press; p 81-82.
- Jones, Jr. J. B. (2001b). Chapter 2. Soil Analysis (Testing). *Laboratory Guide for Conducting Soil Tests and Plant Analysis*. CRCP press; p 68-69.
- Joseph, L., Nazeem, P., Mini, S., Philip, S., and Balachandran, M. (1996). *In vitro* techniques for mass multiplication of black pepper (*Piper nigrum* L.) and ex vitro performance of the plantlets. *J. Plantation Crops*, 511-516.
- Kabata-Pendias, A. (2010). *Trace elements in soils and plants*: CRC press, p. (5) 52.
- Kaipiainen, E. (2009). Parameters of photosynthesis light curve in *Salix dasyclados* and their changes during the growth season. *Russian journal of plant physiology*, 56(4), 445-453.
- Kalyoncu, I., Ersoy, N., Yilmaz, M., and Aydın, M. (2009). Effects of humidity level and IBA dose application on the softwood top cuttings of white mulberry (*Morus alba* L.) and black mulberry (*Morus nigra* L.) types. *African Journal of Biotechnology*, 8(16).
- Karimi, E., Jaafar, H. Z., Ghasemzadeh, A., and Ibrahim, M. H. (2013). Light intensity effects on production and antioxidant activity of flavonoids and phenolic compounds in leaves, stems and roots of three varieties of *Labisia pumila* Benth. *Australian Journal of Crop Science*, 7(7), 1016.



- Karuppusamy, S. (2009). A review on trends in production of secondary metabolites from higher plants by *in vitro* tissue, organ and cell cultures. *Journal of Medicinal Plants Research*, 3(13), 1222-1239.
- Kaur, C., and Kapoor, H. C. (2001). Antioxidants in fruits and vegetables—the millennium's health. *International journal of food science & technology*, 36(7), 703-725.
- Kevers, C., Hausman, J.-F., Faivre-Rampant, O., Evers, D., and Gaspar, T. (1997). Hormonal control of adventitious rooting: progress and questions. *Journal of Applied Botany*, 71(3-4), 71-79.
- Khan, S. R., Rose, R., Haase, D. L., and Sabin, T. E. (2000). Effects of shade on morphology, chlorophyll concentration, and chlorophyll fluorescence of four Pacific Northwest conifer species. *New forests*, 19(2), 171-186.
- Kintzios, S., Drossopoulos, J., and Lymperopoulos, C. (2001). Effect of vitamins and inorganic micronutrients on callus growth and somatic embryogenesis from leaves of *chilli pepper*. *Plant cell, tissue and organ culture*, 67(1), 55-62.
- Kitajima, K., Fox, A. M., Sato, T., and Nagamatsu, D. (2006). Cultivar selection prior to introduction may increase invasiveness: evidence from *Ardisia crenata*. *Biological Invasions*, 8(7), 1471-1482.
- Kochhar, S., Kochhar, V., Singh, S., Katiyar, R., and Pushpangadan, P. (2005). Differential rooting and sprouting behaviour of two *Jatropha* species and associated physiological and biochemical changes. *Current science*, 89(6), 936-939.
- Košir, P., Škof, S., and Luthar, Z. (2004). Direct shoot regeneration from nodes of *Phalaenopsis* orchids. *Acta agriculturae slovenica*, 83(2), 233-242.
- Kovar, J. L., and Kuchenbuch, R. O. (1994). Commercial importance of adventitious rooting to agronomy. In *Biology of adventitious root formation* (pp. 25-35): Springer.
- Kozai, T., Xiao, Y., Nguyen, Q. T., Afreen, F., and Zobayed, S. M. (2005). Photoautotrophic (sugar-free medium) micropropagation systems for large-scale commercialization. *Propagation of ornamental plants*, 5(1), 23-34.
- Kumar, K. S., Dahms, H.-U., Lee, J.-S., Kim, H. C., Lee, W. C., and Shin, K.-H. (2014). Algal photosynthetic responses to toxic metals and herbicides assessed by chlorophyll a fluorescence. *Ecotoxicology and environmental safety*, 104, 51-71.
- Kumar, N. (1999). (02) Betel vine (*Piper betle* L.) Cultivation: A Unique Case of Plant Establishment Under Anthropogenically Regulated Microclimatic Conditions, 34(1), pp.19-32.

- Kumar, N., Misra, P., Dube, A., Bhattacharya, S., Dikshit, M., and Ranade, S. (2010). *Piper betle* Linn. a maligned Pan-Asiatic plant with an array of pharmacological activities and prospects for drug discovery. *Current science*, 99(7), 922-932.
- Kumar, R. N., Chakraborty, S., and Kumar, N. J. (2012). Influence of light and developmental stages on active principles of *Andrographis paniculata* (Burm. f.) Wall, ex Nees. *Indian Journal of Scientific Research*, 3(1), 91.
- Kumaranatunga, K. (2003). Gas chromatographical and antimicrobial studies on *Alpinia calcarata* and *Piper betle* from Sri Lanka. *M. M.Phil. Thesis, University of Kelaniya, Sri Lanka*.
- Labrooy, C. D., Abdullah, T. L., Abdullah, N. A. P., and Stanslas, J. (2016). Optimum shade enhances growth and 5, 7-Dimethoxyflavone accumulation in *Kaempferia parviflora* Wall. ex Baker cultivars. *Scientia Horticulturae*, 213, 346-353.
- Lau, F. C., Shukitt-Hale, B., and Joseph, J. A. (2005). The beneficial effects of fruit polyphenols on brain aging. *Neurobiology of aging*, 26(1), 128-132.
- Lean, L. P., and Mohamed, S. (1999). Antioxidative and antimycotic effects of turmeric, lemon- grass, betel leaves, clove, black pepper leaves and *Garcinia atriviridis* on butter cakes. *Journal of the Science of Food and Agriculture*, 79(13), 1817-1822.
- LeBude, A. V. (2005). Adventitious Rooting and Physiology of Stem Cuttings of Loblolly Pine.
- Lei, D., Chan, C.-P., Wang, Y.-J., Wang, T.-M., Lin, B.-R., Huang, C.-H., et al. (2003). Antioxidative and antiplatelet effects of aqueous inflorescence *Piper betle* extract. *Journal of agricultural and food chemistry*, 51(7), 2083-2088.
- Leifert, C., and Cassells, A. (2001). Microbial hazards in plant tissue and cell cultures. *In Vitro Cellular & Developmental Biology-Plant*, 37(2), 133-138.
- Leifert, C., Waites, W., and Nicholas, J. (1989). Bacterial contaminants of micropropagated plant cultures. *Journal of Applied Microbiology*, 67(4), 353-361.
- Letey, J. (1958). Relationship between soil physical properties and crop production. *In Advances in soil science* (pp. 277-294): Springer.
- Lin, Y. J., and Zhang, Q. (2005). Optimising the tissue culture conditions for high efficiency transformation of indica rice. *Plant cell reports*, 23(8), 540-547.
- Linsmaier, E. M., and Skoog, F. (1965). Organic growth factor requirements of tobacco tissue cultures. *Physiologia plantarum*, 18(1), 100-127.

- Liu, C.-z., Guo, C., Wang, Y.-c., and Ouyang, F. (2002). Effect of light irradiation on hairy root growth and artemisinin biosynthesis of *Artemisia annua* L. *Process Biochemistry*, 38(4), 581-585.
- Lloyd, G., and McCown, B. (1980). Commercially-feasible micropropagation of mountain laurel, *Kalmia latifolia*, by use of shoot-tip culture. *Commercially-feasible micropropagation of mountain laurel, Kalmia latifolia, by use of shoot-tip culture.*, 30, 421-427.
- Lobo, V., Patil, A., Phatak, A., and Chandra, N. (2010). Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacognosy reviews*, 4(8), 118.
- Lovell, P. H., and White, J. (1986). Anatomical changes during adventitious root formation. *New Root Formation in Plants and Cuttings. Martinus Nijhoff Publishers, Dordrecht, The Netherlands*, 111-140.
- Ludwig-Müller, J. (2000). Indole-3-butyric acid in plant growth and development. *Plant Growth Regulation*, 32(2), 219-230.
- Ludwig-Müller, J. (2003). Peroxidase isoenzymes as markers for the rooting ability of easy-to-root and difficult-to-root *Grevillea* species and cultivars of *Protea obtusifolia* (Proteaceae). *In Vitro Cellular & Developmental Biology-Plant*, 39(4), 377-383.
- Ludwig-Müller, J., Vertocnik, A., and Town, C. D. (2005). Analysis of indole-3-butyric acid-induced adventitious root formation on *Arabidopsis* stem segments. *Journal of experimental botany*, 56(418), 2095-2105.
- Macheix, J.-J., and Fleuriet, A. (1990). *Fruit phenolics*: CRC press. Inc. p 1–98.
- Maene, L., and Debergh, P. (1985). Liquid medium additions to established tissue cultures to improve elongation and rooting *in vivo*. *Plant cell, tissue and organ culture*, 5(1), 23-33.
- Mahfouz, S., and Sharaf-Eldin, M. (2007). Effect of mineral vs. biofertilizer on growth, yield, and essential oil content of fennel (*Foeniculum vulgare* Mill.). *International Agrophysics*, 21(4), 361.
- Maisuthisakul, P. (2008). Phenolic antioxidants from betel leaf (*Piper betel* Linn.) extract obtained with different solvents and extraction time. *University of the Thai Chamber of Commerce Journal*, 28(2).
- Maiti, S., Kakam, A. S., Sengupta, K., Punekar, L. K., Das, J. N., Saikia, L., & Mahadeva Reddy, K. (1995). Effect of sources and levels of nitrogen on growth and yield of betelvine (*Piper betle* L.). *JOURNAL OF PLANTATION CROPS*, 23, 122-125.

- Makino, A. (2011). Photosynthesis, grain yield, and nitrogen utilization in rice and wheat. *Plant physiology*, 155(1), 125-129.
- Malaysia, P. (2012). Optimization of spray drying process parameters of *Piper betle* L.(Sirih) leaves extract coated with maltodextrin. *Journal of Chemical and Pharmaceutical Research*, 4(3), 1833-1841.
- Manickavasagam, M., Ganapathi, A., Anbazhagan, V., Sudhakar, B., Selvaraj, N., Vasudevan, A., et al. (2004). Agrobacterium-mediated genetic transformation and development of herbicide-resistant sugarcane (*Saccharum* species hybrids) using axillary buds. *Plant cell reports*, 23(3), 134-143.
- Mantell, S. H., Matthews, J., and McKee, R. (1985). *Principles of plant biotechnology: an introduction to genetic engineering in plants*: Blackwell Scientific Publications, Oxford, UK., 57- 130
- Mathai, C. K. (1983). Growth and yield analysis in black pepper varieties (*Piper nigrum* L.) under different light conditions. *PhD Thesis, University of Agricultural Sciences, Bangalore*.
- Mathews, H. (1987). Morphogenetic responses from in vitro cultured seedling explants of mung bean (*Vigna radiata* L. Wilczek). *Plant cell, tissue and organ culture*, 11(3), 233-240.
- Matkowski, A. (2008). Plant in vitro culture for the production of antioxidants—a review. *Biotechnology advances*, 26(6), 548-560.
- McKey, D., Elias, M., Pujol, B., and Duputié, A. (2010). The evolutionary ecology of clonally propagated domesticated plants. *New Phytologist*, 186(2), 318-332.
- Medany, M., Hassanein, M., and Farag, A. (2008). Effect of black and white nets as alternative covers to sweet pepper production under greenhouses in Egypt. In *International Symposium on Strategies Towards Sustainability of Protected Cultivation in Mild Winter Climate* 807, 121-126.
- Mengel, K., Kirkby, E. A., Kosegarten, H., and Appel, T. (2001). Further elements of importance. In *Principles of plant nutrition*: Springer (pp. 639-655).
- Middleton, E., Kandaswami, C., and Theoharides, T. C. (2000). The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease, and cancer. *Pharmacological reviews*, 52(4), 673-751.
- Mihaljević, I., Dugalić, K., Tomaš, V., Viljevac, M., Pranjić, A., Čmelik, Z., et al. (2013). *In vitro* sterilization procedures for micropropagation of 'Oblačinska' sour cherry. *Journal of Agricultural Sciences, Belgrade*, 58(2), 117-126.

- Miliauskas, G., Venskutonis, P., and Van Beek, T. (2004). Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food chemistry*, 85(2), 231-237.
- Miller, A., and Cramer, M. (2005). Root nitrogen acquisition and assimilation. *In Root Physiology: from Gene to Function*, Springer (pp. 1-36).
- Misra, P., and Shukla, P. K. (2010). In vitro approaches for mass production of medicinal plants. *International E-Publication*.
- Misra, P., Datta, S., Johri, J., Singh, H., and Srivastava, A. (2004). An improved method for *in vitro* large scale propagation of *Piper betle* L. *Journal of plant biochemistry and biotechnology*, 13(2), 161-164.
- Mitra, G. (2017). Essential Plant Nutrients and Recent Concepts about their Uptake. *In Essential Plant Nutrients* (pp. 3-36): Springer.
- M'kada, J., Dorion, N., and Bigot, C. (1991). *In vitro* propagation of *Cistus* × *purpureus* Lam. *Scientia horticultruae*, 46(1-2), 155-160.
- Mohnen, D. (1994). Novel experimental systems for determining cellular competence and determination. *In Biology of adventitious root formation*, Springer (pp. 87-98).
- Mollah, M. (2016). Investigation on the leaf rot and foot and root rot of betel vine (*Piper betle* L.). *In satkhira district of bangladesh*. Doctoral dissertation.
- Molnár, Z., Virág, E., and Ordog, V. (2011). Natural substances in tissue culture media of higher plants. *Acta Biologica Szegediensis*, 55(1), 123-127.
- Mori, K., Goto-Yamamoto, N., Kitayama, M., and Hashizume, K. (2007). Loss of anthocyanins in red-wine grape under high temperature. *Journal of experimental botany*, 58(8), 1935-1945.
- Morte, M. A., and Honrubia, M. (1997). Micropropagation of *Helianthemum almeriense*. *In High-tech and micropropagation*, Springer Berlin Heidelberg VI (pp. 163-177).
- Mosaleeyanon, K., Zobayed, S., Afreen, F., and Kozai, T. (2005). Relationships between net photosynthetic rate and secondary metabolite contents in St. John's wort. *Plant Science*, 169(3), 523-531.
- Mukherjee, K., Paul, P., and Banerjee, E. R. (2014). Free radical scavenging activities of Date Palm (*Phoenix sylvestris*) fruit extracts. *Nat. Prod. Chem. Res*, 2(151), 10.4172.



- Mula, S., Banerjee, D., Patro, B. S., Bhattacharya, S., Barik, A., Bandyopadhyay, S. K., et al. (2008). Inhibitory property of the Piper betel phenolics against photosensitization-induced biological damages. *Bioorganic & medicinal chemistry*, 16(6), 2932-2938.
- Muleo, R., and Thomas, B. (1997). Effects of light quality on shoot proliferation of *Prunus cerasifera* in vitro are the result of differential effects on bud induction and apical dominance. *Journal of Horticultural Science*, 72(3), 483-499.
- Murashige, T., and Skoog, F. (1962). A revised medium for rapid growth and bioassays with *tobacco* tissue cultures. *Physiologia plantarum*, 15(3), 473-497.
- Murashige, T. (1974). Plant propagation through tissue cultures. *Annual review of plant physiology*, 25(1), 135-166.
- Murphy, J., and Riley, J. P. (1962). A modified single solution method for the determination of phosphate in natural waters. *Analytica chimica acta*, 27, 31-36.
- Nagabhushan, M., Amonkar, A., Nair, U., D'Souza, A., and Bhide, S. (1989). Hydroxychavicol: a new anti-nitrosating phenolic compound from betel leaf. *Mutagenesis*, 4(3), 200-204.
- Nagori, K., Singh, M. K., Alexander, A., Kumar, T., Dewangan, D., Badwaik, H., et al. (2011). *Piper betle* L.: A review on its ethnobotany, phytochemistry, pharmacological profile and profiling by new hyphenated technique DART-MS (Direct Analysis in Real Time Mass Spectrometry). *Journal of Pharmacy Research* Vol, 4(9), 2991-2997.
- Naija, S., Elloumi, N., Jbir, N., Ammar, S., and Kevers, C. (2008). Anatomical and biochemical changes during adventitious rooting of apple rootstocks MM 106 cultured *in vitro*. *Comptes rendus biologiques*, 331(7), 518-525.
- Nemeth, G. (1986). Induction of rooting. *In Trees I*, Springer, (pp. 49-64).
- Nishimura, A., Ashikari, M., Lin, S., Takashi, T., Angeles, E. R., Yamamoto, T., et al. (2005). Isolation of a rice regeneration quantitative trait loci gene and its application to transformation systems. *Proceedings of the National Academy of Sciences of the United States of America*, 102(33), 11940-11944.
- Nitsch, J., and Nitsch, C. (1969). Haploid plants from pollen grains. *Science*, 163(3862), 85-87.
- Noor Camellia, N. A., Thohirah, L. A., Abdullah, N. A. P., & Mohd Khidir, O. (2009). Improvement on rooting quality of *Jatropha curcas* using indole butyric acid (IBA). *Research Journal of Agriculture and Biological Sciences*, 5 (4), 338-343.

- O'Brien, T. P., and McCully, M. E. (1981). The study of plant structure: principles and selected methods. *Melbourne, Termarcarphi Pty. Ltd.*
- Oktay, M., Gülçin, İ., and Küfrevioğlu, Ö. İ. (2003). Determination of *in vitro* antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts. *LWT-Food Science and Technology*, 36(2), 263-271.
- Organization, W. H. (2002). WHO (2001). *The world health report*.
- Pan, J., and Guo, B. (2016). Effects of light intensity on the growth, photosynthetic characteristics, and flavonoid content of *Epimedium pseudowushanense* BL Guo. *Molecules*, 21(11), 1475.
- Parr, A. J., and Bolwell, G. P. (2000). Phenols in the plant and in man. The potential for possible nutritional enhancement of the diet by modifying the phenols content or profile. *Journal of the Science of Food and Agriculture*, 80(7), 985-1012.
- Patel, J. B., Patel, V. J., & Patel, J. R. (2006). Influence of different methods of irrigation and nitrogen levels on crop growth rate and yield of maize (*Zea mays* L.). *Ind. J. Crop Sci*, 1(1-2), 175-177.
- Pathak, M. R., and Abido, M. S. (2014). The role of biotechnology in the conservation of biodiversity. *Journal of Experimental Biology*, 2, 4.
- Pati, P. K., Rath, S. P., Sharma, M., Sood, A., and Ahuja, P. S. (2006). In vitro propagation of rose—a review. *Biotechnology advances*, 24(1), 94-114.
- Pawar, S., Kalyankar, V., Dhamangaonkar, B., Dagade, S., and Waghmode, S. (2017). Biochemical profiling of antifungal activity of Betel leaf (*Piper betle* L.) extract and its significance in traditional medicine. *J Adv Res Biotech*, 2(1), 4.
- Perez-Vizcaino, F., and Duarte, J. (2010). Flavonols and cardiovascular disease. *Molecular aspects of medicine*, 31(6), 478-494.
- Pham-Huy, L. A., He, H., and Pham-Huy, C. (2008). Free radicals, antioxidants in disease and health. *International journal of biomedical science: IJBS*, 4(2), 89.
- Philip, V., Joseph, D., Triggs, G., and Dickinson, N. (1992). Micropropagation of black pepper (*Piper nigrum* Linn.) through shoot tip cultures. *Plant Cell Reports*, 12(1), 41-44.
- Pierik, R. (1987). In vitro culture of higher plants Martinus Nijhoff Publishers Dordrecht. Printed in Netherlands, 107.
- Pierik, R. L. M. (1997). *In vitro culture of higher plants*: Springer Science & Business Media.
- Podsędek, A. (2007). Natural antioxidants and antioxidant capacity of Brassica vegetables: A review. *LWT-Food Science and Technology*, 40(1), 1-11.



- Prabhu, M., Platel, K., Saraswathi, G., and Srinivasan, K. (1995). Effect of orally administered betel leaf (*Piper betle* Linn.) on digestive enzymes of pancreas and intestinal mucosa and on bile production in rats. *Indian journal of experimental biology*, 33(10), 752-756.
- Pradhan, D., Biswasroy, P., & Suri, K. A. (2014). Variation in the percentage content of hydroxychavicol in different extracts of *Piper betle* L. by altering the extraction parameters. *Int. J. Adv. Sci. Tech. Res*, 2(4), 517-530.
- Pradhan, D., Suri, K., Pradhan, D., and Biswasroy, P. (2013). Golden Heart of the Nature: *Piper betle* L. *Journal of Pharmacognosy and Phytochemistry*, 1(6).
- Prakash, S., and Van Staden, J. (2007). Micropropagation of *Hoslundia opposita* Vahl—a valuable medicinal plant. *South African Journal of Botany*, 73(1), 60-63.
- Puri, S., and Verma, R. (1996). Vegetative propagation of *Dalbergia sissoo* Roxb. using softwood and hardwood stem cuttings. *Journal of Arid Environments*, 34(2), 235-245.
- Pushparajah, E. (1994). *Leaf analysis and soil testing for plantation tree crops*: ASPAC & FFTC.
- Qaddoury, A., and Amssa, M. (2004). Effect of exogenous indole butyric acid on root formation and peroxidase and indole-3-acetic acid oxidase activities and phenolic contents in date Palm offshoots. *Botanical Bulletin of Academia Sinica*, 45.
- Rahman, S., Rahman, S., Akter, M., Munmun, M., Kalpana, M. A., Jahan, S., ... & Rahmatullah, M. (2012). Scientific validation of folk medicinal uses in Bangladesh of *Piper betle* L. leaves to alleviate pain and lower blood sugar. *Advances in Natural and Applied Sciences*, 6(8), 1496-1503.
- Rai, M. K., Shekhawat, N., Gupta, A. K., Phulwaria, M., Ram, K., and Jaiswal, U. (2011). The role of abscisic acid in plant tissue culture: a review of recent progress. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 106(2), 179-190.
- Raj, S., Srivastava, A., Chowdhury, M., and Johri, J. (2003). Association of *cucumovirus* and potyvirus with betelvine (*Piper betle* L.) as evidenced by ELISA and RT-PCR. *Indian Journal of Experimental Botany*, Vl. 41, pp. 373-276
- Rajasekaran, A., Sivagnanam, G., and Xavier, R. (2008). Nutraceuticals as therapeutic agents: A Review. *Res J Pharm Tech*, 1(4), 328-340.
- Rathee, J. S., Patro, B. S., Mula, S., Gamre, S., and Chattopadhyay, S. (2006). Antioxidant activity of Piper betel leaf extract and its constituents. *Journal of agricultural and food chemistry*, 54(24), 9046-9054.

- Ratnasooriya, W., and Premakumara, G. (1998). Piper betle leaves reversibly inhibits fertility of male rats. *Vidyodaya 1. of Sci, Vol 7*. pp 15- 21
- Razdan, M. (1993). Somatic embryogenesis. *An introduction to plant tissue culture, Oxford & IBH Publi Co Pvt Ltd., New Delhi*, 87-102.
- Richardson, M. D., Karcher, D. E., Hignight, K., & Rush, D. (2008). Drought tolerance and rooting capacity of *Kentucky bluegrass* cultivars. *Crop science*, 48(6), 2429-2436.
- Rintu, D., Shinjini, M., Kaustab, M., Pramathadhip, P., Umesh, P., and Banerjee, E. (2015). Anti-oxidant and anti-inflammatory activities of different varieties of Piper leaf extracts (*Piper betle* L.). *Nutr. Food Sci*, 5(5), 1-15.
- Robert, M. L., Herrera, J., Contreras, F., and Scorer, K. N. (1987). *In vitro* propagation of *Agave fourcroydes* Lem.(Henequen). *Plant Cell, Tissue and Organ Culture*, 8(1), 37-48.
- Rodrigo, R., and Bosco, C. (2006). Oxidative stress and protective effects of polyphenols: comparative studies in human and rodent kidney. *A review. Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 142(3), 317-327.
- Roest, S., & Bokelmann, G. S. (1975). Vegetative propagation of *Chrysanthemum morifolium* Ram. *in vitro*. *Scientia Horticulturae*, 3(4), 317-330.
- Rolland, F., Baena-Gonzalez, E., and Sheen, J. (2006). Sugar sensing and signaling in plants: conserved and novel mechanisms. *Annu. Rev. Plant Biol.*, 57, 675-709.
- Rout, G., Samantaray, S., and Das, P. (2000). *In vitro* manipulation and propagation of medicinal plants. *Biotechnology advances*, 18(2), 91-120.
- Ruby, J. G., Jan, C. H., and Bartel, D. P. (2007). Intronic microRNA precursors that bypass Drosha processing. *Nature*, 448(7149), 83-86.
- Rumary, C., Patel, K. R., and Thorpe, T. A. (1986). Plantlet formation in black and white spruce. II. Histological analysis of adventitious shoot formation *in vitro*. *Canadian journal of botany*, 64(5), 997-1002.
- Saad, A. I., and Elshahed, A. M. (2012). Plant tissue culture media. In Recent advances in plant *in vitro* culture: In Tech.
- Saboonchian, F., Jamei, R., and Sarghein, S. H. (2014). Phenolic and flavonoid content of *Elaeagnus angustifolia* L. (leaf and flower). *Avicenna journal of phytomedicine*, 4(4), 231.
- Sahu, J., and Sahu, R. K. (2013). A review on low cost methods for *in vitro* micropropagation of plant through tissue culture technique. *UK J. Pharm. Biosci*, 1, 38-41.

- Saikia, L., Bhuyan, C., and Dutta, P. (1995). Study on growth, yield and keeping quality of betelvine (*Piper betle* L.) cv. *Local Bangla* as influenced by source and level of nitrogenous fertilizers. *Ind. Cocoa Arec. Spices J*, 19, 46-50.
- Sajc, L., Grubisic, D., and Vunjak-Novakovic, G. (2000). Bioreactors for plant engineering: an outlook for further research. *Biochemical Engineering Journal*, 4(2), 89-99.
- Salamon, I. (2006). Effect of the internal and external factors on yield and qualitative-quantitative characteristics of chamomile essential oil. In *International Symposium on Chamomile Research, Development and Production* 749, (pp. 45-65).
- Salvi, N. D., George, L., and Eapen, S. (2002). Micropropagation and field evaluation of micropropagated plants of turmeric. *Plant Cell, Tissue and Organ Culture*, 68(2), 143-151.
- Samanta, C. (1994). A report on the problems and solutions of betel vine cultivation. *A booklet published by Mr. HR Adhikari: C-2/16, Karunamoyee, Salt Lake City, Kolkata-64 (WB), India*. Document Number)
- Santhakumari, P., Prakasam, A., and Pugalendi, K. (2003). Modulation of oxidative stress parameters by treatment with *Piper betle* leaf in streptozotocin induced diabetic rats. *Indian Journal of pharmacology*, 35(6), 373-378.
- Sarkar, M., Gangopadhyay, P., Basak, B., Chakrabarty, K., Banerji, J., Adhikary, P., et al. (2000). The reversible antifertility effect of *Piper betle* Linn. on Swiss albino male mice. *Contraception*, 62(5), 271-274.
- Schenk, R. U., and Hildebrandt, A. (1972). Medium and techniques for induction and growth of monocotyledonous and dicotyledonous plant cell cultures. *Canadian Journal of Botany*, 50(1), 199-204.
- Scuderi, D., Li Rosi, A., Cassaniti, C., Paratore, A., and Romano, D. (2007). The influence of shading levels on foliage plant growth and quality. In *International Symposium on High Technology for Greenhouse System Management: Greensys2007* 801, 1191-1196.
- Sen, M. K., Nasrin, S., Rahman, S., and Jamal, A. H. M. (2014). *In vitro* callus induction and plantlet regeneration of *Achyranthes aspera* L., a high value medicinal plant. *Asian Pacific journal of tropical biomedicine*, 4(1), 40-46.
- Sharifuddin, H., Fauziah, I., and Zaharah, A. (1990). Technique of soil testing and plant analysis and their utilization for crop production in Malaysia. *Communications in Soil Science & Plant Analysis*, 21(13-16), 1959-1978.
- Sharma, R., and Wakhlu, A. (2001). Adventitious shoot regeneration from petiole explants of *Heracleum candicans* Wall. In *Vitro Cellular & Developmental Biology-Plant*, 37(6), 794-797.

- Sharma, S., Khan, I. A., Ali, I., Ali, F., Kumar, M., Kumar, A., et al. (2009). Evaluation of the antimicrobial, antioxidant, and anti-inflammatory activities of hydroxychavicol for its potential use as an oral care agent. *Antimicrobial agents and chemotherapy*, 53(1), 216-222.
- Sharma, S., Shrivastav, V. K., Shrivastav, A., and Shrivastav, B. (2013). Therapeutic potential of wheat grass (*Triticum aestivum* L.) for the treatment of chronic diseases. *South Asian J Exp Biol*, 3(6), 308-313.
- Shivashankara, K., Mithila, J., and Maiti, S. (2000). Effect of different light intensities on growth and yield of betelvine (*Piper betle* L.). *Journal of Plantation Crops*, 28(3), 196-200.
- Shivashankara, K., Roy, T., and Geetha, G. (2012). Antioxidant capacity, radical scavenging ability, total phenols and flavonoids in three types of betelvine (*Piper betle* L.). *Journal of Spices and Aromatic Crops*, 21(1), 64-67.
- Siddiqui, M. I., and Hussain, S. A. (2007). Effect of indole butyric acid and types of cuttings on root initiation of *Ficus hawaii*. *Sarhad Journal of Agriculture*, 23(4), 919.
- Sidhu, Y. (2011). *In vitro* micropropagation of medicinal plants by tissue culture. *The Plymouth Student Scientist*, 4(1), 432-449.
- Siemann, E., and Rogers, W. E. (2001). Genetic differences in growth of an invasive tree species. *Ecology Letters*, 4(6), 514-518.
- Silber, A., Xu, G., Levkovitch, I., Soriano, S., Bilu, A., and Wallach, R. (2003). High fertigation frequency: the effects on uptake of nutrients, water and plant growth. *Plant and soil*, 253(2), 467-477.
- Simmonds, J. (1983). Direct rooting of micropropagated M26 apple rootstocks. *Scientia horticultrae*, 21(3), 233-241.
- Singh, M., Shakya, S., Soni, V. K., Dangi, A., Kumar, N., and Bhattacharya, S.-M. (2009). The n-hexane and chloroform fractions of *Piper betle* L. trigger different arms of immune responses in BALB/c mice and exhibit antifilarial activity against human lymphatic filarid *Brugia malayi*. *International immunopharmacology*, 9(6), 716-728.
- Sollins, P. (1998). Factors influencing species composition in tropical lowland rain forest: does soil matter? *Ecology*, 79(1), 23-30.
- Somanadhan, B., Varughese, G., Palpu, P., Sreedharan, R., Gudiksen, L., Smitt, U. W., et al. (1999). An ethnopharmacological survey for potential angiotensin converting enzyme inhibitors from Indian medicinal plants. *Journal of ethnopharmacology*, 65(2), 103-112.

- Songsri, P., Jogloy, S., Holbrook, C. C., Kesmala, T., Vorasoot, N., Akkasaeng, C., & Patanothai, A. (2009). Association of root, specific leaf area and SPAD chlorophyll meter reading to water use efficiency of peanut under different available soil water. *Agricultural water management*, 96(5), 790-798.
- Spencer, A., Horvath, N., Gibson, J., Prince, H., Herrmann, R., Bashford, J., et al. (2005). Prospective randomised trial of amifostine cytoprotection in myeloma patients undergoing high-dose melphalan conditioned autologous stem cell transplantation. *Bone marrow transplantation*, 35(10), 971-977.
- Sreekala, G., and Jayachandran, B. (2002). Influence of shade regimes on the physiological parameters of ginger. *Journal of Spices and Aromatic Crops*, 11(1), 30-34.
- Srivastava, L. M. (2002). *Plant growth and development: hormones and environment*: Academic press.
- Srivastava, N., Kamal, B., Sharma, V., Negi, Y. K., Dobriyal, A., Gupta, S., et al. (2010). Standardization of sterilization protocol for micropropagation of *Aconitum heterophyllum*-An endangered medicinal herb. *Academic Arena*, 2(6), 37-42.
- Steffens, B., Wang, J., and Sauter, M. (2006). Interactions between ethylene, gibberellin and abscisic acid regulate emergence and growth rate of adventitious roots in deep water rice. *Planta*, 223(3), 604-612.
- Sugumaran, M., Poornima, M., Venkatraman, S., and Lakshmi, M. (2011). Srinivasansethuvani. Chemical composition and antimicrobial activity of sirugamani variety of *Piper betle* Linn Leaf oil. *Journal of Pharmacy Research*, 4(10), 3424-3426.
- Sulaiman, S. F., Sajak, A. A. B., Ooi, K. L., and Seow, E. M. (2011). Effect of solvents in extracting polyphenols and antioxidants of selected raw vegetables. *Journal of Food Composition and Analysis*, 24(4), 506-515.
- Tan, Y. P., and Chan, E. W. C. (2014). Antioxidant, antityrosinase and antibacterial properties of fresh and processed leaves of *Anacardium occidentale* and *Piper betle*. *Food Bioscience*, 6, 17-23.
- Tchoundjeu, Z., Mpeck, M.-L. N., Asaah, E., and Amougou, A. (2004). The role of vegetative propagation in the domestication of *Pausinystalia johimbe* (K. Schum), a highly threatened medicinal species of West and Central Africa. *Forest Ecology and Management*, 188(1), 175-183.
- Teng, W.-L., Sin, T., and Teng, M.-C. (2002). Explant preparation affects culture initiation and regeneration of *Panax ginseng* and *Panax quinquefolius*. *Plant cell, tissue and organ culture*, 68(3), 233-239.



- Thomas, P. (2004). *In vitro* decline in plant cultures: detection of a legion of covert bacteria as the cause for degeneration of long-term micropropagated triploid watermelon cultures. *Plant cell, tissue and organ culture*, 77(2), 173-179.
- Thomas, T. D., Bhatnagar, A., Razdan, M., and Bhojwani, S. (1999). A reproducible protocol for the production of gynogenic haploids of mulberry, *Morus alba* L. *Euphytica*, 110(3), 169-173.
- Torres, K. C. (1989). Application of Tissue Culture Techniques to Horticultural Crops. *In Tissue Culture Techniques for Horticultural Crops* (pp. 66-69): Springer.
- Tripathi, L., and Tripathi, J. N. (2003). Role of biotechnology in medicinal plants. *Tropical Journal of Pharmaceutical Research*, 2(2), 243-253.
- Usmanghani, K., Saeed, A., and Alam, M. T. (1997). Indusynic medicine. *Department of Pharmacognosy, Faculty of Pharmacy, University of Karachi, Pakistan*, 363-364.
- Valladares, F., Wright, S. J., Lasso, E., Kitajima, K., and Pearcy, R. W. (2000). Plastic phenotypic response to light of 16 congeneric shrubs from a Panamanian rainforest. *Ecology*, 81(7), 1925-1936.
- Vendrame, W., Moore, K. K., and Broschat, T. K. (2004). Interaction of light intensity and controlled release fertilization rate on growth and flowering of two new guinea impatiens cultivars. *HortTechnology*, 14(4), 491-495.
- Verdeil, J.-L., Alemanno, L., Niemenak, N., and Tranbarger, T. J. (2007). Pluripotent versus totipotent plant stem cells: dependence versus autonomy? *Trends in plant science*, 12(6), 245-252.
- Verma, A., Kumar, N., and Ranade, S. (2004). Genetic diversity amongst landraces of a dioecious vegetatively propagated plant, betelvine (*Piper betle* L.). *Journal of Biosciences*, 29(3), 319-328.
- Vieitez, A., Corredoira, E., Ballester, A., Muñoz, F., Durán, J., and Ibarra, M. (2009). *In vitro* regeneration of the important North American oak species *Quercus alba*, *Quercus bicolor* and *Quercus rubra*. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 98(2), 135-145.
- Vinterhalter, D. V. (1989). *In vitro* propagation of green-foliaged *Dracaena fragrans* Ker. *Plant Cell, Tissue and Organ Culture*, 17(1), 13-19.
- Vossen, H., and Wessel, M. (2000). Plant Resources of South-East Asia Stimulants: Backhuys Publisher, Netherlands. 102-106
- Vyas, S. (2004). Influence of light intensity on growth, yield and laws one content of Henna (*Lawsonia inermis* L.). *Indian journal of plant physiology*, 9(3), 275-277.

- Wala, B. B., and Jasrai, Y. T. (2003). Micropropagation of an endangered medicinal plant: *Curculigo orchoides* Gaertn. *Plant Tissue Cult*, 13(1), 13-19.
- Wang, P.-j., and Hu, C.-y. (1984). *In vitro* cloning of the deciduous timber tree *Sassafras randaiense*. *Zeitschrift für Pflanzenphysiologie*, 113(4), 331-335.
- Warren, J. M., Bassman, J. H., Fellman, J. K., Mattinson, D. S., and Eigenbrode, S. (2003). Ultraviolet-B radiation alters phenolic salicylate and flavonoid composition of *Populus trichocarpa* leaves. *Tree Physiology*, 23(8), 527-535.
- Wira, A., Razi, I. M., and Jamil, Z. A. (2011). Composts as additives in coconut coir dust culture for growing rockmelon (*Cucumis melo* L.). *Journal of Tropical Agriculture and Food Science*, 39(2), 229-237.
- Wirotasangthong, M., Inagaki, N., Tanaka, H., Thanakijcharoenpath, W., and Nagai, H. (2008). Inhibitory effects of *Piper betle* on production of allergic mediators by bone marrow-derived mast cells and lung epithelial cells. *International immunopharmacology*, 8(3), 453-457.
- Wolf, B. (1982). A comprehensive system of leaf analyses and its use for diagnosing crop nutrient status. *Communications in Soil Science & Plant Analysis*, 13(12), 1035-1059.
- Xie, B., and Wang, H. (2006). Effects of light spectrum and photoperiod on contents of flavonoid and terpene in leaves of *Ginkgo biloba* L. *Journal of Nanjing Forestry University*, 30, 51-54.
- Xu, G., Levkovitch, I., Soriano, S., Wallach, R., and Silber, A. (2004). Integrated effect of irrigation frequency and phosphorus level on lettuce: P uptake, root growth and yield. *Plant and soil*, 263(1), 297-309.
- Yahya, A., Nor'ain, A. R., and Ab Kahar, S. (2009). Adventitious rooting in microcuttings of selected indigenous landscape trees of Malaysia. *International Journal of Agriculture and Biology*, 11(4), 419-424.
- Yaseen, M., Ahmad, T., Sablok, G., Standardi, A., and Hafiz, I. A. (2013). Role of carbon sources for *in vitro* plant growth and development. *Molecular biology reports*, 40(4), 2837-2849.
- Yuan, L.-Y., Li, W., Liu, G., and Deng, G. (2012). Effects of different shaded conditions and water depths on the growth and reproductive strategy of *Vallisneria spinulosa*. *Pak. J. Bot*, 44(3), 911-918.
- Zakrzewski, P. A. (2002). Bioprospecting or biopiracy? The pharmaceutical industry's use of indigenous medicinal plants as a source of potential drug candidates. *University of Toronto Medical Journal*, 79(3), 252-254.



- Zhao, H., Dong, J., Lu, J., Chen, J., Li, Y., Shan, L., et al. (2006). Effects of extraction solvent mixtures on antioxidant activity evaluation and their extraction capacity and selectivity for free phenolic compounds in barley (*Hordeum vulgare* L.). *Journal of Agricultural and Food Chemistry*, 54(19), 7277-7286.
- Zhou, L. G., and Wu, J. Y. (2006). Development and application of medicinal plant tissue cultures for production of drugs and herbal medicinals in China. *Natural product reports*, 23(5), 789-810.
- Zimmerman, R. H. (1984). Rooting apple cultivars in vitro: Interactions among light, temperature, phloroglucinol and auxin. *Plant cell, tissue and organ culture*, 3(4), 301-311.
- Zlesak, D. C. (2012). Physocarpus plant named 'Donna May': *U.S. Patent No.* pp. 22,634
- Zobel, A., and Clarke, P. (1999). Production of phenolic compounds in seedlings of *Acer saccharum* and *Acer platanoides* in response to UV-A radiation and heavy metals. *Allelopathy J*, 6, 21-34.