



UNIVERSITI PUTRA MALAYSIA

***OIL-BASED NANOEMULSION OF *Metarhizium anisopliae* (Metschn)
SOROKIN TO CONTROL RED PALM WEEVIL
Rhynchophorus ferrugineus (Olivier)***

ALI ZACHI ABDULQADER

FP 2018 81



**OIL-BASED NANOEMULSION OF *Metarhizium anisopliae* (Metschn)
SOROKIN TO CONTROL RED PALM WEEVIL
Rhynchophorus ferrugineus (Olivier)**

By

ALI ZACHI ABDULQADER

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

January 2018

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**OIL-BASED NANOEMULSION OF *Metarhizium anisopliae* (Metschn)
SOROKIN TO CONTROL RED PALM WEEVIL
Rhynchophorus ferrugineus (Olivier)**

By

ALI ZACHI ABDULQADER

January 2018

Chairman : Professor Dzulkhifli Omar, PhD
Faculty : Agriculture

Oil based emulsion formulations of *Metarhizium anisopliae* were prepared, characterised and evaluated for their effectiveness against the larvae and adults of *Rhynchophorus ferrugineus*. Infested *R. ferrugineus* adults were collected from Terengganu, and *M. anisopliae* was isolated by cadavers of red palm weevil adults. Four strains were obtained and identified via morphology and molecular technique as *M. anisopliae*. The virulence of these strains was evaluated against the adult and larvae of *R. ferrugineus* by time exposure mortality bioassay. The strain coded as D1 gave the lowest LT₅₀ values of 7.2 and 5.2 days at the conidia concentration of 10⁶ and 10⁷ spores/mL, respectively against the larvae. While against the adult, the D1 strain also gave the lowest LT₅₀ values of 6 and 5 days at same conidia concentrations. Oil emulsion formulations of the most virulence isolate conidia of *M. anisopliae* were prepared through ternary phase diagram consisting of 20% (w/w) surfactant, 40% (w/w) oil and 40% (w/w) water containing 10⁷ spore/mL. The surfactants and oil were first evaluated for their compatibility with conidia by using direct plating. The effect of the surfactants on conidia germination was evaluated by counting the germination rate of the conidia using a microscope. Agnique PG9116 at 1% concentration gave 87.5 % germination while at 5% surfactant Emereen1604 and EW70 gave 70% conidia germination. In a study of the effect of oils on the conidia germination, glycerin oil gave highest conidia germination rate. Sunflower and glycerin showed less inhibition at 1% concentration with 49.13 and 44.13% growth rate respectively. Palm oil at 5% concentration was the best with 56.88% growth rate. At 10% concentration of oils, soybean and glycerin gave 48.63 and 45.38% growth rate respectively. Eight ternary phase diagram systems were then constructed. The selected systems showed large isotropic regions. They were Agnique PG9116/ glycerin/ water, Emereen1604/ glycerin/ water, Tensiofix 96 DB08/ glycerin/ water, Tensiofix 96 DB10/ glycerin/ water, Tensiofix EW 70/ glycerin/ water, Termul 1284/ glycerin/ water, Tween20/ glycerin/ water and Tween80/ glycerin/ water. Eight oil emulsion formulations were

derived and characterised. All the formulations were stable under centrifuge, storage at 26 ± 1 °C, 60 ± 5 % RH and under high temperature (54 ± 1 °C) for two weeks. On the particle size, seven formulations were in the range below <100 nm sizes indicating that the formulations were in the category of the nanoemulsion. The zeta potential of the formulations ranged between -7.22 to -39.06 mV, the pH ranged from 4 to 6.34, the surface tension ranged from 32.03 to 41.83 mN/m, and the viscosity ranged from 2.40 to 28.8 mPas. In the study on the toxicity of the oil nanoemulsion formulations of *M. anisopliae* conidia against the larvae, the formulation coded as E1604 gave the LT_{50} of 4.90 days while the conidia water suspension gave LT_{50} of 6 days. On adults, the LT_{50} was 2.20 days while the conidia water suspension was 5 days. Effect of oil nanoformulations on the conidia germination on the cuticle of *R. ferrugineus* was also observed, and after 20 hrs., the E1604 showed 55% germination compared to conidia water suspension of 49.8%. The formulation E1604 showed the longest germ tube of 41.34 μ m and full penetration while the conidia water suspension gave 5.28 μ m length of a germ tube. The E1604 recorded 100% cumulative mortality after 6 and 4 days on larvae and adults respectively. The oil nanoemulsion of *M. anisopliae* conidia shows good potential for the sustainable control of both adults and larvae of *R. ferrugineus*.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**NANOEMULSI BERASASKAN MINYAK *Metarhizium anisopliae* (Metschn)
SOROKIN UNTUK MENGAWAL KUMBANG PALMA MERAH
Rhynchophorus ferrugineus (Olivier)**

Oleh

ALI ZACHI ABDULQADER

Januari 2018

Pengerusi : Profesor Dzolkhifli Omar, PhD
Fakulti : Pertanian

Formulasi emulsi berasaskan minyak *Metarhizium anisopliae* disediakan, dicirikan dan dinilai untuk keberkesanannya terhadap larva dan peringkat dewasa *Rhynchophorus ferrugineus*. Peringkat dewasa *R. ferrugineus* dikumpulkan dari Terengganu dan *M. anisopliae* telah diasingkan oleh peringkat dewasa *R. ferrugineus* yang telah mati. Empat strain diperoleh dan dikenal pasti melalui morfologi dan teknik molekul sebagai *M. anisopliae*. Keberkesanan strain ini dinilai terhadap peringkat dewasa dan larva *R. ferrugineus* oleh bioessei mortaliti pendedahan terhadap masa. Strain yang dikodkan sebagai D1 memberikan nilai LT50 terendah sebanyak 7.2 dan 5.2 hari pada kepekatan konidia 106 dan 107 spora / mL masing-masing terhadap larva. Manakala terhadap peringkat dewasa, D1 juga memberikan nilai LT50 paling rendah sebanyak 6 dan 5 hari kepekatan conidia yang sama. Formulasi emulsi minyak yang paling berkesan ialah daripada conidia *M. anisopliae* telah disediakan dengan pembinaan diagram fasa ternari dengan surfaktan 20% (w / w), minyak 40% (w / w) dan air 40% (w / w) yang mengandungi spora 107 / mL. Surfaktan dan minyak mula-mula ditentukan untuk kesesuaian dengan conidia dengan menggunakan secara langsung. Kesan surfaktan pada percambahan konidia ditentukan dengan menggunakan mikroskop untuk menghitung kadar percambahan conidia. Agnique PG9116 pada kepekatan 1% memberikan percambahan 87.5% manakala pada 5% surfactant Emeere 1604 dan EW70 memberikan percambahan conidia 70%. Dalam kajian mengenai kesan minyak pada percambahan konidia, minyak gliserin menunjukkan kadar percambahan konidia tertinggi. Lapan sistem diagram fasa ternari kemudian dibina dan sistem yang dipilih menunjukkan kawasan isotropik yang besar. Ia adalah Agnique PG9116 / gliserin / air, Emereen 1604 / gliserin / air, Tensiofix 96 DB08 / gliserin / air, Tensiofix 96 DB10 / gliserin / air, Tensiofix EW 70 / gliserin / air dan Tween80 / gliserin / air. Lapan formulasi emulsi minyak diperoleh dan dicirikan. Semua formulasi adalah stabil di bawah emparan, penyimpanan pada 26 ±

1 ° C, 60 ± 5% RH dan di bawah suhu tinggi (54 ± 1 ° C) selama dua minggu. Pada saiz zarah, tujuh formulasi adalah dalam julat di bawah saiz <100 nm yang menunjukkan bahawa formulasi berada dalam kategori nanoemulsi. Potensi zeta dari formulasi antara - 7.22 hingga -39.06 mV, pH antara 4 hingga 6.34, ketegangan permukaan antara 32.03 hingga 41.83 mN / m dan kelikatannya antara 2.40 hingga 28.8 mPas. Dalam kajian mengenai ketoksikan formula nanoemulsion minyak *M. anisopliae* conidia terhadap larva, rumusan itu dikodkan sebagai E1604 memberikan LT50 sebanyak 4.90 hari manakala suspensi air conidia memberikan LT50 dari 6 hari. Pada peringkat dewasa, LT50 adalah 2.20 hari manakala suspensi air conidia adalah 5 hari. Kesan formula nano minyak pada percambahan konidia pada kutikel *R. ferrugineus* juga diperhatikan dan selepas 20 jam, E1604 menunjukkan percambahan 55% berbanding dengan penggantungan air konidia sebanyak 49.8%. Perumusan E1604 menunjukkan tiub germanium paling panjang 41.34 µm dan penembusan penuh manakala penggantungan air konidia memberikan panjang 5.28 µm. E1604 mencatatkan kematian kumulatif 100% selepas 6 dan 4 hari pada larva dan dewasa. Emulsi nano minyak *M. anisopliae* conidia menunjukkan potensi yang baik untuk kawalan mampan kedua-dua peringkat dewasa dan larva *R. ferrugineus*.

ACKNOWLEDGEMENTS

بسم الله الرحمن الرحيم

First and foremost, I thank Allah s.w.t. for giving me His blessing and strength throughout this research work. Special thanks to Prof DrDzolkhifli Omar for his guidance, encouragement and assistance throughout myPhD research work. Special thanks also dedicated to Prof Dr Rita Muhamed and DrNorida Binti Mazlan as of my co-supervisor.

Special thanks also go to my father and my mother, thank you for being a loving parent who has always been praying for me during your lifetime. Most profound gratitude even to my wife Yusra Abadulkhilq, thank you for being supportive, beautiful memories in our life, your advice, and your Doa' for my success. Also, special gratitude to my son Sajjad and my daughter Fatima and my son Hassan.

Nor forgotten, thanks to all my colleagues and friends, staffs and technicians at plant protection Department, Faculty of Agriculture, Universiti Putra Malaysia.

-Alhamdulillah-

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Dzolkhifli B Omar, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Rita Muhamad Awang, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Norida Binti Mazlan, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____

Date: _____

Name and Matric No.: Ali Zachy Abdulqader, GS38673

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Professor Dr. Dzolkhifli B Omar

Signature: _____
Name of Member
of Supervisory
Committee: Professor Dr. Rita Muhamad Awang

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Norida Binti Mazlan

TABLE OF CONTENTS

ABSTRACT	Page
ABSTRAK	i
ACKNOWLEDGEMENTS	iii
APPROVAL	v
DECLARATION	vi
LIST OF TABLES	viii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xv
	xviii

CHAPTER

1	INTRODUCTION	1
2	LITERATURE REVIEW	4
2.1	Pesticide Formulation	4
2.2	Formulation of Entomopathogenic Fungi	5
2.2.1	Granular Formulations (G)	5
2.2.2	Sprayable Formulation	6
2.2.3	Aqueous Formulations	6
2.2.4	Oil-Based Formulations	6
2.3	Component of the formulation	7
2.3.1	Carrier	7
2.4	Surfactants	8
2.5	Types of surfactants	9
2.5.1	Anionic surfactants	9
2.5.2	Non-ionic surfactants	9
2.5.3	Cationic surfactants	9
2.5.4	Amphoteric or Zwitterionic surfactants	9
2.6	Formulating and characterization of emulsion formulation	10
2.6.1	Ternary phase diagram	10
2.6.2	Surface tension of emulsion	11
2.7	<i>Metarhizium</i> sp.	11
2.8	Pathogenicity of <i>Metarhizium</i>	12
2.9	Red palm weevil (RPW) <i>Rhynchophorus ferrugineus</i> (Olivier)	14
2.9.1	Life history of <i>R. ferrugineus</i>	16
2.9.2	Damage and Symptoms of <i>R. ferrugineus</i>	17
2.9.3	Control of <i>R. ferrugineus</i>	18

3	IDENTIFICATION AND CHARACTERIZATION OF <i>Metarhizium anisopliae</i>	19
3.1	Introduction	19
3.2	Materials and Methods	19
3.2.1	Collection and Isolation of the fungus	19
3.2.2	Morphological observations	20
3.2.3	Subculture of <i>Metarhizium</i> spp.	20
3.2.4	Molecular characterization of <i>Metarhizium</i> spp.	20
3.2.4.1	DNA Extraction of <i>Metarhizium</i> spp.	20
3.2.4.2	ITS rDNA amplification and sequencing	21
3.2.5	Virulence of <i>M. anisopliae</i> conidia against <i>R. ferrugineus</i>	21
3.2.6	Statistical analysis	22
3.3	Results	22
3.3.1	Isolation Fungus of the study	22
3.3.2	Morphology Observation	22
3.3.3	Molecular identification of the study fungus	23
3.3.4	Virulence of <i>M. anisopliae</i> against <i>R. ferrugineus</i>	25
3.4	Discussion	26
3.5	Conclusion	28
4	FORMULATION OF OIL BASED EMULSION FORMULATION	29
4.1	Introduction	29
4.2	Materials and methods	29
4.2.1	Ingredients	29
4.2.2	Effect of formulation ingredients on the conidia viability	31
4.2.2.1	Effect of surfactants on the conidial viability of <i>M. anisopliae</i>	31
4.2.2.2	Effect of oils on the conidial viability of <i>M. anisopliae</i>	31
4.2.3	Preparation of oil nanoemulsion formulation	31
4.2.3.1	Miscibility test pre-formulation	31
4.2.3.2	Construction of ternary phase diagram	33
4.2.3.3	Selection of formulation composition	34
4.2.4	Characterization of oil nanoemulsion formulation	34
4.2.4.1	Stability of formulations under centrifugation	35
4.2.4.2	Stability of oil emulsion formulations under storage conditions	35
4.2.4.3	Particle size and zeta potential measurements of emulsion	35
4.2.4.4	The pH measurement of emulsions	36
4.2.4.5	Surface tension measurement of emulsions	36
4.2.4.6	Viscosity measurement of emulsions	37
4.2.5	Statistical analysis	38
4.3	Results	38

4.3.1	Effect of surfactants on the conidial viability of <i>M. anisopliae</i>	38
4.3.2	Effects of oils on conidia viability	39
4.3.3	Miscibility Test of the inert ingredients	40
4.3.4	Ternary phase diagram of nanoemulsion system study	43
4.3.5	Points selection	48
4.3.6	Stability of nanoemulsion formulations under centrifuge	49
4.3.7	Thermostability test of nanoemulsion formulations	50
4.3.8	Particle size and Zeta potential	51
4.3.9	The pH measurement of emulsions	52
4.3.10	Surface tension measurement of emulsions	52
4.3.11	Viscosity measurements of emulsions	52
4.4	Discussion	53
4.5	Conclusion	55
5	EFFICACY OF FORMULATED <i>Metarhizium anisopliae</i> AGAINST <i>Rhynchophorus ferrugineus</i>	56
5.1	Introduction	56
5.2	Materials and Methods	57
5.2.1	Effect of the surfactants on the larvae of <i>Rhynchophorus ferrugineus</i>	57
5.2.2	Effect of the surfactants on the adults of <i>R. ferrugineus</i>	57
5.2.3	Effect the oils on <i>R. ferrugineus</i> larvae	57
5.2.4	Effect of the oils on the <i>R. ferrugineus</i> adults	57
5.2.5	Toxicity of the oil nanoemulsion formulations against <i>R. ferrugineus</i>	57
5.2.6	Infection process of <i>M. anisopliae</i> to the larvae and adults of <i>R. ferrugineus</i>	58
5.2.7	Effect of formulations on the infection of <i>M. anisopliae</i> on <i>R. ferrugineus</i> .	58
5.2.8	Statistical analysis	59
5.3	Results	59
5.3.1	Effect of surfactants and oils on <i>R. ferrugineus</i> larvae and adults	59
5.3.2	Toxicity of the oil nanoemulsion formulations against <i>R. ferrugineus</i>	60
5.3.3	Toxicity of the best four formulations on the infection of <i>M. anisopliae</i> on <i>R. ferrugineus</i> .	72
5.4	Discussion	74
5.5	Conclusion	76
6	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH	77
6.1	Recommendations for Future Research	79

REFERENCES	80
APPENDIX	96
BIODATA OF STUDENT	98



LIST OF TABLES

Table		Page
2.1	Types of agrochemicals formulation according to application methods	4
2.2	Surface tensions of common liquids	11
2.3	Scientific names, common names and host of RPW in the world	16
3.1	Blast results of four isolates of <i>Metarhizium</i> isolate based on ITS region	24
3.2	Lethal time analysis of <i>M. anisopliae</i> at 10^6 conidia/ml against <i>R. ferrugineus</i> larvae	25
3.3	Lethal time analysis of <i>M. anisopliae</i> at 10^7 conidia/ml against <i>R. ferrugineus</i> larvae	26
3.4	Lethal time analysis of <i>M. anisopliae</i> at 10^6 conidia/ml against <i>R. ferrugineus</i> adults	26
3.5	Lethal time analysis of <i>M. anisopliae</i> at 10^7 conidia/ml against <i>R. ferrugineus</i> adults	26
4.1	Ingredients used in the ternary phase diagram study	30
4.2	Surfactants, oils and Water grouped in different combinations for phase diagram construction	32
4.3	Conidia germination in different surfactants concentrations	39
4.4	Viability of <i>M. anisopliae</i> conidia in different oils	39
4.5	Miscibility Test between oils and surfactants used based on spontaneous emulsification	42
4.6	Composition of nanoemulsion formulations from the homogeneous phase	44
4.7	Stability test of nanoemulsion formulation at centrifugation and temperature storage 26°C and 54°C .	50
4.8	mean particle size and zeta potential of the formulations	51
4.9	Surface tension values of nanoemulsion	52
5.1	Effect of surfactants and oils on <i>R. ferrugineus</i> larvae and adults after exposure to different concentrations	59
5.2	Mortality time analysis of <i>R. ferrugineus</i> larvae after exposure to eight different nanoemulsion conidial formulations 10^7 conidia/mL.	60
5.3	Mortality time analysis of <i>R. ferrugineus</i> adults after exposure to eight different nanoemulsion conidial formulations 10^7 conidia/mL.	61
5.4	Germ tube of <i>M. anisopliae</i> characterization after 20 hrs. of treatment with the nanoemulsion formulations	61

LIST OF FIGURES

Figure		Page
2.1	General structure of glycerine	8
2.2	Ternary phase diagram system	10
2.3	Life cycle of <i>M. anisopliae</i> and normal infection process	13
2.4	Insect cuticle structure	13
2.5	Life cycle of <i>R. ferrugineus</i>	17
3.1	Infected adults of <i>R. ferrugineus</i> collected from field plantations in Kampung Baru Kuala Abang Dungun Terengganu	20
3.2	The morphological appearance of the four isolates after subculture and they were coded as A1, B1, C1 and D1	23
3.3	Gel electrophoresis is showing bands of PCR products from ITS region and the amplification fragments were approximately 550 bp. M-DNA ladder	24
3.4	Phylogenetic inferred the relationship by Maximum Likelihood of <i>Metarhizium anisopliae</i> isolates compared with accession numbers of other species based on rDNA ITS sequences. Numbers below the branches represent the percentage for each branch in 1000 bootstrap replications.	25
4.1	Ternary phase diagram reading. Point 1 stands for a mixture of 30% oil phase at a fixed ratio (w/w), 30% aqueous phase and 40% surfactant.	34
4.2	(A) The zeta sizer Nano-ZS, Malvern, UK;(B) the capillary zeta potential cell for the Malvern Zetasizer.	35
4.3	Tension meter Model KRUSS Tensiometer K6, UK	37
4.4	Viscometer model RheolabQC	38
4.5	Photographs showed miscibility test after vortex mixture (oil, surfactant and water) a) Sesame oil mixed with 8 surfactants and water W/W; b) Soybean oil mixed with 8 surfactants and water; c) Sunflower oil mixed with 8 surfactants and water w/w; d) Canola oil mixed with 8 surfactants and water W/W; e) Corn oil mixed with 8 surfactants and water W/W; f) Palm oil mixed with 8 surfactants and water W/W; g) Glycerin oil mixed with 8 surfactants and water W/W.	41
4.6	Phase diagram of Agnique PG9116/ Glycerin/ Water system present one phase region 83%	44
4.7	Phase diagram of Emeree1604/ Glycerin/ Water system show 79.5% isotropic region	45
4.8	Phase diagram of Tensiofix DB08/ Glycerin/ Water system presented	45
4.9	Phase diagram of Tensiofix DB08/ Glycerin/ Water system 25% isotropic region, 5% 2ph and 10% nano gel.	46

4.10	Phase diagram of Tensiofix EW70/ Glycerin/ Water system 73% one phase.	46
4.11	Phase diagram of Termul 1284/ Glycerin/ Water system 62% one phase.	47
4.12	Phase diagram of Tween 20/ Glycerin/ Water system showed 80% one phase region.	47
4.13	Phase diagram of Tween 80/ Glycerin/ Water system showed 80% one phase region.	48
4.13	Photographs showed nano emulsion formulations after selected points with conidia and water suspension at 26°C.	49
4.15	Photographs showed stable nanoemulsion formulations after stability test.	50
4.16	Photographs showed nanoemulsion formulations present one phase and transparent at 54°C with droplet diameters of less than 100 nm	51
5.1	SEM micrograph of the adult treatment by the formulation of B08; a) conidia of <i>M. anisopliae</i> not germinated; b) conidia adhered to the seta after 20 hrs.	62
5.2	SEM micrograph of the treatment adult by the formulation of B70; a) conidia of <i>M. anisopliae</i> not germinated after 20 hrs.	62
5.5	SEM micrograph of the treatment adult by the formulation of DB10; a) conidia of <i>M. anisopliae</i> not germinated after 20 hrs.	63
5.4	SEM micrograph of the development of the germ tube at the adult cuticle by the formulation of T20 (a) conidia of <i>M. anisopliae</i> adhered to the body of <i>R. ferrugineus</i> (b) germ tube (c) germ tube penetrate the cuticle of the adult of <i>R. ferrugineus</i> after 20 hrs.	63
5.5	SEM micrograph of the treatment adult by the formulation of Term 1284(a) conidia of <i>M. anisopliae</i> (b) germ tube (c) germ tube penetrate the cuticle of the adult of <i>R. ferrugineus</i> after 20 hrs.	64
5.6	SEM micrograph of the treatment adult by the formulation of T80; a) germinated conidium of <i>M. anisopliae</i> ; b) germ tube; c) germ tube penetrate the cuticle of the adult of <i>R. ferrugineus</i> after 20 hrs.	64
5.7	SEM micrograph of the treatment adult by the formulation of 1604; a) germinated conidia; b) germ tube; c) germination of a germ tube into the cuticle of <i>R. ferrugineus</i> growth of germ tube after 20 hrs.	65
5.8	SEM micrograph of the treatment adult by the formulation Ag 9116 (a) the germinated conidia (b) growth of the germ tube after 20 hrs.	65
5.9	SEM micrograph of control treatment adult (a) germinated conidia of <i>M. anisopliae</i> (b) germ tube penetrate the adult cuticle of <i>R. ferrugineus</i> after 20 hrs.	66

5.10	SEM micrograph of the larval treatment by the formulation of B08; a) conidia of <i>M. anisopliae</i> not germinated.	66
5.11	SEM micrograph of the larval treatment by the formulation of BW70; a) conidia of <i>M. anisopliae</i> not germinated after 20hrs.	67
5.12	SEM micrograph of the larval treatment by the formulation of DB10 (a) conidia of <i>M. anisopliae</i> not germinated.	67
5.13	SEM micrograph of the larva treated by the formulation T20; a) germinated conidia (b) fully formation of the germ tube after 20 hrs.	68
5.14	SEM micrograph of the larva treated by the formulation 1604; a) adhered germinated conidia; b) germ tube with fully formation; c) germ tube penetration the cuticle after 20 h.	68
5.15	SEM micrographs of the larva treated by the formulation T80; a) the germinated conidia; b) growth of the germ tube after 20 h.	69
5.16	SEM micrograph of the larva treated by the formulation Ag 9116 (a) the germinated conidia (b) growth of the germ tube after 20 hrs.	69
5.17	SEM micrograph of the larva treated by the formulation Term 1284; a) the germinated conidia adhered to the cuticle; b) growth of the germ tube penetration the cuticle of the insect after 20 hrs.	70
5.18	SEM micrograph of the larva treated by water suspension (control treatment) (a) the germinated conidia (b) growth of the germ tube after 20 hrs.	70
5.19	SEM micrograph a thickening of the edge of the germ-tube after formation on the cuticle after 20 hrs.	71
5.2	SEM micrograph; a) germ tube with fully formation adhered germinated conidia; b) germ tube penetration the cuticle of the rostrum parts after 20 hr; c) conidium adhered to the rostrum of the <i>R. ferrugineus</i> adult after 20 hrs.	71
5.21	(a) effect of nano formulations against the larva (b) effect of nano formulations against the adult of <i>R. ferrugineus</i>	72
5.22	Toxicity of the best four formulations on <i>R. ferrugineus</i> larvae (a) healthy larvae before treatment with <i>M. anisopliae</i> oil nanoformulation (b) changing the colour to dark (c) appear the mycelium out of the body (c) covering the larva body with <i>M. anisopliae</i> conidia.	73
5.23	Toxicity of the best four formulations on <i>R. ferrugineus</i> adult after applied the oil nanoformulation of <i>M. anisopliae</i> conidia (a) healthy adult of <i>R. ferrugineus</i> before the treatment (b) infected adult of <i>R. ferrugineus</i> with of <i>M. anisopliae</i> after the treatment.	74

LIST OF ABBREVIATIONS

ITS	Internal transcribed spacer
L	liter
G	gram (s)
Mg	milligram(s)
mL	milliliter (s)
cm	centimeter(s)
mm	millimeter (s)
h	hour(s)
%	percent
°	degree
°C	degree(s) Celsius
CRD	Complete Randomized Design
DAT	Day after treatment
S.E	Standard Error
Sec.	second
UPM	Universiti Putra Malaysia
a.i	active ingredient
w/w	weight over weight
rpm	revolutions per minute
no.	number
&	and
<i>i.e</i>	<i>exempli gratia</i> , for example
<i>et al.</i> ,	<i>et alii</i> , and others
RPW	Red Palm Weevil
Dtx	destruxin
♀	male
♂	female
µL	microliter
DNA	Deoxyribonucleic acid
PCR	Polymerase chain reaction
v/v	volume to volume
LT ₅₀	median lethal time
mN/m	milli newton per meter
Mv	millivolt
SEM	scanning electron microscope
UV	ultraviolet
EPF	entomopathogenic fungi
ULV	ultra low volume

CHAPTER 1

INTRODUCTION

The red palm weevil *R. ferrugineus* (Coleoptera: Rhynchophoridae) is a phytophagous insect which feeds explicitly on palm trees. It is a global concern due to its harmful feeding habits within palm trees and further threats to the ornamental and other date palm species. RPW is a severe pest of coconut palm *Cocos nucifera* and was reported from 15% of the coconut growing countries worldwide (Faleiro, 2006). The RPW rapidly spread to date palm-growing countries during the past two decades. In Malaysia; studies show red palm weevil as a pest of palm trees especially the economically important coconut, *C. nucifera*, and the sago, *Metroxylum sagu* (Idris *et al.*, 2014). Plantation of palm oil plays a significant role in the economy of the country as country's GDP relies on it with a maximum percentage. Therefore, red palm weevil is a high threat to its direct effects on people's livelihood in many countries such as Malaysia.

An imperative role of chemical pesticides has been recorded in crop protection programs to serve humankind significantly. An estimated quantity of about 2.5 million tons of pesticides has been used each year at the cost of \$20 billion worldwide. However, the unselective use of these pesticides has resulted in resistance, resurgence and outbreaks of new insect pest species. Furthermore, ground-breaking public exposure of the risk to the environment including benefits organism and public health posed by the frequent use of these chemical threatening the sustainability of ecosystems (Chagnon *et al.*, 2015; Aktaret *et al.*, 2009). There has been a continuous on going endeavour to reduce harmful effects of these pesticides, either by the development of more targeted compounds that exhibit less side effects, abandon of highly hazardous chemicals or by the development of non-chemical methods of pest management (Gilland Garg 2014).

The functional utilisation of biological agents as pesticides have efficiently delivered the satisfactory results in reducing the incidences of insect pests, weeds and diseases for many years. There is a number of entomopathogenic agents that have been documented (Gindin *et al.*, 2006). *M. anisopliae* is one of the worthiest examples in this regard that has been considered as an essential biological agent in reducing the pest population of different insect orders (Zimmermann, 2007).

The fungus was initially isolated from Coleopteran insects (*Anisopliae austriaca*) in 1878 by Leland (2001). It possesses a high rate of germination and responsible for the production of various enzymes that may result in insect toxins (Schrack and Vainstein, 2010) but not infectious or toxic to mammals. Although, entomopathogenic plays a vital part in pest management, but they are still facing many challenges with regard to improving the efficiency of these insect-microbial pesticides formulation, shelf life and compatible with the environment. Therefore, improving the conidial

formulation of *M. anisopliae* is an indispensable step to optimise its biological control strategy.

It has been well documented that the oil in water emulsion enhances the efficacy of entomopathogens against various insects in controlled and uncontrolled conditions (Polar *et al.*, 2005). The oil formulation of the Citowee oil enhanced the fungal virulence on *Eurygaster integriceps* (Sedighi *et al.*, 2013). Also, the oil formulations are useful in strengthening the transfer of conidia to the areas of thinner cuticle (Ibrahim *et al.*, 1999). Similarly, these bio fungicides ensure the greater ability of adherence to the host body and the combination of oil and conidia further assists in protecting the fungus against fast dehydration in low-humidity environments (Bateman *et al.*, 1993), high temperature (McClatchie *et al.*, 1994) and ultraviolet radiation (Moore *et al.*, 1996; Alves *et al.*, 1998; Bateman and Alves, 2002). The preference thus has been given to the use of oil based formulations of fungi that have been shown to be effective in controlling various arthropods in laboratory and field conditions (Bateman *et al.*, 1993; Batta, 2003).

In a recent decade, red palm weevil (RPW) *R. ferrugineus* has become a noxious pest of coconut and date palm trees in most Asian countries. An introduction of *M. anisopliae* has displayed a great ability to control this pest and was found to be the main pathogen against it through using proper formulation and application method (Francardi *et al.*, 2016). It is also suggested that the fungi could spread infecting healthy insects of *R. ferrugineus* by horizontal transmission as the insect is highly promiscuous and live in aggregation (Francardi *et al.*, 2013 and Ll  cer *et al.*, 2013). The oil-based formulation of *Beauveria bassiana* has reported the little effect on RPW (Abdel-Samad *et al.*, 2011) in comparison with *M. anisopliae* that shows the highest efficacy against RPW larvae and adults (Francardi *et al.*, 2012 and 2013). The oil-based formulations of *M. anisopliae* were more effective than aqueous suspensions against eggs, larvae and engorged females of *Rhipicephalus microplus* (Camargo *et al.*, 2012).

The mycopesticide containing *M. anisopliae* has been developed worldwide to control numerous insect pests including *R. ferrugineus*. However, there is lack of information of the *M. anisopliae* formulated as oil emulsion for the control of RPW (Francardi *et al.*, 2016). Therefore, this study was conducted to expand the knowledge on a specific entomopathogenic fungus *M. anisopliae* and to examine a novel approach in preparing oil nanoemulsions formulations as a microbial biopesticide with following objectives.

- 1) To isolate and identify of *M. anisopliae*.
- 2) Prepare and characterize the oil based emulsion formulation of *M. anisopliae*.
- 3) To evaluate the biological effectiveness of oil based formulation against RPW.
- 4) To investigate the effect of the oil emulsion formulation on the penetration of germ tube through RPW cuticle.

It is expected that this study will contribute towards an improved understanding of the fundamental aspects of oil based nano-emulsion of biopesticide. Accordingly, the information obtained can be further exploited for their proper application that will contribute towards the proper pest management of palm oil in Malaysia and other palm oil producing countries.



REFERENCES

- Abdel-Samad, S. S., Mahmoud, B. A., & Abbas, M. S. T. (2011). Evaluation of the Fungus, *Beauveria bassiana* (Bals.) Vuill as a Bio-Control Agent against the Red Palm Weevil, *Rhynchophorus ferrugineus* (Oliv.)(Coleoptera: Curculionidae). *Egyptian Journal of Biological Pest Control*, 21(2), 125.
- Abe, F., Hata, K., & Sone, K. (2009). Life history of the red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae), in Southern Japan. *Florida Entomologist*, 92(3), 421-425.
- Aboofazeli, R., & Lawrence, M. (1994). Investigations into the formation and characterization of phospholipid microemulsions. II. Pseudo-ternary phase diagrams of systems containing water-lecithin-isopropyl myristate and alcohol: influence of purity of lecithin. *International Journal of Pharmaceutics*, 106(1), 51-61.
- Abraham, V. A., Shuaibi, M. A., Faleiro, J. R., Abozuhairah, R. A., & Vidyasagar, P. S. (1998). An integrated management approach for red palm weevil *Rhynchophorus ferrugineus* Oliv. a key pest of date palm in the Middle East. *Journal of Agricultural and Marine Sciences [JAMS]*, 3(1), 77-83.
- Aktar, W., Sengupta, D., & Chowdhury, A. (2009). Impact of pesticides use in agriculture: their benefits and hazards. *Interdisciplinary toxicology*, 2(1), 1-12
- Al-Dosary, N. M. N., Al-Dobai, S., & Faleiro, J. R. (2016). Review on the management of red palm weevil *Rhynchophorus ferrugineus* Olivier in date palm *Phoenix dactylifera* L. *Emirates Journal of Food and Agriculture*, 28(1), 34.
- Alves, R. T., Bateman, R. P., Gunn, J., Prior, C., & Leather, S. R. (2002). Effects of different formulations on viability and medium-term storage of *Metarhizium anisopliae* conidia. *Neotropical Entomology*, 31, 91-99.
- Alves, R.T., Bateman, R.P., Prior, C., Leather, S.R.(1998). Effects of simulated solar radiation on conidial germination of *Metarhizium anisopliae* in different formulations. *Crop Protection*. 17 (8), 675–679.
- Alves, S. B. (1998). Controle Microbiano de Insetos (pp. 1163). Piracicaba: FEALQ.
- Anggraeni, T., & Putra, R. E. (2011). Cellular and humoral immune defenses of *Oxya japonica* (Orthoptera: Acrididae) to entomopathogenic fungi *Metarhizium anisopliae*. *Entomological Research*, 41(1), 1-6.
- Arruda, W., Lübeck, I., Schrank, A., & Vainstein, M. H. (2005). Morphological alterations of *Metarhizium anisopliae* during penetration of *Boophilus microplus* ticks. *Experimental and Applied Acarology*, 37(3), 231-244.

- Azeem, A., Rizwan, M., Ahmad, F. J., Iqbal, Z., Khar, R. K., Aqil, M., & Talegaonkar, S. (2009). Nanoemulsion components screening and selection: a technical note. *AapsPharmscitech*, 10(1), 69-76.
- Baboota, S., Shakeel, F., Ahuja, A., Ali, J., & Shafiq, S. (2007). Design, development and evaluation of novel nanoemulsion formulations for transdermal potential of celecoxib. *Acta pharmaceutica*, 57(3), 315-332.
- Barranco, P., De La Pena, J., & Cabello, T. (1996). El picudorojo de las palmeras, *Rhynchophorus ferrugineus* (Olivier), nuevaplaga en Europa. (Coleoptera, Curculionidae). *Phytoma-España*, 76, 36-40.
- Bateman, R., & Alves, R. (2000). Delivery systems for mycoinsecticides using oil-based formulations. *Aspects of Applied Biology*, 57, 163-170.
- Bateman, R., Carey, M., Moore, D., & Prior, C. (1993). The enhanced infectivity of *Metarhizium flavoviride* in oil formulations to desert locusts at low humidities. *Annals of Applied Biology*, 122(1), 145-152.
- Batta, Y. A. (2003). Production and testing of novel formulations of the entomopathogenic fungus *Metarhizium anisopliae* (Metschnikoff) Sorokin (Deuteromycotina: Hyphomycetes). *Crop Protection*, 22(2), 415-422.
- Batta, Y. A. (2016). Invert emulsion: Method of preparation and application as proper formulation of entomopathogenic fungi. *MethodsX*, 3, 119-127.
- Beattie, G.A.C. & SMITH, D. (1993). Citrus leafminer. Agfact HZ.AE.4, 2nd edition, Sydney: NSW Agriculture and Fisheries, Australia.
- Becker, L. C., Bergfeld, W. F., Belsito, D. V., Hill, R. A., Klaassen, C. D., Liebler, D. C., ... & Gill, L. J. (2015). Safety assessment of glycerin as used in cosmetics. Washington, DC, Cosmetic Ingredient Review, 1-24.
- Benjamin, M. A., Zhioua, E., & Ostfeld, R. S. (2002). Laboratory and field evaluation of the entomopathogenic fungus *Metarhizium anisopliae* (Deuteromycetes) for controlling questing adult *Ixodes scapularis* (Acari: Ixodidae). *Journal of Medical Entomology*, 39(5), 723-728.
- Bernardi, D. S., Pereira, T. A., Maciel, N. R., Bortoloto, J., Viera, G. S., Oliveira, G. C., & Rocha-Filho, P. A. (2011). Formation and stability of oil-in-water nanoemulsions containing rice bran oil: in vitro and in vivo assessments. *Journal of nanobiotechnology*, 9(1), 44.
- Bidochka, M.J., Kamp, A.M. & Amritha de Croos, J.N. (2000). Insect Pathogenic Fungi: From Genes to Populations. Fungal Pathology (ed J. Kronstad), pp. 171-193. Springer Netherlands.

- Bischoff, J. F., Rehner, S. A., & Humber, R. A. (2006). *Metarhizium frigidum* sp. nov.: a cryptic species of *M. anisopliae* and a member of the *M. flavoviride* complex. *Mycologia*, 98(5), 737-745.
- Bischoff, J. F., Rehner, S. A., & Humber, R. A. (2009). A multilocus phylogeny of the *Metarhizium anisopliae* lineage. *Mycologia*, 101(4), 512-530.
- Blanford, S., Jenkins, N. E., Christian, R., Chan, B. H., Nardini, L., Osae, M., ... & Thomas, M. B. (2012). Storage and persistence of a candidate fungal biopesticide for use against adult malaria vectors. *Malaria Journal*, 11(1), 354.
- Bogran, C. E., Ludwig, S., & Metz, B. (2006). Using oils as pesticides. Texas FARMER .Collection.
- Boonme, P., Krauel, K., Graf, A., Rades, T., & Junyaprasert, V. B. (2006). Characterization of microemulsion structures in the pseudoternary phase diagram of isopropyl palmitate/water/Brij 97: 1-butanol. *AapsPharmscitech*, 7(2), E99-E104.
- Bouchemal, K., Briançon, S., Perrier, E., & Fessi, H. (2004). Nano-emulsion formulation using spontaneous emulsification: solvent, oil and surfactant optimisation. *International Journal of pharmaceutics*, 280(1), 241-251.
- Boucias, D. G., & Pendland, J. C. (1991). Attachment of mycopathogens to cuticle. In *The fungal spore and disease initiation in plants and animals* (pp. 101-127). Springer US.
- Braga, G. U., Flint, S. D., Messias, C. L., Anderson, A. J., & Roberts, D. W. (2001). Effects of UVB Irradiance on Conidia and Germinants of the Entomopathogenic Hyphomycete *Metarhizium anisopliae*: A Study of Reciprocity and Recovery. *Photochemistry and Photobiology*, 73(2), 140-146.
- Bukhari, T., Takken, W., & Koenraadt, C. J. (2011). Development of *Metarhizium anisopliae* and *Beauveria bassiana* formulations for control of malaria mosquito larvae. *Parasites & vectors*, 4(1), 23.
- Burgess, H. D. (1998). *Formulation of Microbial Biopesticides*. Kluwer Academic Publishers. London, U.K. 412 pgs.
- Butt, T. M., Greenfield, B. P., Greig, C., Maffei, T. G., Taylor, J. W., Piasecka, J., ... & Quesada-Moraga, E. (2013). *Metarhizium anisopliae* pathogenesis of mosquito larvae: a verdict of accidental death. *PLoS One*, 8(12), e81686.
- Camargo, M. G., Golo, P. S., Angelo, I. C., Perinotto, W. M. S., Sá, F. A., Quinelato, S., & Bittencourt, V. R. E. P. (2012). Effect of oil-based formulations of acaripathogenic fungi to control *Rhipicephalus microplus* ticks under laboratory conditions. *Veterinary Parasitology*, 188(1), 140-147

- Camargo, M. G., Marciano, A. F., Sá, F. A., Perinotto, W. M. S., Quinelato, S., Gôlo, P. S., Bittencourt, V. R. E. P. (2014). Commercial formulation of *Metarhizium anisopliae* for the control of *Rhipicephalus microplus* in a pen study. *Veterinary Parasitology*, 205(1), 271-276.
- Castro, M. J., Ojeda, C., & Cirelli, A. F. (2014). Advances in surfactants for agrochemicals. *Environmental chemistry letters*, 12(1), 85-95.
- Chagnon, M., Kreutzweiser, D., Mitchell, E. A., Morrissey, C. A., Noome, D. A., & Van der Sluijs, J. P. (2015). Risks of large-scale use of systemic insecticides to ecosystem functioning and services. *Environmental Science and Pollution Research*, 22(1), 119-134.
- Chen, F., Wang, Y., Zheng, F., Wu, Y., & Liang, W. (2000). Studies on cloud point of agrochemical microemulsions. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 175(1-2), 257-262.
- Chomczynski, P., & Sacchi, N. (2006). The single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction: twenty-something years on. *Nature protocols*, 1(2), 581.
- Cox, C., & Sorgan, M. (2006). Unidentified inert ingredients in pesticides: implications for human and environmental health. *Environmental health perspectives*, 1803-1806.
- D Gianeti, M., AL Wagemaker, T., C Seixas, V., & MBG Maia Campos, P. (2012). The use of nanotechnology in cosmetic formulations: the influence of vehicle in the vitamin A skin penetration. *Current Nanoscience*, 8(4), 526-534.
- da CR Rodrigues, E., Ferreira, A. M., Vilhena, J. C., Almeida, F. B., Cruz, R. A., Florentino, A. C., ... & Fernandes, C. P. (2014). Development of a larvicidal nanoemulsion with Copaiba (*Copaifera duckei*) oleoresin. *Revista Brasileira de Farmacognosia*, 24(6), 699-705.
- Damalas, C. A., & Eleftherohorinos, I. G. (2011). Pesticide exposure, safety issues, and risk assessment indicators. *International Journal of environmental research and public health*, 8(5), 1402-1419.
- Daoust, R. A., & Roberts, D. W. (1983). Studies on the prolonged storage of *Metarhizium anisopliae* conidia: effect of temperature and relative humidity on conidial viability and virulence against mosquitoes. *Journal of invertebrate pathology*, 41(2), 143-150.
- de Azevedo Ribeiro, R. C., Barreto, S. M. A. G., Ostrosky, E. A., da Rocha-Filho, P. A., Veríssimo, L. M., & Ferrari, M. (2015). Production and Characterization of Cosmetic Nanoemulsions Containing *Opuntia ficus-indica* (L.) Mill Extract as Moisturizing Agent. *Molecules*, 20, 2492-2509.

- de Faria, M. R., & Wraight, S. P. (2007). Mycoinsecticides and mycoacaricides: a comprehensive list with worldwide coverage and international classification of formulation types. *Biological Control*, 43(3), 237-256.
- Dembilio, Ó., & Jacas, J. A. (2011). Basic bio-ecological parameters of the invasive Red Palm Weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae), in *Phoenix canariensis* under Mediterranean climate. *Bulletin of entomological research*, 101(02), 153-163.
- Dembilio, Ó., & Jaques, J. A. (2015). Biology and Management of Red Palm Weevil. In Sustainable Pest Management in Date Palm: Current Status and Emerging Challenges (pp. 13-36). *Springer International Publishing*.
- Dembilio, Ó., Quesada-Moraga, E., Santiago-Álvarez, C., & Jacas, J. A. (2010). Potential of an indigenous strain of the entomopathogenic fungus *Beauveria bassiana* as a biological control agent against the Red Palm Weevil, *Rhynchophorus ferrugineus*. *Journal of invertebrate pathology*, 104(3), 214-221.
- Destéfano, R. H. R., Destéfano, S. A. L., & Messias, C. L. (2004). Detection of *Metarhizium anisopliae* var. *anisopliae* within infected sugarcane borer *Diatraea saccharalis* (Lepidoptera, Pyralidae) using specific primers. *Genetics and Molecular Biology*, 27(2), 245-252.
- Dorner, J.W. & Lamb, M.C. (2006). Development and commercial use of afla-guard®, an aflatoxin biocontrol agent. *Mycotoxin Research*, 21:33-38.
- Driver, F., Milner, R. J., & Trueman, J. W. (2000). A taxonomic revision of *Metarhizium* based on a phylogenetic analysis of rDNA sequence data. *Mycological Research*, 104(2), 134-150.
- Du, Z., Wang, C., Tai, X., Wang, G., & Liu, X. (2016). Optimization and characterization of biocompatible oil-in-water nanoemulsion for pesticide delivery. *ACS Sustainable Chemistry & Engineering*, 4(3), 983-991.
- Echlin, P. (2011). Handbook of sample preparation for scanning electron microscopy and X-ray microanalysis. *Springer Science & Business Media*.
- Ekesi, S., Maniania, N. K., Mohamed, S. A., & Lux, S. A. (2005). Effect of soil application of different formulations of *Metarhizium anisopliae* on African tephritid fruit flies and their associated endoparasitoids. *Biological Control*, 35(1), 83-91.
- ElKichaoui, A. Y., Bara'a, A., & El-Hindi, M. W. (2017). Isolation, Molecular Identification and under Lab Evaluation of the Entomopathogenic Fungi *M. anisopliae* and *B. bassiana* against the Red Palm Weevil *R. ferrugineus* in Gaza Strip. *Advances in Microbiology*, 7(01), 109.

- EPPO (European and Mediterranean Plant Protection Organization). (2008). Data sheet on quarantine pests: *Rhynchophorus ferrugineus*. *EPPO Bull.*, 38, 55-59.
- Erler, F., & Ates, A. O. (2015). Potential of Two Entomopathogenic Fungi, *Beauveria bassiana* and *Metarhizium anisopliae*, as Biological Control Agents Against the *June beetle* (Coleoptera: Scarabaeidae)(vol 15, pg 44, 2015). *Journal of Insect Science*, 15(1).
- Fairhurst, D., & Loxley, A. (2008). Micro-and nano-encapsulation of water-and oil-soluble actives for cosmetic and pharmaceutical applications. Science and Applications of Skin Delivery Systems. *Allured Publishing Corporation*, 313-336.
- Faleiro, J. R. (2006). A review of the issues and management of the red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Rhynchophoridae) in coconut and date palm during the last one hundred years. *International Journal of tropical Insect Science*, 26(03), 135-154.
- Forgiarini, A., Esquena, J., Gonzalez, C., & Solans, C. (2001). Formation of nano-emulsions by low-energy emulsification methods at constant temperature. *Langmuir*, 17(7), 2076-2083.
- Francardi, V., Benvenuti, C., Barzanti, G. P., & Roversi, P. F. (2016). *Metarhizium anisopliae* biopesticides and fungus isolates: control efficacy against *Rhynchophorus ferrugineus* (Olivier)(Coleoptera Dryophthoridae) on different contamination substrata. *Redia*, 98(1), 25-29.
- Francardi, V., Benvenuti, C., Barzanti, G. P., & Roversi, P. F. (2013). Autocontamination trap with entomopathogenic fungi: a possible strategy in the control of *Rhynchophorus ferrugineus* (Olivier)(Coleoptera Curculionidae). *Redia*, 96, 57-67.
- Francardi, V., Benvenuti, C., Roversi, P. F., Rumine, P., & Barzanti, G. (2012). Entomopathogenicity of *Beauveria bassiana* (Bals.) Vuill. and *Metarhizium anisopliae* (Metsch.) Sorokin isolated from different sources in the control of *Rhynchophorus ferrugineus* (Olivier) (Coleoptera Curculionidae). *Redia*, 95, 49-55.
- Fuxa, J.R. & Richter, A.R. (1999). Classical biological control in an ephemeral crop habitat with *Anticarsia gemmatilis* nucleopolyhedrovirus. *Biological Control*, 44: 403–419.
- Gill, H. K., & Garg, H. (2014). Pesticides: environmental impacts and management strategies *Pesticides-Toxic Aspects*: InTech.
- Gindin, G. L. E. V. S. K. I., Levski, S., Glazer, I., & Soroker, V. (2006). Evaluation of the entomopathogenic fungi *Metarhizium anisopliae* and *Beauveria bassiana* against the red palm weevil *Rhynchophorus ferrugineus*. *Phytoparasitica*, 34(4), 370-379.

- Gopal, E. (1968). Principles of emulsion formation (pp. 1-75): *Academic Press: London*.
- Gopal, M., Gupta, A., Sathiamma, B., & Nair, C. R. (2001). Control of the coconut pest *Oryctes rhinoceros* L. using the *Oryctes* virus. *International Journal of Tropical Insect Science*, 21(2), 93-101.
- Gouli, V., Gouli, S., Brownbridge, M., Skinner, M., & Parker, B. L. (2005). Manual for Mass Production of Entomopathogenic fungi in developing countries with particular reference to *Beauveria bassiana* and *Metarhizium anisopliae*. *University of Vermont, Entomology Research Laboratory, UK*.
- Grewal, P.S.; Power, K.T. & Shetler, D.J. (2001). Neonicotinoid insecticides alter diapause behavior and survival of overwintering white grubs (Coleoptera: Scarabaeidae). *Pest Management Science*, 57:852–857.
- Güerri-Agulló, B., Gómez-Vidal, S., Asensio, L., Barranco, P., & Lopez-Llorca, L. V. (2010). Infection of the red palm weevil (*Rhynchophorus ferrugineus*) by the entomopathogenic fungus *Beauveria bassiana*: a SEM study. *Microscopy research and technique*, 73(7), 714-725.
- Gupta, A., Eral, H. B., Hatton, T. A., & Doyle, P. S. (2016). Nanoemulsions: formation, properties and applications. *Soft matter*, 12(11), 2826-2841.
- Hadzir, N. M., Basri, M., Rahman, M. B. A., Salleh, A. B., Rahman, R. N. Z. R. A., & Basri, H. (2013). Phase behaviour and formation of fatty acid esters nanoemulsions containing piroxicam. *AAPS PharmSciTech*, 14(1), 456-463.
- Hajek, A. E., & St. Leger, R. J. (1994). Interactions between fungal pathogens and insect hosts. *Annual review of entomology*, 39(1), 293-322.
- Hajek, A.E. & Delalibera Jr., I. (2010). Fungal pathogens as classical biological control agents against arthropods. *Biological Control*, 55, 1, 147-158, ISSN 1386-6141.
- Hajek, A.E.; McManus, M.L. & Ju'nior, I.D. (2007). A review of introductions of pathogens and nematodes for classical biological control of insects and mites. *Biological Control*, 41: 1-13.
- Hernandez-Marante, D., Folk, F., Sanchez, A., & Fernandez-Escobar, R. (2003). Control of red palm weevil (*Rhynchophorus ferrugineus* Olivier) using trunk injections and foliar sprays. *Boletin de Sanidad Vegetal, Plagas*, 29(4), 563-573.
- Hinaje, M., Ford, M., Banting, L., Arkle, S., & Khambay, B. (2002). An investigation of the ionophoric characteristics of destruxin A. *Archives of biochemistry and biophysics*, 405(1), 73-77.

- Holder, D. J., & Keyhani, N. O. (2005). Adhesion of the entomopathogenic fungus *Beauveria(Cordyceps) bassiana* to substrata. *Applied and environmental microbiology*, 71(9), 5260-5266.
- Hunt, T. R., Moore, D., Higgins, P. M., & Prior, C. (1994). Effect of sunscreens, irradiance and resting periods on the germination of *Metarhizium flavoviride* conidia. *Biological Control*, 39(3), 313-322.
- Hunt, V. L., & Charnley, A. K. (2011). The inhibitory effect of the fungal toxin, destruxin A, on behavioural fever in the desert locust. *Journal of insect physiology*, 57(10), 1341-1346.
- Hussain, A., Tian, M. Y., Ahmed, S., & Shahid, M. (2012). Current status of entomopathogenic fungi as mycoinsecticides and their inexpensive development in liquid cultures. In Zoology. InTech. <https://cdn.intechopen.com/pdfs-wm/32854.pdf>
- Ibrahim, L., Butt, T., Beckett, A., & Clark, S. (1999). The germination of oil-formulated conidia of the insect pathogen, *Metarhizium anisopliae*. *Mycological Research*, 103(07), 901-907.
- Idris, A. B., Mokhtaruddin, H., Zazali, C., Nurul, W., Yaakop, S., & Hazmi, I. R. (2014). The potential of red palm weevil infesting and destroying oil palm industry in Malaysia. *Planter*, 90(1058), 329-335.
- Ince, S., Porcelli, F., & Al-Jboory, I. (2011). Egg laying and egg laying behavior of red palm weevil *Rhynchophorus ferrugineus* (Olivier) 1790 (Coleoptera: Curculionidae). *Agriculture and Biology Journal of North America*, 2(11), 1368-1374.
- Inglis, G. D., Goettel, M. S., Butt, T. M., & Strasser, H. E. R. M. A. N. N. (2001). Use of hyphomycetous fungi for managing insect pests. *Fungi as biocontrol agents*, 23-69.
- Ivanković, T., & Hrenović, J. (2010). Surfactants in the environment. *Arhiv za higijenu rada i toksikologiju*, 61(1), 95-109.
- Jackson, M. A., Dunlap, C. A., & Jaronski, S. T. (2010). Ecological considerations in producing and formulating fungal entomopathogens for use in insect biocontrol. *BioControl*, 55(1), 129-145.
- Jaronski, S. T. (2010). Ecological factors in the inundative use of fungal entomopathogens. *BioControl*, 55(1), 159-185.
- Jin, X., Streett, D. A., Dunlap, C. A., & Lyn, M. E. (2008). Application of hydrophilic-lipophilic balance (HLB) number to optimize a compatible non-ionic surfactant for dried aerial conidia of *Beauveria bassiana*. *Biological Control*, 46(2), 226-233.

- Justin, C., Leelamathi, M., Thangaselvabai, T., & Johnson, S. B. (2008). Bioecology and management of the Red palm weevil, *Rhynchophorus ferrugineus* Oliv.(Coleoptera: Curculionidae) on Coconut–A review. *Agricultural Reviews*, 29(2), 117-124.
- Kaakeh, W., Abou-Nour, M. M., & Khamis, A. A. (2001). Mass rearing of the red palm weevil, *Rhynchophorus ferrugineus* Olivier, on sugarcane and artificial diets for laboratory studies: Illustration of methodology. In *Proc. 2nd International. Conferenc. Date Palms, Al-Ain, UAE* (pp. 25-27).
- Kaaya, G. P., & Hassan, S. (2000). Entomogenous fungi as promising biopesticides for tick control. *Experimental & applied acarology*, 24(12), 913-926.
- Kalra, A. (2013). *Preparation and evaluation of oil-in-water self-nanoemulsifying systems with potential for pulmonary delivery* (Order No. 1537559). Available from ProQuest Dissertations & Theses Global: Health & Medicine. (1364895962).
- Kershaw, M. J., Moorhouse, E. R., Bateman, R., Reynolds, S. E., & Charnley, A. K. (1999). The Role of Destruxins in the Pathogenicity of *Metarhizium anisopliae* for Three Species of Insect. *Journal of Invertebrate Pathology*, 74(3), 213-223.
- Kirubakaran, S. A., Sathish-Narayanan, S., Revathi, K., Chandrasekaran, R., & Senthil-Nathan, S. (2014). Effect of oil-formulated *Metarhizium anisopliae* and *Beauveria bassiana* against the rice leaffolder *Cnaphalocrocis medinalis* Guenée (Lepidoptera: Pyralidae). *Archives of Phytopathology and Plant Protection*, 47(8), 977-992.
- Kogan, M. (1998). Integrated pest management – historical perspectives and contemporary developments. *Annual Review of Entomology*, 43, 243-270, ISSN 0066-4170.
- Koul, O. (2012). Microbial biopesticides: Opportunities and challenges. *Biocontrol News and Information*, 33(2), 1R.
- Lacatusu, I., Badea, N., Murariu, A., & Meghea, A. (2011). The encapsulation effect of UV molecular absorbers into biocompatible lipid nanoparticles. *Nanoscale research letters*, 6(1), 73
- Lacey, L. A. (Ed.). (1997). *Manual of techniques in insect pathology*. Academic Press.
- Lacey, L. A., & Kaya, H. K. (Eds.). (2007). *Field manual of techniques in invertebrate pathology: application and evaluation of pathogens for control of insects and other invertebrate pests*. Springer Science & Business Media.
- Leemon, D. M., & Jonsson, N. N. (2008). Laboratory studies on Australian isolates of *Metarhizium anisopliae* as a biopesticide for the cattle tick *Boophilus microplus*. *Journal of invertebrate pathology*, 97(1), 40-49.

- Leland, J. E. (2001). Environmental-stress tolerant formulations of *Metarhizium anisopliae* var. *acridum* for control of African desert locust (*Schistocerca gregaria*) (Doctoral dissertation, Virginia Polytechnic Institute and State University).
- Lin, J. J., & Lin, S. F. (2003). Size, volume fraction, and nucleation of Stober silica
- Llácer, E., Santiago-Álvarez, C., & Jacas, J. A. (2013). Could sterile males be used to vector a microbiological control agent? The case of *Rhynchophorus ferrugineus* and *Beauveria bassiana*. *Bulletin of entomological research*, 103(2), 241-250.
- LUBILOSA. Biological control. [cited July/27 2007]. Available from <http://www.lubilosa.org>.
- Mahmud, A. I., Farminhao, J., & Viez, E. R. (2015). Red palm weevil (*Rhynchophorus ferrugineus* Olivier, 1790): threat of palms. *Journal of Biological Sciences*, 15(2), 56-67.
- McClatchie, G. V., Moore, D., Bateman, R. P., & Prior, C. (1994). Effects of temperature on the viability of the conidia of *Metarhizium flavoviride* in oil formulations. *Mycological Research*, 98(7), 749-756.
- Ment, D., Gindin, G., Glazer, I., Perl, S., Elad, D., & Samish, M. (2010). The effect of temperature and relative humidity on the formation of *Metarhizium anisopliae* chlamydospores in tick eggs. *Fungal biology*, 114(1), 49-56.
- Merghem, A. (2011). Susceptibility of the red palm weevil, *Rhynchophorus ferrugineus* (olivier) to the green muscardine fungus, *Metarhizium anisopliae* (metsch.) in the laboratory and in palm trees orchards. *Egyptian Journal of Biological Pest Control*, 21(2), 179.
- Meyling, N. V., & Eilenberg, J. (2006). Occurrence and distribution of soil borne entomopathogenic fungi within a single organic agroecosystem. *Agriculture, ecosystems & environment*, 113(1), 336-341.
- Miller, G.T. (2004), Sustaining the Earth, 6th ed. *Thompson Learning, Inc. Pacific Grove, California*.
- Miller, P. C. H., & Ellis, M. B. (2000). Effects of formulation on spray nozzle performance for applications from ground-based boom sprayers. *Crop protection*, 19(8), 609-615.
- Milner, R. J.(1992). Selection and characterization of strains of *Metarhizium anisopliae* for control of soil insects in Australia, in: C. J. Lomer, C. Prior, (Eds.), *Biological Control of Locust and Grasshoppers*. *CAB International, Melksham*, pp. 200-207.

- Mishra, S., Kumar, P., & Malik, A. (2013). Evaluation of *Beauveria bassiana* spore compatibility with surfactants. *World Acad Sci Eng Technol*, 73, 109-113.
- Mizzi, S., Dandria, D., Mifsud, D., & Longo, S. (2009). The Red Palm Weevil, *Rhynchophorus ferrugineus* (Olivier, 1790) in Malta (Coleoptera: Curculionoidea). *Bulletin of the entomological Society of Malta*, 2, 111-121.
- Mochi, D.A., Monteiro, A.C., Machado, A.C.R. & Yoshida, L. (2010) Efficiency of entomopathogenic fungi in the control of eggs and larvae of the horn fly *Haematobia irritans* (Diptera: Muscidae). *Veterinary Parasitology*, 167, 62–66.
- Moino Jr, A., Alves, S. B., Lopes, R. B., Neves, P. M. O. J., Pereira, R. M., & Vieira, S. A. (2002). External development of the entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae* in the subterranean termite *Heterotermes tenuis*. *Scientia agricola*, 59(2), 267-273.
- Moore, D., Bridge, P. D., Higgins, P. M., Bateman, R. P., & Prior, C. (1993). Ultra-violet radiation damage to *Metarhizium flavoviride* conidia and the protection given by vegetable and mineral oils and chemical sunscreens. *Annals of Applied Biology*, 122(3), 605-616.
- Moore, D., Higgins, P., & Lomer, C. (1996). Effects of simulated and natural sunlight on the germination of conidia of *Metarhizium flavoviride* Gams and Rozsypal and interactions with temperature. *Biocontrol Science and Technology*, 6(1), 63-76.
- Morales, D., Gutiérrez, J. M., Garcia-Celma, M. J., & Solans, Y. C. (2003). A study of the relation between bicontinuous microemulsions and oil/water nano-emulsion formation. *Langmuir*, 19(18), 7196-7200.
- Moslim, R., Kamarudin, N., & Wahid, M. B. (2009). Pathogenicity of granule formulations of *Metarhizium anisopliae* against the larvae of the oil palm rhinoceros beetle, *Oryctes rhinoceros* (L.). *Journal of Oil Palm Research*, 21, 602-612.
- Mulqueen, P. (2003). Recent advances in agrochemical formulation. *Advances in colloid and interface science*, 106(1), 83-107.
- Murphy, S. T., & Briscoe, B. R. (1999). The red palm weevil as an alien invasive: biology and the prospects for biological control as a component of IPM. *Biocontrol news and information*, 20, 35N-46N.
- Mwamburi, L. A., Laing, M. D., & Miller, R. M. (2015). Effect of surfactants and temperature on germination and vegetative growth of *Beauveria bassiana*. *Brazilian Journal of Microbiology*, 46(1), 67-74.
- Myers, D. (1990). Surfaces, interfaces and colloids. New York etc.: Wiley-Vch.

- Myers, D. (2005). Surfactant science and technology. John Wiley & Sons. nanoparticles. *Journal of Colloid and Interface Science*, (258), 159.
- Neethirajan, S., & Jayas, D. S. (2011). Nanotechnology for the food and bioprocessing industries. *Food and bioprocess technology*, 4(1), 39-47.
- Ogawa, S., Decker, E. A., & McClements, D. J. (2004). Production and characterization of o/w emulsions containing droplets stabilized by lecithin–chitosan–pectin multilayered membranes. *Journal of agricultural and food chemistry*, 52(11), 3595-3600.
- Paixão, F. R. S., Muniz, E. R., Barreto, L. P., Bernardo, C. C., Mascarin, G. M., Luz, C., & Fernandes, É. K. K. (2017). Increased heat tolerance afforded by oil-based conidial formulations of *Metarhizium anisopliae* and *Metarhizium robertsii*. *Biocontrol Science and Technology*, 27(3), 324-337.
- Pedrini, N., Crespo, R., & Juárez, M. P. (2007). Biochemistry of insect epicuticle degradation by entomopathogenic fungi. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 146(1–2), 124-137.
- Pena, J. E., Osborne, L. S., & Duncan, R. E. (1996). Potential of fungi as biocontrol agents of *polyphagotarsonemus latus* (Acari: Tarsonemidae). *Entomophaga*, 41(1), 27.
- Polar, P., Kairo, M. T., Moore, D., Pegram, R., & John, S. A. (2005). Comparison of water, oils and emulsifiable adjuvant oils as formulating agents for *Metarhizium anisopliae* for use in control of *Boophilus microplus*. *Mycopathologia*, 160(2), 151-157.
- Prior, C., Jollands, P., and Le Patourel, G. (1988). Infectivity of oil and water formulations of *Beauveria bassiana* (Deuteromycotina: Hyphomycetes) to the cocoa weevil pest *Pantorhytes plutus* (Coleoptera: Curculionidae). *Journal of Invertebrate Pathology*, 52: 66-72.
- Quesada-Moraga, E., Santos-Quiros, R., Valverde-Garcia, P., & Santiago-Alvarez, C. (2004). Virulence, horizontal transmission, and sublethal reproductive effects of *Metarhizium anisopliae* (Anamorphic fungi) on the German cockroach (Blattodea: Blattellidae). *Journal of invertebrate pathology*, 87(1), 51-58.
- Rebolleda, S., Sanz, M. T., Benito, J. M., Beltrán, S., Escudero, I., & González San-José, M. L. (2015). Formulation and characterisation of wheat bran oil-in-water nanoemulsions. *Food Chemistry*, 167, 16-23.
- Roberts, D. W., & Leger, R. J. S. (2004). *Metarhizium* spp., cosmopolitan insect-pathogenic fungi: mycological aspects. *Advances in applied microbiology*, 54, 1-70.

- Roland, I., Piel, G., Delattre, L., & Evrard, B. (2003). Systematic characterization of oil-in-water emulsions for formulation design. *International Journal of Pharmaceutics*, 263(1), 85-94.
- Rugman-Jones, P. F., Hoddle, C. D., Hoddle, M. S., & Stouthamer, R. (2013). The lesser of two weevils: molecular-genetics of pest palm weevil populations confirm *Rhynchophorus vulneratus* (Panzer 1798) as a valid species distinct from *R. ferrugineus* (Olivier 1790), and reveal the global extent of both. *PloS one*, 8(10), e78379.
- Sambrook, J., & Russell, D. W. (2006). Purification of nucleic acids by extraction with phenol: chloroform. *Cold Spring Harbor Protocols*, 2006(1), pdb-prot4455.
- Sandrin, T. R., TeBeest, D. O., & Weidemann, G. J. (2003). Soybean and sunflower oils increase the infectivity of *Colletotrichum gloeosporioides* f. sp. *aeschynomene* to Northern Jointvetch. *Biological Control*, 26(3), 244-252.
- Sanjeevani, N. A., & Sakeena, M. H. F. (2013). Formulation and characterization of virgin coconut oil (VCO) based emulsion. *International Journal of Scientific and Research Publications*, 3(12).
- Santi, L., e Silva, L. A. D., da Silva, W. O. B., Corrêa, A. P. F., Rangel, D. E. N., Carlini, C. R., . Vainstein, M. H. (2011). Virulence of the entomopathogenic fungus *Metarhizium anisopliae* using soybean oil formulation for control of the cotton stainer bug, *Dysdercus peruvianus*. *World Journal of Microbiology and Biotechnology*, 27(10), 2297-2303.
- Sarodee, B., Pranab, D., Joyarani, P., Himadri, K., Gogoi, N., Puzari, K. C., & Hazarika, G. N. (2016). SEM study on morphological changes in *Metarhizium anisopliae* infected *Aphis craccivora* Koch. *Journal of Biological Control*, 30(1), 29-33.
- Schrank, A., & Vainstein, M. H. (2010). *Metarhizium anisopliae* enzymes and toxins. *Toxicon*, 56(7), 1267-1274.
- Sedighi, N., Abbasipour, H., Askary, H., & SheikhiGorjan, A. (2013). Effect of oil suspended conidia of *Metarhizium anisopliae* var. *major* on mortality of the sunn pest, *Eurygaster integriceps* Puton (Hemiptera: Scutelleridae). *Archives of Phytopathology and Plant Protection*, 46(2), 128-140.
- Segur, J. B. (1953). Uses of glycerine (pp. 238-330). Reinhold, New York.
- Setya, S., Talegaonkar, S., & Razdan, B. K. (2014). Nanoemulsions: formulation methods and stability aspects. *World J. Pharm. Pharm. Sci*, 3(2), 2214-2228.
- Shafiq, S., Shakeel, F., Talegaonkar, S., Ahmad, F. J., Khar, R. K., & Ali, M. (2007). Development and bioavailability assessment of ramipril nanoemulsion formulation. *European Journal of Pharmaceutics and Biopharmaceutics*, 66(2), 227-243.

- Shah PA, Douro-Kpindou OK, Sidibe A, Daffe CO, Van der Pauuw H, Lomer CJ. (1998). Effects of the sunscreen oxybenzone on field efficacy and persistence of *Metarhizium flavoviride* conidia against *Kraussella amabile* (Orthoptera: Acrididae) in Mali, West Africa. *Biocontrol Science and Technology* 8:357-364
- Shah, P. A., Kooyman, C., & Paraíso, A. (1997). Surveys for fungal pathogens of locusts and grasshoppers in Africa and the Near East. *The Memoirs of the Entomological Society of Canada*, 129(S171), 27-35.
- Shanmugam, A., & Ashokkumar, M. (2014). Ultrasonic preparation of stable flax seed oil emulsions in dairy systems—physicochemical characterization. *Food Hydrocolloids*, 39, 151-162.
- Sharma, N., Madan, P., & Lin, S. (2016). Effect of process and formulation variables on the preparation of parenteral paclitaxel-loaded biodegradable polymeric nanoparticles: A co-surfactant study. *Asian Journal of Pharmaceutical Sciences*, 11(3), 404-416.
- Shokrzadeh, M., & Saravi, S. S. S. (2011). Pesticides in Agricultural Products: Analysis, Reduction, Prevention. INTECH Open Access Publisher.
- Silva, W. O. B., Mitidieri, S., Schrank, A., & Vainstein, M. H. (2005). Production and extraction of an extracellular lipase from the entomopathogenic fungus *Metarhizium anisopliae*. *Process Biochemistry*, 40(1), 321-326.
- Smith, K. E., Wall, R., & French, N. P. (2000). The use of entomopathogenic fungi for the control of parasitic mites, *Psoroptes* spp. *Veterinary parasitology*, 92(2), 97-105.
- Soberón-Chávez, G. (Ed.). (2010). Biosurfactants: from genes to applications (Vol. 20). Springer Science & Business Media.
- Sun, X., Yan, W., Qin, W., Zhang, J., Niu, X., Ma, G., & Li, F. (2016a). Screening of tropical isolates of *Metarhizium anisopliae* for virulence to the red palm weevil *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae). *Springer Plus*, 5(1), 1100.
- Sun, X., Yan, W., Zhang, J., Niu, X., Li, F., Qin, W., & Ma, G. (2016b). Frozen section and electron microscopy studies of the infection of the red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae) by the entomopathogenic fungus *Metarhizium anisopliae*. *Springer Plus*, 5(1), 1748.
- Tadros, T. F. (2006). Applied surfactants: principles and applications. John Wiley & Sons.
- Tadros, T. F. (Ed.). (2013). *Emulsion formation and stability*. John Wiley & Sons.

- Talwar, B. H. (2005). Isolation and characterization of entomopathogenic fungi and their effectiveness (Doctoral dissertation, University of Agricultural Sciences, Dharwad).
- Tangthirasunun, N., Poeaim, S., Soyong, K., Sommartya, P., & Popoonsak, S. (2010). Variation in morphology and ribosomal DNA among isolates of *Metarhizium anisopliae* from Thailand. *Journal of Agricultural Technology*, 2, 317-329.
- Thakore, Y. (2006). The biopesticide market for global agricultural use. *Industrial Biotechnology*, 2(3), 194-208.
- Thomas, M. B., & Read, A. F. (2007). Can fungal biopesticides control malaria?. *Nature Reviews Microbiology*, 5(5), 377-383.
- Toledo, A. V., Lenicov, A. D. R., & Lastra, C. L. (2010). Histopathology caused by the entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae*, in the adult planthopper, *Peregrinus maidis*, a maize virus vector. *Journal of Insect Science*, 10(1), 35.
- Tominack, R. L., & Tominack, R. (2000). Herbicide formulations. *Journal of Toxicology: Clinical Toxicology*, 38(2), 129-135.
- Tong, K., Zhao, C., & Sun, D. (2016). Formation of nanoemulsion with long chain oil by W/O microemulsion dilution method. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 497, 101-108.
- Van Valkenburg, W. (1973). Pesticide formulations. Marcel Dekker.
- Vänninen, I. (1996). Distribution and occurrence of four entomopathogenic fungi in Finland: effect of geographical location, habitat type and soil type. *Mycological Research*, 100(1), 93-101.
- Wai, Y. K., BAKAR, A. A., & Azmi, W. A. (2015). Fecundity, Fertility and Survival of Red Palm Weevil (*Rhynchophorus ferrugineus*) Larvae Reared on Sago Palm. *Sains Malaysiana*, 44(10), 1371-1375.
- Wang, C., & St Leger, R. J. (2007). The MAD1 adhesin of *Metarhizium anisopliae* links adhesion with blastospore production and virulence to insects, and the MAD2 adhesin enables attachment to plants. *Eukaryotic cell*, 6(5), 808-816.
- Wessels, J. G. H. (2000). Hydrophobins, unique fungal proteins. *Mycologist*, 14(4), 153-159.
- White, T. J., Bruns, T., Lee, S. J. W. T., & Taylor, J. L. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications*, 18(1), 315-322.

- Wooster, T. J., Golding, M., & Sanguansri, P. (2008). Impact of oil type on emulsion formation and Ostwald ripening stability. *Langmuir*, 24(22), 12758-12765.
- Wraight, S. P., Jackson, M. A., & De Kock, S. L. (2001). Formulation of Fungal Biocontrol Agents. *Fungi as Biocontrol Agents: Progress Problems and Potential*, 253.
- Xavier-Santos, S., Lopes, R. B., & Faria, M. (2011). Emulsifiable oils protect *Metarhizium robertsii* and *Metarhizium pingshaense* conidia from imbibitional damage. *Biological Control*, 59(2), 261-267.
- Xia, Y., Clarkson, J. M., & Charnley, A. K. (2002). Trehalose-hydrolysing enzymes of *Metarhizium anisopliae* and their role in pathogenesis of the tobacco hornworm, *Manduca sexta*. *Journal of invertebrate pathology*, 80(3), 139-147.
- Zhang, S., Xia, Y. X., Kim, B., & Keyhani, N. O. (2011). Two hydrophobins are involved in fungal spore coat rodlet layer assembly and each play distinct roles in surface interactions, development and pathogenesis in the entomopathogenic fungus, *Beauveria bassiana*. *Molecular microbiology*, 80(3), 811-826.
- Zimmermann, G. (2007). Review on safety of the entomopathogenic fungus *Metarhizium anisopliae*. *Biocontrol Science and Technology*, 17(9), 879-920.