



UNIVERSITI PUTRA MALAYSIA

***RESISTANCE MECHANISM OF *Limnocharis flava* (L.) BUCHENAU TO
ACETOHYDROXY ACID SYNTHASE INHIBITORS***

NORAZUA BINTI ZAKARIA

FP 2018 84



**RESISTANCE MECHANISM OF *Limnocharis flava* (L.) BUCHENAU TO
ACETOHYDROXY ACID SYNTHASE INHIBITORS**

By

NORAZUA BINTI ZAKARIA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

February 2018

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

RESISTANCE MECHANISM OF *Limnocharis flava* (L.) Buchenau TO ACETOHYDROXY ACID SYNTHASE INHIBITORS

By

NORAZUA BINTI ZAKARIA

February 2018

Chairman : Muhammad Saiful Ahmad Hamdani, PhD
Faculty : Agriculture

Limnocharis flava, a problematic weed of rice fields, is reported to have multiple resistance to 2,4-D (synthetic auxin), and bensulfuron-methyl (AHAS inhibitor) due to over-reliance on their use. The present study was conducted to appraise the level of resistance and cross-resistance patterns in *L. flava*, and to elucidate the mechanisms endowing its resistance. Morphological responses between susceptible (S) and resistant (R) populations were evaluated with a view of addressing the species resistance profile. The levels and patterns of resistance in the R population were determined by LD₅₀ and GR₅₀ values. Seven AHAS inhibitors (bensulfuron-methyl, metsulfuron-methyl, pyrazosulfuron-ethyl, pyribenzoxim, imazethapyr, penoxsulam, and bispyribac-sodium) were examined for their efficacy at different dosages against the S and R populations, respectively. It was found that the S population was successfully controlled by all AHAS inhibitors but the R population survived the applications of the first four from seven tested inhibitors, indicating high-level of resistance to bensulfuron-methyl, with various levels of cross-resistance to other AHAS inhibitors. To demonstrate the mechanisms endowing resistance in the R population, molecular investigation was carried out by AHAS amino acids sequence comparison on R plants that survived the inhibitors application. Evidently, the plants were endowed with similar AHAS gene mutation of single nucleotide polymorphism (SNP) (GAC to GAG). This resulted in Asp substitution by Glu at amino acid position 376, suggesting point mutation as the molecular basis of resistance. *In vitro* assays were conducted using standards for AHAS inhibitors consisting of bensulfuron-methyl (99.6%), metsulfuron-methyl (99.2%), pyrazosulfuron-ethyl (99.1%), and pyribenzoxim (99.8%) with seven concentrations (0, 0.001, 0.01, 0.1, 1, 10, and 100 µM). The I₅₀ of *in vitro* AHAS activity showed that R population was >83333-, 398-, 172-, and 48-fold greater than S population for the respective AHAS inhibitors. This suggests stronger cross-resistance to sulfonylureas than to pyrimidinyl (thio) benzoate (pyribenzoxim). The result of non-target site mechanism experiments showed that S

and R populations did not differ significantly in their ability to absorb, translocate, and metabolize bensulfuron-methyl. This indicates that non-target site resistance mechanisms of differential metabolism, absorption, and translocation did not confer resistance in the R population. The morphological and physiological characteristics of AHAS inhibitors on S and R populations were examined using progenies of AHAS herbicide-resistant (R) individuals that survived the application of bensulfuron-methyl (Progeny 1) and pyribenzoxim (Progeny 2). Both R and S plants were compared in terms of fresh weight, dry weight, height, epicuticular wax weight, stomata density, leaf area index, and leaf micromorphology for growth assessment. Net photosynthetic rate, stomatal conductance, intercellular carbon dioxide, and transpiration rate were monitored using Progeny 1 from R population with S plants as control. The results showed that R plants carrying AHAS Asp-376-Glu mutation did not differ in the morpho-physiological characteristics and were comparable to that of S plants. In conclusion, the basis of AHAS-inhibitors resistance in *L. flava* was identified to be due to an Asp-376-Glu mutation that reduced sensitivity of the target site to AHAS inhibitors. The current study presents the first report of resistance mechanism in *L. flava*.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**MEKANISASI RINTANGAN OLEH *Limnocharis flava* (L.) Buchenau
TERHADAP PENGHALANG ACETOHYDROXY ACID SYNTHASE**

Oleh

NORAZUA BINTI ZAKARIA

Februari 2018

Pengerusi : Muhammad Saiful Ahmad Hamdani, PhD
Fakulti : Pertanian

Limnocharis flava, spesies rumpai akuatik bermasalah di sawah padi, telah dilaporkan mempunyai kerintangan berbilang kepada racun rumpai yang berasaskan 2,4-D (auksin sintetik) dan bensulfuron-methyl (penghalang AHAS) oleh kerana pergantungan berlebihan terhadap penggunaannya. Kajian ini dijalankan untuk menilai pola kerintangan serta kerintangan bersilang pada *L. flava* dan untuk mengkaji mekanisma kerintangan tersebut. Reaksi morfologi antara populasi rentan (S) dan rintang (R) telah dikaji dengan tujuan penambahan maklumat dalam profil kerintangan pada spesies tersebut. Paras dan corak kerintangan dalam populasi R telah ditentukan dengan nilai LD₅₀ dan GR₅₀. Tujuh penghalang AHAS (bensulfuron-methyl, metsulfuron-methyl, pyrazosulfuron-ethyl, pyribenzoxim, imazethapyr, penoxsulam, dan bispyribac-sodium) telah dikaji keberkesanannya pada dos yang berbeza terhadap populasi S dan R. Keputusan menunjukkan bahawa populasi S telah dapat dikawal keseluruhannya oleh kesemua penghalang AHAS tetapi populasi R terselamat daripada aplikasi empat penghalang pertama daripada tujuh penghalang AHAS yang diuji, menunjukkan daya ketahanan yang tinggi terhadap bensulfuron-methyl dengan beberapa paras kerintangan terhadap penghalang AHAS yang lain. Untuk mengkaji mekanisma kerintangan dalam populasi R, kajian molekular telah dijalankan dengan menggunakan kaedah perbandingan jujukan asid amino pada AHAS ke atas pokok R yang terselamat daripada aplikasi penghalang AHAS. Semua pokok R didapati mempunyai gen mutase yang serupa pada AHAS melalui polimorfisma nucleotid tunggal (SNP) (GAC to GAG). Ini menghasilkan penggantian Asp oleh Glu pada asid amino berkedudukan 376, sekaligus mentafsirkan mutasi tunggal sebagai asas mekanisma kerintangan. Ujian *in vitro* telah dijalankan dengan menggunakan penghalang AHAS standard yang mengandungi bensulfuron-methyl (99.6%), metsulfuron-methyl (99.2%), pyrazosulfuron-ethyl (99.1%), and pyribenzoxim (99.8%) dalam tujuh kepekatan (0, 0.001, 0.01, 0.1, 1, 10, and 100 µM). Aktiviti I₅₀ bagi *in vitro* AHAS menunjukkan bahawa populasi R adalah >83333-, 398-, 172- dan

48-kali lebih tinggi ketahanannya daripada populasi S untuk setiap penghalang AHAS. Ini menunjukkan bahawa populasi R mempunyai kerintangan bersilang lebih tinggi terhadap sulfonilureas berbanding pyrimidinyl(thio)benzoate (pyribenzoxim). Keputusan eksperimen untuk mekanisme luar tapak-sasaran menunjukkan bahawa tiada perbezaan ketara antara populasi S dan R dalam keupayaan menyerap, mengangkut, dan memetabolisma bensulfuron-methyl. Ini membuktikan bahawa mekanisme luar tapak-sasaran iaitu perbezaan metabolisma, penyerapan, dan translokasi tidak menyumbang kepada kerintangan dalam populasi R. Sifat morfologi dan fisiologi penghalang AHAS ke atas populasi S dan R juga telah dikaji dengan menggunakan generasi rintang (R) daripada individu yang terselamat daripada rawatan bensulfuron-methyl (Progeny 1) dan pyribenzoxim (Progeny 2). Kedua-dua pokok R dan S telah dibandingkan dari segi berat segar, berat kering, ketinggian, berat lilin epikutikular, ketumpatan stomata, indeks keluasan daun, dan mikromorfologi daun bagi menilai tumbesaran. Kadar fotosintesis bersih, konduksian stomata, karbon dioksida antara sel, dan kadar transpirasi telah dipantau pada Progeny 1 dari populasi R dengan pokok S sebagai kawalan. Keputusan menunjukkan bahawa pokok R yang mempunyai mutasi AHAS Asp-376-Glu adalah tidak berbeza pada sifat morfologi apabila dibandingkan dengan pokok S. Pada kesimpulannya, asas kerintangan terhadap penghalang AHAS pada populasi R telah dikenalpasti disebabkan oleh mutasi Asp-376-Glu dalam gen AHAS yang mengurangkan kepekaan tapak-sasaran kepada penghalang AHAS. Kajian ini merupakan kajian pertama yang melaporkan mekanisme kerintangan kepada penghalang AHAS pada *L. flava*.

ACKNOWLEDGEMENTS

In the name of Allah, the Most Gracious and the Most Merciful

Alhamdulillah, I would not have been able to complete my dissertation without help, guidance, support and encouragement from following individuals.

Foremost, I would like to express my deepest gratitude to Dr. Muhammad Saiful Ahmad Hamdani, chairman of the supervisory committee for the continuous support of my Ph.D study and research, for his patience, motivation, enthusiasm, and immense knowledge. His guidance has helped me in all the time of research and writing of this thesis.

Besides my supervisor, I would like to express my sincere thanks to my committee members, Prof. Dr. Abdul Shukor Juraimi, Prof. Dr. Mohd Razi Ismail, and Dr. Khairuddin Abdul Rahim for their encouragement, suggestions and insightful comments in the completion of this project and thesis.

I would like to acknowledge the Department of Crop Science, Faculty of Agriculture, UPM. My graduate experience benefitted greatly from the courses and seminars I took. My sincere thanks to all professors and lecturers in the Faculty of Agriculture, UPM for their encouragement, teaching and suggestion throughout the study period. I would like to offer my special thanks to the staff of Department of Crop Science, Department of Plant Protection, Institute of Tropical Agriculture and Food Security (ITAFoS), Institute of Bioscience, UPM, Malaysian Nuclear Agency, and Malaysian Agriculture Research and Development Institute (MARDI) for the laboratory facilities and their kind assistance.

I am grateful for the funding source that allowed me to pursue my study, the Ministry of Higher Education for providing financial assistance in the form of MyBrain Science to pursue my Ph. D at UPM. My sincere thanks to Long- term Research Grant Scheme (LRGS), Food Security Project, Ministry of Higher Education, Malaysia, and Fundamental Research Grant Scheme (07-01-04-1440FR) under the Ministry of Higher Education, Malaysia, for funding this research. My deepest thanks to International Weed Science Society (IWSS) and Asian-Pacific Weed Science Society (APWSS) for the Travel Award in the 26th Asian-Pacific Weed Science Society Conference held in Kyoto, Japan.

I would like to express the deepest appreciation to my groupmate, friends and family who supported me along the journey. For their support and encouragement, I want to express my sincere appreciation to my parents, Zakaria and Hasnah also my mother in law, Fuzah. Special thanks also to my sister Hazuin and my brothers Faiz and Fahmi. Thanks especially to my husband Adreen for always supporting and cheering me up and stood by me through the good and bad times, and my daughter Adriana Zahra, the love of my life.



I certify that a Thesis Examination Committee has met on 27 February 2018 to conduct the final examination of Norazua binti Zakaria on her thesis entitled "Resistance Mechanism of *Limnocharis flava* (L.) Buchenau to Acetohydroxy Acid Synthase Inhibitors" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Thohirah Lee Abdullah @ Lee Chin Chin, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Mohd Rafii bin Yusop, PhD

Professor
Institute of Tropical Agriculture and Food Security
Universiti Putra Malaysia
(Internal Examiner)

Datuk Mohd Fauzi bin Hj. Ramlan, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Stephen Powles, PhD

Professor
The University of Western Australia
Australia
(External Examiner)



NOR AINI AB. SHUKOR, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 June 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Muhammad Saiful Ahmad Hamdani, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Abdul Shukor Juraimi, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Mohd Razi Ismail, PhD

Professor
Institute of Tropical Agriculture and Food Security
Universiti Putra Malaysia
(Member)

Khairuddin Abdul Rahim, PhD

Director
Division of Agrotechnology and Biosciences
Malaysian Nuclear Agency
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No.: Norazua binti Zakaria, GS39157

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Dr. Muhammad Saiful Ahmad Hamdani

Signature: _____
Name of Member
of Supervisory
Committee: Professor Dr. Abdul Shukor Juraimi

Signature: _____
Name of Member
of Supervisory
Committee: Professor Dr. Mohd Razi Ismail

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Khairuddin Abdul Rahim

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xix
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 Plant acetoxyacid synthase (AHAS) and its characteristics	4
2.2 Acetoxyacid synthase inhibitors	6
2.3 Mode of action of AHAS inhibitors	12
2.3.1 Effects of AHAS resistance on plant growth and fitness	14
2.4 Uses of AHAS inhibitors in agriculture	16
2.5 Selectivity in crops towards AHAS inhibitors	16
2.6 Evolution of resistance to AHAS herbicides in weeds	19
2.7 Mechanisms of resistance to AHAS inhibitors in weed species	20
2.7.1 AHAS target site resistance mechanisms	21
2.7.2 AHAS non-target site metabolic resistance	28
2.8 Rice industry in Malaysia	28
2.8.1 Herbicide resistance in Malaysia	29
2.9 Biology, ecology, and distribution of <i>L. flava</i>	29
2.10 <i>Limnocharis flava</i> in Malaysian rice fields	31
2.10.1 Management <i>L. flava</i>	32
2.10.2 Physical and cultural control	32
2.10.3 Chemical control	33
2.10.4 Natural enemies	34
2.10.5 Management of <i>L. flava</i> in Malaysian rice fields and its impact on AHAS-inhibitors resistance occurrence	34
3 PATTERNS OF RESISTANCE TO AHAS INHIBITORS IN <i>Limnocharis flava</i>	36
3.1 Introduction	36
3.2 Materials and Methods	37
3.2.1 Seed collection and plant materials	37

3.2.2	Whole-plant herbicide dose-response	37
3.3	Statistical Analysis	39
3.4	Results and Discussion	40
3.4.1	Whole-plant dose-response to sulfonylurea herbicides	40
3.4.2	Whole-plant dose-response to pyrimidinyl(thio) benzoate herbicides	48
3.4.3	Whole-plant dose-response to imidazolinone and triazolopyrimidine herbicides	48
3.4.4	Correlation between whole-plant dose-response and possible resistance mechanism(s)	52
3.5	Conclusion	56
4	MOLECULAR BASIS OF RESISTANCE TO AHAS INHIBITORS IN <i>Limnocharis flava</i>	57
4.1	Introduction	57
4.2	Materials and Methods	58
4.2.1	Plant materials	58
4.2.2	<i>De novo</i> RNA sequencing	58
4.2.3	Genomic DNA extraction	60
4.2.4	AHAS gene sequencing	60
4.2.5	<i>In vitro</i> AHAS activity and inhibition assay	61
4.2.6	Statistical Analysis	62
4.3	Results and Discussion	62
4.3.1	AHAS gene sequencing revealed an Asp-376-Glu in R population	62
4.3.2	<i>In vitro</i> AHAS Assay	66
4.4	Conclusion	73
5	INVESTIGATION INTO POSSIBLE NON-TARGET SITE MECHANISMS ENDOWING RESISTANCE TO AHAS INHIBITORS IN <i>Limnocharis flava</i>	74
5.1	Introduction	74
5.2	Materials and Method	76
5.2.1	Plant materials	76
5.2.2	Effect of malathion on bensulfuron-methyl resistant <i>Limnocharis flava</i>	76
5.2.3	Absorption and translocation of bensulfuron-methyl in resistant <i>L. flava</i>	77
5.2.3.1	Sample preparation and extraction	77
5.2.3.2	Recovery studies	78
5.2.3.3	High Performance Liquid Chromatography (HPLC) analysis	78
5.3	Statistical Analysis	79
5.3.1	Effect of malathion on bensulfuron-methyl resistant <i>Limnocharis flava</i>	79

5.3.2	Absorption and translocation of bensulfuron-methyl in <i>L. flava</i>	79
5.4	Results and Discussion	79
5.4.1	Effect of malathion on bensulfuron-methyl resistant <i>Limnocharis flava</i>	79
5.4.2	Recovery studies	85
5.4.3	Absorption of bensulfuron-methyl in <i>L. flava</i>	85
5.4.4	Translocation of bensulfuron-methyl in <i>Limnocharis flava</i>	87
5.5	Conclusion	89
6	MORPHOLOGICAL AND PHYSIOLOGICAL CHARACTERISTICS OF AHAS-INHIBITORS RESISTANT AND SUSCEPTIBLE POPULATIONS OF <i>Limnocharis flava</i>	90
6.1	Introduction	90
6.2	Materials and Methods	92
6.2.1	Plant materials	92
6.2.2	Growth and development of resistant and susceptible populations of <i>L. flava</i>	92
6.2.3	Extraction of epicuticular wax of resistant and susceptible <i>L. flava</i>	93
6.2.4	Stomata counts of resistant and susceptible <i>L. flava</i>	93
6.2.5	Scanning Electron Microscope (SEM)	94
6.3	Statistical Analysis	95
6.4	Results and Discussion	95
6.4.1	Growth and morphological characteristics of susceptible and resistant <i>L. flava</i> populations	95
6.4.2	Physiological responses of susceptible and resistant <i>L. flava</i> populations with bensulfuron-methyl treatment	102
6.5	Conclusion	109
7	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATIONS FOR FUTURE STUDIES	110
	REFERENCES	113
	APPENDICES	128
	BIODATA OF STUDENT	135
	LIST OF PUBLICATIONS	136

LIST OF TABLES

Table	Page
2.1 Common uses of some sulfonylurea (SU), imidazolinone (IMI), triazolopyrimidine, pyrimidinyl(thio)benzoate (PTB) and sulfonyl-aminocarbonyl-triazolinone (SCT)	17
2.2 Factors influencing herbicide resistance evolution in weed populations	20
2.3 Summary of acetohydroxyacid synthase amino acid substitutions in field-evolved resistant weed species worldwide	23
3.1 Herbicides information including active ingredient, percent of active ingredient, trade name, company and rates of herbicides applied in post-emergence herbicide application	38
3.2 Survival rate and dry weight of susceptible (S) and resistant (R) <i>Limnocharis flava</i> treated with AHAS inhibitors with different herbicide rates	41
3.3 Parameters of log-logistic analysis of the AHAS herbicide dose required to cause 50% mortality (LD50) and the resistance index (RI) of susceptible (S) and resistant (R) <i>Limnocharis. flava</i> populations	54
3.4 Parameters of log-logistic analysis of the AHAS herbicide dose required to reduce shoot dry weight (GR50) the resistance index (RI) of susceptible (S) and resistant (R) <i>Limnocharis flava</i> populations	55
4.1 Codons and deduced amino acids at eight sites involved in resistance in the susceptible and resistant accessions of <i>Limnocharis flava</i>	63
4.2 Nucleotide sequence alignment of acetohydroxycyacid synthase And derived amino acid around position 376 AHAS inhibitor susceptible <i>Arabidopsis thaliana</i> (NM_114714) and AHAS inhibitors susceptible (S) and resistant ® <i>Limnocharis flava</i>	64
4.3 Parameters of log-logistic analysis of the AHAS herbicide dose required to cause 50% inhibition of AHAS activity (I ₅₀) and the resistance index (RI) of susceptible (S) and resistant (R) <i>Limnocharis flava</i> populations	67
4.4 Effect of herbicide concentration of bensulfuron-methyl, metsulfuron-methyl, pyrazosulfuron-ethyl, and pyribenzoxim on AHAS activity in <i>Limnocharis flava</i> populations	68

5.1	LD ₅₀ and GR ₅₀ values of susceptible(S) and resistant (R) <i>Limnocharis flava</i> population to bensulfuron-methyl in the absence and presence of cytochrome P450 inhibitor malathion	80
5.2	Survival rate of susceptible (S) and resistant (R) <i>Limnocharis flava</i> treated with bensulfuron-methyl with different herbicide rates	81
5.3	Dry weight of susceptible(S) and resistant (R) <i>Limnocharis flava</i> treated with bensulfuron-methyl in an absent and present of malathion with different herbicide rates	82
5.4	Average Recoveries (%) and Relative Standard Deviation (RSD, %) at bensulfuron-methyl concentration of 0.005, 0.01, and 0.02 mg/ml	85
5.5	Percentage of bensulfuron-methyl that was translocated in <i>Limnocharis flava</i> during 72 hours after treatment. Data was pooled between susceptible and resistant populations	89
6.1	The comparison between susceptible (S) and herbicide-resistant (R) (progeny 1 and progeny 2) <i>Limnocharis flava</i> individuals in terms of fresh weight, dry weight, height, epicuticular weight and leaf area at 4-5-leaf stage	95
6.2	Physiological characteristics of susceptible (S) and herbicide-resistant (R) <i>Limnocharis flava</i> individuals at 4-5-leaf stage 24 h before and after bensulfuron-methyl treatment	105

LIST OF FIGURES

Figure		Page
2.1	Schematic of the branched-chain amino acid (BCAA) biosynthesis pathway	5
2.2	Basic chemical structure of sulfonylurea (SU)	7
2.3	Chemical structure of common sulfonylurea (SU) herbicides	8
2.4	Basic chemical structure of imidazolinone (IMI)	9
2.5	Chemical structure of some imidazolinone (IMI) herbicides	9
2.6	Basic chemical structure of triazolopyrimidine (TP)	10
2.7	Chemical structure of some triazolopyrimidine (TP) herbicides	10
2.8	Chemical structure of some pyrimidinyl(thio)benzoate (PTB) herbicides	11
2.9	Chemical structure of the sulfonyl-aminocarbonyl-triazolinone (SCT) herbicides	11
2.10	Biosynthetic pathway of branched chain amino acids and the site of action of herbicidal inhibitors	13
2.11	Schematic representation of conformational adjustments in the <i>At</i> AHAS herbicide binding sites	14
2.12	Sulphonylurea metabolism in several crops	18
2.13	<i>Limnocharis flava</i> growing in a glasshouse at Universiti Putra Malaysia	32
3.1	Survival (A) and shoot dry weight (B) of the susceptible and resistant populations of <i>Limnocharis flava</i> to bensulfuron-methyl 21 days after treatment	45
3.2	Survival (A) and shoot dry weight (B) of the susceptible and resistant populations of <i>Limnocharis flava</i> to metsulfuron-methyl 21 days after treatment	46
3.3	Survival (A) and shoot dry weight (B) of the susceptible and resistant populations of <i>Limnocharis flava</i> to pyrazosulfuron-ethyl 21 days after treatment	47

3.4	Survival (A) and shoot dry weight (B) of the susceptible and resistant populations of <i>Limnocharis flava</i> to pyribenzoxim 21 days after treatment	49
3.5	Survival (A) and shoot dry weight (B) of the susceptible and resistant populations of <i>Limnocharis flava</i> to bispyribac-sodium 21 days after treatment	50
3.6	Survival (A) and shoot dry weight (B) of the susceptible and resistant populations of <i>Limnocharis flava</i> to imazethapyr 21 days after treatment	51
3.7	Survival (A) and shoot dry weight (B) of the susceptible and resistant populations of <i>Limnocharis flava</i> to penoxsulam 21 days after treatment	52
4.1	DNA sequencing results	65
4.2	<i>In vitro</i> inhibition of AHAS activity by bensulfuron-methyl for susceptible (S) and resistant (R) <i>Limnocharis flava</i> populations	70
4.3	<i>In vitro</i> inhibition of AHAS activity by metsulfuron-methyl for susceptible (S) and resistant (R) <i>Limnocharis flava</i> populations	70
4.4	<i>In vitro</i> inhibition of AHAS activity by pyrazosulfuron-ethyl for susceptible (S) and resistant (R) <i>Limnocharis flava</i> populations	71
4.5	<i>In vitro</i> inhibition of AHAS activity by pyribenzoxim for susceptible (S) and resistant (R) <i>Limnocharis flava</i> populations	71
5.1	Dose-response curves for (A) survival and (B) shoot dry weight of susceptible and resistant <i>Limnocharis flava</i> populations treated with a range of bensulfuron-methyl doses in the absence (-M) and presence (+M) of cytochrome P450 inhibitor malathion (M)	83
5.2	Bensulfuron-methyl absorption in susceptible and resistant <i>Limnocharis flava</i> . The data points are mean \pm standard error. Mean with the same letter is not significantly different at $P < 0.05$	87
6.1	Leaves of susceptible (S) <i>Limnocharis flava</i> plants. The sampled leaves were fourth leaves and were collected at 4-5-leaf stage. Bars: 1 cm	99
6.2	Leaves of herbicide-resistant (R) of Progeny 1 <i>Limnocharis flava</i> plants. The sampled leaves were fourth leaves and were collected at 4-5-leaf stage. Bars: 1 cm	100

- 6.3 Leaves of herbicide-resistant (R) of Progeny 2 *Limnocharis flava* plants. The sampled leaves were fourth leaves and were collected at 4-5-leaf stage. Bars: 1 cm 101
- 6.4 Stomatal density and stomata structure in the epidermis of the fourth leaf of untreated susceptible (S) (a; b) and resistant (R) (c; d) *Limnocharis flava* plants showing opening stomata with no significant difference between both populations without bensulfuron-methyl treatment in terms of stomata density and stomata structure. The structure of the stomata consists of a kidney-shaped epidermal cell with an opening in the centre, which is known as a pore 103
- 6.5 Stomatal density and stomata structure of susceptible (S) (a; b) and resistant (R) (c; d) *Limnocharis flava* plants 24 h after bensulfuron-methyl treatment showing opening stomata of the R plant (d) 24h after bensulfuron-methyl treatment, while S plants showed closed stomata (b). This significant result is due to herbicide application; the R plants were not affected whereas S plants showing the early affected-symptom by the herbicide application. 104

LIST OF ABBREVIATIONS

%	Percent
°C	Degree celsius
m ²	Square meter
cm	Centimeter
ha	Hectare
ml	Milliliter
nm	Nanometer
g	Gram
cm ²	Square centimeter
rpm	Revolution per minute
v/v	Volume per volume
m	Meter
ai	Active ingredient
g ai/ha	Gram a.i. per hectare
S	Susceptible
R	Resistant
TSR	Target site resistance
NTSR	Non-target site resistance
RNA	Ribonucleic acid
DNA	Deoxyribonucleic acid
cDNA	Complementary DNA

LD ₅₀	Lethal dose at 50%
GR ₅₀	Growth reduction at 50%
I ₅₀	Inhibition at 50%
RCBD	Randomized complete block design
HPLC	High Performance Liquid Chromatography
ANOVA	Analysis of variance
SE	Standard error
μM	Micromolar
LSD	Least significant differences
FAD	Flavin adenine dinucleotide
TPP	Thiamine pyrophosphate
Mg ²⁺	Magnesium
SU	Sulfonylurea
IMI	Imidazolinone
TP	Triazolopyrimidine
PTB	Pyrimidinyl(thio)benzoate
SCT	Sulfonyl-aminocarbonyl-triazolinone
HRAC	Herbicide Resistance Action Committee
WSSA	Weed Science Society of America
ng g ⁻¹	Nanogram per gram
ppt	Parts per trillion
AHAS	Acetohydroxyacid synthase

MARDI	Malaysian Agriculture Research and Development Institute
DAS	Days after sowing
IC ₅₀	Inhibitory Concentration of 50%
GST	Glutathione S-Transferase
MADA	Muda Agricultural Development Authority
KADA	Kerian Integrated Agriculture Development Authority
KETARA	North Terengganu Integrated Agriculture Development
Kg/ha	Kilogram per hectare
Ca	Calcium
Fe	Iron
PCR	Polymerase chain reaction
Kb	Kilobases
DTT	Dithiothreitol
PMSF	Phenylmethylsulphonyl fluoride
KH ₂ PO ₄ / K ₂ HPO ₄	Potassium phosphate
(NH ₄) ₂ SO ₄	Ammonium sulphate
ACCase	Acetyl coenzym-A carboxylase
EPSPS	5-enolpyruvylshikimate-3-phosphate synthase
PSII	Photosystem II
HAT	Hours after treatment
SEM	Scanning electron microscope

CHAPTER 1

INTRODUCTION

Agriculture is an important sector of Malaysia's economy. It has contributed significantly to the national Gross Domestic Product (GDP), providing employment for a good percentage of the population. Rubber and oil palm are the two main crops that have dominated the agricultural exports although the Malaysian share of the world's production of these crops has declined over the past years.

Rice is a staple food in the daily diet of Malaysians. It is a symbol of traditional Malaysian culture. Rice plays an important part in the country's agriculture although the overall production has yet to meet the country's needs. According to The National Agrofood Policy of Malaysia, 2011-2020, local rice production should be increased to ensure the country's future demand. Rice production has been facing several constraints. Crop pests including weeds, pathogens, insects, rodents as well as birds have always been acknowledged as among the most devastating factors that significantly reduce production. Among other factors include overuse of pesticides, excessive nitrogen fertilizer applications and unpredictable weather conditions.

Weeds have always been a major pest in most rice production areas, be it in Malaysia or anywhere else in the world. Weeds are generally accepted as plants growing where they are not wanted. Most of the weeds co-evolved with crops and in some cases, they were ancestors of cultivated plant species. The common weeds in rice fields are frequently legacies of previous years' crops which are seeds, rhizomes, tubers and bulb surviving in the soil. The general impact of weeds on rice fields includes increase in production costs as costs of weed control, act as secondary hosts for other insect pests, effect on harvesting and grain quality, and blockage of irrigation canals caused by invasive and persistent aquatic weed species.

To overcome weed problem in rice fields, farmers generally depend on herbicides for their control. Before the introduction and use of modern herbicides, rice farmers depended on hand weeding, ploughing or tillage, crop rotations as the main methods of weed control. However, these were not sustainable in vast fields because of high labour requirements and time-consuming issues. The introduction of herbicides for weed control has a huge relief to farmers all over the world. Among the modern herbicides, synthetic auxin, 2,4-D was developed for the control of broadleaf weeds. Following this introduction, the use of herbicides has been the most rapidly adopted strategy by farmers for the management of weeds in rice as well as in other crops production entities. Without doubt, it has become the most dependable and reliable weed control method (Heap, 2014). In Malaysian rice fields, herbicides have been extensively used for the control of broad-spectrum weed species.

There are numerous herbicide modes of action that have been used to control weeds. One of these most widely used herbicides is acetohydroxyacid synthase (AHAS) herbicides/inhibitors. Currently, there are 26 different herbicide sites of action being documented (Heap, 2017); including having various residual properties, selectivity, application times, and crop-type application suitability. Although herbicides are highly preferred by farmers because of their high efficacy in weed management, over-dependent on and overuse of a particular herbicide or groups of herbicides having similar mode of action is conceivable to result in weed species developing resistance towards the herbicides used. It has been reported that there are 485 cases of weeds resistant to herbicides with 252 weed species reported to have developed resistance to 163 herbicides in 92 crops over 69 countries (Heap, 2017). Among the herbicide groups, AHAS inhibitors top the list with the highest herbicide resistant cases of 159 weed species worldwide, including Malaysia (Heap, 2017).

Demand for herbicides for efficient weed control has always been high in Malaysian rice fields. Unfortunately, continuous dependence on herbicides has caused some weed species to develop resistance to the herbicides used. A total of 16 species have been recorded as herbicide resistant weeds, predominantly in vegetable farms, plantations, and rice fields (Heap, 2017). Of these, 9 species are weeds commonly found in rice field, namely (in the alphabetical order) *Bacopa rotundifolia* (Michx.) Wettst, *Echinochloa crus-galli* (L.) P. Beauv., *Fimbristylis quinquangularis* (Vahl) Kunth, *Leptochloa chinensis* (L.) Nees, *Limnocharis flava* (L.) Buchenau, *Limnophila erecta* Benth, *Oryza sativa* L. complex, *Sagittaria guayanensis* Kunth, and *Sphenoclea zeylanica* Gaertn. (Ruzmi et al., 2017; Dilipkumar et al., 2017). These weed species have been reported either resistant to single herbicide, or cross-resistant to several herbicides having similar mode of action, or multiple resistant to several herbicide modes of action. Among these herbicide-resistant weed species, *Limnocharis flava* has been recognized as one of the most common in Malaysian rice fields (Heap, 2017). This species was reported to have developed the resistance in 1998, and is the first weed species that evolved multiple resistance to both bensulfuron-methyl (AHAS inhibitors) and 2,4-D (synthetic auxins) (Heap, 2017). The species has become invasive in many rice fields, causing farmers to cost thousands of ringgit in terms of control and yield loss through competition, as well as blocking irrigation canals. To date, there has been no record on the resistance in terms of patterns and mechanisms for this aquatic weed species, resulting in lack of information on effective and efficient management of the species.

The present study specifically focused on the resistance evolution across AHAS inhibitors and aimed at defining the mechanisms conferring resistance to these herbicides in *L. flava* populations in Malaysian rice fields. The specific objectives were:

1. To quantify and characterize the level of resistance and cross-resistance patterns to AHAS inhibitors in a resistant *L. flava* population infesting rice fields;
2. To elucidate target-site AHAS gene mutation endowing resistance to AHAS inhibitors at the molecular and enzyme levels of the resistant *L. flava* population;
3. To identify the co-existence of possible non-target site resistance mechanism(s) from resistant *L. flava* individuals carrying no known AHAS gene mutation(s) by comparing the uptake, translocation, and metabolism of AHAS inhibitors with that of a susceptible population;
4. To determine and compare the morphological and physiological responses between susceptible and resistant *L. flava* populations in the presence and absence of AHAS inhibitor.



REFERENCES

- Abhilash, P.C., Singh, N., Sylas, V.P., Kumar, B.A., Mathew, J.C., Satheesh, R. and Thomas A.P. 2008. Eco-distribution mapping of invasive weed *Limnocharis flava* (L.) Buchenau using geographical information system: implications for containment and integrated weed management for ecosystem conservation. *Taiwania*. 53: 30-41.
- Ahmad- Hamdani, M. S., Juraimi, A.S. and Mazlan, N. Herbicide resistant weeds in Malaysian rice fields: will the weedy rice become the next candidate? In 25th Asian Pacific Weed Science Society Conference on Weed Science for Sustainable Agriculture, Environment and Biodiversity. October 13-16, 2015, Hyderabad, India. Proceeding Book (Volume II), pp 13.2015.
- Alcocer- Ruthling, M., Thill D.C. and Mallory- Smith C. 1992a. Monitoring the occurrence of sulfonylurea- resistant prickly lettuce (*Lactuca serriola*). *Weed Tech*. 6:437-440.
- Alcocer-Ruthling, M., Thill D.C. and Shafii B. 1992b. Differential competitiveness of sulfonylurea resistant and susceptible prickly lettuce (*Lactuca serriola*). *Weed Tech*. 6: 303-309.
- Anon. 2008. European Commission. Guidance document on residue analytical methods.
- Ashigh, J., and Tardif, F. J. 2007. An Ala205Val substitution in acetohydroxyacid synthase of eastern black nightshade (*Solanum Ptychanthum*) reduces sensitivity to herbicides and feedback inhibition. *Weed Sci*. 55: 558-565
- Ashigh, J., Corbett, C.A.L., Smith, P.J., Laplante, J. and Tardif, F.J. 2009. Characterization and diagnostic tests of resistance to acetohydroxyacid synthase inhibitors due to an Asp376Glu substitution in *Amaranthus powellii*. *Pestic. Biochem. Physiol*. 95: 38-46.
- Azmi, M., Muhamad, H. and Johnson D, E. 2015. Impact of weedy rice infestation on rice yield and influence of crop establishment technique. In: Proceedings of the 20th Asean-Pacific Weed Science Society Conference, Ho Chi Minh City, Vietnam. Asian Pacific Weed Science Society.
- Bakkali, Y., Ruiz-Santaella, J.P., Osuna, M.D., Wagner, J., Fischer, A. J. and De Prado, R. 2007. Late watergrass (*Echinochloa phyllopogon*): Mechanisms involved in the resistance to fenoxaprop-p-ethyl. *J. Agric. Food Chem*. 55: 4052-4058.
- BASF, 2017. <https://www.agproducts.basf.us/products/other-clearfield-crops.html>. Retrieved 8 November 2017.
- Beckie, H.J. and Tardif, F.J. 2012. Herbicide cross resistance in weeds. *Crop Prot*. 35: 15-28.

- Beckie, H.J., Warwick, S.I., Sauder, C.A., Kelln, G. M. and Lozinski, C. 2012. Acetolactate Synthase Inhibitor-Resistant False Cleavers (*Galium spurium*) in Western Canada. *Weed Technol.* 26: 151–155.
- Bernasconi, P., Woodworth A. R., Rosen B. A., Subramanian M. V., and Siehl D. L. 1995. A naturally occurring point mutation confers broad range tolerance to herbicides that target acetolactate synthase. *J. Biol. Chem.* 270: 17381-17385.
- Bhagat, R.M., Bhuiyan S.I., Moody K. and Estorninos L.E. 1999. Effect of water, tillage and herbicide on ecology of weed communities in intensive wet-seeded rice ecosystem. *Crop Prot.* 18: 293–303.
- Blair, A.M. and Martin, T.D. 1988. A review of the activity, fate, and mode of action of sulfonylurea herbicides. *Pestic. Sci.* 22:195–219.
- Bouzoubaâ, Z., Mousadik, A. El and Belahsen, Y. 2006. Variation in amounts of epicuticular wax on leaves of *Argania spinosa* (L). Skeels. *Acta Bot. Gall.* 153: 167–177.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72: 248–254.
- Brooks, S.J., Panetta, F.D. and Galway, K.E. 2008. Progress towards the eradication of mikania vine (*Mikania micrantha*) and limnocharis (*Limnocharis flava*) in northern Australia. *Invasive Plant Sc and Manag.* 1: 296-303.
- Brosnan, J.T., Vargas, J.J., Breeden, G.K., Grier, L., Aponte, R.A., Tresch, S. and Laforest, M. 2016. A new amino acid substitution (Ala-205-Phe) in acetolactate synthase (ALS) confers broad spectrum resistance to ALS-inhibiting herbicides. *Planta.* 243: 149–159.
- Brown, H.M. 1990. Mode of action , crop selectivity , and soil relations of the sulfonylurea herbicides. *Pestic. Sci.* 29: 263–281.
- Burgos, N.R., Tranel, P.J., Streibig, J.C., Davis, V.M., Shaner, D., Norsworthy, J.K. and Ritz, C. 2012. Review: Confirmation of Resistance to Herbicides and Evaluation of Resistance Levels. *Weed Sci.* 61: 4–20.
- Burks, K.C. 1993. '*Limnocharis flava*: A review of the literature'. Technical Services Section 93-02. Department of Natural Resources, Florida.
- Busi, R. and Powles, S.B. 2009. Evolution of glyphosate resistance in a *Lolium rigidum* population by glyphosate selection at sublethal doses. *Heredity (Edinb).* 103: 318–325.

- Busi, R., Vidotto, F. and Fischer, A.J. 2006. Patterns of resistance to ALS herbicides in smallflower umbrella sedge (*Cyperus Difformis*) and ricefield bulrush (*Schoenoplectus Mucronatus*). *Weed Technol.* 20: 1004–1014.
- CABI, 2013. Invasive Species Compendium. Wallingford, UK: CAB International. www.cabi.org/isc.
- CABI, 2017. Invasive Species Compendium. Wallingford, UK: CAB International. www.cabi.org/isc. Retrieved 8 November 2017.
- Calha, I.M., Osuna, M.D., Serra, C., Moreira, I., De Prado, R. and Rocha, F. 2007. Mechanism of resistance to bensulfuron-methyl in *Alisma plantago-aquatica* biotypes from Portuguese rice paddy fields. *Weed Res.* 47: 231–240.
- Chachalis, D., Reddy, K.N., Elmore, C.D. and Steele, M.L. 2001. Herbicide efficacy, leaf structure, and spray droplet contact angle among Ipomoea species and smallflower morningglory. *Weed Sci.* 49: 628–634.
- Chakraborty, S. and Newton, A.C. 2011. Climate change, plant diseases and food security: An overview. *Plant Pathol.* 60: 2–14.
- Chapin III, F. S., K. Autumn, and F. Pugnaire. 1993. Evolution of suites of traits in response to environmental stress. *Am Nat.* 142: 578-592
- Chauhan B.S. and Johnson D.E. 2010. The role of seed ecology in improving weed management strategies in the tropics. *Adv. Agron.* 105, 221–262.
- Chipman, D., Barak, Z. and Schloss, J. V. 1998. Biosynthesis of 2-aceto-2-hydroxy acids: acetolactate synthases and acetohydroxyacid synthases. *Biochim. Biophys. Acta - Protein Struct. Mol. Enzymol.* 1385: 401–419.
- Coley, P. D., J. P. Bryant, and F. S. Chapin. 1985. Resource availability and plant antiherbivore defense. *Science.* 230: 895-899.
- Cobb, A.H. and Reade, J.P.H. 2010. *Herbicides and Plant Physiology*. 2nd Edition, Wiley-Blackwell, Hoboken, 1-296. <http://dx.doi.org/10.1002/9781444327793>.
- Cruz-Hipolito, H., Domínguez-Valenzuela, J. A., Osuna, M.D. and De Prado, R. 2012. Resistance mechanism to acetyl coenzyme A carboxylase inhibiting herbicides in *Phalaris paradoxa* collected in Mexican wheat fields. *Plant Soil.* 355: 121–130.
- Dayan, F.E., Owens, D.K., Corniani, N., Silva, F.M.L., Watson, S.B., Howell, J. and Shaner, D.L. 2015. Biochemical markers and enzyme assays for herbicide mode of action and resistance studies. *Weed Sci.* 63: 23–63.

- Dal Magro, T., Schaedler, C.E., Fontana, L.C, Agostinetto, D., Vargas, L. 2011. Competitive ability between biotypes of *Cyperus difformis* L. resistant or susceptible to ALS-inhibitor herbicide and those with flooded rice. *Bragantia*. 70: 294 – 301.
- Department of Agriculture, Malaysia. 2013. Overall average yield of paddy by state, Malaysia, 2004–2013. Department of Agriculture, Malaysia, Putrajaya, Malaysia.
- De Prado, R., Osuna, M.D. and Fischer, A.J. 2004. Resistance to ACCase inhibitor herbicides in a green foxtail (*Setaria viridis*) biotype in Europe. *Weed Sci.* 52: 506–512.
- Délye, C. 2012. Unravelling the genetic bases of non-target-site-based resistance (NTSR) to herbicides: A major challenge for weed science in the forthcoming decade. *Pest Manag. Sci.* 69: 176–187
- Délye, C., Jasieniuk, M. and Le Corre, V. 2013. Deciphering the evolution of herbicide resistance in weeds. *Trends Genet.* 29: 649–658.
- Délye, C., Pernin, F. and Scarabel, L., 2011. Evolution and diversity of the mechanisms endowing resistance to herbicides inhibiting acetolactate-synthase (ALS) in corn poppy (*Papaver rhoeas* L.). *Plant Sci.* 180: 333–342.
- Dilipkumar, M., Zainal Abidin, Z., Varghese, Geoge., Burgos, N.R., Chuah T.S. 2017. Proceeding of the 26th Asian- Pacific Weed Science Society Conference, Kyoto, Japan, September 19-22, 2017.
- DOSM. 2015. Statistik kawasan pembangunan pertanian bersepadu. Retrieved from https://www.dosm.gov.my/v1/uploads/files/3_Time%20Series/Malaysia_Time_Series_2015/09Padi.pdf
- Duggleby, R.G., Mccourt, J.A. and Guddat, L.W. 2008. Structure and mechanism of inhibition of plant acetohydroxyacid synthase. *Plant Physiol. Biochem.* 46: 309–324.
- Duggleby, R.G. and Pang, S.S. 2000. Acetohydroxyacid Synthase. *J. Biochem. Mol. Biol.* 33: 1–36.
- Durán-Prado, M., Osuna, M.D., Prado, R. De, Franco, A.R., 2004. Molecular basis of resistance to sulfonylureas in *Papaver rhoeas*. *Pestic. Biochem. Physiol.* 79: 10–17.
- Edwards R. and Dixon DP. 2000. The role of glutathione transferases in herbicides metabolism. In AH Cobb, RC Kirkwood eds, *Herbicides and Their Mechanisms of Action*. Sheffield Academic Press, Sheffield, UK, Pp 38–71

- FAO (Food and Agriculture Organization of the United Nations). 2002. Fao specifications and evaluations for plant protection products. Nations, Food Agric. Organ. United.
- FAO (Food and Agriculture Organization of the United Nations). 2013. Food and Agricultural Commodities Production/Commodities by Country. http://faostat3.fao.org/browse/rankings/commodities_by_country/E
- Fischer, A.J., Bayer, D.E., Carriere, M.D., Ateh, C.M. and Yim, K.O. 2000. Mechanisms of resistance to bispyribac-sodium in an *Echinochloa phyllopogon* accession. *Pestic. Biochem. Physiol.* 68: 156–165.
- Foes, M.J., Liu, L., Vigue, G., Stoller, E.W., Wax, L.M. and Tranel, J. 1999. Weed Science Society of America A kochia (*Kochia scorparia*) biotype resistant to triazine and ALS-inhibiting herbicides. *Weed Sci.* 47: 20–27.
- Gerwick, B. C., Subramanian, M. V. and Loney-Gallant, V. I. 1990. Mechanism of action of the 1,2,4-triazolo[1,5-a] pyrimidines. *Pest Manag. Sci.* 29:357–364.
- Gill, G. S., R. D. Cousens, and M. R. Allan. 1996. Germination, growth and development of herbicide resistant and susceptible populations of rigid ryegrass (*Lolium rigidum*). *Weed Sci.* 44: 252-256
- GISD (Global Invasive Species Database). 2017. Species profile: *Limnocharis flava*. Downloaded from <http://www.iucngisd.org/gisd/species.php?sc=620> on 26-07-2017.
- Gnanamanickam, S. S. 2009. Biological control of bacterial blight of rice, In Biological control of rice diseases. Springer pp. 67-78.
- Grabherr, M., Haas, B., Yassour M, et al. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* 29: 644–652.
- Gressel, J. 2009. Evolving understanding of the evolution of herbicide resistance. *Pest Manag. Sci.* 65: 1164–1173.
- Guttieri, M.J., Eberlein, C.V. and Thill, D. C. 1995. Diverse mutations in the acetolactate synthase gene confer chlorsulfuron resistance in kochia (*Kochia scoparia*). *Weed Sci.* 43:175-178.
- Guttieri, M.J., Eberlein, C. V., Mallory-Smith, C. A., Thill, D.C. and Hoffman, D.L. 1992. DNA Sequence Variation in Domain A of the Acetolactate Synthase Genes of Herbicide-Resistant and -Susceptible Weed Biotypes. *Weed Sci.* 40: 670–677.

- Hakim, M.A., Juraimi, A.S., Ismail, M.R., Hanafi M.M., Selamat, A. 2013. A survey on weed diversity in coastal rice fields of sebarang Perak in Peninsular Malaysia. *J. Anim. Plant Sci.* 23(2): 534-542.
- Halbert, S.E. 1997. Entomology section. Triology 36/1. Florida Department of Agriculture and Consumer Services, Division of Plant Industry. <http://www.freshfromflorida.com/Divisions-Offices/Plant-Industry/Plant-Industry-Publications/Tri-ology-FDACS-DPI>.
- Han, H., Yu, Q., Purba, E., Li, M., Walsh, M., Friesen, S. and Powles, S.B. 2012. A novel amino acid substitution Ala-122-Tyr in ALS confers high-level and broad resistance across ALS-inhibiting herbicides. *Pest Manag. Sci.* 68: 1164–1170.
- Hare, C.J., Chong, W.C., Ooi, G.T., Bhandhufalck, A., Nawsaran, S. and Chanprasit, P. 1989. Sofit Super: broad spectrum weed management for wet sown rice in S.E. Asia. Proceedings of the 12th Asian-Pacific Weed Science Society Conference, Asian-Pacific, eds J.O. Guh, J.Y. Pyon, S.C. Kim, G.P. Chung, S.L. Kwon and J.C. Kim. pp. 165-70. Weed Science Society, Seoul.
- Harker K.N., O', Donovan J.T., Blackshaw R.E., Beckie H.J., Mallory-Smith C., Maxwell B.D. 2012: Our view. *Weed Sci.* 60: 143 – 144.
- Harvey, J., Jr., Dulka, J. and Anderson, J. 1985. Properties of sulfometuron methyl affecting its environmental fate: aqueous hydrolysis and photolysis, mobility and adsorption on soils, and bioaccumulation potential. *J. Agric. Food Chem.* 33:590–596.
- Hatami, Z.M., Gherekhloo, J., Rojano-Delgado, A.M., Osuna, M.D., Alcántara, R., Fernández, P., Sadeghipour, H.R. and De Prado, R. 2016. Multiple Mechanisms Increase Levels of Resistance in *Rapistrum rugosum* to ALS Herbicides. *Front. Plant Sci.* 7: 1–13.
- Heap, I. 2017. The International Survey of Herbicide Resistant Weeds. URL www.weedscience.org. Retrieved 19 January 2017.
- Heap, I. 2016. The International Survey of Herbicide Resistant Weeds. URL www.weedscience.org. Retrieved 30 May 2016.
- Heap, I. 2014. Herbicide resistant weeds. In: Pimentel D., Peshin R. (eds). Integrated Pest Management. New York- Heidelberg-Dordrecht-London, Springer Science: 283 – 300.
- Heap I. and Knight R. 1982. A population of ryegrass tolerant to the herbicide diclofop-methyl. *Aust. J. Agri. Res.* 48: 156 – 157.

- Heap I., Knight R. 1986. The occurrence of herbicide cross resistance in a population of annual ryegrass, *Lolium rigidum*, resistant to diclofop-methyl. *Aust. J. Agri. Res.* 37: 149 – 156.
- Herms, D. A., and W. J. Mattson. 1994. Plant-growth and defense. *Trends Ecol. Evol.* 9: 488.
- Heil, M. 2002. Ecological costs of induced resistance. *Curr. Opin. Plant Biol.* 5: 345–350.
- Hwang, I.T., Lee, K.H., Park, S.H., Lee, B.H., Hong, K.S., Han, S.S. and Cho, K.Y. 2001. Resistance to acetolactate synthase inhibitors in a biotype of *Monochoria vaginalis* discovered in Korea. *Pestic. Biochem. Physiol.* 71: 69–76.
- Ingold, C.T. 1995. An unusual smut fungus: *Doassansiopsis limnocharidis*. *Mycologist.* 9: 58-9.
- Iwakami, S., Watanabe, H., Miura, T., Matsumoto, H. and Uchino, A. 2014. Occurrence of sulfonyleurea resistance in *Sagittaria trifolia*, a basal monocot species, based on target-site and non-target-site resistance. *Weed Biol. Manag.* 14: 43–49.
- Jablonkai, I. 2011. Molecular mechanism of action of herbicides. *4 Herbicides – Mechanisms and Mode of Action*, 3–24.
- Juraimi, A.S., Begum, M., Mohd Yusof, M.N. and Man, A. 2010. Efficacy of herbicides on the control weeds and productivity of direct seeded rice under minimal water conditions. *Plant Prot. Q.* 25: 19–25.
- Juraimi, A. S., Ahmad-Hamdani, M. S., Anuar, A. R., Azmi, M., Anwar, M. P. and Uddin, M. K. 2012a. Effect of water regimes on germination of weed seeds in a Malaysian rice field. *Aust. J. of Crop Sci.* 6(4): 598–605.
- Juraimi, A.S., Begum, M., Parvez Anwar, M., Shari, E.S., Sahid, I. and Man, A. 2012b. Controlling resistant *Limnocharis flava* (L.) Buchenau biotype through herbicide mixture. *J. Food, Agric. Environ.* 10: 1344–1348.
- Kanehisa, M., Araki, M, Goto, S, et al. 2008. KEGG for linking genomes to life and the environment. *Nucleic Acids Res.* 36: D480–D484.
- Karim, R.S.M., Man, A.B. and Sahid, I.B., 2004. REVIEW PAPER: Weed problems and their management in rice fields of Malaysia: An overview. *Weed Biol. Manag.* 4; 177–186.
- Karthigeyan, K., Sumathi, R., Jayanthi, J., Diwakar, P.G. and Lakra, G.S. (2004). *Limnocharis flava* (L.) Buchenau (Alismataceae) –a little known and troublesome weed in Andaman Islands. *Curr Sci.* 87: 140-1.

- Katanyukul, W. and Wirojchewan, T. 1989. Weed control in direct seeded rice with fenoxaprop-P-ethyl. Proceedings of the 12th Asian Pacific Weed Science Society Conference, eds J.O. Guh, J.Y. Pyon, S.C. Kim, G.P. Chung, S.L. Kwon and J.C. Kim. pp. 649-56. Asian-Pacific Weed Science Society, Seoul.
- Kiessling, U., Menck, B.H. and Unglaub, W. 1981. New experiences with bentazon and bentazon combinations for weed control in rice. Proceedings of the 8th Asian-Pacific Weed Science Society Conference, ed. B.V. Venkata Rao, pp. 99-102. Asian-Pacific Weed Science Society, Bangalore.
- Knezevic, S.Z., Streibig, J.C. and Ritz, C. 2007. Utilizing R Software Package for Dose-Response Studies: The Concept and Data Analysis. *Weed Technol.* 21: 840–848.
- Kotalawala, J. 1976. Noxious water vegetation in Sri Lanka: the extent and impact of existing infestations. In 'Aquatic weeds in S.E. Asia', eds C.K. Varshney and J. Rzoska, Dr W. Junk B.V. Publishers. Pp. 51-8.
- Kuk, Y.I., Jung, H. Il, Kwon, O. Do, Lee, D.J., Burgos, N.R. and Guh, J.O. 2003a. Sulfonylurea herbicide-resistant *Monochoria vaginalis* in Korean rice culture. *Pest Manag. Sci.* 59: 949–961.
- Kuk, Y.I., Kim, K.H., Kwon, O. Do, Lee, D.J., Burgos, N.R., Jung, S. and Guh, J.O., 2003b. Cross-resistance pattern and alternative herbicides for *Cyperus difformis* resistant to sulfonylurea herbicides in Korea. *Pest Manag. Sci.* 60: 85–94.
- Lamego, F.P., Charlson, D., Delatorre, C. A., Burgos, N.R. and Vidal, R. A. 2009. Molecular basis of resistance to ALS-Inhibitor Herbicides in greater beggarticks. *Weed Sci.* 57: 474–481.
- Le, D.T., Yoon, M.Y., Kim, Y.T. and Choi, J. D. 2005. Roles of three well-conserved arginine residues in mediating the catalytic activity of tobacco acetohydroxy acid synthase. *J. Biochem.* 138, 35–40.
- Légère, A., Stevenson, F.C., Beckie, H.J., Warwick, S.I., Johnson, E.N., Hrynewich, B. and Lozinski, C. 2013. Growth Characterization of Kochia (*Kochia scoparia*) with Substitutions at Pro 197 or Trp 574 Conferring Resistance to Acetolactate Synthase–Inhibiting Herbicides. *Weed Sci.* 61, 267–276.
- Li, B., Dewey, C. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics.*
- Li, J., Chen, W., Xu, Y. and Wu, X. 2016. Comparative effects of different types of tank-mixed adjuvants on the efficacy, absorption and translocation of cyhalofop-butyl in barnyardgrass (*Echinochloa crus-galli* [L.] Beauv.). *Weed Biol. Manag.* 16: 80–89.

- Li, M., Yu, Q., Han, H., Vila-Aiub, M. and Powles, S.B. 2013. ALS herbicide resistance mutations in *Raphanus raphanistrum*: Evaluation of pleiotropic effects on vegetative growth and ALS activity. *Pest Manag. Sci.* 69: 689–695.
- Liu, W., Bi, Y., Li, L., Yuan, G. and Wang, J. 2013. Molecular basis of resistance to tribenuron in water starwort (*Myosoton aquaticum*) populations from China. *Weed Sci.* 61: 390–395.
- Lu, J., Brigham, C., Plassmeier, J. and Sinskey, A. 2014. Characterization and modification of enzymes in the 2-ketoisovalerate biosynthesis pathway of *Ralstonia eutropha* H16. *Appl. microbiol. and biotechnol.* 99. 10.1007/s00253-014-5965-3.
- Maclean, J.L., Dawe, D.C., Hardy, B. and Hettel, G.P. 2002. Rice Almanac (3rd Edition). Los Banos, Philippines: IRRI.
- Maddison, P.A. 1993. Pests and other fauna associated with plants, with botanical accounts of plants UNDP/ FAO-SPEC Survey of agricultural pests and diseases in the South Pacific, Technical report. Volume 3. Manaaki Whenua Landcare Research, Auckland.
- Mahnken, G.E. and Weber, J.B. 1988. Capillary movement of triasulfuron and chlorsulfuron in Rion sandy loam soil. *Proc. Southern Weed Sci. Soc.* 68:332–336.
- Mallory-Smith, C.A and Retzinger, J. 2003. Revised Classification of Herbicides by Site of Action for Weed Resistance. *Weed Technol.* 17: 605–619.
- Mallory-Smith, C.A., Thill, D.C., Dial, M.J. and Zemetra, R.S. 1990. Inheritance of sulfonylurea herbicide resistance in *Lactuca* spp. *Weed Tech.* 4: 787–790
- Mao, X., Cai, T. and Olyarchuk, J.G., 2005. Automated genome annotation and pathway identification using the KEGG Orthology (KO) as a controlled vocabulary. *BMC Bioinformatics.* 21: 3787–3793.
- Maricle, B.R., Koteyeva, N.K., Voznesenskaya, E. V., Thomasson, J.R. and Edwards, G.E. 2009. Diversity in leaf anatomy, and stomatal distribution and conductance, between salt marsh and freshwater species in the C4 genus *Spartina* (Poaceae). *New Phytol.* 184: 216–233.
- Massa, D. 2011. Investigations on herbicide resistance in *Apera spica-venti* populations 87.
- McCourt, J.A., Pang, S.S., King-Scott, J., Guddat, L.W. and Duggleby, R.G. 2006. Herbicide-binding sites revealed in the structure of plant acetohydroxyacid synthase. *Proc. Natl. Acad. Sci.* 103: 569–573.

- McCourt, J. A. and Duggleby, R.G. 2006. Acetohydroxyacid synthase and its role in the biosynthetic pathway for branched-chain amino acids. *Amino Acids*. 31: 173–210.
- Menne, H. and Kocher, H. 2007. HRAC classification of herbicides and resistance development. In: *Modern Crop Protection Compounds*, W. Kramer and U. Schirmer, Ed., Wiley-VCH Publishing, Weinheim, Germany. Pp: 5-26.
- Merotto, A., Jasieniuk, M., Osuna, M.D., Vidotto, F., Ferrero, A. and Fischer, A.J. 2009. Cross-resistance to herbicides of five ALS-inhibiting groups and sequencing of the ALS gene in *Cyperus difformis* L. *J. Agric. Food Chem.* 57: 1389–1398.
- Miller, J.W. 1997. Plant Pathology section. Tri-ology 36/1. Florida Department of Agriculture and Consumer Services, Division of Plant Industry. <http://www.freshfromflorida.com/Divisions-Offices/Plant-Industry/Plant-Industry-Publications/Tri-ology-FDACS-DPI>.
- Moyer, J. R., Esau, R., and Kozub, G.C. 1990. Chlorsulfuron persistence and response of nine rotational crops in alkaline soils of southern Alberta. *Weed Technol.* 4:543–548
- Oerke, E.C. 2002. Crop losses due to pests in major crops. In *CAB International Crop Protection Compendium 2002. Economic Impact*.
- Orr, H.A. 2010. Fitness and its role in evolutionary genetics. *Nat Rev Genet.* 10: 531–539.
- Osuna, M.D., Vidotto, F., Fischer, A.J., Bayer, D.E., De Prado, R. and Ferrero, A. 2002. Cross-resistance to bispyribac-sodium and bensulfuron-methyl in *Echinochloa phyllopogon* and *Cyperus difformis*. *Pestic. Biochem. Physiol.* 73: 9–17.
- Owen, M.J., Goggin, D.E. and Powles, S.B. 2012. Non-target-site-based resistance to ALS-inhibiting herbicides in six *Bromus rigidus* populations from Western Australian cropping fields. *Pest Manag. Sci.* 68: 1077–1082.
- Pang, S. S., Guddat, L. W. and Duggleby, R. G. 2004. Crystallization of *Arabidopsis thaliana* acetohydroxyacid synthase in complex with the sulfonylurea herbicide chlorimuron ethyl. *Acta Crystallographica Section D: Biological Crystallography.* 60 (1): 153–155.
- Park, K.W., Fandrich, L., Mallory-Smith, C. A. 2004. Absorption, translocation, and metabolism of propoxycarbazone-sodium in ALS-inhibitor resistant *Bromus tectorum* biotypes. *Pestic. Biochem. Physiol.* 79:18–24.

- Parks, K. W., C. A. Mallory-Smith, D. A. Ball and G. W. Mueller-Warrant. 2004. Ecological fitness of acetolactate synthase inhibitor-resistant and -susceptible downy brome (*Bromus tectorum*) biotypes. *Weed Sci.* 52: 768-773.
- Peterson, D.E., Thompson, C.R., Shoup, D.E. and Olson, B.L. 2013. Herbicide Mode of Action. Kansas State Univ. 1–25.
- Pons, T.L. 1982. Factors affecting weed seed germination and seedling growth in lowland rice in Indonesia. *Weed Res.* 22: 155-61.
- Powles, S.B. and Yu, Q. 2010. Evolution in action: plants resistant to herbicides., *Ann Rev Plant Biol.* 61:317-347.
- Praveena, R. and Naseema, A. 2003. Effects of two potent biocontrol agents on water hyacinth. *Int Rice Res Notes.* 28: 40.
- Preston, C., Stone, L.M., Rieger, M.A. and Baker, J. 2006. Multiple effects of a naturally occurring proline to threonine substitution within acetolactate synthase in two herbicide-resistant populations of *Lactuca serriola*. *Pestic. Biochem. Physiol.* 84: 227–235.
- Purrington, C. B. 2000. Costs of resistance. *Curr Opin Plant Biol.* 3: 305-308.
- Rabbani N., Bajwa R. and Javaid A. 2011. Interference of five problematic weed species with rice growth and yield. *Afr. J. Biotechnol.* 10, 1854–1862.
- Rabu, M., Shah, M.D. 2013. Food and livelihood security of the Malaysian paddy farmers. *Econ. Technol. Manag. Rev.* 8, 59–69.
- Riar, D.S., Tehranchian, P., Norsworthy, J., Nandula, V., McElroy, S., Srivastava, V., Chen, S., Bond, J. and Scott, R.C. 2015. Acetolactate synthase-inhibiting herbicide-resistant rice flatsedge (*Cyperus iria*): cross resistance and molecular mechanism of resistance. *Weed Sci.* 63, 748–757.
- Robert, F., Ceccacci, Irde., Cooke, Sanjiva., Constantine, Mark. and Moses, Gene. 2013. *Implementing agriculture for development : World Bank Group agriculture action plan (2013-2015)*. Washington DC; World Bank. <http://documents.worldbank.org/curated/en/331761468152719470/Implementing-agriculture-for-development-World-Bank-Group-agriculture-action-plan-2013-2015>.
- Rost, T.L. and Reynolds, T. 1985. Reversal of chlorsulfuron-induced inhibition of mitotic entry by isoleucine and valine. *Plant Physiol.* 77:481–482.
- Ruzmi, R., Ahmad-Hamdani, M.S. and Bakar, B. 2017. Prevalence of herbicide-resistant weed species in Malaysian rice fields : A review. *Weed Biol. Manag.* 1–14.

- Saari, L. L., Cotterman, J. C. and Thill, D. C. 1994. Resistance to acetolactate synthase inhibiting herbicides. In *Herbicide Resistance in Plants: Biology and Biochemistry*, (S. B. Powles and J. A. M. Holtum, Eds.). Lewis Publishers, Boca Raton, FL, Pp. 83-139.
- Saari, L.L., Cotterman, J.C. and Primiani, M.M. 1990. Mechanism of Sulfonylurea Herbicide Resistance in the Broadleaf Weed, *Kochia scoparia*. *Plant Physiol.* 93: 55–61.
- Sada, Y., Ikeda, H., Kizawa, S. 2013a. Resistance levels of sulfonylurea-resistant *Schoenoplectus juncooides* (Roxb.) Palla with various Pro197 mutations in acetolactate synthase to imazosulfuron, bensulfuron-methyl, metsulfuron-methyl and imazaquin-ammonium. *Weed Biol. Manag.* 13: 53–61.
- Sada, Y., Ikeda, H. and Kizawa, S. 2013b. Rapid diagnosis of sulfonylurea-resistant *Schoenoplectus juncooides* [Roxb.] Palla using polymerase chain reaction-restriction fragment length polymorphism and isogene-specific direct sequencing. *Weed Biol. Manag.* 13: 1–9.
- Sada, Y., Ikeda, H., Yamato, S. and Kizawa, S. 2013c. Characterization of sulfonylurea-resistant *Schoenoplectus juncooides* having a target-site Asp376Glu mutation in the acetolactate synthase. *Pestic. Biochem. Physiol.* 107: 106–111.
- Santel, H.J., Bowden, B.A., Sorensen, V.M., Mueller, K.H. and Reynolds, J. 1999. Flucarbazone-sodium: a new herbicide for grass control in wheat. *Weed Sci. Soc. Am. Abstr* 39:7.
- Schaedler, C.E., Burgos, N.R., Noldin, J.A., Alcober, E.A., Salas, R.A. and Agostinetto, D. 2015. Competitive ability of ALS-inhibitor herbicide-resistant *Fimbristylis miliacea*. *Weed Res.* 55: 482–492.
- Shaner, D.L., Anderson, P.C., Stidham, M.A. 1984. Imidazolinones, Potent inhibitors of acetohydroxyacid synthase. *Plant Physiol.* 76: 545.
- Shaner, D.L. 1999. Resistance to Acetolactate Synthase (ALS) Inhibitors in the United States: History, Occurrence, Detection, and Management. *J. Weed Sci. Technol.* 44: 405–411.
- Shaner, D. L. and Singh, B. K. 1997. Acetohydroxyacid synthase inhibitor. Roe, Burton, and Kuhr (eds.) *Herbicide activity: Toxicology, biochemistry and molecular biology*. IOS Press, Amsterdam Netherlands. Pp:69-110.
- Shaner, D.L. and Reider, M.L. 1986. Physiological responses of corn (*Zea mays*) to AC 243,997 in combination with valine, leucine, and isoleucine. 257: 248–257.

- Shaner, D.L. and Singh, B.K. 1993. Phytotoxicity of Acetohydroxyacid Synthase Inhibitors is Not Due to Accumulation of 2-Ketobutyrate and/or 2-Aminobutyrate. *Plant Physiol.* 103: 1221–1226.
- Shannon, P., Markiel, A. and Ozier, O. 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* 13: 2498–2504.
- Sibony, M., Michel, A., Haas, H.U., Rubin, B. and Hurle, K. 2001. Sulfometuron-resistant *Amaranthus retroflexus*: Cross-resistance and molecular basis for resistance to acetolactate synthase-inhibiting herbicides. *Weed Res.* 41: 509–522.
- Stant, M.Y. 1987. Anatomy of the Butomaceae. *J. Linn. Soc.* 60: 31–60.
- Strauss, S. Y., J. A. Rudgers, and R. E. Irwin. 2002. Direct and ecological costs of resistance to herbivory. *Trends. Ecol. Evol.* 17: 278-285
- Tanaka, Y., 2003. Properties of acetolactate synthase from sulfonylurea-resistant *Scirpus juncooides* Roxb. var. *ohwianus* T. Koyama. *Pestic. Biochem. Physiol.* 77: 147–153.
- Tardif, F.J., Rajcan, I. and Costea, M. 2006. A mutation in the herbicide target site acetohydroxyacid synthase produces morphological and structural alterations and reduces fitness in *Amaranthus powellii*. *New Phytol.* 169 :251–264.
- Tranel, P., Wright, T. and Heap, I. 2017. Mutations in herbicide-resistant weeds to ALS inhibitors. <http://www.weedscience.com>.
- Tranel, P.J. and Wright, T.R. 2002. Resistance of weeds to ALS-inhibiting herbicides: what have we learned? *Weed Sci.* 50: 700–712.
- The National Agrofood Policy of Malaysia, 2011-2020. Retrived from <https://opendevelopmentcambodia.net/tag/national-agro-food-policy-2011-2020>.
- Turner, S.A. 1987. Post-applications movement of sulfometuron methyl from treated rights-of-way areas via wind soil erosion. Proceedings from the Symposium on Environmental Concerns in Rights-of-Way Management. 4: 575–579.
- Usui, K. 2001. Metabolism and selectivity of rice herbicides in plants. *Weed Biol. Manag.* 1: 137–146.
- Vanky, K. 1992. Taxonomical studies on Ustilaginales. IX. *Mycotaxon.* 43: 417-25.
- Veldhuis, L.J., Hall, L.M., O'Donovan, J.T., Dyer, W. and Hall, J.C. 2000. Metabolism-based resistance of a wild mustard (*Sinapis arvensis* L.) biotype to ethametsulfuron-methyl. *J. Agric. Food Chem.* 48: 2986–2990.

- Vila-Aiub, M.M., Neve, P. and Powles, S.B. 2009. Fitness costs associated with evolved herbicide resistance genes in plants. *New Phytol.* 184:751–67.
- Warwick, S.I., Xu, R., Sauder, C. and Beckie, H.J. 2008. Acetolactate synthase target-site mutations and single nucleotide polymorphism genotyping in ALS-resistant Kochia (*Kochia scoparia*). *Weed Sci.* 56: 797–806.
- Waterhouse, B. M. and Mitchell, A. A. 1998. Northern Australia Quarantine Strategy Weeds Target List. AQIS Miscellaneous Publication No. 6/98:110pp
- Weber, J.M. and Brooks, S.J. 2013. The biology of Australian weeds 62. *Limncharis flava*(L.) Buchenau. *Plant Prot. Q.* 28: 101–113.
- Wei, S., Li, P., Ji, M., Dong, Q. and Wang, H. 2015. Target-site resistance to bensulfuron-methyl in *Sagittaria trifolia* L. populations. *Pestic. Biochem. Physiol.* 1–5.
- Westhoven, A. M., G. R. Kruger, C. K. Gerber, J. M. Stachler, M. M. Loux and W. G. Johnson. 2008. Characterization of selected common lambsquarters (*Chenopodium album*) biotypes with tolerance to glyphosate. *Weed Sci.* 56: 685-691.
- Whaley, C.M., Wilson, H.P. and Westwood, J.H. 2007. A new mutation in plant ALS confers resistance to five classes of ALS Inhibiting herbicides. *Weed Sci.* 55: 83–90.
- Whitcomb, C.E. 1999. An introduction to ALS-inhibiting herbicides. *Toxicol. Ind. Health.*15: 232–240.
- White, A.D. 2002. Characterization of acetolactate synthase resistance in common sunflower (*Helianthus annuus*). *Retrospective Theses and Dissertations.* 407. <https://lib.dr.iastate.edu/rtd/407>.
- Yang, Q., Deng, W., Li, X., Yu, Q., Bai, L. and Zheng, M. 2016. Target-site and non-target-site based resistance to the herbicide tribenuron-methyl in flixweed (*Descurainia sophia* L.). *BMC Genomics.* 17: 551–564.
- Young, M., Wakefield, M.J. and Smyth, G.K. 2010. Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome Biol.* doi:10.1186/gb-2010-11-2-r14.
- Yu, Q., Friesen, L.J.S., Zhang, X.Q. and Powles, S.B. 2004. Tolerance to acetolactate synthase and acetyl-coenzyme A carboxylase inhibiting herbicides in *Vulpia bromoides* is conferred by two co-existing resistance mechanisms. *Pestic. Biochem. Physiol.* 78: 21–30.

- Yu, Q., Han, H., Li, M., Purba, E., Walsh, M.J. and Powles, S.B. 2012. Resistance evaluation for herbicide resistance-endowing acetolactate synthase (ALS) gene mutations using *Raphanus raphanistrum* populations homozygous for specific ALS mutations. *Weed Res.* 52: 178–186.
- Yu, Q., Han, H., Vila-Aiub, M.M. and Powles, S.B. 2010. AHAS herbicide resistance endowing mutations: Effect on AHAS functionality and plant growth. *J. Exp. Bot.* 61: 3925–3934.
- Yu, Q. and Powles, S. 2014. Metabolism-based herbicide resistance and cross-resistance in crop weeds: A threat to herbicide sustainability and global crop production. *Plant Physiol.* 166: 1106–1118.
- Yu, Q. and Powles, S.B. 2014. Resistance to AHAS inhibitor herbicides: Current understanding. *Pest Manag. Sci.* 70, 1340–1350.
- Yu, Q., Zhang, X.Q., Hashem, A., Walsh, M.J. and Powles, S.B. 2003. ALS gene proline (197) mutations confer ALS herbicide resistance in eight separated wild radish (*Raphanus raphanistrum*) populations. *Weed Sci.* 51: 831–838.
- Zakaria, N., Ahmad- Hamdani and M., Juraimi, A.S. 2018. Patterns of resistance to AHAS inhibitors in *Limnocharis flava* from Malaysia. *Plant Prot. Sci.* 1:48-59. doi: 10.17221/131/2016-PPS.
- Zhang, J., Jaeck, O., Menegat, A., Zhang, Z., Gerhards, R. and Ni, H. 2013. The Mechanism of Methylated Seed Oil on Enhancing Biological Efficacy of Topramezone on Weeds. *PLoS One.* 8: 2–10.
- Zheng, D., Kruger, G.R., Singh, S., Davis, V.M., Tranel, P.J., Weller, S.C. and Johnson W.G. 2011. Cross-resistance of horseweed (*Conyza canadensis*) populations with three different ALS mutations. *Pest Manag. Sci.* 67: 1486–1492.
- Zhou, Q., Liu, W., Zhang, Y. and Liu, K.K. 2007. Action mechanisms of acetolactate synthase-inhibiting herbicides. *Pestic. Biochem. Physiol.* 89: 89–96.