



UNIVERSITI PUTRA MALAYSIA

***DIFFERENTIAL POTASSIUM UPTAKE AND UTILIZATION EFFICIENCY
OF OIL PALM (*Elaeis guineensis* Jacq.) COMMERCIAL CULTIVARS***

SIM CHOON CHEAK

FP 2018 95



**DIFFERENTIAL POTASSIUM UPTAKE AND UTILIZATION EFFICIENCY
OF OIL PALM (*Elaeis guineensis* Jacq.) COMMERCIAL CULTIVARS**

By

SIM CHOON CHEAK

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of
Doctor of Philosophy**

October 2017

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia.



Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Degree of Doctor of Philosophy.

**DIFFERENTIAL POTASSIUM UPTAKE AND UTILIZATION EFFICIENCY
OF OIL PALM (*Elaeis guineensis* Jacq.) COMMERCIAL CULTIVARS**

By

SIM CHOON CHEAK

October 2017

Chairman : Professor Zaharah Abdul Rahman, PhD
Faculty : Agriculture

Conventional approaches of using higher fertilizer inputs to sustain profitable yields in oil palm plantations can be uneconomical and produce inconsistent results. In addition, the biological potential of attaining much higher oil yield is often limited by marginal environments. Nutrient efficient genotypes could potentially lead to higher productivity when grown on marginal land and eventually improve the sustainable use of resources and production of palm oil. Studies on interaction effects between planting material and nutrient inputs show differential uptake and utilization efficiency between the commercially available oil palm planting materials. The differences in leaf nutrient contents between genotypes and yield response to K fertilizer inputs demonstrated the presence of more efficient uptake characteristics. If such potassium efficient cultivars could be widely adopted, the industry would not only be capable of saving resources but also to increase productivity as well. Potassium use in palm oil production ranges from approximately 13 to 21 kg of palm oil per kg of potassium with varying degree of efficiency depending on planting varieties. The potassium use efficiency could potentially increase by 50 % in the most potassium efficient cultivar. The objectives of this study were (1.) to evaluate the growth response of selected oil palm crosses under K deficient environment and (2.) to estimate potassium use efficiency of different oil palm genotypes as part of the effort to elucidate the physiological mechanism potassium-efficient oil palms. Phenotypic responses of 5 oil palm genotypes with genetic origin from Deli and Nigerian Dura interbred with AVROS, Nigerian and Yangambi Pisiferas grown under deficient and adequate potassium supplies were evaluated. Potassium-efficient genotypes were differentiated in this experiment, where the potassium-efficient genotypes produced higher biomass (by 37.3 %) and had higher potassium uptake activity (by 41.7 %). The efficient genotypes were capable of extracting higher amount of soil potassium (by 95 %) under deficient potassium supplies. The K-efficient genotype was capable of sustain growth and to adapt to potassium-deficient environments. Alterations in rooting behaviour (increasing fine root proliferation) and maintenance of shoot growth (frond production rate) are the primary physical traits of adaptation to

potassium-deficient environment. The ability to remobilize the limiting nutrients from sink tissues to source tissues i.e. from the bole and rachis to the pinnae (the photosynthetically active tissues) and roots (to search for more nutrients allows the plant to further acquire more resources to ensure continuous growth) is also a key trait. Comparative analysis of transcriptomic differences between the efficient and in-efficient genotypes showed significant upregulation of potassium transporters (KUP3, KUP8 and KUP11) in the roots of the K-efficient genotype and genes which confer tolerance to stress, minimizes cellular damage, stress regulation and potassium homeostasis. Traits for potassium efficiency are conferred by the interaction of multiple complex mechanisms, governed by pool of genes controlling the physiological processes of stress regulation, cellular development and metabolite homeostasis. Stress detection and regulating cellular processes to mitigate the effect of stress could be the key in first tolerating and reducing damages to cellular and consequently enhancing the genotype's ability to adapt, absorb and utilize nutrients more effectively.

Abstrak tesis yang telah dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah.

**PERBEZAAN DALAM PENGAMBILAN DAN PENGGUNAAN KALIUM
OLEH KULTIVAR KOMERSIAL KELAPA SAWIT (*Elaeis guineensis* Jacq.)**

Oleh

SIM CHOON CHEAK

Oktober 2017

Pengerusi : Professor Zaharah Abdul Rahman, PhD
Fakulti : Pertanian

Pendekatan konvensional menggunakan kadar pembajaan yang tinggi untuk mengekalkan penghasilan yang menguntungkan oleh industri kelapa sawit sering mencapai keputusan yang tidak konsisten dan tidak ekonomik. Di samping itu, potensi untuk mendapatkan hasil minyak sawit yang tinggi sering dijejaskan oleh keadaan persekitaran yang tidak memberangsangkan. Genotip yang cekap di dalam penggunaan nutrien mampu meningkatkan potensi produktiviti dan mengurangkan penggunaan sumber. Kajian terhadap interaksi di antara genotip dan nutrien menunjukkan perbezaan dalam pengambilan dan penggunaan nutrien antara kultivar komersial. Penanaman kultivar yang cekap dalam penggunaan kalium memerlukan baja kalium yang rendah dan secara tidak langsung mampu meningkatkan hasil sawit nasional serta menjimatkan penggunaan sumber baja. Ketika ini, industri kelapa sawit menghasilkan antara 13 hingga 21 kg minyak sawit untuk setiap kilogram baja kalium diggunakan. Penggunaan kultivar yang cekap dalam kalium mampu meningkatkan hasil sebanyak 50 %. Objektif kajian ini adalah untuk menilai pengaruh kalium dalam tanah terhadap pertumbuhan genotip sawit terpilih dan kecekapan penggunaan kalium oleh baka tanaman sawit ini. Di samping itu, mekanis fisiologi dalam penggunaan kalium akan dikaji. Tindak balas fenotip baka tanaman yang berlainan asal-usulnya iaitu AVROS, Nigeria dan Yangambi dibawah pengaruh kalium telah dikaji. Kadar pengambilan kalium dan pertumbuhan pokok diukur di dalam rumah kaca dengan menggunakan teknik radioisotop. Tindak balas molekular telah diprofilkan dan dikira. Genotip yang ber-kecekapan tinggi berupaya menghasilkan biojisim yang lebih tinggi (> 37.3 %), mempunyai kadar pengambilan kalium yang lebih tinggi (> 41.7 %) dan berupaya mengextrak kandungan kalium yang lebih tinggi (> 95 %) daripada tanah yang mempunyai kandungan kalium yang rendah. Genotip yang sedemikian menyesuaikan diri kepada persekitaran kalium rendah dengan mengubah tingkah laku perakaran, meningkatkan percambahan akar dan mengekalkan nisbah akar dan daun. Tiada variasi atau perubahan morfologi di antara genotip. Analisis perbandingan perbezaan transkrip antara genotip cekap dan cekap memperlihatkan peningkatan besar pengangkut potasium (KUP3, KUP8 dan KUP11) dalam akar genotip dan gen yang efisien yang memberikan toleransi terhadap stres, mengurangkan kerosakan selular, peraturan tekanan dan kalium homeostasis. Ciri-ciri kecekapan kalium diberikan oleh interaksi pelbagai mekanisme kompleks, yang

dikendalikan oleh gen gen yang mengawal proses fisiologi pengawalan tekanan, pembangunan selular dan homeostasis metabolit. Pengesanan tekanan dan mengawal selia proses selular untuk mengurangkan kesan tekanan boleh menjadi kunci dalam menoleransi pertama dan mengurangkan kerosakan kepada selular dan seterusnya meningkatkan keupayaan genotip untuk menyesuaikan, menyerap dan menggunakan nutrien dengan lebih berkesan.



ACKNOWLEDGEMENTS

Special gratitude to my supervisors, Professor Zaharah A.R., Professor Hanafi M., and Professor Siti Nor Akmar for the trust, patience and supervision. Professor Zaharah is extremely crucial to the completion of the thesis and I am proud to be under her academic guidance and motivation.

I would like to thank Pn. Zabedah Tumirin for her support, guidance and assistance in radioisotope experiments. Special appreciation to Dr. Teo T.M. and Dr. Cheng C.R. for their intellectual help and exchange.

I would like to thank my research colleagues, especially, Ms. Tan L.H., Ms. Goh C.Y., Dr. Tasren N.M., Dr. Lim C.K.; and Dr. Lee C.T. of FGV as well as the research staff who ensured the success of this project. Appreciation to Dr. Kee K.K. and Mr. Goh K.J. for allowing me the time and opportunity to complete this study. I gratefully acknowledge Advanced Agriecological Research Sdn. Bhd. and Felda Agricultural Services Sdn. Bhd. for extending their resources in supporting the study.

It has been a long journey. This thesis remains one of the most challenging tasks I have undertaken.

Forever indebted.

Sim Choon Cheak
August 2017

I certify that a Thesis Examination Committee has met on 31 October 2017 to conduct the final examination of Sim Choon Cheak on his thesis entitled "Differential Potassium Uptake and Utilization Efficiency of Oil Palm (*Elaeis guineensis* Jacq.) Commercial Cultivars" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Shamshuddin bin Jusop, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Dato' Mohd Fauzi bin Hj. Ramlan, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Ahmad Husni bin Mohd Haniff, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Alain Rival, PhD

Professor
Cirad Regional Director Office
Indonesia
(External Examiner)



NOR AINI AB. SHUKOR, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 28 December 2017

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Zaharah Bt. Abdul Rahman, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairperson)

Mohamad Hanafi Musa, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Siti Nor Akmar Abdullah, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of the thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification / fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection via Turnitin software.

Signature: _____ Date: _____

Name and Matric No: Sim Choon Cheak (GS29196)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: Zaharah Bt. Abdul Rahman

Signature: _____
Name of
Member of
Supervisory
Committee: Mohamad Hanafi Musa

Signature: _____
Name of
Member of
Supervisory
Committee: Siti Nor Akmar Abdullah

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xvii
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	
2.1 Fertilizer consumption by the oil palm industry	3
2.2 Potassium requirements in oil palm	3
2.2.1 Role of potassium in increasing fresh fruit bunches production and oil biosynthesis	5
2.2.2 Potassium accumulation and remobilization between sink and source organs	7
2.3 Oil palm breeding and selection for improved yield	8
2.4 Oil palm root system and nutrient absorption	9
2.5 Nutrient efficient plants	11
2.5.1 Nutrient efficient model plants	13
2.5.2 Nutrient efficient oil palm	13
2.5.3 Potassium (K) efficient oil palm	13
2.6 Physiological adaptation responsible for Potassium Use Efficiency	17
2.6.1 Biochemical response: exudation of organic compounds to facilitate the release of non- exchangeable soil potassium	18
2.6.2 Morphological response: improved root system of K-efficient genotype	19
2.6.3 Molecular response: increased potassium uptake in roots of K-efficient genotype	20
2.7 Molecular mechanisms of potassium uptake and transport in plants	21
2.7.1 Potassium uptake system in plants: High Affinity Uptake (HAT) vs. Low Affinity Uptake (LAT)	23
2.8 Summary	25

3	POTASSIUM UPTAKE AND UTILIZATION EFFICIENCY OF THE SELECTED ELAEIS GUINEENSIS GENOTYPES	
3.1	Introduction	26
3.2	Materials and Methods	26
3.2.1	<i>Elaeis guineensis</i> genotypes	26
3.2.2	Experimental evaluation and treatments	26
3.2.3	Potassium utilization	27
3.2.3.1	Plant mineral composition and soil analysis	27
3.2.3.2	Nursery management	28
3.2.4	Potassium uptake	29
3.2.4.1	Location of experiment, sand culture and the modified Hoagland solution	30
3.2.4.2	Rubidium-86 as potassium tracer and the Potassium Derived from Fertilizer (KdF)	31
3.2.4.3	Potassium uptake rate of the primary root	31
3.2.5	Statistical analysis	31
3.3	Results	32
3.3.1	Suitability of Rubidium as Potassium tracer	32
3.3.2	Rate of Potassium uptake by the excised roots	33
3.3.3	Dry matter yield, potassium accumulation and the Potassium derived from fertilizer (KdF) in response to potassium supplies	36
3.3.4	Potassium Uptake Efficiency (KUpE) and Potassium Utilization Efficiency (KUtE) of the five oil palm genotypes	38
3.3.5	Genotypic variation in potassium concentrations of tissues and potassium repartition	39
3.3.6	Indicative nutrient uptake and utilization efficiency of other major nutrient elements (N, P and Mg)	41
3.4	Discussions	42
4	GENETIC VARIATION IN THE PHYSICAL ADAPTATION OF OIL PALM TO POTASSIUM SUPPLY	
4.1	Introduction	44
4.2	Materials and Methods	44
4.2.1	Growth and vegetative parameters	44
4.2.2	Soil Plant Analysis Development Chlorophyll (SPAD Chl) index and leaf chlorophyll content	45
4.2.3	Histology studies	45
4.2.4	Statistical analysis	45
4.3	Results	46

4.3.1	Genetic variation in biomass production and resource remobilization under deficient potassium supplies	46
4.3.1.1	Variable responses in the repartitioning of the above ground biomass among the genotypes	49
4.3.1.2	Variation in root growth and adaptation to potassium supplies	49
4.3.2	Changes in leaf growth, production rate and morphology in response to potassium inputs	52
4.3.3	Effect of potassium status on chlorophyll content and photosynthetic rate	54
4.4	Discussion	56
5	MOLECULAR MECHANISM OF OIL PALM IN RESPONSE TO POTASSIUM	
5.1	Introduction	57
5.2	Materials and Methods	57
5.2.1	Plant materials and total RNA extraction	57
5.2.2	Transcriptome analysis: Identification of differentially expressed genes in the two genotypes	58
5.2.3	Transcriptome analysis: Relative quantification of the putative potassium transporter gene	58
5.2.4	Multiple sequence alignment and protein motif identification of the KUP3, KUP8 and KUP11 orthologous gene	59
5.2.5	Statistical analysis	59
5.3	Results	60
5.3.1	Comparative transcriptome analysis: Differentially expressed genes of the potassium efficient and in-efficient genotypes in response to potassium stress	60
5.3.1.1	Commonly expressed gene between the K efficient and in-efficient genotypes.	61
5.3.1.2	Transcripts upregulated by the potassium efficient genotype improving growth and tolerance to potassium stress	62
5.3.1.3	Upregulated transcripts by the K in-efficient, Genotype C	65
5.3.2	Regulation and expression profile of the putative potassium transporters in adaptation to K deficient environment	65

5.3.3	Multiple sequence alignment and protein motif identification of the KUP3, KUP8 and KUP11 orthologous gene	68
5.4	Discussion	71
6	CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH	
6.1	Conclusion	74
6.2	Implication of study and future research	75
	REFERENCES	76
	APPENDICES	90
	BIODATA OF STUDENT	101
	LIST OF PUBLICATIONS	102

LIST OF TABLES

Table	Page
2.1 Main effects of potassium treatments on the oil palm bunch component.	6
2.2 Nutrient uptake (kg) per palm per year.	7
2.3 Main effects of K fertilizer on nutrient content in different palm compartments.	7
2.4 Performance of unselected and selected Duras and Teneras in breeding trials.	8
2.5 Radioactivity counts ($\times 10^5$) of ^{32}P at various positions from P0 after 48 hours.	10
2.6 Yield (vegetative / grain) variation between efficient and non-efficient crop genotype.	12
2.7 Potassium uptake by ten potato genotypes from exchangeable and nonexchangeable soil K in the absence of K fertilizer at 83 days.	19
2.8 Dry weight of different organs of 3 tobacco genotypes and comparison of K^+ uptake.	20
3.1 Lineages of the studied oil palm progenies.	27
3.2 Nutrient input rate and timing of application used in this experiment.	28
3.3 The physical and chemical properties of soils, Typic Kandiodults, used in this experiment.	29
3.4 Nutrient composition of $\frac{1}{2}$ strength modified Hoagland solution No. 2a as carrier solution used in this experiment.	30
3.5 Total dry matter production and accumulated potassium.	36
3.6 Specific activity of ^{86}Rb and the Potassium derived from Fertilizer (KdfF).	38
3.7 A summary of the potassium uptake efficiency (KUpE), potassium utilization efficiency (KUtE) and other nutrient use efficiency parameters of the tested progenies.	39
5.1 Positive up-regulated transcript in expression by the K-efficient, Genotype A under potassium stress	63
5.2 Positive up-regulated transcript in expression by the in-efficient, Genotype C under potassium stress.	65

LIST OF FIGURES

Figure		Page
2.1	Comparison of yield response towards potassium supply in different soil group.	4
2.2	Distribution of roots from one primary root in a 10-year-old palm.	9
2.3	Lignified and unlignified portion of primary root of oil palm.	10
2.4	Differences in P efficiency of wheat and soybean varieties grown on low-P soils.	11
2.5a	Qualitative representation of the genetic effects on the leaf nutrient concentration (%) of major element (a. Nitrogen and Phosphorus).	14
2.5b	Qualitative representation of the genetic effects on the leaf nutrient concentration (%) of major element (b. Potassium and Magnesium).	15
2.6	Average FFB yield (2003 – 2009) of 3 different oil palm genotypes at 3 rates of K ₂ O inputs at optimum N.	16
2.7	FFB yield and K level (2003 – 2009) of 3 different oil palm genotypes.	16
2.8	Potential mechanisms of K uptake and utilization efficiency and their interactions that influence the expression of K-efficient genotype.	17
2.9	Interrelationships of various forms of soil K.	19
2.10	Average K uptake by wheat genotypes growing for 8 weeks in K deficient soil with or without K application.	20
2.11	Movement of inorganic nutrients.	21
2.12	Overview of the various transport processes on the plasma membrane and tonoplast of plant cells.	22
2.13	Models of transporters responsible for mediating potassium transport	24
3.1	Mean of log ₁₀ ⁸⁶ Rb detected in the selected genotype supplied with different carrier solution.	33
3.2	Decay activity of ⁸⁶ Rb in excised root tip of oil palm and the uptake rate within 24 hours after exposure to ⁸⁶ RbCl ₂ solution.	34
3.3	Estimated rate of rubidium uptake at initial uptake prior to saturation phase.	34
3.4	Decay activity of ⁸⁶ Rb profile in the excised non-lignified root tip of the selected genotypes.	35
3.5	Dry matter production of the selected genotypes in response to deficient (-K) and adequate (+K) potassium supplies.	37
3.6	Potassium concentration and remobilization between palm tissues.	40
3.7	Indicative Nitrogen, Phosphorus and Magnesium content of the evaluated genotypes.	41
3.8	Comparison of the “indicative nutrient uptake efficiency” and the “indicative nutrient utilization efficiency” of the major element (Nitrogen, Phosphorus and Magnesium) amongst the tested genotype.	42

4.1	Changes in root to shoot ratio the selected genotypes (mean \pm S.E.) in response to potassium environment.	47
4.2	Dry matter accumulation of the selected genotypes (mean \pm S.E.) in response to potassium inputs at the end of experimental period.	48
4.3	Shoot biomass reallocation to sub-tissues; the bole, rachis and pinnae under different potassium supplies.	50
4.4	Root mass, root growth and F:C ratio in response to potassium inputs.	51
4.5	The frond production rate of the respective genotypes as affected by potassium inputs.	52
4.6	Leaf stomata density of the oil palm genotypes.	53
4.7	Effect of potassium inputs on leaf anatomy of the K efficient and in-efficient oil palm genotype.	54
4.8	Total chlorophyll concentration and net photosynthetic rate in pinnae of the respective oil palm genotypes as per affected by potassium inputs	55
5.1	Common and unique transcripts between Genotype A and Genotype C in response to potassium stress.	60
5.2	Functional classification of the differentially expressed transcripts under potassium deficient environment	61
5.3	Expression change of FAD2 (Unigene ID: At.23898) and SAP5 (Unigene ID: Os.12139) by Genotype A and Genotype C in response to potassium deficient environment.	62
5.4	Relative expression of KUP genes in leaves of different oil palm genotypes ground under potassium deficient environment	66
5.5	Differential expression profile of the KUP 3, KUP 8 and KUP 11 between the potassium efficient oil palm, Genotype A and the inefficient Genotype C.	67
5.6	Unrooted phylogenetic tree constructed and motif distribution of translated KUP3, KUP8 and KUP11 genes obtained from <i>Elaeis guineensis</i> (Eg), <i>Ananas comosus</i> (Ac), <i>Arabidopsis thaliana</i> (At), <i>Musa acuminata</i> (Ma), <i>Oryza sativa</i> (Os), <i>Phoenix dactylifera</i> (Pd) and <i>Zea mays</i> (Zm).	69
5.7	Protein motif sequence alignment of the KUP obtained by MEME.	70
5.8	Molecular responses in adaptation of Genotype A to potassium stress environment.	72

LIST OF ABBREVIATIONS

^{86}Rb	Radioactive isotope of Rubidium
Ca	Calcium
CEC	Cation exchange capacity
CPM	Count per minute
CRD	Complete Randomized Design
DPM	Disintegrations per minute
ICP-OES	Inductively Coupled Plasma - Optical Emission Spectrometry
K	Potassium
KdfF	Potassium derived from fertilizer
KUE	Potassium use efficiency
KUpE	Potassium uptake efficiency
KUtE	Potassium utilization efficiency
Mg	Magnesium
MgUpE	Magnesium uptake efficiency
MgUtE	Magnesium utilization efficiency
N	Nitrogen
NE	Nutrient efficient
NitUpE	Nitrogen uptake efficiency
NitUtE	Nitrogen utilization efficiency
P	Phosphorus
PUpE	Phosphorus uptake efficiency
PUtE	Phosphorus utilization efficiency
RCBD	Randomized Complete Block Design
RNA	Ribonucleic acid



© COPYRIGHT UPM

CHAPTER 1

INTRODUCTION

Occupying merely 4.8 % of the total oil crop area planted worldwide, the oil palm contributed approximately 34.7 % of the global vegetable oil production, ranking it the highest oil producing crop known to-date. At a global production average of 3.8 tonnes of crude palm oil (CPO) per hectare, the palm oil productivity has very much stagnated over the last two decades especially in Malaysia. The stagnation could be attributed to the expansion of oil palm plantation on marginal land where the soil and terrain is least favorable for oil palm cultivation (Jalani *et al.* 2002). The efficient dissemination of selected planting material to end users, particularly to small holders is also a key factor for productivity improvement in the oil palm.

The projected biological potential of oil palm via experimental data indicates that the best progenies could be capable of yielding 11.5 tonnes of CPO per hectare. The highest plantation yield reported to-date achieved an average of about 6.5 tonnes of CPO, which is equivalent to about half of its biological yield potential. The vast gap between this biological yield potential and actual yield attained indicates an immense opportunity for yield improvement. The productivity of CPO increased from 1.3 to 6.5 tonnes per hectare through breeding advances, introduction of pollinating weevils and improved management whilst fertilization alone accounted for 29 % of the increase (Davison, 1993; Corley and Tinker, 2003). Combination of both adequate fertilization regime and improved materials that could thrive under poor and marginal growing conditions could provide viable alternatives in ensuring and improving sustainable production of oil palms grown on those marginal areas.

Amongst all oil crops, oil palm consumed the highest amount of potash fertilizer i.e. about 7 % of total global potash consumption, whilst soybean and other oil crops collectively consumed 12 % (Heffer, 2013). On the contrary, consumption of nitrogenous and phosphorus fertilizer is relatively low with potash constituting almost 60 % of total fertilizer usage for oil palm cultivation. The demand of potassium inputs for growth and oil production increases in parallel to the yield and it is further elevated by losses caused by the degraded land in the tropics. On per hectare basis, the oil palm requires up to 250 kg of potassium annually (Goh *et al.* 1994) and its planting on tropical soils that are inherently low in potassium fertility and prone to potassium leaching (Amberger, 2006) further contributes to high potassium requirement.

Yield responses to potassium fertilization are mostly soil-dependent and are affected by the soils inherent potassium fertility; higher yield response is recorded on soils with low soil potassium reserve and vice versa. Such yield response variations often translate into varying degree of optimal potash fertilization rate which is generally site specific. Generally, less than 1.3 kg K palm⁻¹ yr⁻¹ on fertile coastal soils to as high as 6.0 kg K

$\text{palm}^{-1} \text{ yr}^{-1}$ are needed for maximum yield on sandy textured soils. On most inland sedentary soils, many of the long-term fertilization trial showed an optimum range of 1.8 to 2.3 kg of potassium $\text{palm}^{-1} \text{ yr}^{-1}$.

Realizing such limitations, gradual effort is now being concentrated into improving the oil palm productivity via adopting nutrient efficient genotypes which could potentially lead to higher productivity on marginal land and eventually to a more sustainable use of resources and to the sustainable production of palm oil. The main objectives of this study are (1) to assess the variability of oil palm genotypes in potassium use efficiency and (2) to identify the physiological mechanism underlying the adaptation to potassium-deficient soil and the increase of potassium use efficiency.



REFERENCES

- Ahn S.J., Shin R., Schachtman D.P. (2004) Expression of KT/KUP genes in Arabidopsis and the role of root hairs in K⁺ uptake. *Plant Physiology* 134: 1135-1145.
- Ali L., Rahmatullah, Aziz T., Ashraf M., Maqsood M.A., Kanwal S. (2010) Differential potassium influx influences growth of two cotton varieties in hydroponics. *Pakistan J. Bot.* 42.2: 943-953.
- Amberger A. (2006) Soil fertility and plant nutrition in the tropics and subtropics. International Fertilizer Industry Association. IFA, Paris, France. ISBN 2-9523139-0-3.
- An S.H., Sohn K.H., Choi H.W., Hwang I.S., Lee S.C., Hwang B.K. (2008) Pepper pectin methylesterase inhibitor protein CaPMEI1 is required for anti-fungal activity, basal disease resistance and abiotic stress tolerance. *Planta*. 228: 61–78.
- Andrews M., Sprent J.I., Raven J.A., Eady P.E. (1999) Relationships between shoot to root ratio, growth and leaf soluble protein concentration of *Pisum sativum*, *Phaseolus vulgaris* and *Triticum aestivum* under different nutrient deficiencies. *Plant, Cell and Environment*. 22: 949-958.
- Argen G. and Franklin O. (2003) Root : Shoot ratios, optimization and nitrogen productivity. *Annals of Botany*. 92.6: 795-800.
- Armengaud P., Breitling R., Amtmann A. (2004) The potassium dependent transcriptome of Arabidopsis reveals a prominent role of jasmonic acid in nutrient signaling. *Plant Physiology*. 136: 2556-2576.
- Armengaud P., Breitling R., Amtmann A. (2010) Coronatine-Insensitive 1 (COI1) mediates transcriptional responses of *Arabidopsis thaliana* to external potassium supply. *Molecular Plant*. 3.2: 390-405.
- Ashley M.K., Grant M., Grabov A. (2006) Plant responses to potassium deficiencies: a role for potassium transport proteins. *Journal of Experimental Botany*. 57: 425–436.
- Bailey T.L., Boden M., Buske F.A., Frith M., Grant C.E., Clementi L., Ren J.Y., Li W.W., Noble W.S. (2009) MEME Suite: Tools for motif discovery and searching. *Nucleic Acids Research*. 37: 202-208.
- Baligar V.C., Fageria N.K., He Z.L. (2001) Nutrient use efficiency in plants. *Communications in Soil Science and Plant Analysis*, 32: 7-8, 921-950.
- Bange G. (1979) Comparison of compartmental analysis for Rb⁺ and Na⁺ in low-salt and high-salt barley roots. *Physiol Plant* 46:179 – 183.
- Belknap W.R. and Garbarino J.E. (1996) The role of ubiquitin in plant senescence and stress response. *Trends in Plant Science*. 1.10: 331-335.

- Bennet P.C., Choi W.J., Rogera J.R. (1998) Microbial destruction of feldspars. *Mineral Management*. 8.62: 149-150.
- Benton J.J.Jr. (2001) *Laboratory guide for conducting soil tests and plant analysis*. New York: CRC Press LLC.
- Benton J.J.Jr. (2003). *Agronomic handbook. Management of crops, soils, and their fertility*. CRC Press, New York, USA.
- Bourgis F., Kilaru A., Cao X., Ngando-Ebongue G., Drira N., Ohlrogge J.B., Arondel V. (2011) Comparative transcriptome and metabolite analysis of oil palm and date palm mesocarp that differ dramatically in carbon partitioning. In: *Proceedings of the National Academy of Sciences of the United States of America*. 108.44:18186.
- Callis J. (2014) The ubiquitination machinery of the ubiquitin system. In: *The Arabidopsis Book*. 12: e0174. 1-35. doi: 10.1199/tab.0174.
- Chan P.L., Rose R.J., Abdul M.A.M., Zainal Z., Low E.T.L., Ooi C.L.L., Ooi S.E., Yahya S., Singh R. (2014) Evaluation of reference genes for quantitative real-time PCR in oil palm elite planting materials propagated by tissue culture. *PLOS One*. 9.6: e99774.
- Chen J. and Gabelman W.H. (1995) Isolation of tomato strains varying in potassium acquisition using a sand-zeolite culture system. *Plant Soil* 176 p65-70.
- Chen L. and Hellmann H. (2013) Plant E3 ligases: Flexible enzymes in a sessile world. *Molecular Plant* 6.5: 1388-1404.
- Cheng Y., Qi Y., Zhu Q., Chen X., Wang N., Zhao X., Chen H., Cui X, Xu L., Zhang W. (2009) New changes in the plasma membrane associated proteome of rice roots under salt stress. *Proteomics*. 9: 3100-3114.
- Claasen N. and Steingrobe B. (1999) Mechanistic simulation models for a better understanding of nutrient uptake from soil. In: Rengel Z (ed) *Mineral Nutrition of Crops: Fundamental Mechanisms and Implications*. Food Products Press, Binghamton, NY, 327 – 367.
- Claassen N., Syring K.M., Jungk A. (1986) Verification of a mathematical model by simulating potassium uptake from soil. *Plant Soil*. 95: 209 – 220.
- Conde A., Chaves M.M., Geros H. (2011) Membrane transport, sensing and signaling in plant adaptation to environmental stress. *Plant Cell Physiology*. 52.9: 1583-1602.
- Coolen S., Proietti S., Hickman R., Olivas N.H., Huang P., Van Verk M.C. (2016) Transcriptome dynamics of *Arabidopsis* during sequential biotic and abiotic stresses. *Plant Journal*. 86: 249-267.
- Corley R. H. V. and Tinker P. B. (2003) *The Oil Palm*, 4th edition. Blackwell Science. ISBN: 978-0-470-75036-0

- Cui F., Liu L.J., Zhao Q.Z., Zhang Z.H., Li Q.L., Lin B.Y., Wu Y.R., Tang S.Y., Xie, Q. (2012). Arabidopsis ubiquitin conjugase UBC32 is an ERAD component that functions in brassinosteroid-mediated salt stress tolerance. *Plant Cell*. 24: 233-244.
- Cuin T.A., Pottosin I.I., Shabala S.N. (2008) Mechanisms of potassium uptake and transport in higher plants. In: *Plant Membrane and Vacuolar Transporters* (ed. Jaiwal P.K., Singh R.P. and Dhanker O.P.) CABI, UK.
- Davidson L. (1993) Management for efficient cost-effective and productive plantations. In: *Proceedings of 1991 PORIM International Palm Oil Conference*, [Yusof B. et al. (eds.)]. 153-167.
- Davies C., Shin R., Liu W., Thomas M.R., Schachtman D.P. (2006). Transporters expressed during grape berry (*Vitis vinifera* L.) development are associated with an increase in berry size and berry potassium accumulation. *Journal of Experimental Botany*. 57: 3209-3216.
- Day I.S., Reddy V.S., Ali G.S., Reddy A.S.N. (2002) Analysis of EF-hand-containing proteins in Arabidopsis. *Genome Biology*. 3.10: 56.1 - 56.24.
- Dereeper A., Guignon V., Blanc G., Audic S., Buffet S., Chevenet F., Dufayard J.F., Guindon S., Lefort V., Lescot M., Claverie J.M., Gascuel O. (2008) Phylogeny.Fr: Robust phylogenetic analysis for the non-specialist. *Nucleic Acids Research*. 36: 465-469.
- Dong B., Rengel Z., Graham R.D. (1995) Root morphology of wheat genotypes differing in zinc efficiency. *Journal of Plant Nutrition*. 18.12: 2761-2773.
- Dyson B.C., Allwood J.W., Feil R., Xu Y., Miller M., Bowsher C.G., Goodacre R., Lunn J.E., Johnson G.N. (2015) Acclimation of metabolism to light in Arabidopsis thaliana: the glucose 6-phosphate/phosphate translocator GPT2 directs metabolic acclimation. *Plant, Cell and Environment*. 38: 1404-1417.
- Dyson B.C., Webster R.E., Johnson G.N. (2014) GPT2: a glucose 6-phosphate/phosphate translocator with a novel role in the regulation of sugar signalling during seedling development. *Annals of Botany*. 113: 643-652.
- Epstein, E. and Bloom, A.J. (2004) *Mineral Nutrition of Plants: Principles and perspectives*. 2nd Edition, Sinauer Associates, Inc., Sunderland.
- Epstein, E., D.W. Rains, E. Elzam (1963) Resolution of dual mechanisms of potassium absorption by barley roots. *Proc. Nat. Acad. Sci. USA*. 49: 684-692.
- Fageria N.K. and Baligar V.C. (1994) Screening crop genotypes for mineral stresses. In: *Adaptation of plants to soil stress*, University Nebraska, Lincoln (Ed: Maranville et al.). 152-159.
- Fageria N.K., Baligar V.C., Li Y.C. (2008). The role of nutrient efficient plants in improving crop yields in the twenty first century. *Journal of Plant Nutrition*. 31:1121-1157.

- Fageria, N. K. and Baligar V.C. (1994) Screening crop genotypes for mineral stresses. In: Adaptation of Plants to Soil Stress (Ed: J.W. Maranville et al.) University of Nebraska, Lincoln, NE.
- Fatland B.L., Ke J., Anderson M.D., Mentzen W.I., Cui L.W., Allred C.C., Johnston J.L., Nikolau B.J., Wurtele E.S. (2002) Molecular characterization of a heteromeric ATP-citrate lyase that generates cytosolic acetyl-coenzyme A in Arabidopsis. *Plant Physiology*. 130: 740-756.
- Fatland B.L., Nikolau B.J., Wurtele E.S. (2005) Reverse genetic characterization of cytosolic Acetyl-CoA generation by ATP-Citrate Lyase in Arabidopsis. *The Plant Cell*. 17: 182-203.
- Foster H.L. (2002) Assessment of oil palm fertiliser requirements. In: *The oil palm – management for large and sustainable yields* (Ed. by T.H. Fairhurst & R. Hardter), Potash & Phosphate Institute of Canada (ESEAP), Singapore.
- Foster H.L. and Goh H.S. (1976) Yield response of oil palm to fertilizers in West Malaysia II: Influence of soil and climatic factors. *MARDI Research Bulletin*. 5.1: 6-22.
- Foster H.L. and Prabowo N.E. 2006. Partition and transfer of nutrients in the reserve tissues and leaves of oil palm. In: *Workshop on Nutrient Needs in Oil Palm. A Dialogue Among Experts*. October 17-18, 2006, Singapore 1-17.
- Foster H.L., Chang K.C., Tayeb D.M., Tarmizi A.M., Zin Z.Z.(1985) Oil Palm yield responses to N and K fertilizers in different environments in Peninsular Malaysia. *PORIM* 16:23.
- Foster H.L., Tayeb D.M., Gurmit S. (1988) The effect of fertilisers on oil palm bunch components in Peninsular Malaysia. In: *Proceedings of the 1987 International Oil Palm Conference*, PORIM and Inc. Soc. of Planters, Kuala Lumpur: 294-304.
- Freschet G.T., Swart E.M., Cornelissen J.H.C. (2015) Integrated plant phenotypic responses to contrasting above- and below- ground resources: key roles of specific leaf area and root mass fraction. *New Phytologist*. 206: 1247-1260.
- Gan S.T. (2014) The development and application of molecular markers for linkage mapping and quantitative trait loci analysis of important agronomic traits in oil palm (*Elaeis guineensis* Jacq.). PhD thesis. University of Nottingham, Malaysia.
- Gaydou E.M. and Arrivets J. (1983) Effects of phosphorus, potassium, dolomite, and nitrogen fertilization on the quality of soybean-Yields, proteins, and lipids. *Journal of Agricultural and Food Chemistry*, 31, 765-769.
- Gierth M. and Maser P. (2007) Potassium transporters in plants: Involvement in K⁺ acquisition, redistribution and homeostasis. *FEBS Letters*. 581: 2348-2356.
- Gierth M., Maser P., Schroeder J.I. (2005) The potassium transporter AtHAK5 functions in K⁺ deprivation-induced high affinity K⁺ uptake and AKT1 K⁺ channel

contribution to K⁺ uptake kinetics in Arabidopsis roots. *Plant Physiology*.137:1105-1114.

- Gleeson S.K. (1992) Optimization of tissue nitrogen and root-shoot allocation. *Annals of Botany*. 71: 23-31.
- Goh K.J., Chew P.S., Teo C.B. (1994a) Maximising and maintaining oil palm yields on commercial scale in Malaysia: In: ISP Planters Conference on Managing Oil Palms for Enhanced Profitability (Chee, K.H., ed), ISP, Kuala Lumpur: 121-141.
- Goh K.J., Chew P.S., Teoh K.C. (1994b) K nutrition for mature oil palms in Malaysia. *IPI Research Topics* 17: 36.
- Gomez A.K. and Gomez A.A. (1984) *Statistical procedures for agricultural research*. John Wiley & Sons, New York.
- Gomez-Herreros F., Miguel-Jimenez L., Morillo-Huesca M., Delgado-Ramos L., Munoz-Centeno M., Chavez S. (2012) TFIIS is required for the balanced expression of the genes encoding ribosomal components under transcriptional stress. *Nucleic Acids Research*. 40.14: 6508-6519.
- Grabov A. (2007) Plant KT/KUP/HAK potassium transporter: single family – multiple functions. *Annals of Botany*. 99.6: 1035-1041.
- Guglielmi B., Soutourina J., Esnault C., Werner M. (2007) TFIIS elongation factor and mediator act in conjunction during transcription initiation in vivo. In: *Proceedings of the National Academy of Sciences of the United State of America*, PNAS. 104.41: 16062-16067
- Guo Y., Xiong L., Ishitani M., Zhu J.K. (2002) An Arabidopsis mutation in translation elongation factor 2 causes superinduction of CBF/DREB1 transcription factor genes but blocks the induction of their downstream targets under low temperatures. In: *Proceedings of the National Academy of Sciences of the United State of America*, PNAS. 99.11: 7786-7791.
- Gupta M., Qiu X., Wang L., Xie W., Zhang C., Xiong L., Lian X., Zhang O. (2008) KT/HAK/KUP potassium transporters gene family and their whole-life cycle expression profile in rice (*Oryza sativa*). *Molecular Genetic Genomics*. 280: 437-452.
- Hashim M.A.T. (1999) Potassium requirement of mature oil palm on coastal soils. Phd Thesis. Universiti Putra Malaysia, Serdang, Malaysia.
- Haq M.U. and Mallarino A.P. (2005) Response of soybean grain oil and protein concentration to foliar and soil fertilization. *Journal of Agronomy*. 97: 910–918.
- Heffer P. (2009). Assessment of fertilizer use by crop at the global level (2006/07 – 2007/08). International Fertilizer Industry Association (IFA), France.
- Heffer P. (2013). Assessment of fertilizer use by crop at the global level, 2010-2011. AgCom/13/39-13/111. IFA Paris, France.

- Hiscox J.D. and Israelstam G.F. (1979) A method for the extraction of chlorophyll from leaf tissue without maceration. *Canadian Journal of Botany*. 57.12: 1332-1334.
- Hoagland D.R. and Arnon D.I. (1950) The water-culture method for growing plants without soil. *California Agricultural Experiment Station Circular*. 347: 1 – 32.
- Houston K., Tucker M.R., Chowdhury J., Shirley N., Little A. (2016) The plant cell wall: A complex and dynamic structure as revealed by the responses of genes under stress conditions. *Frontiers in Plant Science*. 7.984.: 1-18.
- Hozain M., Abdelmageed H., Lee J., Kang M., Fokar M, Allen R.D., Holaday A.S. 2012. Over-expressing AtSAP5 in cotton up-regulates putative stress-responsive genes and improves the tolerance to rapidly developing water deficit and moderate heat stress. *Journal of Plant Physiology*. 169: 1261-1270.
- Hsia M.M. and Callis J. (2010) BRIZ1 and BRIZ2 proteins form a heteromeric E3 ligase complex required for seed germination and post-germination growth in *Arabidopsis thaliana*. *The Journal of Biological Chemistry*. 285.47: 37070-37081.
- Huang B. and Eissenstat D.M. (2000) Root plasticity in exploiting water and nutrient heterogeneity. In: *Plant-Environment Interaction* (Ed: Wilkonson R.) Marcel Dekker Inc. New York.
- Inthapanya P., Sipaseuth, Sihavong P., Suhathep V., Chanphengsay M., Fukai S., Basnayake J. (2000). Genotype differences in nutrient uptake and utilization for grain yield production of rainfed lowland rice under fertilized and non-fertilized conditions. *Field Crops Research*. 65: 57-68.
- Jaafar H.Z.E. and Hafiz M.I. (2012) Photosynthesis and quantum yield of oil palm seedlings to elevated carbon dioxide. In: *Advances in Photosynthesis - Fundamental Aspects* (Ed: Najafpour M.). 16: 321-340. ISBN:978-953-307-9288.
- Jacquemard J.C., Ollivier J., Erwanda., Edyana S., Pepep P. (2010). Genetic signature in mineral nutrition in oil palm (*Elaeis guineensis* Jacq.): A new panorama for high yielding materials at low fertilizer cost. In: *International Oil Palm Conference*, 1-3 July 2010, Yogyakarta, Indonesia.
- Jacquemard J.C., Tailliez B., Dadang K., Ouvrier M., Asmady. (2002). Oil Palm (*Elaeis guineensis* Jacq.) nutrition: Planting material effect. In: *International Oil Palm Conference & Exhibition*, 8-12 July 2002, Bali, Indonesia.
- Jalani B.S., Yusuf B., Darus A., Chan K.W., Rajanaidu N. (2002) Prospects of elevating national oil palm productivity: a Malaysian perspective. *Oil Palm Industry Economic Journal*. 2.2: 1-9.
- Jia Y., Yang X., Feng Y., Jilani G. (2008) Differential response of root morphology to potassium deficient stress among rice genotypes varying in potassium efficiency. *Journal of Zhejiang University*. 9.5: 427-434.

- Jones B.Jr. (2003) *Agronomic handbook: management of crops, soils, and their fertility*. CRC Press, Florida.
- Jourdan C and Rey H. (1997) Architecture and development of the oil palm (*Elaeis guineensis* Jacq.) root system. *Plant and Soil*. 189: 33-48.
- Jungk A. (2002) Dynamics of nutrient movement at the soil-root interface. In: *Plant Roots: The Hidden Half*, 3rd edition. (Eds. Waisel et al.) 587-616. Marcel Dekker Inc., New York.
- Kang M., Fokar M., Abdelmageed H., Allen, R.D. (2011). Arabidopsis SAP5 functions as a positive regulator of stress responses and exhibits E3 ubiquitin ligase activity. *Plant Molecular Biology*. 75: 451–466.
- Kang M., Abdelmageed H., Lee S., Reichert A., Mysore K.S., Allen R.D. (2013). AtMBP-1, an alternative translation product of LOS2, affects abscisic acid responses and is modulated by the E3 ubiquitin ligase AtSAP5. *The Plant Journal*. 76: 481–493.
- Karim M., Shafiqur R., Mainur R. (1971) Rb86 as tracer for potassium: I. Uptake of Rb and K by rice plant in nutrient solution. *Plant and Soil*. 35.1: 179-181.
- Kim E.J., Kwak J.M., Uozumi N., Schroeder J.I. (1998) AtKUP1: an Arabidopsis gene encoding high-affinity potassium transport activity. *Plant Cell*. 10.1: 51-62.
- Kiribuchi K., Sugimori M., Takeda M., Otani T., Okada K., Onodera H., Ugaki M., Tanaka Y., Tomiyama-Akimoto C., Yamaguchi T., Nishiyama M., Nojiri H., Yamane H. (2004). RERJ1, a jasmonic acid-responsive gene from rice, encodes a basic helix-loop-helix protein. *Biochemical and Biophysical Research Communication*. 325: 857–863.
- Kosova K., Prasil I.T., Vitamvas P. (2013) Protein contribution to plant salinity response and tolerance acquisition. *International Journal of Molecular Sciences*. 14: 6757-6789.
- Krafczyk I., Trolldenier G., Beringer H. (1984) Soluble root exudates of maize: Influence of potassium supply and rhizosphere microorganisms. *Soil Biology and Biochemistry*. 16.4: 315-322.
- Kronzucker H.J., Szczerba M.W., Britto D.T. (2003) Cytosolic potassium homeostasis revisited: 42K-tracer analysis in *Hordeum vulgare* L. reveals set-point variations in [K⁺]. *Planta*, 217: 540 – 546.
- Kunz H.H., Häusler R.E., Fettke J., Herbst K., Niewiadomski P., Gierth M., Bell K., Steup M., Fluggell U.I., Schneider A. (2010) The role of plastidial glucose-6-phosphate/phosphate translocators in vegetative tissues of *Arabidopsis thaliana* mutants impaired in starch biosynthesis. *Plant Biology*. 12:115–128.
- Läuchli A. and Epstein M. (1970) Transport of potassium and rubidium in plant roots. *Plant Physiology*. 45, 639 – 641.

- Lim C.K., Chong C.Y.W., Lin L.T.C., Ng P.H.C., Goh K.J., Melling L. (2016) Dynamic growth responses of various oil palm planting materials under different rate of fertilizers. In: Proceedings of the 7th International Crop Science Congress. 14th to 19th August. Beijing, China. Abstract No: E1-044.
- Lee C.T. (2004) Effects of genotypes, terrain and irrigation on oil palm yield, and leaf and rachis nutrient concentrations. PhD Thesis. Universiti Putra Malaysia, Serdang, Malaysia.
- Lee D., Redfern O., Orengo C. (2007) Predicting protein function from sequence and structure. *Molecular Cell Biology*. 8: 995-1005.
- Lee S.C., Lan W.Z., Kim B.G., Li L., Cheong Y.H., Pandey G.K., Lu G., Buchanan B.B., Luan S. (2007) A protein phosphorylation/dephosphorylation network regulates a plant potassium channel. In: Proceedings of the National Academy of Sciences of the United States of America. 104: 15959-15964.
- Le-Gall H., Philippe F., Domon J., Gillet F., Pelloux J., Rayon C. (2015) Cell wall metabolism in response to abiotic stress. *Plants*. 4:112-166
- Liao M.T., Fillery I.R.P., Palta J.A. (2004) Early vigorous growth is a major factor influencing nitrogen uptake in wheat. *Functional Plant Biology*. 31: 121-129.
- Liew V.K. (2008) Effects of empty fruit bunches application on oil palm root distribution, proliferation and nutrient uptake. PhD Thesis. Universiti Putra Malaysia, Serdang, Malaysia.
- Liu J. and Zhu J.K. (1998) A calcium sensor homolog required for plant salt tolerance. *Science*. 280: 1943 – 1945.
- Liu L.L., Ren H.M., Chen L.Q. Wang Y., Wu W.H. (2013) A protein kinase, calcineurin B-like protein-interacting protein kinase 9, interacts with calcium sensor calcineurin B-like protein 3 and regulates potassium homeostasis under low potassium stress in Arabidopsis. *Plant Physiology*. 161: 266-277.
- Livak K.J. and Schmittgen T.D. (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods*. 25.4: 402-408.
- Ljungman M. (2007) The transcription stress response. *Cell Cycle*. 6.18: 2252-2256.
- Lu L., Chen Y., Lu L., Lu Y., Li L. (2015) Transcriptome analysis reveals dynamic changes in the gene expression of tobacco seedlings under low potassium stress. *Journal of Genetics*. 94: 397 – 406.
- Maas E.V. and Leggett J.E. (1968) Uptake of ⁸⁶Rb and K by excised maize roots. *Plant Physiology*. 43.12: 2054-2056.
- Maathuis F.J.M., Filatov V., Herzyk P., Krijger G.C., Axelsen K.B., Chen S., Green B.J., Li Y., Madagan K.L., Sanchez-Fernandez R., Forde B.G., Palmgren M.G., Rea P.A., Williams L.E., Sanders D., Amtmann A. (2003) Transcriptome analysis of

root transporters reveals participation of multiple gene families in the response to cation stress. *The Plant Journal*. 35. 675-692.

- Mangeon A., Junqueira R.M., Sachetto-Martins G. (2010) Functional diversity of the plant glycine-rich proteins superfamily. *Plant Signalling & Behaviour*. 5.2: 99-104.
- Mao K., Dong Q., Li C., Liu C., Ma F. (2017) Genome wide identification and characterization of Apple bHLH transcription factors and expression analysis in response to drought and salt stress. *Frontiers in Plant Science*. 8: 480.
- Marschner, H. (1995). *The Mineral Nutrition of Higher Plants* (2nd Edition). Academic Press, USA.
- Marston A.L. (2015) Shugoshins: Tension-sensitive pericentromeric adaptors safeguarding chromosome segregation. *Molecular Cell Biology*. 35:634–648.
- Maser P., Thomine S., Schroeder J.I., Ward J.M., Hirschi K., Sze H., Talke I.N., Amtmann A., Maathuis F.J.M., Sanders D., Harper J.F., Tchieu J., Gribskov M., Persans M.W., Salt D.E., Kim S.A., Guerinot M.L. (2001) Phylogenetic relationships within cation transporter families of Arabidopsis. *Plant Physiology*. 126: 1646-1667.
- Melvin S.G., Lu G.Q., Zhou W.J. (2002). Genotypic variation for potassium uptake and utilization efficiency in sweet potato (*Ipomoea batatas* L.). *Field Crops Research*. 77: 7-15.
- Memon A.R., Saccomani M., Glass A.D.M. (1985) Efficiency of potassium utilization by barley *Hordeum vulgare* varieties the role of subcellular compartmentation. *Journal Experimental Botany*. 36: 1860 – 1876.
- Mi H., Dong Q., Muruganujan A., Gaudet P., Lewis S., Thomas O.D. (2010) PANTHER version 7: Improved phylogenetic trees, orthologs and collaboration with the Gene Ontology Consortium. *Nucleic Acids Research*. 38: 204-210.
- Mochida K. and Shinozaki K. (2011) Advances in omics and bioinformatics tools for system analyses of plant functions. *Plant Cell Physiology*. 52.12: 2017-2038.
- Mohidin H., Hanafi M.M., Rafii Y.M., Abdullah S.N.A., Idris A.S., Man S., Idris J., Sahebi M. (2015) Determination of optimum levels of nitrogen, phosphorus and potassium of oil palm seedlings in solution culture. *Bragantia*. 74.3: 247-254.
- Ng S.K. (2002) Nutrition and nutrient management of oil palm: New thrust for the future perspective. In: Pasricha, N.S., Bansal, S.K. (Eds.), *Potassium for Sustainable Crop Production: International Symposium on the Role of Potassium*. Potash Research Institute of India and International Potash Institute, New Delhi. 415–429.
- Nielsen N.C., Adey A., Stumpf P.K. (1979) Fat metabolism in higher plants: Further characterization of wheat germ Acetyl coenzyme A carboxylase. *Arch. Biochem. Biophys*. 192: 446-456.

- Pan X.W., Li W.B., Zhang Q.Y., Li Y.H., Liu M.S. (2008). Assessment on phosphorus efficiency characteristics of soybean genotypes in phosphorus-deficient soils. *Agricultural Sciences in China*. 7: 958-969.
- Paramanathan S, (2000). *Soils of Malaysia: Their characteristics and identification*, Volume 1. Academy of Sciences Malaysia, Kuala Lumpur, Malaysia.
- Park S.J., Kwak K.J., Oh T.R., Kim Y.O., Kang H. (2009) Cold shock domain proteins affect seed germination and growth of *Arabidopsis thaliana* under abiotic stress conditions. *Plant Cell Physiology*. 50: 869-878.
- Pelechano V., Jimeno-Gonzalez S., Rodriguez-Gil A., Garcia-Martinez J., Perez-Ortin J.E., Chavez S. (2009) Regulon-specific control of transcription elongation across the yeast genome. *PLoS Genetic*. 5: e1000614.
- Peter B.B., Jonathan R.H., Janina J., Rafael L.B., Saroj P., Caroline E.S., Malcolm J.H. (2010). Nitrogen efficiency of wheat: Genotypic and environmental variation and prospects for improvement. *European Journal of Agronomy*. 33: 1-11.
- Pettersson S. and Jensen P. (1983) Variation among species and varieties in uptake and utilization of potassium. *Plant and Soil*. 72.2: 231-237.
- Pettigrew W.T. (2008) Potassium influences on yield and quality production for maize, wheat, soybean and cotton. *Physiologia Plantarum*. 133: 670-681.
- Pilot G., Gaymard F., Mouline K., Cherel I., Sentenac H. (2003) Regulated expression of *Arabidopsis* Shaker K⁺ channel genes involved in K⁺ uptake and distribution in the plant. *Plant Molecular Biology*. 51: 773-787.
- Polley L.D. and Johns W.H. (1979) Rubidium (Potassium) uptake by *Arabidopsis*. *Plant Physiology*. 64: 374-378.
- Prabowo N.E. and Foster H.L. (1998) Variation in oil and kernel extraction rates of oil palms in North Sumatra due to nutritional and climatic factors. In: *International Oil Palm Conference*. Nusa Dua-Bali, Indonesia. September 23-25, 1998.
- Prabowo N.E., Foster H.L., Silalahi A.J. (2006) Recycling oil palm bunch nutrients. In: *International Oil Palm Conference*. June 19 – 23, 2006, Nusa Dua, Bali, Indonesia.
- Qian H., Lu H., Ding H., Lavoie M., Li Y., Liu W., Fu Z. (2015) Analyzing *Arabidopsis thaliana* root proteome provides insights into the molecular bases of enantioselective imazethapyr toxicity. *Scientific Report*. 5: 11975.
- Rankine I.R. and Fairhurst T.H. (1999) *Field Handbook: Oil Palm Series Volume 1ñ Nursery*. (Oil Palm Series), Potash & Phosphate Institute (PPI), Potash & Phosphate Institute of Canada (PPIC) and 4T Consultants (4T), Singapore, 135 p.
- Rengel Z. (2013) *Improving Water and Nutrient-Use Efficiency in Food Production Systems*. John-Wiley and Son. ISBN: 978-0-8138-1989-1.

- Rengel Z. and Damon P.M. (2008) Crops and genotypes differ in efficiency of potassium uptake and use. *Physiologia Plantarum*. 133.4: 624-636.
- Ringli C., Keller B., Ryser U. (2001) Glycine-rich proteins as structural components of plant cell walls. *Cellular and Molecular Life Sciences*. 58: 1430-1441.
- Rivera Y.D.M., Chacon L.M., Bayona C.J., Romero H.M. (2012) Physiological response of oil palm interspecific hybrids (*Elaeis oleifera* H.B.K. Cortes versus *Elaeis guineensis* Jacq.) to water deficit. *Brazilian Journal of Plant Physiology*. 24.4: 273-280.
- Römer W. and Schenk H. (1998). Influence of genotype on phosphate uptake and utilization efficiencies in spring barley. *European Journal of Agronomy*. 8: 215-224.
- Ruan L., Zhang J., Xin X., Zhang C., Ma D., Chen L., Zhao B. (2015) Comparative analysis of potassium deficiency-responsive transcriptomes in low potassium susceptible and tolerant wheat (*Triticum aestivum* L.) *Scientific Reports*. 5: 10090.
- Sachetto-Martins G., Franco L.O., Oliveira D. (2000) Plant glycine-rich proteins: a family or just proteins with a common motif. *Biochimica et Biophysica Acta*. 1942. 1 - 14.
- Samal D., Sadana U.S., Gill A.A.S. (2003) Mechanistic approach to study manganese influx and its depletion in the rhizosphere of wheat and raya. *Communications in Soil Science and Plant Analysis*. 34: 3033-3044.
- Sassi A., Mieulet D., Khan I., Moreau B., Gaillard I., Sentenac H., Very A. (2012). The rice monovalent cation transporter OsHKT2;4: revisited ionic selectivity. *Plant Physiology*. 160: 498–510.
- Schmittgen T.D. and Livak K.J. (2008) Analyzing real-time PCR data by the Comparative CT method. *Nature Protocols*. 3: 1101-1108.
- Seguin P. and Zheng W. (2006) Potassium, phosphorus, sulfur, and boron fertilization effects on soybean isoflavone content and other seed characteristics. *Journal of Plant Nutrition*. 29: 681–698.
- Sievers F., Wilm A., Dineen D., Gibson T.J., Karplus K., Li W.Z., Lopez R., McWilliam H., Remmert M., Soding J., Thompson J.D., Higgins D.G. (2011) Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology*. 7: 539.
- Simkins C.A. and Overdahl C.J. (1982) Fertilizing sunflowers on sandy lands. In: *Better Crops*, Volume LXVI, Potash and Phosphate Institute, USA.
- Singh P. and Blanke M.M. (2000) Deficiency of potassium but not phosphorus enhances root respiration. *Plant Growth Regulation*. 32:77-81.

- Singh R., Ong-Abdullah M., Low E.T.L., Manaf M.A.A., Rosli R., Nookiah R., *et. al.* (2013). Oil palm genome sequence reveals divergence of interfertile species in Old and New worlds. *Nature*. 500:7462, 335-339.
- Song F., Han X., Zhu X., Herbert S.J. (2012) Response to water stress of soil enzymes and root exudates from drought and non-drought tolerant corn hybrids at different growth stages. *Canadian Journal of Soil Science*. 92.3: 501-507.
- Song S.K. and Huang P.M. (1988) Dynamics of potassium release from potassium bearing minerals as influenced by oxalic and citric acids. *Soil Science Society of America Journal*. 52: 383-390.
- Song Z.Z., Ma R.J., Yu M.L. (2015) Genome-wide analysis and identification of KT/HAK/KUP potassium transporter gene family in peach (*Prunus persica*). *Genetics and Molecular Research*. 14.1: 774-787.
- Sparks D.L. (2000) Bioavailability of soil potassium. In: *Handbook of Soil Science*, Section D1.4, pg D-38 to D-53. CRC Press, USA.
- Steingrobe B. (2005) A sensitive analysis for assessing the relevance of fine-root turnover for P and K uptake. *Journal of Plant Nutrition and Soil Science*. 168: 496-502.
- Szczerba M.W., Britto D.T., Kronzucker H.J. (2009) K⁺ transport in plants: Physiology and molecular biology. *Journal of Plant Physiology*. 166.5: 447-466.
- Taiz L. and Zeiger E. (2002) *Plant Physiology* (3rd Edition). Sinauer Associates, Inc. USA.
- Tan C.C. (2008) Nursery practices for production of superior oil palm planting materials. In: *Proc. Seminar on agronomic principles and practices of oil palm cultivation*. Agricultural Crop Trust Preprint. Also published in Goh K.J., Chiu S.B. and Paramanathan S. (eds). 2011. *Agronomic Principles and Practices of Oil Palm Cultivation*. Agricultural Crop Trust: 145-169.
- Terry J.R., Juan P.T., Michael T.R., Fukuta Y., Wissuwa M. (2010). Genotypic variation in grain phosphorus concentration and opportunities to improve P-use efficiency in rice. *Field Crops Research*. 119: 154-160.
- Tirol-Padre A., Ladha J.K., Singh U., Laureles E., Punzalan G., Akita S. (1996) Grain yield performance of rice genotypes at suboptimal levels of soil N as affected by N uptake and utilization efficiency. *Field Crops Research*. 46: 127-143.
- Trehan S.P. and Sharma R.C. (2002) Potassium uptake efficiency of young plants of three potato cultivars as related to root and shoot parameters. *Communications in Soil Science and Plant Analysis*. 33: 1813 - 1823.
- Trehan S.P., Dessougi H.E., Claassen N. (2005) Potassium efficiency of 10 potato cultivars as related to their capability to use non-exchangeable soil potassium by chemical mobilization. *Communications in Soil Science and Plant Analysis*. 36: 1809 - 1822.

- Tu S.X., Guo Z.F., Sun J.H. (2007) Effect of oxalic acid on potassium release from typical chinese soils and minerals. *Pedosphere*. 17.4: 457-466.
- Wang H., Inukai Y., Yamauchi A. (2006) Root development and nutrient uptake. *Critical Reviews in Plant Sciences*. 25: 279-301.
- Wang X., Vignjevic M., Jiang D., Jacobsen S., Wollenweber B. (2014) Improved tolerance to drought stress after anthesis due to priming before anthesis in wheat (*Triticum aestivum* L.) var. Vinjett. *Journal of Experimental Botany*. 65.22: 6441-6456.
- Wang H.Y., Shen Q.H., Zhou J.M., Wang J., Du C.W., Chen X.Q. (2011) Plants use alternative strategies to utilize nonexchangeable potassium in minerals. *Plant Soil*. 343: 209-220.
- Waters S., Gilliam M., Hrmova M. (2013) Plant High-Afinity Potassium (HKT) Transporters involved in salinity tolerance: Structural insight probe difference in ion selectivity. *International Journal of Molecular Sciences*. 14: 7660-7680.
- Weber E.J. (1985) Role of potassium in oil metabolism. In: *Potassium in Agriculture* (Munson R.D., ed). ASA, CSSA and SSSA, Madison, WI, pp 425-442.
- Wellburn A.R. (1994) The spectral determination of chlorophylls a and b, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. *Journal of Plant Physiology*. 144: 307-313.
- White A.C., Rogers A., Rees M., Osborne C.P. (2016) How can we make plants grow faster? A source-sink perspective on growth rate. *Journal of Experimental Botany*. 67.1: 31-45.
- Wu R., Grissom J.E., McKeand S.E., O'Malley D.M. (2004) Phenotypic plasticity of fine root growth increases plant productivity in pine seedlings. *BMC Ecology*. 4: 14.
- Yan X.L., Wu P., Ling H.Q., Xu G.H., Xu F.S., Zhang Q.F. (2006) Plant Nutriomics in China: An Overview. *Annals of Botany* 98: 473-482.
- Yang T.Z., Lu L.M., Xia W., Fan J.H. (2007) Characteristics of potassium-enriched, flue-cured tobacco genotype in potassium absorption, accumulation and inward potassium currents of root cortex. *Agricultural Sciences in China*. 6.12: 1479-1486.
- Yin X. and Vyn T.J. (2003) Potassium placement effects on yield and seed composition of no-till soybean seed in alternate row widths. *Journal of Agronomy*. 95: 126-132.
- Zaharah A.R., Sharifuddin H.A.H., Sahali A.M., Hussein M.M.S. (1989) Fertilizer placement studies in mature oil palm using isotope technique. *Planter*. 65: 384-388.
- Zaharah, A.R., H.A.H. Sharifuddin, A.M. Sahali. (1991) The use of ^{86}Rb as a tracer to quantify potassium uptake by oil palm. *Proceedings of the PORIM International*

Palm Oil Conference: Progress, Prospects and Challenges Towards the 21st Century, September 9-14, 1991, Kuala Lumpur, Malaysia, pp: 499-501.

Zapata, F. and Axmann H. (1995) ^{32}P isotopic techniques for evaluating the agronomic effectiveness of rock phosphate materials. *Fertilizer Research*. 41: 189-195.

Zhao Y., Wang T., Zhang W., Li Xia (2011) SOS3 mediates lateral root development under low salts stress through regulation of auxin redistribution and maxima in *Arabidopsis*. *New Phytologist*. 189: 1122 - 1134

Zhiponova M.K., Morohashi K., Vanhoutte I., Machermer-Noonan K., Revalska M., Montagu M.V., Grotewold E., Russinova E. (2014) Helix-loop-helix/basic helix-loop-helix transcription factor network represses cell elongation in *Arabidopsis* through an apparent incoherent feed-forward loop. In: *Proceedings of the National Academy of Sciences of the United State of America, PNAS*. 111.7: 2824-2829.