



UNIVERSITI PUTRA MALAYSIA

***GENETIC VARIATION AMONG *Stevia rebaudiana* Bertoni GENOTYPES
UNDER LIGHT TREATMENT ON MORPHO-AGRONOMIC TRAITS AND
ISSR MOLECULAR MARKERS***

DHULFIQAR AAMER ABDULAMEER

FP 2018 97



**GENETIC VARIATION AMONG *Stevia rebaudiana* Bertoni GENOTYPES
UNDER LIGHT TREATMENT ON MORPHO-AGRONOMIC TRAITS AND
ISSR MOLECULAR MARKERS**

By

DHULFIQAR AAMER ABDULAMEER

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

February 2018

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

To my lovely country Iraq

To my beloved family who always supported and kept praying for me day and night

to be what I am now

And

To my friends who boosted me all these past years



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

GENETIC VARIATION AMONG *Stevia rebaudiana* Bertoni GENOTYPES UNDER LIGHT TREATMENT ON MORPHO-AGRONOMIC TRAITS AND ISSR MOLECULAR MARKERS

By

DHULFIQAR AAMER ABDULAMEER

February 2018

Chairman : Professor Mohd Rafii Yusop, PhD
Faculty : Agriculture

Stevia rebaudiana Bertoni is a perennial herb with natural sweetening compounds that are 230 times sweeter than sucrose with no calorific value. It belongs to the family *Asteraceae* (Compositae) and is native to Paraguay. It is a good source of carbohydrates, protein, crude fibers and minerals. Stevioside and rebaudioside A are the two major sweetening compounds. *Stevia* is a short day plant which form flowers when day length is less than 12 hours and it is resulting in low biomass production. In this study, a set of stevia genotypes collected from different sources were evaluated for yield components and flowering characteristics both under short day length (12 hours) and long day length (14 hours) environments. Genetic variation among those genotypes was also investigated using ISSR molecular markers. A total of twenty-six stevia genotypes including two hybrids were collected. The propagation of all collected stevia genotypes was achieved through stem micro-cutting within six months. Eight genotypes namely, CHV1, Morita2, MS012, MS012 (4x), SA18, TPU, and hybrids IIUM-F₁ and UiTM-F₁ which had shown good propagation ability under mist-chamber propagation boxes and were evaluated for the agronomic performance under short day length (12 hours) and long day length (14 hours) environments using a split plot design. In general, all the agronomic characteristics of pooled genotypes were increased significantly under the extended light (14 hours). *Stevia* plants under sunlight (12 hours) started to flowering after 52 days after transplanted. On the other hand stevia plants under extended light were not flowered up to 134 days at harvesting stage. Combined biomass fresh weight mean of the eight genotypes was increased 728% under extended light compared to genotypes kept under sunlight. The biomass fresh weight mean and plant height of genotype SA18 was the highest in both extended and sunlight conditions (35.82 g and 3.96 g; 92.47 cm and 24.92 cm respectively). The Pearson correlation coefficient showed that biomass fresh weight had strongly positively correlated with all the

characters except number of branches. Genetic variation and cluster analysis based on ISSR markers showed that genotypes CHV1 and H2 were the most closely related genotypes with the highest similarity index (54%). On the other hand, genotypes MS012 and TPU were the distantly related genotypes with the lowest similarity index (24%). In conclusion, stevia plants kept under extended light showed better performance than under sunlight. Three genotypes, SA18, TPU and CHV1 were shown better performance than other genotypes, and are recommended for further utilization in future stevia breeding programs.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**VARIASI GENETIK KALANGAN GENOTIP *Stevia rebaudiana* Bertoni
DI BAWAH RAWATAN CAHAYA KE ATAS CIRI MORFO-AGRONOMI
DAN PENANDA MOLEKUL ISSR**

Oleh

DHULFIQAR AAMER ABDULAMEER

Februari 2018

Pengerusi : Profesor Mohd Rafii Yusop, PhD
Fakulti : Pertanian

Stevia rebaudiana Bertoni adalah herba saka yang mengandungi kompaun pemanis semulajadi sebanyak 230 kali lebih manis daripada sukrosa dan ianya tanpa nilai kalori. Ia tergolong di bawah keluarga *Asteraceae* (Compositae) dan yang berasal dari Paraguay. Ia merupakan sumber karbohidrat, protein, fiber mentah dan zat galian. Stevioside dan rebaudioside A adalah dua kompaun pemanis utama spesies ini. *Stevia* adalah tumbuhan hari-pendek yang mengeluarkan bunga apabila waktu siang kurang dari 12 jam dan seterusnya ianya mengakibatkan pengeluaran biojisim yang rendah. Dalam kajian ini, satu kumpulan genotip stevia yang dikumpulkan dari beberapa sumber yang berbeza telah dinilai dari segi komponen hasil dan ciri pembungaan di bawah persekitaran jangka siang yang pendek (12 jam) dan jangka siang yang panjang (14 jam). Variasi genetik di kalangan genotip juga telah dikaji dengan menggunakan penanda molikul ISSR. Sejumlah dua puluh enam aksesii stevia termasuk dua hibrid telah dikumpulkan. Pembiakan semua aksesii stevia telah dijalankan melalui keratan mikro batang dalam tempoh enam bulan. Lapan genotip iaitu; CHV1, Morita2, MS012, MS012 (4x), SA18, TPU, dan hibrid IIUM-F₁ dan UiTM-F₁), yang menunjukkan prestasi pembiakan yang baik di bawah kotak pembiakan berkabus, dan telah dinilai dari segi prestasi agronomi di bawah jangka siang yang pendek (12 jam) dan jangka siang yang panjang (14 jam) dalam satu eksperimen faktorial. Secara umumnya, kesemua ciri agronomi bagi kombinasi genotip didapati bertambah di bawah pencahayaan yang dilanjutkan. Pokok stevia di bawah rawatan cahaya (12 jam) mulai berbunga 52 hari selepas penanaman. Sebaliknya pokok di bawah pencahayaan lanjutan, pokok stevia tidak berbunga sehingga 134 hari selepas penanaman pada peringkat penuaian. Berat basah biomass keseluruhan bagi lapan genotip tersebut telah meningkat 728% di bawah pencahayaan lanjutan berbanding pokok di bawah cahaya matahari. Purata berat basah biomass dan ketinggian pokok aksesii SA18 didapati adalah tertinggi di bawah

kedua-dua persekitaran, pencahayaan lanjutan dan cahaya matahari (masing-masing 135.82 g dan 3.96 g; 92.47 cm dan 24.92 cm). Pekali korelasi Pearson menunjukkan berat biomass segar adalah berkorelasi tinggi secara positif dengan semua ciri-ciri kecuali dengan bilangan dahan. Analisa variasi genetik dan kluster berdasarkan penanda ISSR menunjukkan genotip CHV1 and H2 mempunyai perkaitan genotip yang paling rapat dengan indeks kesamaan tertinggi (54%). Sebaliknya, genotip MS012 and TPU adalah genotip yang paling berbeza dengan indeks kesamaan paling rendah (24%). Kesimpulannya, pokok di bawah pencahayaan lanjutan menunjukkan prestasi yang lebih baik daripada pencahayaan matahari. Tiga genotip iaitu SA18, TPU dan CHV1 didapati berprestasi lebih baik daripada semua genotip lain, dan dicadangkan untuk digunakan seterusnya di dalam program pembiakbakaan stevia di masa hadapan.



ACKNOWLEDGEMENTS

Alhamdulillah, In the name of Allah, the Most Gracious and the Most Merciful for the intensities and His blessing in performing this study. Special appreciate goes to my supervisor, Prof. Mohd Rafii Yusop and co-supervisor Assoc Prof. Dr Zulkefly Sulaiman, for their supervision and constant support. Special thanks to Prof. Mohamad Bin Osman for his invaluable helps of constructive comments and suggestion throughout the research activities. I would like to express my appreciation to Dr. Mahmoud Danaie,2 Mr. Mohd Farit, Dr. Parisa Azizi, Dr. Mahbood, Dr. Halimaton and Mr. Shahril for their support and knowledge. My acknowledgement also goes to all the technicians and staffs of Crop Science Department, Faculty of Agriculture for their assistance during this study. Sincere thanks to all my friends especially Firas, Quasy, Ameer, Osama, Qutaiba, Ahmad and Husam for their kindness, helping and moral support during my study.

Last but not least, my deepest gratitude goes to my beloved mother, father and brother for their endless love, helping, support, prayers and encouragement. Also my gorgeous fiancée so thankful and appreciate for her support, love and care. To those who indirectly contributed in this research, your kindness means a lot to me. Thank you so much.

Dhulfiqar Aamer Abdulameer, 2018

I certify that a Thesis Examination Committee has met on 13 July 2018 to conduct the final examination of Dhulfiqar Aamer Abdulameer on his thesis entitled "Genetic Variation Among *Stevia rebaudiana* Bertoni Genotypes Under Light Treatment on Morpho-Agronomic Traits and ISSR Molecular Markers" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Mahmud b Tengku Muda Mohamed, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Noor Azmi Shaharuddin, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Azhar Mohamad, PhD

Senior Lecturer
Malaysian Nuclear Agency
Malaysia
(External Examiner)



RUSLI HAJI ABDULLAH, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 27 September 2018

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Mohd Rafii Yusop, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Zulkefly Sulaiman, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____

Date: _____

Name and Matric No: Dhulfiqar Aamer Abdulameer, GS44809

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____

Name of Chairman
of Supervisory
Committee:

Professor Dr. Mohd Rafii Yusop

Signature: _____

Name of Member
of Supervisory
Committee:

Associate Professor Dr. Zulkefly Sulaiman

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xv
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	3
2.1 Taxonomy and Origin	3
2.2 Classification of Stevia	3
2.3 Genus of Stevia	4
2.4 Geographic range and habitat of stevia	4
2.5 Glycosides and Chemical Constituents	5
2.6 Cytogenetic	5
2.7 Morphology	6
2.8 Flower structure	6
2.9 Seed	7
2.10 Utilization of Stevia	7
2.11 Stevia Product and Commercialization	8
2.12 Chemical Compounds in Stevia	9
2.13 Cultivation	11
2.14 Propagation	11
2.14.1 Environmental Factors	11
2.14.2 Methods of Propagation	11
2.14.2.1 Stem Cutting Propagation	13
2.15 Stevia breeding	13
2.15.1 Stevia breeding in the world	13
2.15.2 Stevia Breeding in Malaysia	14
2.16 Genetic Diversity	17
2.16.1 Morpho Agronomic Diversity	17
2.16.2 Genetic and Metabolic Diversity Using RAPD and HPLC	18
2.17 Extended Day Length	19
2.17.1 Day Length	19
2.17.2 Light	20
2.17.3 Photoperidism	20
2.17.4 Effect of Photoperiods on Growth and Development	20
2.18 PCR-Based Markers	21
2.18.1 Inter-Single Simple Repeat (ISSR) marker	21

3	MATERIALS AND METHODS	23
3.1	Stevia genotypes	23
3.2	Propagation of stevia through stem micro-cuttings	23
3.2.1	Propagation Box	25
3.2.2	Acclimatization of stevia plantlets	25
3.2.3	Selection of genotypes based on number of plantlets produced	26
3.3	Evaluation of the agronomic performance of selected stevia genotypes	26
3.3.1	Lighting treatments	26
3.3.2	Experimental site and design	26
3.3.3	Parameters evaluated	26
3.3.4	Statistical analysis	28
3.4	Molecular diversity among stevia genotypes	28
3.4.1	Plant Materials	28
3.4.2	Samples preparation	28
3.4.3	DNA extraction and quantification	28
3.4.4	Selection and validation of ISSR primers	29
3.4.5	PCR optimization	29
3.4.6	Analysis of polymorphism through gel electrophoresis	30
3.4.7	Analysis of genetic relationship between stevia genotypes	30
4	RESULTS AND DISCUSSION	32
4.1	Propagation and selection of stevia genotypes based on stem micro-cuttings performance	32
4.2	Evaluating the agronomic performance of selected stevia genotypes	33
4.3	Molecular diversity of stevia genotypes	44
4.3.1	ISSR makers screening	44
4.3.2	Genetic variation among stevia genotypes	44
4.3.3	Polymorphism information content (PIC) value	50
5	CONCLUSION AND RECOMMENDATIONS FOR FUTURE STUDIES	53
5.1	Conclusion	53
5.2	Recommendations for future studies	53
	REFERENCES	55
	APPENDICES	69
	BIODATA OF STUDENT	76
	PUBLICATION	77

LIST OF TABLES

Table	Page	
2.1	Ethnomedical use of stevia	8
2.2	Commercially obtainable stevia products in USA market	9
2.3	Amount of sweet glycoside in <i>Stevia rebaudiana</i> leaves (% of the leaves dry weight)	10
2.4	List of stevia genotypes developed	13
2.5	Stevia germplasm available at the International Islamic University of Malaysia (IIUM)	14
2.6	List of stevia collected by MARDI from other countries	15
2.7	Stevia genotypes and area of collection in Malaysia	16
3.1	List of collected stevia genotypes	24
3.2	Eleven ISSR primers used for genetic diversity of stevia genotypes	29
3.3	ISSR primers used for analysis of <i>Stevia rebaudiana</i> Bertoni genotypes with their melting and annealing temperature	30
4.1	Number of propagated plants for each stevia genotype	32
4.2	Percentage of relative efficiency between extended light and sunlight of pooled genotypes on the quantitative traits	34
4.3	Percentage of relative efficiency between extended light and sunlight of different genotypes on the quantitative traits	35
4.4	Analysis of variance on agronomic traits in stevia genotypes	36
4.5	Mean comparison (Means \pm S.D) on agronomic traits in stevia genotypes under sunlight and extended light conditions	39
4.6	Pearson Correlation coefficients among quantitative traits of stevia genotypes pooled data under lightning conditions	41
4.7	Principal components analysis (PCA) for 8 morphological traits among the eight stevia genotypes	43
4.8	ISSR polymorphic primers, number of polymorphic loci, percentage of polymorphic loci, and estimates of variability across eight stevia genotypes	45



© COPYRIGHT UPM

LIST OF FIGURES

Figure		Page
2.1	Three genotypes (MS012, MS007 and MS028) with promising characteristics under Malaysia's condition	15
3.1	Propagation steps for 26 stevia genotypes using mist-chamber propagation box	25
4.1	Dendrogram using average linkage among the selected eight stevia genotypes	42
4.2	Scatter plot of PC1 and PC2 of eight stevia genotypes	43
4.3	Banding patterns of IS57 primer; L: 100 base pair ladder, 1: CHV1, 2: SA18, 3: MS012, 4: 4X, 5: F1, 6: TPU, 7: H2 and 8: Morita	46
4.4	Banding patterns of IS20 primer; L: 100 base pair ladder, 1: CHV1, 2: SA18, 3: MS012, 4: 4X, 5: F1, 6: TPU, 7: H2 and 8: Morita	46
4.5	Banding patterns of IS21 primer; L: 100 base pair ladder, 1: CHV1, 2: SA18, 3: MS012, 4: 4X, 5: F1, 6: TPU, 7: H2 and 8: Morita	47
4.6	Banding patterns of IS20 primer; L: 100 base pair ladder, 1: CHV1, 2: SA18, 3: MS012, 4: 4X, 5: F1, 6: TPU, 7: H2 and 8: Morita	47
4.7	Banding patterns of IS56 primer; L: 100 base pair ladder, 1: CHV1, 2: SA18, 3: MS012, 4: 4X, 5: F1, 6: TPU, 7: H2 and 8: Morita	48
4.8	Banding patterns of IS31 primer; L: 100 base pair ladder, 1: CHV1, 2: SA18, 3: MS012, 4: 4X, 5: F1, 6: TPU, 7: H2 and 8: Morita	48
4.9	Banding patterns of IS16 primer; L: 100 base pair ladder, 1: CHV1, 2: SA18, 3: MS012, 4: 4X, 5: F1, 6: TPU, 7: H2 and 8: Morita	49
4.10	Banding patterns of IS85 primer; L: 100 base pair ladder, 1: CHV1, 2: SA18, 3: MS012, 4: 4X, 5: F1, 6: TPU, 7: H2 and 8: Morita	49
4.11	Dendrogram illustrating genetic relationship among eight stevia genotypes, produced by UPGMA cluster tree analysis (NTSYS pc) calculated from 86 ISSR markers generated from eight ISSR primer	52

LIST OF ABBREVIATIONS

AFLP	Amplified fragment length polymorphisms
bp	Base pair
CV	Coefficient of Variation
DNA	Deoxyribonucleic acid
ISSR	Inter simple sequence repeat
NAA	α - naphthaleneacetic acid
NTSYS-pc	Numerical taxonomy and multivariate analysis system
PCA	Principal component analysis
PCR	Polymerase Chain Reactions
POPGENE	Population genetic analysis
RAPD	Random amplified polymorphic DNA
SSR	Simple sequence repeat
Taq	Thermus aquaticus
TBE	Tris-borate EDTA
UPGMA	Unweighted pair group method based on arithmetic averages

CHAPTER 1

INTRODUCTION

Human interest in natural products is increasing day by day. *Stevia rebaudiana* Bertoni commonly known as candy leaf or sweet leaf (generically as stevia) is an ancient perennial shrub of South America. It belongs to the same family with sunflower (*Asteraceae*) and produces diterpene glycosides that are 300 times lower in calorie sweeteners than saccharose. About 240 stevia species have been identified and characterized, among these species the sweetest essences have successfully been extracted from *Stevia rebaudiana*. Presently, stevia has a great potential as a modern agricultural crop since consumer demand increased for herbal foods, zero-calorie sweetener and appropriate gluco-regulation for consumers afflicted with diabetes and prevention of dental cavities (Thomas, 2010).

The chemical compounds that produce sweetness in stevia are various steviol glycosides while the major components are stevioside and rebaudioside A (Singh and Rao, 2005). Stevioside is the sweet glycoside with the highest concentration found in stevia and contributing about 60-70% of total glycosides (Yadav *et al.*, 2011) but it has licorice aftertaste. However, rebaudioside A contributing about 30-40% of total glycosides in stevia and has no aftertaste (Midmore and Rank, 2002). In addition to sugary compounds, stevia extracts also contain a high level of therapeutic properties that plays an imperative role as antioxidant, antimicrobial, antifungal activities and also serves a good source of fiber and minerals (Abou-Arab *et al.*, 2010).

In Malaysia, stevia was introduced in the 1970s. The initial studies on propagation, cultivation and germplasm evaluation were carried out by Tan *et al.* (2008) under Malaysian Agriculture Research and Development Institute (MARDI). From the study, out of 69 genotypes five genotypes were recommended for local cultivation. However, the major problem for stevia cultivation in Malaysia is photoperiod sensitiveness that affects the early flowering which contributes to low leaf yield and stunted plant growth. Several reports from researchers from Universiti Teknologi MARA (UiTM), Universiti Putra Malaysia (UPM) and International Islamic Universiti Malaysia (IIUM) found that no suitable variety for high biomass production in Malaysia (per comm. Osman 2018). Meanwhile, other countries such as China and Japan, are successful in development of suitable stevia varieties for their local cultivation.

Genetically, *S. rebaudiana* B. is a diploid plant with 22 chromosomes. Several species of stevia have similar characteristics but possess different karyotypes caused by the gain or loss of chromatin due to spontaneous mutation (Frederico and Ruas, 1996). Polyploidy is also recorded in *S. rebaudiana* B. through artificial induction by developing lines containing elevated levels of stevioside (Oliveira *et al.*, 2004). The

stevia has been extensively documented as self-incompatible and pollination assisted by insects (Ramesh *et al.*, 2006). Goettemoeller and Ching (1999) reported that fertile seeds were obtained from self-pollination, however, how much seeds collected was unknown and neither was there any verification of the progeny testing. As a whole, stevia is a highly outcrossing plant that is challenging for breeders to develop inbred lines, whereas, high heterozygosity offers variation to develop hybrid variety.

The agronomical practices for cultivation of stevia needs an optimum temperature of 20-28 °C for good growth, clay sandy soil that rich in organic fertilizers with intensive damping, high illumination and the day length of more than 13 hours per day (Soejarto *et al.*, 1982). The local environment condition of Malaysia can provide the plant with all these conditions except the required day length which leads to early flowering. Because the harvesting of stevia aerial parts need to be carried out before flowering as the level of the sweetness decreases as soon as the plants begin to flower. In addition, the production of stevia varieties with a longer vegetative period in the tropical environment requires a depth study on genetic diversity among the available genotypes.

This study was designed with a set of stevia genotypes collected from different sources evaluated for agronomic traits under short day length (<13 hours) and long day length (>13 hours) conditions. The genetic variation among these genotypes was determined using agronomic characteristics and ISSR molecular markers with the following objectives;

- i. To evaluate the agronomic performance of stevia genotypes collected from different sources under short and long day length environmental conditions.
- ii. To assess the genetic diversity among stevia genotypes using ISSR molecular markers.
- iii. To identify superior genotypes for future stevia breeding programs.

REFERENCES

- Abdel-Tawab, F.M., and Allam, A.W.I (2012). Germplasm Introduction, Collection and Conservation. International conference on new role for the world sugar economy in a changed political and economic environment, 10-13 Nov 2012, Aswan Egypt.
- Abdullateef, R.A., and Osman, M. (2011). Effects of stem cutting types, position and hormonal factors on rooting in *Stevia rebaudiana* Bertoni. *Journal of Agricultural Science*, 4(1), 49.
- Abdullateef, R.A., and Osman, M. (2011). Influence of genetic variation on morphological diversity in accessions of *Stevia rebaudiana* Bertoni. *International Journal of Biology*, 3(3), 66.
- Abdullateef, R.A., and Osman, M. (2011). Studies on effects of pruning on vegetative traits in *Stevia rebaudiana* Bertoni (Compositae). *International Journal of Biology*, 4(1), 146.
- Abdullateef, R.A., and Osman, M. B. (2011). Effects of visible light wavelengths on seed germinability in *Stevia rebaudiana* Bertoni. *International Journal of Biology*, 3(4), 83.
- Abdullateef, R.A., Zakaria, N. H., Hasali, N. H., and Osman, M. (2012). Studies on pollen viability and germinability in accessions of *Stevia rebaudiana* Bertoni. *International Journal of Biology*, 4(3), 72.
- Abou-Arab, A.E., Abou-Arab, A.A., and Abu-Salem, M.F. (2010). Physico-chemical assessment of natural sweeteners steviosides produced from *Stevia rebaudiana* Bertoni plant. *African Journal of Food Science*, 4(5), 269-281.
- Ahmad, L.W.Z., Mohamad, A., Osman, M., Zainuddin, Z., and Khari, F.I.M. (2014). Optimization of ISSR markers for DNA fingerprinting in *Stevia rebaudiana* Bertoni.
- Ahmed, M., Dobberstein, R., and Farnsworth, N. (1980). *Stevia rebaudiana*: Use of p-bromophenacyl bromide to enhance ultraviolet detection of water-soluble organic acids (steviolbioside and rebaudioside B) in high-performance liquid chromatographic analysis. *Journal of Chromatography A*, 192(2), 387-393.
- Ali A, Gull I, Naz S, Afghan S (2010). Biochemical investigation during different stages of in vitro propagation of *Stevia rebaudiana*. *Pak. J. Bot*, 42(4), 2827-2837.
- Al-Turki, T., and Basahi, M.A. (2015). Assessment of ISSR based molecular genetic diversity of Hassawi rice in Saudi Arabia. *Saudi Journal of Biological Sciences*, 22(5), 591-599.

- Armizatul, S.A.H., Azhar, K., and Nazrul, R. (2009). Influence of lengthening day on stevia vegetative growth. *Transaction of the Malaysian Society of Plant Physiology*, 18, 17-18.
- Armizatul, S.A.H., Mohd Aziz, R., Wira, A.B., and Azhar, K. (2010). Effects of Night Break on *Stevia rebaudiana*. *Transaction of the Malaysian Society of Plant Physiology*, 19, 14-17.
- Artkins, R.C. (2010). New Diet revolution. Healthier-Harvest 2009.
- Atteh, J., Onagbesan, O., Tona, K., Buyse, J., Decuyper, E., and Geuns, J. (2011). Empleo potencial de *Stevia rebaudiana* en alimentación animal. *Archivos de zootecnia*, 60(229), 133-136.
- Ahmed B, M. Hossain, R. Islam, A. Kumar Saha, A. Mandal. (2011): A review on natural sweetener plant – stevia having medicinal and commercial importance. *Agonomski Glasnik* 75- 91.
- Bouvet, J. M., Fontaine, C., Sanou, H., & Cardi, C. (2004). An analysis of the pattern of genetic variation in *Vitellaria paradoxa* using RAPD markers. *Agroforestry systems*, 60(1), 61-69.
- Beckmann, J., and Soller, M. (1983). Restriction fragment length polymorphisms in genetic improvement: methodologies, mapping and costs. *Theoretical and Applied Genetics*, 67(1), 35-43.
- Blair, M.W., Panaud, O., and McCouch, S.R. (1999). Inter-simple sequence repeat (ISSR) amplification for analysis of microsatellite motif frequency and fingerprinting in rice (*Oryza sativa* L.). *Theoretical and Applied Genetics*, 98(5), 780-792.
- Bondarev, N.I., Sukhanova, M.A., Reshetnyak, O.V., and Nosov, A.M. (2003). Steviol glycoside content in different organs of *Stevia rebaudiana* and its dynamics during ontogeny. *Biologia Plantarum*, 47(2), 261-264.
- Botstein, D., White, R.L., Skolnick, M., and Davis, R.W. (1980). Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *American Journal of Human Genetics*, 32(3), 314.
- Boussaid, M., Benito, C., Harche, M.K., Naranjo, T., and Zedek, M. (2010). Genetic variation in natural populations of *Stipa tenacissima* from Algeria. *Biochemical Genetics*, 48(9-10), 857-872.
- Brandle, J.E., and Rosa, N. (1992). Heritability for yield, leaf: stem ratio and stevioside content estimated from a landrace cultivar of *Stevia rebaudiana*. *Canadian Journal of Plant Science*, 72(4), 1263-1266.

- Brandle, J.E., Starratt, A.N., and Gijzen, M. (1998). *Stevia rebaudiana*: Its agricultural, biological, and chemical properties. *Canadian Journal of Plant Science*, 78(4), 527-536.
- Brandle, J. E., Starratt, A. N., & Gijzen, M. (2000). *Stevia rebaudiana*. Its biological, chemical and agricultural properties. Agriculture and AgriFood Canada. *Southern Crop Protection and Food Research Centre*, 1391 Sandford St., London, Ontario N5V 4T3. Retrived from www.lni.Unpi.it/stevia/stevia/stevia0005.htm.
- Bülent, K.Ö.S.E. (2014). Effect of light intensity and temperature on growth and quality parameters of grafted vines. *Notulae Botanicae. Horti Agrobotanici Cluj-Napoca*, 42(2), 507.
- Burns, N., Grimwade, B., Ross-Macdonald, P.B., Choi, E.-Y., Finberg, K., Roeder, G.S., and Snyder, M. (1994). Large-scale analysis of gene expression, protein localization, and gene disruption in *Saccharomyces cerevisiae*. *Genes and Development*, 8(9), 1087-1105.
- Chan, P., Xu, D.Y., Liu, J.C., Chen, Y.J., Tomlinson, B., Huang, W.P., and Cheng, J.T. (1998). The effect of stevioside on blood pressure and plasma catecholamines in spontaneously hypertensive rats. *Life sciences*, 63(19), 1679-1684.
- Cho, R.J., Mindrinos, M., Richards, D.R., Sapolsky, R.J., Anderson, M., Drenkard, E., and Theologis, A. (1999). Genome-wide mapping with biallelic markers in *Arabidopsis thaliana*. *Nature Genetics*, 23(2), 203.
- Choi, K.-H., and Kim, K.-J. (2009). Applications of transposon-based gene delivery system in bacteria. *Journal of Microbiology and Biotechnology*, 19(3), 217-228.
- Condit, R., and Hubbell, S.P. (1991). Abundance and DNA sequence of two-base repeat regions in tropical tree genomes. *Genome*, 34(1), 66-71.
- Copes, D.L., and Mandel, N.L. (2000). Effects of IBA and NAA treatments on rooting Douglas-fir stem cuttings. *New Forests*, 20(3), 249-257.
- Crossa, J, and Franco, J. (2004). Statistical methods for classifying genotypes. *Euphytica*, 137(1), 19-37.
- Cruz, C.D., Ferreira, F.M., and Pessoni, L.A. (2011). Biometria aplicada ao estudo da diversidade genética. *Visconde do Rio Branco: Suprema*.
- Davierwala, A., Chowdari, K., Kumar, S., Reddy, A., Ranjekar, P., and Gupta, V. (2000). Use of three different marker systems to estimate genetic diversity of Indian elite rice varieties. *Genetica*, 108(3), 269-284.

- De Klerk, G.J. (2002). Rooting of microcuttings: theory and practice. *In Vitro Cellular and Developmental Biology-Plant*, 38(5), 415-422.
- Devos, K.M., and Gale, M.D. (1997). Comparative genetics in the grasses. *Plant Molecular Biology*, 35(1), 3-15.
- Dos Santos, L.F., de Oliveira, E.J., dos Santos Silva, A., de Carvalho, F.M., Costa, J.L., and Pádua, J.G. (2011). ISSR markers as a tool for the assessment of genetic diversity in Passiflora. *Biochemical Genetics*, 49(7-8), 540-554.
- Duke, J. (1993). *Stevia rebaudiana*. Duke J. ed. *CRC Handbook of Alternative Cash Crops*. CRC Press, 422-424.
- Dwivedi, R.S. (1999). Unnurtured and untapped super sweet non sacchariferous plant species in India. *Current Science*, 76(11), 1454-1461.
- Ferrazzano, G.F., Cantile, T., Alcidi, B., Coda, M., Ingenito, A., Zarrelli, A., and Pollio, A. (2015). Is *Stevia rebaudiana* Bertoni a non cariogenic sweetener? A review. *Molecules*, 21(1), 38.
- Ferri, L.A., Alves- Do- Prado, W., Yamada, S.S., Gazola, S., Batista, M.R., and Bazotte, R.B. (2006). Investigation of the antihypertensive effect of oral crude stevioside in patients with mild essential hypertension. *Phytotherapy Research*, 20(9), 732-736.
- Francisco F., Pereira, G. P., Machado, M. P., Kanis, L. A., & Deschamps, C. (2018). Characterization of *Stevia rebaudiana* Bertoni Accessions Cultived in Southern Brazil. *Journal of Agricultural Science*, 10(3), 353.
- Franco, J., Crossa, J., Ribaut, J.M., Bertran, J., Warburton, M.L., and Khairallah, M. (2001). A method for combining molecular markers and phenotypic attributes for classifying plant genotypes. *Theoretical and Applied Genetics*, 103(6-7), 944-952.
- Franco J, Crossa J, Warburton ML, Taba S (2006) Sampling strategies for conserving maize diversity when forming core subsets using genetic markers. *Crop Science* 46:854–864
- Frederico, A.P., Ruas, P.M., Marin-Morales, M.A., Ruas, C.F., and Nakajima, J.N. (1996). Chromosome studies in some *Stevia*. Cav.(*Compositae*) species from Southern Brazil. *Brazilian Journal of Genetics*, 19(4), 605-609.
- Gawel-Bęben, K., Bujak, T., Nizioł-Lukaszewska, Z., Antosiewicz, B., Jakubczyk, A., Karaś, M., and Rybczyńska, K. (2015). *Stevia rebaudiana* Bert. leaf extracts as a multifunctional source of natural antioxidants. *Molecules*, 20(4), 5468-5486.

- Genger, R.K., Brown, A.H., Nesbitt, K., and Burdon, J.J. (1999). Molecular markers for wild barley-derived scald resistance genes. In *Proc. of the 9th Australian Barley Tech. Symposium* (pp. 1-5).
- Godwin, I.D., Aitken, E.A., and Smith, L.W. (1997). Application of inter simple sequence repeat (ISSR) markers to plant genetics. *Electrophoresis*, 18(9), 1524-1528.
- Goettemoeller, J., and Ching, A. (1999). Seed germination in *Stevia rebaudiana*. *Janick (ed.)*, 510-511.
- Gonçalves, L., Rodrigues, R., Amaral Júnior, A. D., Karasawa, M., & Sudré, C. P. (2009). Heirloom tomato gene bank: assessing genetic divergence based on morphological, agronomic and molecular data using a Ward-modified location model. *Genetics and Molecular Research*, 8(1), 364-374.
- Gower, J.C. (1971). A general coefficient of similarity and some of its properties. *Biometrics*, 857-871.
- Goyal, S.K., and Goyal, R.K. (2010). *Stevia (Stevia rebaudiana)* a bio-sweetener: a review. *International Journal of Food Sciences and Nutrition*.
- Griffin, T.J., and Smith, L.M. (2000). Single-nucleotide polymorphism analysis by MALDI-TOF mass spectrometry. *Trends in Biotechnology*, 18(2), 77-84.
- Gupta, E., Purwar, S., Sundaram, S., and Rai, G.K. (2013). Nutritional and therapeutic values of *Stevia rebaudiana*: A review. *Journal of Medicinal Plants Research*, 7(46), 3343-3353.
- Guruchandran, V., and Sasikumar, C. (2013). Organogenic plant regeneration via callus induction in *Stevia rebaudiana* Bert. *International Journal of Current Microbiology and Applied Sciences*, 2(2), 56-61.
- Halimatun, S.O., Mohamad, O., and Zarina, Z. (2015). Morphological assessment of *Stevia rebaudiana* Bertoni accessions in IIUM's germplasm as initial material for stevia breeding. *Australian Journal of Basic and Applied Sciences*, 9(25 Special), 1-9.
- Han, Y.C., Teng, C.Z., Zhong, S., Zhou, M.Q., Hu, Z.L., and Song, Y.C. (2007). Genetic variation and clonal diversity in populations of *Nelumbo nucifera* (*Nelumbonaceae*) in central China detected by ISSR markers. *Aquatic Botany*, 86(1), 69-75.
- Haq, I.U., Touqeer, A., Hafiz, I.A., and Abbasi, N.A. (2009). Influence of microcutting sizes and IBA concentrations on In vitro rooting of olive cv. 'Dolce Agogia'. *Pakistan Journal Botany*, 41(3), 1213-1222.

- He, F., Kang, D., Ren, Y., Qu, L., Zhen, Y., and Gu, H. (2007). Genetic diversity of the natural populations of *Arabidopsis thaliana* in China. *Heredity*, 99(4), 423-431.
- Heikal, A.H., Badawy, O.M., and Hafez, A.M. (2008). Genetic relationships among some *Stevia* (*Stevia rebaudiana* Bertoni) accessions based on ISSR analysis. *Research Journal of Cell and Molecular Biology*, 2(1), 1-5.
- Horecka, J., and Jigami, Y. (2000). Identifying tagged transposon insertion sites in yeast by direct genomic sequencing. *Yeast*, 16(10), 967-970.
- Huang, Y., Zhang, C.Q., and LI, D.Z. (2009). Low genetic diversity and high genetic differentiation in the critically endangered *Omphalogramma souliei* (*Primulaceae*): Implications for its conservation. *Journal of Systematics and Evolution*, 47(2), 103-109.
- Huda, M., Ahmed, A., Mandal, C., Alam, K.A., Reza, S.H., and Wadud, A. (2007). In vitro morphogenic responses of different explants of *Stevia* (*Stevia rebaudiana* Bert). *International Journal Agriculture Research*, 12, 1006-1013.
- Ingle, M.R. (2008). Effect of growth regulators and environments on rooting of stevia cuttings (*Stevia rebaudiana* Bertoni) (Doctoral dissertation, UAS, Dharwad).
- Jala, A. (2011). Effects of different light treatments on the germination of *Nepenthes mirabilis*. *International Transaction Journal of Engineering, Management, and Applied Sciences and Technologies*, 2(1), 083-091.
- Janke, R. (2004). Farming a few Acres of Herbs: *Stevia*. *Kansas State University*. [Online] Available: <http://www.oznet.ksu.edu>.
- Jeppesen, P.B., Gregersen, S., Alstrup, K.K., and Hermansen, K. (2002). Stevioside induces antihyperglycaemic, insulinotropic and glucagonostatic effects in vivo: studies in the diabetic Goto-Kakizaki (GK) rats. *Phytomedicine*, 9(1), 9-14.
- Jeppesen, P.B., Gregersen, S., Rolfsen, S.E.D., Jepsen, M., Colombo, M., Agger, A., and Hermansen, K. (2003). Antihyperglycemic and blood pressure-reducing effects of stevioside in the diabetic Goto-Kakizaki rat. *Metabolism*, 52(3), 372-378.
- Joshi, S., Gupta, V., Aggarwal, R., Ranjekar, P., and Brar, D. (2000). Genetic diversity and phylogenetic relationship as revealed by inter simple sequence repeat (ISSR) polymorphism in the genus *Oryza*. *Theoretical and Applied Genetics*, 100(8), 1311-1320.
- Kafle, G.G. (2011). Some Studies on the Physiology of *Stevia Rebaudiana* Bertoni. *Faculty of Sciences, Engineering and Health, CQ University Australia*.

- Karimi, M., Hashemi, J., Ahmadi, A., Abbasi, A., Pompeiano, A., Tavarini, S., ... & Angelini, L. G. (2015). Opposing effects of external gibberellin and daminozide on stevia growth and metabolites. *Applied Biochemistry and Biotechnology*, 175(2), 780-791.
- Katayama, O., Sumida, T., Hayashi, H., and Mitsuhashi, H. (1976). The practical application of stevia and research and development data. ISU Company, Japan, 747.
- Kim, M.S., Moore, P.H., Zee, F., Fitch, M.M., Steiger, D.L., Manshardt, R.M., and Ming, R. (2002). Genetic diversity of *Carica papaya* as revealed by AFLP markers. *Genome*, 45(3), 503-512.
- Kinghorn, A.D., Soejarto, D.D., Kennelly, E.J., Kim, D.S., Ohtani, K., and Yamasaki, K. (2002). The genus *Stevia*. *Med. Arom. Plants-Ind. Profiles*, 19, 52-64.
- Kochetov, A. (2004). Growth and yield of *Stevia Rebaudiana* Bertoni under different conditions of lighting. Paper presented at the Agriculture Congress 2004" Innovation towards modernized agriculture".
- Lakshmi, M., Parani, M., Rajalakshmi, S., and Parid, A. (2002). Analysis of species relationship among seven small millets using molecular markers. *Journal of Plant Biochemistry and Biotechnology*, 11(2), 85-91.
- Lamkey, K.R., and Lee, M. (1993). Quantitative genetics, molecular markers, and plant improvement.
- LaPierre, L.M. (2001). Vegetative propagation of *Cecropia obtusifolia* (Cecropiaceae). *Revista De Biología Tropical*, 49(3-4), 973-976.
- Lata, C., Gupta, S., and Prasad, M. (2013). Foxtail millet: a model crop for genetic and genomic studies in bioenergy grasses. *Critical Reviews in Biotechnology*, 33(3), 328-343.
- LeBude, A.V., Blazich, F.A., Walker, L.C., and Robinson, S.M. (2008). Seed germination of two populations of *Rhododendron vaseyi*: Influence of light and temperature. *Journal of Environmental Horticulture*, 26(4), 217-221.
- Lefebvre, V., Goffinet, B., Chauvet, J.C., Caromel, B., Signoret, P., Brand, R., and Palloix, A. (2001). Evaluation of genetic distances between pepper inbred lines for cultivar protection purposes: comparison of AFLP, RAPD and phenotypic data. *Theoretical and Applied Genetics*, 102(5), 741-750.
- Lemus-Mondaca, R., Vega-Gálvez, A., Zura-Bravo, L., and Ah-Hen, K. (2012). *Stevia rebaudiana* Bertoni, source of a high-potency natural sweetener: A comprehensive review on the biochemical, nutritional and functional aspects. *Food Chemistry*, 132(3), 1121-1132.

- Lin, J., Ma, J., Ambrose, M., and Kuo, J. (1997). Chemiluminescent detection of AFLP fingerprints. *Focus*, 19(2), 36-38.
- Lumaret, R. (1988). Adaptive strategies and ploidy levels. *ACTA OECOL.(OECOL. PLANT.)*, 9(1), 83-93.
- Madan S, Ahmad S, Singh G, Kohli K, Kumar Y, Singh R, Garg M (2010): *Stevia rebaudiana* (Bert.) Bertoni-a review. *Indian J Nat Prod Resour.*, 1 (3): 267-286.
- Maiti RK, Purohit SS 2008. Relative sweetness. In: *stevia-a miracle plant for human health. Jodhpur: Agrobios India*, P. 64.
- Manimekalai, R., Nagarajan, P., Bharathi, M., Karun, A., Kumaran, P.M., and Parthasarathy, V.A. (2007). Molecular diversity among south east Asian coconut (*Cocos nudfera* L.) germplasm accessions based on ISSR Markers. *Recent Trends in Horticultural Biotechnology. New India Publishing Agency*, 479-484.
- Marsolais, A.A., Brandle, J., and Sys, E.A. (1998). Stevia plant named 'RSIT 94-751' U.S. Patent Application No. 08/657,463.
- Melis, M.S. (1999). Effects of chronic administration of *Stevia rebaudiana* on fertility in rats. *Journal of Ethnopharmacology*, 67(2), 157-161.
- Metivier, J., and Viana, A.M. (1979). The effect of long and short day length upon the growth of whole plants and the level of soluble proteins, sugars, and stevioside in leaves of *Stevia rebaudiana* Bert. *Journal of Experimental Botany*, 30(6), 1211-1222.
- Midmore, D.J., and Rank, A.H. (2002). A new rural industry stevia to replace imported chemical sweeteners, RIRDC Pub No. W02/22, 55.
- Morita, T., and Bu, Y. (2000). U.S. Patent No. 6,031,157. Washington, DC: U.S. Patent and Trademark Office.
- Nagaraju, J., Kathirvel, M., Kumar, R.R., Siddiq, E., and Hasnain, S.E. (2002). Genetic analysis of traditional and evolved Basmati and non-Basmati rice varieties by using fluorescence-based ISSR-PCR and SSR markers. *Proceedings of the National Academy of Sciences*, 99(9), 5836-5841.
- Nanda, K.K., and Hamner, K.C. (1962). Investigations on the effect of light break on the nature of the endogenous rhythm in the flowering response of *Biloxi soybean*. *Planta*, 58(2), 164-174.
- Niemi, K., Julkunen-Tiitto, R., Tegelberg, R., and Häggman, H. (2005). Light sources with different spectra affect root and mycorrhiza formation in Scots pine in vitro. *Tree Physiology*, 25(1), 123-128.

- Nieto-Lopez, R.M., and Blake, T.K. (1994). Russian wheat aphid resistance in barley: inheritance and linked molecular markers. *Crop Science*, 34(3), 655-659.
- Oliveira, V.M. d., Forni-Martins, E.R., Magalhães, P.M., and Alves, M.N. (2004). Chromosomal and morphological studies of diploid and polyploid cytotypes of *Stevia rebaudiana* Bertoni (Eupatorieae, Asteraceae). *Genetics and Molecular Biology*, 27(2), 215-222.
- Olson, M., Hood, L., Cantor, C., and Botstein, D. (1989). A common language for physical mapping of the human genome. *Science*, 245(4925), 1434-1435.
- Osman, M., Samsudin, N.S., Faruq, G., and Nezhadahmadi, A. (2013). Factors affecting microcuttings of *Stevia* using a mist-chamber propagation box. *The Scientific World Journal*, 1:1-10.
- Pal, P. K., Mahajan, M., Prasad, R., Pathania, V., Singh, B., & Ahuja, P. S. (2015). Harvesting regimes to optimize yield and quality in annual and perennial *Stevia rebaudiana* under sub-temperate conditions. *Industrial Crops and Products*, 65, 556-564.
- Paris, M., Jones, M.G., and Eglinton, J.K. (2002). Genotyping single nucleotide polymorphisms for selection of barley β -amylase alleles. *Plant Molecular Biology Reporter*, 20(2), 149-159.
- Parsons, P. (2003). *Stevia*-Too good to be approved. *Organic NZ*, 50(3), 26.
- Patel, D.K. (2015). Stem cutting propagation for ex-situ conservation of *Stevia rebaudiana* Bertoni in herbal garden. *European Journal of Biotechnology and Bioscience*, 9, 35-42.
- Paterson, I.D., and Zachariades, C. (2013). ISSRs indicate that *Chromolaena odorata* invading southern Africa originates in Jamaica or Cuba. *Biological Control*, 66(2), 132-139.
- Paterson, I.D., Downie, D.A., and Hill, M.P. (2009). Using molecular methods to determine the origin of weed populations of *Pereskia aculeata* in South Africa and its relevance to biological control. *Biological Control*, 48(1), 84-91.
- Patnaik, D., and Khurana, P. (2001). Wheat biotechnology: A mini review. *Electronic Journal of Biotechnology*, 4(2), 7-8.
- Pillay, M., and Myers, G. (1999). Genetic diversity in cotton assessed by variation in ribosomal RNA genes and AFLP markers. *Crop Science*, 39(6), 1881-1886.
- Pon, S.Y., Lee, W.J., and Chong, G.H. (2015). Textural and rheological properties of stevia ice cream. *International Food Research Journal*, 22(4).

- Powell, W., Morgante, M., Andre, C., Hanafey, M., Vogel, J., Tingey, S., and Rafalski, A. (1996). The comparison of RFLP, RAPD, AFLP and SSR (microsatellite) markers for germplasm analysis. *Molecular Breeding*, 2(3), 225-238.
- Rajasekaran, T., Giridhar, P., and Ravishankar, G.A. (2007). Production of steviosides in ex vitro and in vitro grown *Stevia rebaudiana* Bertoni. *Journal of the Science of Food and Agriculture*, 87(3), 420-424.
- Ranjan, R., Jaiswal, J., and Jena, J. (2011). Stevia as a natural sweetener. *International Journal of Research in Pharmacy and Chemistry*, 1(4), 1199-1202.
- Reddy, M.P., Sarla, N., and Siddiq, E. A. (2002). Inter simple sequence repeat (ISSR) polymorphism and its application in plant breeding. *Euphytica*, 128(1), 9-17.
- Reiter, R.S., Williams, J., Feldmann, K. A., Rafalski, J. A., Tingey, S. V., and Scolnik, P. A. (1992). Global and local genome mapping in *Arabidopsis thaliana* by using recombinant inbred lines and random amplified polymorphic DNAs. *Proceedings of the National Academy of Sciences*, 89(4), 1477-1481.
- Robinson, B.L. (1930). Observations on the genus *Stevia*. *Contributions from the Gray Herbarium of Harvard University*, 36-58.
- Rodríguez, J., Rodríguez, P., González, M.E., and Martínez-Gómez, P. (2010). Molecular characterization of *Cuban endemism Carica cubensis* Solms using random amplified polymorphic DNA (RAPD) markers. *Agricultural Sciences*, 1(03), 95.
- Rohlf, F. (2002). NTSYS-pc: numerical taxonomy system, version 2.1 Exeter Publishing. Ltd., Setauket, New York, USA.
- Roldán-Ruiz, I., Dendauw, J., Van Bockstaele, E., Depicker, A., and De Loose, M. (2000). AFLP markers reveal high polymorphic rates in ryegrasses (*Lolium* spp.). *Molecular Breeding*, 6(2), 125-134.
- Runkle, E. S., and Heins, R.D. (2001). Specific functions of red, far red, and blue light in flowering and stem extension of long-day plants. *Journal of the American Society for Horticultural Science*, 126(3), 275-282.
- Saini, N., Jain, N., Jain, S., and Jain, R.K. (2004). Assessment of genetic diversity within and among Basmati and non-Basmati rice varieties using AFLP, ISSR and SSR markers. *Euphytica*, 140(3), 133-146.
- Sarla, N., Bobba, S., and Siddiq, E. (2003). ISSR and SSR markers based on AG and GA repeats delineate geographically diverse *Oryza nivara* accessions and reveal rare alleles. *Current Science*, 84: 683-690.

- Sarwat, M. (2012). ISSR: a reliable and cost-effective technique for detection of DNA polymorphism. *Plant DNA Fingerprinting and Barcoding: Methods and Protocols*, 103-121.
- Sengupta, S., Das, B., Prasad, M., Acharyya, P., and Ghose, T. K. (2013). A comparative survey of genetic diversity among a set of Caricaceae accessions using microsatellite markers. *SpringerPlus*, 2(1), 345.
- Shaffert, E.E. (1994). Chebotar'A.A. (1994). Structure, topography and ontogeny of *Stevia rebaudiana* (Asteraceae) trichomes. *Bot. Zhurn*, 79(4), 38-48.
- Sharma, N., Kaur, R., and Era, V. (2016). Potential of RAPD and ISSR markers for assessing genetic diversity among *Stevia rebaudiana* Bertoni accessions.
- Sharma, N., Kaushal, N., Chawla, A., Mohan, M., Sethi, A., and Sharma, Y. (2006). *Stevia rebaudiana*-A review. *Agrobios Newslett*, 5(7), 46-48.
- Sharma, S., Knox, M., and Ellis, T.N. (1996). AFLP analysis of the diversity and phylogeny of Lens and its comparison with RAPD analysis. *Theoretical and Applied Genetics*, 93(5-6), 751-758.
- Shock, C.C. (1982). Experimental cultivation of Rebaudi's stevia in California. *Univ. California, Davis Agronomy Progress Report* 122.
- Šic Žlabur, J., Voća, S., Dobričević, N., Ježek, D., Bosiljkov, T., Brnčić, M. (2013). *Stevia rebaudiana* Bertoni - A Review of Nutritional and Biochemical Properties of Natural Sweetener. *Agriculturae Conspectus Scientificus*, 78(1), 25-30.
- Singh, S.D., and Rao, G.P. (2005). Stevia: the herbal sugar of 21st century. *Sugar Tech*, 7(1), 17-24.
- Soejarto, D.D., Kinghorn, A.D., and Farnsworth, N.R. (1982). Potential sweetening agents of plant origin. III. Organoleptic evaluation of *Stevia* leaf herbarium samples for sweetness. *Journal of Natural Products*, 45(5), 590-599.
- Southern, E.M. (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *Journal of Molecular Biology*, 98(3), 503-535.
- Spielmeyer, W., Sharp, P.J., and Lagudah, E.S. (2003). Identification and validation of markers linked to broad-spectrum stem rust resistance gene Sr2 in wheat (*Triticum aestivum* L.). *Crop Science*, 43(1), 333-336.

- Srimaroeng, C., Chatsudthipong, V., Aslamkhan, A.G., and Pritchard, J.B. (2005). Transport of the natural sweetener, stevioside, and its aglycone, steviol, by hOAT1 (SLC22A6) and hOAT3 (SLC22A8). *JPET Fast Forward. Laboratory of Pharmacology and Chemistry, National Institute of Environmental Health Sciences, Research Triangle Park, NC*, 10(104.080366).
- Staub, J.E., Serquen, F.C., and Gupta, M. (1996). Genetic markers, map construction, and their application in plant breeding. *Hort Science*, 31(5), 729-741.
- Sumida, T (1980)., Studies on *Stevia rebaudiana* Bertoni as a new possible crop for sweetening resources in Japan. *Journal of the Central Agricultural Experiment Station*, 31: p. p 1-71.
- Sundaresan, V. (1996). Horizontal spread of transposon mutagenesis: new uses for old elements. *Trends in Plant Science*, 1(6), 184-190.
- Taheri, S., Abdullah, T. L., Abdullah, N. A. P., & Ahmad, Z. (2012). Genetic relationships among five varieties of *Curcuma alismatifolia* (Zingiberaceae) based on ISSR markers. *Genetics Molecular Research*, 11(3), 3069-3076.
- Takahashi, K., Matsuda, M., Ohashi, K., Taniguchi, K., Nakagomi, O., Abe, Y., Shigeta, S. (2001). Analysis of anti-rotavirus activity of extract from *Stevia rebaudiana*. *Antiviral Research*, 49(1), 15-24.
- Talbert, L., Blake, N., Chee, P., Blake, T., and Magyar, G.M. (1994). Evaluation of "sequence-tagged-site" PCR products as molecular markers in wheat. *TAG Theoretical and Applied Genetics*, 87(7), 789-794.
- Talha, M., Hussain, I., Ullah, R., and Khan, L. (2012). Analysis of stevioside in *Stevia rebaudiana*. *Journal of Medicinal Plants Research*, 6(11), 2216-2219.
- Tan, S.L., Muhammad Ghawas, M., Mohamad Najib, M.Y., and Zawayi, M. (2008). Preliminary evaluation and selection of stevia under Malaysian conditions. *Journal Tropical Agriculture and Food Science*, 36(2), 000-000.
- Taylor, C., and Barker, N. (2012). Species limits in *Vachellia* (Acacia) karroo (*Mimosoideae: Leguminosae*): Evidence from automated ISSR DNA "fingerprinting". *South African Journal of Botany*, 83, 36-43.
- Taylor, L. (2005). *The Healing Power of Natural Herbs*: Garden City Park, NY: Square One Publishers, Inc. pp(excerpted at web link). ISBN 0-7570-0144-0. <http://rain-tree.com/stevia.htm>.
- Thomas, B., and Vince-Prue, D. (1996). Photoperiodism in plants. *Academic Press*.

- Thomas, J.E., and Glade, M.J. (2010). Stevia: it's not just about calories. *Benefits*, 35, 36.
- Tragoonrung, S., Kanazin, V., Hayes, P., and Blake, T. (1992). Sequence-tagged-site-facilitated PCR for barley genome mapping. *TAG Theoretical and Applied Genetics*, 84(7), 1002-1008.
- Upadhyay, A., Jayadev, K., Manimekalai, R., and Parthasarathy, V. (2004). Genetic relationship and diversity in Indian coconut accessions based on RAPD markers. *Scientia Horticulturae*, 99(3), 353-362.
- Valio, Rocha, R. F. (1977). Effect of photoperiod and growth regulator on growth and flowering of *Stevia rebaudiana Bertoni*. *Japanese Journal of Crop Science*, 46(2), 243-248.
- Van Deynze, A., Sorrells, M., Park, W., Ayres, N., Fu, H., Cartinhour, S., McCouch, S. (1998). Anchor probes for comparative mapping of grass genera. *TAG Theoretical and Applied Genetics*, 97(3), 356-369.
- Venkatachalam, L., Sreedhar, R. V., and Bhagyalakshmi, N. (2008). The use of genetic markers for detecting DNA polymorphism, genotype identification and phylogenetic relationships among banana cultivars. *Molecular Phylogenetics and Evolution*, 47(3), 974-985.
- Vijayanand, V., Senthil, N., Vellaikumar, S., and Paramathma, M. (2009). Genetic diversity of Indian *Jatropha* species as revealed by morphological and ISSR markers. *Journal of Crop Science and Biotechnology*, 12(3), 115-120.
- Vos, P., Hogers, R., Bleeker, M., Reijans, M., Lee, T.V.D., Hornes, M., Kuiper, M. (1995). AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research*, 23(21), 4407-4414.
- Walker, L.C., LeBude, A.V., Blazich, F.A., and Conner, J.E. (2006). Seed germination of pinkshell azalea (*Rhododendron vaseyi*) as influenced by light and temperature. *NCSU Phytotron*, 61.
- Waugh, R., and Power, W. (1992). Using RAPD markers for crop improvement. *Trends in Biotechnology*, 10, 186-191.
- Williams, J.G., Kubelik, A.R., Livak, K.J., Rafalski, J.A., and Tingey, S.V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research*, 18(22), 6531-6535.
- Wolff, K., and Morgan-Richards, M. (1998). PCR markers distinguish *Plantago* major subspecies. *Theoretical and Applied Genetics*, 96(2), 282-286.

- Wongsawad, P., and Peerapornpisal, Y. (2014). Molecular identification and phylogenetic relationship of green algae, *Spirogyra ellipsospora* (*Chlorophyta*) using ISSR and *rbcL* markers. *Saudi Journal of Biological Sciences*, 21(5), 505-510.
- Yadav, A.K., Singh, S., Dhyani, D., and Ahuja, P.S. (2011). A review on the improvement of *Stevia rebaudiana* Bertoni. *Canadian Journal of Plant Science*, 91(1), 1-27.
- Yadav, S., Jain, S., Jain, V., and Jain, R.K. (2007). Genetic analysis of CMS, restorer, hybrid and open-pollinated genotypes of Indian pearl millet [*Pennisetum glaucum* (L.) R. Br.] using ISSR markers. *Indian Journal Biotechnolol*, 6: 205–209.
- Yahya, A., Nor'ain, A.R., and Ab Kahar, S. (2009). Adventitious rooting in microcuttings of selected indigenous landscape trees of Malaysia. *International Journal of Agriculture and Biology*, 11(4), 419-424.
- Yamaguchi, S. (2008). Gibberellin metabolism and its regulation. *Annu. Rev. Plant Biology* 59, 225-251.
- Yao, Y., Ban, M., and Brandle, J. (1999). A genetic linkage map for *Stevia rebaudiana*. *Genome*, 42(4), 657-661.
- Yeh, F.C., Yang, R.C., Boyle, T.B., Ye, Z.H., and Mao, J.X. (1997). POPGENE, the user-friendly shareware for population genetic analysis. *Molecular Biology and Biotechnology Centre, University of Alberta, Canada*, 10, 295-301.
- Huang YS, Guo AG, Qian Y, Chen LY, Gu HF (1995) Studies on the variation of steviosides content and selection of type R-A in *Stevia rebaudiana*. *Journal of Plant Research Environment* 4:28–32.
- Zaidan, L.B., Dietrich, S.M., and Felipe, G. (1980). Effect of photoperiod on flowering and stevioside content in plants of *Stevia rebaudiana* Bertoni. *Japanese Journal of Crop Science*, 49(4), 569-574.
- Zietkiewicz, E., Rafalski, A., and Labuda, D. (1994). Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics*, 20(2), 176-183.