

UNIVERSITI PUTRA MALAYSIA

EVALUATION OF WATER QUALITY USING PATTERN RECOGNITION TECHNIQUE IN MELAKA RIVER BASIN, MALAYSIA

ANG KEAN HUA

FPAS 2018 12



EVALUATION OF WATER QUALITY USING PATTERN RECOGNITION TECHNIQUE IN MELAKA RIVER BASIN, MALAYSIA

By

ANG KEAN HUA

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of Philosophy

May 2018

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

EVALUATION OF WATER QUALITY USING PATTERN RECOGNITION TECHNIQUE IN MELAKA RIVER BASIN, MALAYSIA

By

ANG KEAN HUA

May 2018

Chair Faculty : Faradiella Mohd Kusin, PhD : Environmental Studies

The unorganised expansion and uncontrolled urbanisation development had led to environmental issues involved with river water pollution that occurred within the Melaka River basin. There are three objectives study to solve the water quality of the Melaka River basin through (1) to identify the potential sources of Melaka River pollution in correlation with the temporal land use classification, (2) to determine the land use land cover changes in 2015 and predict the future land use land cover for 2022, and (3) to verify pattern recognition based on the land use classification between 2015 and 2022 in the determination of pollutant sources within the Melaka River. The results indicated HCA have two cluster areas (C1 and C2). The DA indicated 12 variables were found to be the most significant parameters with a high variation in the spatial distribution. The PCA found that the C1 was attributed to contamination from the agricultural and residential activities; while the C2 included the pollution sources from the agricultural, residential, industrial, animal husbandry, and sewage treatment activities. In the Pearson correlation analysis and ANOVA analysis shown Vegetation Area (VA), Non-Industrial Area (NIA), Industrial Area (IA), Open Space Area (OSA), and Farming Area (FA) were correlated with majority of the physico-chemical and biological variables. LULC incorporated with CA-Markov chain model analysis showed NIA (67.50%), IA (29.15%) and FA (3.35%) were increased from 2001 to 2015 and 2022; while the VA (89.59%), OSA (3.32%) and WB (7.08%) continued to decrease. In GIS, the LISA analysis indicates 2015 showed the clustered area of C1 with S1 as the main contributor; while C2 of S5 as the main contributor. The Moran I analysis from high-to-low pollutant sources are NIA (0.80) > VA (0.59) > IA (0.42) > OSA (0.40). In 2022, the contamination from high-to-low are VA (0.59) > NIA (0.58) > IA (0.50) > OSA (0.41) in the Melaka River. The hotspot analysis for year 2015 indicated C1 of S1 as the main contributor; while C2 of S5 as main contributor. General G-statistic analysis from high-to-low are IA (1.099×10^{-3}) > NIA (5.39×10^{-4}) > VA (6.9×10^{-5}) . In 2022, the hotspot analysis in C1 indicated S7 as the main contributor; while C2 remains the same as S5 which is the main contributor to have pollutant sources from high-to-low are IA $(3.012 \times 10^{-3}) > NIA (5.95 \times 10^{-4}) > VA (7.6 \times 10^{-5})$. In conclusion, Moran I and general G-statistic had gained support especially in benefiting from the PCA in recognising the pattern of pollutant sources (e.g. land use classes) with precise details. Simultaneously, LISA and hotspot analysis have an advantage over HCA by recognising the clustered area more specifically, which was based on the pattern of pollutant sources.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

PENILAIAN KUALITI AIR MENGGUNAKAN TEKNIK CORAK PENGIKTIRAFAN DI LEMBANGAN SUNGAI MELAKA, MALAYSIA

Oleh

ANG KEAN HUA

Mei 2018

Pengerusi Fakulti : Faradiella Mohd Kusin, PhD : Pengajian Alam Sekitar

Pembangunan perbandaran yang tidak teratur dan tidak terkawal telah membawa kepada isu-isu alam sekitar yang melibatkan pencemaran air sungai yang berlaku di dalam lembangan Sungai Melaka. Oleh itu, penyelesaian terhadap kualiti air sungai di lembangan Sungai Melaka melalui kajian objektif seperti (1) untuk mengenal pasti potensi sumber pencemar Sungai Melaka dalam korelasi dengan klasifikasi penggunaan tanah sementara, (2) untuk menentukan perubahan guna tanah 2015 dan meramalkan masa depan guna tanah pada 2022, dan (3) untuk mengesahkan pengiktirafan corak berdasarkan klasifikasi guna tanah antara 2015 dan 2022 dalam menentukan sumber pencemar di Sungai Melaka. Dapatan kajian menunjukkan HCA mempunyai dua kawasan cluster (C1 dan C2). DA menunjukkan 12 pembolehubah didapati merupakan parameter yang paling ketara dengan variasi yang tinggi dalam taburan spatial. PCA menunjukkan C1 dikaitkan dengan pencemaran dari aktiviti pertanian dan kediaman; manakala C2 termasuk sumber pencemaran dari aktiviti pertanian, kediaman, perindustrian, penternakan, dan rawatan kumbahan. Dalam analisis korelasi Pearson dan analisis varians (ANOVA) menunjukkan Vegetation Area (VA), Non-Industrial Area (NIA), Industrial Area (IA), Open Space Area (OSA), and Farming Area (FA) mempunyai perhubungan signifikan dengan majoriti pembolehubah fiziko-kimia dan biologi. Seterusnya, LULC menggabungkan analisis model rantaian CA-Markov menunjukkan NIA (67.50%), IA (29.15%), dan FA (3.35%) telah meningkat dalam tempoh 15 tahun (2001 hingga 2015) dan juga luas guna tanah masa depan (2022); manakala VA (89.59%), OSA (3.32%), dan WB (7.08%) pula terus berkurangan. Dalam GIS, analisis LISA tahun 2015 menunjukkan kawasan kluster C1 dengan S1 adalah penyumbang utama; manakala C2 adalah S5 sebagai penyumbang utama. Analisis Moran I dari sumber-sumber pencemar dari tinggi-ke-rendah adalah NIA (0.80) > VA (0.59) > IA (0.42) > OSA (0.40). Pada 2022, pencemaran dari tinggi-ke-rendah adalah VA (0.59) > NIA (0.58) > IA (0.50) > OSA (0.41) di Sungai Melaka. Analisis hotspot tahun

2015 menunjukkan C1 dari S1 adalah sebagai penyumbang utama; manakala C2 ialah S5 sebagai penyumbang utama. Analisis G-statistik umum dari tinggike-rendah adalah IA (1.099 x 10⁻³) > NIA (5.39 x 10⁻⁴) > VA (6.9 x 10⁻⁵). Pada tahun 2022, analisis hotspot di C1 menunjukkan S7 sebagai penyumbang utama; manakala C2 tetap sama dengan S5 sebagai penyumbang utama untuk mempunyai sumber pencemar dari tinggi-ke-rendah ialah IA (3.012 x 10⁻³) > NIA (5.95 x 10⁻⁴) > VA (7.6 x 10⁻⁵). Kesimpulannya, Moran I dan G-statistik banyak menyokong terutama dalam memberi manfaat kepada PCA dalam mengiktiraf corak sumber pencemar (contohnya kelas luas guna tanah) dengan tepat dan terperinci. Pada masa yang sama, analisis LISA dan hotspot mempunyai kelebihan terhadap HCA dengan mengenali kawasan kluster dengan lebih spesifik dan khusus, yang berdasarkan corak sumber pencemar.

ACKNOWLEDGEMENTS

I am very grateful to Buddha for His grace and mercy and for bestowing unto me the strength and blessings to endure until the completion of writing this thesis. Also, I am indebted to my supervisor, Dr. Faradiella Mohd Kusin, for her unfailing supervision and constant support. Her invaluable help in the form of constructive comments and suggestions throughout this thesis work have contributed to the success of this research. I would also like to convey my heartfelt thanks to my co-supervisors, Dr. Sarva Mangala Praveena, Dr. Zulfa Hanan Ashaari, and Dr. Nisfariza Mohd Noor, for their knowledgable advice, guidance, support and endless encouragement by instilling enthusiasm in me to persevere throughout the entire duration of my PhD studies.

At the same time, I would like to extend my grateful thanks to the staff from the Department of Environmental Science, Faculty of Environmental Studies, as well as the School of Graduate Studies (SGS) UPM for their exemplary performance by providing us (graduates students) with important information and reminders regarding our thesis. Needless to say, I would also like to extend my cordial thanks to the Department of Environment (DOE) Malaysia, Department of Irrigation and Drainage (JPS) Malaysia, Department of Town and Country Planning (JPBD) Malaysia, Malaysian Remote Sensing Agency (ARSM), and the United States Geological Survey (USGS) for providing insightful information, cooperation, support and discussions regarding the water quality data and land use data in Melaka River basin.

Above all, a word of thanks to my family who have always stood by me and giving me their wholehearted support and encouragement. Their unwavering support, have given me the strength and confidence to complete this PhD study.

Finally, I hereby acknowledge with full appreciation that this study would not have been realised without the contribution of my field assistants and friends who have given immense contribution, motivation, kindness, and moral support all the way until the completion of this study. Many thanks for their laughter, friendship and great memories. Last but not least, a thousand thanks to the residents and villagers who have given additional information and co-operation when the collection data was carried out within the Melaka River. Hopefully, their collaborative contributions and valuable information will bring to fruition in the near future the aim of this research study which will help improve the future of the water quality that flows along the Melaka River. I certify that a Thesis Examination Committee has met on 4 May 2018 to conduct the final examination of Ang Kean Hua on his thesis entitled "Evaluation of Water Quality Using Pattern Recognition Technique in Melaka River Basin, Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Normala Halimoon, PhD

Senior Lecturer Faculty of Environmental Studies Universiti Putra Malaysia (Chairman)

Ahmad Zaharin Aris, PhD

Professor Faculty of Environmental Studies Universiti Putra Malaysia (Internal Examiner)

Mohd Hasmadi Ismail, PhD

Associate Professor Faculty of Foresty Universiti Putra Malaysia (Internal Examiner)

Helmut Kroiss, PhD

Professor Emeritus Institute for Water Quality, Resources and Waste Management Vienna University of Technology Austria (External Examiner)

RUSLI HAJI ABDULLAH, PhD

Professor and Deputy Dean School of Graduate Studies Universiti Putra Malaysia

Date: 27 September 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Faradiella Mohd Kusin, PhD

Senior Lecturer Faculty of Environmental Studies Universiti Putra Malaysia (Chairman)

Zulfa Hanan Asha'ari, PhD

Senior Lecturer Faculty of Environmental Studies Universiti Putra Malaysia (Member)

Sarva Mangala Praveena, PhD

Associate Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Member)

Nisfariza Mohd Noor, PhD

Senior Lecturer Faculty of Arts and Social Sciences University of Malaya (Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date:

TABLE OF CONTENTS

ABSTRACT

Page

i

ABSTRAK ACKNOWLE APPROVAL DECLARAT LIST OF TA LIST OF FIG LIST OF AB	EDGEI ION BLES BURES BREV	MENTS S IATIONS	iii v vi viii xiii xviii xv xvii
CHAPTER			
1	INTR 1.1 1.2 1.3 1.4 1.5 1.6 1.7 1.8 1.9	DUCTION Introduction Overview of River Water Pollution Overview of the Effect of Land Use Change on Water Quality Problem Statement Objective of Study Research Questions Scope of Work Significance of Study Structure of Thesis	1 1 3 6 7 7 8 9 10
2	LITE	RATURE REVIEW	12
	2.1 2.2 2.3 2.4	Introduction2.1.1Melaka River DevelopmentThe Water Resources2.2.1Physical Parameter2.2.2Chemical Parameter2.2.3Biological Parameter2.2.4The Impact Pollutants2.2.5Environmetric TechniquesThe Land Use and Land ResourcesFundamentals and History of RemoteSensing2.4.1Process and Concept of Remote Sensing2.4.2Land Use Techniques in Water	12 12 13 15 16 18 19 22 41 44 45
	2.5	Quality StudyFundamentals of GeographicalInformation System2.5.1Components of GIS2.5.2The Concept of GIS2.5.3Application of GIS2.5.4Pattern Analysis in Water	47 68 69 71 73 74

	2.6	Quality Study Summary of Literature Review	86
3	MET 3.1 3.2 3.3 3.4 3.5 3.6 3.7	HODOLOGY Introduction Research Material Research Framework 3.3.1 Study Area Water Quality Monitoring Field Sampling and Analysis Data and Data Pre-treatment Interpretation of Water Quality Data 3.7.1 Hierarchical Cluster Analysis	88 88 89 92 94 95 98 99
		 (HCA) 3.7.2 Principal Components Analysis (PCA) 3.7.3 Discriminant Analysis (DA) 3.7.4 Pearson Correlation Coefficient 3.7.5 Prediction of Relation Between Land Use and Water Quality Using Neural Network 	100 100 101 101 102
	3.8	 3.7.6 Water Quality Index (WQI) Application of Remote Sensing 3.8.1 Image Pre-Processing and Land Use Land Cover Classification 	103 104 105
		3.8.2 Accuracy Assessment 3.8.3 Map Observation and Evaluation Process 3.8.4 LULC Change Detection	107 107
		Analysis 3.8.5 Markov Chain Model Analysis 3.8.6 Cellular Automata (CA) 3.8.7 CA-Markov Chain Model 3.8.8 Validating LUL C Prediction	107 108 109 109
		Model 3.8.9 Quality Assurance and Quality Control of Remote Sensing	110
	3.9	Application of Geographical Information System (GIS)	111
		 3.9.1 Morally Analysis and LISA Analysis 3.9.2 Hotspot Analysis 3.9.3 Quality Assurance and Quality 	111 112
4	RES	Control of GIS Analysis	113 115
	4.1 4.2	Introduction Water Quality Analysis 4.2.1.1 River Water Quality	115 115

		4.2.1.2	Assessment Correlation between Physical, Chemical and Biological	115
		4.2.2 4.2.3 4.2.4 4.2.5	Parameters in Melaka River basin Hierarchical Cluster Analysis Discriminant Analysis Principal Component Analysis Correlation between Land Use and Water Quality in Melaka	121 125 126 127
		4.2.6	River Basin Prediction of Land Use Variable Using Multilayered	132
			Neural Network	126
	13	Pomoto	Sensing Analysis	1/1
	4.5	431	Description of Land Lise	141
		4.0.1	Categories	141
		4.3.2	LULC Maps Assessment	142
		4.3.3	Validating LULC Prediction	
			Model	143
		4.3.4	LULC Change Detection	144
	4.4	Geograp	hical Information System	
		Analysis		157
		4.4.1	Moran I Analysis	157
		4.4.2	General G-statistic Analysis	165
		4.4.3	LISA Analysis	174
		4.4.4	Hotspot Analysis	189
	4.5	Summar	y of Major Findings	203
5	CONC			205
0	5 1	Introduct	ion	205
	5.2	Summar	v of the Major Findings	205
	0.2	5.2.1	Identification of the Pollution	200
			Sources in Melaka River	205
		5.2.2.	Land Use Land Cover Change	
			and Future Prediction of LULC	206
		5.2.3	Pattern Recognition Analysis of	
			Land Use Classification	206
	5.3	Recomm	nendations for Future Work	207
REFERENCES/BIBLIOGRAPHY				208
APPENDICE	S			222
BIODATA O	F STU	DENT		228
LIST OF PU	BLICA	TIONS		229

C

LIST OF TABLES

Table		Page
1.1	Water Pollution Source by Sector in 2012	
	(Malaysia)	5
2.1	Category of Water Resources	13
2.2	The Major Categories and Impact of Pollutants	19
2.3	Environmetric Techniques in Water Quality Studies	25
2.4 2.5	The Main Part of Spectrum	44
2.5	LULC Techniques in Water Quality Study	40
2.0	Hardware Software and Users	69
2.8	GIS Operation	71
2.9	Pattern Analysis in Water Quality Studies	78
3.1	River Physical and Functional Changes	93
3.2 (i)	Coordinate of Sampling Station	95
3.2 (ii)	Recommended Storage Condition for Certain	
	Analysis of Water Sample	96
3.3	Classes Delineated on the Basis of Supervised	
	Classification	106
4.1	Mean (and Standard Deviation) Values of Water	
	Quality Data Along Melaka River from Years 2001	110
1 2 (i)	National Water Quality Standard for Malaysia	119
4.2 (I)	NWOS	120
4.2 (ii)	Water Quality Classification According to National	120
()	Water Quality Standard (NQWS) for Malaysia	121
4.2 (iii)	PCC between Physical, Chemical and Biological	
	Parameter in Melaka River Basin	124
4.3	Classification Matrix for DA of Spatial Variation in	
	Melaka River Basin	126
4.4	Varimax Rotation PCs for Water Quality Data	
	Based on Two Clusters within Melaka River Basin	130
4.5 (1)	Pearson Correlation Coefficient and Coefficient of	
	Quality Variables in 2001 to 2015	126
4.5 (ii)	The Prediction Performance of R2 and RMSE for	130
4.5 (II)	Multi-Lavered Perceptron of Feed-Forward Neural	
	Network	137
4.5 (iii)	The Land Use Categories in 2015	142
4.5 (iv)	Comparison of Actual and Projected LULC types in	
	2015	143
4.6	Area (ha) of LULC type in Melaka River Basin for	
. –	2001, 2008, 2015 and 2022	146
4./	I ransition Probability of Area and Matrix Calculated	140
4.0	using Land-use Maps of 2001-2008	146
4.0	using Land use Maps of 2008 2015	146
49	Transition Probability of Area and Matrix Calculated	140
1.0	ranomore robubling of Area and Mathy Odloulated	

	using Land-use Maps of 2015-2022	147
4.10	Rate of Losses, Gains, and Net Changes of LULC Areas (ha)	148
4.11	Moran Index Analysis for Vegetation Area, Non- Industrial Area, Industrial Area, Open Space Area and Farming Area between the year of 2015 to	
	2022	159
4.12	General G-Statistic Analysis for Vegetation Area, Non-Industrial Area, Industrial Area, Open Space Area and Farming Area between the year of 2015	
	to 2022	168
4.13	LISA Analysis for Vegetation Area, Non-Industrial Area, Industrial Area, Open Space Area and	
	Farming Area between the year of 2015 to 2022	178
4.14	Hotspot Analysis for Vegetation Area, Non- Industrial Area, Industrial Area, Open Space Area and Farming Area, between the year of 2015 to	
	2022	192

C

LIST OF FIGURES

	LIST OF FIGURES	
Figure		Page
1.0	Tree Analysis of Land Use Change in Melaka River	
	Basin	4
1.1	River Water Quality Status in Malaysia (2005-2012)	5
2.1	Independent and Dependent Variable	24
2.2	Symptoms of Problems of Pressure on Land and	
2.2	Resources	43
2.3	Electromagnetic Spectrum	45
2.5	GIS Components	69
2.6	The Concept of GIS	71
2.7	GIS Applied in Various Activities	74
3.1	Flow Chart of Research Framework	91
3.2	Sampling Stations Along Melaka River Basin	94
3.3	A Schematic Diagrams of The Final Neural Network	102
4.1 (a)	Solide	117
41(b)	Mean Data for pH Temperature Salinity Total	117
(0)	Suspended Solid, and Turbidity	118
4.1 (c)	Mean Data for BOD, COD, DO, and NH3N	118
4.1 (d)	Mean Data for Trace Metals of As, Hg, Cd, Cr, Pb,	
	and Zn	118
4.1 (e)	Mean Data for Total Coliform and Escherichia	110
12	Colliorm	119
4.2	Clusters using Ward Linkage Method in HCA	126
4.3	Box and Whisker Plots of Some Parameters	
	Separated from DA Associated with Water Quality	
	Data of Melaka River	127
4.4 (a)	Cluster 1 with 6 Principal Components	131
4.4 (b)	Cluster 2 with 8 Principal Components	131
4.4 (CT)	Percentron of Feed Forward Neural Network	
	between Land-use and Water Quality	138
4.4 (c2)	The Performance of RMSE for Multi-Lavered	100
	Perceptron of Feed-Forward Neural Network	
	between Land Used and Water Quality	138
4.4 (d1)	Prediction for Vegetation Area with Water Quality	
4.4.(-10)	Variables	139
4.4 (d2)	Prediction for Non-Industrial Area with Water	120
4 4 (d3)	Prediction for Industrial Area with Water Quality	155
(00)	Variables	140
4.4 (d4)	Prediction for Open Space Area with Water Quality	
	Variables	140
4.4 (d5)	Prediction for Farming Area with Water Quality	
	Variables	141

G

4.5 (a)	LULC Actual Maps of 2001	149
4.5 (b)	LULC Actual Maps of 2008	150
4.5 (c)	LULC Actual Maps of 2015	151
4.5 (d)	LULC Simulated Map of 2015	152
4.5 (e)	LULC Simulated Map of 2022	153
4.6	Moran's Index Analysis of VA2015, VA2022,	
	NIA2015, NIA2022, IA2015, IA2022, OSA2015,	
	OSA2022, FA2015, FA2022	160
4.7	General G Analysis of VA2015, VA2022, NIA2015,	
	NIA2022, IA2015, IA2022, OSA2015, OSA2022,	
	FA2015, FA2022	169
4.8	LISA Index Analysis of VA2015, VA2022, NIA2015,	
	NIA2022, IA2015, IA2022, OSA2015, OSA2022,	
	FA2015, FA2022	179
4.9	Hotspot Analysis of VA2015, VA2022, NIA2015,	
	NIA2022, IA2015, IA2022, OSA2015, OSA2022,	
	FA2015, FA2022	193
	Reading produced produced in the second seco	

C

LIST OF ABBREVATIONS

Area of Interest American Public Health Association Cellular Automata-Markov Discriminant Analysis Discriminant Function Department of Environment Geographical Information System
Hectare
Hierarchical Cluster Analysis
High-Pollution Sources
Inductive Coupled Plasma-Mass Spectrometry
Local Indicators of Spatial Association
Land Use Land Cover
Millions of Litres per Day
Moderate-Pollution Sources
Malaysian Remote Sensing Agency
National Water Quality Standard
Principal Component Analysis
Quality Control and Quality Assurance
Region of Interest
Remote Sensing
Standard Deviation
Sub-Indexes
Sampling Station
United Nationals Educational, Scientific and Cultural Organization
United States Geological Survey
Varimax Factor
Water Quality Index

CHAPTER 1

INTRODUCTION

1.1 Introduction

Malaysia achieved the best performance in the Environmental Performance Index (EPI) which was recognised as the best in the Asia-Pacific region and ranked 25th out of a total of 132 countries worldwide (MRPE Official Portal, 2012). New Zealand took the top position followed by Japan and Malaysia in the Asia-Pacific region. The EPI for Malaysia is equivalent to the EPI for the best-performing group of countries such as Germany, Iceland, Finland, Denmark, Belgium, and Japan. Among the various aspects that were evaluated ranged from climate change, agriculture, fisheries, forestry, air and water pollution (MRPE Official Portal, 2012). Inevitably, water pollution is considered an important aspect that requires attention which would affect the EPI performance.

Generally, Malaysia is made up of thirteen states, in which Melaka is the smallest state after Perlis and Penang. The Melaka state is located in the southern region of the Peninsular Malaysia and is bordered by Negeri Sembilan to the north and Johor to the south. The Melaka state also known as a state of traditions and was listed as a UNESCO World Heritage Site since 7th July, 2008 (UNESCO Official Portal, 2016). This has provided immense economic development and job opportunities through various sectors such as tourism, commercial and industrial activities. However, the rapid development and urbanisation have indirectly affected the political, cultural, and environmental aspects in the Melaka state.

Inadvertently, unorganised expansion and uncontrolled urbanisation development have led to several water pollution issues within the Melaka River (Rosli et al., 2015). As highlighted earlier, the rapid development and urbanisation that took place in the Melaka state have indirectly resulted in drastic and extreme land use changes within a few years back. In other words, forest land are being victimised for deforestation to transform into agriculture, urban, industrial, as well as farmland, which are important to enhance the quality of human life. This circumstance is taking place especially in Kampung Batu Berendam sub-basin in the urban area after it was recognised as the world tourism centre, where majority of the land resources was in the process of conversion into other classes of land use for human activity purposes (Figure 1.0). Due to the increasing demand for land resources within a period of 5 years after 2007 (Daneshmand et al., 2011), the development for urbanisation has also increased and is suspected to extend northwards into Kampung Cheng, Kampung Tualang and Kampung Harmoni Belimbing Dalam sub-basin area. Since various activities are carried out along the Melaka River and within the Melaka River basin, this has indirectly caused river water pollution (Hamid *et al.*, 2016; Rosli *et al.*, 2015; Baharuddin *et al.*, 2014; Mustpha *et al.*, 2013; Lim *et al.*, 2012; Juahir *et al.*, 2011) and has impacted the riverine ecosystem (Govorov *et al.*, 2016; Sun, 2015; Zhai *et al.*, 2014). This was supported by previous studies of Bu *et al.* (2014), Fucik *et al.* (2014) and Kibena *et al.* (2014), who emphasised that any changes in land use will affect the environmental ecosystem, especially the quality of river water.

Until today, the development of Melaka State is still in line with its mission to be a developed country by 2020. In line with the vision, the state government had planned to reconstruct the Melaka State into three phases, namely (i) the urban areas with residential and commercial activities, (ii) sub-urban areas with industrial activities, as well as (iii) rural areas with agriculture and farming activities. Continuous development could cause pollution in the Melaka River to increase, as the circumstances are connected with the illegal vegetation and farming activities, rapid and uncontrolled development of residential and commercial activities, failure to maintain and manage sewage treatment plant, as well as focusing less on industrial activities. This would probably lead to the inability to control the unceasing pollution due to the ignorance of the danger of pollution on waste dumping, legal law and regulation. These are unavoidable in order to prevent direct discharge of wastewater, as 40% of household, 30% of industrial, and 30% of agriculture are connected with sewage network in the Melaka River basin (Figure 1.0).

In this case, the river water will experience high level of organic and inorganic pollution. This condition would increase when wastewater from factories and household are discharged directly into the river without undergoing any treatment. Similarly, there are also issues where wastewater treated in plants does not meet the environmental standards. In other words, the river water quality can be deteriorated due to majorly by anthropogenic activities and a minor contribution from natural activities (Al-badaii *et al.*, 2016; Hamid *et al.*, 2016; Herojeet *et al.*, 2016). This means that any changes of land resources from the natural into constructed environment (El-Zeiny and El-Kafrawy, 2016; Mishra and Rai, 2016; Yildiz and Doker, 2016) will directly or/and indirectly affect the good and healthy quality of water condition.

Apart from reducing the water quality level, the riverine ecosystem will be harmful as some organic pollution will lead to the spread of disease resulting to the death of fishes. For instance, several issues on the pollution of the Melaka River had been reported for the death of aquatic fisheries due to lack of oxygen and disease spreading within the river water (PPPP Official Portal, 2015; Sinar Official Portal, 2012; Nasbah, 2010; Jabar, 2010), where such circumstances would cause a decline in the fish species and families resulting to extinction. Simultaneously, the water diseases in the river would be harmful to the aquatic species, which are considered as food supply in the food chain. This will directly or indirectly transfer the disease to the human community upon consumption of the food. Therefore, drastic land use changes could lead to the

destruction and impairment of the river water quality as well as to animal and human which are fully dependent on the environment for survival.

In view of this, the present study intends to investigate the extent to which the Melaka River had been contaminated. By taking into account the potential sources of water pollution in relation to the changes in previous and current land uses, future prediction of the affected areas would be possible. This is especially important for future development planning and decision making in order to avoid any overexploitation of the existing land use.

1.2 Overview of River Water Pollution

Every day, 2 million tons of industrial and agricultural discharges were released into the world water, of which the estimated amount of wastewater produced annually is about 1,500 millions of litres per day (MLD) (UNWWAP, 2009). The National Geographic Portal (2016) reported that developing countries produced 70% of industrial wastes that were dumped untreated into waters. The deterioration in the quality of the river water was due to growing population, rapid urban development, anthropogenic inputs (e.g. municipal and industrial wastewater discharges, agricultural runoff) and natural processes (e.g. chemical weathering and soil erosion) (Figure 1.0) (Holloway *et al.*, 1998; Singh *et al.*, 2011; Shin *et al.*, 2013) that have threatened human and ecological health, availability of drinking water, and further economic development (Houser and Richardson, 2010; Li and Zhang, 2010; Morse and Wolheim, 2014).



The contamination of river water is no exception for a developing country like Malaysia. According to the Department of Environment (2012), among the 473 rivers monitored, only 278 (59%) were found to be clean, 161 (34%) were slightly polluted and 34 (7%) were already polluted (Figure 1.1). Apparently, within 8 years from 2005 to 2012, the total number of rivers that have a clean condition had decreased by about 60 rivers. The trend indicates that the quality of clean river water is decreasing, whereas a higher percentage of the rivers had changed into slightly polluted or polluted river. Notwithstanding, generally the status of the river water pollution can be associated with the level of water quality indicators such as biochemical oxygen demand (BOD), ammoniacal nitrogen (NH₃N), and suspended solids (SS). For instance, a high BOD can be attributed to inadequate treatment of sewage or effluent from agro-based and manufacturing industries. NH₃N mainly comes from livestock farming and domestic sewage; while SS is mainly due to improper earthworks and land clearing activities. Apparently, most of the factors that have led to river water pollution are caused by human activities as shown in Table 1.1.



Figure 1.1: River water quality status in Malaysia (2005-2012) (Source: DOE Malaysia Report, 2012)

Ible 1.1: Water Pollution Source by Sector in 2012 (Malaysia)					
Type of So	ources	No. of Sources	Percentage		
Manufacturing	Industries	4,595	0.276		
Agro-based Ir	ndustries				
(a) Rub	ber Mill	72	0.004		
(b) Palm	n Oil Mill	436	0.026		
Animal Farm (Pi Sewage Treatn	ig Farming) nent Plant	754	0.045		
(a) P	ublic	5,800	0.349		
(b) Pr	rivate	4,083	0.246		

Ta

(c) Individual Septic Tank	1,449,383	87.190		
(d) Communal Septic Lank	3,631	0.218		
Food Services Establishment	192,710	11.593		
Wet Market	865	0.053		
Source: DOE, 2012				

1.3 Overview of the Effect of Land Use Change on Water Quality

Melaka state is experiencing rapid economic growth with various kinds of transformation through industrialisation and urbanisation. This has somehow resulted to an overexploitation of natural resources and massive increase in the discharge of pollutants into the waterways. This is especially pronounced in areas of dense population with rapid population growth and rapid changes in land use land cover (LULC). Among the most affected areas in the state of Melaka are within the Melaka River basin; potentially due to increased industrial and residential expansion. Many researchers agreed that uncontrolled and unmanageable land use development would seriously pollute the river water quality (Bu et al., 2014; Fucik et al., 2014; Gyawali et al., 2013). Generally, both industrial and domestic wastewater discharged by human activities may cause serious water quality degradation such as water eutrophication, algal blooms, loss of ecological function and other problems (Zhao et al., 2012). Since industrial wastewater has complex chemical components, and is difficult to treat, degrade and purify, it could be more harmful than the domestic wastewater. Apparently, the impact of industrialisation and urbanisation on water quality is inevitable. However, careful management of current and future land use may negate the impact and lessen the extent of contamination on our waterways.

1.4 Problem Statement

Rapid population growth, unorganised expansion, and uncontrolled urbanisation development has led to various environmental issues in the Melaka state (Rosli et al., 2015). Among the apparent environmental issues which was reported to the public and the state authorities are cases of water pollution in the river. In recent years, several cases involving river water pollution have been reported on Melaka River (Nasbah, 2010; Jabar, 2010). Various efforts have been carried out to reduce and prevent continuous pollution in the river, e.g. through the beautification of the Melaka River which is a synchronised effort with the state tourism sector (Sinar Official Portal, 2012). Even with multiple beautification and rehabilitation programmes in place, the Melaka River continues to be polluted. For instance, numerous cases have been reported on aquatics death and the situation is still worrying (PPPP Official Portal, 2015; Zakinan, 2015). Therefore, it is important that extensive monitoring and assessment of the river water quality be carried out to identify potential sources of pollution across the river basin (Hamid et al., 2016; Albadaii et al., 2016; Rosli et al., 2015; Baharuddin et al., 2014).

In most previous studies, the water quality was analysed using statistical data analysis. The analysis confirmed that the water quality was affected due to the temporal and spatial variations that have possibly caused the contamination (Hamid et al., 2016; Al-badaii et al., 2016; Aris et al., 2013; Juahir et al., 2011). According to a report by DOE in 2012, the river water pollution as indicated by the values of Biochemical Oxygen demand (BOD), Suspended Solid (SS), and Ammoniacial-NitrogenBOD, (NH₃N), was suspected to have come from industrial, residential, agriculture, as well as farming activities (Mishra and Rai, 2016; Parsa et al., 2016; Yulianto et al., 2016; Wilson, 2015; Fucik et al., 2014; Gyawali et al., 2013). These are being continuously investigated to find an appropriate solution towards water quality destruction of the Melaka River basin (Rosli et al., 2015; Daneshmend et al., 2011). In other words, land use classes are anticipated to have a relationship with the water quality in order to sustain the level of ecological health (Govorov et al., 2016; Shi et al., 2016; Fucik et al., 2014; Kibena et al., 2014). Therefore, if the land use changes in the future, the level of contamination will also be subjected to changes. However, there are uncertainties on what the impact would be. Hence, several studies have suggested that land uses are assumed to influence the river water quality and are appropriate to be included as part of the analysis (El-Zeiny and El-Kafrawy, 2016; Kibena et al., 2014), rather than be dependent on the water quality data when determining the source of the pollutants (Hamid et al., 2016; Al-badaii et al., 2016; Rosli et al., 2015; Baharuddin et al., 2014). Therefore, by suggesting land use as a variable, this study is conducted to determine with the relation between land use and water quality in the Melaka River basin using Geographical Information System (GIS) and remote sensing techniques.

1.5 Objectives of Study

This study aims to identify the pollutant sources in the Melaka River using the LULC change model and pattern analysis. The specific objectives can be described as follows;

- (a) To identify the potential sources of Melaka River pollution in relation with temporal land use classification;
- (b) To determine the land use land cover changes in 2015 and predict the land use land cover for 2022;
- (c) To verify the pattern recognition based on land use classification between 2015 and 2022 in the determination of the pollutant source within the Melaka River.

1.6 Research Questions

This study will attempt to identify the sources of pollution within a catchment scale based on the changes in river water quality and corresponding land uses. The underlying research questions are:

 What is the extent of river pollution occurrence in the Melaka River? Is there any relation between land use classification and water quality variables?

- What are the changes from previous, current and future land use in relation to water quality in a particular area?
- What is the pattern of pollutant source in conjunction with the land use that may impact the river water quality in the Melaka River?

1.7 Scope of Work

River pollution is reported yearly worldwide (Al-badaii *et al.*, 2016; Baharuddin *et al.*, 2014; Mustapha *et al.*, 2013; Gazza *et al.*, 2012; Juahir *et al.*, 2011). According to DOE (2012), Melaka River was listed in the report as having pollution in the country. This condition was supported by Daneshmand *et al.* (2011) and the river water quality is expected to continue decreasing (Rosli *et al.*, 2015). As has been proven, daily news had reported that some aquatic species were found dead (PPPP Official Portal, 2015; Zakinan, 2015) in the Melaka River. Apparently, the quality of the riverine ecosystem has been greatly affected. This condition reflects that contamination might have occurred in the Melaka River basin.

Applied environmetric techniques using HCA, DA and PCA were popularly used by researchers to investigate and identify the source of pollution (Hamid et al., 2016; Al-badaii et al., 2016; Aris et al., 2013; Juahir et al., 2011) in the Melaka River basin (Rosli et al., 2015; Daneshmand et al., 2011). In general, river water was investigated of its pollution level based on the the physicochemical parameters (i.e. pH, temperature, electrical conductivity (EC), salinity, turbidity, total suspended solid (TSS), dissolved solids (DS), dissolved oxygen (DO), biochemical oxygen demand (BOD), chemical oxygen demand (COD) and ammoniacal-nitrogen (NH₃N), biological indicators (i.e. total coliform and Escherichia coliform) as well as trace elements (i.e. mercury, cadmium, chromium, arsenic, zinc, lead, and iron). Pollution can be anticipated from anthropogenic activities (such as industrial, municipal and domestic waste, agricultural runoff, animal farm, land clearing) and/or natural processes (e.g. erosion and runoff) (Al-badaii et al., 2016; Hamid et al., 2016; Mustapha et al., 2013; Juahir et al., 2011). Therefore, environmetric techniques are suitable for application in the Melaka River basin as these techniques are important for collating information regarding the pollutant sources.

In general, the water pollution in the Melaka River basin continues to occur until today. Therefore, suggestion to incorporate land use and water quality data has become the main subject in this study. In particular, the pollutant sources are suspected to be linked with land use classes. The land use classes are able to change temporarily over a period of time (Mishra and Rai, 2016; Parsa *et al.*, 2016; Yulianto *et al.*, 2016; Wilson, 2015). In other words, land use classes possess connectivity with human activities, where this condition requires particular focus on the land use change (El-Zeiny and El-Kafrawy, 2016; Kibena *et al.*, 2014) especially when involving extensive development. For example, the activities that is associated with industrial, residential, commercial, transportation, and sewage treatment plant (Wilson, 2015; Fucik *et al.*, 2014; Gyawali *et al.*, 2013). Therefore, the techniques in GIS (e.g. Moran I

and LISA, as well as general G-statistic and Hotspot analysis) have become the major tools to pattern the pollutant source in a river basin. Accordingly, this study uses the land use classes and water quality data to determine the pollutant source based on the monitored station along the Melaka River. Apparently, the application of GIS in the river water quality analysis had been widely used in the discovery of the contamination (Sun, 2015; Alqadi *et al.*, 2014; Lezzaik and Milewski, 2014), and additional input from the land use data (El-Zeiny and El-Kafrawy, 2016; Kibena *et al.*, 2014; Chang, 2008) will extensively support the findings that is produced from the environmetric technique approach.

1.8 Significance of Study

Environmetric techniques analysing the spatial variability of large and complex river water quality data provide meaningful information by identifying, delineating, and differentiating contaminants resulting from various sources (Hamid *et al.*, 2016; Al-badaii *et al.*, 2016; Aris *et al.*, 2013; Juahir *et al.*, 2011). Considering the methods applied such as HCA, DA, and PCA in the techniques, the findings offer reliable classification and serve as an excellent tool for water resources control and management (Hamid *et al.*, 2016; Al-badaii *et al.*, 2015; Baharuddin *et al.*, 2014). These techniques allow the optimisation of a monitoring programme by decreasing the number of sampling stations, number of parameters monitored, as well as serving as a cost-effective method in the water quality assessment (Papaioannou *et al.*, 2010a). Therefore, this study applies environmetric techniques into water quality data analysis to visualise the condition of the pollutant source in the Melaka River basin.

Further discussion on the river pollution has become the main point of study for majority of the researchers. The pollutant source is typically related with river water quality and the land use change (Mishra and Rai, 2016; Parsa et al., 2016; Yulianto et al., 2016; Wilson, 2015; Fucik et al., 2014; Gyawali et al., 2013). The land use in LULC analysis is anticipated to be linked with water quality in the river (Wilson, 2015; Fucik et al., 2014; Gyawali et al., 2013). This condition may continue to affect the river in the future, -if it has connection with the land use (Mishra and Rai, 2016; Parsa et al., 2016; Yulianto et al., 2016). Additional evidence from El-Zeiny and El-Kafrawy (2016), as well as Kibena et al. (2014) indicated that the land use may be considered as a variable (e.g. independent) in the analysis, whenever land use has relation with the water quality data. Hence, this study is carried out to evaluate the land use classes that potentially impact the water quality in the river. Therefore, this study is focused on the previous and current land use changes while predicting the future trend in relation to the water quality in the Melaka River basin. On the other hand, GIS was typically included for assessment of the water quality data (Sun, 2015; Alqadi et al., 2014; Lezzaik and Milewski, 2014). Previous studies suggested that pattern analysis of GIS (Moran I and LISA, general G-statistic and Hotspot) are favourable for inclusion into the river water quality study to interpret the pollutant source in a particular area (Shi et al., 2016; Govorov et al., 2016; Zhai et al., 2014; Zhao et al., 2012). Therefore, this study is carried

out to incorporate the land use classes and water quality analysis to recognise the pattern of pollutant source in the Melaka River basin. This study applies land use classes as independent variable and water quality data as dependent variable.

1.9 Structure of Thesis

The content of this thesis is presented in five chapters as follows;

(i) Chapter One: Introduction

In this chapter, an overview of the river water pollution in Malaysia is presented. This chapter also highlights the problem statement of the contamination that has occurred in the Melaka River. This will be further explained through the objectives of the study, thus strengthening the significance of this study. Literature reviews and research findings are presented in the following chapters.

(ii) Chapter Two: Literature Review

The literature review explains the previous studies on river water quality including the fundamentals of water quality characteristics (i.e. physicochemical and biological parameters), as well as the impact of water pollution. The conceptual and fundamental methods and techniques applied in the methodologies will also be described, which include the environmetric technique, LULC change models, and pattern analysis models.

(iii) Chapter Three: Methodology

The methodology section describes the research methodology that includes primary and secondary data collection. The collection and analysis of data for river water quality is presented alongside the land use data acquisition. In addition, the concepts of environmetric models, LULC change models, and pattern analysis models are also explained in this chapter.

(iv) Chapter Four: Results and Discussion

This chapter reports on the major findings of the study. This begins with a discussion and interpretation of the river water quality assessment of the Melaka River. It is followed by the results of the land use and land cover changes across the study area. This includes the discussion on previous and current land uses and future prediction changes, as well as the pattern analysis based on the land use classification.



(v) Chapter Five: Conclusion and Recommendation

Chapter Five summarises the overall research findings and the conclusion drawn from the study. This section also provides some recommendations for further improvement for future research.



REFERENCES

- Abdi, H. & William, L.J. (2010). Principal component analysis. *Wiley Interdisciplinary Reviews: Computational Statistics*, 2(4), 433-459.
- Abdulkareem, J. H., Pradhan, B., Sulaiman, W. N. A., & Jamil, N. R. (2017). Prediction of spatial soil loss impacted by long-term landuse/land-cover change in a tropical watershed. *Geoscience Frontiers*.
- ABS [Australian Bureau of Statistics] (2003). A Guide to Interpreting Time Series Monitoring Trends.
- Al-Badaii, F., Halim, A. A., & Shuhaimi-Othman, M. (2016). Evaluation of Dissolved Heavy Metals in Water of the Sungai Semenyih (Peninsular Malaysia) using Environmetric Methods. Sains Malaysiana, 45(6), 841-852.
- Alberto, W. D., del Pilar, D. M., Valeria, A. M., Fabiana, P. S., Cecilia, H. A., & de los Ángeles, B. M. (2001). Pattern Recognition Techniques for the Evaluation of Spatial and Temporal Variations in Water Quality. A Case Study:: Suquía River Basin (Córdoba–Argentina). Water research, 35(12), 2881-2894.
- Alqadi, K. A., Kumar, L., & Khormi, H. M. (2014). Mapping hotspots of underground water quality based on the variation of chemical concentration in Amman, Zarqa and Balqa regions, Jordan. *Environmental earth sciences*,71(5), 2309-2317.
- American Public Health Association (APHA) (2005). Standard Methods for the Examination of Water and Wastewater. 21st ed. Washington: American Water Works Association, Water Environment Federation.
- Anselin, L. (1995). Local indicators of spatial association LISA. *Geographical analysis*, 27(2), 93-115.
- Aris, A. Z., Praveena, S. M., Isa, N. M., Lim, W. Y., Juahir, H., Yusoff, M. K., & Mustapha, A. (2013). Application of environmetric methods to surface water quality assessment of Langkawi Geopark (Malaysia). *Environmental Forensics*, 14(3), 230-239.
- Aris, A. Z., Praveena, S. M., Abdullah, M. H., & Radojevic, M. (2012). Statistical approaches and hydrochemical modelling of groundwater system in a small tropical island. *Journal of Hydroinformatics*, 14(1), 206-220.
- Aronoff, S. (1989) *Geographical information systems: A management perspective*. Ottawa, WDL Publications.
- Arsanjani, J. J., Helbich, M., Kainz, W., & Boloorani, A. D. (2013). Integration of logistic regression, Markov chain and cellular automata models to simulate urban expansion. *International Journal of Applied Earth Observation and Geoinformation*, 21, 265-275.

Avvannavar, S. M., & Shrihari, S. (2008). Evaluation of water quality

index for drinking purposes for river Netravathi, Mangalore, South India. *Environmental Monitoring and Assessment*, *143*(1-3), 279-290.

- Azid, A., Juahir, H., Toriman, M.E., Kamarudin, M.K.A., Saudi, A.S.M., Hasnam, C.N.C., Aziz, N.A.A., Azaman, F., Latif, M.T., Zainuddin, S.F.M. and Osman, M.R. (2014). Prediction of the level of air pollution using principal component analysis and artificial neural network techniques: A case study in Malaysia. *Water, Air, & Soil Pollution, 225*(8), 1-14.
- Azid, A., Juahir, H., Latif, M. T., Zain, S. M., & Osman, M. R. (2013). Feed-forward artificial neural network model for air pollutant index prediction in the southern region of Peninsular Malaysia. *Journal of Environmental Protection*, *4*(12), 1-10.
- Badreldin, N., & Goossens, R. (2014). Monitoring land use/land cover change using multi-temporal Landsat satellite images in an arid environment: a case study of El-Arish, Egypt. *Arabian Journal of Geosciences*, 7(5), 1671-1681.
- Baharuddin, N., NOR'ASHIKIN, S. A. I. M., & Zain, S. M. (2014). Characterization of Spatial Patterns in River Water Quality Using Chemometric Techniques. *Sains Malaysiana*, *43*(9), 1355-1362.
- Bai, Y., Wang, M., Peng, C., & Alatalo, J. M. (2016). Impacts of urbanization on the distribution of heavy metals in soils along the Huangpu River, the drinking water source for Shanghai. *Environmental Science and Pollution Research*, 23(6), 5222-5231.
- Balzter, H. (2000). Markov chain models for vegetation dynamics. *Ecological Modelling*, 126(2), 139-154.
- Barcelo, D. & Petrovic, M. (2011). Ebro River Basin: The Handbook of Environmental Chemistry. Heidelberg Germany: Springer, p. 431.
- Bateni, F., Fakheran, S., & Soffianian, A. (2013). Assessment of land cover changes & water quality changes in the Zayandehroud River Basin between 1997–2008. *Environmental monitoring and assessment*, *185*(12), 10511-10519.
- Behera, D. M., Borate, S. N., Panda, S. N., Behera, P. R., & Roy, P. S. (2012): Modelling and analyzing the watershed dynamics using Cellular Automata (CA)–Markov model–A geo-information based approach, *Journal of earth system science*, 121(4), 1011-1024.
- Bierman, P., Lewis, M., Ostendorf, B., & Tanner, J. (2011). A review of methods for analysing spatial and temporal patterns in coastal water quality. *Ecological Indicators*, *11*(1), 103-114.
- Bolstad, P. V., & Swank, W. T. (1997). Cumulative impacts of landuse on water quality in a southern Appalachian watershed. *JAWRA Journal of the American Water Resources Association*, *33*(3), 519-533.
- Boyacioglu, H. (2012). Utilization of environmetric & index methods as

water quality comparative assessment tools focusing on heavy metal content. *Archives of Environmental Protection*, *38*(3), 17-28.

- Boyacioglu, H., & Boyacioglu, H. (2008). Water pollution sources assessment by multivariate statistical methods in the Tahtali Basin, Turkey. *Environmental Geology*, *54*(2), 275-282.
- Boyacioglu, H., & Boyacioglu, H. (2007). Surface water quality assessment by environmetric methods. *Environmental monitoring and assessment*, *131*(1-3), 371-376.
- Brown, S. D., Sum, S. T., & Despagne, F. (1996). Chemometrics. *Analytical Chemistry*, 68, 21R–61R.
- Brown, S. D., Blank, T. B., Sum, S. T., & Weyer, L. G. (1994). Chemometrics. *Analytical Chemistry*, *66*, 315R–359R
- Bu, H., Meng, W., Zhang, Y., & Wan, J. (2014). Relationships between land use patterns and water quality in the Taizi River basin, China. *Ecological Indicators*, *41*, 187-197.
- Campbell, J.B. & Wynne, R.H. (Jun 21, 2011) *Introduction to Remote Sensing*. 5th Edition. The Guilford Press, New York, London. 667p.
- Chang, H. (2008). Spatial analysis of water quality trends in the Han River basin, South Korea. *Water research*, *42*(13), 3285-3304.
- Chang, K. S. (2008) Introduction to Geographic Information Systems. Forth edition. McGraw-Hill International Edition.
- Chatfield, C. (2004). The Analysis of Time Series: An Introduction. 6th edition. Chapman & Hall/CRC.
- Chen, J., & Lu, J. (2014). Effects of land use, topography and socio economic factors on river water quality in a mountainous watershed with intensive agricultural production in East China. *PloS one*, 9(8), e102714.
- Chua, Y. P. (2011). Kaedah dan statistik penyelidikan: kaedah penyelidikan. Mcgraw Hill Education.
- Cigizoglu, H. K., & Kisi, Ö. (2006). Methods to improve the neural network performance in suspended sediment estimation. *Journal* of hydrology, 317(3-4), 221-238.
- Cliff, A. D. A. D. C., & Ord, J. K. (1973). *Spatial autocorrelation* (No. 04; QA278. 2, C5.). New York: Methuen.
- Colwell, R.N. (1966) Uses and limitation of multispectral Remote Sensing. In *Proceedings of the Fourth Symposium on Remote Sensing of Environment*, pp. 71-100. Ann Arbor: Institute of Science and Technology, University of Michigan.
- Daneshmand, S., Huat, B. B., Moayedi, H., & Ali, T. A. M. (2011). Study on water quality parameters of Linggi and Melaka rivers catchments in Malaysia. *Engineering Journal*, 15(4), 41-52.
- Davis, A. P., & McCuen, R. H. (2005). Stormwater management for smart growth. Springer Science & Business Media.
 Department of Environment Malaysia (DOE) (2012). Malaysia

Environmental Quality Report 2012. Department of Environment, Ministry of Natural Resources and Environment, Kuala Lumpur, Malaysia.

Department of the Environment (1987) *Handling Geographic Information.* Report of the Committee of Enquiry chaired by Lord Chorley. HMSO, London.

- Department of Irrigation and Drainage Malaysia (DIDM) (2009). Study of the river water quality trends and indexes in Peninsular Malaysia. *Water Resources Publication*, No.2
- Di Gregorio, A., & Jansen, L. J. (1998). Land Cover Classification System (LCCS): classification concepts and user manual. *FAO*, *Rome*.
- Dominick, D., Juahir, H., Latif, M. T., Zain, S. M., & Aris, A. Z. (2012). Spatial assessment of air quality patterns in Malaysia using multivariate analysis. *Atmospheric Environment*, 60, 172-181.
- Environmental Literacy Council (ELC) (2016). Land Use. Retrieved from https://enviroliteracy.org/land-use/
- El-Asmar, H. M., Hereher, M. E., & El Kafrawy, S. B. (2013). Surface area change detection of the Burullus Lagoon, North of the Nile Delta, Egypt, using water indices: A remote sensing approach. *The Egyptian Journal of Remote Sensing and Space Science*, *16*(1), 119-123.
- El-Kawy, O. A., Rød, J. K., Ismail, H. A., & Suliman, A. S. (2011). Land use and land cover change detection in the western Nile delta of Egypt using remote sensing data. *Applied Geography*, *31*(2), 483-494.
- El-Zeiny, A., & El-Kafrawy, S. (2016). Assessment of water pollution induced by human activities in Burullus Lake using Landsat 8 operational land imager and GIS. *The Egyptian Journal of Remote Sensing and Space Science*.
- Fan, F., Wang, Y., & Wang, Z. (2008). Temporal and spatial change detecting (1998-2003) and predicting of land use and land cover in Core corridor of Pearl River Delta (China) by using TM and ETM+ images. *Environmental Monitoring and Assessment*, 137(1-3), 127-147.
- FAO/UNEP (1997). Negotiating a Sustainable Future for Land. Structural and Institutional Guidelines for Land Resources Management in the 21st Century. FAO/UNEP, Rome.
- Fecht, D., Hansell, A. L., Morley, D., Dajnak, D., Vienneau, D., Beevers, S., Toledano, M.B., Kelly, F.J., Anderson, H.R. & Gulliver, J. (2016). Spatial and temporal associations of road traffic noise and air pollution in London: Implications for epidemiological studies. *Environment international*, 88, 235-242.
- Fučík, P., Novák, P., & Žížala, D. (2014). A combined statistical approach for evaluation of the effects of land use, agricultural and urban activities on stream water chemistry in small tile-

drained catchments of south Bohemia, Czech Republic. *Environmental Earth Sciences*, 72(6), 2195-2216.

- Gao, J. (2009). *Digital analysis of Remote Sensing imagery*. McGraw Hill, New York. 646p.
- Garnier, J., Brion, N., Callens, J., Passy, P., Deligne, C., Billen, G., Servais, P., & Billen, C. (2013). Modeling historical changes in nutrient delivery and water quality of the Zenne River (1790s– 2010): the role of land use, waterscape and urban wastewater management. *Journal of marine systems*, *128*, 62-76.
- Gazzaz, N. M., Yusoff, M. K., Ramli, M. F., Aris, A. Z., & Juahir, H. (2012). Characterization of spatial patterns in river water quality using chemometric pattern recognition techniques. *Marine Pollution Bulletin*, 64(4), 688-698.
- Getis, A., & Ord, J. K. (1992). The analysis of spatial association by use of distance statistics. *Geographical analysis*, 24(3), 189-206.
- Gleick, P. H. (1993). *Water in crisis: a guide to the world's fresh water resources*. Oxford University Press, Inc..
- Govorov, M., Putrenko, V., & Gienko, G. (2016). Exploring Spatial Patterns of Uranium Distribution in Ukraine.
- Gyawali, S., Techato, K., Monprapussorn, S., & Yuangyai, C. (2013). Integrating Land Use and Water Quality for Environmental based Land Use Planning for U-tapao River Basin, Thailand. *Procedia-Social and Behavioral Sciences*, *91*, 556-563.
- Haining, R. (2003) *Spatial Data Analysis: Theory and Practice.* Cambridge, England: Cambridge University Press.
- Hamid, A., Bhat, S. A., Bhat, S. U., & Jehangir, A. (2016). Environmetric techniques in water quality assessment and monitoring: a case study. *Environmental Earth Sciences*, 75(4), 1-13.
- Herojeet, R., Rishi, M. S., Lata, R., & Sharma, R. (2016). Application of environmetrics statistical models and water quality index for groundwater quality characterization of alluvial aquifer of Nalagarh Valley, Himachal, India. Sustainable Water Resources Management, 2(1), 39-53.
- Heywood, I., Cornelius, S., & Carver, S. (2002) *An introduction to Geographical Information Systems*, 2nd ed. Pearson Education Limited, England.
- Holloway, J. M., Dahlgren, R. A., Hansen, B., & Casey, W. H. (1998). Contribution of bedrock nitrogen to high nitrate concentrations in stream water. *Nature*, *395*(6704), 785-788.
- Houser, J. N., & Richardson, W. B. (2010). Nitrogen and phosphorus in the Upper Mississippi River: transport, processing, and effects on the river ecosystem. *Hydrobiologia*, *640*(1), 71-88.
- Huang, J., Li, Q., Pontius Jr, R. G., Klemas, V., & Hong, H. (2013). Detecting the dynamic linkage between landscape characteristics and water quality in a subtropical coastal

watershed, South *management*, *51*(1), 32-44.

Southeast

China. Environmental

Idris, N. (2010). *Penyelidikan dalam pendidikan*. McGraw Hill (Malaysia).

Ielpo, P., Cassano, D., Lopez, A., Pappagallo, G., Uricchio, V. F., & De Napoli, P. A. (2012). Source apportionment of groundwater pollutants in Apulian agricultural sites using multivariate statistical analyses: case study of Foggia province. *Chemistry Central Journal*, 6(2), 1.

Iglewicz, B., & Hoaglin, D. C. (1993). *How to detect and handle outliers* (Vol. 16). Milwaukee, WI: ASQC Quality Press.

Indah Water Official Portal (2016). Sewage Facts. Retrieved from <u>http://www.iwk.com.my/v/knowledgearena/ammonia</u>

Jabar, B.H. (August 09, 2010). Melaka: Longkang tercemar. *Berita Harian Online*. Retrieved from <u>http://www.bharian.com.my/bharian/articles/Melaka_Longkanterc</u> <u>emar/Article/</u>

Jain, S. K., & Singh, V. P. (2003). *Water resources systems planning* and management (Vol. 51). Elsevier.

Johnson, R. A., & Wichern, D. W. (1992). *Applied multivariate statistical analysis* (Vol. 4). Englewood Cliffs, NJ: Prentice hall.

- Joshi, P. K., Kumar, M., Paliwal, A., Midha, N., & Dash, P. P. (2009). Assessing impact of industrialization in terms of LULC in a dry tropical region (Chhattisgarh), India using remote sensing data and GIS over a period of 30 years. *Environmental monitoring and assessment*, *149*(1-4), 371-376.
- Juahir, H., Zain, S. M., Yusoff, M. K., Hanidza, T. T., Armi, A. M., Toriman, M. E., & Mokhtar, M. (2011). Spatial water quality assessment of Langat River Basin (Malaysia) using environmetric techniques. *Environmental monitoring and* assessment, 173(1-4), 625-641.

Junninen, H., Niska, H., Tuppurainen, K., Ruuskanen, J., &

Kolehmainen, M. (2004). Methods for imputation of missing values in air quality data sets. *Atmospheric Environment*, *38*(18), 2895-2907.

Kenney, J. & Keeping, E. (1962). Mathematics of statistics. Princeton, NJ. 3rd ed. Van Nostrand, p.223.

Kibena, J., Nhapi, I., & Gumindoga, W. (2014). Assessing the

relationship between water quality parameters and changes in landuse patterns in the Upper Manyame River, Zimbabwe. *Physics and Chemistry of the Earth, Parts A/B/C*, 67, 153-163.

Kim, J.O., & Mueller, C.W. (1987). Introduction to factor analysis: what it is and hot to do it. Quantitative applications in the social sciences series. Newbury Park: Sage University Press.

Koomen, E., & Borsboom-van Beurden, J. (2011). Land-use modelling

in planning practice (pp. XVI-214). Springer.

- Lakide, V. (2009). Classification of synthetic aperture radar images using particle swarm optimization technique (Doctoral dissertation, National Institute of Technology Rourkela).
- Langland, M., & Cronin, T. (Eds.). (2003). A Summary Report of Sediment Processes in Chesapeake Bay and Watershed. In Water Resources Investigations Report 03-4123. New Cumberland, PA: U S Geological Survey
- Lee, C. S. L., Li, X., Shi, W., Cheung, S. C. N., & Thornton, I. (2006). Metal contamination in urban, suburban, and country park soils of Hong Kong: a study based on GIS and multivariate statistics. *Science of the Total Environment*, *356*(1), 45-61.
- Lezzaik, K., & Milewski, A. M. (2014, October). A Global Hot Spot Analysis (Getis Ord Gi*) of Groundwater Storage Change using GRACE Satellite and GIS-Based Spatial Statistical Analysis. In 2014 GSA Annual Meeting in Vancouver, British Columbia.
- Li, S., & Zhang, Q. (2010). Spatial characterization of dissolved trace elements and heavy metals in the upper Han River (China) using multivariate statistical techniques. *Journal of Hazardous Materials*, 176(1), 579-588.
- Li, S., Gu, S., Tan, X., & Zhang, Q. (2009). Water quality in the upper Han River basin, China: the impacts of land use/land cover in riparian buffer zone. *Journal of hazardous materials*, *165*(1), 317-324.
- Li, W., Xu, B., Song, Q., Liu, X., Xu, J., & Brookes, P. C. (2014a). The identification of 'hotspots' of heavy metal pollution in soil–rice systems at a regional scale in eastern China. *Science of the Total Environment*, 472, 407-420.
- Li, Y. L., Liu, K., Li, L., & Xu, Z. X. (2012). Relationship of land use/cover on water quality in the Liao River basin, China. *Procedia Environmental Sciences*, *13*, 1484-1493.
- Li, Q., Song, J., Wang, E., Hu, H., Zhang, J., & Wang, Y. (2014b). Economic growth and pollutant emissions in China: a spatial econometric analysis. *Stochastic environmental research and risk assessment*, *28*(2), 429-442.
- Lim, W. Y., Aris, A. Z., & Praveena, S. M. (2013). Application of the chemometric approach to evaluate the spatial variation of water chemistry and the identification of the sources of pollution in Langat River, Malaysia. *Arabian Journal of Geosciences*, *6*(12), 4891-4901.
- Lim, W. Y., Aris, A. Z., & Zakaria, M. P. (2012). Spatial variability of metals in surface water and sediment in the Langat River and geochemical factors that influence their water-sediment interactions. *The Scientific World Journal*, 2012.
- Liu, C. W., Lin, K. H., & Kuo, Y. M. (2003). Application of factor analysis

in the assessment of groundwater quality in a blackfoot disease area in Taiwan. *Science of the Total Environment*, *313*(1), 77-89.

- Liu, G., Bi, R., Wang, S., Li, F., & Guo, G. (2013). The use of spatial autocorrelation analysis to identify PAHs pollution hotspots at an industrially contaminated site. *Environmental monitoring and assessment*, *185*(11), 9549-9558.
- Liu, Z., Li, Y., & Li, Z. (2009). Surface water quality and land use in Wisconsin, USA–a GIS approach. *Journal of Integrative Environmental Sciences*, 6(1), 69-89.
- Mahmud, A., & Achide, A. S. (2012). Analysis of Land Use/Land Cover Changes to Monitor Urban Sprawl in Keffi-Nigeria. *Environmental Research Journal*, 6(2), 129-134.
- Mahvi A.H. & Razazi M. (2005). Application of polyelectrolyte in turbidity removal from surface water, Am. J. Appl. Sci. **2**: 397
- Melaka State Government Official Portal (2016). District & Land Office. Retrieved from <u>http://www.melaka.gov.my/en/government/others/district-and-</u> land-offices
- Maillard, P., & Santos, N. A. P. (2008). A spatial-statistical approach for modeling the effect of non-point source pollution on different water quality parameters in the Velhas river watershed– Brazil. *Journal of Environmental Management*, *86*(1), 158-170.
- Massart, D. L., Vandeginste, B. G. M., Buydens, L. M. C., De Jong, S., Lewi, P. J., & Smeyers-Verbeke, J. (1997). Handbook of chemometrics and qualimetrics: Data handling in science and technology (Parts A and B, Vols. 20A and 20B). Elsevier: Amsterdam.
- McCutcheon, S. C., Martin, J. L., & Barnwell Jr, T. O. (1992). Water quality: in Handbook of Hydrology, Maidment, DR. Maidment. McGraw-Hill Inc., New York
- McGarigal, K., Cushman, S., & Stafford, S. (2000). Multivariate Statistics for Wildlife and Ecology Research-Springer-Verlag. New York.
- Mengistu, D. A., & Salami, A. T. (2007). Application of remote sensing and GIS inland use/land cover mapping and change detection in a part of south western Nigeria. *African Journal of Environmental Science and Technology*, *1*(5), 99-109.
- Miller, G. T., & Spoolman, S. (2012). *Environmental science*. Cengage Learning. Ministry of Resource Planning and Environment (MRPE) Official Portal (2012). *Indeks Prestasi Alam Sekitar Malaysia antara terbaik di dunia*. Retrieved from <u>http://www.kpps.sarawak.gov.my/modules/web/pages.php?lang</u> <u>=en&mod=news&sub=news view&menu id=0&sub id=109&nid</u> <u>=26&m=&y</u>=

Mishra, V. N., & Rai, P. K. (2016). A remote sensing aided multi-layer

perceptron Markov chain analysis for land use and land cover change prediction in Patna district (Bihar), India. *Arabian Journal of Geosciences*, 9(4), 1-18.

- Morse, N. B., & Wollheim, W. M. (2014). Climate variability masks the impacts of land use change on nutrient export in a suburbanizing watershed. *Biogeochemistry*, *121*(1), 45-59.
- Murray, A. T., McGuffog, I., Western, J. S., & Mullins, P. (2001). Exploratory spatial data analysis techniques for examining urban crime implications for evaluating treatment. *British Journal of criminology*, *41*(2), 309-329.
- Mustapha, A., Aris, A. Z., Juahir, H., Ramli, M. F., & Kura, N. U. (2013). River water quality assessment using environmentric techniques: case study of Jakara River Basin. *Environmental Science and Pollution Research*, 20(8), 5630-5644.
- Mustapha, A., & Aris, A. Z. (2012). Spatial aspects of surface water quality in the Jakara Basin, Nigeria using chemometric analysis. *Journal of Environmental Science and Health, Part A*, 47(10), 1455-1465.
- Mustapha, A., Aris, A. Z., Ramli, M. F., & Juahir, H. (2012). Spatial temporal variation of surface water quality in the downstream region of the Jakara River, north-western Nigeria: A statistical approach. *Journal of Environmental Science and Health, Part A*, 47(11), 1551-1560.
- National Geographic Portal (2016). Water Pollution. Retrieved from http://environment.nationalgeographic.com/environment/freshwa ter/pollution/
- Nasbah, N.N. (January 23, 2010). Sungai Melaka Tercemar. *Utusan* Online. Retrieved from <u>http://ww1.utusan.com.my/utusan/info.asp?y=2010&dt=0123&se</u> c=Selatan&pg=ws_01.htm
- Otokunefor, T. V., & Obiukwu, C. (2005). Impact of refinery effluent on the physicochemical properties of a water body in the Niger delta. *Applied ecology and environmental research*, *3*(1), 61-72.
- Otukei, J. R., & Blaschke, T. (2010). Land cover change assessment using decision trees, support vector machines and maximum likelihood classification algorithms. *International Journal of Applied Earth Observation and Geoinformation*, *12*, S27-S31.
- Papaioannou, A., Dovriki, E., Rigas, N., Plageras, P., Rigas, I., Kokkora, M., & Papastergiou, P. (2010a). Assessment and modelling of groundwater quality data by environmetric methods in the context of public health. *Water resources management*, 24(12), 3257-3278.

Papaioannou, A., Mavridou, A., Hadjichristodoulou, C., Papastergiou,

P., Pappa, O., Dovriki, E., & Rigas, I. (2010b). Application of multivariate statistical methods for groundwater physicochemical and biological quality assessment in the context of public

health. *Environmental monitoring and assessment*, 170(1-4), 87-97.

- Parsa, V. A., Yavari, A., & Nejadi, A. (2016). Spatio-temporal analysis of land use/land cover pattern changes in Arasbaran Biosphere Reserve: Iran. *Modeling Earth Systems and Environment*, 2(4), 178.
- Persona Metro Official Portal (2016). Rehabilitation and Beautification of Sungai Melaka. Retrieved from http://www.pesona.com.my/sungai-melaka.php
- Pontius Jr, R. G., & Millones, M. (2011). Death to Kappa: birth of quantity disagreement and allocation disagreement for accuracy assessment. *International Journal of Remote Sensing*, 32(15), 4407-4429.
- PPPP (Persatuan Pengguna Pulau Pinang) Oficial Portal (April 23, 2015). CAP kesal sungai Melaka teruk tercemar. Retrieved from <u>http://www.pengguna.org.my/index.php/pembangunan/alam-sekitar/884-cap-kesal-sungai-melaka-teruk-tercemar</u>
- Proakis, J. & Manolakis, D. (2007). Tratamiento digital de senales. Pearson Education, Vol. 1, 4th Ed. Espana.
- Rizo-Decelis, L. D., Pardo-Igúzquiza, E., & Andreo, B. (2017). Spatial prediction of water quality variables along a main river channel, in presence of pollution hotspots. *Science of The Total Environment*, 605, 276-290.
- Rosli, S. N., Aris, A. Z., & Majid, N. M. (2015). Spatial variation assessment of Malacca River water quality using multivariate statistical analysis. *Malaysian Applied Biology*, 44(1), 13-18.
- Rügner, H., Schwientek, M., Beckingham, B., Kuch, B., & Grathwohl, P. (2013). Turbidity as a proxy for total suspended solids (TSS) and particle facilitated pollutant transport in catchments. *Environmental earth sciences*, 69(2), 373-380.
- Ruiz-Luna, A., & Berlanga-Robles, C. A. (2003). Land use, land cover changes and coastal lagoon surface reduction associated with urban growth in northwest Mexico. *Landscape Ecology*, *18*(2), 159-171.
- Salleh, D. (2002). Sungai sebagai alternatif pelancongan di Negeri Kedah. In: *Kedah 100 tahun 1900-200 isu-isu politik dan sosioekonomi*, 227-236. Penerbit Universiti Utara Malaysia, Sintok.
- Said, A., Stevens, D. K., & Sehlke, G. (2004). An innovative index for evaluating water quality in streams. *Environmental management*, *34*(3), 406-414.
- Shi, P., Zhang, Y., Li, Z., Li, P., & Xu, G. (2017). Influence of land use and land cover patterns on seasonal water quality at multispatial scales. *Catena*, *151*, 182-190

Shi, W., Xia, J., & Zhang, X. (2016). Influences of anthropogenic

activities and topography on water quality in the highly regulated Huai River basin, China. *Environmental Science and Pollution Research*, 1-15.

- Shazili, N. A. M., Yunus, K., Ahmad, A. S., Abdullah, N., & Rashid, M. K. A. (2006). Heavy metal pollution status in the Malaysian aquatic environment. *Aquatic Ecosystem Health & Management*, 9(2), 137-145.
- Shin, J. Y., Artigas, F., Hobble, C., & Lee, Y. S. (2013). Assessment of anthropogenic influences on surface water quality in urban estuary, northern New Jersey: multivariate approach. *Environmental monitoring and assessment*, *185*(3), 2777-2794.
- Shrestha, S., & Kazama, F. (2007). Assessment of surface water quality using multivariate statistical techniques: A case study of the Fuji river basin, Japan. *Environmental Modelling & Software*, 22(4), 464-475.
- Simeonov, V., Einax, J., Stanimirova, I., & Kraft, J. (2002). Environmetric modelling and interpretation of river water monitoring data. *Analytical and bioanalytical chemistry*, 374(5), 898-905.
- Sinar Official Portal (April 15, 2012). Sungai tercemar berubah wajah. *Sinar Online*. Retrieved from <u>http://www.sinarharian.com.my/edisi/melaka-ns/sungai-</u> <u>tercemar-berubah-wajah-1.40085</u>
- Singh, A. K. (2003). Modelling land use land cover changes using cellular automata in a geo-spatial environment. International Institute for Geo-Information Science and Earth Observation. Vol. Master of Science, Enscheda, the Netherlands.
- Singh, K. P., Malik, A., Singh, V. K., Mohan, D., & Sinha, S. (2005). Chemometric analysis of groundwater quality data of alluvial aquifer of Gangetic plain, North India. *Analytica Chimica Acta*, 550(1), 82-91.
- Singh, K. P., Malik, A., Mohan, D., & Sinha, S. (2004). Multivariate statistical techniques for the evaluation of spatial and temporal variations in water quality of Gomti River (India)—a case study. *Water research*, *38*(18), 3980-3992.
- Singh, K. P., Basant, N., & Gupta, S. (2011). Support vector machines in water quality management. *Analytica chimica acta*, 703(2), 152-162.
- Singh, S., Raju, N. J., & Nazneen, S. (2015). Environmental risk of heavy metal pollution and contamination sources using multivariate analysis in the soils of Varanasi environs, India. *Environmental monitoring and assessment*,187(6), 1-12.
- Sinha, P., & Kumar, L. (2013): Markov land cover change modelling using pairs of time series satellite images. *Photogramm. Eng. Remote Sens*, 79 (11): 11.

- Subedi, P., Subedi, K., & Thapa, B. (2013). Application of a hybrid cellular automaton Markov (CA-Markov) Model in land-use change prediction: a case study of saddle creek drainage Basin, Florida. *Applied Ecology and Environmental Sciences*, *1*(6), 126-132.
- Sun, G., Hoff, S. J., Zelle, B. C., & Smith, M. A. (2008). Development and comparison of backpropagation and generalized regression neural network models to predict diurnal and seasonal gas and PM10 concentrations and emissions from swine buildings. In 2008 Providence, Rhode Island, June 29–July 2, 2008 (p. 1). American Society of Agricultural and Biological Engineers.
- Sun, L. (2015). Environmental baseline evaluation of lead in shallow groundwater based on statistical and spatial outlier identification. *Chinese Journal of Geochemistry*, *34*(3), 416-421.
- Sun, Z., Ma, R., & Wang, Y. (2009). Using Landsat data to determine land use changes in Datong basin, China. *Environmental* geology, 57(8), 1825-1837.
- Sundaray, S. K. (2010). Application of multivariate statistical techniques in hydrogeochemical studies—a case study: Brahmani–Koel River (India). *Environmental monitoring and assessment*, 164(1-4), 297-310.
- Umar, A., Umar, R., & Ahmad, M. S. (2001). Hydrogeological and hydrochemical framework of regional aquifer system in Kali-Ganga sub-basin, India. *Environmental Geology*, *40*(4-5), 602-611.
- UNESCO Official Portal (2016). *Melaka and George Town, Historic Cities of the Straits of Malacca.* Retrieved from <u>http://whc.unesco.org/en/list/1223</u>
- Viera, A. J., & Garrett, J. M. (2005). Understanding interobserver agreement: the kappa statistic. *Fam Med*, *37*(5), 360-363.
- Vrebos, D., Beauchard, O., & Meire, P. (2017). The impact of land use and spatial mediated processes on the water quality in a river system. *Science of The Total Environment*, 601, 365-373.
- Wahl, M. H., McKellar, H. N., & Williams, T. M. (1997). Patterns of nutrient loading in forested and urbanized coastal streams. *Journal of Experimental Marine Biology and Ecology*, 213(1), 111-131.
- Wang F., Ding Y., Ge L., Ren H., & Ding L. (2010). Effect of high strength ammonia nitrogen acclimation on sludge activity in sequencing batch reactor, J. Environ. Sci. 22 (11): 1683.
- Weng, Q.H. (2010). *Remote Sensing and GIS integration*. McGraw-Hill, New York. 424p.
- White, R., & Engelen, G. (2000). High-resolution integrated modelling of the spatial dynamics of urban and regional systems. *Computers, environment and urban systems*, 24(5), 383-400.
- Wilson, C. O. (2015). Land use/land cover water quality nexus:

quantifying anthropogenic influences on surface water quality. *Environmental monitoring and assessment*, 187(7), 1-23.

- World Health Organization (WHO) (2014). Global burden of disease: Impacts of poor water, sanitation and hygiene. Retrieved from <u>http://www.who.int/water_sanitation_health/gbd_poor_water/en/</u>
- WHO (2004). *Guidelines for Drinking-Water Quality*, vol. 1, World Health Organization, Geneva, Switzerland, 3rd edition.
- WSDE (Washington State Department of Ecology) (2002). Setting standards for the bacteriological quality of Washington's surface water: Draft discussion paper and literature summary, Publication No. 00-10-072. Washington DC.
- Ye, B., & Bai, Z. (2007, August). Simulating land use/cover changes of Nenjiang County based on CA-Markov model. In *International Conference on Computer and Computing Technologies in Agriculture* (pp. 321-329). Springer US.
- Yetilmezsoy, K., & Demirel, S. (2008). Artificial neural network (ANN) approach for modeling of Pb (II) adsorption from aqueous solution by Antep pistachio (Pistacia Vera L.) shells. *Journal of hazardous materials*, *153*(3), 1288-1300.
- Yildiz, S., & Doker, M. F. (2016). Monitoring urban growth by using segmentation classification of multispectral Landsat images in Izmit, Turkey. *Environmental monitoring and assessment*, 188(7), 1-12.
- Yu, S., Xu, Z., Wu, W., & Zuo, D. (2016). Effect of land use types on stream water quality under seasonal variation and topographic characteristics in the Wei River basin, China. *Ecological indicators*, 60, 202-212.
- Yulianto, F., Prasasti, I., Pasaribu, J. M., Fitriana, H. L., Haryani, N. S., & Sofan, P. (2016). The dynamics of land use/land cover change modeling and their implication for the flood damage assessment in the Tondano watershed, North Sulawesi, Indonesia. *Modeling Earth Systems and Environment*, 2(1), 1-20.
- Zakinan, N.F. (Disember 30, 2015). Kekurangan oksigen punca ikan mati di sungai Melaka. *Berita Harian Online*. Retrieved from http://www.bharian.com.my/node/109663
- Zhai, X., Xia, J., & Zhang, Y. (2014). Water quality variation in the highly disturbed Huai River Basin, China from 1994 to 2005 by multistatistical analyses. *Science of the Total Environment*, *496*, 594-606.
- Zhao, J., Lin, L., Yang, K., Liu, Q., & Qian, G. (2015). Influences of land use on water quality in a reticular river network area: A case study in Shanghai, China. *Landscape and Urban Planning*, *137*, 20-29.
- Zhao, X., Huang, X., & Liu, Y. (2012). Spatial autocorrelation analysis of

Chinese inter provincial industrial chemical oxygen demand discharge. *International journal of environmental research and public health*, 9(6), 2031-2044.

- Zheng, W., Li, X., Yin, L., & Wang, Y. (2015). Spatiotemporal heterogeneity of urban air pollution in China based on spatial analysis. *Rendiconti Lincei*, 1-6.
- Zou, B., Peng, F., Wan, N., Mamady, K., & Wilson, G. J. (2014). Spatial cluster detection of air pollution exposure inequities across the United States. *PloS one*, *9*(3), e91917.

