



**UNIVERSITI PUTRA MALAYSIA**

***HYDROGEOCHEMISTRY OF THE LOWER KELANTAN BASIN,  
MALAYSIA***

**ANUAR BIN SEFIE**

**FPAS 2018 13**



**HYDROGEOCHEMISTRY OF THE LOWER KELANTAN BASIN,  
MALAYSIA**

By

**ANUAR BIN SEFIE**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfillment of the Requirements for the degree of Master of Science**

**May 2018**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

## HYDROGEOCHEMISTRY OF THE LOWER KELANTAN BASIN, MALAYSIA

By

**ANUAR BIN SEFIE**

May 2018

**Chairman: Ahmad Zaharin Aris, PhD**  
**Faculty: Environmental Studies**

Groundwater plays a crucial role in supporting the physical and economic development of any region, especially in areas that are experiencing shortage of surface water resources. Due to industrialization, population growth and intensive agriculture, the dependence on groundwater resources has increased dramatically from time to time resulting in quantity and quality deterioration, consequently, affecting human health and plant growth. A comprehensive hydrogeochemical study is essential in understanding the evolution of groundwater in a multi-layered aquifer system. The aims of this study were to construct a conceptual model, identify the factors controlling the chemistry of the groundwater, assess the hydrogeochemical facies of the groundwater, and evaluate the suitability of the groundwater for drinking and agricultural purposes. 101 groundwater samples were collected and analyzed for physico-chemical parameters such as the pH, the electrical conductivity (EC), the total dissolved solid (TDS),  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{HCO}_3^-$ ,  $\text{CO}_3^{2-}$ ,  $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{Fe}^{2+}$  and  $\text{Mn}^{2+}$ . Based on lithological data correlation of exploration wells, three aquifer layers were delineated; a shallow, an intermediate and a deep. PCA demonstrated that Factor 1 (VF1) for the shallow aquifer represents 41.23% of the total variance. The variables  $\text{Na}^+$ ,  $\text{Cl}^-$ , the TDS, the EC,  $\text{Mg}^{2+}$ , and  $\text{SO}_4^{2-}$  showed strong positive factor loadings and indicate the influence of remnant saltwater in the aquifer system. Factor 2 (VF2) which accounts for 13.23% of the total variance exhibited a strong positive loading for  $\text{HCO}_3^-$  and suggests an origin from the weathering of carbonate. Factor 3 (VF3) was associated with 9.33% of the total variance and had strong positive loadings for  $\text{Fe}^{2+}$  and  $\text{Mn}^{2+}$ , while  $\text{NO}_3^-$  showed a strong positive loading in Factor 4 (VF4) representing 7.23% of the variance. The  $\text{Fe}^{2+}$  and  $\text{Mn}^{2+}$  in Factor 3 (VF3) are attributed to weathering of iron and manganese bearing minerals, whereas  $\text{NO}_3^-$  is assigned to anthropogenic sources like sewage and fertilizer usage. The four principal components (VF1, VF2, VF3 and VF4) account for 71.11% of the total variance for the shallow aquifer. FA for the intermediate aquifer resulted in three significant factors accounting for 75.92% of the total variance. Factor 1 (VF1) demonstrated strong positive loadings for the TDS, the EC,  $\text{Cl}^-$ ,  $\text{Na}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$  and  $\text{Ca}^{2+}$  representing 53.73% of total

variance, and attributed dissolution of minerals from seawater remnant. Factor 2 (VF2) which explained 14.55 % of the variance showed a strong positive loading for pH and a strong negative loading for  $\text{Fe}^{2+}$ . This indicated the pH was affected due to enhanced dissolution and  $\text{Fe}^{2+}$  was assigned to weathering of iron bearing minerals. The variable of  $\text{SO}_4^{2-}$  showed a strong negative loading in Factor 3 (VF3) that explained 7.64% of the total variance, and it is attributed to oxidation and dissolution of sulfate-rich marine deposits. For the deep aquifer, Factor 1 (VF1) showed strong positive loadings for  $\text{Cl}^-$ , the EC, the TDS,  $\text{Na}^+$ ,  $\text{Fe}^{2+}$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$ , and a strong negative loading for pH, accounting for 55.08% of the total variance. Factor 2 (VF2) accounted for 12.13% of the total variance, with strong positive loadings for carbonate that are explained by calcium carbonate weathering processes. Factor 3 (VF3) showed strong positive loadings for pH,  $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Fe}^{2+}$  and represented 8.39% of the total variance. This was attributed to the dissolution and weathering of rock forming minerals. Factor 4 (VF4) representing 7.38% of the total variance showed a strong negative loadings for  $\text{SO}_4^{2-}$  reflecting oxidation and dissolution of sulfate-rich marine deposits. The abundance of the major cations followed the order:  $\text{Na}^+ > \text{Ca}^{2+} > \text{Mg}^{2+} > \text{K}^+$ , while that of anions was  $\text{HCO}_3^- > \text{Cl}^- > \text{SO}_4^{2-} > \text{CO}_3^{2-}$ . The prominent groundwater facies in the shallow aquifer are Ca- $\text{HCO}_3$  and Na- $\text{HCO}_3$  and indicate the presence of fresh water and mix-water types. The Na- $\text{HCO}_3$  and Na-Cl are the main water types in the intermediate aquifer and exhibit the presence of mix-water and saltwater types. The Na- $\text{HCO}_3$  is the dominant water type in the deep aquifer and is probably due to the consumption of  $\text{H}^+$  in the chemical weathering of feldspars to clay minerals. The Gibbs diagram stipulates that the groundwater in the aquifers are mostly affected by rock dominance, regardless of aquifer depth. Based on US Salinity Laboratory (USSL) diagram, the shallow groundwater samples fall in the C1-C3 and S1, and is suitable for irrigation of most crops. Samples from the intermediate aquifer plot in the C1-C4 and S1-S4 and are not suitable for irrigation of crops. The samples from deep aquifer fall in the C1-C3 and S1 and some samples are unsuitable for irrigation of crops. Some groundwater samples taken from the intermediate and the deep aquifers that fall into S4 zone are categorized as very high salinity hazard samples. The groundwater samples obtained from areas located more than 8 km from coast for all aquifers are suitable for crop irrigation. A high SAR value in the groundwater leads to the formation of an alkaline soil which reduces soil permeability due to hardening and compaction when dry. The physical structure of the soil crumbles and renders it unsuitable for plant growth. The study reveals that the groundwater quality from the shallow and the deep aquifers are better for domestic and agricultural purposes relative to groundwater from the intermediate aquifer for aquifer located less than 8 km from coast. These results provide a baseline data and offers insights on the status of groundwater quality in the Lower Kelantan Basin. The study also determined the factors controlling the quality of groundwater in each aquifer. Groundwater quality in the Lower Kelantan Basin is governed by anthropogenic activities and natural factors.

**Keywords:** Lower Kelantan Basin, conceptual model, multivariate statistical analyses, hydrogeochemical facies, saturation index, groundwater quality assessment

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

## HIDROGEOKIMIA DI KAWASAN LEMBANGAN KELANTAN HILIR, MALAYSIA

Oleh

**ANUAR BIN SEFIE**

Mei 2018

**Pengerusi: Ahmad Zaharin Aris, PhD**  
**Fakulti: Pengajian Alam Sekitar**

Air tanah memainkan peranan penting dalam menyokong pembangunan fizikal dan ekonomi di sesuatu kawasan terutamanya di kawasan yang mengalami kekurangan sumber air permukaan. Akibat daripada aktiviti perindustrian, pertumbuhan penduduk dan pertanian secara intensif, kebergantungan kepada sumber air tanah telah meningkat secara mendadak dari semasa ke semasa yang menyebabkan kemerosotan kuantiti dan kualiti yang seterusnya akan menjejaskan kesihatan manusia dan tumbesaran tanaman. Kajian hidrogeokimia secara menyeluruh adalah penting dalam memahami evolusi air tanah dalam sistem akuifer pelbagai lapisan. Tujuan kajian ini adalah untuk membina model konseptual, mengenalpasti faktor yang mengawal geokimia air tanah, menilai hidrogeokimia air tanah dan kesesuaiannya untuk kegunaan minuman dan pertanian. Sebanyak 101 sampel air tanah telah diambil dan dianalisis bagi parameter fiziko-kimia seperti pH, kekonduksian elektrik (EC), jumlah pepejal terlarut (TDS),  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{HCO}_3^-$ ,  $\text{CO}_3^{2-}$ ,  $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{Fe}^{2+}$  dan  $\text{Mn}^{2+}$ . Berdasarkan korelasi data litologi telaga eksplorasi, sebanyak tiga lapisan akuifer utama dikenalpasti; akuifer cetek, pertengahan dan dalam. Analisis komponen prinsip (PCA) menunjukkan Faktor 1 (VF1) bagi akuifer cetek mewakili 41.23% daripada jumlah varian. Pembolehubah  $\text{Na}^+$ ,  $\text{Cl}^-$ , TDS, EC,  $\text{Mg}^{2+}$  dan  $\text{SO}_4^{2-}$  mempunyai beban positif tinggi yang menunjukkan pengaruh tinggalan air masin dalam sistem akuifer. Faktor 2 (VF2), menyumbang 13.23% daripada jumlah varian dan mempunyai beban positif yang tinggi bagi  $\text{HCO}_3^-$  di mana ia terhasil daripada proses perlarutan karbonat. Faktor 3 (VF3) mewakili 9.33% daripada jumlah varian yang mempunyai beban positif bagi  $\text{Fe}^{2+}$  dan  $\text{Mn}^{2+}$ , manakala  $\text{NO}_3^-$  mempunyai beban positif yang tinggi bagi Faktor 4 (VF4) iaitu mewakili 7.23% varian.  $\text{Fe}^{2+}$  dan  $\text{Mn}^{2+}$  bagi Faktor 3 (VF3) adalah disebabkan oleh luluhawa mineral besi dan mangan manakala  $\text{NO}_3^-$  berpunca daripada aktiviti manusia seperti sisa kumbahan dan penggunaan baja. Empat komponen prinsip (VF1, VF2, VF3 dan VF4) mencakupi 71.11% daripada jumlah varian bagi akuifer cetek. FA bagi akuifer pertengahan menjana tiga faktor penting yang menyumbang 75.92% daripada jumlah varian. Faktor 1 (VF1) mempunyai beban positif yang tinggi bagi TDS, EC,  $\text{Cl}^-$ ,  $\text{Na}^+$ ,  $\text{Mg}^{2+}$ ,

$K^+$  dan  $Ca^{2+}$  mewakili 53.73% daripada jumlah varian yang disebabkan oleh perlarutan mineral kesan tinggalan air laut kuno. Faktor 2 (VF2) mewakili 14.55% daripada jumlah varian dan mempunyai beban positif yang tinggi bagi pH dan berkadar songsang dengan  $Fe^{2+}$ . Ini menunjukkan penurunan nilai pH akan meningkatkan perlarutan  $Fe^{2+}$  hasil proses luluhawa mineral besi. Pembolehubah  $SO_4^{2-}$  menunjukkan beban negatif yang tinggi bagi Faktor 3 (VF3) yang mewakili 7.64% daripada jumlah varian dan ia dikaitkan dengan proses pengoksidaan dan perlarutan enapan marin yang kaya sulfat. Bagi akuifer dalam, Faktor 1 (VF1) menunjukkan beban positif yang tinggi bagi  $Cl^-$ , EC, TDS,  $Na^+$ ,  $Fe^{2+}$ ,  $K^+$ ,  $Mg^{2+}$  dan  $Ca^{2+}$ , dan beban negatif tinggi bagi pH dengan menyumbang 55.08%. Faktor 2 (VF2) menyumbang 12.13% daripada jumlah varian dengan bebanan positif yang tinggi bagi karbonat yang menjelaskan proses luluhawa kalsium karbonat. Faktor 3 (VF3) menunjukkan beban positif tinggi bagi pH,  $Na^+$ ,  $K^+$  dan  $Fe^{2+}$  yang mewakili 8.39% daripada jumlah varian. Ini mungkin disebabkan oleh perlarutan dan luluhawa mineral batuan. Faktor 4 (VF4) yang mewakili 7.38% daripada jumlah varian menunjukkan beban negatif tinggi bagi  $SO_4^{2-}$  yang juga menggambarkan proses pengoksidaan dan perlarutan enapan marin kaya sulfat. Kelimpahan kation utama mengikut susunan ialah  $Na^+ > Ca^{2+} > Mg^{2+} > K^+$ , sementara anion adalah  $HCO_3^- > Cl^- > SO_4^{2-} > CO_3^{2-}$ . Fasies air tanah yang utama bagi akuifer cetek adalah Ca- $HCO_3$  dan Na- $HCO_3$  yang menunjukkan kehadiran air tawar dan campuran air tawar-masin. Na- $HCO_3$  dan Na-Cl adalah jenis air utama dalam akuifer pertengahan yang menunjukkan percampuran air tawar-masin. Na- $HCO_3$  adalah jenis air yang dominan bagi akuifer dalam yang mungkin disebabkan oleh penggunaan  $H^+$  dalam luluhawa kimia feldspar kepada mineral lempung. Gambarajah Gibbs menunjukkan air tanah dipengaruhi oleh dominasi batuan tanpa mengira kedalaman akuifer. Berdasarkan kepada gambarajah US Salinity Laboratory (USSL) pula, semua sampel air tanah bagi akuifer cetek berada dalam zon C1-C3 dan S1 yang sesuai digunakan bagi kebanyakan jenis tanaman untuk tujuan pengairan. Sampel bagi akuifer pertengahan pula, sebahagian besar sampel berada dalam zon C1-C4 dan S1-S4 di mana ia tidak sesuai untuk pengairan tanaman. Manakala sebahagian sampel air tanah daripada akuifer dalam berada dalam zon C1-C3 dan S1 tidak sesuai untuk pengairan tanaman. Beberapa sampel air tanah yang diperolehi dari akuifer pertengahan dan dalam berada dalam zon S4 dan dikategorikan sebagai bahaya kemasinan yang sangat tinggi. Semua sampel air tanah yang diambil dari semua akuifer dan berada di kawasan lebih 8 km dari garis pantai adalah sesuai untuk pengairan tanaman. Nilai nisbah serapan sodium (SAR) yang tinggi di dalam air tanah akan memberi kesan kepada pembentukan tanah beralkali yang mengurangkan kebolehtelapan tanah kerana pengerasan dan pepadatan apabila kering. Struktur fizikal tanah akan rosak menjadikannya tidak sesuai untuk pertumbuhan tanaman. Kajian ini menunjukkan kualiti air tanah dari akuifer cetek dan dalam adalah sesuai untuk tujuan domestik dan pertanian berbanding dengan akuifer pertengahan terutama yang terletak kurang 8 km dari garis pantai. Hasil kajian ini menyediakan data asas dan memberi gambaran yang lebih jelas mengenai status semasa kualiti air tanah di Lembangan Kelantan Hilir. Ia juga dapat menentukan faktor utama yang mengawal kualiti air bagi setiap akuifer. Ini dapat disimpulkan bahawa kualiti air tanah di Lembangan Kelantan Hilir dikawal oleh aktiviti antropogenik dan juga faktor semula jadi.

**Kata kunci:** Lembangan Kelantan Hilir, model konseptual, analisis statistik multivariat, fasies hidrogeokimia, indeks ketepuan, penilaian kualiti air tanah

## ACKNOWLEDGEMENTS

In the name of Allah, the most beneficent, the most eternally merciful. Alhamdulillah, I am grateful and thankful to the Almighty Allah S.W.T for giving me strength, courage, patience and determination in preparing and completing this research successfully. I would like to express my deepest gratitude to my supervisor, Prof. Dr. Ahmad Zaharin Aris for his patience, continuous supervision, assistance, motivation, invaluable advices, guidance and suggestions. I would also like to thank the other my committee members, Associate Prof. Dr. Mohammad Firuz Ramli and Dr. Saim Suratman for giving valuable suggestions and continuous encouragement during the preparation of this thesis. I would like to express my honest thanks to Dr. Tahoora Sheikhy Narany and Dr. Hafizan Juahir for their valuable guide, advices and helps.

In addition, I wish to thank all the government agencies especially Tuan Haji Mohamad Yusof bin Che Sulaiman from Minerals and Geosciences Department (MGD) of Kelantan, Air Kelantan Sdn Bhd (AKSB), Chemistry Department of Malaysia, Ministry of Agriculture (MOA), Malaysia Meteorological Department (MMD), Drainage and Irrigation Department (DID) and Department of Environment (DOE) that have been assisting me in groundwater sampling and analyses activities. They also provided secondary groundwater quality, land use, hydrological and environmental data as well as groundwater analyses throughout the study. I would also like to acknowledge the Ministry of Natural Resources and Environment (NRE) for financing this research. My deepest appreciation to the Public Services Commission of Malaysia for the financial support.

I would like to thank all the research officers and assistants from National Hydraulic Research Institute of Malaysia (NAHRIM) for their continuous support, reminders, encouragement, advice, ideas, assistances and for lending their hands especially Mr. Ismail Tawnie, Mr. Mohd Khairul Nizar Shamsuddin, Mr. Azrul Normi Idris, Mr. Syaiful Bahren Saadudin, Mr. Arshad Osman, Mr. Mohammad Huzaifah Wahap and Mr. Mohd Ali Sarudin who have helped and accompanied in groundwater sampling and *in-situ* analyses.

Last but not least, I wish to thank Mr. Ridzuan Isa and all my friends at the Faculty of Environmental Studies, particularly Ms. Wee Sze Yee, Tuan Mohd Fauzan, Dr. Noorain, Dr. Shahcristiani, Ms. Farhana, Ms. Hanisah, Ms. Afifah, Ms. Nasyitah, Dr. Looi Ley Juen and Ms. Lim Ai Phing. My appreciation is extended to my beloved parents, Haji Sefie Ibrahim and Hajjah Halimah Awang for their constant support, encouragement and prayers. Lastly, the warmest thank to my dearest wife, Mrs Haslinda Hasanuddin for her sincere support, advice and inspiration. There are no words that I can give, other than how thankful I am for all the assistance in making this research a reality. I believe only Allah can repay the kindness that each one have given me in order for my dreams to come true. Honestly, these experiences have been nothing but the best in my life. Thank you to all that have directly or indirectly contributed to the completion of this research.



I certify that a Thesis Examination Committee has met on 31 May 2018 to conduct the final examination of Anuar bin Sefie on his thesis entitled "Hydrogeochemistry of the Lower Kelantan Basin, Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

**Normala binti Halimoon, PhD**

Senior Lecturer  
Faculty of Environmental Studies  
Universiti Putra Malaysia  
(Chairman)

**Wan Nor Azmin bin Sulaiman, PhD**

Professor  
Faculty of Environmental Studies  
Universiti Putra Malaysia  
(Internal Examiner)

**Ismail bin Yusoff, PhD**

Professor  
University of Malaya  
Malaysia  
(External Examiner)



**RUSLI HAJI ABDULLAH, PhD**  
Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 30 July 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Ahmad Zaharin Aris, PhD**

Professor  
Faculty of Environmental Studies  
Universiti Putra Malaysia  
(Chairman)

**Mohammad Firuz Ramli, PhD**

Associate Professor  
Faculty of Environmental Studies  
Universiti Putra Malaysia  
(Member)

**Saim Suratman, PhD**

Director  
Hydrogeology Research Centre  
National Hydraulic Research Institute of Malaysia (NAHRIM)  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by Graduate Student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Anuar bin Sefie (GS35394)

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to

Signature: \_\_\_\_\_

Name of  
Chairman of  
Supervisory  
Committee:

Professor Dr. Ahmad Zaharin Aris

Signature: \_\_\_\_\_

Name of  
Member of  
Supervisory  
Committee:

Associate Professor Dr. Mohammad Firuz Ramli

Signature: \_\_\_\_\_

Name of  
Member of  
Supervisory  
Committee:

Dr. Saim Suratman

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiii
<b>LIST OF FIGURES</b>	xiv
<b>LIST OF ABBREVIATIONS</b>	xvii
<b>CHAPTER</b>	<b>1</b>
<b>1 INTRODUCTION</b>	<b>1</b>
1.1 Background of the Study	1
1.2 Problem Statement	2
1.3 Objectives of the Study	3
1.4 Scope of the Study	3
1.5 Significance of the Study	4
1.6 Research Framework of the Study	5
1.7 Thesis Outline	7
<b>2 LITERATURE REVIEW</b>	<b>8</b>
2.1 Introduction	8
2.2 Geology and Hydrogeology	9
2.3 Groundwater Utilisation	18
2.4 Sources of Groundwater Contamination	20
2.4.1 Geogenic Sources	20
2.4.2 Seawater Intrusion	21
2.4.3 Agricultural Sources	22
2.4.4 Municipal and Industrial Sources	23
2.5 Effects of Groundwater Contamination and Over Abstraction	23
2.5.1 Human Health	24
2.5.2 Plant Growth	25
2.5.3 Over-Abstraction of Groundwater	26
2.6 The Importance of Groundwater Quality Assessment	26
2.7 Methods Used for Groundwater Quality Assessment	27
2.7.1 Saturation Index	27
2.7.2 Geographic Information System (GIS)	28
2.7.3 Application of Graphical Representations	28
2.7.4 Multivariate Statistical Analysis	29

<b>3</b>	<b>MATERIAL AND METHODS</b>	<b>31</b>
3.1	Description of the Study Area	31
3.1.1	Location	31
3.1.2	Drainage	31
3.1.3	Topography	34
3.1.4	Land Use	34
3.1.5	Climate	35
3.2	Materials and Methods	38
3.2.1	Aquifer Classification and Extent	38
3.2.2	Groundwater Sampling and Analytical Procedures	40
3.2.3	Groundwater Quality Assessment	42
3.2.4	Groundwater Quality Distribution Maps	42
3.3	Data Pre-treatment and Analyses	43
3.3.1	Pearson Correlation	45
3.3.2	Discriminant Analysis (DA)	45
3.3.3	Principle Component Analysis (PCA)/Factor Analysis (FA)	46
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	<b>48</b>
4.1	Geological Setup and Factors Controlling the Variations in Groundwater Quality	48
4.1.1	Geological Conceptual Model	48
4.1.2	Correlation Coefficient	54
4.1.3	Discriminant Analysis (DA)	58
4.1.4	Principle Component Analysis (PCA)/Factor Analysis (FA)	60
4.2	Groundwater Hydrogeochemistry	64
4.2.1	Geochemical Evaluation	64
4.2.2	Hydrochemical Facies	86
4.2.3	Saturation Index (SI) and Mineral Equilibrium	90
4.2.4	Gibbs Classification	94
4.3	Groundwater Quality Assessment for Drinking and Agricultural Irrigation Purposes	96
4.3.1	Groundwater Quality Assessment for Drinking Purposes	96
4.3.2	Groundwater Suitability for Irrigation Purposes	99
4.3.2.1	Sodium Adsorption Ratio (SAR)	99
4.3.2.2	Sodium Percentage (Na%)	100
4.3.2.3	Residual Sodium Carbonate (RSC)	103
4.3.2.4	Permeability Index (PI)	103
4.3.2.5	Magnesium adsorption ratio (MAR)	105
4.3.2.6	Kelly's Ratio (KR)	105
4.4	Discussion on the Hydrogeochemistry of the Lower Kelantan Basin	105

<b>5</b>	<b>CONCLUSION AND RECOMMENDATION</b>	109
5.1	Conclusions	109
5.2	Recommendations for Future Research	112

	<b>REFERENCES</b>	114
	<b>APPENDICES</b>	143
	<b>BIODATA OF STUDENTS</b>	152
	<b>LIST OF PUBLICATIONS/PRESENTED PAPERS</b>	153



## LIST OF TABLES

Table		Page
2.1	Summary of hydrogeology and hydrogeochemical studies in Lower Kelantan Basin	14
2.2	Total groundwater abstraction by AKSB wellfields in 2012	20
2.3	Some sources of ions in groundwater	21
3.1	The proportion of land use types in the Lower Kelantan Basin	35
4.1	Correlation coefficient of major cations and anions of groundwater in shallow aquifer	55
4.2	Correlation coefficient of major cations and anions of groundwater in intermediate aquifer	56
4.3	Correlation coefficient of major cations and anions of groundwater in deep aquifer	57
4.4	The confusion matrix for the aquifer classification in the Lower Kelantan Basin (DA-deep aquifer, IA-intermediate aquifer and SA-shallow aquifer)	60
4.5	Factor loading after Varimax rotation, eigenvalues and cumulative variance values of shallow aquifer (n = 427)	61
4.6	Factor loading after Varimax rotation, eigenvalues and cumulative variance values of intermediate aquifer (n = 237)	62
4.7	Factor loading after Varimax rotation, eigenvalues and cumulative variance values of deep aquifer (n = 412)	63
4.8	Descriptive statistics for hydrochemical parameters for aquifers in the Lower Kelantan Basin	83
4.9	Summary of the saturation indices for minerals for different groundwater aquifers using the PHREEQC software	91
4.10	Comparison of the hydrochemical parameters in different aquifers with drinking water standards from the WHO (2011) and the MOH (2004)	98
4.11	Groundwater quality classification for irrigation	107
4.12	The similarities, differences and new findings of the study	108



## LIST OF FIGURES

Figure		Page
2.1	Geological map of the study area	10
2.2	The total groundwater production for potable use from 1990-2012	19
3.1	Map showing the location of the study area	32
3.2	Major river basins in the study area	33
3.3	The land use map of the study area	36
3.4	Yearly rainfall recorded from 1989-2016 at the study area	37
3.5	Cross section line of exploration wells	39
3.6	The location of groundwater sampling wells	41
3.7	The scheme adopted for research on the hydrogeochemistry of the Lower Kelantan Basin	44
4.1	Cross section line of exploration wells	49
4.2	Geological cross section A-A'	50
4.3	Geological cross section B-B'	50
4.4	Geological cross section C-C'	50
4.5	Geological cross section D-D'	51
4.6	Geological cross section E-E'	51
4.7	Geological cross section F-F'	51
4.8	Geological fence diagram of the Lower Kelantan Basin	52
4.9	Geological conceptual model of the Lower Kelantan Basin	53
4.10	The discrimination of the aquifers in the Lower Kelantan Basin (DA-deep aquifer, IA-intermediate aquifer and SA-shallow aquifer)	59
4.11	Variation of pH values with depth for aquifers in the Lower Kelantan Basin showing a decreasing trend	65

4.12	pH values in groundwater samples taken from (a) shallow, (b) intermediate and (c) deep aquifers	66
4.13	Spatial distribution of EC for the groundwater samples taken from (a) shallow, (b) intermediate and (c) deep aquifers	67
4.14	TDS concentration in groundwater samples taken from (a) shallow, (b) intermediate and (c) deep aquifers	68
4.15	Ca <sup>2+</sup> concentration in groundwater samples from (a) shallow, (b) intermediate and (c) deep aquifers	70
4.16	Concentration of Mg <sup>2+</sup> in groundwater samples from (a) shallow, (b) intermediate and (c) deep aquifers	71
4.17	Na <sup>+</sup> concentration in groundwater samples from (a) shallow, (b) intermediate and (c) deep aquifers	73
4.18	Concentration of K <sup>+</sup> in groundwater samples from (a) shallow, (b) intermediate and (c) deep aquifers	74
4.19	Cl <sup>-</sup> concentration in groundwater samples from (a) shallow, (b) intermediate and (c) deep aquifer	75
4.20	CO <sub>3</sub> <sup>2-</sup> concentration in groundwater samples taken from (a) shallow, (b) intermediate and (c) deep aquifers	76
4.21	HCO <sub>3</sub> <sup>-</sup> concentration in groundwater sample from (a) shallow, (b) intermediate and (c) deep aquifers	77
4.22	SO <sub>4</sub> <sup>2-</sup> concentration in (a) shallow, (b) intermediate and (c) deep aquifers	79
4.23	The distribution of NO <sub>3</sub> <sup>-</sup> concentration in (a) shallow, (b) intermediate and (c) deep groundwater aquifers	80
4.24	Fe <sup>2+</sup> concentration distribution in (a) shallow, (b) intermediate and (c) deep aquifers	81
4.25	Mn <sup>2+</sup> distribution in (a) shallow, (b) intermediate and (c) deep aquifers	82
4.26	Box and whisker plots for groundwater samples collected from different aquifers in the Lower Kelantan Basin (DA-deep aquifer, IA-intermediate aquifer and SA-shallow aquifer)	84
4.27	Piper trilinear diagram of groundwater samples collected from shallow (SA), intermediate (IA) and deep (DA) aquifers	87

4.28	Spatial distribution of hydrochemical facies in (a) shallow, (b) intermediate and (c) deep aquifers	88
4.29	The hydrogeological profile in relation to EC and water types of the study area (cross section line E-E' as indicated in Figure 4.5)	89
4.30	Saturation indices (anhydrite, aragonite, calcite, dolomite and halite) of groundwater samples collected from the shallow, the intermediate and the deep aquifers	92
4.31	Saturation indices (goethite, $\text{Fe}(\text{OH})_3$ , hematite and magnetite) of groundwater samples collected from the shallow, intermediate and deep aquifers	93
4.32	Gibbs diagrams showing the mechanism governing the hydrogeochemistry of groundwater (a) major cations versus TDS and (b) major anions versus TDS	95
4.33	Groundwater classification for irrigation based on the SAR and the Salinity Hazard	100
4.34	Wilcox diagram for the groundwater samples from the shallow (SA), the intermediate (IA) and the deep (DA) aquifers	102
4.35	Doneen classification for irrigation water based on the permeability index (PI)	104

## LIST OF ABBREVIATIONS

a.s.l.	Above sea level
AKSB	Air Kelantan Sdn Bhd
ANOVA	Analysis of Variance
BGR	Bundesanstalt Für Geowissenschaften Und Rohstoffe
DA	Discriminant Analysis
DEM	Digital Elevation Data
DID	Department of Irrigation and Drainage of Malaysia
EIA	Environmental Impact Assessment
GDP	Gross Domestic Product
GIS	Geographic Information System
GSD	Geological Survey Department
HCW	Horizontal Collector Well
IADA	Integrated Agricultural Development Area
KADA	Kemubu Agricultural Development Authority
Kbg	Kubang
Kg	Kampung
Km	Kilometers
KR	Kelly's Ratio
LOD	Limit of Detection
m	Meter
MAR	Magnesium Adsorption Ratio
MGD	Minerals and Geoscience Department
MLD	Million Litres per Day
MLR	Multiple Linear Regression
MMD	Malaysia Meteorological Department
MOH	Ministry of Health Malaysia
NAHRIM	National Hydraulic Research Institute of Malaysia
NEPU	National Economic Planning Unit
NWRS	National Water Resources Study
PCA	Principal Component Analysis
PI	Permeability Index
PWD	Public Works Department
RSC	Residual Sodium Carbonate
SAR	Sodium Adsorption Ratio
SI	Saturation Index
Tg	Tanjung
UNEP	United Nations Environment Program
UNESCO	United Nations Educational, Scientific and Cultural Organization
USEPA	United States Environmental Protection Agency
USGS	United States Geological Survey
USSL	United States Salinity Laboratory
VF	Varimax Factors
WHO	World Health Organisation

# CHAPTER 1

## INTRODUCTION

### 1.1 Background of the Study

Groundwater is a vital source of fresh water in many countries such as India, China, USA, Pakistan, Iran, Bangladesh, Mexico and Saudi Arabia (Van der Gun, 2012). Nearly one-third of the global population relies on it to meet domestic, industrial and agricultural demands (UNEP, 2013). Over the past few decades, the great competition for economic expansion resulted in rapid population growth, urbanization, industrialization and intensive agricultural activities. These changes orchestrated significant demands on land use and food production, increasing the demand for groundwater as a freshwater resource (Singh et al., 2015). Such demand inevitably leads to the exploitation of invaluable resources in ways that deteriorate water quality.

The quality of groundwater is controlled by reactions associated with natural processes and anthropogenic influences (Zuane, 1990). The negative effects of poor water quality on human health and the environment have increasingly been prominent in the eyes of the public. The preservation of groundwater quality is important because around 80% of water-borne diseases in developing countries are related to poor water quality (Olajire and Imeokparia, 2001). An understanding of groundwater chemistry, the hydrogeological evolution and the factors controlling the groundwater quality is crucial to determine its suitability for domestic, industrial and agricultural uses (Mokhtar et al., 2009; Department of Water, 2010; Alaya et al., 2014; Fijani et al., 2017).

The state of Kelantan ranks first in the consumption of groundwater in Malaysia (Manap et al., 2013). The total groundwater abstraction for public water supply in 2012 was 160.08 million liters per day (MLD), amounting to approximately 41% of the total abstraction for public water supply in Kelantan (AKSB, 2012). About 80% of the inhabitants in the region depend on groundwater, from either dug or tube wells. The Lower Kelantan Basin is a densely populated region, with high development activities due to its fertile soils, a low relief, a shallow water table and the widespread occurrence of aquifers with sufficient and accessible water. Groundwater quality is generally better than that of surface water, making it suitable to meet the increasing global water demand (Lan et al., 2015). It also requires minimal treatment, the development for water supply is low and it is scarcely affected by prolonged drought associated with climate change.

## 1.2 Problem Statement

Population growth, urban expansion and agricultural activities stress the water resources and pose threats to groundwater quality. In recent decades, the competition for water increased significantly due to the needs of the rising population. Since water supply is vital in any region, an understanding of groundwater resources is essential for the planning of future developments (Kudoda and Abdalla, 2015).

It is estimated that one billion people rely on groundwater for daily needs in Asia (Sivasankar and Gomathi, 2009). In Malaysia, groundwater utilization is expected to increase by 63% between 2000 and 2050 (Manap et al., 2013). The average public water supply coverage in Malaysia for 2014 was 94.7%, with 96.9% and 90.7% reported for urban and rural areas, respectively. The public water supply coverage by state is Johor (99.8%), Kedah (98.2%), Labuan (100%), Malacca (100%), Negeri Sembilan (99.9%), Penang (99.9%), Pahang (98.0%), Perak (99.6%), Perlis (99.5%), Sabah (82.0%), Sarawak (93.5%), Selangor (99.8%) and Terengganu (96.0%). However, 59.9% of the population in Kelantan depends on treated water from the public water supply (ASM, 2015).

The dependence of the inhabitants on groundwater is significant due to the unsatisfactory quality of the public water supply. Low water pressure, poor water quality such as odor, and frequent disruption of supply are some of the problems. These problems are attributed to the lack of funds to upgrade the water supply infrastructure and augment the water supply coverage (Association of Water and Energy Research of Malaysia, 2011).

Most inhabitants in the Lower Kelantan Basin obtain water from an unconfined shallow aquifer with a depth of less than 6 m, because it is cost effective and provides adequate supply throughout the year. Several studies indicate that the intermediate aquifer with depths of 20-40 m is brackish to saline and not suitable for drinking and agriculture (Abdul Rashid, 1989; Ang and Ismail, 1995; Azmi, 2004).

Since more than 95% of groundwater wellfields are in the study area, large-scale abstraction to meet high demand can cause lowering of the regional water level and induce deeper and spread-out cones of depression. Lowering of regional water level poses a further threat to the groundwater quality because it favors seawater intrusion (Aris et al., 2010). The understanding of a groundwater system is complicated by aquifer heterogeneity, groundwater flow and chemical evolution (Han et al., 2009).

Groundwater quality is a preoccupation (WHO, 2011; Vasanthavigar et al., 2012a; Bouderbala, 2015; Nag and Das, 2017) because poor water quality severely affects human health and crop growth. Accurate groundwater quality data is indispensable since the study area is exposed to contamination from the surface activities like sewerage, waste disposal, agricultural and industrial. Previous studies in the area

(Sofner, 1989; Suratman, 1997) were localized and restricted to parts of the urban and the wellfield areas, and focused on elevated concentrations of iron ( $\text{Fe}^{2+}$ ), electrical conductivity (EC), chloride ( $\text{Cl}^-$ ), manganese ( $\text{Mn}^{2+}$ ) and nitrate ( $\text{NO}_3^-$ ).

The Lower Kelantan Basin is located downstream of the Kelantan River and the river was considered as sufficient for freshwater because it is part of the largest river basin originating upstream of the mountainous region of the Titiwangsa Range. The encroachment of seawater into the Kelantan River which is observed up to 13 km from the estuary, subsequently restricted the utilization of the surface water resource (Chong, 1974).

The Department of Irrigation (DID, 2000) estimated that the total groundwater discharge from shallow and deep aquifers was approximately 488 MLD with allowable abstraction of 148 MLD. Since the total groundwater abstraction in 2012 of over 160 MLD (AKSB, 2012) exceeded the recharge capacity, monitoring on groundwater pumping activities is needed to prevent saltwater intrusion into freshwater aquifers. The study area is vulnerable to saltwater intrusion because it is a flat and low-lying coastal plain with limited drainage infrastructure (Tran et al., 2012). Despite the existence of several studies on groundwater in the Lower Kelantan Basin, few exist on aquifer continuity, the extent of remnant saline water hydrogeochemistry, factors controlling the groundwater quality, and groundwater suitability.

### **1.3 Objectives of the Study**

The objectives of this study were:

- i. To construct a conceptual model of the groundwater aquifer system.
- ii. To identify the factors contributing to the variations in groundwater chemistry.
- iii. To assess the hydrogeochemical characteristics in different aquifers (shallow, intermediate and deep aquifers).
- iv. To evaluate the groundwater suitability and potential usage of different aquifers for drinking and agricultural purposes.

### **1.4 Scope of the Study**

This study covers the following:

- i. Generating data (primary data) from groundwater sampling of existing tube wells.
- ii. Five new tube wells were constructed to obtain soils for description, creation of geological cross sections, and construction of a conceptual hydrogeological model. The constructed tube wells were also used for *in-situ* measurements as well as groundwater sampling.

- iii. Groundwater sampling was conducted primarily to evaluate the physico-chemical parameters of the water that were not covered in earlier studies and to complement secondary data.
- iv. This study involved acquiring and reviewing available secondary data including groundwater quality, geophysical, lithologic, land use, digital elevation model (DEM), rainfall and hydrology data.
- v. The study integrated the primary and secondary data, performed further treatment and analyses using a multivariate statistical approach to examine the relationship between the variables and the factors controlling the quality of groundwater in each aquifer.

### **1.5 Significance of the Study**

Water supply coverage represents the proportion of residents with access to a safe drinking water supply. The water operator capability should be extended to service all residents. For instance, the water supply coverage in Kelantan was 59.9% in 2014, whereas other states had already attained more than 90% coverage (ASM, 2015). This distinct difference was due to either the enhanced availability of groundwater or the direct abstraction of water from rivers, which made a portion of the population less dependent on public water supply services. However, an insufficient water supply, low water pressure, frequent water disruptions, and poor water quality could also contribute to low water supply coverage.

Since groundwater is important for drinking, industrial and agricultural purposes, the assessment of its suitability for different purposes is vital. This study will help in characterizing the groundwater quality and evolution of the aquifer systems, as well as develop conceptual hydrogeological models. The findings will establish a water quality baseline data and elucidate the suitability for drinking and agricultural purposes of different aquifers.

This study will also assess the factors controlling the groundwater quality in different aquifers. It will enable the delineation of the extent of saltwater remnants and groundwater contamination sources (anthropogenic and natural). Furthermore, this study will facilitate the estimating of the scale or corrosive potential of groundwater sources. This study aims to estimate a suitable drilling depth to aid assessments of groundwater potential for the quantitative valuation of water resources, therefore reducing the financial burden of soil investigations.

This study can also easily evaluate the spatial distribution of groundwater of various qualities in different aquifers for domestic, industrial, and agricultural use in the study area. This information helps stakeholders to take appropriate actions because of the problems corrosive water causes when it reacts with pipes and plumbing fixtures.



## 1.6 Research Framework of the Study

The research framework developed for the hydrogeochemistry study of the Lower Kelantan Basin is depicted in Figure 1.1.





**Figure 1.1: Research framework for the hydrogeochemistry study of the Lower Kelantan Basin.**

## 1.7 Thesis Outline

The body of this thesis consists of five chapters with information on the hydrogeochemistry, the suitability of the groundwater for drinking and agricultural purposes, and the factors affecting the quality of the groundwater. The chapters, from the introduction to the conclusion are as follows:

- i. Chapter 1 is an introduction with a study background, objectives, a problem statement and significance of the study.
- ii. Chapter 2 provides a comprehensive review of literature related to pollution of groundwater, the sources and effects on human health and plant growth based on comparisons with standards or guidelines, and the application of graphical representations and multivariate statistical analysis on the groundwater quality data.
- iii. Chapter 3 comprises the materials and methods including groundwater sampling, *in-situ* and laboratory analysis, and data interpretation.
- iv. Chapter 4 elaborates on the factors controlling variations in groundwater quality using multivariate statistical analysis, hydrogeochemistry, groundwater evaluation and its suitability for drinking and agricultural uses.
- v. Chapter 5 summarizes and concludes on the findings and recommendations made on appropriate measures for managing groundwater resources in a sustainable manner.

## REFERENCES

- Abdul Rashid, B. (1989). Groundwater monitoring system in the eastern Sungai Kelantan Delta, Kelantan. Geological Survey of Malaysia, Report GPH 05/1989, 32p.
- Abdul Rashid, B. (1994). Distribution of the nitrate levels in shallow groundwater in Kelantan River Basin: An Overview. In *Proceedings of 25<sup>th</sup> Annual Geological Conference 1994*. Geological Survey of Malaysia, pA07/1-A07/22.
- Abdullah, M. H., Mokhtar, M., Musta, B., & Aris, A. Z. (2008). Hydrochemical analyses of a disturbed aquifer of a small island in Malaysia. *Fresenius Environmental Bulletin*, 17(12 A), 2043-2051.
- Adhikary, P. P., Chandrasekharan, H., Chakraborty, D., & Kamble, K. (2010). Assessment of groundwater pollution in West Delhi, India using geostatistical approach. *Environmental Monitoring and Assessment*, 167(1-4), 599-615.
- Afzal Hossain, A. F. M. (1989). Conjunctive use of surface and groundwater in North Kelantan. Malaysia. Ph.D. Thesis, Universiti Kebangsaan Malaysia, Bangi, Malaysia, 487p.
- Aghazadeh, N., & Mogaddam, A. (2011). Investigation of hydrochemical characteristics of groundwater in the Harzandat aquifer, Northwest of Iran. *Environmental Monitoring and Assessment*, 176(1-4), 183-195.
- Aghazadeh, N., & Mogaddam, A. A. (2010). Assessment of groundwater quality and its suitability for drinking and agricultural uses in the Oshnavieh area, Northwest of Iran. *Journal of Environmental Protection*, 1(01), 30-40.
- Ahamed, A. J., Loganathan, K., & Jayakumar, R. (2015). Hydrochemical characteristics and quality assessment of groundwater in Amaravathi river basin of Karur district, Tamil Nadu, South India. *Sustainable Water Resources Management*, 1(3), 273-291.
- Ahn, H. I., & Chon, H. T. (1999). Assessment of groundwater contamination using geographic information systems. *Environmental Geochemistry and Health*, 21(3), 273-289.
- Air Kelantan Sendirian Berhad (AKSB) (2012). *The total groundwater production for potable use from 1990-2012*. Air Kelantan Sdn Bhd, Kuala Lumpur, Malaysia.
- Al-Amry, A. S. (2008). Hydrogeochemistry and groundwater quality assessment in an arid region: a case study from Al Salameh Area, Shabwah, Yemen. In *The 3rd International*
- Alaya, M. B., Saidi, S., Zemni, T., & Zargouni, F. (2014). Suitability assessment of deep groundwater for drinking and irrigation use in the Djeffara aquifers (Northern Gabes, south-eastern Tunisia). *Environmental Earth Sciences*, 71(8), 3387-3421.
- Alexakis, D. (2011). Assessment of water quality in the Messolonghi-Etoliko and Neochorio region (West Greece) using hydrochemical and statistical analysis methods. *Environmental Monitoring and Assessment*, 182(1), 397-413.
- Alkarkhi, A. F., Ahmad, A., Ismail, N., Mat Easa, A., & Omar, K. (2009). Assessment of surface water through multivariate analysis. *Journal of Sustainable Development*, 1(3), 27.
- Almasri, M. N., & Kaluarachchi, J. J. (2007). Modeling nitrate contamination of groundwater in agricultural watersheds. *Journal of Hydrology*, 343(3), 211-229.

- Al-Omran, A. M., Al-Harbi, A. R., Wahb-Allah, M. A., Nadeem, M., & Al-Eter, A. (2010). Impact of irrigation water quality, irrigation systems, irrigation rates and soil amendments on tomato production in sandy calcareous soil. *Turkish Journal of Agriculture and Forestry*, 34(1), 59-73.
- Amer, A. M. (1995). Saltwater intrusion in coastal aquifers. *Water Resources Management*, 2, 521-529.
- American Public Health Association (APHA) (1995). *Standard methods for the examination of water and wastewater*. American Public Health Association, American Water Works association, Water Environment Federation, Washington.
- Amiri, V., Sohrabi, N., & Dadgar, M. A. (2015). Evaluation of groundwater chemistry and its suitability for drinking and agricultural uses in the Lenjanat plain, central Iran. *Environmental Earth Sciences*, 74(7), 6163-6176.
- Amores, M. J., Verones, F., Raptis, C., Juraske, R., Pfister, S., Stoessel, F., Antón, A., Castells, F., & Hellweg, S. (2013). Biodiversity impacts from salinity increase in a coastal wetland. *Environmental Science & Technology*, 47(12), 6384-6392.
- Anbazhagan, S., & Nair, A. M. (2005). Geographic information system and groundwater quality mapping in Panvel Basin, Maharashtra, India. *Environmental Geology*, 45(6), 753-761.
- Ang, N. K. & Ismail, C. M. (1996). *Groundwater monitoring report Peninsular Malaysia 1995 (Kelantan)*. Geological Survey of Malaysia, Report GPH 4/96.
- Ang, N. K. & Kwan, T. S. (1978). *Hydrogeological investigation-proposed DID Water Management Centre, Kg. Panji, Kota Bharu, Kelantan*. Geological Survey of Malaysia, Report GPH 1/1978, 7p.
- Angelopoulos, K., Spiliopoulos, I. C., Mandoulaki, A., Theodorakopoulou, A., & Kouvelas, A. (2009). Groundwater nitrate pollution in northern part of Achaia Prefecture. *Desalination*, 248(1-3), 852-858.
- Appelo, C. A. J., & Postma, D. (2004). *Geochemistry, groundwater and pollution*. CRC press.
- Argamasilla, M., Barberá, J. A., & Andreo, B. (2017). Factors controlling groundwater salinization and hydrogeochemical processes in coastal aquifers from southern Spain. *Science of The Total Environment*, 580, 50-68.
- Aris, A. Z., Abdullah, M. H., & Kim, K. W. (2007). Hydrogeochemistry of groundwater in Manukan Island, Sabah. *The Malaysian Journal of Analytical Sciences*, 11(2), 407-413.
- Aris, A. Z., Abdullah, M. H., Kim, K. W., & Praveena, S. M. (2009). Hydrochemical changes in a small tropical island's aquifer: Manukan Island, Sabah, Malaysia. *Environmental Geology*, 56(8), 1721-1732.
- Aris, A. Z., Abdullah, M. H., Praveena, S. M., Yusoff, M. K., & Juahir, H. (2010). Extenuation of saline solutes in shallow aquifer of a small tropical island: a case study of Manukan Island, North Borneo. *Environment Asia*, 3(Special issue), 84-92.
- Arnous, M. O., Aboulela, H. A., & Green, D. R. (2011). Geo-environmental hazards assessment of the north western Gulf of Suez, Egypt. *Journal of Coastal Conservation*, 15(1), 37-50.

- Arunprakash, M., Giridharan, L., Krishnamurthy, R. R., & Jayaprakash, M. (2013). Impact of urbanization in groundwater of south Chennai City, Tamil Nadu, India. *Environmental Earth Sciences*, 71(2), 947-957.
- Asabere, B., & Bempah, K. (2013). Groundwater quality assessment for drinking and irrigation purposes in Obuasi municipality of Ghana, a preliminary study. *Research Journal of Environmental and Earth Sciences*, 5(1), 6-17.
- Ashraf, M., & Saeed, M. (2006). Effect of improved cultural practices on crop yield and soil salinity under relatively saline groundwater applications. *Irrigation and Drainage Systems*, 20, 111-124.
- ASM. (2015). *Study on the Current Issues and Needs for Water Supply and Wastewater Management in Malaysia. Volume 2*. Academy of Sciences Malaysia.
- Association of Water and Energy Research Malaysia (AWER) (2011). *National Water Services Industry Restructuring-Kelantan State Case Study*. Retrieved from: [http://www.awer.org.my/publication/report/national\\_water\\_services\\_industry\\_restructuring\\_kelantan\\_case\\_study\\_eng.pdf](http://www.awer.org.my/publication/report/national_water_services_industry_restructuring_kelantan_case_study_eng.pdf)
- Ayazi, M. H., Pirasteh, S., Arvin, A. K. P., Pradhan, B., Nikouravan, B., & Mansor, S. (2010). Disasters and risk reduction in groundwater: Zagros Mountain Southwest Iran using geoinformatics techniques. *Disaster Advances* 3(1), 51-57.
- Ayers, R. S., & Westcot, D. W. (1985). Water quality for agriculture. FAO Irrigation and drainage paper 29 Rev. 1. *Food and Agricultural Organization. Rome*.
- Azmi, I. (2004). *Groundwater monitoring report of North Kelantan (1989-1999)*. Department of Mineral and Geosciences Malaysia, JMG.KEL 01/04 (HYDRO 01), 39p
- Babu, N. B., Gajendran, C., Hameed, A. S., & James, E. J. (2015). Appraisal of groundwater quality for drinking and irrigation purposes in Nambiyar river basin, Tamil Nadu, India. *Water Resources*, 42(4), 553-562.
- Back, W., & Custodio, E. (1985). Chemical aspects of Groundwater. *Hydrogeology, éd., Csallany, CS and Kanivetskv, RA, McGraw-Hill Company*.
- Back, W., Baedecker, M. J., & Wood, W. W. (1993). Scales in chemical hydrogeology: A historical perspective. *Regional Ground Water Quality (William Alley, editor): Van Nostrand Reinhold, New York, NY*, 111-128.
- Baghvand, A., Nasrabadi, T., Bidhendi, G. N., Vosoogh, A., Karbassi, A., & Mehrdadi, N. (2010). Groundwater quality degradation of an aquifer in Iran central desert. *Desalination*, 260(1), 264-275.
- Baharuddin, M. F. T., Taib, S., Hashim, R., Abidin, M. H. Z., & Rahman, N. I. (2013). Assessment of seawater intrusion to the agricultural sustainability at the coastal area of Carey Island, Selangor, Malaysia. *Arabian Journal of Geosciences*, 6(10), 3909-3928.
- Balasubramanian, N., Sivasubramanian, P., Soundranayagam, J. P., Chandrasekar, N., & Gowtham, B. (2015). Groundwater classification and its suitability in Kadaladi, Ramanathapuram, India using GIS techniques. *Environmental Earth Sciences*, 74(4), 3263-3285.
- Ballukraya, P. N., & Ravi, R. (1998). Natural fresh-water ridge as barrier against sea-water intrusion in Chennai city. *Geological Society of India*, 52(3), 279-286.

- Banat, K. M., Howari, F. M., & Al-Hamad, A. A. (2005). Heavy metals in urban soils of central Jordan: should we worry about their environmental risks? *Environmental Research*, 97(3), 258-273.
- Barker, A. P., Newton, R. J., Bottrell, S. H., & Tellam, J. H. (1998). Processes affecting groundwater chemistry in a zone of saline intrusion into an urban sandstone aquifer. *Applied Geochemistry*, 13(6), 735-749.
- Barros, R., Isidoro, D., & Aragüés, R. (2012). Irrigation management, nitrogen fertilization and nitrogen losses in the return flows of La Violada irrigation district (Spain). *Agriculture, Ecosystems & Environment*, 155, 161-171.
- Bashir, E., Naseem, S., Hanif, H., & Pirzada, T. (2013). Geochemical study of groundwater of Uthal and Bela areas, Balochistan and its appraisal for drinking and irrigation water quality. *International Journal of Agriculture and Environmental Research*, 2, 1-13.
- Batayneh, A., & Zumlot, T. (2012). Multivariate statistical approach to geochemical methods in water quality factor identification; application to the shallow aquifer system of the Yarmouk basin of north Jordan. *Research Journal of Environmental and Earth Sciences*, 4(7), 756-768.
- Bengraïne, K., & Marhaba, T. F. (2003). Using principal component analysis to monitor spatial and temporal changes in water quality. *Journal of Hazardous Materials*, 100(1), 179-195.
- Bennetts, D. A., Webb, J. A., Stone, D. J. M., & Hill, D. M. (2006). Understanding the salinisation process for groundwater in an area of south-eastern Australia, using hydrochemical and isotopic evidence. *Journal of Hydrology*, 323(1), 178-192.
- Berg, M., Trang, P. T. K., Stengel, C., Buschmann, J., Viet, P. H., Van Dan, N., Giger, W., & Stüben, D. (2008). Hydrological and sedimentary controls leading to arsenic contamination of groundwater in the Hanoi area, Vietnam: the impact of iron-arsenic ratios, peat, river bank deposits, and excessive groundwater abstraction. *Chemical Geology*, 249(1), 91-112.
- Bina Runding Sdn. Bhd. (1986). Preliminary proposal for groundwater monitoring at Tanjung Mas Wellfields, Kelantan. Bina Runding Sdn. Bhd. in association with John Taylor & Sons. (International), 9p.
- Birkinshaw, S. J., & Ewen, J. (2000). Nitrogen transformation component for Shetran catchment nitrate transport modelling. *Journal of Hydrology*, 230(1), 1-17.
- Blood, D. C., Radostits, O. M., Henderson, J. A., Arundel, J. A. & Gay, C. C. (1983). *Veterinary Medicine. Bailliere Tindall. London, UK.*
- Bosch, J. H. A. (1986). *Young Quaternary sediments in the coastal plain of Kelantan, Peninsular Malaysia.* Geological Survey of Malaysia, Quaternary Geology Section.
- Bouderbala, A. (2015). Assessment of groundwater quality and its suitability for agricultural uses in the Nador Plain, North of Algeria. *Water Quality, Exposure and Health*, 7(4), 445-457.
- Boyce, J. S., Muir, J., Edwards, A. P., Seim, E. C., & Olson, R. A. (1976). Geologic nitrogen in Pleistocene loess of Nebraska. *Journal of Environmental Quality*, 5(1), 93-96.
- Boyd, C. E. (2015). *Water Quality: An Introduction.* Springer.

- Bozdağ, A., & Göçmez, G. (2013). Evaluation of groundwater quality in the Cihanbeyli basin, Konya, Central Anatolia, Turkey. *Environmental Earth Sciences*, 69(3), 921-937.
- Bradley, S. L., Milne, G. A., Horton, B. P., & Zong, Y. (2016). Modelling sea level data from China and Malay-Thailand to estimate Holocene ice-volume equivalent sea level change. *Quaternary Science Reviews*, 137, 54-68.
- Brindha, K., Vaman, K. N., Srinivasan, K., Babu, M. S., & Elango, L. (2014). Identification of surface water-groundwater interaction by hydrogeochemical indicators and assessing its suitability for drinking and irrigational purposes in Chennai, Southern India. *Applied Water Science*, 4(2), 159-174.
- Burow, K. R., Jurgens, B. C., Belitz, K., & Dubrovsky, N. M. (2013). Assessment of regional change in nitrate concentrations in groundwater in the Central Valley, California, USA, 1950s–2000s. *Environmental Earth Sciences*, 69(8), 2609-2621.
- Canter, L. W. (1996). *Nitrates in groundwater*. CRC press.
- Chandrasekar, N., Selvakumar, S., Srinivas, Y., Wilson, J. J., Peter, T. S., & Magesh, N. S. (2014). Hydrogeochemical assessment of groundwater quality along the coastal aquifers of southern Tamil Nadu, India. *Environmental Earth Sciences*, 71(11), 4739-4750.
- Chapelle, F. H. (2001). *Ground-water Microbiology and Geochemistry*. Wiley, New York.
- Chen, J., Wu, H., & Qian, H., (2016). Groundwater Nitrate Contamination and Associated Health Risk for the Rural Communities in an Agricultural Area of Ningxia, Northwest China. *Exposure and Health*, 1-11.
- Chen, K., Jiao, J. J., Huang, J., & Huang, R. (2007). Multivariate statistical evaluation of trace elements in groundwater in a coastal area in Shenzhen, China. *Environmental Pollution*, 147(3), 771-780.
- Cheong, J. Y., Hamm, S. Y., Lee, J. H., Lee, K. S., & Woo, N. C. (2012). Groundwater nitrate contamination and risk assessment in an agricultural area, South Korea. *Environmental Earth Sciences*, 66(4), 1127-1136.
- Chidambaram, S., Karmegam, U., Prasanna, M. V., Sasidhar, P., & Vasanthavigar, M. (2011). A study on hydrochemical elucidation of coastal groundwater in and around Kalpakkam region, Southern India. *Environmental Earth Sciences*, 64(5), 1419-1431.
- Chidambaram, S., Kumar, G. S., Prasanna, M. V., Peter, A. J., Ramanathan, A. L., & Srinivasamoorthy, K. (2009). A study on the hydrogeology and hydrogeochemistry of groundwater from different depths in a coastal aquifer: Annamalai Nagar, Tamilnadu, India. *Environmental Geology*, 57(1), 59-73.
- Chong, F. S., Pfeiffer, D. & Tiedemann, O. (1976). *Hydrogeological report of Tumpat, Kelantan*. Geological Survey of Malaysia, Report GPH 2/1976, 8p.
- Chong, F. S., Pfeiffer, D. and Tiedemann, O. (1977). *Hydrogeological report of Kota Bharu, Kelantan City Waterworks*. Geological Survey of Malaysia, Report GPH 11/1977, 19p.
- Chong, F. S., Pfeiffer, D., Sidhu, H. S. & Tiedemann, O. (1974). *Groundwater exploration in Kota Bharu, Report No. II*. Geological Survey of Malaysia, Report GPH 2/1974, 7p.



- Cloutier, V., Lefebvre, R., Therrien, R., & Savard, M. M. (2008). Multivariate statistical analysis of geochemical data as indicative of the hydrogeochemical evolution of groundwater in a sedimentary rock aquifer system. *Journal of Hydrology*, 353(3), 294-313.
- Cruz, J. V., & Silva, M. O. (2000). Groundwater salinization in Pico Island (Azores, Portugal): origin and mechanisms. *Environmental Geology*, 39(10), 1181-1189.
- Dalton, M. G., & Upchurch, S. B. (1978). Interpretation of hydrochemical facies by factor analysis. *Ground Water*, 16(4), 228-233.
- Davraz, A., & Özdemir, A. (2014). Groundwater quality assessment and its suitability in Çeltikçi plain (Burdur/Turkey). *Environmental Earth Sciences*, 72(4), 1167-1190.
- Department of Environment of Malaysia (DOE). (1994). *Groundwater quality monitoring network in Malaysia*. Final Report, Vol. I-II.
- Department of Environment of Malaysia (DOE). (2009). *2009 Annual Report*. Department of Environment, Ministry of Natural Resources and Environment.
- Department of Statistics Malaysia (2010). *Population Distribution and Basic Demographic Characteristic Report 2010*. Department of Statistics, Kuala Lumpur.
- Department of Water (2010). *Hydrogeochemical assessment of the Superficial aquifer, Perth metropolitan area*. Hydrogeological record series report no. HG37. St. Georges Terrace, Perth Western Australia, 174p.
- Deutsch, W. J., & Siegel, R. (1997). *Groundwater geochemistry: fundamentals and applications to contamination*. CRC press.
- DID. (2000). *National Water Resources Study (2000-2050) of Peninsular Malaysia, Volume 18 (Kelantan)*. Ministry of Natural Resources and Environment Malaysia, Kuala Lumpur. p300.
- DID. (2011). *Review of the national water resources study (2000-2050) and formulation of national water resources policy, Volume 10*. Ministry of Natural Resources and Environment Malaysia, Kuala Lumpur. p237.
- Dillon, P. J., Close, M. E., & Scott, R. I. (1989). Diffuse-Source Nitrate Contamination of Groundwater in New Zealand and Australia. In *Hydrology and Water Resources Symposium 1989: Comparisons in Austral Hydrology; Preprints of Papers* (p. 351). Institution of Engineers, Australia.
- Doneen, L. D. (1975). Water quality for irrigated agriculture. In *Plants in saline environments* (pp. 56-76). Springer Berlin Heidelberg.
- Drever, J. I. (1988). *The geochemistry of natural waters* (Vol. 437). Englewood Cliffs: Prentice Hall.
- Dun, Y., Tang, C., & Shen, Y. (2014). Identifying interactions between river water and groundwater in the North China Plain using multiple tracers. *Environmental earth sciences*, 72(1), 99-110.
- Durov, S. A. (1948). Natural waters and graphic representation of their composition. In *Dokl Akad Nauk SSSR* (Vol. 59, No. 3, pp. 87-90).
- Dwivedi, A. K., & Vankar, P. S. (2014). Source identification study of heavy metal contamination in the industrial hub of Unnao, India. *Environmental Monitoring and Assessment*, 186(6), 3531-3539.
- Earth Summit (1992). Programme of action for sustainable development (174-175). *Agenda 21*, UNCED, Rio de Janeiro, Brazil

- Eaton, F. M. (1950). Significance of carbonates in irrigation waters. *Soil Science*, 69(2), 123-134.
- Edmunds, W. M., Guendouz, A. H., Mamou, A., Moulla, A., Shand, P., & Zouari, K. (2003). Groundwater evolution in the Continental Intercalaire aquifer of southern Algeria and Tunisia: Trace element and isotopic indicators. *Applied Geochemistry*, 18, 805-822.
- Edmunds, W. M., Ma, J., Aeschbach-Hertig, W., Kipfer, R., & Darbyshire, D. P. F. (2006). Groundwater recharge history and hydrogeochemical evolution in the Minqin Basin, North West China. *Applied Geochemistry*, 21(12), 2148-2170.
- El Kashouty, M., & Abdel-Lattif, A. (2011). Multivariate statistical analyses of the groundwater system in Darb El-Arbaein, Southwestern Desert, Egypt. *Arabian Journal of Geosciences*, 4(7-8), 1117-1130.
- ENEX. (1977). *The Kelantan river basin study*. The Government of Malaysia and the New Zealand Ministry of Foreign Affairs, Main Report, Hydrology, Vol. 1.
- Esmaeili, A., & Moore, F. (2012). Hydrogeochemical assessment of groundwater in Isfahan province, Iran. *Environmental Earth Sciences*, 67(1), 107-120.
- Fabro, A. Y. R., Ávila, J. G. P., Alberich, M. V. E., Sansores, S. A. C., & Camargo-Valero, M. A. (2015). Spatial distribution of nitrate health risk associated with groundwater use as drinking water in Merida, Mexico. *Applied Geography*, 65, 49-57.
- Faniran, J. A., Ngceba, F. S., Bhat, R. B., & Oche, C. Y. (2001). An assessment of the water quality of the Isinuka springs in the Transkei region of the Eastern Cape, Republic of South Africa. *Water SA*, 27(2), 241-250.
- Farnham, I. M., Johannesson, K. H., Singh, A. K., Hodge, V. F., & Stetzenbach, K. J. (2003). Factor analytical approaches for evaluating groundwater trace element chemistry data. *Analytica Chimica Acta*, 490(1), 123-138.
- Faure, G. (1998). *Principles and applications of geochemistry*, 2<sup>nd</sup> Edn. Englewood Cliffs: Prentice Hall.
- Fehdi, C., Rouabhia, A., Baali, F., & Boudoukha, A. (2009). The hydrogeochemical characterization of Morsott-El Aouinet aquifer, northeastern Algeria. *Environmental Geology*, 58(7), 1611-1620.
- Fijani, E., Moghaddam, A. A., Tsai, F. T. C., & Tayfur, G. (2017). Analysis and assessment of hydrochemical characteristics of Maragheh-Bonab plain aquifer, northwest of Iran. *Water Resources Management*, 31(3), 765-780.
- Fipps, G. (2003). Irrigation water quality standards and salinity management strategies. *Texas FARMER Collection*
- Flathe, H. (1974). *Geoelectrical investigation, Kota Bharu/Kelantan, Malaysia*. Bundesanstalt Für Geowissenschaften Und Rohstoffe (BGR), Germany. 9p.
- Forman, D. (1989). Are nitrates a significant risk factor in human cancer? *Cancer Surveys*, 8(2), 443-458.
- Foster, S., Chilton, J., Moench, M., Cardy, F., & Schiffler, M. (2015). *Groundwater in rural development: facing the challenges of supply and resource sustainability*. Water P-notes, no. 19. World Bank, Washington, DC.
- Freeze, R. A. & Cherry, J. A. (1979). *Groundwater*, Prentice-Hall, Englewood Cliffs, New Jersey, 376p.

- Gárfias, J., Arroyo, N., & Aravena, R. (2010). Hydrochemistry and origins of mineralized waters in the Puebla aquifer system, Mexico. *Environmental Earth Sciences*, 59(8), 1789-1805.
- Garg, V. K., Suthar, S., Singh, S., Sheoran, A., & Jain, S. (2009). Drinking water quality in villages of southwestern Haryana, India: assessing human health risks associated with hydrochemistry. *Environmental Geology*, 58(6), 1329-1340.
- Ghabayen, S. M., McKee, M., & Kemblowski, M. (2006). Ionic and isotopic ratios for identification of salinity sources and missing data in the Gaza aquifer. *Journal of Hydrology*, 318(1), 360-373.
- Ghodeif, K. O., Arnous, M. O., & Geriesh, M. H. (2013). Define a protected buffer zone for Ismailia Canal, Egypt using geographic information systems. *Arabian Journal of Geosciences*, 6(1), 43-53
- Ghosh, N., Virk, P., & Rishi, M. S. & Kamaldeep (2010). Study of groundwater quality in parts of district Patiala, Punjab, India with special reference to nitrate pollution. *International Journal of Environmental Science*, 1(2), 175-182.
- Gibbs, R. J. (1970). Mechanisms controlling world water chemistry. *Science*, 170(3962), 1088-1090.
- Gimenez, E., & Morell, I. (1997). Hydrogeochemical analysis of salinization processes in the coastal aquifer of Oropesa (Castellon, Spain). *Environmental Geology*, 29(1-2), 118-131.
- Giridharan, L., Venugopal, T., & Jayaprakash, M. (2008). Evaluation of the seasonal variation on the geochemical parameters and quality assessment of the groundwater in the proximity of River Cooum, Chennai, India. *Environmental Monitoring and Assessment*, 143(1), 161-178.
- Giridharan, L., Venugopal, T., & Jayaprakash, M. (2009). Identification and evaluation of hydrogeochemical processes on river Cooum, South India. *Environmental Monitoring and Assessment*, 162, 277-289.
- Gnanachandrasamy, G., Ramkumar, T., Venkatramanan, S., Anithamary, I., & Vasudevan, S. (2012). GIS based hydrogeochemical characteristics of groundwater quality in Nagapattinam district, Tamil Nadu, India. *Carpathian Journal of Earth and Environmental Sciences*, 7(3), 205-210.
- Gnanachandrasamy, G., Ramkumar, T., Venkatramanan, S., Vasudevan, S., Chung, S. Y., & Bagyaraj, M. (2015). Accessing groundwater quality in lower part of Nagapattinam district, Southern India: using hydrogeochemistry and GIS interpolation techniques. *Applied Water Science*, 5(1), 39-55.
- Goyal, S. K., Chaudhary, B. S., Singh, O., Sethi, G. K., & Thakur, P. K. (2010). GIS based spatial distribution mapping and suitability evaluation of groundwater quality for domestic and agricultural purpose in Kaithal district, Haryana state, India. *Environmental Earth Sciences*, 61(8), 1587-1597.
- Guendouz, A., Moulla, A. S., Edmunds, W. M., Zouari, K., Shand, P., & Mamou, A. (2003). Hydrogeochemical and isotopic evolution of water in the Complexe Terminal aquifer in the Algerian Sahara. *Hydrogeology Journal*, 11(4), 483-495.
- Güler, C., & Thyne, G. D. (2004). Hydrologic and geologic factors controlling surface and groundwater chemistry in Indian Wells-Owens Valley area, southeastern California, USA. *Journal of Hydrology*, 285(1), 177-198.

- Güler, C., Thyne, G. D., McCray, J. E., & Turner, K. A. (2002). Evaluation of graphical and multivariate statistical methods for classification of water chemistry data. *Hydrogeology Journal*, 10(4), 455-474.
- Guo, H., & Wang, Y. (2004). Hydrogeochemical processes in shallow quaternary aquifers from the northern part of the Datong Basin, China. *Applied Geochemistry*, 19(1), 19-27.
- Gupta, S. K., & Gupta, I. C. (1987). Management of saline soils and waters.
- Gupta, S., Mahato, A., Roy, P., & Datta, J. K. (2008). Geochemistry of groundwater, Burdwan District, West Bengal, India. *Environmental Geology*, 53, 1271-1282.
- Halim, M. A., Majumder, R. K., Nessa, S. A., Hiroshiro, Y., Sasaki, K., Saha, B. B., Saepuloh, A., & Jinno, K. (2010a). Evaluation of processes controlling the geochemical constituents in deep groundwater in Bangladesh: spatial variability on arsenic and boron enrichment. *Journal of Hazardous Materials*, 180(1), 50-62.
- Halim, M. A., Majumder, R. K., Nessa, S. A., Oda, K., Hiroshiro, Y., & Jinno, K. (2010b). Arsenic in shallow aquifer in the eastern region of Bangladesh: insights from principal component analysis of groundwater compositions. *Environmental Monitoring and Assessment*, 161(1-4), 453-472.
- Hallberg, G. R. (1986). From hoes to herbicides-agriculture and groundwater quality. *Journal of Soil and Water Conservation*, 41, 357-364.
- Han, D., Liang, X., Jin, M., Currell, M. J., Han, Y., & Song, X. (2009). Hydrogeochemical indicators of groundwater flow systems in the Yangwu River alluvial fan, Xinzhou Basin, Shanxi, China. *Environmental Management*, 44(2), 243-255.
- Haryono, A. (1997). *Penggunaan kaedah geofizik dan geokimia untuk mengesan penerobosan air masin ke dalam akuifer pantai, di utara Kelantan*. PhD thesis, National University of Malaysia, Malaysia
- Helena, B., Pardo, R., Vega, M., Barrado, E., Fernandez, J. M., & Fernandez, L. (2000). Temporal evolution of groundwater composition in an alluvial aquifer (Pisuerga River, Spain) by principal component analysis. *Water research*, 34(3), 807-816.
- Hem, J. D. (1985). *Study and interpretation of the chemical characteristics of natural water* (Vol. 2254). Department of the Interior, US Geological Survey.
- Hem, J. D. (1989). *Study and interpretation of the chemical characteristics of natural water*. US Geological Survey, Water-Supply Paper 2254, United States Department of the Interior. United States Government Printing Office, Washington, D.C., 263p.
- Hem, J. D. (1991). *Study and interpretation of the chemical characteristics of natural water*. 2254, Scientific Publishers, Jodhpur, India, 263p.
- Herojeet, R., Rishi, M. S., Lata, R., & Sharma, R. (2016). Application of environmetrics statistical models and water quality index for groundwater quality characterization of alluvial aquifer of Nalagarh Valley, Himachal Pradesh, India. *Sustainable Water Resources Management*, 2(1), 39-53.
- Hiscock, K. M. (2011). Groundwater in the 21st century-meeting the challenges. In *Sustaining Groundwater Resources* (pp. 207-225). Springer Netherlands.
- Hornsby, A. G. (1990). Pollution and public health problems related to irrigation. In: *Steward, B.A. and Nielson, D.R. (eds.), Irrigation of Agricultural Crops. American Society of Agronomy, Madison, Wisconsin, USA. Agronomy Series, No. 30, p1176-1187.*

- Hosseinifard, S. J., & Aminiyan, M. M. (2015). Hydrochemical characterization of groundwater quality for drinking and agricultural purposes: a case study in Rafsanjan plain, Iran. *Water Quality, Exposure and Health*, 7(4), 531-544.
- Houatmia, F., Azouzi, R., Charef, A., & Bédir, M. (2016). Assessment of groundwater quality for irrigation and drinking purposes and identification of hydrogeochemical mechanisms evolution in Northeastern, Tunisia. *Environmental Earth Sciences*, 75(9), 746.
- Hounslow, A. (1995). *Water quality data: analysis and interpretation*. CRC press.
- Howard, J., & Merrifield, M. (2010). Mapping groundwater dependent ecosystems in California. *PLoS One*, 5(6), e11249.
- Howari, F. M., & Banat, K. M. (2002). Hydrochemical characteristics of Jordan and Yarmouk river waters: effect of natural and human activities. *Journal of Hydrology and Hydromechanics*, 50(1), 50-64.
- Hundal, H. (2011). Geochemistry and assessment of hydrogeochemical processes in groundwater in the southern part of Bathinda District of Punjab, northwest India. *Environmental Earth Sciences*, 64(7), 1823-1833.
- Hussin, N. H., Yusoff, I., Tahir, W. Z. W. M., Mohamed, I., Ibrahim, A. I. N., & Rambli, A. (2016). Multivariate statistical analysis for identifying water quality and hydrogeochemical evolution of shallow groundwater in Quaternary deposits in the Lower Kelantan River Basin, Malaysian Peninsula. *Environmental Earth Sciences*, 75(14), 1081.
- Isa, N. M., & Aris, A. Z. (2014). Application of Geochemical and Geostatistical Analyses in Observing the Controlling Factors of Groundwater Compositions. In *From Sources to Solution* (pp. 171-174). Springer, Singapore.
- Isa, N. M., Aris, A. Z., & Sulaiman, W. N. A. W. (2012). Extent and severity of groundwater contamination based on hydrochemistry mechanism of sandy tropical coastal aquifer. *Science of the Total Environment*, 438, 414-425.
- Ishaku, J. M. (2011). Assessment of groundwater quality index for Jimeta-Yola area, Northeastern Nigeria. *Journal of Geology and Mining Research*, 3(9), 219-231.
- Islam, M. A., Zahid, A., Rahman, M. M., Rahman, M. S., Islam, M. J., Akter, Y., Shammi, M., Bodrud-Doza, M., & Roy, B. (2017). Investigation of groundwater quality and its suitability for drinking and agricultural use in the south central part of the coastal region in Bangladesh. *Exposure and Health*, 9(1), 27-41.
- Jackson, J. E. (2005). *A user's guide to principal components* (Vol. 587). John Wiley & Sons.
- Jalali, M. (2009). Geochemistry characterization of groundwater in an agricultural area of Razan, Hamadan, Iran. *Environmental Geology*, 56(7), 1479-1488.
- Jamshidzadeh, Z., & Mirbagheri, S. A. (2011). Evaluation of groundwater quantity and quality in the Kashan Basin, Central Iran. *Desalination*, 270(1), 23-30.
- Jassas, H., & Merkel, B. (2015). Assessment of hydrochemical evolution of groundwater and its suitability for drinking and irrigation purposes in Al-Khazir Gomal Basin, Northern Iraq. *Environmental Earth Sciences*, 74(9), 6647-6663.
- Javadi, A. A., Abd-Elhamid, H. F., & Farmani, R. (2012). A simulation-optimization model to control seawater intrusion in coastal aquifers using abstraction/recharge wells. *International Journal for Numerical and Analytical Methods in Geomechanics*, 36(16), 1757-1779.

- Jeevanandam, M., Kannan, R., Srinivasalu, S., & Rammohan, V. (2007). Hydrogeochemistry and groundwater quality assessment of lower part of the Ponnaiyar River Basin, Cuddalore district, South India. *Environmental Monitoring and Assessment*, 132(1-3), 263.
- Jeevanandam, M., Nagarajan, R., Manikandan, M., Senthilkumar, M., Srinivasalu, S., & Prasanna, M. V. (2012). Hydrogeochemistry and microbial contamination of groundwater from Lower Ponnaiyar basin, Cuddalore district, Tamil Nadu, India. *Environmental Earth Sciences*, 67(3), 867-887.
- Jehangir, A., Tanveer, A., Yousuf, A. R., Masood, A., & Naqash, A. H. (2011). Geochemistry and irrigation quality of groundwater along river Jhelum in south Kashmir, India. *Recent Research in Science and Technology*, 3(6).
- Jeong, C. H. (2001). Effect of land use and urbanization on hydrochemistry and contamination of groundwater from Taejon area, Korea. *Journal of Hydrology*, 253(1), 194-210.
- Jiang, G., Liu, D., Song, K., Wang, Z., Zhang, B., & Wang, Y. (2010). Application of multivariate model based on three simulated sensors for water quality variables estimation in Shitoukoumen reservoir, Jilin province, China. *Chinese Geographical Science*, 20(4), 337-344.
- Jiang, Y., & Yan, J. (2010). Effects of land use on hydrochemistry and contamination of Karst groundwater from Nandong underground river system, China. *Water, Air, & Soil Pollution*, 210(1-4), 123-141.
- Johnson, C. C. (1979). Land application of waste-An accident waiting to happen. *Ground Water*, 17(1), 69-72.
- Johnson, R. A., & Wichern, D. W. (2002). *Applied multivariate statistical analysis* (Vol. 5, No. 8). Upper Saddle River, NJ: Prentice hall.
- Joshi, A., & Seth, G. (2011). Hydrochemical profile for assessing the groundwater quality of Sambhar Lake City and its Adjoining area. *Environmental Monitoring and Assessment*, 174(1), 547-554.
- Ju, X. T., Kou, C. L., Zhang, F. S., & Christie, P. (2006). Nitrogen balance and groundwater nitrate contamination: comparison among three intensive cropping systems on the North China Plain. *Environmental Pollution*, 143(1), 117-125.
- Juahir, H., Zain, S. M., Khan, R. A., Yusoff, M. K., Mokhtar, M. B., & Toriman, M. E. (2009). Using chemometrics in assessing Langat River water quality and designing a cost-effective water sampling strategy. *Maejo International Journal of Science and Technology*, 3(1), 26-42.
- Juahir, H., Zain, S. M., Yusoff, M. K., Hanidza, T. I. T., Armi, A. S. M., Toriman, M. E., & Mokhtar, M. (2011). Spatial water quality assessment of Langat River Basin (Malaysia) using environmetric techniques. *Environmental Monitoring and Assessment*, 173(1), 625-641.
- Kaiser, H. F. (1960). The application of electronic computers to factor analysis. *Educational and Psychological Measurement*, 20(1), 141-151.
- Kalpana, L., & Elango, L. (2013). Assessment of groundwater quality for drinking and irrigation purposes in Pambar river sub-basin, Tamil Nadu. *Indian Journal of Environmental Protection*, 33(1), 1-8.

- Kannel, P. R., Lee, S., Kanel, S. R., & Khan, S. P. (2007). Chemometric application in classification and assessment of monitoring locations of an urban river system. *Analytica Chimica Acta*, 582(2), 390-399.
- Karanth, K. R. (1987). *Ground water assessment: development and management*. Tata McGraw-Hill Education.
- Kelley, W. P. (1963). Use of Saline Irrigation Water. *Soil Science*, 95(6), 385-391.
- Kelly, W. P. (1940). Permissible composition and concentration of irrigated waters. *Proceedings of the ASCF*, 607.
- Khalifa, I. H., & Arnous, M. O. (2012). Assessment of hazardous mine waste transport in west central Sinai, using remote sensing and GIS approaches: a case study of Um Bogma area, Egypt. *Arabian Journal of Geosciences*, 5(3), 407-420.
- Khodapanah, L., Sulaiman, W. N. A., & Khodapanah, N. (2009). Groundwater quality assessment for different purposes in Eshtehard District, Tehran, Iran. *European Journal of Scientific Research*, 36(4), 543-553.
- Kim, J. O., & Mueller, C. W. (1987). Introduction to factor analysis: what it is and how to do it. Quantitative applications in the social sciences series.
- Kim, K., Kim, H. J., Choi, B. Y., Kim, S. H., Park, K. H., Park, E., Koh, D. C. & Yun, S. T. (2008). Fe and Mn levels regulated by agricultural activities in alluvial groundwaters underneath a flooded paddy field. *Applied Geochemistry*, 23(1), 44-57.
- Kim, K., Rajmohan, N., Kim, H. J., Hwang, G. S., & Cho, M. J. (2004). Assessment of groundwater chemistry in a coastal region (Kunsan, Korea) having complex contaminant sources: a stoichiometric approach. *Environmental Geology*, 46(6-7), 763-774.
- Kim, R. H., Kim, J. H., Ryu, J. S., & Chang, H. W. (2006). Salinization properties of a shallow groundwater in a coastal reclaimed area, Yeonggwang, Korea. *Environmental Geology*, 49(8), 1180-1194.
- Kim, Y., Lee, K. S., Koh, D. C., Lee, D. H., Lee, S. G., Park, W. B., Koh, G. W. & Woo, N. C. (2003). Hydrogeochemical and isotopic evidence of groundwater salinization in a coastal aquifer: a case study in Jeju volcanic island, Korea. *Journal of Hydrology*, 270(3), 282-294.
- Klammler, G., Kupfersberger, H., Rock, G., & Fank, J. (2013). Modeling coupled unsaturated and saturated nitrate distribution of the aquifer Westliches Leibnitzer Feld, Austria. *Environmental Earth Sciences*, 69(2), 663-678.
- Klimas, A., & Gregorauskas, M. (2002). Groundwater abstraction and contamination in Lithuania as geoindicators of environmental change. *Environmental Geology*, 42(7), 767-772.
- Knapp, M. F. (2005). Diffuse pollution threats to groundwater: a UK water company perspective. *Quarterly Journal of Engineering Geology and Hydrogeology*, 38(1), 39-51.
- Koopmans, B. N. (1972). Sedimentation in the Kelantan delta (Malaysia). *Sedimentary Geology*, 7(1), 65-84.
- Krishnamurthy, J., & Srinivas, G. (1995). Role of geological and geomorphological factors in ground water exploration: a study using IRS LISS data. *International Journal of Remote Sensing*, 16(14), 2595-2618.

- Krumbein, W. C., & Graybill, F. A. (1965). *An introduction to statistical models in geology* (Vol. 475). New York: McGraw-Hill.
- Kudoda, A. M., & Abdalla, O. A. (2015). Hydrochemical characterization of the main aquifers in Khartoum, the capital city of Sudan. *Environmental Earth Sciences*, 74(6), 4771-4786.
- Kumar, P. J. S., Elango, L., & James, E. J. (2014). Assessment of hydrochemistry and groundwater quality in the coastal area of South Chennai, India. *Arabian Journal of Geosciences*, 7(7), 2641-2653.
- Kumar, P. S., Jegathambal, P., & James, E. J. (2011). Multivariate and geostatistical analysis of groundwater quality in Palar river basin. *International Journal of Geology*, 4, 108-119.
- Kumar, P., Kumar, A., Singh, C. K., Saraswat, C., Avtar, R., Ramanathan, A. L., & Herath, S. (2016). Hydrogeochemical Evolution and Appraisal of Groundwater Quality in Panna District, Central India. *Exposure and Health*, 8(1), 19-30.
- Kumar, S. K., Bharani, R., Magesh, N. S., Godson, P. S., & Chandrasekar, N. (2014). Hydrogeochemistry and groundwater quality appraisal of part of south Chennai coastal aquifers, Tamil Nadu, India using WQI and fuzzy logic method. *Applied Water Science*, 4(4), 341-350.
- Kumar, S. K., Chandrasekar, N., Seralathan, P., Godson, P. S., & Magesh, N. S. (2012). Hydrogeochemical study of shallow carbonate aquifers, Rameswaram Island, India. *Environmental Monitoring and Assessment*, 184(7), 4127-4138.
- Kumar, S. K., Rammohan, V., Sahayam, J. D., & Jeevanandam, M. (2009). Assessment of groundwater quality and hydrogeochemistry of Manimuktha River basin, Tamil Nadu, India. *Environmental Monitoring and Assessment*, 159(1), 341-351.
- Kura, N. U., Ramli, M. F., Ibrahim, S., Sulaiman, W. N. A., & Aris, A. Z. (2014). An integrated assessment of seawater intrusion in a small tropical island using geophysical, geochemical, and geostatistical techniques. *Environmental Science and Pollution Research*, 21(11), 7047-7064.
- Kura, N. U., Ramli, M. F., Ibrahim, S., Sulaiman, W. N. A., Aris, A. Z., Tanko, A. I., & Zaudi, M. A. (2015a). Assessment of groundwater vulnerability to anthropogenic pollution and seawater intrusion in a small tropical island using index-based methods. *Environmental Science and Pollution Research*, 22(2), 1512-1533.
- Kura, N. U., Ramli, M. F., Sulaiman, W. N. A., Ibrahim, S., & Aris, A. Z. (2015b). An overview of groundwater chemistry studies in Malaysia. *Environmental Science and Pollution Research*, 1-19.
- Laaksoharju, M., Skarman, C., & Skarman, E. (1999). Multivariate mixing and mass balance (M3) calculations, a new tool for decoding hydrogeochemical information. *Applied Geochemistry*, 14(7), 861-871.
- Laftouhi, N. E., Vanclooster, M., Jalal, M., Witam, O., Aboufirassi, M., Bahir, M., & Persoons, É. (2003). Groundwater nitrate pollution in the Essaouira Basin (Morocco). *Comptes Rendus Geoscience*, 335(3), 307-317.
- Lakshmanan, E., Kannan, R., & Kumar, M. S. (2003). Major ion chemistry and identification of hydrogeochemical processes of ground water in a part of Kancheepuram district, Tamil Nadu, India. *Environmental Geosciences*, 10(4), 157-166.



- Lan, Y., Jin, M., Yan, C., & Zou, Y. (2015). Schemes of groundwater exploitation for emergency water supply and their environmental impacts on Jiujiang City, China. *Environmental Earth Sciences*, 73(5), 2365-2376.
- Langenegger, O., & Mundial, B. (1987). Groundwater quality in rural areas of Western Africa. In *Groundwater quality in rural areas of Western Africa*. World Bank. Regional Water and Sanitation Group.
- Laurent, M., François, A., & Marie, M. J. (2010). Assessment of groundwater quality during dry season in Southeastern Brazzaville, Congo. *International Journal of Applied Biology and Pharmaceutical Technology*, 1(3), 762-769.
- Lawrence, A., Gooddy, D., Kanatharana, P., Meesilp, W., & Ramnarong, V. (2000). Groundwater evolution beneath Hat Yai, a rapidly developing city in Thailand. *Hydrogeology Journal*, 8(5), 564-575.
- Ledoux, E., Gomez, E., Monget, J. M., Viavattene, C., Viennot, P., Ducharne, A., Benoit, M., Mignolet, C., Schott, C. & Mary, B. (2007). Agriculture and groundwater nitrate contamination in the Seine basin. The STICS–MODCOU modelling chain. *Science of the Total Environment*, 375(1), 33-47.
- Lee, S. M., Min, K. D., Woo, N. C., Kim, Y. J., & Ahn, C. H. (2003). Statistical models for the assessment of nitrate contamination in urban groundwater using GIS. *Environmental Geology*, 44(2), 210-221.
- Lee, Y. W., Dahab, M. F., & Bogardi, I. (1992). Nitrate risk management under uncertainty. *Journal of Water Resources Planning and Management*, 118(2), 151-165.
- Li, P., Wu, J., & Qian, H. (2013). Assessment of groundwater quality for irrigation purposes and identification of hydrogeochemical evolution mechanisms in Pengyang County, China. *Environmental Earth Sciences*, 69(7), 2211-2225.
- Li, P., Wu, J., Qian, H., Lyu, X., & Liu, H. (2014). Origin and assessment of groundwater pollution and associated health risk: a case study in an industrial park, northwest China. *Environmental Geochemistry and Health*, 36(4), 693-712.
- Li, P., Wu, J., Qian, H., Zhang, Y., Yang, N., Jing, L., & Yu, P. (2016). Hydrogeochemical characterization of groundwater in and around a wastewater irrigated forest in the southeastern edge of the Tengger desert, Northwest China. *Exposure and Health*, 1-18.
- Li, X., Zhang, L., & Hou, X. (2008). Use of hydrogeochemistry and environmental isotopes for evaluation of groundwater in Qingshuihe Basin, northwestern China. *Hydrogeology Journal*, 16(2), 335-348.
- Lin, C. Y., Abdullah, M. H., Praveena, S. M., Yahaya, A. H. B., & Musta, B. (2012). Delineation of temporal variability and governing factors influencing the spatial variability of shallow groundwater chemistry in a tropical sedimentary island. *Journal of Hydrology*, 432, 26-42.
- Liu, C. W., Lin, C. N., Jang, C. S., Ling, M. P., & Tsai, J. W. (2011). Assessing nitrate contamination and its potential health risk to Kinmen residents. *Environmental Geochemistry and Health*, 33(5), 503-514.
- Liu, C. W., Lin, K. H., & Kuo, Y. M. (2003). Application of factor analysis in the assessment of groundwater quality in a blackfoot disease area in Taiwan. *Science of the Total Environment*, 313(1), 77-89.

- Lloyd, J. W. (1965). The hydrochemistry of the aquifers of north-eastern Jordan. *Journal of Hydrology*, 3(3-4), 319-330.
- Lloyd, J. W., & Heathcote, J. A. A. (1985). Natural inorganic hydrochemistry in relation to ground water.
- Logeshkumaran, A., Magesh, N. S., Godson, P. S., & Chandrasekar, N. (2015). Hydro-geochemistry and application of water quality index (WQI) for groundwater quality assessment, Anna Nagar, part of Chennai City, Tamil Nadu, India. *Applied Water Science*, 5(4), 335-343.
- Lu, Y., Tang, C., Chen, J., & Yao, H. (2016). Assessment of major ions and heavy metals in groundwater: a case study from Guangzhou and Zhuhai of the Pearl River Delta, China. *Frontiers of Earth Science*, 10(2), 340-351.
- Ma, F. S., Yang, Y. S., Yuan, R. M., Cai, Z., & Pan, S. (2007). Study of shallow groundwater quality evolution under saline intrusion with environmental isotopes and geochemistry. *Environmental Geology*, 51(6), 1009-1017.
- Ma, J., Ding, Z., Wei, G., Zhao, H., & Huang, T. (2009). Sources of water pollution and evolution of water quality in the Wuwei basin of Shiyang river, Northwest China. *Journal of Environmental Management*, 90(2), 1168-1177.
- MacDonald, S. (1967). The geology and mineral resources of North Kelantan and North Terengganu. Geological Survey of Malaysia, Memoir 10, 202p.
- Machiwal, D., Jha, M. K., & Mal, B. C. (2011). GIS-based assessment and characterization of groundwater quality in a hard-rock hilly terrain of Western India. *Environmental Monitoring and Assessment*, 174(1), 645-663.
- Mahlknecht, J., Steinich, B., & De León, I. N. (2004). Groundwater chemistry and mass transfers in the Independence aquifer, central Mexico, by using multivariate statistics and mass-balance models. *Environmental Geology*, 45(6), 781-795.
- Mahvi, A. H., Nouri, J., Babaei, A. A., & Nabizadeh, R. (2005). Agricultural activities impact on groundwater nitrate pollution. *International Journal of Environmental Science & Technology*, 2(1), 41-47.
- Malaysia Meteorological Department (MMD) (2017). *Monthly rainfall data of Kota Bharu from 1989-2017*. Malaysia Meteorological Department, Kuala Lumpur.
- Manap, M. A., Nampak, H., Pradhan, B., Lee, S., Sulaiman, W. N. A., & Ramli, M. F. (2014). Application of probabilistic-based frequency ratio model in groundwater potential mapping using remote sensing data and GIS. *Arabian Journal of Geosciences*, 7(2), 711-724.
- Manap, M. A., Sulaiman, W. N. A., Ramli, M. F., Pradhan, B., & Surip, N. (2013). A knowledge-driven GIS modeling technique for groundwater potential mapping at the Upper Langat Basin, Malaysia. *Arabian Journal of Geosciences*, 6(5), 1621-1637.
- Marinov, I., & Marinov, A. M. (2014). A coupled mathematical model to predict the influence of nitrogen fertilization on crop, soil and groundwater quality. *Water Resources Management*, 28(15), 5231-5246.
- Martínez, D., & Bocanegra, E. (2002). Hydrogeochemistry and cation-exchange processes in the coastal aquifer of Mar Del Plata, Argentina. *Hydrogeology Journal*, 10(3), 393-408.
- McGarigal, K., Cushman, S., & Stafford, S. G. (2000). *Multivariate statistics for wildlife and ecology research* (Vol. 279). New York: Springer.

- Melloul, A., & Collin, M. (1992). The 'principal components' statistical method as a complementary approach to geochemical methods in water quality factor identification; application to the Coastal Plain aquifer of Israel. *Journal of Hydrology*, 140(1), 49-73.
- Menció, A., & Mas-Pla, J. (2008). Assessment by multivariate analysis of groundwater-surface water interactions in urbanized Mediterranean streams. *Journal of Hydrology*, 352(3), 355-366.
- Minerals and Geoscience Department (MGD) (2008). *Geological map State of Peninsular Malaysia (First Edition, 2008)*. Ministry of Natural Resources and Environment. Kuala Lumpur, Malaysia.
- Ministry of Agriculture (MOA) (2010). *Land use map of Peninsular Malaysia*. Department of Agriculture. Kuala Lumpur, Malaysia.
- Ministry of Health (MOH). (2004). *National Guidelines for Drinking Water Quality*. Ministry of Health, Kuala Lumpur.
- Moghaddam, A. A., & Fijani, E. (2009). Hydrogeologic framework of the Maku area basalts, northwestern Iran. *Hydrogeology Journal*, 17(4), 949.
- Mohamad, D. B. (1983). Report on the environmental isotopic investigations in Lower Kelantan River Basin, Malaysia-Part III. IAEA Regional Cooperative Agreement-MAL 2623/R1/AG, 11p.
- Mohamad, D. B., Ali, R. M., Chong, F. S., Yurtsever, Y., Gonfiantini, R. and Calf, G. (1987). Study of groundwater recharge in shallow and deep aquifers in Lower Kelantan River Basin with environmental isotopes, Malaysia. In *Isotope Techniques in Water Resources Development; IAEA- Proceeding*; pp758-763; Vienna
- Mohapatra, P. K., Vijay, R., Pujari, P. R., Sundaray, S. K., & Mohanty, B. P. (2011). Determination of processes affecting groundwater quality in the coastal aquifer beneath Puri city, India: a multivariate statistical approach. *Water Science and Technology*, 64(4), 809-817.
- Mokhtar, M. B., Aris, A. Z., Abdullah, M. H., Yusoff, M. K., Abdullah, M., Idris, Abdullah, M. P, Idris, A. R & Raja Uzir, R. I (2009). A pristine environment and water quality in perspective: Maliau Basin, Borneo's mysterious world. *Water and Environment Journal*, 23(3), 219-228.
- Mondal, N. C., Singh, V. P., Singh, S., & Singh, V. S. (2011). Hydrochemical characteristic of coastal aquifer from Tuticorin, Tamil Nadu, India. *Environmental Monitoring and Assessment*, 175(1-4), 531-550.
- Montcoudiol, N., Molson, J., & Lemieux, J. M. (2015). Groundwater geochemistry of the Outaouais Region (Québec, Canada): a regional-scale study. *Hydrogeology Journal*, 23(2), 377-396.
- Morris, B. L., Lawrence, A. R., Chilton, P. J. C., Adams, B., Calow, R. C., & Klinck, B. A. (2003). *Groundwater and its susceptibility to degradation: a global assessment of the problem and options for management* (Vol. 3). United Nations Environment Programme.
- Mukherjee, A., Fryar, A. E., & Howell, P. D. (2007). Regional hydrostratigraphy and groundwater flow modeling in the arsenic-affected areas of the western Bengal basin, West Bengal, India. *Hydrogeology Journal*, 15(7), 1397.

- Murad, A. A., & Krishnamurthy, R. V. (2004). Factors controlling groundwater quality in Eastern United Arab Emirates: a chemical and isotopic approach. *Journal of Hydrology*, 286(1), 227-235.
- Nag, S. K., & Das, S. (2017). Assessment of groundwater quality from Bankura I and II Blocks, Bankura District, West Bengal, India. *Applied Water Science*, 1-16.
- Nag, S. K., & Ghosh, P. (2013). Variation in groundwater levels and water quality in Chhatna Block, Bankura district, West Bengal-a GIS approach. *Journal of the Geological Society of India*, 81(2), 261-280.
- Nagarajan, R., Rajmohan, N., Mahendran, U., & Senthamilkumar, S. (2010). Evaluation of groundwater quality and its suitability for drinking and agricultural use in Thanjavur city, Tamil Nadu, India. *Environmental Monitoring and Assessment*, 171(1-4), 289-308.
- Nagaraju, A., Kumar, K. S., & Thejaswi, A. (2014). Assessment of groundwater quality for irrigation: a case study from Bandalamottu lead mining area, Guntur District, Andhra Pradesh, South India. *Applied Water Science*, 4(4), 385-396.
- Naik, P. K., Awasthi, A. K., Anand, A. V. S. S., & Behera, P. N. (2009). Hydrogeochemistry of the Koyna River basin, India. *Environmental Earth Sciences*, 59(3), 613-629.
- Nandimandalam, J. R. (2012). Evaluation of hydrogeochemical processes in the Pleistocene aquifers of middle Ganga Plain, Uttar Pradesh, India. *Environmental Earth Sciences*, 65(4), 1291-1308.
- Narany, T. S., Aris, A. Z., Sefie, A., & Keesstra, S. (2017). Detecting and predicting the impact of land use changes on groundwater quality, a case study in Northern Kelantan, Malaysia. *Science of The Total Environment*, 599, 844-853.
- Narany, T. S., Sefie, A., & Aris, A. Z. (2018). The long-term impacts of anthropogenic and natural processes on groundwater deterioration in a multilayered aquifer. *Science of The Total Environment*, 630, 931-942.
- Nas, B., & Berktaş, A. (2010). Groundwater quality mapping in urban groundwater using GIS. *Environmental Monitoring and Assessment*, 160(1), 215-227.
- National Economic Planning Unit. (2006). *9<sup>th</sup> Malaysia Plan*. Kuala Lumpur: NEPU, Malaysia
- National Research Council. (1995). *Nitrate and nitrite in drinking water*. National Academies Press.
- Nawang, W. M. (1988). River-aquifer interrelation near public groundwater intake in North Kelantan, Malaysia. In *Proceedings of the VIth IWRA World Congress on Water Resources: Water for World Development*, May 29-June 3, 1988, Ottawa, Canada, Vol. II, p138-148.
- Nawang, W. M. and Kishi, Y. (1990). Saltwater encroachment prediction in multilayered aquifer at Tanjung Mas, Kelantan. In *Proceedings of the 2<sup>nd</sup> International Groundwater Conference*, IGW2-90, Jun. 25-29, 1990, Kota Bharu, Kelantan, Malaysia.
- Nazzal, Y., Ahmed, I., Al-Arifi, N. S., Ghrefat, H., Batayneh, A., Abuamarah, B. A., & Zaidi, F. K. (2015). A combined hydrochemical-statistical analysis of Saq aquifer, northwestern part of the Kingdom of Saudi Arabia. *Geosciences Journal*, 19(1), 145-155.

- Nematollahi, M. J., Ebrahimi, P., Razmara, M., & Ghasemi, A. (2016). Hydrogeochemical investigations and groundwater quality assessment of Torbat-Zaveh plain, Khorasan Razavi, Iran. *Environmental Monitoring and Assessment*, 188(1), 1-21.
- Nghi, T., Toan, N. Q., Minh, N. D., & Van Vuong, N. (1991). Quaternary sedimentation of the principal deltas of Vietnam. *Journal of Southeast Asian Earth Sciences*, 6(2), 103-110.
- Nguyen, T. T., Kawamura, A., Tong, T. N., Nakagawa, N., Amaguchi, H., & Gilbuena Jr, R. (2015). Hydrogeochemical assessment of groundwater quality during dry and rainy seasons for the two main aquifers in Hanoi, Vietnam. *Environmental Earth Sciences*, 73(8), 4287-4303.
- Nickson, R. T., McArthur, J. M., Shrestha, B., Kyaw-Nyint, T. O., & Lowrt, D. (2005). Arsenic and other drinking water quality issues, Muzaffargarh District, Pakistan. *Applied Geochemistry*, 20, 55-68.
- Nishanthiny, S. C., Thushyanthy, M., Barathithasan, T., & Saravanan, S. (2010). Irrigation water quality based on hydro chemical analysis, Jaffna, Sri Lanka. *American-Eurasian Journal of Agricultural & Environmental Sciences*, 7(1), 100-102.
- Nisi, B., Bucciatti, A., Vaselli, O., Perini, G., Tassi, F., Minissale, A., & Montegrossi, G. (2008). Hydrogeochemistry and strontium isotopes in the Arno River Basin (Tuscany, Italy): constraints on natural controls by statistical modeling. *Journal of Hydrology*, 360(1), 166-183.
- Noor, I. B. M. (1980). *Prefeasibility study of potential groundwater development in Kelantan, Malaysia*. Ph.D. Thesis. University of Birmingham, United Kingdom, 427p.
- Noor, I. B. M. (1985). Groundwater management of Kelantan Aquifer, Malaysia. Groundwater and Mineral Exploration (GAME), Bangkok, Thailand.
- Nur, A., & Ayuni, K. N. (2011). Hydro-geophysical study of michika and environs, northeast Nigeria. *International Journal of Physical Sciences*, 6(34), 7816-7827.
- Okiongbo, K. S., & Douglas, R. K. (2015). Evaluation of major factors influencing the geochemistry of groundwater using graphical and multivariate statistical methods in Yenagoa city, Southern Nigeria. *Applied Water Science*, 5(1), 27-37.
- Olajire, A. A., & Imeokparia, F. E. (2001). Water quality assessment of Osun River: Studies on inorganic nutrients. *Environmental Monitoring and Assessment*, 69(1), 17-28.
- Olobaniyi, S. B., & Owoyemi, F. B. (2006). Characterization by factor analysis of the chemical facies of groundwater in the deltaic plain-sands aquifer of Warri, Western Niger Delta Nigeria. *African Journal of Science and Technology*, 7(1), 73-81.
- Osman, R., Saim, N., Juahir, H., & Abdullah, M. P. (2012). Chemometric application in identifying sources of organic contaminants in Langat river basin. *Environmental Monitoring and Assessment*, 184(2), 1001-1014.
- Oyarzún, R., Jofré, E., Morales, P., Maturana, H., Oyarzún, J., Kretschmer, N. & Aravena, R. (2015). A hydrogeochemistry and isotopic approach for the assessment of surface water-groundwater dynamics in an arid basin: the Limarí watershed, North-Central Chile. *Environmental Earth Sciences*, 73(1), 39-55

- Pacheco, F. A., & Szocs, T. (2006). "Dedolomitization reactions" driven by anthropogenic activity on loessy sediments, SW Hungary. *Applied Geochemistry*, 21(4), 614-631.
- Parkhurst, D. L., & Appelo, C. A. J. (1999). User's guide to PHREEQC (Version 2): A computer program for speciation, batch-reaction, one-dimensional transport, and inverse geochemical calculations.
- Pazand, K. (2016). Geochemical and statistical evaluation of groundwater in Razan basin, Western Iran. *Carbonates and Evaporites*, 31(2), 179-185.
- Pazand, K., & Javanshir, A. R. (2013). Hydrogeochemistry and arsenic contamination of groundwater in the Rayen area, southeastern Iran. *Environmental Earth Sciences*, 70(6), 2633-2644.
- Petrides, B., & Cartwright, I. (2006). The hydrogeology and hydrogeochemistry of the Barwon Downs Graben aquifer, southwestern Victoria, Australia. *Hydrogeology Journal*, 14(5), 809-826.
- Pfeiffer, D. & Chong, F. S. (1974). *Groundwater exploration in Kota Bharu, Report No. I*. Geological Survey of Malaysia, Report GPH 1/1974, 8p.
- Pfeiffer, D. & Tiedemann, O. (1986). Hydrogeological study and the development of groundwater in Kelantan Coastal Plain, Malaysia. *Natural Resources & Development*, 24, p72-82.
- Piper, A. M. (1944). A graphic procedure in the geochemical interpretation of water-analyses. *Transactions American Geophysical Union*, 25(6), 914-928.
- Prasad, R. K., Mondal, N. C., Banerjee, P., Nandakumar, M. V., & Singh, V. S. (2008). Deciphering potential groundwater zone in hard rock through the application of GIS. *Environmental Geology*, 55(3), 467-475.
- Prasanna, M. V., Chidambaram, S., & Srinivasamoorthy, K. (2010a). Statistical analysis of the hydrogeochemical evolution of groundwater in hard and sedimentary aquifers system of Gadilam river basin, South India. *Journal of King Saud University-Science*, 22(3), 133-145.
- Prasanna, M. V., Chidambaram, S., Gireesh, T. V., & Ali, T. J. (2011b). A study on hydrochemical characteristics of surface and sub-surface water in and around Perumal Lake, Cuddalore district, Tamil Nadu, South India. *Environmental Earth Sciences*, 63(1), 31-47.
- Prasanna, M. V., Chidambaram, S., Hameed, A. S., & Srinivasamoorthy, K. (2011a). Hydrogeochemical analysis and evaluation of groundwater quality in the Gadilam river basin, Tamil Nadu, India. *Journal of Earth System Science*, 120(1), 85-98.
- Prasanna, M. V., Chidambaram, S., Shahul Hameed, A., & Srinivasamoorthy, K. (2010b). Study of evaluation of groundwater in Gadilam basin using hydrogeochemical and isotope data. *Environmental Monitoring and Assessment*, 168(1), 63-90.
- Praveena, S. M., Abdullah, M. H., Bidin, K., & Aris, A. Z. (2011). Understanding of groundwater salinity using statistical modeling in a small tropical island, East Malaysia. *The Environmentalist*, 31(3), 279.
- Qiu, J. (2010). China faces up to groundwater crisis. *Nature*, 466(7304), 308.
- Raghunath, H. M. (1987). *Ground water*. New Age International.
- Rajmohan, N., & Elango, L. (2005). Distribution of iron, manganese, zinc and atrazine in groundwater in parts of Palar and Cheyyar river basins, South India. *Environmental Monitoring and Assessment*, 107, 115-131.

- Rajmohan, N., Al-Futaisi, A., & Al-Touqi, S. (2009). Geochemical process regulating groundwater quality in a coastal region with complex contamination sources: Barka, Sultanate of Oman. *Environmental Earth Sciences*, 59(2), 385-398.
- Rajmohan, N., Elango, L., & Elampoornam, T. (1997). Seasonal and spatial variation in magnesium and chloride concentration in ground waters of Nagai Quaid E Milleth district in Tamil Nadu. *Indian Journal of Environmental Protection*, 17(6), 448-453.
- Raju, N. J. (2006). Iron contamination in groundwater: A case from Tirumala-Tirupati environs, India. *The Researcher*, 1(1), 28-31.
- Raju, N. J., Ram, P., & Dey, S. (2009). Groundwater quality in the lower Varuna River basin, Varanasi district, Uttar Pradesh, India. *Journal of Geological Society of India*, 73, 178-192.
- Raju, N. J., Shukla, U. K., & Ram, P. (2011). Hydrogeochemistry for the assessment of groundwater quality in Varanasi: a fast-urbanizing center in Uttar Pradesh, India. *Environmental Monitoring and Assessment*, 173(1), 279-300.
- Rao, N. S. (2002). Geochemistry of groundwater in parts of Guntur district, Andhra Pradesh, India. *Environmental Geology*, 41(5), 552-562.
- Rao, N. S. (2006). Seasonal variation of groundwater quality in a part of Guntur District, Andhra Pradesh, India. *Environmental Geology*, 49(3), 413-429.
- Rao, N. S. (2008). Iron content in groundwaters of Visakhapatnam environs, Andhra Pradesh, India. *Environmental Monitoring and Assessment*, 136(1-3), 437-447.
- Rao, N. S., Rao, P. S., & Varma, D. D. (2013). Spatial variations of groundwater vulnerability using cluster analysis. *Journal of the Geological Society of India*, 81(5), 685-697.
- Rao, N. S., Rao, P. S., Reddy, G. V., Nagamani, M., Vidyasagar, G., & Satyanarayana, N. L. V. V. (2012). Chemical characteristics of groundwater and assessment of groundwater quality in Varaha River Basin, Visakhapatnam District, Andhra Pradesh, India. *Environmental Monitoring and Assessment*, 184(8), 5189-5214.
- Rao, P. N., Rao, G. B., Subrahmanyam, A., Rao, N. S., & Paul, K. J. (2015). Chemistry of groundwater in Erravagu sub-basin, Guntur district, Andhra Pradesh. *Journal of Applied Geochemistry*, 17(1), 56.
- Rasouli, F., Pouya, A. K., & Cheraghi, S. A. M. (2012). Hydrogeochemistry and water quality assessment of the Kor-Sivand Basin, Fars province, Iran. *Environmental Monitoring and Assessment*, 184(8), 4861-4877.
- Ravikumar, P., & Somashekar, R. K. (2011). Geochemistry of groundwater, Markandeya River Basin, Belgaum district, Karnataka State, India. *Chinese Journal of Geochemistry*, 30(1), 51-74.
- Ravikumar, P., & Somashekar, R. K. (2015). Principal component analysis and hydrochemical facies characterization to evaluate groundwater quality in Varahi river basin, Karnataka state, India. *Applied Water Science*, 1-11.
- Ravikumar, P., Somashekar, R. K., & Angami, M. (2011). Hydrochemistry and evaluation of groundwater suitability for irrigation and drinking purposes in the Markandeya River basin, Belgaum District, Karnataka State, India. *Environmental Monitoring and Assessment*, 173(1), 459-487.
- Re, V., Sacchi, E., Mas-Pla, J., Menció, A., & El Amrani, N. (2014). Identifying the effects of human pressure on groundwater quality to support water management

- strategies in coastal regions: a multi-tracer and statistical approach (Bou-Areg region, Morocco). *Science of the Total Environment*, 500, 211-223.
- Reghunath, R., Murthy, T. S., & Raghavan, B. R. (2002). The utility of multivariate statistical techniques in hydrogeochemical studies: an example from Karnataka, India. *Water Research*, 36(10), 2437-2442.
- Reimann, C., & Filzmoser, P. (2000). Normal and lognormal data distribution in geochemistry: death of a myth. Consequences for the statistical treatment of geochemical and environmental data. *Environmental Geology*, 39(9), 1001-1014.
- Reisenhofer, E., Adami, G., & Favretto, A. (1996). Heavy metals and nutrients in coastal, surface seawaters (Gulf of Trieste, Northern Adriatic Sea): an environmental study by factor analysis. *Fresenius' Journal of Analytical Chemistry*, 354(5), 729-734.
- Rey, J., Martínez, J., Barberá, G. G., García-Aróstegui, J. L., García-Pintado, J., & Martínez-Vicente, D. (2013). Geophysical characterization of the complex dynamics of groundwater and seawater exchange in a highly stressed aquifer system linked to a coastal lagoon (SE Spain). *Environmental Earth Sciences*, 70(5), 2271-2282.
- Rhoades, J. D., Kandiah A., & Mashali A. M. (1992). *The use of saline waters for crop production*. Rome: Food Agriculture Organization of the United Nations (FAO), No. 48.
- Richards, L. A. (1947). Diagnosis and Improvement of Saline and Alkaline Soils. *Soil Science*, 64, 432.
- Richards, L. A. (1954). *Diagnosis and improvement of saline and alkali soils* (Vol. 78, No. 2, p. 154). LWW.
- Rina, K., Datta, P. S., Singh, C. K., & Mukherjee, S. (2013). Characterization and evaluation of processes governing the groundwater quality in parts of the Sabarmati basin, Gujarat using hydrochemistry integrated with GIS. *Hydrological Processes*, 26(10), 1538-1551.
- Romic, M., & Romic, D. (2003). Heavy metal distribution in agricultural topsoils in urban area. *Environmental Geology*, 43, 795-805.
- Roslan, I., Shamshuddin, J., Fauziah, C. I., & Anuar, A. R. (2010). Occurrence and properties of soils on sandy beach ridges in the Kelantan-Terengganu Plains, Peninsular Malaysia. *Catena*, 83(1), 55-63.
- Rouabhia, A., Baali, F., & Fehdi, C. (2010). Impact of agricultural activity and lithology on groundwater quality in the Merdja area, Tebessa, Algeria. *Arabian Journal of Geosciences*, 3(3), 307-318.
- Rouabhia, A., Baali, F., Fehdi, C., Kherici, N., & Djabri, L. (2009). Hydrochemical and isotopic investigation of a sandstone aquifer groundwater in a semi-arid region, El Ma El Abiod, Algeria. *Environmental Geology*, 57(8), 1699-1705.
- Sadashivaiah, C., Ramakrishnaiah, C. R., & Ranganna, G. (2008). Hydrochemical analysis and evaluation of groundwater quality in Tumkur Taluk, Karnataka State, India. *International Journal of Environmental Research and Public Health*, 5(3), 158-164.
- Saghavani, S. R., Yusoff, I., Tahir, W. Z. W. M., & Othman, Z. (2014). Comparison of water table fluctuation and chloride mass balance methods for recharge estimation in a tropical rainforest climate: a case study from Kelantan River catchment, Malaysia. *Environmental Earth Sciences*, 73(8), 4419-4428.



- Saim, N., Osman, R., Spian, D. R. S. A., Jaafar, M. Z., Juahir, H., Abdullah, M. P., & Ab Ghani, F. (2009). Chemometric approach to validating faecal sterols as source tracer for faecal contamination in water. *Water Research*, 43(20), 5023-5030.
- Saleem, A., Dandigi, M. N., & Kumar, K. V. (2012). Correlation-regression model for physico-chemical quality of groundwater in the South Indian city of Gulbarga. *African Journal of Environmental Science and Technology*, 6(9), 353-364.
- Saleh, A., Al-Ruwaih, F., & Shehata, M. (1999). Hydrogeochemical processes operating within the main aquifers of Kuwait. *Journal of Arid Environments*, 42(3), 195-209.
- Samsudin, A. R., Haryono, A., Hamzah, U., & Rafek, A. G. (2008). Salinity mapping of coastal groundwater aquifers using hydrogeochemical and geophysical methods: a case study from north Kelantan, Malaysia. *Environmental Geology*, 55(8), 1737-1743.
- Sankaramakrishnan, N., Sharma, A. K., & Iyengar, L. (2008). Contamination of nitrate and fluoride in ground water along the Ganges Alluvial Plain of Kanpur district, Uttar Pradesh, India. *Environmental Monitoring and Assessment*, 146(1-3), 375-382.
- Sarbu, C., & Pop, H. F. (2005). Principal component analysis versus fuzzy principal component analysis: a case study: the quality of Danube water (1985-1996). *Talanta*, 65(5), 1215-1220.
- Saxena, V. K., Mondal, N. C., & Singh, V. S. (2004). Evaluation of hydrogeochemical parameters to delineate fresh groundwater zones in coastal aquifers. *Journal of Applied Geochemistry*, 6, 245-254.
- Schirmer, M., Leschik, S., & Musolff, A. (2013). Current research in urban hydrogeology-a review. *Advances in Water Resources*, 51, 280-291.
- Schnallinger, B. and Sofner, B. (1994). *Hydrochemistry, Isotopes Composition Hydrochemistry, Isotope Composition (18O, D) and Genesis of the Groundwater in the Lower Kelantan River Basin, Malaysia*. Technical Cooperation Project Report No. 8822.25.6, BGR, Hannover, 19p.
- Schwartz, F. W., & Zhang, H. (2003). *Groundwater*. John Wiley & Sons Inc., New York.
- Sefie, A., Aris, A. Z., Ramli, M. F., Narany, T. S., Shamsuddin, M. K. N., Saadudin, S. B., & Zali, M. A. (2018). Hydrogeochemistry and groundwater quality assessment of the multilayered aquifer in Lower Kelantan Basin, Kelantan, Malaysia. *Environmental Earth Sciences*, 77(10), 397.
- Senthilkumar, M., & Elango, L. (2013). Geochemical processes controlling the groundwater quality in lower Palar river basin, southern India. *Journal of Earth System Science*, 122(2), 419-432.
- Seong, K. T., Pfeiffer, D. & Tiedmann, O. (1977). *Hydrogeological Report Bachok, Kelantan*. Geological Survey of Malaysia, Report GPH 3/1977, 6p.
- Sepakat Setia Perunding. (1990). *Preliminary environmental impact assessment for North Kelantan Water Supply with Groundwater Sources*. Consultant Report to Jabatan Kerja Raya Kelantan, December 1990.
- Shamrukh, M., Corapcioglu, M. Y., & Hassona, F. A. (2001). Modeling the effect of chemical fertilizers on ground water quality in the Nile Valley Aquifer, Egypt. *Ground Water*, 39(1), 59-67.

- Shamshuddin, J., Kang, M. S. K. E., Fauziah, C. I., & Panhwar, Q. A. (2013). On the pyritization of the coastal sediments in the Malay Peninsula during the Holocene and its effects on soil. *Malaysian Journal of Soil Science*, 17, 1-15.
- Shanmugasundharam, A., Kalpana, G., Mahapatra, S. R., Sudharson, E. R., & Jayaprakash, M. (2017). Assessment of Groundwater quality in Krishnagiri and Vellore Districts in Tamil Nadu, India. *Applied Water Science*, 7(4), 1869-1879.
- Sharma, M. L., Kin, P. G., Mok, J. Y. M., & Gerritse, R. G. (1991). Nitrate and phosphate leaching beneath irrigated horticultural systems. In *Modelling the Fate of Chemicals in the Environment, Proceedings of a national workshop, ANU, Canberra*.
- Sheikhy Narany, T., Ramli, M. F., Aris, A. Z., Sulaiman, W. N. A., Juahir, H., & Fakharian, K. (2014). Identification of the hydrogeochemical processes in groundwater using classic integrated geochemical methods and geostatistical techniques, in Amol-Babol plain, Iran. *The Scientific World Journal*, 2014.
- Shin, C. F., Preiffer, D., Sidhu, H. S. & Tiedmann, O. (1974). Groundwater exploration in Kota Bharu. Geological Survey of Malaysia, Report No. 11, 7p.
- Shrestha, S., & Kazama, F. (2007). Assessment of surface water quality using multivariate statistical techniques: A case study of the Fuji river basin, Japan. *Environmental Modelling & Software*, 22(4), 464-475.
- Shrestha, S., & Muangthong, S. (2014). Assessment of surface water quality of Songkhram River (Thailand) using environmetric techniques. *International Journal of River Basin Management*, 12(4), 341-356.
- Siddiqi, M., Kumar, R., Fazili, Z., Spiegelhalder, B., & Preussmann, R. (1992). Increased exposure to dietary amines and nitrate in a population at high risk of oesophageal and gastric cancer in Kashmir (India). *Carcinogenesis*, 13(8), 1331-1335.
- Simeonov, V., Stratis, J. A., Samara, C., Zachariadis, G., Voutsas, D., Anthemidis, A., Sofoniou, M., & Kouimtzis, T. (2003). Assessment of the surface water quality in Northern Greece. *Water research*, 37(17), 4119-4124.
- Simsek, C., & Gunduz, O. (2007). IWQ Index: a GIS-integrated technique to assess irrigation water quality. *Environmental Monitoring and Assessment*, 128(1), 277-300.
- Singaraja, C., Chidambaram, S., Anandhan, P., Prasanna, M. V., Thivya, C., Thilagavathi, R., & Sarathidasan, J. (2014a). Hydrochemistry of groundwater in a coastal region and its repercussion on quality, a case study-Thoothukudi district, Tamil Nadu, India. *Arabian Journal of Geosciences*, 7(3), 939-950.
- Singaraja, C., Chidambaram, S., Prasanna, M. V., Thivya, C., & Thilagavathi, R. (2014b). Statistical analysis of the hydrogeochemical evolution of groundwater in hard rock coastal aquifers of Thoothukudi district in Tamil Nadu, India. *Environmental Earth Sciences*, 71(1), 451-464.
- Singh, A. K., Mondal, G. C., Singh, T. B., Singh, S., Tewary, B. K., & Sinha, A. (2012). Hydrogeochemical processes and quality assessment of groundwater in Dumka and Jamtara districts, Jharkhand, India. *Environmental Earth Sciences*, 67(8), 2175-2191.
- Singh, A. K., Tewary, B. K., & Sinha, A. (2011). Hydrochemistry and quality assessment of groundwater in part of Noida metropolitan city, Uttar Pradesh. *Journal of the Geological Society of India*, 78(6), 523-540.

- Singh, K. P., Malik, A., & Sinha, S. (2005). Water quality assessment and apportionment of pollution sources of Gomti river (India) using multivariate statistical techniques-a case study. *Analytica Chimica Acta*, 538(1), 355-374.
- Singh, K. P., Malik, A., Mohan, D., & Sinha, S. (2004). Multivariate statistical techniques for the evaluation of spatial and temporal variations in water quality of Gomti River (India)-a case study. *Water Research*, 38(18), 3980-3992.
- Singh, K. P., Malik, A., Singh, V. K., Mohan, D., & Sinha, S. (2005). Chemometric analysis of groundwater quality data of alluvial aquifer of Gangetic plain, North India. *Analytica Chimica Acta*, 550(1), 82-91.
- Singh, N., Singh, R. P., Kamal, V., Sen, R., & Mukherjee, S. (2015). Assessment of hydrogeochemistry and the quality of groundwater in 24-Parganas districts, West Bengal. *Environmental Earth Sciences*, 73(1), 375-386.
- Singh, P. K., Malik, M., Mohan, D., Singh, V. K., & Singh, S. (2006). Evaluation of groundwater quality in northern Indo-Gangetic alluvium region. *Environmental Monitoring and Assessment*, 112, 211-230.
- Singhal, B. B. S., & Gupta, R. P. (2010). *Applied hydrogeology of fractured rocks*. Springer Science & Business Media.
- Sivasankar, V., & Gomathi, R. (2009). Fluoride and other quality parameters in the groundwater samples of Pettaivaithalai and Kulithalai areas of Tamil Nadu, Southern India. *Water Quality, Exposure and Health*, 1(2), 123-134.
- Sofner, B. (1989). Groundwater monitoring and groundwater protection in North Kelantan and Perlis, Malaysia. Technical Cooperation Project Report No. 88.2224.6, BGR, Hannover, 25p.
- Sofner, B. (1992). *Technical Cooperation, Project No. 88.2225.6: Groundwater Monitoring and Groundwater protection at the Geological Survey Department (GSD) of Malaysia: report on an assignment to the GSD from 7.3.-17.6.1992; BGR; Hannover*.
- Sophocleous, M. (2002). Interactions between groundwater and surface water: the state of the science. *Hydrogeology journal*, 10(1), 52-67.
- Sprinkle, C. L. (1989). Geochemistry of the Floridan aquifer system in Florida and in parts of Georgia, South Carolina, and Alabama. *US Geological Survey Professional Paper*, 1403(1).
- Srinivas, Y., Aghil, T. B., Oliver, D. H., Nair, C. N., & Chandrasekar, N. (2015). Hydrochemical characteristics and quality assessment of groundwater along the Manavalakurichi coast, Tamil Nadu, India. *Applied Water Science*, 1-10.
- Srinivas, Y., Oliver, D. H., Raj, A. S., & Chandrasekar, N. (2013). Evaluation of groundwater quality in and around Nagercoil town, Tamilnadu, India: an integrated geochemical and GIS approach. *Applied Water Science*, 3(3), 631-651.
- Srinivasamoorthy, K., Chidambaram, S., Prasanna, M. V., Vasanthavihar, M., Peter, J., & Anandhan, P. (2008a). Identification of major sources controlling groundwater chemistry from a hard rock terrain-a case study from Mettur taluk, Salem district, Tamil Nadu, India. *Journal of Earth System Science*, 117(1), 49-58.
- Srinivasamoorthy, K., Chidambaram, S., Vasanthavigar, M., Prasanna, M. V., & John Peter, A. (2008b). Geochemistry of fluorides in groundwater: Salem District, Tamilnadu, India. *Journal of Environmental Hydrology*, 1, 16-25.

- Srinivasamoorthy, K., Vasanthavigar, M., Chidambaram, S., Anandhan, P., Manivannan, R., & Rajivgandhi, R. (2012). Hydrochemistry of groundwater from Sarabanga minor basin, Tamilnadu, India. *Proceedings of the International Academy of Ecology and Environmental Sciences*, 2(3), 193.
- Srinivasamoorthy, K., Vijayaraghavan, K., Vasanthavigar, M., Rajivgandhi, R., & Sarma, V. S. (2011). Integrated techniques to identify groundwater vulnerability to pollution in a highly industrialized terrain, Tamilnadu, India. *Environmental Monitoring and Assessment*, 182(1-4), 47-60.
- Stiff Jr, H. A. (1951). The interpretation of chemical water analysis by means of patterns. *Journal of Petroleum Technology*, 3(10), 15-3.
- Stites, W., & Kraft, G. J. (2001). Nitrate and chloride loading to groundwater from an irrigated north-central US sand-plain vegetable field. *Journal of Environmental Quality*, 30(4), 1176-1184.
- Subba Rao, N. (2008). Factors controlling the salinity in groundwater in parts of Guntur district, Andhra Pradesh, India. *Environmental Monitoring and Assessment*, 138(1), 327-341.
- Subramani, T., Elango, L., & Damodarasamy, S. R. (2005). Groundwater quality and its suitability for drinking and agricultural use in Chithar River Basin, Tamil Nadu, India. *Environmental Geology*, 47(8), 1099-1110.
- Sun, L., & Gui, H. (2015). Hydro-chemical evolution of groundwater and mixing between aquifers: a statistical approach based on major ions. *Applied Water Science*, 5(1), 97-104.
- Sundaram, B., Feitz, A., Caritat, P. D., Plazinska, A., Brodie, R., Coram, J., & Ransley, T. (2009). Groundwater sampling and analysis—a field guide. *Geoscience Australia, Record*, 27, 95.
- Sundaray, S. K. (2010). Application of multivariate statistical techniques in hydrogeochemical studies-a case study: Brahmani-Koel River (India). *Environmental Monitoring and Assessment*, 164(1-4), 297-310.
- Suratman, S. (1997). *Groundwater protection in North Kelantan, Malaysia: An integrated mapping approach using modelling and GIS* (Ph.D. Thesis, University of Newcastle Upon Tyne).
- Suswanto, T., Shamshuddin, J., Syed Omar, S. R., Mat, P., & Teh, C. B. S. (2007). Effects of lime and fertiliser application in combination with water management on rice (*Oryza sativa*) cultivated on an acid sulfate soil. *Malaysian Journal of Soil Science*, 11, 1-16.
- Suthar, S., Bishnoi, P., Singh, S., Mutiyar, P. K., Nema, A. K., & Patil, N. S. (2009). Nitrate contamination in groundwater of some rural areas of Rajasthan, India. *Journal of Hazardous Materials*, 171(1), 189-199.
- Swan, A. R., & Sandilands, M. (1995). Introduction to geological data analysis. In *International Journal of Rock Mechanics and Mining Sciences and Geomechanics Abstracts* (Vol. 8, No. 32, p. 387A).
- Szabolcs, I., & Darab, C. (1964). The influence of irrigation water of high sodium carbonate content of soils. In *Proceedings of 8<sup>th</sup> international congress of ISSS, Trans, II* (pp. 803-812).

- Talukder, M. S. U., Shirazi, S. M., Niazuddin, M., & Ghosh, A. K. (1999). Groundwater quality and its impact on soil and crop. *Bangladesh Journal of Agricultural Sciences*, 26(3), 153-60.
- Tanabe, S., Saito, Y., Quang, L. V., Hanebuth, T. J. J., Quang L. N. & Kitamura, A. (2006). Holocene evolution of the Song Hong (Red River) delta system, northern Vietnam. *Sediment Geol* 187:29–61
- Tauler, R., Peré-Trepat, E., Lacorte, S., & Barceló, D. (2004). *Chemometrics modelling of environmental data* (Doctoral dissertation, International Environmental Modelling and Software Society).
- Tayfur, G., Kirer, T., & Baba, A. (2008). Groundwater quality and hydrogeochemical properties of Torbalı Region, Izmir, Turkey. *Environmental Monitoring and Assessment*, 146(1), 157-169.
- Teh, T. S. (1976). Evolution of the Older Permatang Series of Kelantan. *Geographica*, 11, pp. 20-29
- Teikeu, W. A., Meli'i, J. L., Nouck, P. N., Tabod, T. C., Nyam, F. E. A., & Aretouyap, Z. (2016). Assessment of groundwater quality in Yaoundé area, Cameroon, using geostatistical and statistical approaches. *Environmental Earth Sciences*, 75(1), 21.
- Teikeu, W. A., Njandjock, P. N., Ndougsa-Mbarga, T., & Tabod, T. C. (2012). Geoelectric investigation for groundwater exploration in Yaoundé area, Cameroon. *International Journal of Geosciences*, 3(03), 640.
- Thakur, J. K., Thakur, R. K., Ramanathan, A. L., Kumar, M., & Singh, S. K. (2011). Arsenic contamination of groundwater in Nepal-an overview. *Water*, 3(1), 1-20.
- Thivya, C., Chidambaram, S., Singaraja, C., Thilagavathi, R., Prasanna, M. V., Anandhan, P., & Jainab, I. (2013). A study on the significance of lithology in groundwater quality of Madurai district, Tamil Nadu (India). *Environment, Development and Sustainability*, 15(5), 1365-1387.
- Thivya, C., Chidambaram, S., Thilagavathi, R., Prasanna, M. V., Singaraja, C., Nepolian, M., & Sundararajan, M. (2014). Identification of the geochemical processes in groundwater by factor analysis in hard rock aquifers of Madurai District, South India. *Arabian Journal of Geosciences*, 7(9), 3767-3777
- Thivya, C., Chidambaram, S., Thilagavathi, R., Prasanna, M. V., Singaraja, C., Adithya, V. S., & Nepolian, M. (2015). A multivariate statistical approach to identify the spatio-temporal variation of geochemical process in a hard rock aquifer. *Environmental Monitoring and Assessment*, 187(9), 1-19.
- Tijani, M. N. (1994). Hydrogeochemical assessment of groundwater in Moro area, Kwara state, Nigeria. *Environmental Geology*, 24(3), 194-202.
- Tijani, M. N. (2004). Evolution of saline waters and brines in the Benue-Trough, Nigeria. *Applied Geochemistry*, 19(9), 1355-1365.
- Tiwari, A. K., & Singh, A. K. (2014). Hydrogeochemical investigation and groundwater quality assessment of Pratapgarh district, Uttar Pradesh. *Journal of the Geological Society of India*, 83(3), 329-343.
- Tiwari, K. K., Prasad, R. N., Chandra, R., & Mondal, N. C. (2012). Geochemical parameters for assessment of groundwater quality around urban and suburban areas of Dausa city in Rajasthan, India. *Journal of Applied Geochemistry*, 14(2), 184-193.

- Tizro, A. T., & Voudouris, K. (2008). Groundwater quality in the semi-arid region of the Chahardouly basin, West Iran. *Hydrological Processes*, 22(16), 3066-3078.
- Todd, D. K. (1980). Groundwater Hydrology, 2nd edn. New York: John Wiley and Sons, (p. 535).
- Touhari, F., Meddi, M., Mehaiguen, M., & Razack, M. (2015). Hydrogeochemical assessment of the Upper Cheliff groundwater (North West Algeria). *Environmental Earth Sciences*, 73(7), 3043-3061.
- Trabelsi, R., Abid, K., & Zouari, K. (2012). Geochemistry processes of the Djeffara palaeogroundwater (Southeastern Tunisia). *Quaternary International*, 257, 43-55.
- Tran, L. T., Larsen, F., Pham, N. Q., Christiansen, A. V., Tran, N., Vu, H. V., Tran, L. V., Hoang, H. V. & Hinsby, K. (2012). Origin and extent of fresh groundwater, salty paleowaters and recent saltwater intrusions in Red River flood plain aquifers, Vietnam. *Hydrogeology Journal*, 20(7), 1295-1313.
- UNEP. (2013). Earthscan, Global Environmental Outlook, United Kingdom: 2000
- UNESCO. (2006), UNESCO water portal newsletter no. 161: Water-related diseases. Available at: <http://www.unesco.org/water/news/newsletter/161.shtml>.
- USEPA. (1993). Wellhead protection: a guide for small communities.
- USSL. (1954). Diagnosis and improvement of salinity and alkaline soil. USDA Hand Book no. 60, Washington.
- Utom, A. U., Odoh, B. I., & Egboka, B. C. (2013). Assessment of hydrogeochemical characteristics of groundwater quality in the vicinity of Okpara coal and Obwetti fireclay mines, near Enugu town, Nigeria. *Applied Water Science*, 3(1), 271-283.
- Valdes, D., Dupont, J. P., Laignel, B., Ogier, S., Leboulanger, T., & Mahler, B. J. (2007). A spatial analysis of structural controls on Karst groundwater geochemistry at a regional scale. *Journal of Hydrology*, 340(3), 244-255.
- Van der Gun, J. (2012). *Groundwater and global change: trends, opportunities and challenges*. UNESCO
- Van Ryneveld, M. B., & Fourie, A. B. (1997). A strategy for evaluating the environmental impact of on-site sanitation systems. *Water SA*, 23(4), 279-291.
- Varol, S., & Davraz, A. (2014). Assessment of geochemistry and hydrogeochemical processes in groundwater of the Tefenni plain (Burdur/Turkey). *Environmental Earth Sciences*, 71(11), 4657-4673.
- Vasanthavigar, M., Srinivasamoorthy, K., & Prasanna, M. V. (2012a). Evaluation of groundwater suitability for domestic, irrigational, and industrial purposes: a case study from Thirumanimuttar river basin, Tamilnadu, India. *Environmental Monitoring and Assessment*, 184(1), 405-420
- Vasanthavigar, M., Srinivasamoorthy, K., Ganthi, R. R., Vijayaraghavan, K., & Sarma, V. S. (2012b). Characterisation and quality assessment of groundwater with a special emphasis on irrigation utility: Thirumanimuttar sub-basin, Tamil Nadu, India. *Arabian Journal of Geosciences*, 5(2), 245-258.
- Vasanthavigar, M., Srinivasamoorthy, K., Vijayaraghavan, K., Ganthi, R. R., Chidambaram, S., Anandhan, P., Manivannan, R. & Vasudevan, S. (2010). Application of water quality index for groundwater quality assessment: Thirumanimuttar sub-basin, Tamilnadu, India. *Environmental Monitoring and Assessment*, 171(1-4), 595-609.

- Vega, M., Pardo, R., Barrado, E., & Debán, L. (1998). Assessment of seasonal and polluting effects on the quality of river water by exploratory data analysis. *Water Research*, 32(12), 3581-3592
- Venkatramanan, S., Chung, S. Y., Kim, T. H., Prasanna, M. V., & Hamm, S. Y. (2015a). Assessment and distribution of metals contamination in groundwater: a case study of Busan city, Korea. *Water Quality, Exposure and Health*, 7(2), 219-225.
- Venkatramanan, S., Chung, S. Y., Ramkumar, T., Gnanachandrasamy, G., Vasudevan, S., & Lee, S. Y. (2015b). Application of GIS and hydrogeochemistry of groundwater pollution status of Nagapattinam district of Tamil Nadu, India. *Environmental Earth Sciences*, 73(8), 4429-4442.
- Voudouris, K. S., Lambrakis, N. J., Papatheothorou, G., & Daskalaki, P. (1997). An application of factor analysis for the study of the hydrogeological conditions in Plio-Pleistocene aquifers of NW Achaia (NW Peloponnesus, Greece). *Mathematical Geology*, 29(1), 43-59.
- Wakida, F. T., & Lerner, D. N. (2005). Non-agricultural sources of groundwater nitrate: a review and case study. *Water research*, 39(1), 3-16.
- Wen, X. H., Wu, Y. Q., & Wu, J. (2008). Hydrochemical characteristics of groundwater in the Zhangye Basin, Northwestern China. *Environmental Geology*, 55(8), 1713-1724.
- WHO (2004). *Guidelines for drinking-water quality* (Vol. 1). World Health Organization.
- WHO (2006). *Guidelines for Drinking-water Quality*. 3<sup>rd</sup> Edition. 1<sup>st</sup> Addendum to Vol. 1. WHO, Geneva.
- WHO (2011). *Guidelines for drinking water quality*, 4<sup>th</sup> Edition. World Health Organization, Geneva, 978-92-4-154815-1
- Wilcox, L. V. (1948). *The quality of water for irrigation use*. Washington, DC: U.S. Department of Agriculture.
- Wilcox, L. V. (1955). *Classification and use of irrigation waters*. USDA, Circular 969, Washington, DC
- Wilcox, L. V. (1995). Classification and use of irrigation waters (p. 19). Washington: US Department of Agriculture.
- Wu, E. M. Y., & Kuo, S. L. (2012). Applying a multivariate statistical analysis model to evaluate the water quality of a watershed. *Water Environment Research*, 84(12), 2075-2085.
- Xenidis, A., Papassiopi, N., & Komnitsas, K. (2003). Carbonate-rich mining tailings in Lavrion: risk assessment and proposed rehabilitation schemes. *Advances in Environmental Research*, 7(2), 479-494.
- Yanggen, D. A., & Born, S. M. (1990). Protecting groundwater quality by managing local land use. *Journal of Soil and Water Conservation*, 45(2), 207-210.
- Yidana, S. M., Banoeng-Yakubo, B., & Akabzaa, T. M. (2010). Analysis of groundwater quality using multivariate and spatial analyses in the Keta basin, Ghana. *Journal of African Earth Sciences*, 58(2), 220-234.
- Yongsi, H. N. (2011). Access and management of drinking water in developing cities: evidence from Yaoundé (Cameroon). *Research Journal of Environmental Sciences*, 5(2), 124.
- Zakaria, A. S. (1975). The Geomorphology of Kelantan Delta (Malaysia). *Catena*, 2, 337-349.

- Zhang, X., Qian, H., Chen, J., & Qiao, L. (2014). Assessment of groundwater chemistry and status in a heavily used semi-arid region with multivariate statistical analysis. *Water*, 6(8), 2212-2232.
- Zhou, F., Huang, G. H., Guo, H., Zhang, W., & Hao, Z. (2007). Spatio-temporal patterns and source apportionment of coastal water pollution in eastern Hong Kong. *Water Research*, 41(15), 3429-3439.
- Zhu, C., & Schwartz, F. W. (2011). Hydrogeochemical processes and controls on water quality and water management. *Elements*, 7(3), 169-174.
- Zuane, J. D. (1990). *Drinking water quality of standards and control*. New York: Van Nostr

