



UNIVERSITI PUTRA MALAYSIA

***EFFECT OF NORTHEAST MONSOON ON COPPER SPECIATION AT
PULAU PERHENTIAN, TERENGGANU, MALAYSIA***

EDRIANNA GODON

FPAS 2018 15



**EFFECT OF NORTHEAST MONSOON ON COPPER SPECIATION AT
PULAU PERHENTIAN, TERENGGANU, MALAYSIA**

By

EDRIANNA GODON

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

December 2017

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**EFFECT OF NORTHEAST MONSOON ON COPPER SPECIATION AT
PULAU PERHENTIAN, TERENGGANU, MALAYSIA**

By

EDRIANNA GODON

December 2017

Chair : Khairul Nizam bin Mohamed @ Mohd Ramli,, PhD
Faculty : Environmental Studies

Copper (Cu) speciation was studied in the seawater at Pulau Perhentian, Terengganu, Malaysia in two different seasons, the pre- and post-Northeast monsoon. The existence of organic Cu (II) complexing ligands were determined by the application of CLE-AdCSV to determine Cu bioavailability and toxicity in the area. The existence of organic ligands (CuL) in the area helps to keeps the solubility of Cu and maintains its concentration under the toxicity level. The development method of CLE-AdCSV for Cu(II) speciation was optimized by analysing the seawater samples collected from Pulau Perhentian. Multiple series of sampling were carried out in pre- and post-monsoon season to identify the possible effect of the Northeast monsoon to the Cu(II) speciation in the area. The profile of seawater samples was analyzed from selected stations at Pulau Perhentian, Terengganu for Cu(II) speciation determination in order to determine the concentration level of Cu(II) in dissolved seawater. The analysis showed that >99.5% of total dissolved Cu (dCu) was bound to Cu (II) complexing ligands in this area. The ratio of CuL/dCu was analysed to see the ligand distributions, which showed their saturation rate with dissolved Cu. For both Nov 2015 and 2016, the CuL/dCu ratio showed low ratio (1.00-1.90, 1.00-1.20) in open ocean (St. 1 and St. 2), and varied ratios were observed in near-shore stations at St. 3, St. 4 and St. 5 (1.10-4.30). Meanwhile for both April 2016 and 2017, the ratios were in low ranges (1.02-2.31, 1.00-1.60), especially in April 2017. This finding indicated a saturated Cu-ligand complexation in the area. The concentration of natural organic Cu(II)-ligand binding (CuL) and its conditional stability constant ($\text{Log } K'_{\text{CuL}}$) in each sample was determined by using salicylaldoxime (SA) as competitive ligand. Our present data indicated the presence of strong natural organic Cu(II) complexing ligands in our seawater samples. Its distribution was similar to the dissolved Cu (dCu) throughout the water column, where the concentrations increased from the surface (3 m) to the middle layer (15m), before decreasing at the bottom layer. The $\text{Log } K'_{\text{CuL}}$ was more than 12.00 throughout the water column and indicated the presence of strong natural organic ligands in our study area. Preliminary data for Cu

speciation in pre- and post-Northeast monsoon at Pulau Perhentian for two years (2015/2016 and 2016/2017) were presented here. All the results observed in saturation rate, complexation percentage, and the conditional stability constant, $\text{Log } K'_{\text{CuL}}$ and the pCu data suggested that the dissolved Cu in the study area are taken up by marine organisms, thus preventing the free Cu^{2+} ions to be produced. The complexing capacity of the species in the area were also highlighted in the results, which gave an indication of the ability of the organisms to absorb dissolved Cu and render it non-toxic, by complexing with dissolved Cu. In addition, the presence of dominant diatoms species might had also influence Cu bioavailability in the area. This initial study on Cu(II) speciation has suggested that more than 99% of dissolved Cu was complexed to the natural organic ligands which prevents the toxicity of Cu(II) to the marine organisms.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

KESAN MONSUN LAUT KEPADA KOMPLEKSASI KUPRUM DI PULAU PERHENTIAN, TERENGGANU

Oleh

EDRIANNA GODON

Disember 2017

Pengerusi : Khairul Nizam bin Mohamed @ Mohd Ramli, PhD
Fakulti : Pengajian Alam Sekitar

Kajian kompleksasi Kuprum (Cu) dijalankan di air laut Pulau Perhentian, Terengganu, Malaysia pada dua musim yang berlainan, iaitu sebelum dan selepas Monsun Barat Daya. Kehadiran ligan (Cu) organik dikaji melalui aplikasi CLE-AdCSV untuk mengukur bioavailabiliti dan toksisiti Cu di kawasan kajian. Ligan organik ini membantu mengekalkan kelarutan Cu dan seterusnya menghalang toksisiti Cu. Kaedah CLE-AdCSV kemudian dibangunkan dan dioptimumkan sesuai dengan keadaan air laut di tempat kajian. Kemudian beberapa siri sampling dijalankan untuk mengambil sampel air laut pada sebelum dan selepas monsun untuk mengkaji kesan monsun Barat Daya kepada kompleksasi Cu. Profil daripada sampel air laut telah dianalisis dari beberapa stesen di Pulau Perhentian, Terengganu untuk kajian kompleksasi Kuprum (II) untuk mengenalpasti tahap kepekatan Cu(II) didalam air laut. Hasil analisis menunjukkan bahawa >99.5% daripada keseluruhan Cu terlarut adalah terikat kepada ligan kompleks di kawasan tersebut. Nisbah CuL/dCu kemudian dianalisis untuk melihat taburan ligan dan kadar tepu diantara CuL dan dCu. Untuk kedua2 Nov 2015 dan 2016, nisbah CuL/dCu adalah rendah (1.00-1.90, 1.00-1.20) di kawasan laut terbuka (St.1 dan 2) dan bervariasi di kawasan berdekatan pantai St. 3, 4 dan 5 (1.10-4.30). Sementara itu bagi April 2016 dan 2017, nisbah didapati dalam julat yang rendah (1.02-2.31, 1.00-1.60), terutama bagi April 2017. Ini menunjukkan bahawa kompleksasi ligan-Cu adalah tepu bagi kawasan berkenaan. Kepekatan ligan organik Kuprum (II) semulajadi (CuL) dan *conditional stability constant* ($\log K'_{CuL}$) di dalam setiap sampel diukur dengan menggunakan kaedah *competitive ligand exchange-adsorptive cathodic stripping voltammetry* (CLE-AdCSV) dengan ligan pesaing *salicylaldoxime*, (SA). Data terkini menunjukkan kehadiran ligan organik Kuprum (II) semulajadi di dalam semua sampel air laut. Taburannya adalah sama dengan Kuprum terlarut (dCu) disemua kolum air, dimana kepekatannya meningkat dari permukaan (3m) ke lapisan tengah (15m), sebelum menurun ke lapisan bawah (30m). Bacaan $\log K'_{CuL}$ adalah lebih daripada 12.00 disemua kolum air, memberi gambaran kehadiran ligan organik semulajadi yang kuat di

kawasan kajian ini. Kajian awal dalam analisis kompleksasi Kuprum(II) mencadangkan bahawa lebih daripada 99% Kuprum terlarut telah dikompleks dengan ligan semulajadi organik yang menghalang toksisiti Kuprum(II) kepada organisma marin. Data awal untuk kompleksasi Cu sebelum dan selepas Monsun Barat Daya di Pulau Perhentian untuk dua tahun, (2015/ 2016 dan 2016/2017) dibincangkan disini. Semua keputusan yang diperoleh dari kadar tepu, peratus kompleksasi dan *conditional stability constant*, $\text{Log } K'_{\text{CuL}}$ dan data pCu menunjukkan bahawa Kuprum terlarut di kawasan kajian digunakan oleh organisma marin, lalu menghalang pembentukan ion Cu^{2+} . Kapasiti kompleksasi oleh sepsis di kawasan tersebut juga dibincangkan, dimana hasil kajian memberi indikasi abiliti organisma untuk menyerap Kuprum terlarut dan menghalang toksisiti. Sebagai tambahan, kehadiran diatom sebagai spesis phytoplankton dominan juga mungkin mempengaruhi bioavailabiliti Kuprum di kawasan ini.

ACKNOWLEDGEMENTS

Philippians 4:13

My heartfelt gratitude to my outstanding, one and the only supervisor, Dr. Khairul Nizam Mohamed, for the countless hours of refining my works, especially during the last hours, pushing my limits and challenging me every day for better performance and self-improvement, for never getting upset whenever I made mistakes but kept on teaching me the right way of doing my research, for his endless support, guidance and valuable advices, but most importantly, for being a great teacher and friend throughout these semesters. This study wouldn't be possible without him and I was blessed to be his student.

I extend my appreciation to my supervisory committee, Dr Ferdius and Prof Suhaimi for their support and advices, for allocating their precious time to help me in improving my research works.

I'm eternally indebted to my dear family especially my parents, Mr. Edwin Korok Godon and Mdm. Irene Rounah Asang for their love, prayers, constant encouragement and support. To Dolores, Clare, Jane, Imelda, Cornelea, Bryan, Veralyn, Brymen, Brywon, Brywin, Anne, Ivanka, all my nieces and nephews, I love you all so much. You are God's greatest gifts to me. I'm more than blessed to have all of you beside me. Your encouragement and conviction will always inspire me.

To my CSSUPM family, the Legend Tepung group, all those strong ladies who had my back when I was in need of shelter and comfort; Clodenna, Cassie, Cynthia, Nazerita, Pris, Josie, to Clarence and Alex for their help, thank you for everything that you guys did for me. Hugs!

To my lab mates in Makmal Penyelidikan Kimia JSAS from 2015 to 2017, especially Adillah Surimin, thank you for all the experiences and laughter that we shared, they will forever be treasured. Special acknowledgment is also sent for all lab assistants, who had generously helped me throughout my study.

Last but certainly not the least; I am most grateful to the love of my life, Jcrechi Japial, my flashlight. Through all thick and thin, he was there with me. I can never thank him enough for every sacrifice that he had done all this time.

I certify that a Thesis Examination Committee has met on 12 December 2017 to conduct the final examination of Edrianna Godon on her thesis entitled "Effects of Northeast Monsoon on Copper Speciation at Pulau Perhentian, Terengganu, Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Normala binti Halimoon, PhD
Senior Lecturer
Faculty of Environmental Studies
Universiti Putra Malaysia
(Chairman)

Yusran bin Sulaiman, PhD
Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Che Abd Rahim Mohamed, PhD
Professor
Universiti Kebangsaan Malaysia
Malaysia
(External Examiner)



RUSLI HAJI ABDULLAH, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 30 July 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Khairul Nizam bin Mohamed @ Mohd Ramli, PhD

Senior Lecturer
Faculty of Environmental Studies
Universiti Putra Malaysia
(Chairman)

Ferdius @ Ferdaus binti Mohamat Yusuff, PhD

Senior Lecturer
Faculty of Environmental Studies
Universiti Putra Malaysia
(Member)

Suhaimi Bin Suratman, PhD

Associate Professor
Institute of Oceanography and Environment (INOS)
Universiti Malaysia Terengganu
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: _____

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of Chairman of
Supervisory
Committee: _____

Signature: _____
Name of Member of
Supervisory
Committee: _____

Signature: _____
Name of Member of
Supervisory
Committee: _____

Signature: _____
Name of Member of
Supervisory
Committee: _____

TABLE OF CONTENTS

	Page	
ABSTRACT	i	
ABSTRAK	iii	
ACKNOWLEDGEMENTS	v	
APPROVAL	vi	
DECLARATION	viii	
LIST OF TABLES	xiii	
LIST OF FIGURES	xiv	
LIST OF ABBREVIATIONS	xviii	
CHAPTER		
1	INTRODUCTION	1
	1.1 Introduction	4
	1.2 Problem Statement	4
	1.3 Research Objectives	4
	1.4 Research Questions	5
	1.5 Scope of Study	5
	1.6 Significance of Study	5
	1.7 Thesis Organization	6
2	LITERATURE REVIEW	8
	2.1 Introduction	8
	2.2 The Development of Cu Speciation by CLE-AdCSV	8
	2.3 Cu Toxicity and Bioavailability in Seawater	11
	2.4 Production of Natural Ligands by Phytoplankton	13
	2.5 Complexation of Trace Metal with Organic Ligand	15
	2.6 The Monsoon Season in Southeast Asia	16
	2.7 Speciation Studies in Other Regions	16
	2.8 Current Speciation Study in Malaysia	17
3	MATERIALS AND METHODS / METHODOLOGY	19
	3.1 Cleaning Processes	19
	3.1.1 Teflon fluorinated ethylene propylene (FEP) (Nalgene)	19
	3.1.2 Low density polyethylene bottles (Nalgene)	19
	3.2 Sampling	19
	3.3 Determination of dissolved Fe, Al and Cu by using column pre-concentration and separation method (Chelex-100 resin exchange resin)	22

3.3.1	Column set up and purification procedure	22
3.3.2	Pre-concentration and separation procedure	22
3.4	Chemicals Preparation	23
3.4.1	Borate buffer (H_3BO_3)	23
3.4.2	2-(2-Thiazolylazo)-p-cresol (TAC) stock solution	23
3.4.3	Cu(II) and SA Stock Solution	23
3.4.4	Sample preparation	24
3.5	Determination of total dissolved copper concentration	25
3.6	Determination of natural organic Cu(II) binding ligands	26
3.7	Voltammetry procedure	27
3.8	Theory of Cu Speciation with Salicylaldoxime (SA)	28
3.9	Measurement of Total Copper, Cupric Ion and Ligand Concentration	28
4	DISTRIBUTION OF DISSOLVED Fe, Al and Cu AT PULAU PERHENTIAN, TERENGGANU	31
4.1	Introduction	31
4.2	Materials and methods	33
4.3	Results and Discussion	34
4.3.1	Surface In-situ Parameters	34
4.3.2	Salinity, Conductivity and Turbidity Profiles for Pre-and Post-Monsoon	36
4.3.3	Surface Distribution of Metals	40
4.3.4	Depth Profiles for Al, Fe and Cu	42
4.3.4.1	Fe Distribution	42
4.3.4.2	Al Distribution	43
4.3.4.3	Cu Distribution	44
4.4	Conclusion	46
4.5	Acknowledgement	46
5	DETERMINATION OF COPPER (II) COMPLEXATION IN SEAWATER AT PULAU PERHENTIAN, TERENGGANU	47
5.1	Introduction	47
5.2	Materials and Method	47
5.3	Results and Discussion	48
5.3.1	Determination of natural organic ligand	48
5.3.2	Distribution of Cu (II) speciation	51
5.4	Conclusion	56
5.5	Acknowledgment	56

6	EFFECTS OF NORTHEAST MONSOON ON COPPER (Cu) SPECIATION AND DISTRIBUTION AT PULAU PERHENTIAN, TERENGGANU	57
	6.1 Introduction	57
	6.2 Material and Method	58
	6.3 Results and Discussion	58
	6.3.1 Physicochemical Characteristics of the Water Column	59
	6.3.2 Distribution of natural organic Cu(II)- ligands binding (CuL)	63
	6.3.3 Possible source of natural organic ligands	72
	6.3.4 Toxicity and bioavailability level at Pulau Perhentian	75
	6.4 Conclusion	76
7	CONCLUSION	77
	REFERENCES	79
	APPENDICES	89
	BIODATA OF STUDENT	99
	LIST OF PUBLICATIONS	100

LIST OF TABLES

Table		Page
4.1	Statistical analysis showing the p -values and the r2 values for all dissolved metals against salinity in pre-and post-monsoon season.	37
5.1	Recovery data for titration method accuracy	51
5.2	Cu speciation (CuL) data at Perhentian Island, Terengganu in Nov 2015. Total dissolved Cu concentration ([dCu]), total Cu binding ligand ([CuL]) and stability constant of Cu ligand (log K) were determined, and free Cu binding-ligand([L'] = [CuL] – [dCu]), $\alpha_{\text{Cu organic}} = [L'] \times K$, $[p\text{Cu}] = -\log \{d\text{Cu} / (\alpha_{\text{FCu organic}} - \alpha_{\text{Cu inorganic}})\}$ were calculated.	54
5.3	Comparison of Cu(II) speciation at Perhentian Island during this study and other coastal waters studies.	55
6.4	Cu speciation data at Pulau Perhentian, Terengganu in Nov 2015 (Pre-monsoon) and April 2016 (Post-monsoon). Dissolved Cu concentration ([dCu]), total Cu binding ligand ([CuL]) and stability constant of Cu ligand (log K) were determined, and free Cu binding-ligand([L'] = [CuL] – [dCu]) are presented.	66
6.5	Cu speciation data at Pulau Perhentian, Terengganu in Nov 2016 (Pre-monsoon) and April 2017 (Post-monsoon). Dissolved Cu concentration ([dCu]), total Cu binding ligand ([CuL]) and stability constant of Cu ligand (log K) were determined, and also the free Cu binding-ligand ([L'] = [CuL] – [dCu]).	67
6.6	Cu speciation and comparison at Pulau Perhentian and other coastal areas.	72

LIST OF FIGURES

Figure		Page
1.1	Sizes for various fractions in natural water sample and the typical constituents of the fractions. Typical filtrations use either a 0.2 or 0.45 μ m pore size	1
1.2	Schematic showing the main sources of particulate and dissolved trace metals in seawater are dust, rivers and shelf sediments, whereas hydrothermal inputs are significantly less. After introduction into sea water, the metals undergo biogeochemical cycling, which causes fractionation and vertical transport from surface to deep waters. Additionally, the metals are transported and distributed by surface and deep-water circulation	2
1.3	Southeast Asia region diagram	3
2.1	Chemical structures of a selected AdCSV ligands. The AdCSV measurement approach using the ligands involves: (a) reduction of the metal; (b) catalytic hydrogen formation; (c) reduction of ligand	9
2.2	The crystal structure of the parent compound salicylaldehyde, SA	10
2.3	The voltammetry instrument used in speciation analysis.	11
2.4	The Biotic Ligand Model schematic diagram explaining the processes of Cu-complexation in the marine environment	13
2.5	Adapted schematic diagram from Morel and Price (2003) shows seven examples of ligands productions by marine plankton; [1] CuY: Cu-peptide complexes released by coccolithophorids (Francis and Tebo, 2001); [2] CuZ: unidentified Cu ligand complex released by <i>Synechococcus</i> (Martin et al., 1994; Tebo et al., 1997); [3] L, <i>Prochlorococcus</i> released an unidentified Co complexing agent (Boyd, 2000); [4] heterotrophic marine bacteria oxidize Mn(II), forming a MnO ₂ casing around the cell (Lane and Morel, 2000); [5] During Fe uptake, diatoms extracellularly reduce Fe(III) ligand complexes (Berman-Frank et al., 2001; Kustka et al., in press); [6] sid, siderophore released by cyanobacteria and heterotrophic bacteria (Lee et al., 1996); [7] CdX: phytochelatin-Cd complex released by diatoms (Herbik et al., 2002).	14

3.1	Location of our sampling stations at Pulau Perhentian, Terengganu, Peninsular Malaysia during sampling activity. Each station was marked with the GPS coordinate indicated as S1 to S5.	20
3.2	Flow chart of research methodology	21
3.3	The diagram of sample preparation	25
3.4	The current (nA) plotted versus the total amount of Cu (nM) from St. 5 (6m depth). The first few points (black dots) of the titration indicated that about half of the natural organic complex dissociated due to the competition with the added ligand (SA). The concentration of added ligand and Cu(II) complex is directly related to the peak heights (ip) of the voltammetric measurements	27
4.1	Surface in-situ parameters for all stations during October 2014 and April 2015	35
4.2	Salinity profiles for both pre-and post-monsoon seasons.	37
4.3	Conductivity profiles for both pre-and post-monsoon seasons.	38
4.4	Turbidity profiles for both pre-and post-monsoon seasons.	39
4.5	Surface (3 m depth) distribution of dissolved Fe, Al and Cu during Oct 2014 and April 2015 at Pulau Perhentian, Terengganu.	39
4.6	Surface (3 m depth) distribution of dissolved Fe during Oct 2014 and April 2015 at Pulau Perhentian, Terengganu.	40
4.7	Surface (3 m depth) distribution of dissolved Al during Oct 2014 and April 2015 at Pulau Perhentian, Terengganu	41
4.8	Surface (3 m depth) distribution of dissolved Cu during Oct 2014 and April 2015 at Pulau Perhentian, Terengganu	41
4.9	Vertical distribution of Fe in October 2014 and April 2015	42
4.10	Vertical distribution of Al in October 2014 and April 2015	43
4.11	Vertical distribution of Cu in October 2014 and April 2015	44

5.1	(A) Voltammogram of Cu (SA) peak under differential pulse mode from seawater sample from St. 2 (6 m depth) at Pulau Perhentian, Terengganu. Voltammetric parameters: deposition time 120s; deposition potential -1.1 V and start potential -0.15 V. (B) The current (nA) plotted versus the total amount of Cu (nM) from St. 5 (6 m depth). The first few points (black square) of the titration indicated that about half of the natural organic complex dissociated due to the competition with the added ligand (SA). The concentration of added ligand and Cu (II) complex is directly related to the peak heights (i_p) of the voltammetric measurements.	49
5.2	(A) Comparison of UV irradiated (straight line) and non-UV (curved) seawater sample from St.2 at Pulau Perhentian, Terengganu. (B) Linearization of the titration data from St. 5 (6 m depth) by using van den Berg linearization formula to determine the complexing ligand concentrations and conditional stability constant.	50
5.3	The concentration of copper complexing ligands (CuL) and dissolved copper (dCu) at all stations at Pulau Perhentian, Terengganu. The CSV measurements were carried out using ligand competition with SA of 25 μ M.	52
6.1	The salinity profile at our stations during pre-and post-monsoon seasons.	60
6.2	pH profile at our stations during pre-and post-monsoon seasons.	61
6.3	Conductivity profile at our stations during pre-and post-monsoon seasons.	62
6.4	Dissolved oxygen profile at our stations during pre-and post-monsoon seasons.	63
6.5	The concentration of copper complexing ligands (CuL) and dissolved copper (dCu) at all stations at Pulau Perhentian, Terengganu during pre-monsoon (Nov 2015) and post-monsoon (April 2016).	68
6.6	The concentration of copper complexing ligands (CuL) and dissolved copper (dCu) at all stations at Perhentian Island, Terengganu on pre-monsoon (Nov 2016) and post-monsoon (April 2017).	70
6.7	The time series graph of monthly Chlorophyll- α values from January 2012 – December 2014	73

LIST OF APPENDICES

Appendix		Page
A	The raw data generated from the CLE-AdCSV by using the voltammetry, collected in the Excel spreadsheet.	89
B	The formulae and calculations by using Van den Berg/Ruzic Linearization	90
C	The linearized data obtained from Van den Berg/Ruzic formulae.	91
D	Regression analysis for pre- and post-monsoon season with dissolved Fe	92
E	The regression analysis for effect of Northeast monsoon season to Cu-ligand concentration	93
F	The in-situ parameters data during our sampling during Oct 2014 (Pre) and Apr 2015 (Post) at our stations at Pulau Perhentian, Terengganu. All the readings are from mean calculations.	94
G	Concentration of dissolved metals in Oct 2014 (Pre) and April 2015 (Post)	95
H	In-situ parameter data taken on pre (Oct 2014) and post monsoon (April 2015) season.	96
I	In-situ parameter data taken for pre (Nov 2015) and post-monsoon season (April 2016).	97
J	In-situ parameter data taken for pre (Nov 2016) and post-monsoon season (April 2017).	98

LIST OF ABBREVIATIONS

CLE-AdCSV	Competitive Ligand Exchange–Adsorptive Cathodic Stripping Voltammetry
K_{CuL}	Conditional Stability Constant
SA	Salicylaldoxime
CuL	Cu-ligand complex
dCu	Dissolved Cu
IUPAC	International Union of Pure and Applied Chemistry
GC-ICP-MS	Gas Chromatography - Inductively Coupled Plasma Mass Spectrometry
ASV	Anodic Stripping Voltammetry
AdCSV	Adsorptive Cathodic Stripping Voltammetry
LDPE	Low Density Polyethylene
PTFE	Polytetrafluoroethylene
NEM	Northeast Monsoon
SWM	Southwest Monsoon
DOC	Dissolved Organic Carbon
DOM	Dissolved Organic Matter
SPM	Suspended Particulate Matter
FEP	Fluorinated Ethylene Propylene
CSC	Conditional Stability Constant

CHAPTER 1

INTRODUCTION

1.1 Introduction

Some trace metals such as copper (Cu), iron (Fe), zinc (Zn), manganese (Mn) are vital for the organisms' growth by behaving as micronutrients for primary producers. In marine environment, usually these trace metals are found in low concentration (picomolar to nanomolar), where their roles are important in ocean biogeochemistry (Sunda, 2012). Marine organisms such as phytoplankton, macro algae bacterioplankton and fungi also require the micronutrients for many enzymatic reactions (Vraspir and Butler, 2009).

Basic physical differentiation of trace metals is either in particulate or dissolved forms, but they can also be found in other chemical and physical forms. The widely used definition to differentiate between both is by using the 0.4 μm or 0.2 μm filter size (Fig 1.1). The particulate forms of the metals involve those which are soaked to particles surfaces, absorbed in the biogenic origin particles or in the meld of minerals. Meanwhile, dissolved forms are those of potentially colloids or soluble trace metals fixation. Colloids, both organic and inorganic in nature, are produced abiotically or biotically and involve small organisms such as bacteria and viruses, and are frequently defined as the >0.02 and <0.4 μm fraction (Aiken *et al.*, 2011).




Figure 1.1: Sizes for various fractions in natural water sample and the typical constituents of the fractions. Typical filtrations use either a 0.2 or 0.45 μm pore size.

(Source: Aiken *et al.* (2011) *Environmental Science & Technology* 45: 3196-201, Copyright (2011), American Chemical Society).

Ocean biogeochemical cycle generally includes the transformations that occur in a substance as it goes through Earth's lithosphere, biosphere, hydro-sphere, and atmosphere (Raven *et al.*, 2006). The flux and subsequent rate, as well as the composition of trace metals are important to understand their biogeochemistry in the ocean. Atmospheric and riverine inputs are known as the two major sources of trace metals in the surface ocean, as shown in Fig. 1.2 (Donat and Bruland, 1994). Industrial and domestic waste, urban runoff and sewerage discharges are also possible to introduce the trace metals in aquatic environment but usually point sources are the ones that lead to increased concentrations (Whitby *et al.*, 2017).




Figure 1.2: Diagram of dissolved and particulate trace metals origins in seawater which include rivers, shelf sediment and dust, but less impact from hydrothermal inputs. After the trace metals reach the sea water, biogeochemical cycling occurs, and causes vertical transport and fractionation from surface to bottom waters. In addition, the metals are distributed or transported by the water circulation (Frank, 2011).

The first-hand input of this metals to open ocean is usually affected by deposition of atmospheric dust and long-range transport. While volcanic activity and continental dust serve as the natural sources, waste incineration and fossil fuel burning are mostly the anthropogenic sources (Chester and Jickells, 2012). The movement of dust particles in the atmosphere is significantly dependent on the condition of the grassland, forests, lands or desert in an area, which in turn are affected by climate, weathering process or geological aspects (Bruland and Lohan, 2003)

The Southeast Asian countries (Fig. 1.3), are governed by monsoon season, which is defined as a 'huge-scale seasonal turn-around of wind order' (Serreze and Barry, 2010). These areas greet maximums summer rainfall and nearly twofold rainfall maximums. The season does not only affect Asian countries, but also goes past the tropical latitudes and can also impact other non-monsoonal regions (Serreze and Barry, 2010).

The two-major monsoon order are distinctly named the southwest monsoon (SWM) or the summer monsoon, and northeast monsoon (NEM) or the winter monsoon. NEM starts from November to March, and SWM starts from late May to September. October is the changing month from SWM to NEM seasons (Cruz *et al.*, 2012).




Figure 1.3: Southeast Asia region (Kripalani and Kulkarni, 1997).

In Peninsular Malaysia, the NEM maintains the easterly or north-easterly winds of 10 to 20 knots but the winds may achieve up to 30 knots or more during periods of strong cold surges. Light and inconsistent winds are observed during the two inter-monsoon seasons but the monsoon carries winds that affect the vertical and upwelling process in the seawater because of the movement from the wind stress curl between air and the seawater surface (Adiana *et al.*, 2014; Mohamed and Amil, 2015). Previous study by Adiana *et al.* (2014) only highlighted the impact of Northeast monsoon on metals distribution in sediment, but not in dissolved or complexed phase. This condition triggers the need to conduct the study to see the relationship between the monsoon season to the trace metal distribution and speciation processes, as well as the impact on the biogeochemistry cycling in the water column, especially Cu, due to its toxicity and bioavailability functions.

1.2 Problem Statement

A number of heavy metals data in the South China Sea (SCS) had already been published elsewhere, but the database lists at www.elsevier.com/scopus shows that from 1968 to 2012, approximately only ten percent of studies on heavy metals have been discussed for southern part of SCS (Adiana *et al.*, 2014).

More importantly, most of the studies focus on marine organisms and sediments (Shazili *et al.* 2006) and lack of analysis was done on particulate and dissolved metals in seawater because of their trace concentrations (Jiménez *et al.* 2002). Unfortunately, speciation data has not been available in the country yet, due to the lack of research in this area. Reactivity and bioavailability of metals are determined by its speciation process and affect their biogeochemical cycling in the marine environment, hence speciation analysis is necessary in order to understand its reactivity in the environment such as its toxicity (for Cu) and biological availability.

Pulau Perhentian's status as a Marine Park in the area highlights the needs to conduct the study as an indication of its ecosystem health. This area is affected by the Northeast monsoon season, which carry wind over about maximum of 30 knots and affects the vertical and upwelling activities thus lifting nutrient-rich water to the ocean surface water (Mohamed and Amil, 2015). Generally, trace metals speciation in the upper layer is governed by the complexation of strong organic ligands (Bruland *et al.*, 1991), produced by phytoplankton in the water column (Croot *et al.*, 2000). Since speciation is controlled by the natural ligands produced by phytoplankton, an analysis is required to determine the effect of monsoon season to metal speciation at the study area.

1.3 Research Objectives

The overall aim of this research is to study the effect of Northeast monsoon on bioavailability of Cu to understand the effect of seasonal monsoon in the biogeochemical cycle of Cu at Pulau Perhentian, Malaysia.

The definite objectives are:

1. To study the distribution of total dissolved Cu at Pulau Perhentian, Malaysia in order to identify the influence of Northeast monsoon.
2. To determine Cu(II) speciation at Pulau Perhentian, Malaysia to by modifying the existing method to measure the Cu-ligand complex and the conditional stability constant in the seawater.
3. To quantify the degree of Cu(II) speciation to understand its toxicity threats by measuring the amount of free cupric ions, Cu^{2+} , saturation rate and its conditional stability constant.

1.4 Research Questions

Based on the research objectives above, the underlying research questions are as follows:

1. What are the role or effect(s), if any, of the Northeast monsoon on total dissolved Cu distribution at Pulau Perhentian?
2. Is there any presence of natural Cu(II) organic ligands in dissolved phase at Pulau Perhentian?
3. What are the degree of Cu(II) speciation and its toxicity threats, if any, during pre-and post-monsoon season at Pulau Perhentian?

1.5 Scope of study

This study generally covers the area as follows:

- 1- This study focuses on Cu speciation at Pulau Perhentian during pre-and post-monsoon season by using the current method, competing ligand exchange-adsorptive cathodic stripping voltammetry (CLE-AdCSV) as an initial study of its kind in the area. CLE-AdCSV is a very sensitive voltammetric technique that is accurate for detection limit of the order 10^{-11} M for dissolved copper concentration in water samples. Data collection was taken both during pre-and post-monsoon to compare the results obtained in lab analysis. The influencing parameters include the dissolved Cu concentration, total Cu concentration, natural ligands concentration, as well as the conditional stability constant, $\log K$, to quantify the free cupric ions concentration, Cu^{2+} in the water sample. The stability constant was influenced by the chemical properties of both metal ions and organic ligands. The concentration of organic ligand in seawater was among important parameter to know the ecological roles of these ligands and their possible origins.
- 2- This study also involves the identification of toxicity threats (if any), posed by free Cu^{2+} ions in the water column as a result of weak speciation process. Toxicity levels are determined by the pCu activity and the free Cu^{2+} concentrations in the sample, in which are quantified by using van den Berg linearization formula, after the lab analysis using CLE-AdCSV.

1.6 Significance of study

Copper speciation analysis is necessary to understand its reactivity in marine environment, for example its toxicity and bioavailability, as well as the geochemical

behavior of its species. Diatoms are among vital phytoplankton responsible for most marine photosynthesis, thus playing significant role in ocean carbon cycling.

As one of the essential nutrients for biological functions, concentration and speciation, trace metal Cu may directly influence the distribution of phytoplankton species in the ocean. Utilization and biological uptake of this element is related to its chemical interactions, which include interactions with the phytoplankton themselves. Interestingly, any changes to carbonate, hydrogen and hydroxide ions concentrations in carbon pump process may directly affect the speciation of metal complexes with dissolved organic matter (Stockdale *et al.*, 2016).

As a Marine Park itself, the health of the ecosystem at Pulau Perhentian requires a close monitoring of its trace metals' concentrations, especially Cu, as it can affect both in bioavailability and toxicity to microorganisms in the area. Previously, a limited condition of phytoplankton growth during pre-northeast monsoon was reported in a study (Mohamed and Amil, 2015). This raises a question whether the phytoplankton can produce strong natural organic ligands during the season, and whether the concentration of natural ligands are sufficient for Cu complexation, thus eliminating the toxicity threat it brings.

By using the current technology, this study hopes to provide a preliminary data for future comparisons of Cu distribution and speciation in seawater for pre-and post-NEM in southern SCS.

1.7 Thesis organization

Following this chapter, the remaining chapters of the thesis are outlined below:

Chapter 2 (Literature Review) details on the topics covered throughout the study that consists of the fundamentals background of Cu toxicity and bioavailability in seawater, the history of Cu speciation by other methods and by CLE-AdCSV, the production of natural ligands by phytoplankton, and Cu(II) speciation with natural ligands.

Chapter 3 (Methodology) presents the research design for this study in detail. The laboratory analysis in this study consist of sample treatment and preparation, as well as using CLE-AdCSV. The data collected experiments were analyzed using van den Berg linearization formula after the analysis.

Chapter 4, 5 and 6 (Results and Discussion) discuss the findings of the distribution of dissolved Fe, Al and Cu; the determination of Cu speciation by using CLE-AdCSV; and finally, the effect of Northeast monsoon on Cu speciation at Perhentian Island, Terengganu. All the findings are presented in profile graphs for easy comparisons and

discussions. This chapter also highlights the answers for all the research questions in this study.

Chapter 7 (Conclusions) summarizes the major findings of the study in the fulfilment of the research objectives. Recommendations for future studies in this field are also included.



REFERENCES

- Abualhaija, M. and Van den Berg, C.M.G. (2014). Chemical speciation of iron in seawater using catalytic cathodic stripping voltammetry with ligand competition against salicylaldehyde. *Mar. Chem.* 164, 60–74.
- Achterberg, E.P., Holland, T.W., Bowie, A.R., Mantoura, R.F.C. and Wordfold, P.J. (2001). Determination of iron in seawater. *Anal Chim Acta* 442, 1–14.
- Adiana, G., Shazili, N. A. M., Marinah, M. A. and Bidai, J. (2014). Effects of northeast monsoon on trace metal distribution in the South China Sea Off Peninsular Malaysia, *Environ Monit Assess* 186:421–431.
- Adiana, G., Shazili, N.A.M., Bidai, J. and Shaari, H. (2016). Special Issue Number 1: The International Seminar on The Straits of Malacca and the South China Sea: *J. Sustain. Sci. Manage* 93-106.
- Agusa, T., Kunito, T., Yasunaga, G., Iwata, H., Subramanian, A., Ismail, A. and Tanabe, S. (2005). Concentrations of trace elements in marine fish and its risk assessment in Malaysia. *Mar. Pol. Bul.* 51: 896–911.
- Ahner, B.A., Kong, S. and Morel, F.M.M. (1995). Phytochelatin production in marine algae. 1. An interspecies comparison. *Limnol. Oceanogr* 40, 649–657
- Aiken, G. R., Hsu-Kim, H. and Ryan, J. N. (2011) *Environmental Science & Technology* 45: 3196-201.
- APHA. (2005). Standard methods for the examination of water and wastewater, 21st edn. American Public Health Association, Washington, DC
- APHA. (1992). Standard Methods for the Examination of Water and Wastewater. 18th Edition, American Public Health Association (APHA), American Water Works Association (AWWA) and Water Pollution Control Federation (WPCF), Washington DC.
- Averill, B. A. and Eldredge, P. (2016). Solubility and Complexation Equilibria. Retrieved from website URL [https://chem.libretexts.org/Textbook_Maps/General_Chemistry_Textbook_Maps/Map%3A_Chemistry_\(Averill_and_Eldredge\)/17%3A_Solubility_and_Complexation_Equilibria](https://chem.libretexts.org/Textbook_Maps/General_Chemistry_Textbook_Maps/Map%3A_Chemistry_(Averill_and_Eldredge)/17%3A_Solubility_and_Complexation_Equilibria)
- Azhar, N. I. and Mohamed, K. N. (2016). Determination of natural organic Fe (III) binding ligand at Pulau Perhentian. *In-house Seminar of the Chemical Oceanography Laboratory* (Issue 1, Vol IV) 12/2016, ISSN - 0127 – 9629.
- Baron, M., Arellano, J.B. and Gorge, J.L. (1995). Copper and photosystem II: a controversial relationship. *Physiol. Plant* 94, 147–180.
- Brand, L.E., Sunda, W.G. and Guillard, R.R.L. (1986). Reduction of marine phytoplankton reproduction rates by copper and cadmium. *J. Exp. Mar. Biol. Ecol.* 96, 225–250.

- Barber, R. T. and Ryther, J. H. (1969). Organic chelators: Factors affecting production in the Cromwell Current upwelling, *J. Exp. Mar. Biol. Ecol* 3, 191-199.
- Batley, G. E. and Florence, T. M. (1976). Determination of the Chemical Forms of Dissolved Cadmium, Lead and Copper in Seawater. *Mar. Chem* 4, 347.
- Bruland, K. W. (1989). Complexation of zinc by natural organic ligands in the central North Pacific. *Limnol. Oceanogr* 34, 269–285.
- Bruland, K. W. and Rue, E. L. (2001). *Analytical methods for determination of concentrations and speciation of iron*. In the Biogeochemistry of Iron in Seawater (eds. D. R. Turner and K. A. Hunter). Wiley, Chichester 255–289.
- Bruland, K.W., Donat, J.R. and Hutchins, D.A. (1991). Interactive influences of bioactive trace metals on biological production in oceanic waters. *Limnol. Oceanogr.* 36, 1555–1577.
- Buck, K.N. and Bruland, K.W. (2007). The physicochemical speciation of dissolved iron in the Bering Sea, Alaska. *Limnol. Oceanogr* 52 (5), 1800–1808.
- Boyd, P. W. and Ellwood, M. J. (2010). The biogeochemical cycle of iron in the ocean, *Nat. Geosci*; 3, 675–682.
- Campos, M. L. A. M. and Van den Berg C. M. G. (1994). Determination of copper complexation in sea water by cathodic stripping voltammetry and ligand competition with salicylaldoxime. *Anal. Chim. Acta*; 284, 481–496.
- Campbell, P.G.C. (1995). *Interactions between trace metals and aquatic organisms: a critique of the free-ion activity model*. In: Tessier, A., Turner, D.R. (Eds.), *Metal Speciation and Bioavailability in Aquatic Systems*. IUPAC Series on Analytical and Physical Chemistry of Environmental Systems. John Wiley & Sons, Chichester, 45–102.
- Chapman, C.S., Capodaglio, G., Turetta, C. and Van den Berg, C.M.G. (2009). Benthic fluxes of copper, complexing ligands and thiol compounds in shallow lagoon waters. *Mar Environ. Res* 67, 17–24
- Cruz, F.T., Narisma, T.G., Villafuerte II, M.Q., Cheng-Chua, K.U. and Olaguera, L.M. (2012). A climatological analysis of the southwest monsoon rainfall in the Philippines. *Atmospheric Research* 122, 609-616.
- Charriau, A., Ludovic, L., Gao, Y., Leermakers, M., Baeyens, W., Ouddane, B. and Billon, G. (2011). Trace metal behaviour in riverine sediments: Role of organic matter and sulphides. *Applied Geochemistry* 26, 80–90.
- Coale, K. H. and Bruland, K. W. (1990). Spatial and temporal variability in copper complexation in the North Pacific. *Deep-Sea Res* 37, 317–33.
- Coale, K. H. and Bruland, K. W. (1988). Copper complexation in the northeast Pacific. *Limnol. Oceanogr* 33, 1084–1101.

- Coale, K.H., Johnson, K.S., Fitzwater, S.E., Gordon, R.M., Tanner, S., Chavez, F.P., Ferioli, L., Sakamoto, C., Rogers, P., Millero, F., Steinberg, P. and Nightingale, P. (1996). A massive phytoplankton bloom induced by an ecosystem scale iron fertilization experiment in the equatorial Pacific Ocean. *Nature* 383, 495–501.
- Croot, P.L., Moffett, J.W. and Brand, L. (2000). Production of extracellular Cu complexing ligands by eucaryotic phytoplankton in response to Cu stress. *Limnol. Oceanogr* 45, 619–627.
- Croot, P.L. (2003). Seasonal cycle of copper speciation in Gullmar Fjord, Sweden. *Limnol. Oceanogr* 48(2), 764–776.
- Cuong, D.T. and Obbard, J.P. (2006) Metal speciation in coastal marine sediments from Singapore using a modified BCR-sequential extraction procedure, In *Applied Geochemistry*, Volume 21, Issue 8, 2006, Pages 1335-1346, ISSN 0883-2927.
- Clean Water Team (CWT) (2004). *Electrical conductivity/salinity Fact Sheet*, FS-3.1.3.0(EC). in: *The Clean Water Team Guidance Compendium for Watershed Monitoring and Assessment, Version 2.0*. Division of Water Quality, California State Water Resources Control Board (SWRCB), Sacramento, CA.
- Duinker, J.C. and Kramer, C.J.M. (1977). An experimental study on the speciation of dissolved zinc, cadmium, lead and copper in river Rhine and North Sea water by differential pulse anodic stripping voltametry. *Mar. Chem* 5: 207- 228.
- Donat, J.R. and van den Berg, C.M.G. (1992). A new cathodic stripping voltammetry method for determining organic complexation in seawater. *Mar. Chem* 38: 69-90.
- Donat, J.R., Statham, P.J. and Bruland, K.W. (1986). An evaluation of a C-18 solid phase extraction technique for isolating metal-organic complexes from central North Pacific Ocean waters. *Mar. Chem* 18: 85-99.
- Ellwood, M. J. and Van den Berg, C. M. G. (2001). Determination of organic complexation of cobalt in seawater by cathodic stripping voltammetry. *Mar. Chem* 75, 49–68.
- Fairbrother, A., R. Wenstel, K. Sappington, and W. Wood. (2007). Framework for metals risk assessment. *Ecotoxicology and Environmental Safety* 68: 145-227.
- Fein, J. B., Daughney, C. J., Yee, N. and Davis, T. (1997). A chemical equilibrium model for metal adsorption onto bacterial surfaces. *Geochim Cosmochim Acta*; 61, 3319.
- Francis, A. J. (1999). *Biomediation of Contaminated Soils*, Agronomy Monograph No. 37, USA; 239.
- Flegal, A.R., Rivera-Duarte, I., Ritson, P.I., Scelfo, G.M., Smith, G.J., Gordon, M.R. and Sandoz-Wilhelmy, S.A. (1996). *Metal contamination in San Francisco*

- Bay waters: historic perturbations, contemporary concentrations, and future considerations.* In: Hollibaugh, J.T. (Ed.), San Francisco Bay: The Ecosystem. Pacific Division of the American Association for the Advancement of Science, San Francisco, 173–188.
- Fortune, J. and Mauraud, N. (2015). *Effect of tide on water quality of Jones Creek, Darwin Harbour.* Report No. 02/2015D. Department of Land Resource Management, Aquatic Health Unit. Palmerston, NT.
- Gee, A.K. and Bruland, K.W. (2002). Tracing Ni, Cu, and Zn kinetics and equilibrium partitioning between dissolved and particulate phases in South San Francisco Bay, California, using stable isotopes and high resolution inductively coupled plasma mass spectrometry. *Geochim Cosmochim Acta* 66 (17), 3063–3083.
- Gerringa, L.J.A., Veldhuis, M.J.W., Timmermans, K.R., Sarthou, G. and de Baar, H.J.W. (2006). Co-variance of dissolved Fe-binding ligands with phytoplankton characteristics in the Canary Basin. *Mar. Chem* 102 (3–4), 276–290.
- Gledhill, M. and Van den Berg, C. M. G. (1994). Determination of complexation of iron(III) with natural organic complexing ligands in seawater using cathodic stripping voltammetry. *Mar. Chem* 47, 41.
- Gordon, A.S., Donat, J.R., Kango, R.A., Dyer, B.J. and Stuart, L.M. (2000). Dissolved copper-complexing ligands in cultures of marine bacteria and estuarine water. *Mar. Chem.* 70, 149–160.
- Gonzalez-Davila, M., Santana-Casiano, J. M., Perez-Pena, J. and Millero, F. J. (1995). *Binding of Cu(II) to the surface and exudates of the alga Dunaliella tertiolecta in seawater.* In Environmental Science & Technology 2nd Edn Vol. 29, 289-301.
- Graeme, E. B. (1989). *Trace Element Speciation Analytical Methods and Problems.* Published by CRC Press. ISBN 10: 0849347122.
- Helios-Rybicka, E. and Kyziol, J. (1991). Clays and clay minerals as the natural barriers for heavy metals in pollution mechanisms— Illustrated by Polish rivers and soils. *Mitt.österr. geol Ges* 83, 163–176.
- Hirose, K. (2007). Metal-organic matter interaction: Ecological roles of ligands in oceanic DOM. *Appl. Geochem* 22:1636-1645.
- Hirose, K., Dokiya, Y. and Sugimura, Y. (1982). Determination of conditional stability constants of organic copper and zinc complexes dissolved in seawater using ligand exchange method with EDTA. *Mar. Chem* 11, 343–354.
- Hirose, K. (1994). Conditional stability constants of organic metal complexes in seawater: Past and present, and a simple coordination chemistry model. *Anal. Chim. Acta* 284, 621–634.

- Huang, T.C., Lay, K.C. and Tong, S.R. (1991). Resetting the endogenous circadian N₂-fixing rhythm of the prokaryote *Synechococcus* RF-1. *Bot Bull Acad Sin* 32, 129-133.
- Irwandi, J. and Farida, O. (2009). Mineral and heavy metal contents of marine fin fish in Langkawi Island, Malaysia. *Int Food Res J* 16:105-112.
- IUPAC Chemical data series. (1979). *Stability constants of metal-ion complexes, Part B: Organic ligands*, Pergamon Press, Oxford 225 29.
- Jean, N., Dumont, E., Durrieu, G., Balliau, T., Jamet, J-L. and Personnic, S. (2012). Protein expression from zooplankton communities in a metal contaminated NW Mediterranean coastal ecosystem. *Mar. Environ. Res* 80, 12–26.
- Johnson, K.S., Gordon, R.M. and Coale, K.H. (1997). What controls dissolved iron concentrations in the world ocean? *Mar. Chem.* 57 (3–4), 137–161.
- Kristen, N.B., John, R.M., Rossb, A., Flegalc, R. and Bruland, K W. (2006). A review of total dissolved copper and its chemical speciation in San Francisco Bay, California, *Environmental Research* 105(1):5-19.
- Kraepiel, A. M. L., Chiffoleau, J. F., Martin, J. M. and Morel, F. M. M. (1997). Geochemistry of trace metals in the Gironde estuary. *Geochimica et Cosmochimica Acta* 61, 1421–1436.
- Kozelka, P.B. and Bruland, K. W. (1998). Chemical Speciation of Dissolved Cu, Zn, Cd, Pb in Narragansett Bay, Rhode Island. *Mar. Chem* 60 267.
- Kim, K. T., Kim, E. S., Cho, S. R., Park, J. K., Ra, K. T. and Lee, J. M. (2010). *Distribution of Heavy Metals in the Environmental Samples of the Saemangeum Coastal Area, Korea*. Coastal Environmental and Ecosystem Issues of the East China Sea, Eds., A. Ishimatsu and H.-J. Lie, 71–90.
- Leal, M.F.C., Vasconcelos, M.T.S.D. and Van den Berg, C.M.G. (1999). Copper-induced release of complexing ligands similar to thiols by *Emiliania huxleyi* in seawater cultures. *Limnol. Oceanogr.* 44, 1750–1762.
- Lewis, B. L., Holt, P. D., Taylor, S. W., Wilhelm, S. W., Trick, C. G., Butler, A. and Luther, G. W. III, Voltammetric estimation of iron (III) thermodynamic stability constant for catecholate siderophores isolated from marine bacteria and cyanobacteria. *Mar. Chem* (1995); 50, 179.
- Li, L., Liu, J., Wang, X. and Shi, X. (2015). Dissolved trace metals distributions and Cu speciation in the southern Bohai Sea China. *Mar Chem* 172, 34– 35.
- Li, L., Pala, F., Haskins, J., Sukola, K. and Wallace, G.T. (2011). Measurement of free Cu ion activity in seawater using a passive-equilibrium sonic- assisted free ion recorder (SAFIR). *Environ. Sci. Technol* 5, 5660–5667.
- Li, Y, and Mohamed, K. N. (2016). *Nutrients enrichment experiment on seawater samples at Pulau Perhentian, Terengganu*. Universiti Putra Malaysia.

- Li, L., Wang, X., Liu, J. and Shi, X. (2017). Dissolved trace metal (Cu, Cd, Co, Ni, Ag) distribution and Cu speciation in the Southern Yellow Sea and Bohai Sea, China. *Journal of Geophysical Research: Oceans* doi: 10.1002/2016JC012500
- Mahmood, A., Abualhaija, M. M., van den Berg, C. M. G., & Sander, S. G. (2015). Organic speciation of dissolved iron in estuarine and coastal waters at multiple analytical windows. *Mar Chem* 177, 706-719.
- Martin, J.H. and Gordon, R.M. (1988). Northeast Pacific iron distributions in relation to phytoplankton productivity. *Deep Sea Res. Part A*, 35, 177- 196.
- Markai, S., Montavon, G., Andres, Y. and Grambow, B. (2003). Complexation studies of Eu(III) with alumina-bound polymaleic acid: effect of organic polymer loading and metal ion concentration, *Appl. Radiat. Isot* 58, 16.
- Miller, L. A. and Bruland, K. W. (1994). Determination of copper speciation in marine waters by competitive ligand equilibration/liquid-liquid extraction: an evaluation of the technique. *Anal. Chim. Acta* 284, 573.
- Moffett, J.W. and Zika, R.G. (1987). Solvent extraction of copper acetylacetonate in studies of copper (II) speciation in seawater. *Mar. Chem* 301-313.
- Moffett, J. W., Brand, L. E. and Zika, R. G. (1990). Distribution and potential sources and sinks of copper chelators in the Sargasso Sea. *Deep-Sea Res* 37, 27–36.
- Moffett, J.W., Brand, L.E., Croot, P.L. and Barbeau, K.A. (1997). Cu speciation and cyanobacterial distribution in harbors subject to anthropogenic Cu inputs. *Limnol. Oceanogr* 42,789–799.
- Moffett, J. W., Brand, L. E. and Zika, R. G. (1990). Distribution and potential sources and sinks of copper chelators in the Sargasso Sea. *Deep-Sea Res* 37, 27–36.
- Moffett, J.W., Brand, L.E., Croot, P.L. and Barbeau, K.A. (1997). Cu speciation and cyanobacterial distribution in harbors subject to anthropogenic Cu inputs. *Limnol. Oceanogr.* 42,789–799.
- Moffett, J.W., Zika, R.G. and Brand, L.E. (1990). Distribution and potential sources and sinks of copper chelators in the Sargasso Sea. *Deep-Sea Res.* 37, 27–36.
- Moffett, J. W. and Brand, L. E. (1996). Production of strong, extracellular Cu chelators by marine cyanobacteria in response to Cu stress. *Limnol. Oceanogr* 41, 388–395.
- Mohamed, K. N., Steigenberger, S., Nielsdottir, M. C., Gledhill, M. and Achterberg, E. P. Dissolved iron(III) speciation in the high latitude North Atlantic Ocean. *Deep Sea Res Part I. Oceanographic Research Papers* (2011); 58, (11), 1049-1059.

- Nakajima, A., Horikoshi, T. and Sakaguchi, T. (1979). Studies on the accumulation of heavy metal elements in biological systems X. Uptake of copper ion by green microalgae. *Agric. Biol. Chem* 43, 1455.
- Ozaki, T., Kimura, T., Ohnuki, T. and Francis, A. J. (2005). Associations of Eu(III) with gram-negative bacteria *alcaligenesfaecalis*, *schwanella putrefaciens*, and *paracoccus* denitrificans. *J. Nucl. Radiochem. Sci* 6, 73.
- Panganelli, F., Papini, M. P., Toro, L., Trifoni, M. and Veglio F. (2000). Biosorption of metal ions on *Arthrobacter sp.*: biomass characterization and biosorption modelling. *Environ. Sci. Technol* 34, 2773.
- Posacka, A. M., Semeniuk, D. M., Whitby, H., van den Berg, C. M. G., Cullen, J. T., Orians, K., & Maldonado, M. T. (2017). Dissolved copper (dCu) biogeochemical cycling in the subarctic Northeast Pacific and a call for improving methodologies. *Marine Chemistry*, 196, 47-61.
- Peers, G. and Price, N.M. (2006). Copper-containing plastocyanin used for electron transport by an oceanic diatom. *Nature* 441 (7091), 341–344.
- Pribil, S. and Marvan, P. (1976). Algological Studies. *Arch. Hydrobiol. Suppl* 49, 214.
- Plavsic, M., Krznicaric, D. and Branica, M. (1982). Determination of the apparent copper complexing capacity of seawater by anodic stripping voltammetry. *Mar. Chem* 11, 17–31.
- Peason, R. G. (1963). Hard and soft acids and bases. *J. Am. Chem. Soc* 85, 3533.
- Plette, A. C. C., Beneditti, M. F. and Van Reimsdijk, W. H. (1996). Competitive binding of protons, calcium, cadmium, and zinc to isolated cell walls of a Gram-positive soil bacterium. *Environ. Sci. Technol* 30, 1902.
- Raven, P. H. and Linda R. B. (2006). *Environment*. Hoboken, NJ: John Wiley and Sons.
- Reid, R. T., Live, D. H., Faulkner, D. J. and Butler, A. (1993). A siderophore from a marine bacterium with an exceptional ferric ion affinity constant. *Nature* 366, 455.
- Rue, E. L. and Bruland, K. W. (1995). Complexation of iron (III) by natural organic ligands in the central North Pacific as determined by a new competitive ligand equilibration/adsorptive cathodic stripping voltammetry method. *Mar. Chem* 50, 117–138.
- Ruzic, I. (1982). Theoretical aspects of the direct titration of natural waters and its information yield for trace metal speciation. *Anal Chem. Acta* 140:99- 113.
- Sakaguchi, T. and Nakajima, A. (1991). *Mineral Bioprocessing*, ed. R. W. Smith and M. Misra, The Minerals, Metals & Materials Society; 309.
- Santos-Echeandia, J., Laglera, L.M., Prego, R. and Van den Berg, C.M.G. (2008). Dissolved copper speciation behaviour during estuarine mixing in the San Simon Inlet (wet season, Galicia). Influence of particulate matter. *Estuarine, Coastal and Shelf Science* 76, 447–453.

- Schreiber, E., Harshman, K., Kemler, I., Malipiero, U., Schaffner, W. and Fontana, A. (1990). Astrocytes and glioblastoma cells express novel octamer-DNA binding proteins distinct from the ubiquitous Oct-1 and B cell type Oct-2 proteins. *Nucleic Acids Research* 18 (18) : 5495-5503.
- Serreze, M.C. and Barry, R.G. (2010). *Climate change*. In: Barry, R.G., Chorley, R.J. (Eds.), *Atmosphere, Weather and Climate*. Routledge, Oxon.
- Sergio, C. (1996). *Element Speciation in Bioinorganic Chemistry*. Wiley-Interscience. ISBN, 0471576417.
- Stumm, W. and Morgan, J. J. (1996). *Chemical equilibria and rates in natural waters*. Aquatic Chemistry. Wiley Interscience, New York.
- Sugimura, Y., Suzuki, Y. and Miyake, Y. (1978). Chemical forms of minor metallic elements in the ocean. *J. Oceanogr. Soc. Jpn* 34, 93.
- Sunda, W. G. and Ferguson, R. L. (1983). *Trace Metals in Sea Water*. ed. C. S. Wong, E. Boyle, K. Bruland, J. Burton, and E. D. Goldberg, Plenum, New York 871.
- Sunda, W.G., Tester, P.A. and Huntsman, S.A. (1987). Effects of cupric and zinc ion activities on the survival and reproduction of marine copepods. *Mar Biol* 94, 203–210.
- Sunda, W.G. and Guillard, R.R.L. (1976). The relationship between cupric ion activity and the toxicity of copper to phytoplankton. *J. Mar. Res.* 34 (4), 511–529.
- Sunda, W.G. and Huntsman, S.A. (1983). Effect of competitive interactions between manganese and copper on cellular manganese and growth in estuarine and oceanic species of the diatom *Thalassiosira*. *Limnol. Oceanogr.* 28 (5), 924–934.
- Sunda, W.G. and Huntsman, S.A. (1998). Interactions among Cu^{2+} , Zn^{2+} , and Mn^{2+} in controlling cellular Mn, Zn, and growth rate in the coastal alga *Chlamydomonas*. *Limnol. Oceanogr.* 43 (6), 1055–1064.
- Sunda W. G. and Huntsman, S. A. (1991). The use of chemiluminescence and ligand competition with EDTA to measure copper concentration and speciation in seawater. *Mar. Chem* 36, 136.
- Sunda, W.G. (1994). *Trace metal/phytoplankton interactions in the sea*. In: Bidoglio, G., Stumm, W. (Eds.), *Chemistry of Aquatic Systems: Local and Global Perspectives*. Kluwer Academic, Dordrecht, 213–247.
- Sunda, W. G. (2012). *Feedback interactions between trace metal nutrients and phytoplankton in the ocean*. *Frontiers in Microbiology*.
- Sunda, W.G. and Huntsman, S.A. (1995). Regulation of copper concentration in the oceanic nutricline by phytoplankton uptake and regeneration cycles. *Limnol. Oceanogr* 40 (1), 132–137.

- Talley, L. D. (2002). *Salinity patterns*. Encyclopedia of Global Environmental Change, vol. 1, M. C. MacCracken and J. S. Perry, editors, John Wiley and Sons.
- Thuroczy, C.E., Gerringa, L.J.A., Klunder, M.B., Laan, P. and de Baar H.J.W. (2011). Observation of consistent trends in the organic complexation of dissolved iron in the Atlantic sector of the Southern Ocean. *Deep-Sea Research II*, 58 (2695–2706).
- Thuroczy, C. E., Alderkamp, A. C. P., Laan, L. J. A., Gerringa, M. M., Mills, G. L., Van, D., de Baar, H. J. W. and Arrigo, K.R. (2012). Key role of organic complexation of iron in sustaining phytoplankton blooms in the Pine Island and Amundsen Polynyas (Southern Ocean), *Deep Sea Res., Part II*, 71-76, 49–60.
- Sang, Y. W., Adam, M. K.M., Fakaruddin, F. J., Saleh, F. Z., Chang, N. K., Yunus, F. and Abdullah, M. H. (2016). *Review of The November 2015 – March 2016 Northeast Monsoon in Malaysia*. Malaysian Meteorological Department, MOSTI Research Publication 4/2016.
- Stryer, L. (1981). *Biochemistry*. H. Freeman, San Francisco
- U.S. EPA (2006). National recommended water quality criteria. Washington DC: Environmental Protection Agency (EPA). URL [Accessed: 2.05.2016].
- Van den Berg, C.M.G. and Nimmo M. (1987). Direct determination of uranium in water by cathodic stripping voltammetry. *Anal Chem* 59, 924.
- Van den Berg, C.M.G. (2006). Chemical speciation of iron in seawater by cathodic stripping voltammetry with dihydroxy naphthalene. *Anal. Chem* 78 (1), 156–163.
- Van den Berg, C.M.G. (1982). Determination of copper complexation with natural organic ligands in sea water by equilibration with MnO₂. I. Theory. *Mar. Chem* 11: 307-322.
- Van den Berg, C.M.G. (1984). Determination of the complexing capacity of copper(II) with natural organic ligands in seawater by cathodic stripping voltammetry of copper-catechol complex ions. *Mar. Chem* 15: 1-18.
- Van den Berg, C.M.G., Merks, A.G.A. and Duursma, E.K. (1987). Organic complexation and its control of the dissolved concentration of copper and zinc in Scheld Estuary. *Estuar. Coastal Shelf Sci* 24, 785.
- Van den Berg, C.M.G. (1984). Determination of the complexing capacity of copper(II) with natural organic ligands in seawater by cathodic stripping voltammetry of copper-catechol complex ions. *Mar. Chem* 15: 1-18.
- Van den Berg, C.M.G. (2006). Chemical speciation of iron in seawater by cathodic stripping voltammetry with dihydroxy naphthalene. *Anal. Chem* 78 (1), 156–163.

- Vraspir, J. M. and Butler, A. (2009) *Chemistry of Marine Ligands and Siderophores*. In *Annual Review of Marine Science*, 1. Annual Review of Marine Science. Annual Reviews, Palo Alto, 43-63.
- Wang, J. (1996). *Laboratory Techniques in Electroanalytical Chemistry*. In P.T. Kissinger, W.R. Heineman (Eds.), Marcel Dekker, New York 719–738.
- Wang, L., Li, J., Lu, H., Gu, Z., Rioual, P., Hao, Q., Mackay, A.W., Jiang, W., Cai, B., Xu, B. and Chu, G. (2012). The East Asian winter monsoon over the last 15,000 years: its links to high-latitudes and tropical climate systems and complex correlation to the summer monsoon. *Quaternary Science Reviews* 32, 131-142.
- Westerlund, S. and Öhman, P. (1991). Cadmium, copper, cobalt, nickel, lead, and zinc in the water column of the Weddell Sea, Antarctica. *Geochimica et Cosmochimica* 55, 2127–2146.
- Williams, P.M. (1969). The association of copper with dissolved organic matter in sea water. *Limnol. Oceanogr* 14:156.
- Whitby, H., Hollibaugh, J. T. and van den Berg, C. M. G. (2017). Chemical Speciation of Copper in a Salt Marsh Estuary and Bioavailability to Thaumarchaeota. *Frontiers in Marine Science*, 4.
- Wu, J. and Luther, G. W. III. (1995). Complexation of iron(III) by natural organic ligands in the Northwest Atlantic Ocean by a competitive ligand equilibration method and a kinetic approach. *Mar. Chem.* 50, 159–179.
- Yusoff, F. M., Zubaidah, M. S., Matias, H. B. and Kwan, T. S. (2002). Phytoplankton succession in intensive marine shrimp culture ponds treated with a commercial bacterial product. *Aquaculture Research*. 33(4):269–278.
- Yoshida, T., Ozaki, T., Ohnuki, T. and Francis, A. J. (2004). Adsorption of rare earth elements by γ -Al₂O₃ and *Pseudomonas fluorescens* cells in the presence of desferrioxamine B: Implication of siderophores for Ce anomaly. *Chem. Geol.*, 212, 239.
- Zhou, X. and Wangersky, P. J. (1989). Production of copper-complexing organic ligands by the marine diatom *Phaeodactylum tricorutum* in a cage culture turbidostat. *Mar. Chem.* 26: 239–259.