



UNIVERSITI PUTRA MALAYSIA

***PRODUCTION, PURIFICATION, AND CHARACTERIZATION OF
HALOTOLERANT LIPASE FROM *Leuconostoc mesenteroides* SUBSP.
mesenteroides ATCC 8293***

NURFADHILAH HIDAYAH EKO SUKOHIDAYAT

FSTM 2018 22



**PRODUCTION, PURIFICATION, AND CHARACTERIZATION OF
HALOTOLERANT LIPASE FROM *Leuconostoc mesenteroides* SUBSP.
mesenteroides ATCC 8293**

By

NURFADHILAH HIDAYAH EKO SUKOHIDAYAT

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

April 2018

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**PRODUCTION, PURIFICATION, AND CHARACTERIZATION OF
HALOTOLERANT LIPASE FROM *Leuconostoc mesenteroides* SUBSP.
mesenteroides ATCC 8293**

By

NURFADHILAH HIDAYAH EKO SUKOHIDAYAT

April 2018

Chairman : Associate Professor Mehrnoush Amid, PhD
Faculty : Food Science and Technology

Halotolerant lipases are essential in several industries, particularly food industry due to their ability to withstand different salt concentrations. Food industry usually involves food fermentation process, which requires the enzymes to be active and stable in the presence of high salt concentration in order to carry out the process efficiently. To date, only few halotolerant lipases have been discovered and characterized. Thus, the present work was conducted in order to produce, purify, and characterize halotolerant lipases from *L. mesenteroides* subsp. *mesenteroides* ATCC 8293.

Production of extracellular lipase from *L. mesenteroides* subsp. *mesenteroides* ATCC 8293 was optimized according to culture conditions (carbon sources, nitrogen sources, emulsifier, and surfactants) and physical parameters (temperature and pH) using the shake-flask fermentation system. It was observed that lipase production was optimized when the following culture composition was used: peptone, 0.2% (w/v), olive oil, 1.0% (v/v), monopotassium phosphate (KH_2PO_4) 0.5g, dipotassium phosphate (K_2HPO_4) 0.5g, sodium chloride (NaCl) 0.1g, calcium chloride (CaCl_2) 0.1g, magnesium sulfate heptahydrate ($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$) 0.5g, gum arabic, 0.1% (w/v) and Tween 80, 0.1% (v/v) per liter. The best physical parameters for maximum lipase production were at 30°C and pH 6.

Purification of lipases produced from *L. mesenteroides* subsp. *mesenteroides* ATCC 8293 was conducted using a novel aqueous two-phase system (ATPS) composed of Triton X-100 and maltitol. Although, the search for alternative components of ATPS has been widely investigated, little attention has been given to polyols as a component of a two-phase system. The ability of polyols to mimic the structure of water and

maintain an artificial sphere of hydration around the macromolecules makes them a potential candidate as a phase separating agent in ATPS. Initially, the phase diagram of this system was constructed using different concentrations of Triton X-100 and maltitol based on turbidimetric titration method. The partitioning of lipases was then optimized according to several parameters, which were pH, temperature, and crude load. It was observed that lipases preferentially migrated to the Triton X-100 rich phase. Optimum lipase partitioning was achieved in ATPS at tie-line length of 46.4%, crude load of 20% at a temperature of 30°C and pH 8 respectively. Purification of lipases using this system resulted in high lipase purification factor of 17.28 and lipase yield of 94.7%. The purified lipase showed a prominent band on SDS-PAGE with an estimated molecular weight of 50 kDa. Hence, this study demonstrated that Triton X-100 and maltitol could be potentially used as an alternative ATPS in order to efficiently purify valuable enzymes.

The purified lipases were then characterized on the basis of temperature, pH, and the presence of surfactants, metal ions as well as different salt concentrations. Optimum lipase activity was observed at 37°C and pH 8 respectively. The purified lipase was also observed to be stable at temperature range of 30-60°C and pH range of 6-11. Lipase exhibited enhanced activity in the presence of non-ionic surfactants, which were Triton X-100, Tween 80 and Tween 20, with increased activity up to 40%. Conversely, the presence of SDS, an anionic surfactant, inhibited 78% of the lipase activity. Metals ions such as Na⁺, Mg²⁺, K⁺ and Ca²⁺ stimulated lipase activity whereas divalent ions such as Zn⁺ and Cu²⁺ significantly reduced lipase activity by almost 50%. It was also observed that lipase activity was remarkably enhanced beyond 100% in the presence of different salt concentrations (0-10% w/v), thus confirming that lipase from *L. mesenteroides* subsp. *mesenteroides* ATCC 8293 is a halotolerant enzyme.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGHASILAN, PENULENAN DAN PENCIRIAN LIPASE
HALOTOLERAN DARIPADA *Leuconostoc mesenteroides* SUBSP.
mesenteroides ATCC 8293**

Oleh

NURFADHILAH HIDAYAH EKO SUKOHIDAYAT

April 2018

Pengerusi : Profesor Madya Mehrnoush Amid, PhD
Fakulti : Sains dan Teknologi Makanan

Lipase halotoleran adalah penting dalam beberapa industri, terutama industri makanan disebabkan kemampuannya untuk bertahan daripada kepekatan garam yang berbeza. Industri makanan lazimnya melibatkan proses penapaian makanan yang memerlukan enzim menjadi aktif dan stabil dengan kehadiran kepekatan garam yang tinggi untuk menjalankan proses dengan efisien. Sehingga kini, hanya sedikit lipase halotoleran yang telah ditemui dan dicirikan. Oleh itu, kajian ini dijalankan untuk menghasilkan, menulen dan mencirikan lipase halotoleran dari *L. mesenteroides* subsp. *mesenteroides* ATCC 8293.

Penghasilan lipase ekstrasel dari *L. mesenteroides* subsp. *mesenteroides* ATCC 8293 dioptimumkan mengikut keadaan kultur (sumber karbon, sumber nitrogen, pengemulsi, dan surfaktan) dan parameter fizikal (suhu dan pH) menggunakan sistem penapaian kelalang-goncang. Didapati bahawa penghasilan lipase adalah optimum apabila komposisi kultur berikut digunakan: pepton, 0.2% (w/v), minyak zaitun, 1.0% (v/v), monopotassium fosfat (KH_2PO_4) 0.5g, dipotassium fosfat (K_2HPO_4) 0.5g, natrium klorida (NaCl) 0.1g, kalsium klorida (CaCl_2) 0.1g, magnesium sulfat heptahidrat ($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$) 0.5g, gum arabik, 0.1% (w/v) dan Tween 80, 0.1% (v/v) seliter. Parameter fizikal terbaik untuk pengeluaran lipase yang maksimum adalah pada suhu 30°C dan pH 6.

Penulenan lipase yang dihasilkan dari *L. mesenteroides* subsp. *mesenteroides* ATCC 8293 telah dijalankan menggunakan sistem dua-fasa akueus (SDFA) yang terdiri daripada Triton X-100 dan maltitol. Walaupun pencarian komponen alternatif SDFA telah dikaji secara meluas, perhatian yang sedikit telah diberikan kepada poliol sebagai

ACKNOWLEDGEMENTS

All praises to Allah, the Most Merciful and Most Beneficent, for giving me strength, health and wisdom throughout my research journey.

I would like to express my deep gratitude to my project supervisor Dr. Mehrnoush Amid for giving me the opportunity to work with her and sharing her invaluable knowledge and passion in research. Her willingness to invest her time and efforts in guiding me in the course of my studies has been very much appreciated.

It is my pleasure to extend my appreciation to my co-supervisor, Associate Professor Badlishah Sham Baharin, for his continuous support and guidance. I am extremely thankful to Professor Dato Dr. Mohd Yazid Abdul Manap who took keen interest in the progress of my research and assisted me all along, until the completion of this project by providing all the necessary helps despite his busy schedule. I am privileged to be able to work under him.

I owe my deepest gratitude to my parents, Eko Sukohidayat and Norizan Alias for their encouragement and prayers throughout this challenging journey. Without each of them, I'd be nowhere near the person I am and the person I'm still working on becoming. I would not forget to remember my siblings Fikriyah, Nurul, and Farah for providing me with laughter and humor. Their humors helped me a lot in relieving some stress and lifted my spirit on my worse days.

Last but not least, I am grateful and fortunate to have Khaleeda, Aliaa, Aqilah, Atiqah, Muhaini, Fara and Farhana as my friends during my research journey. Their constant supports, bickering and failed jokes always make my days better.

I certify that a Thesis Examination Committee has met on..... to conduct the final examination of Nurfadhilah Hidayah Eko Sukohidayat on her thesis entitled “Production, purification, and characterization of halotolerant lipase from *Leuconostoc mesenteroides* subsp. *mesenteroides* ATCC 8293” in accordance with the Universities and University College Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U(A) 106] 15 March 1998. The committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Name of Chairperson,

Title (e.g., Professor/ Associate Professor/ Ir; omit if irrelevant)

Name of Faculty

Universiti Putra Malaysia
(Chairman)

Name of Examiner 1,

Title (e.g., Professor/ Associate Professor/ Ir; omit if irrelevant)

Name of Faculty

Universiti Putra Malaysia
(Internal Examiner)

Name of Examiner 2,

Title (e.g., Professor/ Associate Professor/ Ir; omit if irrelevant)

Name of Faculty

Universiti Putra Malaysia
(Internal Examiner)

Name of External Examiner,

Title (e.g., Professor/ Associate Professor/ Ir; omit if irrelevant)

Name of Department and/or Faculty

Name of Organisation (University/Institute)

Country

(External Examiner)

AZIZAH ABDUL HAMID, PhD

Professor and Deputy Dean School
of Graduate Studies
Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Mehrnoush Amid, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Mohd Yazid Abd Manap, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Badlishah Sham Baharin, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____

Name of Chairman
of Supervisory
Committee:

Associate Professor Dr. Mehrnoush Amid

Signature: _____

Name of Member
of Supervisory
Committee:

Professor Dr. Mohd Yazid Abd Manap

Signature: _____

Name of Member
of Supervisory
Committee:

Associate Professor Dr. Badlishah Sham Baharin

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF APPENDICES	xvi
LIST OF ABBREVIATIONS	xvii
 CHAPTER	
1 INTRODUCTION	1
 2 LITERATURE REVIEW	4
2.1 Introduction to lipases	4
2.1.1 Lipase catalytic properties	4
2.1.2 Halotolerant and halophilic lipases	5
2.2 Sources of lipases	6
2.2.1 <i>Leuconostoc mesenteroides</i>	7
2.2.1.1 <i>L. mesenteroides</i> subsp. <i>mesenteroides</i> ATCC 8293	8
2.3 Application of microbial lipases	9
2.3.1 Lipase in food industry	9
2.3.2 Lipase in detergent industry	10
2.3.3 Lipase in biomedical industry	11
2.3.4 Lipase in leather industry	12
2.3.5 Lipase in cosmetics and perfume industry	12
2.3.6 Lipase in waste and sewage treatment	13
2.4 Factors affecting microbial lipase production	13
2.4.1 Physical factors	13
2.4.2 Nutritional factors	14
2.5 Purification of lipases	17
2.5.1 Conventional purification of microbial lipases	17
2.5.2 Problems associated with conventional purification techniques	18
2.6 Alternative purification technique: Aqueous two-phase system	18
2.6.1 Theoretical background of ATPS	19
2.6.1.1 Phase formation	19
2.6.1.2 Phase diagram	20
2.6.1.3 Binodal curve	20
2.6.1.4 Tie-line length (TLL)	21
2.6.2 Factors affecting protein partitioning in ATPS	22
2.6.2.1 Effect of system pH on protein partitioning	22

2.6.2.2	Effect of system temperature on protein partitioning	22
2.6.2.3	Effect of concentration of crude load on protein partitioning	22
2.6.3	Types of components used in ATPS	23
2.7	Characterization of microbial lipases	24
2.7.1	pH and temperature optima	25
2.7.2	pH and temperature stability	28
2.7.3	Effect of metal ions	28
2.7.4	Effect of organic solvents	34
2.7.5	Effect of surfactants and other additives	34
3	MATERIALS AND METHODS	35
3.1	Materials	35
3.2	Culture maintenance	35
3.3	Inoculum preparation	35
3.4	Production medium	35
3.5	Enzyme extraction	36
3.6	Lipase activity assay	36
3.7	Protein concentration determination	36
3.8	Optimization of nutritional factors for lipase production	36
3.8.1	Effect of carbon sources on lipase production	36
3.8.2	Effect of emulsifier on lipase production	37
3.8.3	Effect of nitrogen sources on lipase production	37
3.8.4	Effect of surfactants on lipase production	37
3.9	Optimization of physical parameters for lipase production	38
3.9.1	Effect of incubation temperature on lipase production	38
3.9.2	Effect of initial pH on lipase production	38
3.10	Purification of lipases using Aqueous Two-Phase System (ATPS)	38
3.10.1	Construction of Triton X-100/Maltitol phase diagram	38
3.10.1.1	Binodal curve	38
3.10.1.2	Tie line-lengths (TLL)	39
3.10.2	Lipase purification using Triton X-100/Maltitol system	40
3.10.3	Evaluation parameters for lipase purification	40
3.10.4	Determination of lipase molecular weight using SDS-PAGE	41
3.11	Characterization of lipase	42
3.11.1	Effect of temperature on lipase activity and stability	42
3.11.2	Effect of pH on lipase activity and stability	42
3.11.3	Effect of surfactant agent on lipase activity	42
3.11.4	Effect of metal ions on lipase activity	42
3.11.5	Effect of salt concentration on lipase stability	43
3.12	Statistical Design and Analysis	43

4	RESULTS AND DISCUSSION	44
4.1	Optimization of nutritional factors on lipase production	44
4.1.1	Effect of carbon sources on lipase production	44
4.1.2	Effect of emulsifier on lipase production	46
4.1.3	Effect of nitrogen sources on lipase production	48
4.1.4	Effect of surfactants on lipase production	49
4.2	Optimization of physical parameters on lipase production	51
4.2.1	Effect of incubation temperature on lipase production	52
4.2.2	Effect of initial pH on lipase production	54
4.3	Purification of lipase using novel aqueous two phase system composed of Triton X-100 and maltitol.	56
4.3.1	Phase diagram and TLL	56
4.3.2	Effect of crude feedstock concentration on lipase partitioning	61
4.3.3	Effect of pH on lipase partitioning	62
4.3.4	Effect of temperature on lipase partitioning	64
4.3.5	Recovery of lipase in Triton X-100/maltitol ATPS	66
4.4	Characterization of purified lipase	67
4.4.1	Effect of temperature on purified lipase activity	67
4.4.2	Effect of temperature on purified lipase stability	68
4.4.3	Effect of pH on purified lipase activity	70
4.4.4	Effect of pH on purified lipase stability	71
4.4.5	Effect of surfactant agents on purified lipase activity	73
4.4.6	Effect of metal ions on purified lipase activity	75
4.4.7	Effect of salt concentration on purified lipase stability	76
5	SUMMARY, CONCLUSION AND RECOMMENDATION	78
5.1	Summary and Conclusion	78
5.2	Recommendation	79
	REFERENCES	80
	APPENDICES	94
	BIODATA OF STUDENT	97
	PUBLICATION	98

LIST OF TABLES

Table		Page
2.1	pH and temperature optima and stability of selected microbial lipases	26
2.2	Characteristics of microbial lipases in the presence of metal ions, organic solvents, surfactants and other additives	30
4.1	Weight fraction composition for the TLLs, at the top (T) and bottom phase (B), and initial biphasic composition of the mixture (M), composed of Triton X – 100 ([Tri X-100]) and maltitol ([Maltitol]) and the corresponding lipase selectivity and purification factor (PF) at the respective TLLs	60

LIST OF FIGURES

Figure	Page
2.1 Reactions catalyzed by lipases	5
2.2 Schematic representation of a phase diagram for an ATPS	21
4.1 Effect of carbon sources on lipase production by <i>L. mesenteroides</i> subsp. <i>mesenteroides</i> ATCC 8293	45
4.2 Production of lipases in the presence or absence of gum arabic in the culture media	47
4.3 Effect of nitrogen sources on lipase production by <i>L. mesenteroides</i> subsp. <i>mesenteroides</i> ATCC 8293	49
4.4 Effect of various surfactants on lipase production from <i>L. mesenteroides</i> subsp. <i>mesenteroides</i> ATCC 8293	51
4.5 Effect of culture temperature on lipase production by <i>L. mesenteroides</i> subsp. <i>mesenteroides</i> ATCC 8293	53
4.6 Effect of initial pH on lipase production by <i>L. mesenteroides</i> subsp. <i>mesenteroides</i> ATCC 8293	55
4.7 Turbidometric titration method for the construction of binodal curve	57
4.8 Phase diagram for ATPS composed of Triton X-100, maltitol and water at room temperature	58
4.9 Effect of crude load on lipase selectivity and yield	62
4.10 Influence of pH on lipase partitioning	64
4.11 Effect of system temperature on lipase partitioning	65
4.12 SDS-PAGE analysis of lipase recovered using Triton X-100/maltitol ATPS	66
4.13 Effect of temperature on the activity lipase produced by <i>L. mesenteroides</i> subsp. <i>mesenteroides</i> ATCC 8293	68
4.14 Effect of temperature on the stability of lipase produced by <i>L. mesenteroides</i> subsp. <i>mesenteroides</i> ATCC 8293	69
4.15 Effect of pH on the activity of purified lipase	71
4.16 Effect of pH on the stability of purified lipase	72

4.17	Effect of surfactants on purified lipase	74
4.18	Effect of 5 mM metal ions on purified lipase	76
4.19	Effect of various NaCl concentrations on lipase activity	77



LIST OF APPENDICES

Appendix	Page
A Standard Curve of BSA For Protein Concentration Determination	94
B Gel Preparation For SDS-PAGE Analysis	95
C Protein Molecular Marker Used in SDS-PAGE Analysis	96



LIST OF ABBREVIATIONS

3-D	3- Dimensional
ANOVA	Analysis of Variance
APS	Ammonium Persulfate
ATCC	American Type Collection Culture
ATPS	Aqueous Two-Phase System
BSA	Bovine Serum Albumin
C	Carbon Chain
CTAB	Cetrimonium Bromide
DEAE	Diethylaminoethyl
DMSO	Dimethyl Sulfoxide
E.C.	Enzyme Commission
EDTA	Ethylenediaminetetraacetic Acid
EMC	Enzyme-Modified Cheese
Exp	Exponential
FDA	Food and Drug Association
FF	Fast Flow
g	Gram
g	Gravitational Force
GRAS	Generally Regarded As Safe
HCl	Hydrochloric Acid
H ₂ O ₂	Hydrogen Peroxide
LAB	Lactic Acid Bacteria

K_M	Michaelis Constant
kDa	Kilo Dalton
M	Molar
mg	Milligram
Min	Minute
mL	Millilitre
mM	MilliMolar
MRS	De Man, Rogosa and Sharpe
N	Normality
NaOCl ₂	Sodium Chlorite
NaOH	Sodium Hydroxide
nm	Nano Metre
PEG	Polyethylene Glycol
pI	Isoelectric Point
PMSF	Phenylmethane Sulfonyl Fluoride
pNPP	Para-Nitrophenyl Palmitate
PUFA	Polyunsaturated Fatty Acid
Rpm	Revolution per Minute
SD	Standard Deviation
SDS	Sodium Dodecyl Sulphate
SDS-PAGE	Sodium Dodecyl Sulphate-Polyacrylamide Gel Electrophoresis
<i>Sn</i>	Stereospecifically Numbered
TEMED	Tetramethylethylenediamine

TLL	Tie-Line Length
Tris	Trisaminomethane
U	Unit
μL	Micro Litre
μmol	Micro Mol
U/mg	Unit per Milligram
U/mL	Unit per Millilitre
USD	United State Dollar
V	Volt
V_{max}	Maximum Rate of Reaction
v/v	Volume per Volume
VOC	Volatile Organic Compound
w/v	Weight per Volume
w/w	Weight per Weight
Y_{T}	Top Phase Yield

CHAPTER 1

INTRODUCTION

Enzymes have become an important catalyst in the biotechnology industry as the demand for industrial enzymes are increasing throughout the years. The global market for industrial enzymes in 2015 was estimated at USD 4.94 billion and is expected to increase 6% annually to reach USD 6.99 billion by 2021 (Business Wire, 2016). Hydrolytic enzymes such as proteases, carbohydrases and lipases dominate the total sale of global enzymes. Recently, lipases have garnered particular attention in the biotechnology industry owing to their multifaceted properties. The global lipase market is expected to increase with an average of 6.5% annually between 2015 and 2020 to reach USD 590.5 million by 2020 (Markets and Markets, 2015).

Lipases (triacylglycerol acylhydrolase E.C. 3.1.1.3) are a group of enzymes that are able to catalyze the hydrolysis of triglycerides into glycerol and fatty acids at lipid-water interface (Emtenani *et al.*, 2013). However, the catalytic reaction of lipases largely depends on the interfacial area, as lipases found in non-aqueous environment, are able to reverse the reaction, which leads to the synthesis of triglycerides from glycerol and free fatty acid (Papagora *et al.*, 2013). Due to their versatile reactions in both aqueous and non-aqueous media, lipases have become one of the commonly used hydrolytic enzymes in the enzyme-related industries such as food, pharmaceutical, detergent, cosmetic and paper manufacturing (Ma *et al.*, 2006; Sharma *et al.*, 2002).

Lipases are ubiquitous as they can be found in plants, animals and microorganisms. However, lipases from microorganisms are found to be more suitable for biotechnology and commercial applications compared to animal and plant lipases due to the various advantages of microbial lipases such as short production time, large quantities production, simple extraction procedure and cost effectiveness (Rabbani *et al.*, 2013). Although there are many microbial lipases currently available for the use of various industries, they are still insufficient to fulfill the operating requirements of most industries because lipases produced by different microbes will have different optimal reaction conditions. These conditions can greatly influence the applications of lipases in the industries.

Currently, limited researches regarding production and characterization of halotolerant lipases have become a common limitation for lipases used in industrial applications (Asoodeh & Ghanbari, 2013). Halotolerant enzymes are able to tolerate and remain active over a wide range of salt concentration. These enzymes are particularly essential in food industries. Lipases are commonly used in food fermentation, which is one of the common processes in food industry. This process involves salting step, in which high concentration of salt is present (Esteban-Torres *et al.*, 2015). Hence, lipases used for this process are required to be halotolerant so that they can remain active and stable at high salt concentration in order to carry out the

fermentation process efficiently. Thus, it is important to find novel microbial enzyme sources for the production of lipases with suitable characteristics to meet the requirements of particular industries.

Leuconostoc mesenteroides is a gram positive and facultative anaerobe bacterium. It belongs to the group of lactic acid bacteria and widely used in the production of various fermented foods such as kimchi, sauerkraut, dairy and bread dough. Several lactic acid bacteria have been reported to produce lipase (Esteban-Torres *et al.*, 2015; Ramyasree & Dutta, 2013). Although many studies have been done on the production of lipase from lactic acid bacteria, lipase production by *L. mesenteroides* has never been reported. Compared to other lactic acid bacteria, *L. mesenteroides* can tolerate fairly high concentration of salt and sugar. *L. mesenteroides* is also non-pathogenic and classified as generally regarded as safe (GRAS) organism by the U.S Food and Drug Administration (FDA). These characteristics are essential for the commercial applications of lipases in various industries.

Although the demand for enzymes is growing, the high cost of enzymes is a major concern for the enzyme-related industries. Enzymes are costly due to the complexity and difficulty of their downstream processing. The downstream processing of enzymes usually involves purification step, which accounts for more than 70% of total production cost of the enzymes (Raja *et al.*, 2012). The purification methods traditionally used in the industries include several steps such as ultrafiltration, precipitation (salting out) and chromatography (ion exchange, hydrophobic interaction, size exclusion, and affinity), which resulted in the loss of product in each steps (Benavides *et al.*, 2011). Furthermore, these conventional methods of purification are expensive, multistep, and discontinuous as well as labor and time extensive. Due to these characteristics, the conventional purification methods are found to be difficult to scale up, thus making them uneconomical at large-scale. Hence, there is a need to develop quick, cost-effective, efficient and scalable purification technique with a potential for continuous operations, which can improve the yield and purity of the biological product.

Aqueous two-phase system (ATPS) is an alternative purification technique that meets all the aforementioned criteria. ATPS was introduced during the mid-1950's by Albertsson (Srinivas, 2000), however its potential as an efficient purification method has been realized only recently. ATPS is conducted by mixing two different water-soluble compounds in water until the limiting concentrations of these two compounds have been exceeded, which then results in the formation of two immiscible aqueous two phases. When the mixture containing biomolecule of interest is added into this system, the target biomolecule will be partitioned and concentrated into one of the phases and the contaminants will be partitioned into the other phase (Raja *et al.*, 2012). Due to its water-rich environment, ATPS serves as a friendly and gentle environment for biological materials. Besides, the parameters of the system can be controlled and manipulated in order to achieve optimum partitioning (Amid *et al.*, 2011).

Furthermore, this system is economical, continuous, simple, rapid, and easy to scale up.

The search for novel lipase-producing bacteria should be continuously pursued because currently there is limited availability of lipase possessing satisfactory operating characteristics. Besides, the high cost of lipases has become a significant concern for the biotechnological industries. Therefore, there is a need to develop inexpensive, simple, continuous and scalable purification strategy to cater to industries' needs of a fast and economic downstream process. Furthermore, this process should provide a biocompatible environment for lipase in order to improve the yield and purity of the lipase. Thus, this research was undertaken with the following objectives:

- I. To optimize production of extracellular lipases from *Leuconostoc mesenteroides* subsp. *mesenteroides* ATCC 8293.
- II. To purify extracellular lipases produced by *L. mesenteroides* subsp. *mesenteroides* ATCC 8293 using novel aqueous system (ATPS) composed of Triton X-100 and maltitol.
- III. To characterize the properties of the purified lipases.

REFERENCES

- Adan, A. (2009). *Isolation And Identification Of A Lipase Producing Psychotropic Bacteria From Soil: Cloning And Partial Characterization Of Its Lipase*. Master Thesis. Izmir Institute of Technology.
- Amid, M., Manap, M. Y., Hussin, M., & Mustafa, S. (2015). A novel aqueous two phase system composed of surfactant and xylitol for the purification of lipase from pumpkin (*Cucurbita moschata*) seeds and recycling of phase components. *Molecules*, 20(6), 11184–11201.
- Amid, M., Murshid, F. S., Manap, M. Y., & Hussin, M. (2015). A novel aqueous micellar two-phase system composed of surfactant and sorbitol for purification of pectinase enzyme from *Psidium guajava* and recycling phase components. *BioMed Research International*, ID 815413.
- Amoozegar, M. A., Salehghamari, E., Khajeh, K., Kabiri, M., & Naddaf, S. (2008). Production of an extracellular thermohalophilic lipase from a moderately halophilic bacterium, *Salinivibrio* sp. strain SA-2. *Journal of Basic Microbiology*, 48(3), 160–167.
- Andualema, B., & Gessesse, A. (2012). Microbial Lipase and Their Industrial Applications: Review. *Biotechnology*, 11(3), 100–118.
- Aravindan, R., Anbumathi, P., & Viruthagiri, T. (2007). Lipase applications in food industry. *Indian Journal of Biotechnology*, 6(April), 141–158.
- Asenjo, J. A., & Andrews, B. A. (2011). Aqueous two-phase systems for protein separation: A perspective. *Journal of Chromatography A*, 1218(49), 8826–8835.
- Asenjo, J. A., & Andrews, B. A. (2012). Aqueous two-phase systems for protein separation: Phase separation and applications. *Journal of Chromatography A*, 1238, 1–10.
- Asoodeh, A., & Ghanbari, T. (2013). Characterization of an extracellular thermophilic alkaline esterase produced by *Bacillus subtilis* DR8806. *Journal of Molecular Catalysis B: Enzymatic*, 85–86, 49–55.
- Aulakh, S. S., & Prakash, R. (2010). Optimization of medium and process parameters for the production of lipase from an oil-tolerant *Aspergillus* sp. (RBD-01). *Journal of Basic Microbiology*, 50(1), 37–42.
- Ayaz, B., Ugur, A., & Boran, R. (2015). Purification and characterization of organic solvent-tolerant lipase from *Streptomyces* sp. OC119-7 for biodiesel production. *Biocatalysis and Agricultural Biotechnology*, 4(1), 103–108.

- Bae, J., Kim, I., Lee, K., Hou, C. T., & Kim, H. (2017). Molecular cloning and characterization of a novel cold-active lipase from *Pichia lynferdii* NRRL Y-7723. *Biocatalysis and Agricultural Biotechnology*, 11, 19–25.
- Balaji, L., & Jayaraman, G. (2014). Metal ion activated lipase from halotolerant *Bacillus* sp. VITL8 displays broader operational range. *International Journal of Biological Macromolecules*, 67, 380–386.
- Balasubramaniam, D. (2003). *Lysozyme Separation From Tobacco Extract By Aqueous Two-Phase Extraction*. Master Thesis. Virginia Polytechnic Institute and State University.
- Barros, M., Fleuri, L. F., & Macedo, G. A. (2010). Seed lipases: sources, applications and properties - a review. *Brazilian Journal of Chemical Engineering*, 27(1), 15–29.
- Baynes, J. W., & Dominiczak, M. W. (2014). *Medical Biochemistry* (4th ed). Philadelphia, US: Saunders.
- Benavides, J., Rito-Palomares, M., & Asenjo, J. A. (2011). Aqueous Two-Phase Systems. *Comprehensive Biotechnology*, 2, 697–713.
- Beşel, E. (2003). *Use Of Triton X-114A Aqueous Two Phase System For Recovery Of Mushroom (Agaricus Bisporus) Polyphenoloxidase*. Master Thesis. The Middle East Technical University.
- Bhargavi, P. L., Manjushri, R., & Neelakanta, R. (2010). Lipase production by lactic acid bacteria in submerged and solid state fermentation. *BTAIJ*, 4(3), 126–129.
- Bora, L., & Bora, M. (2012). Optimization of extracellular thermophilic highly alkaline lipase from thermophilic *Bacillus* sp. isolated from hot spring of Arunachal Pradesh, India. *Brazilian Journal of Microbiology*, 30–42.
- Bose, A., & Keharia, H. (2013). Production, characterization and applications of organic solvent tolerant lipase by *Pseudomonas aeruginosa* AAU2. *Biocatalysis and Agricultural Biotechnology*, 2(3), 255–266.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1–2), 248–254.
- Business Wire. (2016). Global Industrial Enzymes Market Worth USD 6.99 Billion by 2021 - Analysis, Technologies & Forecasts Report 2016-2021 - Vendors: Novozymes, DuPont, DSM - Research and Markets. <http://www.businesswire.com/news/home/20161121005632/en/Global-Industrial-Enzymes-Market-Worth-USD-6.99> (accessed 13th February 2017).

- Cabral, J. M. S. (2007). Cell Separation: Fundamentals, Analytical and Preparative Methods. In T. Scheper (Ed.), *Advances in Biochemical Engineering/Biotechnology* (Vol. 106). New York: Springer.
- Cammarota, M. C., & Freire, D. M. G. (2006). A review on hydrolytic enzymes in the treatment of wastewater with high oil and grease content. *Bioresource Technology*, 97(17), 2195–2210.
- Carvalho, T., Finotelli, P. V., Bonomo, R. C. F., Franco, M., & Amaral, P. F. F. (2017). Evaluating aqueous two-phase systems for *Yarrowia lipolytica* extracellular lipase purification. *Process Biochemistry*, 53, 259–266.
- Chakraborty, K., & Paulraj, R. (2009). Purification and biochemical characterization of an extracellular lipase from *Pseudomonas fluorescens* MTCC 2421. *Journal of Agricultural and Food Chemistry*, 57(9), 3859–3866.
- Charoenpanich, J., Suktanarag, S., & Toobbucha, N. (2011). Production of a thermostable lipase by *Aeromonas* sp. EBB-1 isolated from marine sludge in Angsila, Thailand. *ScienceAsia*, 37(2), 105–114.
- Chinedu, S. M., & Emmanuel, E. (2014). Isolation of microorganisms associated with deterioration of tomato (*Lycopersicon esculentum*) and pawpaw (*Carica papaya*) fruits. *International Journal of Current Microbiology and Applied Sciences*, 3(5), 501–512.
- Choudhury, P., & Bhunia, B. (2015). Industrial application of lipase: a review. *Biopharm Journal*, 1(2), 41–47.
- Cornish-Bowden, A. (2014). *Principles of enzyme kinetics*. London, UK: Butterworth.
- Daoud, L., Kamoun, J., Ali, M. B., Jallouli, R., Bradai, R., Mechichi, T., ... Aloulou, A. (2013). Purification and biochemical characterization of a halotolerant *Staphylococcus* sp. extracellular lipase. *International Journal of Biological Macromolecules*, 57, 232–237.
- Darvishi, F., Destain, J., Nahvi, I., Thonart, P., & Zarkesh-Esfahani, H. (2011). High-level production of extracellular lipase by *Yarrowia lipolytica* mutants from methyl oleate. *New Biotechnology*, 28(6), 756–760.
- De Almeida, A. F., Tauk-Tornisielo, S. M., & Carmona, E. C. (2013). Acid lipase from *Candida viswanathii*: production, biochemical properties, and potential application. *BioMed Research International*, 2013, 1–10.
- de Brito Cardoso, G., Souza, I. N., Mourão, T., Freire, M. G., Soares, C. M. F., & Lima, Á. S. (2014). Novel aqueous two-phase systems composed of acetonitrile and polyols: Phase diagrams and extractive performance. *Separation and Purification Technology*, 124, 54–60.

- de Morais Junior, W. G., Kamimura, E. S., Ribeiro, E. J., Pessela, B. C., Cardoso, V. L., & de Resende, M. M. (2016). Optimization of the production and characterization of lipase from *Candida rugosa* and *Geotrichum candidum* in soybean molasses by submerged fermentation. *Protein Expression and Purification*, 123, 26–34.
- Dhiman, S., & Chapadgaonkar, S. S. (2013). Optimization of lipase production medium for a bacterial isolate. *International Journal of ChemTech Research*, 5(6), 2837–2843.
- Dimic, G. (2006). Characteristics of the *Leuconostoc mesenteroides* subsp. *mesenteroides* strains from fresh vegetables. *APTEFF*, 192(37), 3–11.
- Domínguez, A., Deive, F. J., Sanromán, M. A., & Longo, M. A. (2003). Effect of lipids and surfactants on extracellular lipase production by *Yarrowia lipolytica*. *Journal of Chemical Technology and Biotechnology*, 78(11), 1166–1170.
- Dreyer, S. E. (2008). *Aqueous Two-Phase Extraction Of Proteins And Enzymes Using Tetraalkylammonium-Based Ionic Liquids*. PhD Thesis. University of Rostock.
- Emtenani, S., Asoodeh, A., & Emtanani, S. (2013). Molecular cloning of a thermo-alkaliphilic lipase from *Bacillus subtilis* DR8806: expression and biochemical characterization. *Process Biochemistry*, 48(11), 1679–1685.
- Enger, E. D., Ross, F. C., & Bailey, D. B. (2012). *Concepts in Biology* (14th ed.). New York, USA: McGraw-Hill.
- Ertuğrul, S., Dönmez, G., & Takaç, S. (2007). Isolation of lipase producing *Bacillus* sp. from olive mill wastewater and improving its enzyme activity. *Journal of Hazardous Materials*, 149(3), 720–724.
- Esakkiraj, P., Antonyraj, C. B., Meleppat, B., Ankaiah, D., Ayyanna, R., Ahamed, S. I. B., & Arul, V. (2017). Molecular characterization and application of lipase from *Bacillus* sp. PU1 and investigation of structural changes based on pH and temperature using MD simulation. *International Journal of Biological Macromolecules*, 103, 47–56.
- Esteban-Torres, M., Mancheño, J. M., de las Rivas, B., & Muñoz, R. (2015). Characterization of a halotolerant lipase from the lactic acid bacteria *Lactobacillus plantarum* useful in food fermentations. *LWT - Food Science and Technology*, 60(1), 246–252.
- Florou-Paneri, P., Christaki, E., & Bonos, E. (2013). Lactic Acid Bacteria As Source Of Functional Ingredients. In J. . Kongo (Ed.), *Lactic Acid Bacteria-R& D for Food, Health and Livestock Purposes* (pp. 589–614). InTech.

- Ganasen, M., Yaacob, N., Raja Abd Rahman, R. N. Z., Leow, A. T. C., Basri, M., Salleh, A. B., & Mohamad Ali, M. S. (2016). Cold-adapted organic solvent tolerant alkalophilic family I.3 lipase from an *Antarctic Pseudomonas*. *International Journal of Biological Macromolecules*, 92, 1266–1276.
- Gaoa, X.-G., Cao, S.-G., & Zhang, K.-C. (2000). Production, properties and application to nonaqueous enzymatic catalysis of lipase from a newly isolated *Pseudomonas* strain. *Enzyme and Microbial Technology*, 27(1–2), 74–82.
- Gaur, R., Gupta, A., & Khare, S. K. (2008). Purification and characterization of lipase from solvent tolerant *Pseudomonas aeruginosa* PseA. *Process Biochemistry*, 43, 1040–1046.
- Gobbetti, M., Fox, P. F., & Stepaniak, L. (1997). Isolation and characterization of a tributyrin esterase from *Lactobacillus plantarum* 2739. *Journal of Dairy Science*, 80(12), 3099–3106.
- Goel, A. (2006). *Surface Chemistry*. New Delhi, India: Discovery Publishing House.
- Golaki, B. P., Aminzadeh, S., Karkhane, A. A., Yakhchali, B., Farrokh, P., Khaleghinejad, S. H., ... Mehrpooyan, S. (2015). Cloning, expression, purification, and characterization of lipase 3646 from thermophilic indigenous *Cohnella* sp. A01. *Protein Expression and Purification*, 109, 120–126.
- Gravie, E. I. (1976). Hybridization between the deoxyribonucleic acids of some strains of heterofermentative lactic acid bacteria. *International Journal of Systematic Bacteriology*, 26(2), 116–122.
- Gravie, E. I. (1983). *Leuconostoc mesenteroides* subsp. *cremoris* (Knudsen and Sgrensen) comb. nov. and *Leuconostoc mesenteroides* subsp. *dextranicum* (Beijerinck) comb. nov. *International Journal of Systematic Bacteriology*, 33(1), 118–119.
- Gricajeva, A., Bendikien, V., & Kalediene, L. (2016). Lipase of *Bacillus stratosphericus* L1: cloning, expression and characterization. *International Journal of Biological Macromolecules*, 92, 96–104.
- Gupta, R., Gupta, N., & Rathi, P. (2004). Bacterial lipases: an overview of production, purification and biochemical properties. *Applied Microbiology and Biotechnology*, 64(6), 763–781.
- Gururaj, P., Ramalingam, S., Devi, G. N., & Gautam, P. (2016). Process optimization for production and purification of a thermostable, organic solvent tolerant lipase from *Acinetobacter* sp. AU07. *Brazilian Journal of Microbiology*, 47(3), 647–657.

- Guzmán, M. N. De, Vargas, V. a, Antezana, H., & Svoboda, M. (2008). Lipolytic enzyme production by halophilic / halotolerant microorganisms isolated from Laguna Verde , Bolivia. *Revista Boliviana de Quimica*, 25(1), 14–23.
- Hasan, F., Shah, A., Javed, S., & Hameed, A. (2010). Enzymes used in detergents: lipases. *African Journal of Biotechnology*, 9(31), 4836–4844.
- Hong-wei, Y., Jun, H., Ning, L., Xiao-sha, Q., & Ying-min, J. (2009). Fermentation performance and characterization of cold-adapted lipase produced with *Pseudomonas* Lip35. *Agricultural Sciences in China*, 8(8), 956–962.
- Hontebeyrje, M., & Gasser, F. (1977). Deoxyribonucleic acid homologies in the Genus *Leuconostoc*. *International Journal of Systematic Bacteriology*. 27(1), 9-14.
- Immanuel, G., Esakkiraj, P., Jebadhas, A., Iyapparaj, P., & Palavesam, A. (2008). Investigation of lipase production by milk isolate *Serratia rubidaea*. *Food Technology and Biotechnology*, 46(1), 60–65.
- Javed, S., Azeem, F., Hussain, S., Rasul, I., Siddique, M. H., Riaz, M., ... Nadeem, H. (2017). Bacterial lipases: a review on purification and characterization. *Progress in Biophysics and Molecular Biology*, 1–12.
- Ji, X., Li, S., Wang, B., Zhang, Q., Lin, L., Dong, Z., & Wei, Y. (2015). Expression , purification and characterization of a functional , recombinant , cold-active lipase (LipA) from psychrotrophic *Yersinia enterocolitica*. *Protein Expression and Purification*, 115, 125–131.
- Joseph, B., Upadhyaya, S., & Ramteke, P. (2011). Production of cold-active bacterial lipases through semisolid state fermentation using oil cakes. *Enzyme Research*, 2011, 6.
- Kambourova, M., Kirilova, N., Mandeva, R., & Derekova, A. (2003). Purification and properties of thermostable lipase from a thermophilic *Bacillus stearothermophilus* MC 7. *Journal of Molecular Catalysis B: Enzymatic*, 22(5–6), 307–313.
- Kamini, N. R., Fujii, T., Kurosu, T., & Iefuji, H. (2000). Production, purification and characterization of an extracellular lipase from the yeast, *Cryptococcus* sp. S-2. *Process Biochemistry*, 36(4), 317–324.
- Kanlayakrit, W., & Boonpan, A. (2007). Screening of halophilic lipase-producing bacteria and characterization of enzyme for fish sauce quality improvement. *Kasetsart Journal Natural Science*, 41(3), 576–585.
- Katz, M., Medina, R., Gonzalez, S., & Oliver, G. (2002). Esterolytic and lipolytic activities of lactic acid bacteria isolated from ewe's milk and cheese. *Journal of Food Protection*, 65(12), 1997–2001.

- Keskin, S. Ö., Sumnu, G., & Sahin, S. (2004). Usage of enzymes in a novel baking process. *Nahrung - Food*, 48(2), 156–160.
- Kiran, G. S., Lipton, A. N., Kennedy, J., Dobson, A. D. W., & Selvin, J. (2014). A halotolerant thermostable lipase from the marine bacterium *Oceanobacillus* sp. PUMB02 with an ability to disrupt bacterial biofilms. *Bioengineered*, 5(5), 305–
- Kumar, R., Sharma, A., Kumar, A., & Singh, D. (2012). Lipase from *Bacillus pumilus* RK31: production, purification and some properties. *World Applied Sciences Journal*, 16(7), 940–948.
- Kwon, C. H., Lee, J. H., Kim, S. W., & Kang, J. W. (2009). Lipase-catalyzed esterification of (S)-naproxen ethyl ester in supercritical carbon dioxide. *Journal of Microbiology and Biotechnology*, 19(12), 1596–1602.
- Lau, H. L., Ariff, A., Woo, K. K., Ling, T. C., & Hii, S. L. (2011). Production and optimization of alkalostable lipase by alkalophilic *Burkholderia cenocepacia* ST8. *African Journal of Biotechnology*, 10(36), 7002–7009.
- Lee, S. Y., Khoiroh, I., Ling, T. C., & Show, P. L. (2017). Enhanced recovery of lipase derived from *Burkholderia cepacia* from fermentation broth using recyclable ionic liquid/polymer-based aqueous two-phase systems. *Separation and Purification Technology*, 179, 152–160.
- Li, M., Yang, L., Xu, G., & Wu, J. (2016). Cloning and characterization of a novel lipase from *Stenotrophomonas maltophilia* GS11 : The first member of a new bacterial lipase family XVI. *Journal of Biotechnology*, 228, 30–36.
- Lima, V. M. G., Krieger, N., Sarquis, M. I. M., Mitchell, D. a, Ramos, L. P., & Fontana, J. D. (2003). Effect of nitrogen and carbon sources on lipase production by *Penicillium aurantiogriseum*. *Food Technology and Biotechnology*, 41(2), 105–110.
- Lipu, L. R., Li, C., Li, Q., Zhang, Y., Gong, Z., Ren, S., ... Xie, J. (2017). Characterization and function of *Mycobacterium tuberculosis* H37Rv lipase Rv1076 (LipU). *Microbiological Research*, 196, 7–16.
- Liu, Z.-Q., Zheng, X.-B., Zhang, S.-P., & Zheng, Y.-G. (2012). Cloning, expression and characterization of a lipase gene from the *Candida antarctica* ZJB09193 and its application in biosynthesis of vitamin A esters. *Microbiological Research*, 167(8), 452–460.
- Long, K. (2009). *Unlocking the Miracle of Lipases*. Serdang: MARDI.

- Lopes, M. de F. S., Leitão, A. L., Regalla, M., Figueiredo Marques, J. J., Carrondo, M. J. T., & Crespo, M. T. B. (2002). Characterization of a highly thermostable extracellular lipase from *Lactobacillus plantarum*. *International Journal of Food Microbiology*, 76(1–2), 107–115.
- Lovrien, R., & Matulis, D. (2005). Assays for total protein. In *Current protocols in microbiology*.
- Lukaszewicz, M., Jablonski, S., & Krasowska, A. (2013). Characterisation of alkaline lipase from an arctic yeast strain *Rhodospiridium Babjevae* Bd19. *European Scientific Journal*, 441–449.
- Lule, V. K., Singh, R., Pophaly, S. D., Poonam, & Tomar, S. K. (2016). Production and structural characterisation of dextran from an indigenous strain of *Leuconostoc mesenteroides* BA08 in Whey. *International Journal of Dairy Technology*, 69(4), 520–531.
- Ma, J., Zhang, Z., Wang, B., Kong, X., Wang, Y., Cao, S., & Feng, Y. (2006). Overexpression and characterization of a lipase from *Bacillus subtilis*. *Protein Expression and Purification*, 45(1), 22–29.
- Mahdi, B. A., Bhattacharya, A., & Gupta, A. (2012). Enhanced lipase production from *Aeromonas* sp. S1 using Sal deoiled seed cake as novel natural substrate for potential application in dairy wastewater treatment. *Journal of Chemical Technology and Biotechnology*, 87(3), 418–426.
- Maia, M. M. D., Heasley, A., de Moraes, M. M., Melo, E. H. M., Moraes Jr, M. A., Ledingham, W. M., & Lima Filho, J. L. (2001). Effect of culture conditions on lipase production by *Fusarium solani* in batch fermentation. *Bioresource Technology*, 76(1), 23–27.
- Markets and Markets. (2015). Lipase Market by Source, Application, by Geography - Global Forecast to 2020. <https://www.marketsandmarkets.com/Market-Reports/lipase-market-205981206.html> (accessed 15 February 2017).
- Masomian, M. (2007). *Characterization of a Thermostable Lipase From Aneurinibacillus Thermoaerophilus Strain HZ*. Master Thesis. Universiti Putra Malaysia.
- Massadeh, M. I., & Sabra, F. M. (2011). Production and characterization of lipase from *Bacillus stearothermophilus*. *African Journal of Biotechnology*, 10(61), 13139–13146.
- Massadeh, M., Sabra, F., Dajani, R., & Arafat, A. (2012). Purification of lipase enzyme produced by *Bacillus Stearothermophilus* HU1. *International Conference on Eco-Systems and Biological Sciences*, 34–37.

- Matthews, A., Grimaldi, A., Walker, M., Bartowsky, E., Grbin, P., & Jiranek, V. (2004). Lactic acid bacteria as a potential source of enzymes for use in vinification. *Applied and Environmental Microbiology*, 70(10), 5715–5731.
- Mehrnoush, A., Sarker, M. Z. I., Mustafa, S., & Mohd Yazid, A. M. (2011). Direct purification of pectinase from mango (*Mangifera Indica* Cv. *Chokanan*) peel using a PEG/salt-based aqueous two phase system. *Molecules*, 16(10), 8419–8427.
- Merchuk, J. C., Andrews, B. a, & Asenjo, J. a. (1998). Aqueous two-phase systems for protein separation studies on phase separation. *Journal of Chromatography. B*, 711(1–2), 285–293.
- Minuth, T. (2000). Extraction of Amphiphilic Proteins Using Detergent-Based Aqueous Two-Phase Systems. In R. Hatti-kaul (Ed.) *Aqueous Two-Phase Systems: Methods and Protocols* (pp. 291–302). Totowa, NJ: Humana Press.
- Mohamed, S. A., Abdel-Mageed, H. M., Tayel, S. A., El-Nabrawi, M. A., & Fahmy, A. S. (2011). Characterization of *Mucor racemosus* lipase with potential application for the treatment of cellulite. *Process Biochemistry*, 46(3), 642–648.
- Moosavi-Nasab, M., Alahdad, Z., & Nazemi, S. H. (2009). Characterization of the dextran produced by *Leuconostoc mesenteroides* from date fruit extract. *Iran Agricultural Research*, 27(1–2), 1–10.
- Nicanuzia, J., Aparecida, J., Cruz, B., & Pastore, G. M. (2006). Characterization of alkaline lipase from fusarium oxysporum and the effect of different surfactants and detergents on the enzyme activity. *Brazilian Journal of Microbiology*, 37, 505–509.
- Nikita, C., & Hemangi, D. (2012). Isolation , identification and characterization of lactic acid bacteria from dairy sludge sample. *Journal of Environmental Research And Development*, 17(1), 234–244.
- Olusesan, A. T. (2010). *Production, Purification and Characterization of Thermostable Lipase From an Extremophilic Bacillus Subtilis NS 8*. Master Thesis. Universiti Putra Malaysia.
- Olusesan, A. T., Azura, L. K., Forghani, B., Bakar, F. A., Mohamed, A. K. S., Radu, S., ... Saari, N. (2011). Purification, characterization and thermal inactivation kinetics of a non-regioselective thermostable lipase from a genotypically identified extremophilic *Bacillus subtilis* NS 8. *New Biotechnology*, 28(6), 738–745.
- Ooi, C. W. (2011). *Recovery of Extracellular Lipase From Burkholderia SP. ST8 In Aqueous Two-Phase Systems*. PhD Thesis. Universiti Putra Malaysia.

- Ozgen, M., Attar, A., Elalmis, Y., Birbir, M., & Yucel, S. (2016). Enzymatic activity of a novel halotolerant lipase from *Haloarcula hispanica* 2TK2. *Polish Journal of Chemical Technology*, 18(2), 3–8.
- Padmapriya, B., Rajeswari, T., Noushida, E., Sethupalan, D. G., & Venil, C. K. (2011). Production of lipase enzyme from *Lactobacillus* spp. and its application in the degradation of meat. *World Applied Sciences Journal*, 12(10), 1798–1802.
- Papagora, C., Roukas, T., & Kotzekidou, P. (2013). Optimization of extracellular lipase production by *Debaryomyces hansenii* isolates from dry-salted olives using response surface methodology. *Food and Bioproducts Processing*, 91(4), 413–420.
- Pérez, D., Martín, S., Fernández-Lorente, G., Filice, M., Guisán, J. M., Ventosa, A., ... Mellado, E. (2011). A novel halophilic lipase, LipBL, showing high efficiency in the production of eicosapentaenoic acid (EPA). *PLoS ONE*, 6(8), 1–11.
- Rabbani, M., Bagherinejad, M. R., Sadeghi, H. M., Shariat, Z. S., Etemadifar, Z., Moazen, F., ... Zaghian, S. (2013). Isolation and characterization of novel thermophilic lipase-secreting bacteria. *Brazilian Journal of Microbiology*, 44(4), 1113–1119.
- Rahman, R. N. Z. A., Baharum, S. N., Salleh, A. B., & Basri, M. (2006). S5 Lipase: an organic solvent tolerant enzyme. *Journal of Microbiology*, 44(6), 583–590.
- Raja, S., Murty, V. R., Thivaharan, V., Rajasekar, V., & Ramesh, V. (2012). Aqueous two phase systems for the recovery of biomolecules – a review. *Science and Technology*, 1(1), 7–16.
- Rajesh, E. M., Arthe, R., Rajendran, R., Balakumar, C., Pradeepa, N., & Anitha, S. (2010). Investigation of lipase production by *Trichoderma Reesei* and optimization of production parameters. *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 9(7), 1177–1189.
- Ramakrishnan, V., Goveas, L. C., Suralikerimath, N., Jampani, C., Halami, P. M., & Narayan, B. (2016). Extraction and purification of lipase from *Enterococcus faecium* MTCC5695 by PEG/phosphate aqueous-two phase system (ATPS) and its biochemical characterization. *Biocatalysis and Agricultural Biotechnology*, 6, 19–27.
- Ramani, K., Chockalingam, E., & Sekaran, G. (2010). Production of a novel extracellular acidic lipase from *Pseudomonas gessardii* using slaughterhouse waste as a substrate. *Journal of Industrial Microbiology and Biotechnology*, 37(5), 531–535.

- Ramyasree, S., & Dutta, J. R. (2013). The effect of process parameters in enhancement of lipase production by co-culture of lactic acid bacteria and their mutagenesis study. *Biocatalysis and Agricultural Biotechnology*, 2(4), 393–398.
- Rashmi, B. S., & Gayathri, D. (2014). Partial purification, characterization of *Lactobacillus* sp. G5 lipase and their probiotic potential. *International Food Research Journal*, 21(5), 1737–1743.
- Ray, A. (2012). Application of lipase in industry. *Asian Journal of Pharmaceutical Technology*, 2(2), 33–37.
- Saengsanga, T., Siripornadulsil, W., & Siripornadulsil, S. (2016). Molecular and enzymatic characterization of alkaline lipase from *Bacillus amyloliquefaciens* E1PA isolated from lipid-rich food waste. *Enzyme and Microbial Technology*, 82, 23–33.
- Sajur, S. A., Saguir, F. M., & Manca de Nadra, M. C. (2007). Effect of dominant specie of lactic acid bacteria from tomato on natural microflora development in tomato purée. *Food Control*, 18(5), 594–600.
- Sarwat, F., Qader, S. A. U., Aman, A., & Ahmed, N. (2008). Production & characterization of a unique dextran from an indigenous *Leuconostoc mesenteroides* CMG713. *International Journal of Biological Sciences*, 4(6), 379–386.
- Saxena, R. K., Sheoran, A., Giri, B., & Davidson, W. S. (2003). Purification strategies for microbial lipases. *Journal of Microbiological Methods*, 52(1), 1–18.
- Schindler, J., & Nothwang, H. G. (2006). Aqueous polymer two-phase systems: Effective tools for plasma membrane proteomics. *Proteomics*, 6(20), 5409–5417.
- Sethi, B. K., Nanda, P. K., & Sahoo, S. (2016). Characterization of biotechnologically relevant extracellular lipase produced by *Aspergillus terreus* NCFT 4269 . 10. *Brazilian Journal of Microbiology*, 47(1), 143–149.
- Sharma, C., Sharma, P., & Kanwar, S. (2012). Optimization of production conditions of lipase from *B. licheniformis* MTCC-10498. *Research Journal of Recent Sciences*, 1(7), 25–32.
- Sharma, D., Kumbhar, B. K., Verma, A. K., & Tewari, L. (2014). Optimization of critical growth parameters for enhancing extracellular lipase production by alkalophilic *Bacillus* sp. *Biocatalysis and Agricultural Biotechnology*, 3(4), 205–211.

- Sharma, P., Sharma, N., Pathania, S., & Handa, S. (2017). Purification and characterization of lipase by *Bacillus methylotrophicus* PS3 under submerged fermentation and its application in detergent industry. *Journal of Genetic Engineering and Biotechnology*.
- Sharma, R., Chisti, Y., & Banerjee, U. C. (2001). Production, purification, characterization, and applications of lipases. *Biotechnology Advances*, 19, 627–662.
- Sharma, R., Soni, S. K., Vohra, R. M., Gupta, L. K., & Gupta, J. K. (2002). Purification and characterisation of a thermostable alkaline lipase from a new thermophilic *Bacillus* sp. RSJ-1. *Process Biochemistry*, 37(10), 1075–1084.
- Shi, Q. (2010). *Selection and Partial Characterization of Lipases From Raw Milk Bacterial Isolates for*. Master Thesis. University of Helsinki.
- Show, P. L., Tan, C. P., Anuar, M. S., Ariff, A., Yusof, Y. A., Chen, S. K., & Ling, T. C. (2012). Primary recovery of lipase derived from *Burkholderia cenocepacia* strain ST8 and recycling of phase components in an aqueous two-phase system. *Biochemical Engineering Journal*, 60, 74–80.
- Silva, W. O. B., Mitidieri, S., Schrank, A., & Vainstein, M. H. (2005). Production and extraction of extracellular lipase from the entomopathogenic fungus *Metarhizium anisopliae*. *Process Biochemistry*, 40, 321–326.
- Sivaramakrishnan, R., & Incharoensakdi, A. (2016). Purification and characterization of solvent tolerant lipase from *Bacillus* sp. for methyl ester production from algal oil. *Journal of Bioscience and Bioengineering*, 121(5), 517–522.
- Srinivas, N. D. (2000). *Aqueous Two-Phase Extraction For The Downstream Processing Of Enzymes*. PhD Thesis. University of Mysore.
- Sunna, A., Hunter, L., Hutton, C. A., & Bergquist, P. L. (2002). Biochemical characterization of a recombinant thermoalkalophilic lipase and assessment of its substrate enantioselectivity. *Enzyme and Microbial Technology*, 31(4), 472–476.
- Tamilarasan, K., & Kumar, M. D. (2012). Purification and characterization of solvent tolerant lipase from *Bacillus sphaericus* MTCC 7542. *Biocatalysis and Agricultural Biotechnology*, 1(4), 309–313.
- Tan, T., Zhang, M., Wang, B., Ying, C., & Deng, L. (2003). Screening of high lipase producing *Candida* sp. and production of lipase by fermentation. *Process Biochemistry*, 39(4), 459–465.
- Tanasupawat, S., Phoottosavako, M., & Keeratipibul, S. (2015). Characterization and lipolytic activity of lactic acid bacteria isolated from Thai fermented meat. *Journal of Applied Pharmaceutical Science*, 5(3), 6–12.

- Tayyab, M., Rashid, N., & Akhtar, M. (2011). Isolation and identification of lipase producing thermophilic *Geobacillus* sp. SBS-4S: cloning and characterization of the lipase. *Journal of Bioscience and Bioengineering*, 111(3), 272–278.
- Thakur, V., Tewari, R., & Sharma, R. (2014). Evaluation of production parameters for maximum lipase production by *P. stutzeri* MTCC 5618 and scale-up in bioreactor. *Chinese Journal of Biology*, 2014, 14.
- Toole, G., & Toole, S. (2004). *Essential AS biology for OCR*. Oxford, UK: Nelson Thornes.
- Tripathi, R., Singh, J., Bharti, R. K., & Thakur, I. S. (2014). Isolation, purification and characterization of lipase from *Microbacterium* sp. and its application in biodiesel production. *Energy Procedia*, 54, 518–529.
- Ulker, S., & Karaoglu, S. A. (2012). Purification and characterization of an extracellular lipase from *Mucor hiemalis* f. *corticola* isolated from soil. *Journal of Bioscience and Bioengineering*, 114(4), 385–390.
- Ungcharoenwiwat, P., & H-kittikun, A. (2015). Purification and characterization of lipase from *Burkholderia* sp. EQ3 isolated from wastewater from a canned fish factory and its application for the synthesis of wax esters. *Journal of Molecular Catalysis. B, Enzymatic*, 115, 96–104.
- Uppada, S. R. (2015). *Optimization Study Of Lipase From Lactic Acid Bacteria And Synthesis Of Flavor Esters*. PhD Thesis. Birla Institute of Technology and Science, Pilani.
- Wang, Y., Luo, D., Zhao, Y., Tian, S., Deng, W., Li, C., & Ma, L. (2017). High-level expression and characterization of solvent-tolerant lipase. *Journal of Bioscience and Bioengineering*, 1–7.
- Wu, Y. T., Pereira, M., Venâncio, A., & Teixeira, J. (2001). Separation of endopolygalacturonase using aqueous two-phase partitioning. *Journal of Chromatography A*, 929(1–2), 23–29.
- Xin, J., & Jiang, J. (2015). Lipase-Catalyzed Synthesis of S-Naproxenol oleins, (Emim), 236–239.
- Yadav, K. S., Adsul, M., Bastawde, K., Jadhav, D., Thulasiram, H., & Gokhale, D. (2011). Differential induction, purification and characterization of cold active lipase from *Yarrowia lipolytica* NCIM 3639. *Bioresource Technology*, 102(22), 10663–10670.
- Yang, W., He, Y., Xu, L., Zhang, H., & Yan, Y. (2016). A new extracellular thermo-solvent-stable lipase from *Burkholderia ubonensis* SL-4: identification, characterization and application for biodiesel production, 126, 76–89.

- Yen, Y. S. (2007). *Screening , Purification And Characterization Of Extracellular Lipase Produced By Pediococcus acidilactici UB6 Isolated From Malaysian Fermented Foods*. Master Thesis. Universiti Putra Malaysia.
- Yılmaz, D. E., & Sayar, N. A. (2015). Enzymatic organic solvent stable lipase from *Cryptococcus diffluens* D44 isolated from petroleum sludge. *Journal of Molecular Catalysis B: Enzymatic*, 122, 72–79.
- Zarevúcka, M. (2012). Olive Oil as Inductor of Microbial Lipase. In B. Dimitrios (Ed.), *Olive Oil - Constituents, Quality, Health Properties and Bioconversions* (pp. 457–470). InTech.
- Zhang, H., Zhang, F., & Li, Z. (2009). Gene analysis, optimized production and property of marine lipase from *Bacillus pumilus* B106 associated with South China Sea sponge *Halichondria rugosa*. *World Journal of Microbiology and Biotechnology*, 25(7), 1267–1274.
- Zheng, Y.-Y., Guo, X.-H., Song, N., & Li, D.-C. (2011). Thermophilic lipase from *Thermomyces lanuginosus*: gene cloning, expression and characterization. *Journal of Molecular Catalysis. B, Enzymatic*, 69(3–4), 127–132.