



**UNIVERSITI PUTRA MALAYSIA**

***ANTIMICROBIAL ACTIVITY, PHYTOCHEMICAL AND TOXICITY  
ANALYSES OF SALAM [*Syzygium polyanthum* (Wight) Walp.] LEAF  
EXTRACT AND ITS APPLICATION IN FOOD***

**SUZITA BINTI RAMLI**

**FSTM 2018 28**



**ANTIMICROBIAL ACTIVITY, PHYTOCHEMICAL AND TOXICITY  
ANALYSES OF SALAM [*Syzygium polyanthum* (Wight) Walp.] LEAF  
EXTRACT AND ITS APPLICATION IN FOOD**

By

**SUZITA BINTI RAMLI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia  
in Fulfillment of the Requirement for the Degree of Doctor of Philosophy**

**March 2018**

## COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Doctor of Philosophy

**ANTIMICROBIAL ACTIVITY, PHYTOCHEMICAL AND TOXICITY  
ANALYSES OF SALAM [*Syzygium polyanthum* (Wight) Walp.] LEAF  
EXTRACT AND ITS APPLICATION IN FOOD**

By

**SUZITA BINTI RAMLI**

**March 2018**

**Chairman : Associate Professor Yaya Rukayadi, PhD**

**Faculty : Food Science and Technology**

Food products can be subjected to contaminate by bacteria and fungi. The growth of this microorganisms in food products can cause foodborne illness. To overcome this problem, the prevention should be done at the early stage of food processing such as sanitizing. Commonly, chemicals sanitizer had been apply in food industry. However, application of this chemicals for long term was affected human health. Therefore, development of natural sanitizer derived from plant sources are gaining more attention nowadays. In this study, the antimicrobial activity of salam [*Syzygium polyanthum* (Wight) Walp.] leaves extract was evaluated against 17 types of pathogenic microorganisms including *Escherichia coli* O157:H7, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella* Typhimurium, *Staphylococcus aureus*, *Vibrio cholerae*, *Vibrio parahaemolyticus*, *Aspergillus flavus*, *Aspergillus niger*, *Rhizopus oligosporus*, *Rhizopus oryzae*, *Candida albicans*, *Candida krusei*, *Candida glabrata* and *Candida parapsilosis* in order to determine the ability of *S. polyanthum* as natural food sanitizer. The susceptibility test showed that all tested pathogenic bacteria were inhibited by *S. polyanthum* extract, with the range of inhibition zone between 6.67 to 9.67 mm. The extract could inhibit the growth of *L. monocytogenes* and *S. aureus* with MIC of 0.63 mg/mL, meanwhile MIC of the extract against others pathogens were 1.25 mg/mL. *L. monocytogenes* can be killed completely at MBC value of 0.63 mg/mL, *S. aureus*, *S. Typhimurium*, *V. cholerae* and *V. parahaemolyticus* were at 1.25 mg/mL, while the other strains were at MBC value of 2.5 mg/mL. Time-kill curve study showed that *E. coli* O157:H7, *L. monocytogenes*, *P. aeruginosa*, *S. aureus*, *V. cholerae* and *V. parahaemolyticus* can be killed by *S. polyanthum* extract at 4× MIC for 4 h, 4× MIC for 1 h, 4× MIC for 1 h, 4× MIC for 4 h, 4× MIC for 4 h, 4× MIC for 4 h, respectively. However, the population of *K. pneumoniae*, *P. mirabilis* and *S. Typhimurium* showed 3 Log reduction after treated at 4× MIC for 4 h. All filamentous fungi species showed 6.5 mm in inhibition zone, while 1.25 mg/mL and 5.0 mg/mL

for their MIC and MFC, respectively. In qualitative analysis of inhibition germination conidia, all the tested fungi species showed no growth after treated with extract started at  $1\times$  MIC for 9 days. Based on quantitative analysis, using  $4\times$  MIC for 24 h, the percentage of conidia germination were fully inhibited for *A. flavus* and *R. oryzae* (0%), meanwhile, for *R. oligosporus* and *A. niger* were reduced to 1% and 13%, respectively. Inhibition zone for *Candida* species were between 7.00-7.67 mm. For MIC and MBC, all *Candida* species can both inhibited and killed completely at range 0.63-1.25 mg/mL. *C. albicans*, *C. glabrata* and *C. parapsilosis* can be killed by *S. polyanthum* extract at  $4\times$  MIC for 4 h,  $4\times$  MIC for 2 h,  $4\times$  MIC for 4 h or  $2\times$  MIC for 4 h while the population of *C. krusei* reduced about 3 Log reduction after treated for  $4\times$  MIC for 4 h. Cell constituents release analysis and observation by using scanning electron microscope showed altering in cell wall linearity, cells ruptured and leaked of the cytoplasm. Generally, the antimicrobial activities of *S. polyanthum* extract were not affected by different pH and temperatures. GC-MS analysis identified the presence of active compounds which responsible to contribute antimicrobial properties in *S. polyanthum* extract included pyrogallol, phytol, hexadecanoic acid,  $\alpha$ -Tocopherol and  $\beta$ -Sitosterol while gallic acid, bergenin, quercetin 3-(6"-galloyl)galactoside), madecassic acid, quillaic acid and asiatic acid were detected by using LC-MS. The toxicity study by using brine shrimp assay demonstrated that *S. polyanthum*, extract was not toxic to *Artemia salina* with  $LC_{50}$  was 75.85 mg/mL. Generally the significant reduction of natural microflora in tested food samples were started at 0.50% (v/v) of extract at 5 min. During storage, 5% (v/v) showed better effect in controlling the microbial survival throughout the storage time. Physical characteristics in term of colour, texture and odour, all the treated samples until the highest concentration of extract [5% (v/v)] showed not significantly different and this finding also parallel to sensory acceptability where 5% of extract was accepted by the panelists. In conclusion, *S. polyanthum* extract exhibited antimicrobial activity, thus it can be developed as natural sanitizer for washing raw food materials and prevent the food spoilage during storage.

**Keyword:** Antimicrobial activity, *S. polyanthum* leaves, pathogenic microorganisms, toxicity, natural sanitizer.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**AKTIVITI ANTIMIKROBIAL, FITOKIMIA DAN TOKSISITI ANALISIS  
EKTRAK DAUN SALAM [*Syzygium polyanthum* (Wight) Walp.] DAN  
PENGUNAANNYA DI DALAM MAKANAN**

Oleh

**SUZITA BINTI RAMLI**

**Mac 2018**

**Pengerusi : Profesor Madya Yaya Rukayadi, PhD**  
**Fakulti : Sains dan Teknologi Makanan**

Produk makanan mempunyai risiko untuk terdedah dengan pencemaran bakteria dan kulat. Pertumbuhan mikroorganisma di dalam produk makanan boleh menyebabkan penyakit bawaan makanan. Untuk mengatasi masalah ini, langkah pencegahan harus dilakukan pada peringkat permulaan semasa pemprosesan makanan seperti mencuci dengan ejen pembersih. Kebiasaannya, ejen pembersih kimia digunakan di dalam industri makanan. Walaubagaimanapun, penggunaan bahan kimia ni untuk jangka masa yang panjang akan menjejaskan kesihatan manusia. Oleh itu, penghasilan ejen pembersih semulajadi daripada sumber tumbuh-tumbuhan semakin mendapat perhatian masa kini. Dalam kajian ini, aktiviti antimikrobial ekstrak daun salam [*Syzygium polyanthum* (Wight) Walp.] dinilai terhadap 17 jenis mikroorganisma patogenik termasuk *Escherichia coli* O157:H1, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella* Typhimurium, *Staphylococcus aureus*, *Vibrio cholerae*, *Vibrio parahaemolyticus*, *Aspergillus flavus*, *Aspergillus niger*, *Rhizopus oligosporus*, *Rhizopus oryzae*, *Candida albicans*, *Candida krusei*, *Candida glabrata* dan *Candida parapsilosis* untuk menentukan keupayaan ekstrak *S. polyanthum* sebagai ejen pembersih makanan semulajadi. Hasil kajian menunjukkan bahawa semua patogen bawaan makanan yang diuji, terencat apabila didedahkan kepada ekstrak *S. polyanthum*, dengan lingkungan zon perencatan antara 6.67 - 9.76 mm. Ekstrak *S. polyanthum* boleh menghalang pertumbuhan *L. monocytogenes* dan *S. aureus* dengan MIC, 0.63 mg/mL, sementara itu MIC ekstrak terhadap patogen yang lain adalah 1.25 mg/mL. *L. monocytogenes* boleh dibunuh sepenuhnya pada nilai MBC 0.63 mg/mL, *S. aureus*, *S. Typhimurium*, *V. cholerae* dan *V. parahaemolyticus* adalah 1.25 mg/mL, manakala patogen jenis lain adalah pada nilai 2.5 mg/mL. Analisis keluk-masa pembunuhan menunjukkan bahawa, *E. coli* O157: H7, *L. monocytogenes*, *P. aeruginosa*, *S. aureus*, *V. cholerae* dan *V. parahaemolyticus* boleh dibunuh oleh ekstrak *S. polyanthum* pada 4× MIC untuk 4 jam, 4× MIC untuk selama 1 jam, 4× MIC selama 1 jam, 4× MIC untuk 4 jam,

4× MIC untuk 4 jam, 4× MIC untuk 4 jam, masing-masing. Walau bagaimanapun, populasi *K. pneumoniae*, *P. mirabilis* dan *S. Typhimurium* hanya menurun sebanyak 3 Log selepas dirawat di 4× MIC untuk 4 jam. Semua spesies kulat menunjukkan 6.5 mm untuk zon perencatan, manakala 1.25 mg/mL dan 5.0 mg/mL untuk MIC dan MFC, masing-masing. Analisis kualitatif melalui pemerhatian visual untuk perencatan percambahan konidia, secara amnya, semua spesies kulat yang diuji menunjukkan tiada pertumbuhan selepas dirawat dengan ekstrak bermula pada 1× MIC selama 9 hari. Berdasarkan analisis kuantitatif, menggunakan 4× MIC selama 72 jam, peratusan percambahan konidia telah menurun kepada 0% untuk *A. flavus* dan *R. oryzae*, sedangkan percambahan *R. oligosporus* dan *A. niger* adalah 1% dan 13% masing-masing. Zon perencatan untuk *Candida* spp. adalah di antara 7.00-7.67 mm. Untuk MIC dan MBC, *Candida* spp. boleh direncat dan dibunuh sepenuhnya pada 0.63-1.25 mg/mL. *C. albicans*, *C. glabrata* dan *C. parapsilosis* boleh dibunuh oleh ekstrak *S. polyanthum* pada 4× MIC untuk 4 jam, 4× MIC untuk 2 jam, 4× MIC untuk 4 jam atau 2× MIC untuk 4 jam manakala populasi *C. krusei* hanya berkurang sebanyak 3 Log selepas dirawat selama 4× MIC untuk 4 jam. Analisis pelepasan sel dan pemerhatian menggunakan SEM pada wakil patogen yang dirawat dengan ekstrak *S. polyanthum* pada nilai MIC menunjukkan perubahan dalam kelicinan dinding sel, sel pecah dan kebocoran sitoplasma. Secara umumnya, aktiviti antimikrobial ekstrak *S. polyanthum* tidak terjejas dengan ketara oleh pelbagai nilai pH dan suhu. Analisis GC-MS mengenal pasti kehadiran sebatian aktif yang menjadikan *S. polyanthum* bersifat antimikrobial termasuk pyrogallol, phytol, asid hexadecanoic,  $\alpha$ -Tocopherol dan  $\beta$ -Sitosterol sementara asid gallic, bergenin, quercetin 3-(6"-galloyl)galactoside, asid madecassic, asid quillaic and asid asiatic telah dikenal pasti menggunakan LC-MS. Kajian menunjukkan bahawa ekstrak *S. polyanthum* tidak toksik kepada *Artemia salina* dengan nilai  $LC_{50}$  ialah 75.85 mg/mL. Umumnya, penurunan populasi mikroflora semulajadi bermula pada ekstrak kepekatan 0.50% (v/v) dalam tempoh 5 min. Untuk menyimpan makanan, 5% (v/v) ekstrak telah menunjukkan kesan terhadap perencatan pertumbuhan mikroorganisma. Ciri-ciri fizikal dari segi warna, tekstur dan bau, sampel makanan yang dirawat dengan ekstrak yang berkepekatan paling tinggi [5% (v/v)] menunjukkan tidak signifikan dan ini selari dengan penerimaan ahli panel semasa saringan deria. Kesimpulannya, ekstrak *S. polyanthum* mempunyai aktiviti antimikrobial, oleh itu ia boleh dibangunkan sebagai ejen pembasuh semula jadi untuk membasuh bahan makanan mentah dan untuk mencegah kerosakan semasa penyimpanan makanan.

**Kata kunci:** Aktiviti antimikrobial, *S. polyanthum*, mikroorganisma patogenik, toksisiti, ejen pembasuh semulajadi.

## ACKNOWLEDGEMENTS

Alhamdulillah, thank you Allah, for the strength that He has given to me, for the wisdom that He granted me, for the unconditional love that He shown me until I am able to persue and completed my Doctor of Philosophy. Without You, I would never have the perserverance to make it until the end.

I would like to dedicate my heartfelt to Prof. Madya Dr. Yaya Rukayadi as my supervisor, for the guidance, encouragement and constructive suggestions during the entire study. A millions thanks to Prof. Dr. Son Radu and Prof. Dr. Khozirah Shaari, who has contributed tremendously of their time and expertise in this research.

Sincere thanks and appreciations are extended to all support staff of Microbiology and Biochemistry Laboratory, Faculty of Food Science and Technology for their assistants during my laboratory work. Thousands thanks also goes to my lab mates for their helping and support throughout my study.

Words could not describe my deepest appreciation to my mum, Mrs. Fatimah Mohd Jasin and all my family members for their love, encouragements and understandings. Millions thanks extended to my lovely husband, Mr. Roslan Ramli, who always be my side along my study journey. Not forgotten to my late father, Mr. Ramli Martin, your loving and encouragement always in my mind. Lastly, thank you very much again to all of you that had been involved in my research. I will remember you all for the rest of my life.



I certify that a Thesis Examination Committee has met on 15 March 2018 to conduct the final examination of Suzita binti Ramli on her thesis entitled "Antimicrobial Activity, Phytochemical and Toxicity Analyses of Salam [*Syzygium polyanthum* (Wight) Walp.] Leaf Extract and its Application in Food" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

**Azizah binti Abdul Hamid, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Rosfarizan binti Mohamad, PhD**

Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Muhajir bin Hamid, PhD**

Associate Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Kalidas Shetty, PhD**

Professor  
North Dakota State University  
United States  
(External Examiner)



---

**NOR AINI AB. SHUKOR, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 24 May 2018

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Yaya Rukayadi, PhD**

Associate Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Son Radu, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Member)

**Khozirah binti Shaari, PhD**

Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Name and Matric No.: Suzita binti Ramli, GS 43671

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of  
Chairman of  
Supervisory  
Committee: Associate Professor Dr. Yaya Rukayadi

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: Professor Dr. Son Radu

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: Professor Dr. Khozilah binti Shaari

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiv
<b>LIST OF FIGURES</b>	xvi
<b>LIST OF ABBREVIATIONS</b>	xix
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
1.1 Background	1
1.2 Problem statements	2
1.3 Objectives	2
<b>2 LITERATURE REVIEW</b>	<b>4</b>
2.1 Foodborne disease	4
2.2 Foodborne pathogens and food spoilage microorganisms	4
2.3 Sources of foodborne pathogens and food spoilage microorganisms	5
2.3.1 Poultry	6
2.3.2 Seafood	6
2.3.3 Dairy product	7
2.3.4 Fruit and vegetables	8
2.4 Pathogenic microorganisms associated with food	9
2.5 Food preservatives	17
2.6 Food sanitizer	18
2.7 Bioactive compounds responsible in antimicrobial activity	19
2.7.1 Eugenol	19
2.7.2 Citral	20
2.7.3 Tannin	20
2.7.4 Flavonoid	20
2.8 Medicinal plant	22
2.9 <i>Syzygium polyanthum</i> (Wight) Walp.	23
2.9.1 Biology of <i>S. polyanthum</i>	23
2.9.2 Traditional uses	25
2.9.3 Phytochemical properties	25
2.9.4 Biological properties	25
2.10 Safety and toxicity of medicinal plant	26

<b>3</b>	<b>MATERIALS AND METHODS</b>	<b>28</b>
3.1	Plant sample collection	28
3.2	Preparation of crude leaves extract	28
3.3	Microbial strains and maintenance	29
3.4	Preparation of media and chemicals	30
3.4.1	Preparation of microbiological agar/broth	30
3.4.2	Preparation of phosphate buffer saline	30
3.4.3	Preparation of chlorhexidine	31
3.4.4	Preparation of amphotericin B	31
3.4.5	Preparation of dimethyl sulfoxide (DMSO)	31
3.4.6	Preparation of antimicrobial agents	31
3.4.7	Preparation of inoculums	31
3.5	Disc diffusion test	32
3.6	Minimum inhibition concentration (MIC)	32
3.7	Minimum bactericidal concentration/ minimum fungicidal concentration (MBC/MFC)	33
3.8	Time kill curve assay	33
3.9	Inhibition conidial germination assay	33
3.9.1	Qualitative analysis	33
3.9.2	Quantitative analysis	34
3.10	Scanning electron microscope (SEM)	34
3.11	Cell constituents' release analysis	34
3.12	Stability of <i>S. polyanthum</i> extract at different pH and temperature	35
3.13	Identification and quantification of active compounds in <i>S. polyanthum</i> extract	35
3.13.1	Gas chromatography-mass spectrometry (GC-MS)	35
3.13.2	Liquid chromatography-mass spectrometry (LC-MS)	35
3.14	Toxicity analysis of <i>S. polyanthum</i> extract	36
3.14.1	Hatching of brine shrimp ( <i>Artemia salina</i> sp.)	36
3.14.2	Brine shrimp lethality assay	36
3.15	Effect of <i>S. polyanthum</i> extract to natural microorganisms in raw food samples	36
3.15.1	Raw food materials	36
3.15.2	Preparation of <i>S. polyanthum</i> extract for treatment solutions	37
3.15.3	Preparation of selective media	37
3.15.4	Washing treatment of raw food materials with <i>S. polyanthum</i> extract solutions	37
3.15.5	Effect of <i>S. polyanthum</i> extract washing treatment at different storage	37
3.15.6	Microbiology analysis	38
3.15.7	Colour analysis	38
3.15.8	Texture analysis	38
3.16	Evaluation of sensory attributes acceptability of treated raw food materials	39
3.17	Statistical analysis	39

<b>4</b>	<b>RESULTS AND DISCUSSION</b>	<b>40</b>
4.1	Yield of extract	40
4.2	Antimicrobial activity of <i>S. polyanthum</i> extract against pathogenic microorganism	41
4.2.1	Antibacterial activity of <i>S. polyanthum</i> extract against pathogenic bacteria	41
4.2.1.1	Disc diffusion test	41
4.2.1.2	Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC)	43
4.2.1.3	Time-kill curve analysis of <i>S. polyanthum</i> extract on pathogenic bacteria	45
4.2.2	Antifungal activity of <i>S. polyanthum</i> extract against food spoilage fungi	49
4.2.2.1	Disc diffusion test	49
4.2.2.2	Minimum inhibition concentration (MIC) and minimum fungicidal concentration (MFC)	51
4.2.2.3	Inhibition conidia germination	52
4.2.2.4	Time kill curve of <i>Candida</i> species	59
4.3	Effect of <i>S. polyanthum</i> extract on the viability and cellular injury to the structures of <i>K. pneumoniae</i> , <i>S. aureus</i> , <i>A. niger</i> and <i>C. albicans</i> .	62
4.3.1	Morphology changes observation by using scanning electron microscope (SEM)	62
4.3.1.1	<i>Klebsiella pneumoniae</i>	62
4.3.1.2	<i>Staphylococcus aureus</i>	62
4.3.1.3	<i>Aspergillus niger</i>	65
4.3.1.4	<i>Candida albicans</i>	67
4.3.2	Cell constituents' releases analysis	69
4.4	Stability of <i>S. polyanthum</i> extract at different pH and temperature	71
4.4.1	Stability of <i>S. polyanthum</i> extract at different pH	71
4.4.2	Stability of <i>S. polyanthum</i> extract at different temperatures	73
4.5	Determination of phytochemical compounds in <i>S. polyanthum</i> extract	75
4.5.1	Gas Chromatography-Mass Spectrometry (GC-MS) based profile of bioactive compounds in <i>S. polyanthum</i> extract	75
4.5.2	Liquid Chromatography-Mass Spectrometry (LC-MS) based profile of bioactive compounds in <i>S. polyanthum</i> extract	84
4.6	Toxicity of <i>S. polyanthum</i> extract on <i>Artemia salina</i>	90
4.7	Application of <i>S. polyanthum</i> extract as natural sanitizer on raw food materials	93
4.7.1	Natural microbial populations detected in chicken meat, shrimp, cherry tomato and grapes.	93

4.7.2	Effect of different concentration and exposure time of <i>S. polyanthum</i> extract on the number of natural microbial populations in raw food materials by soaking treatment	96
4.7.2.1	Microbiology analysis	96
4.7.2.2	Colour analysis	104
4.7.2.3	Texture analysis	112
4.7.3	Effect of soaking treatment with <i>S. polyanthum</i> extract on natural flora of food samples at different storage time.	114
4.7.3.1	Chicken meat	115
4.7.3.2	Shrimp	118
4.7.3.3	Cherry tomato	120
4.7.3.4	Grapes	123
4.8	Sensory attribute evaluation of treated chicken meat, shrimp, cherry tomato and grapes with <i>S. polyanthum</i> extract	127
<b>5</b>	<b>SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	131
5.1	Summary	131
5.2	Conclusion	132
5.3	Recommendations for future research	133
	<b>REFERENCES</b>	134
	<b>APPENDICES</b>	173
	<b>BIODATA OF STUDENT</b>	182
	<b>LIST OF PUBLICATIONS</b>	183



## LIST OF TABLES

Table		Page
2.1	Pathogenic microorganisms associated with food	11
2.2	Classification of <i>S. polyanthum</i>	24
3.1	Pathogenic bacteria strains	29
3.2	Food spoilage fungi strains	29
3.3	List of media	30
4.1	Total yield of <i>S. polyanthum</i> extract	40
4.2	Inhibition zone of <i>S. polyanthum</i> leaves extract against pathogenic bacteria	43
4.3	Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of <i>S. polyanthum</i> extract against pathogenic bacteria	44
4.4	Time kill curve assay of pathogenic bacteria	45
4.5	Inhibition zone of <i>S. polyanthum</i> extract against food spoilage fungi	50
4.6	Minimum inhibitory concentration (MIC) and minimum fungicidal concentration (MFC) of <i>S. polyanthum</i> L. extract against food spoilage fungi and <i>Candida</i> species	51
4.7	Time kill curve analysis of <i>Candida</i> species	59
4.8	Stability of <i>S. polyanthum</i> extract at different pH on the MIC and MBC of pathogenic bacteria	72
4.9	Stability of <i>S. polyanthum</i> extract at different pH on the MIC and MFC of food spoilage fungi	73
4.10	Stability of <i>S. polyanthum</i> extract at different temperatures on the MIC and MBC of pathogenic bacteria	74
4.11	Stability of <i>S. polyanthum</i> extract at different temperatures on the MIC and MFC of food spoilage fungi	75
4.12	Identification of phytochemical compounds in <i>S. polyanthum</i> leaves extract by using GC-MS	78
4.13	Bioactive compounds which had antimicrobial properties in <i>S. polyanthum</i> extract	79
4.14	Identification of phytochemical compounds in <i>S. polyanthum</i> leaves extract by using LC-MS	86
4.15	Identification of bioactive compounds which had antimicrobial properties in <i>S. polyanthum</i> extract using LC-MS	88

4.16	Toxicity of <i>S. polyanthum</i> extract using brine shrimp lethality assay	92
4.17	Total plate count and number of natural microbial detected in raw food samples using different selective media agar (Log <sub>10</sub> CFU/mL)	95
4.18	Effects of different concentrations and exposure times of <i>S. polyanthum</i> extract on natural microbial in chicken meat (Log <sub>10</sub> CFU/mL)	98
4.19	Effects of different concentrations and exposure times of <i>S. polyanthum</i> extract on natural microbial in shrimp (Log <sub>10</sub> CFU/mL)	100
4.20	Effects of different concentration and exposure time of <i>S. polyanthum</i> extract on natural microbial in cherry tomato (Log <sub>10</sub> CFU/mL)	102
4.21	Effects of different concentrations and exposure times of <i>S. polyanthum</i> extract on natural microbial in grapes (Log <sub>10</sub>	103
4.22	Colour analysis on effect of different concentrations of <i>S. polyanthum</i> extract on the chicken meat	106
4.23	Colour analysis on effect of different concentrations of <i>S. polyanthum</i> extract on shrimp	107
4.24	Colour analysis on effect of different concentrations of <i>S. polyanthum</i> extract on the cherry tomato	109
4.25	Colour analysis on effect of different concentrations of <i>S. polyanthum</i> extract on grapes	111
4.26	Texture analysis on effect of different concentrations of <i>S. polyanthum</i> extract on chicken meat	112
4.27	Texture analysis on effect of different concentrations of <i>S. polyanthum</i> extract on shrimp	113
4.28	Texture analysis on effect of different concentrations of <i>S. polyanthum</i> extract on cherry tomato	113
4.29	Texture analysis on effect of different concentrations of <i>S. polyanthum</i> extract on grapes	114
4.30	Sensory attributes acceptability of treated chicken meat with <i>S. polyanthum</i> extract	127
4.31	Sensory attributes acceptability of treated shrimp with <i>S. polyanthum</i> extract	128
4.32	Sensory attributes acceptability of treated cherry tomato with <i>S. polyanthum</i> extract	129
4.33	Sensory attributes acceptability of treated grapes with <i>S. polyanthum</i> extract	129

## LIST OF FIGURES

Figure	Page
2.1	23
2.2	24
3.1	28
4.1	49
4.2	53
4.3	54
4.4	55
4.5	56
4.6	58
4.7	61
4.8	63
4.9	64

4.10	Scanning electron micrograph of <i>Asp. niger</i> [(a), (c) non-treated; (b), (d), after treated with <i>S. polyanthum</i> extract for 24 h]	66
4.11	Scanning electron micrograph of <i>C. albicans</i> [(a) non-treated; (b) after treated with <i>S. polyanthum</i> extract for 24 h]	68
4.12	Cell constituent release analysis of (a) <i>K. pneumoniae</i> , (b) <i>S. aureus</i> , (c) <i>A. niger</i> and (d) <i>C. albicans</i> after treated with <i>S. polyanthum</i> extract at concentration of 0× MIC, 1× MIC, 2× MIC and 4× MIC, respectively.	70
4.13	Chomatogram of phytochemical compounds of <i>S. polyanthum</i> extract	77
4.14	Mass fragmentation pattern of pyrogallol (above), compared with reference (below)	80
4.15	Mass fragmentation pattern of hexadecanoic acid (above), compared with reference (below)	81
4.16	Mass fragmentation pattern of phytol (above), compared with reference (below)	82
4.17	Mass fragmentation pattern of α-Tocopherol (above), compared with reference (below)	83
4.18	Mass fragmentation pattern of β-Sitosterol (above), compared with reference (below)	84
4.19	Chromatogram of compounds in <i>S. polyanthum</i> extract in positive ion mode	85
4.20	Chromatogram of compounds in <i>S. polyanthum</i> extract in negative ion mode	85
4.21	Standard curve of brine shrimp lethality assay after treated with <i>S. polyanthum</i> extract for 24 h	91
4.22	Standard curve of brine shrimp lethality assay after treated with potassium dichromate for 24 h	92
4.23	Representative the (a) total bacterial, (b) yeast and mold and (c) <i>E. coli</i> counts on the chicken meat exposed to different concentration of <i>S. polyanthum</i> extract during storage at temperature $4 \pm 2^\circ\text{C}$ for 5 days. (■) Control, (◆) 1% extract and (▲) 5% extract	116
4.24	Representative the (a) total bacterial, (b) yeast and mold and (c) <i>E. coli</i> counts on the chicken meat exposed to different concentration of <i>S. polyanthum</i> extract during storage at temperature $-18 \pm 2^\circ\text{C}$ for 21 days (■) Control, (◆) 1% extract and (▲) 5% extract	117
4.25	Representative the (a) total bacterial, (b) yeast and mold, (c) <i>E. coli</i> and (d) <i>Vibrio</i> spp. counts on the shrimp exposed to different concentration of <i>S. polyanthum</i> extract during storage at temperature $4 \pm 2^\circ\text{C}$ for 5 days (■) Control, (◆) 1% extract and (▲) 5% extract	118

4.26	Representative the (a) total bacterial, (b) yeast and mold, (c) <i>E. coli</i> and (d) <i>Vibrio</i> spp. counts on the shrimp exposed to different concentration of <i>S. polyanthum</i> extract during storage at temperature $-18 \pm 2^{\circ}\text{C}$ for 21 days (■) Control, (◆) 1% extract and (▲) 5% extract	119
4.27	Representative the (a) total bacterial, (b) yeast and mold, (c) <i>E. coli</i> and (d) <i>Pseudomonas</i> spp. counts on cherry tomato exposed to different concentration of <i>S. polyanthum</i> extract during storage at room temperature $27 \pm 2^{\circ}\text{C}$ for 5 days (■) Control, (◆) 1% extract and (▲) 5% extract	121
4.28	Representative the (a) total bacterial, (b) yeast and mold, (c) <i>E. coli</i> and (d) <i>Pseudomonas</i> spp. counts on cherry tomato exposed to different concentration of <i>S. polyanthum</i> extract during storage at $4 \pm 2^{\circ}\text{C}$ for 15 days (■) Control, (◆) 1% extract and (▲) 5% extract	122
4.29	Representative the (a) total bacterial, (b) yeast and mold, (c) <i>E. coli</i> and (d) <i>S. aureus</i> counts on grapes exposed to different concentration of <i>S. polyanthum</i> extract during storage at room temperature $27 \pm 2^{\circ}\text{C}$ for 5 days (■) Control, (◆) 1% extract and (▲) 5% extract	123
4.30	Representative the (a) total bacterial, (b) yeast and mold, (c) <i>E. coli</i> and (d) <i>S. aureus</i> counts on grapes exposed to different concentration of <i>S. polyanthum</i> extract during storage at $4 \pm 2^{\circ}\text{C}$ for 15 days (■) Control, (◆) 1% extract and (▲) 5% extract	125

## LIST OF ABBREVIATIONS

ATCC	American Type Culture Collection
ANOVA	Analysis of variance
BP	Baird-Parker
CFU	Colony forming unit
CHX	Chlorhexidine
CLSI	Clinical and Laboratory Standards Institute
DMSO	Dimethylsulfoxide
GC-MS	Gas Chromatography – Mass Spectrometry
GRAS	Generally Regarded as Safe
IBS	Institute of Bioscience
LC	Lethality concentration
LC-MS	Liquid Chromatography – Mass Spectrometry
MBC	Minimum Bactericidal Concentration
MFC	Minimum Fungicidal Concentration
MHB	Mueller Hinton broth
MIC	Minimum Inhibitory Concentration
MOH	Ministry of Health
m/z	Mass/charge ratio
NIST	National Institute of Standards and Technology
OD	Optical density
PBS	Phosphate buffered saline
PCA	Plate Count Agar
PDA	Potato Dextrose Agar
PDB	Potato Dextrose Broth
Ppm	parts per million
Rpm	Revolutions per minute
SDA	Sabouraud Agar
SDB	Sabouraud Broth
SEM	Scanning Electron Microscopy
spp.	Species
TCBS	Thiosulfate citrate bile salts sucrose

UPM

Universiti Putra Malaysia

WHO

World Health organization

XLD

Xylose lysine deoxylate



# CHAPTER 1

## INTRODUCTION

### 1.1 Background

The problem of foodborne pathogens is one of major public concerns in both developed and developing countries worldwide. They become a threat to food safety especially in developing countries with poor hygiene and sanitation facilities. The rise of foodborne pathogens are responsible for millions of infectious gastrointestinal disease cases each year which increase the cost of medical care and causing loss of productivity (Bloomfield et al., 2007). Symptoms of foodborne diseases were include nausea, vomit, diarrhea and fever, whereas severe foodborne diseases may lead to death (Gandhi and Chikindas, 2007).

In France, foodborne incidence was 1210 cases per 100,000 inhabitants, meanwhile in United Kingdom and United State were 2600 cases and more than 25,000 cases per 100,000 inhabitants, respectively (Teisl and Roe, 2010). Besides that, in Australia, about 5.4 million cases were reported with 15,000 people got hospitalized and 120 deaths annually (Soon et al., 2011). In the United States, the reported foodborne cases have shown an increase of approximately 48 million illnesses with 128,000 hospitalizations while death cases was about 3,000 each year (Scallan et al., 2011). In Malaysia, the incidence rate of reported food poisoning in 2013 is 47.79 per 100,000 populations, with mortality rate of 0.04% (MOH, 2014). There also food poisoning cases among pupils in school has increased by 57% in four months in 2016 compared to 2015 (MOH, 2016).

Generally, foodborne illnesses occur are due to improper food handling and poor hygienic practices, especially during food preparation and storage period (Scott, 2003). Common examples of widely reported foodborne pathogens include *Bacillus cereus*, *Campylobacter*, *Clostridium*, *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Salmonella* and *Staphylococcus aureus* (Singh et al., 2003). In addition, deterioration of food produce by spoilage fungi at various levels of raw food processing and production affect the supply chains of agri-food supply to the customers. Its has been estimated 5-10% of food lossess were caused by fungal food spoilage (Pitt and Hocking, 2009). Various reports claimed *Alternaria* spp., *Aspergillus* spp., *Botrytis cinerea*, *Cladosporium* spp., *Fusarium* spp., *Penicillium* spp., and *Rhizopus* spp. related to the spoilage of perishable products (Tournas, 2005; Pitt and Hocking, 2009; Birhanu et al., 2014; Khokhar and Bajwa, 2014).

Food contamination can occur during post-harvest processing or even earlier which is during pre-harvesting period due to the contamination from a farm environment, water irrigation systems, manure and so on (Doyle and Erickson, 2006). Without proper



decontamination techniques, pathogens will keep on growing especially during food storage period and during food preparation caused by cross contamination (Scott, 2003).

## 1.2 Problem statements

Eventhough, there are many food decontamination treatments and also food preservatives that have been applied to eliminate or substantially decrease bacterial populations both in fresh produce and raw poultries including salt, sugar, chlorine, organic acids, hydrogen peroxide, ozonation and irradiation (Leistner, 2002), however, there was important to eliminate or reduce the microbial population at the early stage of food processing. Therefore, washing the food materials with any food sanitizer was crucial especially for ready to eat food such as fruits and vegetables. There was several chemical sanitizer that had been applied in food industry including chlorine, hypochlorite, iodine, quaternary ammonium, hydrogen proxide and fatty acid sanitizer. However, this chemical can be harmful to health and also to the environment (Neo et al., 2013). In addition, washing with tap or chlorinated water is the common decontamination practice used at home. However, this practice is being questioned due to the formation of carcinogenic by-products such as trihalomethanes when chlorine reacts with organic matter such as decayed leaves (Chang and Fang, 2007). Besides that, organic substance in foods can be oxidized by chlorine whereas in water carcinogenic and mutagenic by product such as haloforms and haloacetic acids can be formed (Perez-Gregorio et al., 2011).

Due to disadvantage of chemical food sanitizer, study on natural antimicrobial agents from plants is the alternative way. Furthermore, commercial natural food sanitizer is still scarce. Plant antimicrobials are gaining wide interest because most of them are classified as Generally Recognized as Safe (GRAS) status which has been discovered to have higher levels of food safety (Alzarokay and Nakahara, 2003). Many studies have reported that most of the medicinal plants, including spices and herbs have strong antimicrobial compounds (Limsuwan et al., 2009).

## 1.3 Objectives

A study was undertaken to discover the antimicrobial activity of *Syzygium polyanthum* extract against several types of pathogenic microorganisms and also to measure the ability of this plant extract as natural food sanitizer. The specific objectives were:

1. To determine antimicrobial activity of *S. polyanthum* leaves extract against pathogenic microorganisms.
2. To analyse modes of action of *S. polyanthum* leaves extract by cell constituents release analysis and observation using scanning electron microscope (SEM).

3. To determine the stability of *S. polyanthum* extract at different pHs and temperatures.
4. To identify the present of bioactive compounds in *S. polyanthum* leaves extract by Gas Chromatography Mass - Spectrometry (GC-MS) Liquid Chromatography Mass - Spectrometry (LC-MS).
5. To determine toxicity of *S. polyanthum* leaves extract by using brine shrimp (*Artemia salina*) lethality assay.
6. To examine the effect of *S. polyanthum* leaves extract on natural microflora in food samples at different concentrations of sanitizer solution and different storage times and also its sensory attributes acceptability.



## REFERENCES

- Abadias, M., Canamas, T.P., Asensio, A., Anguera, M. and Vinas, I. (2006). Microbial quality of commercial 'Golden Delicious' apples throughout production and shelf-life of Lleda (Catalonia, Spain). *International Journal of Food Microbiology*. 108: 404-409.
- Abdumumeen, H. A., Risikat, A. N. and Sururah, A. R. (2012). Food: Its preservatives, additives and applications. *International Journal of Chemical and Biochemical Sciences*. 1:36-47.
- Abd Rahim, E. N. A., Ismail, A., Omar, M. N., Rahmat, U. N. and Wan Ahmad, W. A. N. (2018). GC-MS analysis of phytochemical compounds in *Syzygium polyanthum* leaves extracted using ultrasound-assisted method. *Pharmacognosy Journal*. 10(1):110-119.
- Abdullah, E., Raus, R.A. and Jamal, P. (2012). Extraction and evaluation of antibacterial activity from selected flowering plants. *American Medical Journal*. 3(1): 27-32.
- Abee, T., Krockel, L. and Hill, C. (1995). Bacteriocins: modes of action and potentials in food preservation and control of food poisoning. *International Journal of Food Microbiology*. 28(2): 169-185.
- Adcock H. Pharmageddon. (2002). Is it too late to tackle growing resistance to anti-infectives? *Pharmacology Journal*. 269: 59-60.
- Addgene. (2014). Creating bacterial glycerol stocks for long-term storage of plasmids. Retrieved 28 May 2017 from <https://www.addgene.org/plasmids-protocols/create-glycerol-stocks/>.
- Afolayan, A.J. and Meyer, J.J. (1997). The antimicrobial activity of 3, 5, 7-trihydroxyflavone isolated from the shoots of *Helichrysum aureonitens*. *Journal of Ethnopharmacology*. 57: 177-81.
- Ahlawat, K. S., Gulia, N. and Khatkar, B. S. (2013). Effect of plant maturity on leaf growth, yield and physicochemical properties of aloe vera. *Food analysis*. 20(1): 22-24.
- Ahmad, I. and Beg, A.Z. (2001). Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *Journal of Ethnopharmacology*. 74: 113-123.
- Ahmed, A., Labu, Z.K., Dey, S.K., Hira, A., Howlader, M.S.I., Mohamed Hemayet Hossain, M.H., Roy, J. (2015). Phytochemical screening, antibacterial and cytotoxic activity of different fractions of *Xylocarpus mekongensis* Bark. *Ibnosina Journal of Medicine and Biomedical Sciences*. 5: 206-213.

- Akbar, A. and Anal, A.K. (2013). Prevalence and antibiogram study of *Salmonella* and *Staphylococcus aureus* in poultry meat. *Asian Pasific Journal of Tropical Biomedicine*. 3(2): 163-168.
- Akbar, A. and Anal, A. K (2011). Food safety concerns and food-borne pathogens, *Salmonella*, *Escherichia coli* and *Campylobacter*. *Journal of Biology*. 1(1): 5-17.
- Akhtar, N., Anjum, T. and Jabeen. R. (2013). Isolation and identification of storage fungi from citrus sampled from major growing areas of Punjab, Pakistan. *International Journal of Agriculture and Biology*. 15(6): 1283-1288.
- Akhter, H. and Khan S. A. (2012). Effect of gamma irradiation on the quality (colour, firmness and total soluble solid) of tomato (*Lycopersian esculentum* Mil.) stored at different temperature. *Asian Journal of Agricultural Research*. 6(1): 12-20.
- Alastruey-Izquierdo, A., Melhem, M. S. C., Bonfietti, L. X. and rodriguez-tudela, J. (2015). Susceptibility test for fungi: clinical and laboratorial correlations in medical mycology. *Journal of the Sao Paulo Institute of Tropical Medicine*. 57(19): 57-64.
- Al-Dughaym, A. M. and Altabari. A. F. (2010). Safety and quality of some chicken meat products in Al-Ahsa markets-Saudi Arabia. *Saudi Journal of Biological Sciences*. 17: 37-42.
- Al-Shammari, L.A., Hassan, W.H.B. and Al-Youssef, H.M. (2012). Chemical composition and antimicrobial activity of the essential oil and lipid content of *Carduus pycnocephalus* L. growing in Saudi Arabia. *Journal of Chemical and Pharmaceutical Research*. 4(2): 1281-1287.
- Ali, A.M., Mooi, L.Y., Yih, K.Y., Norhanom, A.W., Saleh, K.M., Lajis, N.H., Yazid, A.M.; Ahmad, F.B.H. and Prasad, U. (2000). Anti-tumor promoting activity of some Malaysian traditional vegetable (*Ulam*) extracts by immunoblotting analysis of raji cells. *Natural Product Sciences*. 6(3): 147-150.
- Alvarez-Ordóñez, A., Fernández, A., Bernardo, A. and Lopez, M. (2010). Acid tolerance in *Salmonella* Typhimurium induced by culturing in the presence of organic acids at different growth temperatures. *Food Microbiology*. 27: 44-49.
- Alves, T.M.A., Silva, A.F., Brandao, M., Grandi, T.S.M., Smania, E.F., Smania Jr., A. and Zani, C.L. (2000). Biological screening of Brazilian medicinal plants. *Memorias do Instituto Oswaldo Cruz*. 95: 367-373.
- Alwash, M.S., Ibrahim, N. and Ahmad, W.Y. (2013). Identification and mode of action of antibacterial components from *Melastoma malabathricum* Linn. leaves. *American Journal of Infectious Diseases*. 9(2): 46-58.

- Alzarokey, N.S. and Nakahara, K. (2003). Antibacterial activity of extracts from some edible plants commonly consumed in Asia. *International Journal of Food Microbiology*. 80: 223-230.
- Andrews, J.M. (2001). Determination of minimum inhibitory concentration. *Journal of Antimicrobial Chemotherapy*. 1: 5-16.
- Anees, A. M., Ravi, S. and Ghogare, P. (2015). Studies on antimicrobial activity of spices and effect of temperature and pH on its antimicrobial properties. *IOSR Journal of Pharmacy and Biological Sciences*. 10(1): 99-102.
- Anna, M., Dara, K., Hickey, I., Mercedes, A.G. and Martin, W. (2013). Evaluation of Antimicrobial activities of commercial herb and spice extracts against selected food-borne bacteria. *Journal of Food Research*. 2: 431-437.
- Annegowda, H.V., Bhat, R., Min-Tze, L., Karim, A.A. and Mansor, S.M. (2012). Influence of sonication treatments and extraction solvents on the phenolics and antioxidants in star fruits. *Journal of Food Science and Technology*. 49(4): 510-514.
- Arabshahi-D.S., Vishalakshi Devi, D. and Urooj, A. (2007). Evaluation of antioxidant activity of some plant extracts and their heat, pH and storage stability. *Food Chemistry*. 100: 1100-1105.
- Archer, D.L. (2004). Freezing: an underutilized food safety technology?. *International Journal of Food Microbiology*. 90: 127-138.
- Arif, K. M. T., Datta, R. K., Golam Sarower, M. and Hossain, S. J. (2014). Comparison of phenolics and antibacterial activity of commonly used antidiabetic medicinal plants in Bangladesh. *Pharmacology Online*. 1: 7-15.
- Ariviani, S. (2010). Pengaruh diet ekstrak buah salam [*Syzygium polyanthum* (wight.) walp] terhadap kadar glukosa darah dan profil lipid (total kolesterol, HDL-C, LDL-C, dan trigliserida) pada tikus wistar. *Jurnal Teknologi Hasil Pertanian*, 3(1): 21-27.
- Arslanyolu, M. and Erdemgil, F. Z. (2006). Evaluation of the antibacterial activity and toxicity of isolated arctiin from the seeds of *Centaurea sclerolepis*. *Journal of Pharmacy and Pharmacology*. 35: 103-109.
- Asao, T., Kumeda, Y. and Kawai (2003). An extensive outbreak of staphylococcal food poisoning due to low-fat milk in Japan: estimation of enterotoxin A in the incriminated milk and powdered skim milk. *Epidemiology and Infection*. 130: 33-40.
- Asuquo, E G. and Udobi, C. E. (2016). Antibacterial and toxicity studies of the ethanol extract of *Musa paradisiaca* leaf. *Microbiology, Parasitology and Virology*. 2: 1-10.

- Atanda, O.O., Akpan, I. and Oluwafemi, F. (2007). The potential of some spice essential oils in the control of *Aspergillus parasiticus* CFR 223 and aflatoxin production. *Food Control*. 18: 601-607.
- Atanassova, V. and C. Ring. (1999). Prevalence of *Campylobacter* spp. in poultry and poultry meat in Germany. *International Journal of Food Microbiology*. 51: 187-190.
- Aydogdu, H. and Gucer, Y. (2009). Microfungi and mycotoxins of grapes and grape products. *Trakia Journal of Sciences*. 7(2): 211-214.
- Ayyanar, M. and Subash-Babu, P. (2012). *Syzygium cumini* (L.) Skeels: A review of its phytochemical constituents and traditional uses. *Asian Pacific Journal of Tropical Biomedicine*. 2(3): 240-246.
- Bacteriological Analytical Manual (BAM). (1998). U. S. Food and Drug Administration. 8th Edition, Revision A.
- Baird-Parker, A. C. (1994). Foods and microbiological risks. *Microbiology*. 140: 687-695.
- Banerjee, J. and Narendhirakannan R.T. (2011). Phytochemical analyses, antibacterial, *in vitro* antioxidant and cytotoxic activities of ethanolic extract of *Syzygium cumini* (L.) seed extract. *International Journal of Pharmaceutical Sciences and Research*. 2(7): 1799-1806.
- Barbanti, D. and Pasquini, M. (2005). Influence of cooking conditions on cooking loss and tenderness of raw and marinated chicken breast meat. *LWT- Food Science and Technology*. 38: 895-901.
- Basma, A.A., Zuraini, Z. and Sasidharan, S. (2011). A transmission electron microscopy study of the diversity of *Candida albicans* induced by *Euphorbia hirta* L. leaf extract *in vitro*. *Asian Pacific Journal of Tropical Biomedicine*. 1(1): 20-22.
- Baumler, E.R., Carrin, M.E., Carelli, A.A. (2017). Diffusion of tocopherols, phospholipids and sugars during oil extraction from sunflower collets using ethanol as solvent. *Journal of Food Engineering*. 194: 1-8.
- Begum, M.F., Mahal and Alam, M.S. (2010). Inhibition of spore germination and mycelial growth of three fruit rot pathogens using some chemical fungicides and botanical extracts. *Journal of Earth Science*. 5: 23-27.
- Behling, R.G., Eifert, J., Erickson, M. C., Gurtler, J.B., Kornacki, J.L., Radcliff, E.L. R., Ryser, E.T., Stawick, B. and Yan, Z. (2010). Principles of microbiological troubleshooting in the industrial. *Food Processing Environment, Food Microbiology and Food Safety*, J.L. Kornacki (ed.), 5-63.

- Beleneva, I.A. (2011). Incidence and characteristics of *Staphylococcus aureus* and *Listeria monocytogenes* from the Japan and South China seas. *Marine Pollution Bulletin*. 62: 382-387.
- Beltran, D., Selma, M.V., Tudela, J.A. and Gil, M.I. (2005). Effect of different sanitizers on microbial and sensory quality of fresh-cut potato strips stored under modified atmosphere or vacuum packaging. *Postharvest Biology and Technology*. 37: 37-46.
- Benjamin, L., Atwill, E.R., Jay-Russell, M., Cooley, M., Carychao, D., Gorski, L. and Mandrell, R.E. (2013). Occurrence of generic *Escherichia coli*, *E. coli* O157 and *Salmonella* spp. in water and sediment from leafy green produce farms and streams on the Central California coast. *International Journal of Food Microbiology*. 165: 65-76.
- Bennis, S., Chami, F., Chami, N., Rhayour, K., Tantaoui-Elaraki, A. and Remmal, A. (2004). Eugenol induces damage of bacterial and fungal envelope. *Moroccan Journal of Biology*. 1: 31-38.
- Beuchat, L.R., Nail, B.V., Adler, B.B. and Clavero, M.R.S. (1998). Efficacy of spray application of chlorinated water in killing pathogenic bacteria on raw apples, tomatoes and lettuce. *Journal of Food Protection*. 61: 1305-1311.
- Beuchat, L.R. (2002). Ecological factors influencing survival and growth of human pathogens on raw fruits and vegetables. *Microbes and Infection*. 4: 413-423.
- Birhanu, S., Akhtar, M.S., and Muleta, D. (2014). Management of post-harvest fruit spoilage fungi by some potential spice extracts. *Archives of Phytopathology and Plant Protection*. 47(17): 2124-2140.
- Biswas, D. and Roymon, M. G. (2013). LC/ TOF/ ESI/ MS based detection of bioactive compounds present in leaf and bark extract of *Acacia Arabica*. *Recent Research in Science and Technology*. 5(2): 37-40.
- Bhunias, A. (2008). Foodborne microbial pathogens: mechanisms and pathogenesis. Springer, New York, USA.
- Bhunias, A.K. and Lathrop, A. (2003). Pathogen detection, food-borne. McGraw Hill Yearbook of Science and Technology. The McGraw- Hill Companies, Inc. 320-323.
- Bloomfield, S.F., Aiello, A.E., Cookson, B., O' Boyle, C. and Larson, E.L. (2007). The effectiveness of hand hygiene procedures in reducing the risks of infections in home and community settings including handwashing and alcohol-based hand sanitizers. *American Journal of Infection Control*. 35(10): 27-64.

- Bommanavar, S.B., Gugwad, S. and Malik, N. (2017). Phenotypic switch: The enigmatic white-gray-opaque transition system of *Candida albicans*. 21(1): 82-86.
- Bordini, B.M.E., Ristori, C.E., Jakabi, M. and Gelli, D.S. (2007). Incidence, internalization and behavior of *Salmonella* in mangoes, var. Tommy Atkins. *Food Control*. 18: 1002-1007.
- Borges, A., Ferreira, C., Saavedra, M.J. and Simoes, M. (2013). Antibacterial activity and mode of action of ferulic and gallic acids against pathogenic bacteria. *Microbial Drug Resistance*. 19(4):256-65.
- Botey-Salo, P., Anyogu, A., Varnam, A.H. and Sutherland, J.P. (2012). Survival of inoculated *Salmonella* on the shell of hens' eggs and its potential significance. *Food Control*. 28: 463-469.
- Brackett, R.E. (1992). Shelf stability and safety of fresh produce as influenced by sanitation and disinfection. *Journal of Food Protection*. 55: 808-814.
- Brasil, I.M., Gomes, C., Puerta-Gomez, A., Castell-Perez, M.E. and Moreira, R.G. (2012). Polysaccharide-based multi-layered antimicrobial edible coating enhances quality of fresh-cut papaya. *LWT-Food Science and Technology*. 47: 39-45.
- Burt, S. (2004). Essential oils: Their antibacterial properties and potential applications in foods - A review. *International Journal of Food Microbiology*. 94: 223-253.
- Burnham, V.E., Janes, M.E., Jakus, L. A., Supan, J., De Paola, A. and Bell, J. (2009). Growth and survival differences of *Vibrio vulnificus* and *Vibrio parahaemolyticus* strains during cold storage. *Journal of Food Science*. 74: 314-318.
- Cafarchia, C., De Laurentis, N., Milillo, M.A., Losacco, V., Puccini, V. (2000). Antifungal activity of *Apulia region propolis*. *Parasitology*. 41: 587-90.
- Casey, G. D and Dobson. A. D. W. (2003). Molecular detection of *Candida krusei* contamination in fruit juice using the citrate synthase gene *cs1* and a potential role for this gene in the adaptive response to acetic acid. *Journal of Applied Microbiology*. 95: 13-22.
- Centres for Disease Control and Prevention (CDC). (2011). Investigation update: outbreak of *Shiga toxin-producing E. coli* O104 (STEC O104:H4) infections associated with travel to Germany.
- Chahyadi, A., Hartati, R., Wirasutisna, K. R. and Elfahmi. (2014). *Boesenbergia pandurata* Roxb., An Indonesian Medicinal Plant: Phytochemistry, Biological activity, plant biotechnology. *Procedia Chemistry*. 13: 13-37.



- Chang, J.M. and Fang, T.J. (2007). Survival of *Escherichia coli* O157:H7 and *Salmonella enterica* serovars Typhimurium in iceberg lettuce and the antimicrobial effect of rice vinegar against *E. coli* O157:H7. *Food Microbiology*. 24(7-8): 745-751.
- Chen, J.H. and Hotchkiss, J.H. (2003). *Listeria monocytogenes*: low levels equals low risk. *Journal of Food Protection*. 66: 570-7.
- Chen, X. J., Meng, Z. Y., Wang, P., Lu, C. L., Yang, Y. Z., Zhang, L., Liu, L., Chen, S. (2015). Evaluation of household cleaning methods for reducing chlorantraniliprole residues on cowpea fruits. *Journal of Agricultural Science*. 7(9): 129-137.
- Clinical and Laboratory Standards Institute (CLSI). (2008). Reference method for broth dilution antifungal susceptibility testing of yeasts; approved standard-third edition; CLSI document M27 - A3. Clinical and Laboratory Standards Institute, Wayne, PA, USA.
- Clinical and Laboratory Standards Institute (CLSI). (2012). Reference method for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard-ninth edition; CLSI document M07-A9. Clinical and Laboratory Standards Institute, Wayne, PA, USA.
- Cohen, N., Ennaji, H., Bouchrif, B., Hassar, B. and Karib, H. (2007). Comparative study of microbiological quality of raw poultry meat at various seasons and for different slaughtering processes in Casablanca (Morocco). *Journal of Applied Poultry Research*. 16:502-508.
- Cooper, K. E., Davies, J. and Wiseman, J. (2005). An investigation of an outbreak of food poisoning associated with organisms of the *Proteus* group. *The Journal of Pathology*. 52(1): 91-98.
- Corbo, M.R., Bevilacqua, A., Campaniello, D.D., Amato, D., Speranza, B. and Sinigaglia, M. (2009). Prolonging microbial shelf life of foods through the use of natural compounds and non-thermal approaches-a review. *International Journal of Food Science and Technology*. 44: 223-241.
- Coro, F.A.G., Youssef, E.Y. and Shimokomaki, M. (2003). Age related changes in poultry breast meat collagen pyridinoline and texture. *Journal of Food Biochemistry*. 26: 533-541.
- Correa-Royero, J., Tangarife, V., Duran, C., Stashenko, E. and Mesa-Arango, A. (2010). *In vitro* antifungal activity and cytotoxic effect of essential oils and extracts of medicinal and aromatic plants against *Candida krusei* and *Aspergillus fumigatus*. *Brazilian Journal of Pharmacognocny*. 20(5): 734-741.
- Cowan, M.M. (1999). Plant products as antimicrobial agents. *Clinical microbiology reviews*. 12(4): 564-582.

- da Silva, C.B., Guterres, S.S., Weisheimer, V. and Schapoval, E.E.S (2008). Antifungal activity of the lemongrass oil and citral against *Candida* spp. *The Brazilian Journal of Infectious Diseases*. 12(1):63-66.
- da Silva, S. L, de Oliveira, V. G., Yano, T., Nunomura, R. C. S (2009). Antimicrobial activity of bergenin from *Endopleura uchi* (Huber) Cuatrec. *Acta Amazonica*. 39(1): 187-192.
- Dall'Agno, R., Ferraz, A., Bernardi, A.P. (2003). Antimicrobial activity of some *Hypericum* species. *Phytomedicine*. 10: 511-6.
- Daniel, D.S., Ng, Y.K., Chua, E.L., Arumugam, Y., Wong, Y.L and Kumaran, J.V. (2013). Isolation and identification of gastrointestinal microbiota from the short-nosed fruit bat *Cynopterus brachyotis brachyotis*. *Microbiology Research*. 168(8): 485-496.
- Daniels, N.A., Ray, B., Easton, A.N., Marano, N.N., Kahn, E., McShan, A.L., Del Rosario, L., Baldwin, T., Kingsley, M.A., Pühr, N.D., Wells, J.G. and Angulo, F. J. (2000). Emergence of new *Vibrio parahaemolyticus* serotype in raw oysters: A prevention quandary. *Journal of American Medical Association*. 284:1541-1545.
- de Billerback, V.G., Roques, C.G., Bessiere, J.M., Fonvieille, J.L., Dargent, R. (2001). Effect of *Cymbopogon nardus* (L.) W. Watson essential oil on the growth and morphogenesis of *Aspergillus niger*. *Canada Journal of Microbiology*. 47: 9-17.
- de Jonghe, V., Coorevits, A., Van Hoorde, K., Messens, W., Van Landschoot, A., De Vos, P. and Heyndrickx, M. (2010). Influence of storage conditions on the growth of *Pseudomonas* species in refrigerated raw milk. *Applied and Environmental Microbiology*. 77(2): 460-470.
- de Lucca, A.J. (2007). Harmful fungi in both agriculture and medicine. *American Journal of Mycology*. 24: 3-13.
- de Toledo, L.G., Ramos, M.A.D.S., Sposito, L., Castilho, E.M., Pavan, F.R., Lopes, E.D.O., Zocolo, G.J., Silva, F.A.N., Soares, T.H., Santos, A.G.D., Bauab, S.M. and Almeida, M.T.G.D. (2016). Essential oil of *Cymbopogon nardus* (L.) rendle: A strategy to combat fungal infections caused by *Candida* species. *International Journal of Molecular Science*. 17(1251): 1-16.
- Derakhshan, S., Sattari, M. and Bigdeli, M. (2008). Effect of subinhibitory concentrations of cumin (*Cuminum cyminum* L.) seed essential oil and alcoholic extract on the morphology, capsule expression and urease activity of *Klebsiella pneumoniae*. *International Journal of Antimicrobial Agents*. 32(5): 432-436.

- Derrickson-Tharrington, E., Kendall, P.A. and Sofos, J.N. (2005). Inactivation of *Escherichia coli* O157:H7 during storage or drying of apple slices pre-treated with acid solutions. *International Journal of Food Microbiology*. 99: 79-89.
- Devi, K.P., Nisha, S.A., Sakthivel, R., Pandia, S.K. (2010). Eugenol (an essential oil of clove) acts as an antibacterial agent against *Salmonella typhi* by disrupting the cellular membrane. *Journal of Ethnopharmacology*. 130: 107-115.
- Dewanti, S. and Wahyudi, M.T. (2011). Antibacterial activity of bay leaf infuse (*Folia Syzygium polyanthum* WIGHT) to *Escherichia coli* in-vitro. *Jurnal Medika Planta*. 1: 78-81.
- Deza, M.A., Araujo, M. and Garrido, M.J. (2005). Inactivation of *Escherichia coli*, *Listeria monocytogenes*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* on stainless steel and glass surfaces by neutral electrolyzed water. *Letters in Applied Microbiology*. 40: 341-346.
- Diaz-Visurraga, J. Cardenas, G. and Garcia, A. (2010). Morphological changes induced in bacteria as evaluated by electron. *Microscopy: Science, Technology, Applications and Education*. 307-315.
- Ding, T., Rahman, S.M.E. and Oh, D. (2011). Inhibitory effects of low concentration electrolyzed water and other sanitizers against foodborne pathogens on oyster mushroom. *Food Control*. 22: 318-322.
- Djoukeng, J. D., Abou-Mansour, E., Tabacchi, R., Tapondjou, A. L., Bouda, H. and Lontsi, D. (2005). Antibacterial triterpenes from *Syzygium guineense* (Myrtaceae). *Journal of Ethnopharmacology*. 101(1-3): 283-286.
- Doughari, J. H. and Manzara, S. (2008). *In vitro* antibacterial activity of crude leaf extracts of *Mangifera indica* Linn. *African Journal of Microbiology Research*. 2: 067-072.
- dos Santos, D.C. and Marin, J.M. (2005). Isolation of *Candida* spp. from mastitic bovine milk in Brazil. *Mycopathologia*. 159(2): 251-253.
- Doyle, M. E., Hartmann, F. A. and Wong, A. C. L. (2011). White paper on sources of methicillin-resistant *Staphylococcus aureus* (MRSA) and other methicillin-resistant staphylococci: Implications for our food supply?. *Fri Food Safety Reviews*. Food Research Institute, Madison.
- Doyle, M.P. and Erickson, M.C. (2006). Reducing the carriage of foodborne pathogens in livestock and poultry. *Poultry Science*. 85(6): 960-973.
- Duarte, M.C.P., Figueira, G. M., Sartoratto, A., Rehder, V.L.C. and Delarmelina, C. (2005). Anti-*Candida* activity of Brazilian medicinal plants. *Journal of Ethnopharmacology*. 97: 305-311.

- Dubal, Z.B., Paturkar, A.M., Waskar, V.S., Zende, R.J., Latha, C., Rawool, D.B. and Kadam, M.M. (2004). Effect of food grade organic acids on inoculated *S. aureus*, *L. monocytogenes*, *E. coli* and *S. Typhimurium* in sheep/goat meat stored at refrigeration temperature. *Meat Science*. 66: 817-821.
- Dubash, T., Gupta, S., Prakash, P. Y. and Bairy, I. (2010). Isolation of yeasts from various food products and detection of killer toxin activity *in vitro*. *Journal of Scientific Research*. 2(2): 407-411.
- Durairaj, S., Srinivasan, S. and Lakshmanaperumalsamy, P. (2009). *In vitro* antibacterial activity and stability of garlic extract at different pH and temperature. *Electronic Journal of Biology*. 5(1): 5-10.
- Echerenwa, M.C. and Umechuruba, C.I. (2004). Post-harvest fungal diseases of pawpaw (*Carica papaya* L.) fruits and seeds in Nigeria. *Global Journal of Pure and Applied Science*. 10(1): 69-73.
- Ekwenchi, M.M., Oluigbo, J. and Akpuaka A. (2014). Antibacterial activity of n-hexane extract of *Ocimum gratissimum* leaves. *Journal of Applied Chemistry*. 7: 6-10.
- El-Farmawi., E. Olama, D.Z. and Holail, H. (2014). The antibacterial effect of some natural bioactive materials against *Klebsiella pneumoniae* and MRSA. *International Journal of Current Microbiology and Applied Science*. 3: 576-588.
- Eluchie, C.N., Oranusi, S., Akujobi C.O. and Alagbaoso, S.O. (2016). Effect of phytol on dehydrogenase activity of bacterial isolates from grilled meat. *American Journal of Food Science and Technology*. 4(1): 1-6.
- Espinoza, N. A. (2013). Development of safe and ready to eat frozen oyster products using microwave steam-venting technology. Louisiana State University and Agricultural and Mechanical College. Doctoral dissertations.
- Ezat, S.W.P., Netty, D. and Sangaran, G. (2013). Paper review of factors, surveillance and burden of foodborne disease outbreak in Malaysia. *Malaysian Journal of Public Health Medicine*. 13(2): 98-105.
- Fallah, A.A., Saei-Dehkordi, S. and Rahnama, M. (2010). Enhancement of microbial quality and inactivation of pathogenic bacteria by gamma irradiation of ready-to-cook. Iranian barbecued chicken. *Radiation Physics and Chemistry*. 79: 1073–1078.
- Fatoki, O. A and Onifade, D. A. (2013). Use of plant antimicrobials for food preservation. *International Journal of Veterinary and Agricultural Engineering*, 7(12): 1110-1113.

- Fearnley, E., Raupach, J., Lagala, F. and Cameron, S. (2011). *Salmonella* in chicken meat, eggs and humans; Adelaide, South Australia, 2008. *International Journal of Food Microbiology*. 146: 219-227.
- Fei, L. V., Liang, H., Yuan, Q., Li, C. (2011). *In vitro* antimicrobial effects and mechanism of action of selected plant essential oil combinations against four food-related microorganisms. *Food Research International*. 44: 3057–3064.
- Ferguson, D.D., Scheftel, J., Cronquist, A., Smith, K., Woo-Ming, A. and Anderson, A. (2005). Temporally distinct *Escherichia coli* O157 outbreaks associated with alfalfa sprouts linked to a common seed source – Colorado and Minnesota. *Epidemiology Infections*. 133: 439-447.
- Fernandez-Delgado, M., Garcia-Amado, M.A., Contretras, M., Incani, R. N., Chirinos, H., Rojas, H. and Suarez, P. (2015). Survival, induction and resuscitation of *Vibrio cholerae* from the viable but non-culturable state in the Southern Caribbean Sea. *Revista De Medicina Tropical De Sao Paulo*. 57(1): 21-26.
- Ferreira-Dias, S., Valente, D.G. and Abreu. J.M.F. (2003). Comparison between ethanol and hexane for oil extraction from *Quercus suber* L. fruits. *Grasas y Aceites*. 54(4): 378-383.
- Fidel, P.L., Vazquez, J.A. and Sobel, J. D. *Candida glabrata*: Review of epidemiology, pathogenesis, and clinical disease with comparison to *C. albicans*. *Clinical Microbiology review*. 12(1): 80-96.
- Fishburn, J. D., Tang, Y and Frank, J.F. (2012). Efficacy of various consumer-friendly produce washing technologies in reducing pathogens on fresh produce. *Food Protection Trends*. 32(8): 456-466.
- Foerster, S., Unemo, M., Hathway, L. J., Low, N and Althus, C. L (2016). Time – kill curve analysis and pharmacodynamics modelling for *in vitro* evaluation of antimicrobials against *Neisseria gonorrhoeae*. *BMC Microbiology*. 16(216): 1-11.
- Food and Drug Administration (FDA). (2006). Fresh and frozen seafood: selecting and serving it safely. FDA, Rockville, MD. <http://www.fda.gov/Food/ResourcesForYou/Consumers/ucm077331.htm>.
- Foster, J.W. (1995). Low pH adaptation and the acid tolerance response of *Salmonella* Typhimurium. *Criticals Review in Microbiology*. 21(4): 215-37.
- Fu, L., Lu, W.Q. and Zhou, X.M. (2016). Phenolic compounds and *in vitro* antibacterial and antioxidant activities of three tropic fruits: persimmon, guava, and sweetsop. *BioMed Research International*. 2016: 1-9.

- Galeotti, F., Barile, E., Curir, P., Dolci, M. and Lanzotti, V. (2008). Flavonoids from carnation (*Dianthus caryophyllus*) and their antifungal activity. *Phytochemistry Letters*. 1: 44-48.
- Gamboa-Angulo, M. M., Cristobal-Alejo, J., Medina-Baizabal, I.L., Chi-Romero, F., Mendez-Gonzalez, R., Sima-Polanco, P. and May-Pat, F. (2008). Antifungal properties of selected plants from the Yucatan Peninsula, Mexico. *World Journal Microbial Biotechnology*. 24: 1955-1959.
- Gandhi, M. and Chikindas, M.L. (2007). *Listeria*: A foodborne pathogen that knows how to survive. *International Journal of Food Microbiology*. 113: 1-15.
- Gangoue-Pieboji, J., Eze, N., Djintchui, A.N., Ngameni, B., Tsabang, N., Pegnyemb, D.E. Biyiti, L., Ngassam, P., Koulla-Shiroa, P and Galleni, M. (2009). The *in vitro* antimicrobial activity of some traditionally used medicinal plants against beta-lactam-resistant bacteria. *The Journal of Infection in Developing Countries*. 3(9): 671-680.
- Garcia-Cela, E., Crespo-Sempere, A., Ramos, A. J., Sanchis, V. and Marin, S. (2014). Ecophysiological characterization of *Aspergillus carbonarius*, *Aspergillus tubingensis* and *Aspergillus niger* isolated from grapes in Spanish vineyards. *International Journal of Food Microbiology*. 173(3): 89-98.
- Garo, E, Eldridge, G.R., Goering, M.G., Pulcini, E. D., Hamilton, M. A. and Costerton, J.W. (2007). Asiatic acid and corosolic acid enhance the susceptibility of *Pseudomonas aeruginosa* biofilms to tobramycin. *Antimicrobial Agents and Chemotherapy*. 51: 1813-7.
- Garnier, L., Valence, F. and Mounier. J. (2017). Diversity and control of spoilage fungi in dairy products: An Update. *Microorganisms*. 5(42): 1-33.
- Geiges O. (1996). Microbial processes in frozen food. *Advances in Space Research*. 18(12): 109-118.
- Ghenghesh, K.S., Belhaj, K., El-Amin, W.B., El-Nefathi, S.E. and Zalmum, A. (2005). Microbiological quality of fruit juices sold in Tripoli-Libya. *Food Control* 16: 855-858.
- Gil, M.I., Selma, M.V., Lopez-Galvez, F. and Allende, A. (2009). Fresh-cut product sanitation and wash disinfection: problems and solutions. *International Journal of Food Microbiology*. 134(1-2): 37-45.
- Gill, C.O. and Badoni. M. (2005). Recovery of bacteria from poultry carcasses by rinsing, swabbing or excision of skin. *Food Microbiology*. 22: 101-107.
- Gill, A.O. and Holley, R.A. (2004). Mechanisms of Bactericidal Action of Cinnamaldehyde against *Listeria monocytogenes* and of Eugenol against *L.*

*Monocytogenes* and *Lactobacillus sakei*. *Applied and Environmental Microbiology*. 70(10): 5750-5.

- Glare, T.R., Gartrell, B.D., Brookes, J.J. and Perrot, J.K. (2014). Isolation and identification of *Aspergillus* spp. from brown kiwi (*Apteryx mantelli*) nocturnal house in New Zealand. *Avian Diseases*. 58: 16-24.
- Gomez, F., Igual, M., Camacho, M. M. and Pagan. M. J. (2013). Effect of the addition of plant extracts on the microbiota of minimally processed strawberry jam and its physicochemical and sensorial properties. *CyTA-Journal of Food*. 11(2): 171-178.
- Gomez-Lopez, V. M., Lannoo, A. S., Gil, M. I. and Allende A. (2014). Minimum free chlorine residual level required for the inactivation of *Escherichia coli* O157:H7 and trihalomethane generation during dynamic washing of fresh-cut spinach. *Food Control*. 42: 132-138.
- Goyal, A., Singh, J., Jain, S. and Pathak, D. P. (2009). Phytochemicals as potential antimutagens. *International Journal of Pharmaceutical Research and Development*. ISSN 0974-9446.
- Greig, J.D. and Ravel, A. (2009). Analysis of foodborne outbreak data reported internationally for source attribution. *International Journal of Food Microbiology*. 130: 77-87.
- Guilherme F. and Niede. A. (2007). Antimicrobial activity of *Syzygium polyanthum* (Myrtaceae) leaves extract. *Brazilian Journal of Microbiology*. 38:21.
- Guo, B.L., Li, W.K., Yu, J.G. and Xiao, P.G. (1996). Brevicornin, a flavonol from *Epimedium brevicornum*. *Phytochemistry*. 41(3): 991-992.
- Guo, Y., Zhou, H., Qin, L., Pang, Z., Qin, T., Ren, H., Pan, Z. and Zhou, K. (2016). Frequency, antimicrobial resistance and genetic diversity of *Klebsiella pneumoniae* in food samples. *Plos One*. 11(4): 1-13.
- Gupta, A., Mahajan, S., Sharma, R. (2015). Evaluation of antimicrobial activity of *Curcuma longa* rhizome extract against *Staphylococcus aureus*. *Biotechnology Reports*. 6: 51-55.
- Gutierrez, J., Bourke, P., Lonchamp, J. and Barry-Ryan, C. (2009). Impact of plant essential oils on microbiological, organoleptic and quality markers of minimally processed vegetables. *Innovative Food Science and Emerging Technologies*. 10: 195-202.
- Guynot, M.E., Marin, S., Setu, L., Sanchis, V. and Ramos, A.J. (2005). Screening for antifungal activity of some essential oils against common spoilage fungi of bakery products. *Food Science and Technology International*. 11(1): 25-32.

- Guzel-Seydim, Z. B., Greene, A. K. and Seydim, A. C. (2004). Use of ozone in the food industry. *LWT - Food Science and Technology*. 37: 453-460.
- Hamad, A., Mahardika, M.G.P., Istifah, I. and Hartanti, D. (2016). Antimicrobial and volatile compounds study of four spices commonly used in Indonesian culinary. *Journal of Food and Pharmaceutical Sciences*. 4: 1-5.
- Hamidi, M. R., Jovanova, B. and Panovska, T. K. (2014). Toxicological evaluation of the plant products using brine shrimp (*Artemia salina* L.) model. *Macedonian pharmaceutical bulletin* 60(1): 9-18.
- Har, L.W. and Ismail, I.S. (2012). Antioxidants activity, total phenolics, and total flavonoids of *Syzygium polyanthum* (Wight) wip leaves. *Journal of Applied Research on Medicinal and Aromatic Plants*. 2(2): 219-228.
- Harborne, J.B. and Williams, C. A. (2000). Advances in flavonoid research since 1992. *Phytochemistry*. 55: 481-504.
- Hartmann, M., Berditsch, M., Hawecker, J., Ardakani, M. F., Gerthsen, D. and Ulrich, A.S. (2010). Damage of the bacterial cell envelope by antimicrobial peptides gramicidin S and PGLa as revealed by transmission and scanning electron microscopy. *Antimicrobial Agents and Chemotherapy*. 54(8): 3132-3142.
- Hassan, K.I. and Ali, A.A. (2014). Rapid detection system of *Salmonella Typhi* in some commercial brands of milk in Sulamani of Iraq. *International Journal of Advanced Research*. 2(5): 1063-1067.
- Haute, V., Lopez-Galvez, Gomez-Lopez, V.M., Eriksson, M., Devlieghere, F., Allende, A., Sampers, I. (2015). Methodology for modeling the disinfection efficiency of fresh-cut leafy vegetables wash water applied on peracetic acid combined with lactic acid. *International Journal of Food Microbiology*. 208: 102-113.
- Hedayati, M.T., Pasqualotto, A.C., Warn, P.A., Bowyer, P. and Denning, D.W. (2007). *Aspergillus flavus*: human pathogen, allergen and mycotoxin producer. *Microbiology*. 153: 1677-1692.
- Heinitz, M.L., Ruble, R.D., Wagner, D.E. and Tatini, S.R. (2000). Incidence of *Salmonella* in fish and seafood. *Journal of Food Protection*. 63:579-59.
- Hemalata, V. B and Virupakshaiah, D. B. M. (2016). Isolation and Identification of food borne pathogens from Spoiled food samples. *International Journal of Current Microbiology and Applied Science*. 5(6): 1017-1025.
- Heredia, A., Barrera, C. and Andre's, A. (2007). Drying of cherry tomato by a combination of different dehydration techniques. Comparison of kinetics and other related properties. *Journal of Food Engineering*. 80: 111-118.



- Hong, C.K., Seo, Y.H., Choi, C.M., Hwang, I.S. and Kim, M.S. (2012). Microbial quality of fresh vegetables and fruits in Seoul, Korea. *Journal of Food Hygiene and Safety*. 27: 24-29.
- Huang, Y.R., Hung, Y. C., Hsu, S. Y., Huang, Y. W. and Hwang. D. F (2008). Application of electrolyzed water in the food industry. *Food Control*. 19:329-345.
- Humphries, R.M. and Linscott. A.J. (2015). Laboratory diagnosis of bacterial gastroenteritis. *Clinical Microbiology Review*. 28(1): 3-31.
- Hwang, C.A. (2016). Effect of mayonnaise pH and storage temperature on the behavior of *Listeria monocytogenes* in ham salad and potato salad. *Journal of Food Protection*. 68(8): 1628-1634.
- Indira, G. (2014). *In vitro* antifungal susceptibility testing of five antifungal agents against *Dermatophytic species* by CLSI (M38-A) microdilution method. *Clinical Microbiology*. 3: 301-305.
- Ismail, A., Mohamed, M., Sulaiman, S.A. and Wan Ahmad, W.A.N. (2013). Autonomic nervous system mediates the hypotensive effects of aqueous and residual methanolic extracts of *Syzygium polyanthum* (Wight) Walp. var. *polyanthum* leaves in anaesthetized rats. *Evidence-Based Complementary and Alternative Medicine*.
- Issa-Zaccharia, A., Kamitani, Y., Morita, K. and Iwasaki, K. (2009). Decontamination of ready-to-eat Japanese mustard green (*Brassica japonica*) from *Escherichia coli* using slightly acidic electrolyzed water. *African Journal of Microbiology Research*. 3(10): 649-653.
- Iwamoto, M., Ayers, T., Mahon, B.E. and Swerdlow, D.L. (2010). Epidemiology of seafood-associated infections in the United States. *Clinical Microbiology Reviews*. 23(2): 399-411.
- Jadhav, V., Kalase, V. and Pati, P. (2014). GC-MS analysis of bioactive compounds in methanolic extract of *Holigarna grahamii* (wight) Kurz. *International Journal of Herbal Medicine*. 2 (4): 35-39.
- Jahan, S. (2012). Epidemiology of foodborne illness, scientific, health and social aspects of the food industry, Dr. Benjamin Valdez (Ed.), ISBN: 978-953-307-916-5.
- James, T. J. and Dubery, I. A. (2009). Pentacyclic triterpenoids from the medicinal herb, *Centella asiatica* (L.) *Urban Molecules*. 14: 3922-3941.
- Jami, M., Ghanbari, M., Zunabovic, M., Domig, K. J. and Kneifel, W. (2014). *Listeria monocytogenes* in aquatic food products - A review. *Comprehensive Reviews in Food Science and Food Safety*. 13.

- Janisiewicz, W.J., Conway, W.S., Brown, M.W., Sapers, G.M., Fratamico, P. and Buchanan, R.L. (1999). Fate of *Escherichia coli* O157:H7 on fresh-cut apple tissue and its potential for transmission by fruit flies. *Applied and Environmental Microbiology*. 65(1): 1-5.
- Jay, M.J. (2000). Food Analysis: Theory and Practices. 3<sup>rd</sup> Edition. New Delhi: CBS Publishers and Distributors.
- Jennesen, J., Neilsen, K.F., Houbraken, J., Lyhne, L., Schnurer, J., Frisad, J.C. and Samson, R.A. (2005). Secondary metabolite and mycotxin production by *Rhizopus* microspores group. *Journal of agricultural and Food Chemistry*. 53: 1833-1840.
- Jin, J.K., Adams, D.O., Ko, Y., Yu, C. W. and Lin, C.H. (2004). Aviglycines and propargylglycine inhibit conidial germination and mycelia growth of *Fusarium oxysporum* F. sp. luffae. *Mycopathologia*, 158: 369-375.
- Johannes, L. (2010). Shiga toxins, from cell biology to biomedical applications. *National Review of Microbiology*. 8: 105-116.
- Jung, Y., Jang, H. and Matthews, K.R. (2014). Effect of the food production chain from farm practices to vegetable processing on outbreak incidence. *Microbiology Biotechnology*. 7(6): 517-527.
- Juntheikki, M.R. and Ritta, J.T. (2000). Inhibition of  $\beta$ - Glucosidase and esterase by tannins from betula, salix and pinus species. *Journal of Chemical Ecology*. 26(5): 1151-1165.
- Kakde, U.B. and Kakde, H.U. (2012). Incidence of post-harvest disease and airborne fungal spores in a vegetable market. *Acta Botanica Croatica*. 71(1): 147-157.
- Kala, K., Antony, V. T., Sheemole, M.S and Asha, S. (21016). Analysis of bioactive compounds present in *Syzygium caryophyllatum* (L.) Alston fruit. *International Journal of Pharmaceutical Sciences Review and Research*. 36(42): 239-243.
- Kamba, A.S., Hassan, L.G., 2010. Antibacterial screening and brine shrimp (*Artemia salina*) toxicity over *Securidaca longepedunculata* (Polygalaceae) root bark. *Asean Journal of Pharmaceutical Science*. 1: 85-95.
- Kanatt, S.R., Chander, R. and Sharma, A. (2010). Antioxidant and antimicrobial activity of pomegranate peel extract improves the shelf-life of chicken products. *International Journal of Food Science and Technology*. 45: 216-222.
- Karsha, P. V. and Lakshmi, O. B. (2010). Antibacterial of black pepper (*Piper nigrum* Linn.) with special references to its mode of action on bacteria. *Indian Journal of Natural Products and Resources*. 1(2): 213-215.

- Karunai Raj, M., Duraipandiyan, V., Agastian, P. and Ignacimuthu, S. (2012). Antimicrobial activity of bergenin isolated from *Peltophorum pterocarpum* DC. flowers. *Asian Pacific Journal of Tropical Biomedicine*. 2012: 901-904.
- Keraita, B., Konradsen, F., Drechsel, P. and Abaido, R. (2007). Reducing microbial contamination on wastewater-irrigated lettuce by cessation of irrigation before harvesting. *Tropical Medicinal International Health*. 12: 8-14.
- Keskin, D. and Ekmekci, S. (2008). Investigation of the incidence of *Pseudomonas aeruginosa* in foods and the effect of salt and pH on *P. aeruginosa*. *Hacettepe Journal of Biology and Chemistry*. 36(1): 36-41.
- Khadka, R.B., Marasini, M., Rawal, R., Gautam, D.M. and Acedo Jr, A.L. (2017). Effects of variety and postharvest handling practices on microbial population at different value chain of fresh tomato (*Solanum lycopersicum*) in Western Terai of Nepal. *BioMed Research International*. 2017: 1-6.
- Khalil, A.A., Rahman, U.U., Khan, M.R., Sahar, A., Mehmood, T. and Khan, M. (2017). Essential oil eugenol: sources, extraction technique and nutraceutical perspectives. *RSC advances*. 7: 32669.
- Khokhar, I. and Bajwa, R. (2014). Prevalence of post-harvest rot of fruits and vegetables by *Penicillium* species. *International Journal of Advanced Research in Biological Sciences*. 1(9): 14-19.
- Kim, J.B., Kim, J.M., Cho, S.H., Oh, H.S., Choi, N.J. and Oh, D.H. (2011). Toxin genes profiles and toxin production ability of *Bacillus cereus* isolated from clinical and food samples. *Journal of Food Science*. 76(1): 26-29.
- Klaiber, R.G., Baur, S., Wolf, G., Hammes, W.P. and Carle, R. (2005). Quality of minimally processed carrots as affected by warm water washing and chlorination. *Innovative Food Science and Emerging Technologies*. 6: 351-362.
- Ko, T.F., Weng, Y. M, and Robin, Y. Y. (2002). Squelene content and antioxidant activity of *Terminalia cattappa* leaves and seeds. *Journal of Agricultural and Food Chemistry*. 50(9): 5343-5348.
- Kocacaliskan, I., Talan, I., Terzi, I. (2006). Antimicrobial activity of catechol and pyrogallol as allelochemicals. *Journal of Bioscience*. 61(9-10): 639-642.
- Konate, K. and Souza, A. (2010). Polyphenol contents, antioxidant and anti-inflammatory activities of six malvaceae species traditionally used to treat hepatitis B in Burkina Faso. *European Journal Science Research*. 44(4): 570-580.

- Koutsoumanis, K.P., Kendall, P.A. and Sofos, J. N. (2003). Effect of food processing-related stresses on acid tolerance of *Listeria monocytogenes*. *Applied and Environmental Microbiology*. 69(12): 7514-7516.
- Koseki, S., Yoshida, K., Isobe, S. and Itoh, K. (2004). Efficacy of acidic electrolyzed water for microbial decontamination of cucumbers and strawberries. *Journal of Food Protection*. 67: 1247-1251.
- Kotzekidou, P., Giannakidis, P. and Boulamatsis, A. (2008). Antimicrobial activity of some plant extracts and essential oils against foodborne pathogens *in vitro* and on the fate of inoculated pathogens in chocolate. *Food Science and Technology*. 41(1): 119-127.
- Krishnan, K.R., James, F. and Mohan, A. (2016). Isolation and characterization of n-hexadecanoic acid from *Canthium parviflorum* leaves. *Journal of Chemical and Pharmaceutical Research*. 8(8): 614-617.
- Kroupitski, Y., Pinto, R., Brandl, M.T., Belausov, E. and Sela, S. (2009). Interactions of *Salmonella enterica* with lettuce leaves. *Journal of Applied Microbiology*. 106: 1876-1885.
- Kruk, Z. A., Yun, H., Rutley, D. L., Lee, E. J., Kim, Y. J. and Jo, C. (2011). The effect of high pressure on microbial population, meat quality and sensory characteristics of chicken breast fillet. *Food Control*. 22: 6-12.
- Kumar, H.N.H., Jha, A.K., Taneja, K.K., Kabra, K. and Sadiq, H.M. (2013). A Study on consumer awareness, safety perceptions & practices about food preservatives and flavouring agents used in packed/canned foods from South India. *National Journal of Community Medicine*. 4(3): 402.
- Kumudavally, K.V., Tabassum, A., Radhakrishna, K. and Bawa, A.S. (2011). Effect of ethanolic extract of clove on the keeping quality of fresh mutton during storage at ambient temperature ( $25 \pm 2^\circ\text{C}$ ). *Journal of Food Science and Technology*. 48(4): 466-471.
- Kung, V.L., Ozer, E.A. and Hauser, A.R. (2010). The accessory genome of *Pseudomonas aeruginosa*. *Microbiology and Molecular Biology Reviews*. 74(4): 621-641.
- Kusuma I. W., Harlinda, K., Enos, T. A., Farida, A., Min, Y., Kim, J. and Kim, Y. (2011). Biological activity and phytochemical analysis of three Indonesian medicinal plants, *Murraya koenigii*, *Syzygium polyanthum* and *Zingiber Purpurea*. *Journal of Acupuncture Meridian Studies*. 4 (1): 75-79.
- Kwiecinski, J., Eick, S. and Wojcika, K. (2009). Effects of tea tree (*Melaleuca alternifolia*) oil on *Staphylococcus aureus* in biofilms and stationary growth phase. *International Journal of Antimicrobial Agents*. 33(4): 343-347.

- Kwon, J. H. and Lee, C. J. (2006). *Rhizopus* soft rot on pear (*Pyrus serotina*) caused by *Rhizopus stolonifer* in Korea. *Mycobiology*. 34(3): 151-153.
- Lado, B. and Yousef A.E. (2007). Characteristics of *Listeria monocytogenes* important to food processors. In *Listeria, listeriosis and food safety*, ed. E.T. Ryser, and E.H. Marth. 157-213. Boca Raton: Taylor and Francis Group.
- Laffey, S. F. and G. Butler. (2005). Phenotype switching affects biofilm formation by *Candida parapsilosis*. *Microbiology*. 151: 1073-1081.
- Lambert, P.A. (2002). Cellular impermeability and uptake of biocides and antibiotics in Gram-positive bacteria and mycobacteria. *Journal of Applied Microbiology*. 92: 46-54.
- Lazarova, R. and Krystallis, A. (2010). Consumer's perception of food quality and its relation to the choice of food. Department of Marketing and Statistics. Aarhus Universitet.
- Lau, K.Y., Zainin, N.S., Abas, F. and Rukayadi, Y. (2014). Antibacterial and sporicidal activity of *Eugenia polyantha* Wight against *Bacillus cereus* and *Bacillus subtilis*. *International Journal of Current Microbiology and Applied Sciences*. 3(12): 499-510.
- Lawrie, R.A. and Ledward, D.A. (2006). *Lawrie's Meat Science: Seventh English*, edition ed. Cambridge England: Woodhead Publishing Limited.
- Leifert, C., Ball, K., Volakakis, N. and Cooper, J.M. (2008). Control of enteric pathogens in ready-to-eat vegetable crops in organic in low input production system: a HACCP-based approach. *Journal of Applied Microbiology*. 105(4): 931-950.
- Lee, M. (2009). Basic skills in interpreting laboratory data: fourth edition. American Society of Health-System Pharmacist, Inc. US.
- Leistner, L. (2000). Basic aspects of food preservation by hurdle technology. *International Journal of Food Microbiology*. 55: 181-186.
- Lekhnath, K. and Cheng, J. L. (2013). Toxicity and repellency of compounds from Clove (*Syzygium aromaticum*) to red imported fire ants *Solenopsis Invicta* (Hymenoptera: Formicidae). *Journal of Economic Entomology*. 106(1): 131-135.
- Lena, D. and Anustri, T. (2013). Antimicrobial activity of eugenol and cinnamaldehyde against extended spectrum beta lactamase producing *enterobacteriaceae* by *in vitro* and molecular docking analysis. *Journal of Phytochemistry*. 5(6): 527-536.

- Levin, R. E. (2006). *Vibrio parahaemolyticus*, a notably lethal human pathogen derived from seafood: A review of its pathogenicity, characteristics, subspecies characterization, and molecular methods of detection. *Food Biotechnology*. 20: 93-128.
- Li, R., Wu, X., Yin, X., Liang, J. N and Li, M. (2014). The natural product citral can cause significant damage to the hyphal cell walls of *Magnaporthe grisea*. *Molecules*. 19: 10279-10290.
- Limsuwan, S., Subhadhirasakul, S. and Voravuthikunchai, S. P. (2009). Medicinal plants with significant activity against important pathogenic bacteria. *Pharmaceutical Biology*. 47(8): 683-689.
- Lin, Y.T., Labbe, R.G. and Shetty, K. (2004). Inhibition of *Listeria monocytogenes* in fish and meat systems by use of oregano and cranberry phytochemical synergies. *Applied and Environmental Microbiology*. 70: 5672-5678.
- Little, C.L., Sagoo, S. K., Gillespie, I.A., Grant, K. and Mc Lauchlin, J. (2009). Prevalance and level of *Listeria monocytogenes* and other *Listeria* species in selected retail ready-to-eat foods in the United Kingdom. *Journal of Food Protection*. 72: 1869-1877.
- Liu, W. H., Liua, T.C. and Mong, M. C. (2015). Antibacterial effects and action modes of asiatic acid. *Biomedicine*. 5(3): 22-29.
- Liu, Y., Fan, X., Chen, Y.R. and Thayer, D.W. (2003). Changes in structure and color characteristics of irradiated chicken breasts as a function of dosage and storage time. *Meat Science*. 63: 301-307.
- Lopez- Meneses, A. K., Plascencia-Jatomea, M., Lizardi-Mendoza, J., Rosas-Burgos, E. C., Luque-Alcaraz, A. G. and Cortez-Rocha, M. O. (2015). Antifungal and antimycotoxigenic activity of essential oils from *Eucalyptus globulus*, *Thymus capitatus* and *Schinus molle*. *Food Science and Technology*. 35(4): 664-671.
- Loziene, K., Sakal, J., Pask, A. and Venskutonis, P. R. (2008). Anti-*Candida* activity of *Thymus pulegioides* (Lamiaceae) essential oils depends on the plant chemotype. *Herba Polonica*. 54: 80-92.
- Lutz, C., Erken, M., Noorian, P., Sun, S. and McDougald, D. (2016). Environmental reservoirs and mechanisms of persistence of *Vibrio cholerae*. *Frontier in Microbiology*. 4(375): 1-15.
- Lynch, M.F., Tauxe, R.V. and Hedberg, C.W. (2009). The growing burden of foodborne outbreaks due to contaminated fresh produce: risks and opportunities. *Epidemiology of Infections*. 137: 307-315.

- Lungu, B., Ricke, S.C. and Johnson, M.G. (2009). Growth, survival, proliferation and pathogenesis of *Listeria monocytogenes* under low oxygen or anaerobic conditions: A review. *Anaerobe*. 15: 7-17.
- Mahmood, A.K., Khan, M.S., Khan, M.A. and Bilal, M. (2014). Prevalence of *Salmonella* in diarrheic adult goats in field conditions. *The Journal of Animal and Plant Sciences*. 24(1): 98-102.
- Mahfuzul, H.M., Barib, M.L., Vijay, K. and Kawamoto, S. (2008). Antimicrobial activity of cloves and cinnamon extracts against food borne pathogens and spoilage bacteria, and inactivation of *Listeria monocytogenes* in ground chicken meat with their essential oils. *International Union of Pure and Applied Chemistry, International Rice Research Institute*. 72: 9-21.
- Maldonado, V.A., Glode, M.P. and Bhatia, J. (2014). Consumption of raw and unpasteurized milk and milk products by pregnant women and children. *Pediatrics*. 133(1): 134-139.
- Maldovan, M., Bogdan, C., Ursul, I., Ileana, M. I. and Crişanm, M. (2016). Evaluation of the efficacy and characterization of an anti-acne cream containing herbal extracts. *Pharmacia*. 64(2): 217-223.
- Mamman, P. H., Mshelia, W. P., Susbatrus, S. C. and Sambo. K. (2013). Antibacterial effects of crude extract of *Azadirachta indica* against *Escherichia coli*, *Salmonella* spp. and *Staphylococcus aureus*. *International Journal of Medicine and Medical Sciences*. 5: 14-18.
- Manaharan, T., Chakravarthi, S., Radhakrishnan, A. K. and Palanisamy, U. D. (2014). *In vivo* toxicity evaluation of a standardized extract of *Syzygium aqueum* leaf. *Toxicology Reports*. 1:718-725.
- Manavathu, E. K., Cutright, J. and Chandrashekar, P. H. (1999). Comparative study of susceptibilities of germinated and ungerminated conidia of *Aspergillus fumigatus* to various antifungal agents. *Journal of Clinical Microbiology*. 37: 858-861.
- Mandal, S., Deb Mandal, M., Pal, N.K. and Saha, K. (2011). Inhibitory and killing activities of black tea (*Camellia sinensis*) extract against *Salmonella enterica* serovar Typhi and *Vibrio cholera* O1 biotype E1 Tor serotype Ogawa isolates. *Jundishapur Journal of Microbiology*. 4(2): 115-121.
- Manimaran, U., Shakila, R. J., Sivaraman, B., Shalini, R., Jeyasekaran, G. and Shanmugam, S. A. (2016). Biochemical quality changes during iced storage of Indian octopus (*Cistopus indicus*). *Journal of Food Quality*. 39 (2016): 487-495.

- Maria, I. G., Maria, V. S., Lopez-Galvez, F. and Allende, A. (2009). Fresh-cut product sanitation and wash water disinfection: Problems and solutions. *International Journal of Food Microbiology*. 134: 37-45.
- Margerat, E., Shailaja, M. and Roa V.V. (2015). Evaluation of antioxidant activity in different parts of *Syzygium cumini* (Linn.). *International Journal of Current Microbiology*. 4(9): 372-379.
- Maroszynska, M., Kunicka-Styczynska, A., Rajkowska, K., Maroszynska, I. (2013). Antibiotics sensitivity of *Candida* clinical and food-borne isolates. *Acta Biochimica Polonica*. 60(4): 719-724.
- Marriott, R.J. (2010). Greener chemistry preparation of traditional flavour extracts and molecules. *Agro Food Industry Hi-Tech*. 21(2): 46-48.
- Martinez, M.J., Betancourt, J., Alonso-Gonzalez, N. and Jauregui, A. (1996). Screening of some Cuban medicinal plants for antimicrobial activity. *Journal of Ethnopharmacology*. 52: 171-174.
- Masoko, P., Picard, J., Howard, R.L., Mampuru, L.J. and Eloff, J.N. (2010). *In vivo* antifungal effect of *Combretum* and *Terminalia* species extracts on cutaneous wound healing in immunosuppressed rats. *Journal of Pharmaceutical and Biologocal Science*. 48: 621-32.
- McDonnell, G. and Russell, A.D. (1999). Antiseptics and disinfectants: activity, action and resistance. *Clinical Microbiology Reviews*. 12(1): 147-79.
- Mead, P. S., Slutsker, L., Dietz, V., McCaig, L.F., Breese, J.S., Shapiro, C., Griffin, P. M. and Tauxe, R.V. (1999). Food related illness and death in the United States. *Emergence of Infectious Diseases*. 5: 607-625.
- Meussen, B. J., de Graaff, L. H., Sanders, J. P. M. and Weusthuis, R. A. (2012). Metabolic engineering of *Rhizopus oryzae* for the production of platform chemicals. *Applied Microbiology and Biotechnology*. 94: 875-886.
- Meyer, B.N., Ferrigni, N.R., Putnam, J.E., Jascobsen, L.B., Nichols, D.E. and McLaughlin, J.L. (1982). Brine shrimp: A convenient general bioassay for active plant constituents. *Plant Medica*. 45(1): 31-34.
- Mhone, T.A., Matope, G. and Saidi, P.T. (2012). Detection of *Salmonella* spp., *Candida albicans*, *Aspergillus* spp., and antimicrobial residues in raw and processed cow milk from selected smallholder farms of Zimbabwe. *Veterinary Medicine International*. 2012: 1-5.
- Michael, A.S., Thompson, C.G. and Abramovitz, M. (1956). *Artemia salina* as a test organism for a bioassay. *Science*. 123: 464.



- Ministry of Health, Malaysia (MOH), (2014). Incidence rate and mortality rate of communicable diseases, 2013.
- Ministry of Health, Malaysia (MOH), (2016). Annual reports 2016, Malaysia.
- Miron, D., Battisti, F., Silva, F.K., Lana, A.D., Pippi, B., Casanova, B., Gnoatto, S., Fuentefria, A., Mayorga, P. and Schapoval, E.E.S. (2014). Antifungal activity and mechanism of action of monoterpenes against dermatophytes and yeasts. *Brazillian Journal of Pharmacognosy*. 24: 660-667.
- Molan, P.C. (1992). The antibacterial activity of honey-the nature of antibacterial activity. *Bee World*. 73(1): 5-28.
- Molero, G., Diez-Orejas, R., Navarro-Garcia, F., Monteoliva, L., Pla, J., Gil, C., Sanchez-Perez, M. and Nombela, C. (1998). *Candida albicans*: genetics, dimorphism and pathogenicity. *International Microbiology*. 1: 95-106.
- Morshedy, A.E.M.A. and Sallam, K.I. (2009). Improving the microbial quality and shelf life of chicken carcasses by trisodium phosphate and lactic acid dipping. *International Journal of Poultry Science* 8: 645-650.
- Mpuchane, S.F., and Gashe, B.A. (1996). Presence of *Escherichia coli*, *Klebsiella pneumoniae* and *Enterobacter* species in dried bush okra (*Corchorus olitorius*) and African spider herb (*Cleome gynandra*). *Food Control*. 7: 169-172.
- Mrityunjay, A., Kaniz, F., Fahmida J., Shanzida, J.S., Md. Aftab, U. and Rashed, N. Prevalence of *Vibrio cholerae* in different food samples in the city of Dhaka, Bangladesh. *International Food Research Journal*. 20(2): 1017-1022.
- Mueller, M., Pena, A. and Derendorf, H. (2004). Issues in pharmacokinetics and pharmacodynamics of anti-infective agents: kill curves versus MIC. *Antimicrobial Agents and Chemotherapy*. 48(2): 369-377.
- Muniandy, K., Hassan, Z. and Isa, M.H.M. (2014). Antimicrobial activity of ethanolic extract of *Coleus aromaticus* against common wound pathogens. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 5(3): 1286.
- Murali Krishna, T., Meena, G., Kavya, T., Someshwar, C., Soumya, J., Aswaq A., Rajender, V. and Rajesh, G. G. (2013). *In vitro* determination of antioxidant and anti-bacterial activities of *Vitex Negundo* Linn. *International Journal of Pharmacology and Biological Sciences*. 4 (1): 121-127.
- Musa, A.A. (2012). Cytotoxicity activity and phytochemical screening of *Cochlospermum tinctorium* Perr Ex A. rich rhizome. *Journal of Applied Pharmacy and Sciences*. 2: 155-159.

- Mustafa, R.A., Abdul Hamid, A., Mohamed, S. and Abu Bakar, F. (2010). Total phenolic compounds, flavonoids and radical scavenging activity of 21 selected tropical plants. *Journal of Food Science*. 75(1).
- Nair, G.B., Ramamurthy, T., Bhattacharya, S. K., Dutta, B., Takeda, Y. and Sack, D. A. (2007). Global dissemination of *Vibrio parahaemolyticus* serotype O3:K6 and its serovariants. *Clinical Microbiology Reviews*. 20: 39-48.
- Nadeem, S.G., Shafiq, A., Hakim, S.T., Anjum, Y. and Kazm, S.U. (2013). Effect of growth media, pH and temperature on yeast to hyphal transition in *Candida albicans*. *Open Journal of Medical Microbiology*. 3:185-192.
- Nakayama, S., Morita, T., Negishi, H., Ikegami, T., Sakaki, K. and Kitamoto, D. (2008). *Candida krusei* produces ethanol without production of succinic acid; A potential advantage for ethanol recovery by evaporation membrane separation. *FEMS Yeast Research*. 8(5). 706-714.
- Namuli, A., Abdullah, N., Sieo, C. C., Zuhainis, S. W. and Oskoueian, E. (2011). Phytochemical compounds and antibacterial activity of *Jatropha curcas* Linn. extracts. *Journal of Medicinal Plants Research*. 5(16): 3982-3990.
- National Committee for Clinical Laboratory Standards. (NCCLS). (2004). Reference method for antifungal disk diffusion susceptibility testing of yeasts; approved guideline. NCCLS document M44-A. National Committee for Clinical and Laboratory Standards, Wayne, PA, USA.
- Ndjateu, F.S.T., Tsafack, R.B.N., Nganou, B.K., Awouafack, M.D., Wabo, H.K., Tene, M, Tane, P. and Eloff, J.N. (2014). Antimicrobial and antioxidant activities of extracts and ten compounds from three Cameroonian medicinal plants: *Dissotis perkinsiae* (Melastomaceae), *Adenocarpus mannii* (Fabaceae) and *Barteria fistulosa* (Passifloraceae). *Journal of Botany*. 91: 37-42.
- Nebahat, O., Gulmez, M., Vatansever, L. and Guven, A. (2008). Application of antimicrobial ice for extending shelf life of fish. *Journal of Food Protection*. 71(1): 218-222.
- Negi, P.S. (2012). Plant extracts for the control of bacterial growth: Efficacy, stability and safety issues for food application. *International Journal of Food Microbiology*. 156: 7-17.
- Nelson, E.J., Harris, J.B., Jr, J.G., Calderwood, S.B. and Camilli, A. (2013). Cholera transmission: the host, pathogen and bacteriophage dynamic. *Nature Review Microbiology*. 7(10): 1-24.
- Neo, S.Y., Lim, P.Y., Phua, L.K., Khoo, G.H., Kim, S.J., Lee, S.C. and Yuk, H.G. (2013). Efficacy of chlorine and peroxyacetic acid on reduction of natural microflora, *Escherichia coli* O157: H7, *Listeria monocytogenes* and *Salmonella* spp. on mung bean sprouts. *Food Microbiology*. 36: 475-480.

- Neto, O.C.F., Filho, R.A.C.P., Barrow, P. and Junior A.B. (2010). Sources of human non-typhoid salmonellosis: A Review. *Brazilian Journal of Poultry Science*. 12(1): 1-11.
- Newkirk, R., Hedberg, C. and Bender, J. (2011). Establishing a milkborne disease outbreak profile: potential food defense implications. *Foodborne Pathogens Diseases*. 8(3): 433-437.
- Nitiema, L.W., Savadogo, A., Simpore, J., Dianou, D. and Traore, A.S. (2012). *In vitro* antimicrobial activity of some phenolic compounds (coumarin and quercetin) against gastroenteritis bacterial strains. *International Journal of Microbiological Research*. 3(3): 183-187.
- Noorlis, A., Ghazali, F. M., Cheah, Y. K., Tuan Zainazor, T. C., Ponniah, J., Tunung, R., Tang, J. Y. H., Nishibuchi, M., Nakaguchi, Y. and Son, R. (2011). Prevalence and quantification of *Vibrio* species and *Vibrio parahaemolyticus* in freshwater fish at hypermarket level. *International Food Research Journal* 18: 689-695.
- Normanno, G., La-Salandra, G., Dambrosio, A., Quaglia, N. C., Corrente, M., Parisi, A., Santagada, G., Firinu, A., Crisetti, E. and Celano, G.V. (2007). Occurrence, characterization and antimicrobial resistance of enterotoxigenic *Staphylococcus aureus* isolated from meat and dairy products. *International Journal of Food Microbiology*. 115: 290-296.
- Northcutt, J.K., Buhr, R.J., Young, L.L., Lyon, C.E. and Ware, G.O. (2001). Influence of age and postchill carcass aging duration on chicken breast fillet quality. *Poultry Science*. 80: 808-812.
- Noveriza, R. and Miftakhurohmah. (2010). Efektivitas ekstrak metanol daun salam (*Eugenia Polyantha*) dan daun jeruk purut (*Cytrus histrix*) sebagai antijamur pada pertumbuhan *Fusarium Oxysporum*. *Jurnal Littri*. 16 (1): 6 -11.
- Nyenje, M.E., Odjadjare, C.E., Tanih, N.F., Green, E. and Ndip, R.N. (2012). Foodborne pathogens recovered from ready-to-eat foods from roadside cafeterias and retail outlets in Alice, Eastern Cape Province, South Africa: public health implications. *International Journal of Environmental Research Public Health*. 9: 2608-2619.
- Obire, O., Okigbo, R. N. and Minimah, T. R. (2013). Mycoflora of fresh shrimps (*Penaeus aztecus*) from different markets in Port Harcourt, Nigeria. *African Journal of Food Sciences*. 7(8): 204-209.
- Odeyemi, O. A. (2016). Incidence and prevalence of *Vibrio parahaemolyticus* in seafood: a systematic review and meta-analysis. *Springer Plus*. 5(464): 1-17.

- Odiba, J.O., Aliyu M.M., Halimatu, S.H., Sani, M.Y. and Ephraim, I.O. (2014). Antimicrobial activity of isolated Stigmast- 5-en-3 $\beta$ -ol ( $\beta$ -Sitosterol) from honeybee propolis from North-Western, Nigeria. *International Journal of Pharma Sciences and Research*. 5(12): 345-350.
- Olayujigbe, O.O., Oluremi, B.B. and Umaru, D.G. (2006). Bacterial spoilage of fresh meat in some selected lagos markets. *Ife Journal of Science*. 8(2): 193-198.
- Oliver, S.P., Jayarao, B.M. and Almeida, R.A. (2005). Foodborne pathogens in milk and the dairy farms environment: food safety and public health implications. *Foodborne Pathogens and Disease*. 2(2): 302-307.
- Olsen, S.J., Aucott, J.N. and Swerdlow, D.L. (2002). Food poisoning. Infections of the gastrointestinal tract, 2nd ed. Lippincott Williams and Wilkins, Philadelphia, PA. 199-214.
- Omidbeygi, M., Barzegar, M., Hamidi, Z. and Naghdibadi, H. (2007). Antifungal activity of thyme, summer savory and clove essential oils against *Aspergillus flavus* in liquid medium and tomato paste. *Food Control*. 18: 1518-1523.
- Omokaro, O. and Minimah, T.R. (2013). Mycoflora of fresh shrimps (*Penaeus aztecus*) from different markets in Port Harcourt Nigeria. *e-Journal of Science & Technology*. 8(4):87-95.
- Orhan, D.D., Ozcelik, B., Ozgen, S. and Ergun, F. (2010). Antibacterial, antifungal and antiviral activities of some flavonoids. *Microbiological Research*. 165: 496-504.
- Otang, W.M., Grierson, D.S., Ndip, R.N. (2011). The effect of the acetone extract of *Arctotis arctotoides* (Asteraceae) on the growth and ultrastructure of some opportunistic fungi associated with HIV/AIDS. *International Journal of Molecular Science*. 12: 9226-9235.
- Othman, M., Wiart, H.S., Khoo, C., Lim, T.J. and Ting, K.N. (2011). Optimal methods for evaluating antimicrobial activities from plant extracts. *Journal of Microbiological Methods*. 84: 161-166.
- Othman, A., Mukhtar, N.J., Ismail, N.S. and Sui, K.C. (2014). Phenolics, flavonoids content and antioxidant activities of 4 Malaysian herbal plants. *International Food Research Journal*. 21(2): 759-766.
- Oussalah, M., Caillet, S., Saucier, L. and Lacroix, M. (2006). Antimicrobial effects of selected plant essential oils on the growth of four pathogenic bacteria: *E. coli* O157:H7, *Salmonella* Typhimurium, *Staphylococcus aureus* and *Listeria monocytogenes*. *Food Control*. 18(5): 414-420.
- Pag, U., Oedenkoven, M., Sass, V., Shai, Y., Shamova, O., Antcheva, N., Tossi, A. and Sahl. H.G. (2008). Analysis of *in vitro* activities and modes of action of

synthetic antimicrobial peptides derived from an alpha-helical 'sequence template.' *Journal of Antimicrobial and Chemotherapy*. 61:341-352.

- Palacio, A.D, Bettucci, L. and Dinorah, P. (2016). *Fusarium* and *Aspergillus* mycotoxins contaminating wheat silage for dairy cattle feeding in Uruguay. *Brazilian Journal of Microbiology*. 47:1000-1005.
- Parekh, J., Jadeja, D. and Chanda, S. (2005). Efficacy of aqueous and methanol extracts of some medicinal plants for potential antibacterial activity. *Turkish Journal of Biology*. 29: 203-210.
- Park, M., Gwak, K., Yang, I., Kim, K., Jeung, E. and Chang, J. (2009). Effect of citral, eugenol, nerolidol and  $\alpha$ -terpineol on the ultrastructural changes of *Trichophyton mentagrophytes*. *Fitoterapia*. 80(5):290-6.
- Parnell, T.L., Harris, L.J. and Suslow, T.V. (2005). Reducing *Salmonella* on cantaloupes and honeydew melons using wash practices applicable to postharvest handling, foodservice and consumer preparation. *International Journal of Food Microbiology*. 99: 59-70.
- Paul, S., Dubey, R.C., Maheswari, D.K. and Kang, S.C. (2011). *Trachyspermum ammi* (L.) fruit essential oil influencing on membrane permeability and surface characteristics in inhibiting food-borne pathogens. *Food Control*. 22: 725-731.
- Peles, F., Wagner, M., Varga, L., Hein, I., Rieck, P., Gutser, K., Kereszturi, P., Kardos, G., Turcsanyi, I., Beri, B. and Szabo, A. (2007). Characterization of *Staphylococcus aureus* strains isolated from bovine milk in Hungary. *International Journal of Food Microbiology*. 118: 186-193.
- Penduka, D. and Okoh, A.I. (2012). *In vitro* antilisterial properties of crude methanol extracts of *Garcinia kola* (heckel) seeds. *The Scientific World Journal*. 12: 6952-6965.
- Perez-Gregorio, M. R., Gonzalez-Barreiro, C., Rial-Otero, R., Simal-Gándara, J. (2011). Comparison of sanitizing technologies on the quality appearance and antioxidant levels in onion slices. *Food Control*. 22: 2052-2058.
- Perrone, G., Susca, A., Cozzi, G., Ehrlich, I.K., Varga, J., Frisvad, J.C., Meijer, M., Noonim, P., Mahakarnchanakul, W. and Samson, R.A. (2007). Biodiversity of *Aspergillus* species in some important agricultural products. *Studies in Mycology*. 59: 53-66.
- Perumal, S., Mahmud, R., Piaru, S.P., Cai, L. W. and Ramanathan, S. (2012). Potential antiradical activity and cytotoxicity assessment of *Ziziphus mauritiana* and *Syzygium polyanthum*. *International Journal of Pharmacology*. 8(6): 535-541.
- Philippe, S., Souaibou, F., Jean-Pierre, N., Brice, F., Paulin, A. and Issaka, Y. (2012). Chemical composition and *in vitro* antifungal activity of *Zingiber officinale*

essential oil against foodborne pathogens isolated from a traditional cheese wagashi produced in Benin. *International Journal of Biosciences*. 2: 22-28.

- Phillips, C.A. (2001). *Arcobacter* spp in food: isolation, identification and control. *Trend in Food Science and Technology*. 12(8): 263-275.
- Pisutthanan, S., Plianbangchang, P., Pisutthanan, N., Ruanruay, S., Muanrit, O. (2004). Brine shrimp lethality activity of Thai medicinal plants in the family Meliaceae. *Naresuan University Journal*. 12: 13-18.
- Pitt, J.I. and Hocking, A.D. (2009). *Fungi and Food Spoilage* (3<sup>rd</sup> edition). New York: Springer.
- Pizato, S., Cortez-Vega, W.R. and Prentice, C. (2015). Quality assessment of cooked chicken breast meat at different storage temperatures. *International Food Research Journal*. 22(1): 143-154.
- Powell, S. and Attwell, R.W. (1998). An evaluation of the collection and analysis of epidemiological data for support of food safety control systems. *Journal of Food Protection*. 61: 1170-1174.
- Prabhakar, K., Sathish Kumar, L. and Sajit Khan, A.K. (2008). Antifungal activity of plant extracts against *Candida* species from oral lesions. *Indian Journal of Pharmaceutical Sciences*. 70(60): 801-803.
- Pretti, R. (2012). Cytotoxicity of *Aspergillus* fungi as a potential infectious threat. Insight and Control of Infectious Disease in Global Scenario. *In Tech*. (231-248). Croatia.
- Pui, C.F., Wong, W.C., Chai, L.C., Lee, H.Y., Noorlis, A., Zainazor, T.C.T., Tang, J.Y.H., Ghazali, F.M., Cheah, Y.K., Nakaguchi, Y., Nishibuchi, M. and Radu, S. (2011). Multiplex PCR for the concurrent detection and differentiation of *Salmonella* spp., *Salmonella* Typhi and *Salmonella* Typhimurium. *Tropical Medicine and Health*. 39(1): 9-15.
- Puspanadan, S., Afsah-Hejri, L., John, Y.H.T., Rukayadi, Y., Loo, Y.Y., Nillian, E., and Son, R. (2013). Characterization of extended-spectrum  $\beta$ -lactamases (ESBLs) producers in *Klebsiella pneumoniae* by genotypic and phenotypic method. *International Food Research Journal*. 20(3): 1479-1483.
- Puupponen-Pimia, R., Nohynek, L., Meier, C., Kahkonen, M., Heinonen, M. and Hopia, A. (2001). Antimicrobial properties of phenolic compounds from berries. *Journal of Applied Microbiology*. 90: 494-507.
- Quarenghi, M.V, Tereschuk, M.L, Baigori, M.D. and Abdala, L.R. (2000). Antimicrobial activity of flowers from *Anthemis cotula*. *Fitoterapia*. 71:710-712.

- Quiroga, E.N., Sampietro, A.R., and Vattuone, M.A. (2001). Screening antifungal activities of selected medicinal plants. *Journal of Ethnopharmacology*. 74: 89-96.
- Rabbani, G.H. and Greenough, W.B. (1999). Food as a vehicle of transmission of cholera. *Journal of Diarrhoeal Diseases Research*. 17(1): 1-9.
- Rafiqul Islam, M. Shahidul, A., Ziaur Rahman, M., Chowdhury, S. P., Begum, M. S., Nargis Akhter, M.S. Alam, S., Han, K. D. and Lee, M. W. (2003). Effects of plant extracts on conidial germination, mycelial growth and sporulation of fungi isolated from poultry feed. *Mycobiology*. 31(4): 221-225.
- Rajabi, S., Ramazani, A., Hamidi, M. and Naji, T. (2015). *Artemia salina* as a model organism in toxicity assessment of nanoparticles. *DARU Journal of Pharmaceutical Sciences*. 23: 20-21.
- Rajeh, M.A.B., Zuraini, Z., Sasidharan, S., Latha, L.Y. and Amutha, S. (2010). Assessment of *Euphorbia hirta* L. leaf, flower, stem and root extracts for their antibacterial and antifungal activity and brine shrimp lethality. *Molecules*. 15(9): 6008-6018.
- Ramachandran, S., Vamsikrishna, M., Gowthami, K.V., Heera, B. and Dhanaraju, M. D. (2011). Assessment of cytotoxic activity of *Agave cantula* using brine shrimp (*Artemia salina*) lethality bioassay. *Asian Journal of Scientific Research*. 4(1): 90-94.
- Rawat, S. (2015). Food Spoilage: Microorganisms and their prevention. *Asian Journal of Plant Science and Research*. 5(4): 47-56.
- Razzaq, R., Farzana, K., Mahmood, S. and Murtaza, G. (2014). Microbiological analysis of street vended vegetables in Multan city, Pakistan: A public health concern. *Pakistan Journal of Zoology*. 46(4): 1133-1138.
- Ribes, J. A., Vanover-Sams, C. L. and Baker, D. J. (2000). Zygomycetes in human disease. *Clinical Microbiology Review*. 13(2): 236-301.
- Ricardo, E., Miranda, I. M., Faria-Ramos, I., Silva, R. M., Rodrigues, A. G. and Pina-Vaz, I. (2014). *In vivo* and *in vitro* acquisition of resistance to voriconazole by *Candida krusei*. *Antimicrobial Agents and Chemotherapy*. 58(8): 4604-4611.
- Rittner, H. (1992). Extraction of vegetable oils with ethyl alcohol. *Oleagineux*. 47: 29-42.
- Rivera, E. V. (2005). A review of chemical disinfection methods for minimally processed leafy vegetables. Kansas State University. Manhattan, Kansas, United State. Master Thesis.

- Roberts, D. and Greenwood, M. (2008). Practical Food Microbiology. Wiley Online Library. 3<sup>rd</sup> edition. Blackwell Publishing Ltd. Online ISBN: 9780470757512.
- Rocha, R.P., Evandro, C.M., Barbosa, L.C.A., dos Santos, R.H.S., Cecon, P.R., Dallacort, R. and Santi, A. (2014). Influence of plant age on the content and composition of essential oils of *Cymbopogon citratus* (DC.) Stapf. *Journal of Medicinal Plant Research*. 8(37): 1121-1126.
- Rukayadi, Y. and Hwang, J.K. (2006). *In vitro* activity of xanthorrhizol against *Streptococcus mutans* biofilms. *Letters in Applied Microbiology*. 42: 400-404.
- Rukayadi, Y. and Hwang, J. K. (2007). *In vitro* antimycotic activity of xanthorrhizol isolated from *Curcuma xanthorrhiza* Roxb. against opportunistic filamentous fungi. *Phytotherapy Research* 23: 434-438.
- Rukayadi, Y., Shim, J.S. and Hwang, J.K. (2008). Screening of Thai medicinal plants for antiCandidal activity. *Mycoses*. 51(4): 308-312.
- Rukayadi, Y., Lau, K.Y., Zainin, N.S., Zakaria, M. and Abas, F. (2013). Screening of antimicrobial activity of tropical edible medicinal plant extracts against five standard microorganisms for natural food preservatives. *International Food Research Journal*. 20(5): 2905-2910.
- Sabota, J. M., Hoppes, W. L., Ziegler, J. R., du Pont, H, Mathewson, J and Rutecki, G. W. (1998). A new variant of food poisoning: enteroinvasive *Klebsiella pneumoniae* and *Escherichia coli* sepsis from a contaminated hamburger. *American Journal of Gastroenterology*. 93(1): 118-119.
- Sabu, A., Sarita, S., Pandey, A., Bogar, B., Szakacs, G. and Soccol, C. R. (2002). Solid-state fermentation for production of phytase by *Rhizopus oligosporus*. *Applied Biochemistry and Biotechnology*. 102(1-6): 251-260.
- Saftner, R.A., Bai, J.H., Abbott, J.A. and Lee, Y.S. (2003). Sanitary dips with calcium propionate, calcium chloride or a calcium amino acid chelate maintain quality and shelf stability of fresh-cut honeydew chunks. *Postharvest Biology and Technology*. 29: 257-269.
- Sahgal, G., Ramanathan, S., Sasidharan, S., Mordi, M.N., Ismail, S. and Mansor, S.M. (2010). Brine shrimp lethality assay and acute oral toxicity studies on *Swietenia mahagoni* (Linn.) Jacq. seed methanolic extract. *Pharmacognosy Research*. 2(4): 215-220.
- Salama, H. M.H. and Marraiki, N. (2010). Antimicrobial activity and phytochemical analyses of *Polygonum aviculare* L. (Polygonaceae), naturally growing in Egypt. *Saudi Journal of Biological Sciences*. 7(12): 185-189.



- Salehan, N.M., Meon, S. and Ismail, I.S. (2013). Antifungal activity of *Cosmos caudatus* extracts against seven economically important plant pathogens. *International Journal of Agriculture and Biology*. 15: 864-870.
- Samaranayake, Y.H., Wu, P. C., Samaranayake, L.P. and Ho, P. L. (1994). The relative pathogenicity of *Candida krusei* and *Candida albicans* in the rat oral mucosa. *Journal of Medical Microbiology*. 47: 1047-1057.
- Sandvig, K., Grimmer, S., Lauvrak, S.U., Torgersen, M.L, Skretting, G., Van Deurs, B. and Iversen, T.G. (2002). Pathways followed by ricin and Shiga toxin into cells. *Histochemistry and Cell Biology*. 117: 131-141.
- Santarelli, R. L, Pierre, F. and Corpet, D. E. (2009). Processed meat and colorectal cancer: a review of epidemiologic and experimental evidence. *Nutrition and Cancer*. 60(2): 131-144.
- Sapers, G.M. (2003). Washing and sanitizing raw materials for minimally processed fruits and vegetables. In Novak, J.S., G.M. Sapers, and V.K. Juneja. (Eds). *Microbial Safety of Minimally Processed Foods*. Boca Raton, FL: CRC Press: 221-253.
- Sariga, C. D., Shakila, R. and Kothai, S. (2015). Isolation, characterization and quantification of bergenin from *Syzygium cumini* stem bark. *International Journal of Pharmacy*. 6(2): 108-110.
- Sarker, S.D. and Nahar, L. (2012). Natural products isolation: methods in molecular biology. *Springer Science and Business Media*. 864: 33-34.
- Sautron, C. and Cock, I. (2014). Antimicrobial activity and toxicity of *Syzygium australe* and *Syzygium leuhmannii* fruit extracts. *Pharmacognosy Communications*. 4(1): 53-60.
- Scallan, E., Hoekstra, R.M., Angulo, F.J., Tauxe, R.V., Widdowson, M., Roy, S.L., Jones, J.L. and Griffin, P.M. (2011). Foodborne illness acquired in the United States – major pathogens. *Emerging Infectious Diseases*. 17(1): 7-15.
- Schlesser, J.E., Gerdes, R., Ravishankar, S., Madsen, K., Mowbray, J. and Teo, A.Y. (2006). Survival of a five-strain cocktail of *Escherichia coli* O157:H7 during the 60-day aging period of cheddar cheese made from unpasteurized milk. *Journal of Food Protection*. 69(5): 990-998.
- Schuler, G. A., Nolan, M. P., Reynolds, A. E. and W.C. Hurst. (1999). Cleaning, sanitization and pest control in food processing, storage and service area. University of Georgia College of Agricultural and Environmental Sciences and the U.S. Department of Agriculture cooperating. Bulletin 927. *Food Science*. 11-1.

- Schuster, E., Dunn-Coleman, N., Frisvad, J. C. and van Dijk, P. W. M. (2002). On the safety of *Aspergillus niger* – A review. *Applied Microbiology and Biotechnology*. 59: 426-435.
- Scorzoni, L., Benaducci, T., Almeida, A.M.F., Silva, D.H.S., Bolzani, V.S., Mendes-Giannini, M.J.S. (2007). Comparative study of disk diffusion and microdilution methods for evaluation of antifungal activity of natural compounds against medical yeasts *Candida* spp and *Cryptococcus* sp. *Journal of Basic and Applied Pharmaceutical Sciences*. 28(1): 25-34.
- Scott, E. (2003). Food safety and foodborne disease in 21<sup>st</sup> century homes. *The Canadian Journal of Infectious Diseases*. 14(5): 277-280.
- Selma, N., Silva, G., Iracelle C. and Borges, H. (2012). The toxicity evaluation of *Syzygium cumini* leaves in rodents. *Revista Brasileira de Farmacognosia*. 22:121-125.
- Sengun, I.Y. and Karapinar, M. (2004). Effectiveness of lemon juice, vinegar and their mixture in the elimination of *Salmonella* Typhimurium on carrots (*Daucus carota* L.). *International Journal of Food Microbiology*. 96: 301-305.
- Sen, A., Dhavan, P., Shukla, K. K, Singh, S., Tejovathi, G. (2012) Analysis of IR, NMR and antimicrobial activity of  $\beta$ -sitosterol isolated from *Momordica charantia*. *Science Secure Journal of Biotechnology*. 1(1):9-13.
- Setiawan, C. P. (2002). Effect of chemical and physical treatment of the antimicrobial activity of leaves (*Syzygium polyanthum* (Wight) Walp). Thesis. Faculty of Agricultural Technology, Bogor Agricultural University, Bogor.
- Shaik, M.A, Aleem, A.K. and Irshad, A. (2005). Antimicrobial activities of eugenol and cinnamaldehyde against the human gastric pathogen *Helicobacter pylori*. *Annals of Clinical Microbiology and Antimicrobials*. 4: 2025
- Sharma, N. and Tripathi, A. (2008). Effects of *Citrus sinensis* (L.) Osbeck epicarp essential oil on growth and morphogenesis of *Aspergillus niger* (L.) Van Tieghem. *Microbiological Research*. 163(3): 337-344.
- Shaw, I.C. (2012). Food Safety : The Science of Keeping Food Safe. Retrieved from <http://www.ebilib.com> pp. 306- 334.
- Shi, C., Song, K., Zhang, X., Sun, Y., Sui, Y., Chen, Y., Jia, Z., Sun, H., Sun, Z. and Xia, X. (2016). Antimicrobial activity and possible mechanism of action of citral against *Cronobacter sakazakii*. *PLoS One*. 11(7): 1-12.
- Shukla, Y. N., Srivastava, A., Kumar, S. and Kumar, S. (1999). Phytotoxic and antimicrobial constituents of *Argyreia* species and *Oenothera biennis*. *Journal of Ethnopharmacology*. 67:241-245.

- Sims, C. R., Paetznick, V. L., Rodriguez, J. R., Chen, E. and Ostrosky-Zeichner, L. (2006). Correlation between microdilution, E-test, and disk diffusion methods for antifungal susceptibility testing of Posaconazole against *Candida* spp. *Journal of Clinical Microbiology*. 4(6): 2105-2108.
- Singh, G. and Kumar, P. (2013). Evaluation of antimicrobial efficacy of flavonoids of *Withania Somnifera* L. *Indian Journal of Pharmaceutical Science*. 73(4): 473-478.
- Singh, N., Singh, R.K., Bhunia, A.K. and Stroshine, R.L. (2003). Efficacy of chlorine dioxide, ozone and thyme essential oil or sequential washing in killing *Escherichia coli* O157:H7 on lettuce and baby carrots. *Lebensmittel-Wissenschaft and Technology*. 35: 720-729.
- Skrinja, M. M. and Nemet, N. T. (2009). Antimicrobial effects of spices and herbs essential oils. *Acta Periodica Technologica*. 50: 195-209.
- Soderstrom, A., Osterberg, P., Lindqvist, A., Jonsson, B., Lindberg, A. and Blide, U. S. (2008). A large *Escherichia coli* O157 outbreak in Sweden associated with locally produced lettuce. *Foodborne Pathogens Disease*. 5: 339-349.
- Solomon, E.B., Yaron, S. and Matthews, K.R. (2002). Transmission of *Escherichia coli* O157:H7 from contaminated manure and irrigation water to lettuce plant tissue and its subsequent internalization. *Applied of Environmental Microbiology*. 68: 397-400.
- Soon, J.M., Singh, H. and Baines, R. (2011). Foodborne diseases in Malaysia: A review. *Food Control*. 22(6): 823-830.
- Souza, E.L., Stamford, T.L.M. and Lima, E.D.O. (2006). Sensitivity of spoiling and pathogen food-related bacteria to *Origanum vulgare* L. (Lamiaceae) essential oil. *Brazillian Journal of Microbiology*. 37(4): 527-532.
- Soylu, S. (2006). Accumulation of cell-wall bound phenolic compounds and phytoalexin in *Arabidopsis thaliana* leaves following inoculation with pathovers of *Pseudomonas syringae*. *Plant Sciences*. 170: 942-952.
- Spacil, Z., Novakova, L. and Solich, P. (2010). Comparison of positive and negative ion detection of tea catechins using tandem mass spectrometry and ultrahigh performance liquid chromatography. *Food Chemistry*. 123: 535-541.
- Srivastava, R.P. and Kumar, S. (2002). *Fruit and Vegetable Preservation: Principles and Practices*. Lucknow: International Book Distributing Co.
- Stivarius, M.R., Pohlman, F.W., McElyea, K.S. and Apple, J.K. (2002). Microbial, instrumental color and sensory color and odor characteristics of ground beef produced from beef trimmings treated with ozone or chlorine dioxide. *Meat Science*. 60: 299-305.

- Stockinger Lab. (2001). Glycerol stocks. Retrieved 5 January 2016 from [www.oardc.ohio-state/stockingerlab/](http://www.oardc.ohio-state/stockingerlab/).
- Stopforth, J.D., Ikeda, J.S., Kendall, P.A. and Sofos, J.N. (2004). Survival of acid-adapted or non adapted *Escherichia coli* O157:H7 in apple wounds and surrounding tissue following chemical treatments and storage. *International Journal Food Microbiology*. 90: 51-61.
- Subashini, M. S. and Rajendran, P. (2015). *In vitro* screening of anti HBV and anti HIV properties of *Gymnema sylvestre* R. Br leaves from Kolli Hills, Tamilnadu, India. *International Journal of Current Microbiology and Applied Sciences*. 4(3): 542-547.
- Sudjana, A.N., D' Rozario, D., Ryan, B., Rasool, L., Ng, J., Islam, N., Riley, T.V. and Hammer, K. A. (2008). Antimicrobial activity of commercial *Olea europaea* (olive) leaf extract. *International Journal of Antimicrobial Agents*. 33(5): 461-463.
- Su, Y.C. and Liu, C. (2007). *Vibrio parahaemolyticus*: a concern of seafood safety. *Food Microbiology* 24: 549-558.
- Suharti, S., Banowati, A., Hermana, W. and Wiryawan, K. G. (2008). Komposisi dan kandungan kolesterol karkas ayam broiler diare yang diberi tepung daun salam (*Syzygium polyanthum* Wight) dalam ransum. *Media Peternakan*. 31(2): 138-145.
- Suleiman, E. A and Ahmed Mohamed, E. (2014). *In vitro* activity of *Lawsonia inermis* (Henna) on some pathogenic fungi. *Journal of Mycology*. 2014: 1-5.
- Sultana, B., Anwar, F. and Ashraf, M. (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. *Molecules*. 14(6): 2167-2180.
- Sumono, A. and Wulan, A.S. (2008). The use of bay leaf (*Eugenia polyantha* Wight) in dentistry. *Dental Journal*. 41(3): 147-150.
- Sun, X., Jin, Q. and Li, X. (2016). Physical quality changes of pre-cooked shrimps during frozen storage. *Advances in Engineering Research*. 63: 76-81.
- Supardy, R., Afifah, N., Ibrahim, D., Sulaiman, S.F. and Zakaria, N.Z. (2012). Inhibition of *Klebsiella pneumoniae* ATCC 13883 by hexane extract of *Halimeda discoidea* (Decaisne) and the identification of its potential bioactive compounds. *Journal of Microbiology and Biotechnology*. 22(6): 872-881.
- Suraj, S. K. and Chaterjee, P. (2015). Antifungal activity of 50% aqueous-ethanolic extract of leaves of *Calotropis procera* R. Br. *International Journal of Scientific and Research Publications*. 5(11): 230-235.

- Sutrisna E. M., Tanti Azizah, S. and Lutfiana Dewi, I. (2016). Antidiabetic activity of ethanolic extract of *Eugenia polyantha* wight leaf from Indonesia in diabetic rat wistar strain induced by alloxan. *Asian Journal of Pharmaceutical and Clinical Research*. 9(1): 374-376.
- Swamy, T.A., Obey, J. and Mutuku, N.C. (2013). Phytochemical analysis of *Vernonia adoensis* leaves and roots used as a traditional medicinal plant in Kenya. *International Journal of Pharmaceutical and Biological*. 3: 46-52.
- Swamy, M. K., Sinniah, U. R and Akhtar, M. S. (2015). *In vitro* pharmacological activities and GC-MS analysis of different solvent extracts of *Lantana camara* leaves collected from tropical region of Malaysia. *Evidence-Based Complementary and Alternative Medicine*. 2015: 1-9.
- Syahmi, A.R.M., Vijayarathna, S., Sasidharan, S., Latha, L.Y., Kwan, Y.P., Lau, Y.L., Shin, L.N. and Chen, Y. (2010). Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* Jacq. (oil palm leaf) methanol extract. *Molecules*. 15: 8111-8121.
- Talcott, S.T., Percival, S.S., Pittet-Moore, J. and Celoria, C. (2003). Phytochemical composition and antioxidant stability of fortified yellow passion fruit (*Passiflora edulis*). *Journal of Agricultural and Food Chemistry*. 51: 935-941.
- Talimkhani, A. and Mashak, Z. (2017). Prevalence and genotyping of *Helicobacter pylori* isolated from meat, milk and vegetable in Iran. *Jundishapur Journal of Microbiology*. 10(11): 2-6.
- Tamang, J. P., Dong-Hwa, S., Su-Jin, J. and Soo-Wan, C. (2016). Functional properties of microorganisms in fermented food. *Frontier in Microbiology*. 7(578): 1-13.
- Tauxe, R.V. (2002). Emerging foodborne pathogens. *International Journal of Food Microbiology*. 78(1-2): 31-41.
- Teisl, M.F. and Roe, B.E. (2010). Consumer willingness-to-pay to reduce the probability of retail foodborne pathogen contamination. *Food policy*. 35: 521-530.
- Toa, N., Jia, L. and Zhou, H. (2014). Anti-fungal activity of *Citrus reticulata* Blanco essential oil against *Penicillium italicum* and *Penicillium digitatum*. *Food Chemistry*. 153:265-271.
- Tornuk, F., Cankurt, H., Ozturk, I., Sagdic, O., Bayram, O. and Yetim, H. (2011). Efficacy of various plant hydrosols as natural food sanitizers in reducing *Escherichia coli* O157:H7 and *Salmonella* Typhimurium on fresh cut carrots and apples. *International Journal of Food Microbiology*. 148: 30-35.

- Tournas, V.H. (2005). Spoilage of vegetable crops by bacteria and fungi and related health hazards. *Critical Reviews in Microbiology*. 31(1): 33-44.
- Trofa, D., Gacser, A. and Nosanshuk, J.D. (2008). *Candida parapsilosis*, an emerging fungal pathogen. *Clinical Microbiology Review*. 21(4): 606-625.
- Tseng, Y.C., Xiong, Y.L. and Webster, C.D. (2005). The preservation of the quality of the muscle in frozen Australian red claw crayfish (*Cherax quadricarinatus*) by pre-storage anti-oxidant dipping treatments. *International Journal of Food Science & Technology*. 40(8): 841-848.
- Tsuchiya, H. (2001). Stereo specificity in membrane effects of catechins. *Chemico-Biological Interactions*. 134(1): 41-54.
- Udoh. I.P., Eleazar, C.I., Ogeneh, B.O. and Ohanu, M.E. (2015). Studies on fungi responsible for the spoilage/deterioration of some edible fruits and vegetables. *Advances in Microbiology*. 5: 285-290.
- Ukuku, D.O. and Fett, W.E. (2004). Method of applying sanitizers and sample preparation affects recovery of native microflora and *Salmonella* on whole cantaloupe surfaces. *Journal of Food Protection*. 67: 999-1004.
- Ultee, A., Bennik, M. H. J. and Moezelaar, R. (2002). The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Applied and Environmental Microbiology*. 68(4): 1561-1568.
- Ulusoy, S., Bolgelmez-Tinaz, G. and Secilmis-Canbay, H. (2009). Tocopherol, carotene, phenolic contents and antibacterial properties of rose essential oil, hydrosol and absolute. *Current Microbiology*. 59(5):554-8.
- Valdez. B. (2012). Scientific, Health and Social Aspects of the Food Industry. In Tech. ISBN 978-953-307-916-5.
- Van Acker, F.A.A., Schouten, O., Haenen, G.R.M.M., Va der Vijgh, W.J.F. and Bast, A. (2000). Flavonoids can replace  $\alpha$ -tocopherol as an antioxidant. *Federal of European Biochemical Societies Letters*. 473: 145-148.
- Ventaka raman, B. Samuel, L. A., Pardha Saradhi, M., Narashimha Rao B., Naga Vashmi Krishna A., Sudhakar, M. and Radhakrishnan, T. M. (2012). Antibacterail, antioxidant activity and GCMS analysis *Eupatorium odoratum*. *Asian Journal of Pharmaceutical and Clinical Research*. 5(2): 99-106.
- Vilgis, T.A. (2013). Texture, taste and aroma: multi-scale materials and the gastrophysics of food. *Flavour*. 2(12): 2-5.

- Vimal, M., Vijaya, P.P., Mumtaj, P. and Seema Farhath, M.S. (2013). Antibacterial activity of selected compounds of essential oils from indigenous plants. *Journal of Chemical and Pharmaceutical Research*. 5(1): 248-253.
- Vinha, A. F., Barreira<sup>1</sup>, S. V. P., Castro<sup>1</sup>, A., Costa, C. and Oliveira. M. B. P.P (2013). Influence of the storage conditions on the physicochemical properties, antioxidant activity and microbial flora of different tomato (*Lycopersicon esculentum* L.) cultivars. *Journal of Agricultural Science*. 5(2): 119-128.
- Viswanathan, P. and Kaur, R. (2001). Prevalence and growth of pathogens on salad vegetables, fruits and sprouts. *International Journal of Hygiene and Environmental Health*. 203(3): 205-213.
- Wachter, G.A, Hoffmann, J.J, Furbacher, T., Blake, M.E. and Timmermann, B.N. (1999). Antibacterial and antifungal flavanones from *Eysenhardtia texana*. *Phytochemistry*. 52: 1469-1471.
- Wahjuni, S. and Wita, I. W. (2017). Hypoglycemic and antioxidant effects of *Syzygium polyanthum* leaves extract on alloxan induced hyperglycemic Wistar rats. *Bali Medicinal Journal*. 3(3): 113-16.
- Wang, Y., Zhang, S., Yu, J., Zhang, H., Yuan, Z., Sun, Y. and Song, H. (2010). An outbreak of *Proteus mirabilis* food poisoning associated with eating stewed pork balls in brown sauce, Beijing. *Food Control*. 21(3): 302-305.
- Wani, N.S., Bhalerao, A.K., Ranaware, V.P. and Zanje, R. (2013). Formulation and evaluation of herbal sanitizer. *International Journal of PharmTech Research*. 5: 40-43.
- Westrell, T., Ciampa, N., Boelaert, F., Helwigh, B., Korsgaard, H. and Chriel, M. (2009). Zoonotic infections in Europe in 2007: A summary of the EFSA-ECDC annual report. *Euro Surveill*. 14.
- Whitman, W.C., Johnson, W.M. and Tomczyk, J.A. (2005). Refrigeration and Air Conditioning Technology. 5<sup>th</sup> Edition. New Delhi: Thomson Publications.
- Wiedmann, M. and Roberts, A.J. (2003). Pathogen, host and environmental factors contributing to the pathogenesis of listeriosis. *Cellular Molecular Life Science*. 60: 904-918.
- Wilson, L. (2011). On-farm Food Safety: Cleaning and Sanitizing Guide. *Food Science and Human Nutrition*. pp. 1-4.
- Witkowska, A.M., Hickey, D.K., Alonso-Gomez, M. and Wilkinson, M. (2013). Antimicrobial activity of commercial herb and spices extracts against selected foodborne bacteria. *Journal of Food Research*. 2(4): 37-54.

- Wong, K.C., Hag Ali, D.M. and Boey, P.L. (2012). Chemical constituents and antibacterial activity of *Melastoma malabathricum* L. *Natural Product Research*. 26: 609-18.
- World Health Organization (WHO). (1996). International stability testing: guidelines for stability of pharmaceutical products containing well established drug substances in conventional dosage forms. Switzerland: WHO.
- Wu, J., Lin, L. and Chau, F. T. (2001). Ultrasound-assisted extraction of ginseng saponins from ginseng roots and cultured ginseng cells. *Ultrason Sonochem*. 8(4): 347-352.
- Wu, S., Wu, Q., Zhang, J., Chen, M. and Yan, Z. (2015). Prevalence, antibiotic resistance and genetic diversity of *Listeria monocytogenes* isolated from retail ready-to-eat foods in China. *Food Control*. 47: 340-347.
- Yamaguchi, M.U., Rampazzo, R.C.P., Yamada- Ogatta, S.F., Nakamura, C., V.N., Ueda-Nakamura, T. and Dias, B.P. (2007). Yeasts and filamentous fungi in bottled mineral water and tap water from municipal supplies. *Brazilian Archives of Biology and Technology*. 50: 1-9.
- Yaashikaa, P.R., Saravanan, A. and Kumar, P.S. (2016). Isolation and identification of *Vibrio cholerae* and *Vibrio parahaemolyticus* from prawn (*Penaeus monodon*) seafood: Preservation strategies. *Microbial Pathogenesis*. 99: 5-13.
- Yang, S.E., Yu, R.C. and Chou, C.C. (2001). Influence of holding temperature on the growth and survival of *Salmonella* spp. and *Staphylococcus aureus* and the production of staphylococcal enterotoxin in egg products. *International Journal of Food Microbiology*. 63: 99-107.
- Yoshiki, K., Hui-Kang, W., Tsuneatsu, N., Susumu, K., Ichiro, Y., Toshihiro, F., Takashi, Y., Cosentino, L.M., Mutsuo, K., Hikaru, O., Yasumasa, I., Chang-Qi Hu, Yeh, E. and Kuo-Hsiung, L. (1998). Anti- AIDS agents. 30. Anti-HIV activity of oleanolic acid, pomolic acid and structurally related triterpenoids. *Journal of Natural Products*. 61: 1090-1095.
- Yousef, A.E. and Balasubramaniam, V.M. (2013). Physical methods of food preservations. *Food Microbiology, Fundamental and Frontiers*, 4<sup>th</sup> Edition. American Society for Microbiology Press. Washington, D. C.
- Yu, T. and Jiang, X. (2014). Prevalence and characterization of *Listeria monocytogenes* isolated from retail food in Henan, China. *Food Control*. 37: 228-231.
- Yusoff, N.A.H., Noor, N.F. and Rukayadi, Y. (2015). Effects of *Cosmos caudatus* Kunth. (*Ulam raja*) extract on microflora in raw chicken meat. *International Journal of Current Microbiology and Applied Sciences*. 4(2): 426-435.



- Yuste, J. and Fung, D.Y.C. (2002). Inactivation of *Listeria monocytogenes* Scott A 49594 in apple juice supplemented with cinnamon. *Journal of Food Protection*. 65: 1663-1666.
- Zainin, N.S., Lau, K.Y., Zakaria, M., Son, R., Abdull Razis, A.F. and Rukayadi, Y. (2013). Antibacterial activity of *Boesenbergia rotunda* (L.) Mansf. A. extract against *Escherichia coli*. *International Food Research Journal*. 20(6): 3319-3323.
- Zarinah, Z., Maaruf, A. G., Nazaruddin, R., Wong, W. W. W. and Xuebing, X. (2014). Antioxidant, antimicrobial activity and *in-vitro* cytotoxicity screening study of Pili nut oil. *International Food Research Journal*. 21(1): 309-316.
- Zhang, H. N., Hou, P. B., Chen, Y. Z., Yu, M., Li, X. P., Hui, L. V., Mei, W. Tan, H. L. and Bi, Z. W. (2016). Prevalence of foodborne pathogens in cooked meat and seafood from 2010 to 2013 in Shandong Province, China. *Iran Journal of Public Health*. 45(12): 1577-1585.
- Zhang, J., Sun, H., Chen, S., Zeng, L., and Wang, T. (2017). Anti-fungal activity, mechanism studies on  $\alpha$ -Phellandrene and nonanal against *Penicillium cyclopium*. *Botanical Studies*. 58(13): 2-9.
- Zhang, S. and J.M. Faber. (1996). The effects of various disinfectants against *Listeria monocytogenes* on fresh-cut vegetables. *Food Microbiology*. 13: 311- 321.
- Zhao, C., Beilei, G., de Villena, J., Sudler, R., Yeh, E., Zhao, S., White, D. G., Wagner, D. and Meng, J. (2001). Prevalence of *Campylobacter* spp., *Escherichia coli*, and *Salmonella* serovars in retail chicken, turkey, pork, and beef from the Greater Washington, D.C., Area. *Applied and Environmental Microbiology*. 67(12): 5431-5436.
- Zheng, W.F., Tan, R.X., Yang, L. and Liu, Z.L. (2000). Two flavones from *Artemisia giraldii* and their antimicrobial activity. *Planta Medica*. 62: 160-2.
- Zhou, K., Zhou, W., Li, P., Liu, G., Zhang, J. and Dai, Y. (2008). Mode of action of pentocin, an antilisteria bacteriocin produced by *Lactobacillus pentosus* from Chinese traditional ham. *Food Control*. 19: 817-822.
- Zhou, G.H., Xu, X.L. and Liu, Y. (2010). Preservation technologies for fresh meat - A review. *Meat Science*. 86: 119-128.
- Zulfa, Z., Chia, C. T. and Rukayadi, Y. (2016). *In vitro* antimicrobial activity of *Cymbopogon citratus* (lemongrass) extracts against selected foodborne pathogens. *International Food Research Journal*. 23(3): 1262-1267.