

Diversity of macrobenthos in the mangrove forest of Kuala Sibuti, Miri, Sarawak

Abu Hena M.K.^{a,*}, Idris M.H.^b, Khairul R.M.Y.^a, Bhuiyan M.K.A.^c, Nesarul M.H.^a and Uttam K.^a

^aDepartment of Animal Science and Fishery, Faculty of Agriculture and Food Sciences, Universiti Putra Malaysia Bintulu Sarawak Campus, 97008 Bintulu, Sarawak, Malaysia.

^bSchool of Fisheries and Aquaculture Sciences, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu Darul Iman, Malaysia.

^cFaculty of Marine and Environmental Sciences, University of Cadiz, Puerto Real Campus, Puerto Real, Spain.

Abstract

The diversity of mangrove macrobenthos assemblages at mangrove forest of Kuala Sibuti, Miri, Sarawak was investigated for northeast monsoon and inter monsoon season. The aim of this study was to determine the diversity and composition of macrobenthos in the Kuala Sibuti mangrove forest. Sampling was carried out at 3 different locations during each monsoon. In macrobenthos assemblages, bivalves (49%) was the most diverse and abundant followed by gastropods (23%), polychaetes (17%) and crustaceans (10%). A total of 25 species of macrobenthos have been recorded during the study period. Studies found that sediment texture plays an important role in the ecology of benthic invertebrates. Polychaete was found be influenced by the percentage of soil texture, while temperature and pH have had little influence on the macrobenthos assemblages in this mangrove forest ecosystem.

Keywords: Mangrove, macrobenthos, Kuala Sibuti, Sarawak.

*Corresponding author: hena71@yahoo.com