



UNIVERSITI PUTRA MALAYSIA

***EPIDEMIOLOGY AND ETIOLOGY OF ganoderma UPPER AND BASAL
STEM ROT IN OIL PALM (elaeis guineensis JACQ.) ON PEAT IN
SARAWAK, MALAYSIA***

MOHD RAKIB MOHD RASHID

FSPM 2015 2



**EPIDEMIOLOGY AND ETIOLOGY OF *Ganoderma* UPPER AND BASAL
STEM ROT IN OIL PALM (*Elaeis guineensis* JACQ.) ON PEAT IN
SARAWAK, MALAYSIA**

By

MOHD RAKIB BIN MOHD RASHID

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of
Philosophy**

March 2015

All material contained within the thesis, including without limitation text, logo, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Doctor of Philosophy

EPIDEMIOLOGY AND ETIOLOGY OF *Ganoderma* UPPER AND BASAL STEM ROT IN OIL PALM (*Elaeis guineensis* JACQ.) ON PEAT IN SARAWAK, MALAYSIA

By

MOHD RAKIB BIN MOHD RASHID

March 2015

Chairman : Joseph Bong Choon Fah, PhD
Faculty : Faculty of Agriculture and Food Sciences (Bintulu)

Oil palm (*Elaeis guineensis*) is the world's most important oil producing crop which is mostly cultivated in Southeast Asia. The sustainability of oil palm is threatened by *Ganoderma* species that cause upper stem rot (USR) and basal stem rot (BSR). Basal stem rot is a prominent disease of oil palm, while little is known about USR. Lack of knowledge in terms of epidemiology and etiology of the diseases may lead to inaccurate disease control strategies. A study was conducted in oil palm plantations on peat in Sarawak, Malaysia, namely, Sessang Estate in Betong and Sungai Balim Estate in Miri. A total of 46 isolates of *Ganoderma* species from the USR- and BSR-infected palms were identified using multiplex PCR, and their *in-vitro* cultural and basidiospore morphological characteristics were investigated. It was found that both USR and BSR were associated with similar pathogens, where *G. zonatum* (71.7%) was dominant, followed by *G. boninense* (26.1%) and *G. miniatocinctum* (2.2%). This suggests that *G. zonatum* may have played a more vital role in the epidemiology of the disease than previously believed. All samples were found incompatible based on somatic compatibility test which indicated genetic heterogeneity of *Ganoderma* species in oil palm plantation and was suggestive of disease spread via spore dispersal that generated new genetically distinct individuals. There were also significant variations within and between *Ganoderma* species and no distinct relationship were found between species, disease types or geographical origins in terms of their cultural morphology and basidiospore characteristics. This suggests that any of the *Ganoderma* isolates found in this study have similar possibility of either causing USR or BSR. The pathogenic capability and aggressiveness of *Ganoderma* species were assessed based on external and internal infection symptoms using artificial inoculation on oil palm seedlings throughout 24 weeks period. All 46 *Ganoderma* samples tested were confirmed to be pathogenic to oil palm. Among them, *G. zonatum* isolated from USR-infected palms showed superior aggressiveness as compared with others with average foliar symptoms severity (FSS), disease severity index (DSI), stem bole lesion and primary root decay up to 33.18%, 43.78%, 42.20% and 48.55%, respectively. Hence, besides the prominent *G. boninense* and BSR, more studies should emphasize on *G. zonatum* and USR for more effective disease management since it was found as an increasingly important pathogen and disease in oil palm

plantations, especially on peat areas in Sarawak. The infection of USR and BSR were mainly confirmed based on appearance of *Ganoderma* basidiomata on the stem, while the point of infection on the stem was the distinctive characteristic between them, where infection of *Ganoderma* species about one meter above ground indicated USR, while that at the base indicating BSR. USR and BSR disease coexisted in both sites, with lower USR (1.14-1.32%) as compared with BSR (3.48-4.01%) in Betong, while higher USR (3.66-6.02%) as compared with BSR (0.87-3.78%) in Miri throughout the study period. This also indicated that *Ganoderma* species infection in Miri was relatively higher than that in Betong. Geostatistical analysis of the *Ganoderma* species distribution revealed generally weak spatial dependence ($>75\%$), which indicated that the distribution was random, and related to *Ganoderma* species spread via basidiospore dispersal at greater distance, instead of root-to-root infection. It was revealed that spatial distribution of *Ganoderma* species was at relatively greater density in Miri (15.1-100%) as compared with Betong (15.1-30%). Although the distribution of *Ganoderma* species were random, the hotspot patterns generated allow site specific disease management as in precision agriculture and aid in sampling programmes for further investigations on the factors attributed to the patterns. Studies on macro- and micronutrients composition in relation to spatial distribution of *Ganoderma* species revealed that the foliar macro- and micronutrient elements affected the infection of *Ganoderma* species. Based on the four trials conducted at the two study sites, it was found that higher level of phosphorus (P), and lower and deficient level of copper (Cu) and zinc (Zn) in oil palms were associated with higher *Ganoderma* infection. In addition, lower and deficient level of Cu and Zn also could be related to higher *Ganoderma* infection in Miri as compared with Betong. Furthermore, there was no distinct factor in terms of soil and foliar chemical properties between USR and BSR. Therefore, this finding could be used as general guide to control *Ganoderma* species by giving more concern on manipulation of P, Cu and Zn level in oil palm plantations generally, and specifically in the two sites in this study for site-specific disease management as in precision agriculture.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan ijazah Doktor Falsafah

**EPIDEMIOLOGI DAN ETIOLOGI REPUT ATAS DAN PANGKAL BATANG
Ganoderma DALAM KELAPA SAWIT (*Elaeis guineensis* JACQ.) PADA
GAMBUT DI SARAWAK, MALAYSIA**

Oleh

MOHD RAKIB BIN MOHD RASHID

Mac 2015

Pengerusi : Joseph Bong Choon Fah, PhD
Fakulti : Fakulti Sains Pertanian dan Makanan (Bintulu)

Kelapa sawit (*Elaeis guineensis*) adalah tanaman penghasil minyak terpenting di dunia yang kebanyakannya ditanam di Asia Tenggara. Kelestarian kelapa sawit diancam oleh species *Ganoderma* yang menyebabkan reput atas batang (USR) dan reput pangkal batang (BSR). Reput pangkal batang adalah penyakit terkenal pada kelapa sawit, manakala sedikit yang diketahui tentang USR. Kekurangan pengetahuan dari segi epidemiologi dan etiologi penyakit-penyakit tersebut boleh menyebabkan strategi kawalan penyakit yang kurang tepat. Satu kajian telah dijalankan di ladang-ladang kelapa sawit di tanah gambut di Sarawak, Malaysia, iaitu, Estet Sessang di Betong dan Estet Sungai Balim di Miri. Sejumlah 46 pencilan spesies *Ganoderma* daripada pokok-pokok sawit dijangkiti USR dan BSR telah dikenal pasti menggunakan multiplex PCR, dan ciri-ciri morfologinya *in-vitro* kultur dan basidiospora telah dikaji. Didapati bahawa kedua-dua USR dan BSR dikaitkan dengan pathogen yang sama, di mana *G. zonatum* (71.7 %) adalah dominan, diikuti oleh *G. boninense* (26.1 %) dan *G. miniatocinctum* (2.2%). Ini menunjukkan bahawa *G. zonatum* mungkin memainkan peranan yang lebih penting dalam epidemiologi penyakit tersebut daripada yang dipercayai sebelum ini. Semua sampel didapati tidak serasi berdasarkan ujian keserasian somatic yang menunjukkan kepelbagaian genetik spesies *Ganoderma* di ladang kelapa sawit dan menandakan penyakit disebarkan melalui penyebaran spora yang menghasilkan individu baru yang genetiknya berbeza. Terdapat juga perbezaan yang ketara dalam dan antara spesies *Ganoderma* dan tidak ada hubungan yang berbeza ditemui antara spesies, jenis penyakit atau asal-usul geografi dari segi ciri-ciri morfologi kultur dan basidiospora mereka. Ini menunjukkan bahawa mana-mana pencilan *Ganoderma* yang ditemui dalam kajian ini mempunyai kemungkinan yang sama, iaitu samada menyebabkan USR atau BSR. Keupayaan patogenik dan keagresifan spesies *Ganoderma* dinilai berdasarkan gejala jangkitan luaran dan dalaman menggunakan inokulasi buatan pada anak pokok kelapa sawit sepanjang tempoh 24 minggu. Kesemua 46 sampel *Ganoderma* yang diuji disahkan sebagai patogen kepada kelapa sawit. Antara mereka, pencilan *G. zonatum* dari pokok sawit yang dijangkiti USR menunjukkan keagresifan unggul berbanding yang lain-lain dengan tahap gejala foliar (FSS), indeks tahap penyakit (DSI), reput batang dan akar utama, masing-masing sehingga 33.18 %, 43.78 %, 42.20 % dan

48.55 %. Justeru, selain *G. boninense* dan BSR yang terkenal, lebih banyak perhatian perlu diberikan untuk kajian terhadap *G. zonatum* dan USB untuk pengurusan penyakit yang lebih berkesan kerana ia telah didapati sebagai patogen dan penyakit yang semakin penting di lading-ladang kelapa sawit, terutama di kawasan-kawasan tanah gambut di Sarawak. Jangkitan USB and BSR disahkan terutamanya berdasarkan kemunculan basidiomata *Ganoderma* pada batang, manakala bahagian jangkitan pada batang adalah ciri tersendiri yang membezakan di antara penyakit tersebut, di mana jangkitan species *Ganoderma* kira-kira satu meter dari aras tanah menunjukkan USB, manakala di pangkal menunjukkan BSR. Penyakit USB dan BSR wujud bersama di kedua-dua tapak kajian, dengan USB lebih rendah (1.14-1.32%) berbanding BSR (3.48-4.01%) di Betong, manakala USB lebih tinggi (3.66-6.02%) berbanding BSR (0.87-3.78%) di Miri sepanjang tempoh kajian. Ini juga menunjukkan bahawa serangan species *Ganoderma* di Miri adalah lebih tinggi berbanding di Betong. Analisis geostatistik pada taburan species *Ganoderma* umumnya menunjukkan pergantungan lemah spatial (>75%), yang menunjukkan bahawa taburannya adalah rawak dan dikaitkan dengan perebakan species *Ganoderma* melalui penyebaran basidiospore pada jarak yang lebih jauh, dan bukannya jangkitan akar-ke-akar. Ditunjukkan bahawa taburan species *Ganoderma* di Miri adalah lebih padat (15.1-100%) berbanding di Betong (15.1-30%). Walaupun taburan spesies *Ganoderma* adalah rawak, corak titik panas yang dijana membolehkan pengurusan penyakit tapak spesifik seperti dalam pertanian persis dan membantu dalam program persampelan untuk kajian lanjut pada factor-faktor yang dikaitkan dengan corak-corak taburan tersebut. Kajian ke atas komposisi makro dan mikro-nutrien berhubung dengan taburan species *Ganoderma* menunjukkan unsur-unsur makro- dan mikronutrien foliar mempengaruhi serangan species *Ganoderma*. Berdasarkan empat percubaan yang telah dijalankan di dua tapak kajian, didapati bahawa tahap fosforus (P) yang lebih tinggi, dan tahap kuprum (Cu) dan zink (Zn) yang lebih rendah dan kekurangannya dalam kelapa sawit dikaitkan dengan serangan *Ganoderma* yang lebih tinggi. Sebagai tambahan, tahap yang lebih rendah dan kekurangan Cu dan Zn juga boleh dikaitkan dengan serangan *Ganoderma* yang lebih tinggi di Miri berbanding di Betong. Tambahan pula, tidak terdapat sebarang factor berbeza dari segi sifat-sifat kimia foliar dan tanah antara USB dan BSR. Oleh itu, penemuan ini boleh digunakan sebagai panduan umum untuk mengawal spesies *Ganoderma* dengan memberi perhatian yang lebih pada manipulasi P, Cu dan Zn tahap di ladang-ladang kelapa sawit secara amnya, dan khususnya di dua tapak kajian ini untuk pengurusan penyakit khusus kawasan seperti dalam pertanian persis.

ACKNOWLEDGEMENTS

In the name of God, the most gracious, the most merciful. My deepest gratitude goes to my respected supervisory committee chairperson, Assoc. Prof. Dr. Joseph Bong Choon Fah, and other members, namely, Dr. Khairulmazmi Ahmad and Dr. Idris Abu Seman for their expertise, advice, guidance, contributions, supports and encouragements during my study. I also would like to express my gratitude to my beloved family, especially my father, Mr. Mohd Rashid and mother, Mrs. Hasina for their prayers and endless love. Not to forget, thanks to all the supportive staffs in Universiti Putra Malaysia Bintulu Campus (UPMKB), Malaysian Palm Oil Board (MPOB), and Sarawak Oil Palm Berhad (SOPB). Special thanks to all my colleagues and friends for being my “backbone” throughout my study.

I also would like to thank the MPOB under the Malaysian Ministry of Plantation Industries and Commodities (MPIC) for financial assistance through research grant, and the Malaysian Ministry of Science, Technology and Innovation (MOSTI) for National Science Fellowship (NSF) scholarship award.

I certify that a Thesis Examination Committee has met on 10 March 2015 to conduct the final examination of Mohd Rakib bin Mohd Rashid on his thesis entitled "Epidemiology and Etiology of *Ganoderma* Upper and Basal Stem Rot in Oil Palm (*Elaeis guineensis* Jacq.) on Peat in Sarawak, Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Ahmed Osumanu Haruna, PhD

Associate Professor
Faculty of Agriculture and Food Sciences
Universiti Putra Malaysia
(Chairman)

Zainal Abidin bin Mior Ahmad, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Julie Flood, PhD

Commonwealth Agricultural Bureau International (CABI)
United Kingdom
(External Examiner)

ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Joseph Bong Choon Fah, PhD

Associate Professor
Faculty of Agriculture and Food Sciences
Universiti Putra Malaysia
(Chairman)

Khairulmazmi bin Ahmad, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Idris bin Abu Seman, PhD

Head of Ganoderma and Disease Research for Oil Palm Unit
Biology Research Division
Malaysian Palm Oil Board (MPOB)
(Member)

BUJANG KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: 25 May 2015

Name and Matric No.: Mohd Rakib bin Mohd Rashid / GS25424

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: Joseph Bong Choon Fah

Signature: _____
Name of
Member of
Supervisory
Committee: Khairulmazmi bin Ahmad

Signature: _____
Name of
Member of
Supervisory
Committee: Idris bin Abu Seman

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xvii
LIST OF PLATES	xxii
LIST OF ABBREVIATIONS	xxiv
 CHAPTER	
 1 INTRODUCTION	 1
1.1 Background	1
1.2 Justification	2
1.3 Objectives	3
 2 LITERATURE REVIEW	 4
2.1 The Oil Palm	4
2.1.1 The origin of oil palm	4
2.1.2 Oil palm as oil producing crop	4
2.1.3 Morphology of oil palm	5
2.1.4 Nutrient content of oil palm	8
2.2 Overview of Oil Palm Industry in Malaysia	9
2.3 Oil Palm Cultivation on Peat Land	12
2.3.1 Distribution of peat land and oil palm cultivation on peat	12
2.3.2 Management of peat soil for oil palm cultivation	12
2.3.3 Nutrient content of peat soil	15
2.4 Oil Palm Diseases	16
2.4.1 Basal stem rot (BSR)	16
2.4.2 Upper stem rot (USR)	18
2.5 <i>Ganoderma</i> Species	19
2.5.1 Taxonomy of <i>Ganoderma</i> species	19
2.5.2 Genetic diversity of <i>Ganoderma</i> species	19
2.5.3 Morphological diversity of <i>Ganoderma</i> species	21
2.5.4 Detection and identification of <i>Ganoderma</i> species in oil palm	22
2.5.5 Pathological studies of <i>Ganoderma</i> species in oil palm	23
2.6 Relationship between Nutrients and Incidence of <i>Ganoderma</i> Species in Oil Palm	24

3	GENERAL MATERIALS AND METHODS	26
3.1	Description of Study Sites	26
3.2	Studies on the <i>Ganoderma</i> species of USR and BSR	34
3.2.1	Identification of USR- and BSR-infected oil palms	34
3.2.2	Isolation and collection of <i>Ganoderma</i> species	34
3.2.3	Identification of <i>Ganoderma</i> species	34
3.2.4	Genetic compatibility assessment of <i>Ganoderma</i> species	34
2.2.5	<i>In-vitro</i> cultural and basidiospore morphological characteristics of <i>Ganoderma</i> species	34
3.2.6	Pathogenicity and aggressiveness of <i>Ganoderma</i> species	35
3.3	Studies on the Occurrence of USR and BSR, and the Distribution (Hotspot Pattern) of <i>Ganoderma</i> Species	35
3.4	Peat Soils and Foliar Sampling, and Chemical Analysis	35
4	GENETIC AND MORPHOLOGICAL DIVERSITY OF <i>Ganoderma</i> spp. ISOLATED FROM UPPER- AND BASAL STEM ROT INFECTED OIL PALM (<i>Elaeis guineensis</i>)	36
4.1	Introduction	36
4.2	Materials and Methods	37
4.2.1	Sample collection of <i>Ganoderma</i> species	37
4.2.2	Isolation of <i>Ganoderma</i> species on <i>Ganoderma</i> selective medium (GSM)	37
4.2.3	Identification of <i>Ganoderma</i> species using multiplex PCR	38
4.2.4	Assessment of genetic similarity of <i>Ganoderma</i> species using somatic compatibility test	39
4.2.5	Assessment of <i>in-vitro</i> cultural characteristics of <i>Ganoderma</i> species	39
4.2.6	Assessment of basidiospore characteristics of <i>Ganoderma</i> species	40
4.3	Results and Discussion	43
4.3.1	Genetic diversity of <i>Ganoderma</i> species	43
4.3.2	<i>In-vitro</i> cultural and basidiospore morphological characteristics of <i>Ganoderma</i> species	47
4.4	Conclusions	55

5	AGGRESSIVENESS OF <i>Ganoderma</i> spp. ISOLATED FROM UPPER- AND BASAL STEM ROT INFECTED OIL PALMS (<i>Elaeis guineensis</i>) ON OIL PALM SEEDLINGS	56
5.1	Introduction	56
5.2	Materials and Methods	57
5.2.1	<i>Ganoderma</i> samples and planting materials	57
5.2.2	Artificial inoculation of <i>Ganoderma</i> species on oil palm seedlings	57
5.2.3	Assessment of pathogenicity trial	58
5.2.4	Data analysis	60
5.3	Results and Discussion	60
5.3.1	External and internal symptoms of <i>Ganoderma</i> -infected oil palm seedlings	60
5.3.2	Aggressiveness of different <i>Ganoderma</i> isolates from USR and BSR-infected oil palms	65
5.4	Conclusions	82
6	SYMPTOMATOLOGY, OCCURRENCE AND SPATIO-TEMPORAL DISTRIBUTION OF UPPER AND BASAL STEM ROT, AND HOTSPOT ANALYSIS OF <i>Ganoderma</i> spp. OF THE DISEASES IN OIL PALM (<i>Elaeis guineensis</i>)	83
6.1	Introduction	83
6.2	Materials and Methods	84
6.2.1	Study site	84
6.2.2	Field symptomatology of USR and BSR	84
6.2.3	Data collection	84
6.2.4	Hotspot analysis	84
6.3	Results and Discussion	86
6.3.1	Field symptomatology of USR and BSR	86
6.3.2	Occurrence and Spatio-temporal distribution of USR and BSR	88
6.3.3	Hotspot pattern of <i>Ganoderma</i> species distribution associated with USR and BSR	96
6.4	Conclusions	101
7	MACRO- AND MICRONUTRIENTS IN RELATION TO SPATIAL DISTRIBUTION OF <i>Ganoderma</i> spp. IN OIL PALM (<i>Elaeis guineensis</i>) PLANTATIONS ON PEAT	102
7.1	Introduction	102
7.2	Materials and Methods	103
7.2.1	Peat soil and foliar samples collection and preparation	103

	7.2.2	Soil and foliar chemical properties analysis	103
	7.2.3	Data analysis	109
7.3		Results and Discussion	109
	7.3.1	Nitrogen (N)	109
	7.3.2	Phosphorus (P)	112
	7.3.3	Potassium (K)	114
	7.3.4	Calcium (Ca)	117
	7.3.5	Magnesium (Mg)	119
	7.3.6	Copper (Cu)	122
	7.3.7	Zinc (Zn)	124
	7.3.8	Boron (B)	127
	7.3.9	Manganese (Mn)	129
	7.3.10	Iron (Fe)	132
	7.3.11	General discussion on the relationship between soil and foliar chemical properties to <i>Ganoderma</i> species distribution	134
	7.3.12	Comparison between USR- and BSR-infected palms in terms of soil and foliar chemical properties	135
7.4		Conclusions	140
8		SUMMARY, GENERAL CONCLUSIONS AND RECOMMENDATIONS FOR FUTURE RESEARCH	141
		REFERENCES	143
		APPENDICES	162
		BIODATA OF STUDENT	184
		LIST OF PUBLICATIONS	185

LIST OF TABLES

Table		Page
2.1	World's major vegetable oil producing crops (million tonnes)	5
2.2	Yield of oil producing crops (tonnes per hectare per year)	5
2.3	Nutrient concentration in frond 17 associated with deficiency, optimum and excess in mature oil palms	9
2.4	Crude palm oil (CPO) production by country (million tonnes)	10
2.5	Malaysia: Oil palm planted area by states (hectare)	10
2.6	Malaysia: Change in planted area (hectare) across three years (2010 to 2013)	11
2.7	Sarawak: Oil palm planted area by Division (hectare)	11
2.8	General peat soil nutrient content in oil palm plantations	15
2.9	Basal stem rot (BSR) incidence in relation to palm age in Peninsular Malaysia	17
2.10	Basal stem rot (BSR) incidence in oil palm in relation to previous crop	18
3.1	Description summaries of study sites in Betong and Miri	30
3.2	Fertilization programme in Betong study plot over 2010-2012	32
3.3	Fertilization programme in Miri study plot over 2010-2012	33
4.1	List of <i>Ganoderma</i> samples from USR- and BSR- infected oil palms in Betong and Miri	37
4.2	Cultural characters and their corresponding codes used to describe <i>Ganoderma</i> isolates for assessment of <i>in-vitro</i> cultural characteristics	40

4.3	<i>In-vitro</i> cultural characteristics of <i>Ganoderma</i> isolates from upper stem rot (USR) and basal stem rot (BSR) infected oil palms in Betong assessed on potato dextrose agar (PDA)	48
4.4	<i>In-vitro</i> cultural characteristics of <i>Ganoderma</i> isolates from upper stem rot (USR) and basal stem rot (BSR) infected oil palms in Miri assessed on potato dextrose agar (PDA)	49
4.5	Basidiospore characteristics of <i>Ganoderma</i> isolates from upper stem rot (USR) and basal stem rot (BSR) infected palms in Betong	52
4.6	Basidiospore characteristics of <i>Ganoderma</i> isolates from upper stem rot (USR) and basal stem rot (BSR) infected palms in Miri	53
5.1	Numerical disease classes and their corresponding symptoms on oil palm seedling	59
5.2	Disease incidence (DI) of <i>Ganoderma</i> isolates from upper stem rot (USR) infected palms in Betong and Miri on oil palm seedlings	66
5.3	Disease incidence (DI) of <i>Ganoderma</i> isolates from basal stem rot (BSR) infected palms in Betong and Miri on oil palm seedlings	67
5.4	Foliar symptoms severity (FSS) of <i>Ganoderma</i> isolates from upper stem rot (USR) infected palms in Betong and Miri on oil palm seedlings	69
5.5	Foliar symptoms severity (FSS) of <i>Ganoderma</i> isolates from basal stem rot (BSR) infected palms in Betong and Miri on oil palm seedlings	70
5.6	Disease severity index (DSI) of <i>Ganoderma</i> isolates from upper stem rot (USR) infected palms in Betong and Miri on oil palm seedlings	71
5.7	Disease severity index (DSI) of <i>Ganoderma</i> isolates from basal stem rot (BSR) infected palms in Betong and Miri on oil palm seedlings	72
5.8	Stem bole and primary root necrosis at 24 weeks after inoculation with <i>Ganoderma</i> isolates from upper stem rot (USR) infected palms in Betong and Miri on oil palm seedlings	73

5.9	Stem bole and primary root necrosis at 24 weeks after inoculation with <i>Ganoderma</i> isolates from basal stem rot (BSR) infected palms in Betong and Miri on oil palm seedlings	74
5.10	Pearson's correlation analysis between variables tested to assess disease infection of <i>Ganoderma</i> species on oil palm seedlings	81
6.1	Summary of spherical semivariogram model of the hotspot analysis for <i>Ganoderma</i> species distribution in Betong and Miri	97
7.1	Summary of soil chemical properties analytical method	106
7.2	Summary of foliar chemical properties analytical method	107

LIST OF FIGURES

Figure		Page
2.1	Oil palm seedling (2 months after planting). SL: soil level, S: shell, Co: cotyledon (haustorium), P: plumule, R: radicle, Pr: primary root, Ad: adventitious roots	6
2.2	Leaf or frond of oil palm. (1) Cross-section of rachis. (2) Cross-section of petiole. AD: adaxial face, AB: abaxial face, LF: lateral face, SP: spine. (3) Cross-section of leaf viewed end-on, showing two-ranked insertion of leaflets. (4) Diagram of oil palm leaf (frond). TL: terminal pair of ovate leaflets, LL: leaflets, RA: rachis, VL: leaflets with vestigial laminae, PE: petiole, SP: spine. (5) Central portion of rachis from above, showing irregular leaflet insertion. BS: basal swelling, LR: lower rank leaflet, LF: lateral face. (6) Developing leaf, enclosed within apex of palm. LB: leaf base, FL: future leaf, SH: leaf sheath completely encircling apex	7
2.3	Phyllotaxis of oil palm fronds showing spiral arrangement of the fronds (Frond 1, 9, 17, 25 and 33)	8
2.4	Distribution of oil palm and peat in Sarawak, Malaysia	13
2.5	Distribution of peat lands in Sarawak	14
2.6	General life cycle of fruiting body forming basidiomycete	20
2.7	General macro-morphological characteristics of <i>Ganoderma</i>	22
3.1	Study sites located in Betong and Miri, Sarawak, Malaysia	27
3.2	Map of the plantation in Betong with 15 hectares of study plot	28
3.3	Map of the plantation in Miri with 16 hectares of study plot	29
3.4	Daily mean precipitation and temperature for two years period (year 2011 to 2012) obtained from (A) Sibul weather station and (B) Miri weather station	31
4.1	Species distribution of <i>Ganoderma</i> in the study sites. (A) In Betong, (B) In Miri, (C) Overall average	45

4.2	Dendrogram (UPGMA) generated from 25 cultural morphological characters of <i>Ganoderma</i> species isolated from upper stem rot (USR) and basal stem rot (BSR) infected oil palms in the plantation in Betong and Miri using simple matching coefficient	50
4.3	Distribution of spore shape index (SSI) on spore length for 46 isolates of <i>Ganoderma</i> species isolated from upper stem rot (USR) and basal stem rot (BSR) infected palms in Betong and Miri	54
5.1	Illustration showing diagram of the artificial inoculation of <i>Ganoderma</i> by rubber wood block (RWB) sitting technique on oil palm seedling	58
5.2	Mean (\pm standard error) percentage of disease incidence (DI) of all <i>Ganoderma</i> -inoculated oil palm seedlings based on four different evaluations. Means with different alphabet were significantly different at $p < 0.05$ by DNMR	62
5.3	Mean (\pm standard error) foliar symptoms severity (FSS) at four weeks intervals throughout 24 weeks after inoculation with <i>G. zonatum</i> and <i>G. boninense</i> from upper stem rot (USR) and basal stem rot (BSR) infected oil palms. (A) Between <i>G. zonatum</i> from USR and BSR, (B) Between <i>G. boninense</i> from USR and BSR, (C) Between <i>G. zonatum</i> and <i>G. boninense</i> from USR, (D) Between <i>G. zonatum</i> and <i>G. boninense</i> from BSR	77
5.4	Mean (\pm standard error) disease severity index (DSI) at four weeks intervals throughout 24 weeks after inoculation with <i>G. zonatum</i> and <i>G. boninense</i> from upper stem rot (USR) and basal stem rot (BSR) infected oil palms. (A) Between <i>G. zonatum</i> from USR and BSR, (B) Between <i>G. boninense</i> from USR and BSR, (C) Between <i>G. zonatum</i> and <i>G. boninense</i> from USR, (D) Between <i>G. zonatum</i> and <i>G. boninense</i> from BSR	78
5.5	Mean (\pm standard error) stem bole necrosis and primary root necrosis at 24 weeks after inoculation with <i>G. zonatum</i> and <i>G. boninense</i> from upper stem rot (USR) and basal stem rot (BSR) infected oil palms. (A) Between <i>G. zonatum</i> from USR and BSR, (B) Between <i>G. boninense</i> from USR and BSR, (C) Between <i>G. zonatum</i> and <i>G. boninense</i> from USR, (D) Between <i>G. zonatum</i> and <i>G. boninense</i> from BSR	79
6.1	Scheme of semivariogram used for hotspot analysis	85

6.2	Upper stem rot (USR) and basal stem rot (BSR) occurrence in oil palm plantation at 6 months interval in (A) Betong in Jan. 2011, Jul. 2011 and Jan. 2012, (B) Miri in Mar. 2011, Sep. 2011 and Mar. 2012	89
6.3	Distribution of upper stem rot (USR) and basal stem rot (BSR) infected palms in January 2011 in Betong	90
6.4	Distribution of upper stem rot (USR) and basal stem rot (BSR) infected palms in July 2011 in Betong	91
6.5	Distribution of upper stem rot (USR) and basal stem rot (BSR) infected palms in January 2012 in Betong	92
6.6	Distribution of upper stem rot (USR) and basal stem rot (BSR) infected palms in March 2011 in Miri	93
6.7	Distribution of upper stem rot (USR) and basal stem rot (BSR) infected palms in September 2011 in Miri	94
6.8	Distribution of upper stem rot (USR) and basal stem rot (BSR) infected palms in March 2012 in Miri	95
6.9	Spatial pattern of <i>Ganoderma</i> species based on hotspot analysis in Betong in (A) January 2011, (B) July 2011, (C) January 2012	99
6.10	Spatial pattern of <i>Ganoderma</i> species based on hotspot analysis in Miri in (A) March 2011, (B) September 2011, (C) March 2012	100
7.1	Samples location in Betong. (A) In January 2011, (B) In January 2012	104
7.2	Samples location in Miri. (A) In March 2011, (B) In March 2012	105
7.3	Means (\pm standard error) of soil total N between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	110
7.4	Means (\pm standard error) of foliar (frond 17) total N between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	110
7.5	Means (\pm standard error) of soil exchangeable P between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	113
7.6	Means (\pm standard error) of foliar (frond 17) total P between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	113

7.7	Means (\pm standard error) of soil exchangeable K between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	115
7.8	Means (\pm standard error) of foliar (frond 17) total K between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	116
7.9	Means (\pm standard error) of soil exchangeable Ca between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	118
7.10	Means (\pm standard error) of foliar (frond 17) total Ca between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	118
7.11	Means (\pm standard error) of soil exchangeable Mg between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	120
7.12	Means (\pm standard error) of foliar (frond 17) total Mg between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	121
7.13	Means (\pm standard error) of soil exchangeable Cu between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	123
7.14	Means (\pm standard error) of foliar (frond 17) total Cu between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	123
7.15	Means (\pm standard error) of soil exchangeable Zn between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	125
7.16	Means (\pm standard error) of foliar (frond 17) total Zn between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	126
7.17	Means (\pm standard error) of soil exchangeable B between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	128
7.18	Means (\pm standard error) of foliar (frond 17) total B between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	128
7.19	Means (\pm standard error) of soil exchangeable Mn between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	130

7.20	Means (\pm standard error) of foliar (frond 17) total Mn between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	131
7.21	Means (\pm standard error) of soil exchangeable Fe between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	132
7.22	Means (\pm standard error) of foliar (frond 17) total Fe between uninfected and infected area, and its average in Betong and Miri in 2011 and 2012	133
7.23	Means (\pm standard error) of chemical properties of peat soils between upper stem rot (USR) and basal stem rot (BSR) infected palms in Betong and Miri on 2011 and 2012. (A) Total N, (B) Exchangeable P, (C) Exchangeable K, (D) Exchangeable Ca, (E) Exchangeable Mg, (F) Exchangeable Cu, (G) Exchangeable Zn, (H) Exchangeable B, (I) Exchangeable Mn, (J) Exchangeable Fe	136
7.24	Means (\pm standard error) of chemical properties of foliar (frond 17) between upper stem rot (USR) and basal stem rot (BSR) infected palms in Betong and Miri on 2011 and 2012. (A) Total N, (B) Total P, (C) Total K, (D) Total Ca, (E) Total Mg, (F) Total Cu, (G) Total Zn, (H) Total B, (I) Total Mn, (J) Total Fe	138

LIST OF PLATES

Plate		Page
4.1	White colony with presence of a brown halo around it after three to four days indicates successful isolation of <i>Ganoderma</i> mycelia on <i>Ganoderma</i> selective medium (GSM)	38
4.2	Non-colonized rubber wood block (RWB) of size 6 x 6 x 12 cm	41
4.3	Fully colonized rubber wood block (RWB) of size 6 x 6 x 12 cm with <i>Ganoderma</i> after nine weeks of incubation	41
4.4	Basidiospores of <i>Ganoderma</i> species under compound light microscope (1000X magnification)	42
4.5	DNA fragments yielded at 900 bp (internal control), 656 bp (<i>G. boninense</i>), 415 bp (<i>G. zonatum</i>) and 242 bp (<i>G. miniatocinctum</i>) indicating the identity of <i>Ganoderma</i> isolates from upper stem rot (USR) and basal stem rot (BSR) infected palms in Betong and Miri using multiplex PCR. (A) Isolates G1 to G23, and (B) Isolates G24 to G46	44
4.6	Compatibility test. (A) Compatible reaction between same isolate (control) of G29 (<i>G. zonatum</i>) merged into a single colony, (B) Incompatible reaction between G2 (<i>G. boninense</i>) and G3 (<i>G. zonatum</i>) showing formation of inhibition zone (arrow), (C) Incompatible reaction between G2 (<i>G. boninense</i>) and G9 (<i>G. boninense</i>) showing formation of barrage (arrow), (D) Incompatible reaction between G26 (<i>G. zonatum</i>) and G34 (<i>G. zonatum</i>) showing formation of barrage (arrow)	46
5.1	Plating of necrotic roots and stem bole tissues of the <i>Ganoderma</i> -inoculated oil palm seedling onto <i>Ganoderma</i> selective medium (GSM) yielded white colony with presence of brown halo around it (arrows)	62
5.2	Infection progression of <i>Ganoderma</i> -inoculated oil palm seedlings exhibits external symptoms. (A) Healthy seedling with no sign of infection, (B) Seedling started to show leaf chlorosis or yellowing of the leaf either with or without fungal mass, (C) More leaves became chlorotic and eventually became necrotic (dead leaves), and fungal mass	63

appeared at the base of a seedling's bole (arrow), (D) Severe leaf chlorosis and necrosis, (E) Basidioma of *Ganoderma* (arrow) formed with severe leaf symptoms on a dying seedling, (F) Desiccated dead seedling with well-developed *Ganoderma* basidioma (arrow)

- 5.3 Oil palm seedlings exhibit internal symptoms of the bole (stem) and root (root base). (A) Control (uninoculated) sample with healthy bole with no sign of necrosis (arrow), (B) Control (uninoculated) sample with healthy primary root base with no sign of necrosis, (C) *Ganoderma*-inoculated sample with necrosis of internal bole tissue (arrow), (D) *Ganoderma*-inoculated sample with necrosis of primary root base (arrows) 64
- 6.1 Field symptoms of upper stem rot (USR) and basal stem rot (BSR). (A) Standing USR-infected palm showing *Ganoderma* basidiomata (gb) at upper portion of the palm's stem, (B) Standing BSR-infected palm showing multiple *Ganoderma* basidiomata (gb) at the base or bole of a palm (C) Collapsed USR-infected palm fractured at the upper stem (fc) leaving the crown on the ground (cr), (D) Multiple *Ganoderma* basidiomata (gb) at the fractured point of USR-infected palm, (E) Cross section of USR-infected palm showing decayed stem tissue (dt) occupied almost half the area, and (F) Cross section of USR-infected palm showing healthy stem tissue at the base (ht) of palm 87

LIST OF ABBREVIATIONS

8-MOP	8-methoxypsoralen
AAS	Atomic absorption spectrometer
AFLP	Amplified fragment length polymorphism
ANOVA	Analysis of variance
AUDPC	Area under disease progress curve
bp	Base pair
BSR	Basal stem rot
CIRP	Christmas Island Rock Phosphate
CPO	Crude palm oil
D×P	Dura x Pisifera or; Tenera
DI	Disease incidence
DNA	Deoxyribonucleic acid
DNMRT	Duncan's new multiple range test
DPO	Dual priming oligonucleotide
DSI	Disease severity index
FFB	Fresh fruit bunch
FMP	Fused magnesium phosphate
FSS	Foliar symptoms severity
GIS	Geographical Information System
GLM	General linear model
GPS	Global Positioning System
GSM	Ganoderma selective medium
IPM	Integrated pest management
kPa	Kilopascal
ME	Malt extract

MEA	Malt extract agar
MOP	Muriate of potash
MPOB	Malaysian Palm Oil Board
MVSP	Multivariate statistical package
NK Mix	Nitrogen and potassium mixture
PCR	Polymerase chain reaction
PDA	Potato dextrose agar
PDB	Potato dextrose broth
R^2	Coefficient of determination
RAMS	Random amplified microsatellite
RAPD	Random amplified polymorphic DNA
RCBD	Randomised complete block design
RFLP	Restriction fragment length polymorphisms
RWB	Rubber wood block
SAS	Statistical Analysis System
SOA	Ammonium sulphate
SOPB	Sarawak Oil Palm Berhad
SSI	Spore shape index
UPGMA	Unweighted pair group method of arithmetic averages
UPMKB	Universiti Putra Malaysia Bintulu Campus
USR	Upper stem rot
WAI	Week after inoculation



CHAPTER 1

INTRODUCTION

1.1 Background

Oil palm (*Elaeis guineensis*) is the world's most important oil producing crop as it supplies 65.29 million tonnes, which is approximately 39% of the world vegetable oils and fats, and it is also the most efficient oil producing crop as compared with other major oil producing crops (Kurki *et al.*, 2006; USDA, 2014). Currently, Malaysia is one of the world's largest producers of palm oil contributing 30% towards world palm oil supplies (USDA, 2014). In Malaysia, the oil palm plantations reached 5.23 million hectares with an output of 19.32 million tonnes of crude palm oil (CPO), where the states of Sabah and Sarawak are the largest producers with recent trend showing rapid expansion of plantation on peat in Sarawak (MPOB, 2013). Hence, the utilization of peat lands in Sarawak for oil palm cultivation has become increasingly important (ASOS, 2011; Lim *et al.*, 2012).

The major threats to the sustainability of oil palm are pests and diseases. Currently, it is known that the most prominent disease in oil palm plantations across the Southeast Asia region is basal stem rot (BSR) which is caused by a pathogenic basidiomycete fungus, *Ganoderma boninense* (Flood *et al.*, 2000; Corley and Tinker, 2003; Rees *et al.*, 2012; Wong *et al.*, 2012). The disease reduced the yield and shortening the economic life of an oil palm. Moreover, *Ganoderma* species are also found to be closely associated with upper stem rot (USR) in oil palm (Hasan *et al.*, 2005; Pilotti, 2005; Rees *et al.*, 2012). The infection of *Ganoderma* in USR is on the upper portion of the stem or trunk of an oil palm, instead of at the base as in BSR.

Upper stem rot of oil palm in Malaysia has been noticed since 1937 and it was reported to usually occur on deep peat and inland valley soils. Upper stem rot is not considered as a major disease of oil palm at the earlier time (Thompson, 1937) but recently, USR has begun to gain more attention when a few cases of the disease were observed to severely infect the oil palm plantations with the presence of *Ganoderma* species in Sabah, Malaysia (Abdullah *et al.*, 1999), Papua New Guinea (Pilotti, 2005) and Indonesia (Rees *et al.*, 2012).

Over the years, researchers and oil palm planters have put much effort into the management strategies of the diseases caused by *Ganoderma* species through various cultural, chemical and biological aspects of the pest and host. Some of the disease control strategies are through the development of bio-control agents, disease resistant planting materials, methods of pesticides application, cover crops, fertilizers input, early detection of pathogens, and various aspects of estate sanitation (Chung, 2011; Cooper *et al.*, 2011; Hushiarian *et al.*, 2013; Naher *et al.*, 2013). However, until today days, *Ganoderma* species remains the major threats to the sustainability of oil palm.

1.2 Justification

Implementation of effective control strategies on the stem rot diseases and its causative agents in oil palm require for better understandings on the basic information of the diseases and pathogens. Such information include the etiological aspects in terms of symptomology and occurrence of the diseases, and species distribution and aggressiveness of the pathogens, and epidemiological aspects in terms of mode of disease spread and the environmental factors associated with the diseases.

Although it is commonly known that BSR is caused by *Ganoderma boninense*, little is known of the threats by other species of *Ganoderma* in oil palm plantations such as *G. zonatum* and *G. miniatocinctum* that are also implicated in BSR (Idris *et al.*, 2000a; Wong *et al.*, 2012). Moreover, both USR and BSR are associated with similar pathogens, but little is known about the distinctive characteristics between them. In addition, there is little information known about USR due to the lack of comprehensive studies on the disease in terms of epidemiological and etiological aspects. In some cases, USR is misinterpreted as BSR due to lack of information on the disease symptoms. The role of USR and other species of *Ganoderma* (i.e. *G. zonatum* and *G. miniatocinctum*) that may devastate the oil palm industry should not be overlooked by the researchers as well as the planters because in the long run this disease and pathogen may destroy the estates and affect oil palm industries seriously.

The infection mode of *Ganoderma* species in BSR were reported to initiate at the root of an oil palm, and the infection could spread to other healthy neighbouring palms through root-to-root contact (Flood *et al.*, 2005; Khairudin and Chong, 2008). However, the root-to-root infection mode is considered uncommon since *Ganoderma* species in oil palm plantations are genetically heterogeneous which suggests the spread of *Ganoderma* species through basidiospores (Miller *et al.*, 1999; Pilotti *et al.*, 2003; Sanderson, 2005). In addition, USR spread from ground level such as root-to-root contact and infection from inoculum source in the ground was obviously not possible. Hence, the infection modes of *Ganoderma* species are discussed in this study because it is crucial information for effective disease control strategies.

There were several reports on the aggressiveness of *Ganoderma boninense* of basal stem rot (BSR) in oil palms (Rees *et al.*, 2007; Sariah *et al.*, 2007; Chan *et al.*, 2011; Kok *et al.*, 2013). However, until recently, there was no serious attempt on comparative study of *G. boninense*, *G. zonatum* and *G. miniatocinctum*. Moreover, aggressiveness of *Ganoderma* isolates from USR-infected oil palms remains unknown. Difference in aggressiveness among *Ganoderma* species from different sources (USR and BSR) may play a vital role in the intensity of the diseases.

Furthermore, little was known about the status of the diseases and pathogens in Sarawak or Borneo since most of the previous reports were based on studies in Peninsular Malaysia and other countries (Turner, 1981; Ariffin *et al.*, 1989; Idris *et al.*, 2000a; Hasan *et al.*, 2005; Pilotti, 2005; Khairudin and Chong, 2008; Rees *et al.*, 2012). The state of Sarawak, which has the major distribution of peat lands in Malaysia has become an important area for oil palm cultivation (ASOS, 2011; Lim *et al.*, 2012). One of the major drawbacks of oil

palm cultivation on peat is nutrient unavailability due to nutrients fixation onto the organic matter (Fageria *et al.*, 2002; Hasnol *et al.*, 2011; Lim *et al.*, 2012) that may lead to unhealthy oil palm which later leads to increased susceptibility of disease infection (Marschner, 1995). Hence, one of the important environmental factors that could be related to the diseases epidemiology is the nutrients availability.

1.3 Objectives

The objectives of this research are to:

- i. Investigate the genetic and morphological diversity of *Ganoderma* species associated with USR and BSR in oil palm.
- ii. Evaluate the aggressiveness of *Ganoderma* species isolated from USR- and BSR-infected oil palms.
- iii. Investigate disease symptoms, occurrence and spatio-temporal distribution of USR and BSR, and hotspot analysis of *Ganoderma* species of the diseases in oil palm.
- iv. Investigate the relationship between oil palm nutrients status and infection by *Ganoderma* species in plantation on peat.

REFERENCES

- Abdullah, F., S.B. Liew and N. Malik. 1999. Upper stem rot of oil palms (*E. guineensis*) in Langkon, Sabah. In: *Sustainable Crop Protection Practices in the Next Millennium*. eds. Sidek, Z., S.L. Bong, S.K. Vijaya, C.A. Ong and A.K. Husan, pp. 101-103. Malaysia: Malaysian Plant Protection Society.
- Abdullah, F., 2000. Spatial and sequential mapping of the incidence of basal stem rot of oil palm (*Elaeis guineensis*) on a former coconut (*Cocos nucifera*) plantation. In: *Ganoderma Diseases of Perennial Crops*. eds. Flood, J., P.D. Bridge and M. Holderness, pp. 183-194. United Kingdom, Wallingford: CABI Publishing.
- Adaskaveg, J.E. and R.L. Gilbertson. 1989. Cultural studies of four North American species in the *Ganoderma lucidum* complex with comparisons to *G. lucidum* and *G. tsugae*. *Mycological Research* **92**: 182-191.
- Agnieszka, G.G. 2010. The occurrence of *Ganoderma* spores in the air and its relationship with meteorological factors. *ACTA Agrobotanica* **63**: 93-97.
- Agriculture Statistics of Sarawak (ASOS). 2011. Oil Palm. Sarawak, Malaysia. Department of Agriculture, Sarawak, Malaysia. http://www.doa.sarawak.gov.my/modules/web/download_show.php?id=1110. Retrieved 24 March 2014.
- Agrios, G.N. 2005. *Plant Pathology (5th ed.)*. USA: Elsevier Academic Press, 952 p.
- Ahmad, R., M. Arshad, A. Khalid and Zahir. Z.A. 2008. Effectiveness of organic-/bio-fertilizer supplemented with chemical fertilizers for improving soil water retention, aggregate stability, growth and nutrient uptake of maize (*Zea mays* L.). *Journal of Sustainable Agriculture* **31**: 57-77.
- Aholoukpe, H., B. Dubos, A. Flori, P. Deleporte, G. Amadji, J.L. Chotte and D. Blavet. 2013. Estimating aboveground biomass of oil palm: Allometric equations for estimating frond biomass. *Forest Ecology and Management* **292**: 122-129.
- Andriesse, J.P. 1988. *Nature and Management of Tropical Peats*, *FAO Soils Bulletin* 59. Rome: FAO, 179 p.
- Anshari, G.Z., M. Afifudin, M. Nuriman, E. Gusmayanti, L. Arianie, R. Susana, R.W. Nusantara, J. Sugardjito and A. Rafiastanto. 2010. Drainage and land use impacts on changes in selected peat properties and peat degradation in West Kalimantan Province, Indonesia. *Biogeosciences* **7**: 3403-3419.

- Ariffin, D., A.S. Idris and H.A. Halim. 1989. Significance of the black line within oil palm tissue decay by *Ganoderma boninense*. *Elaeis* **1**: 11-16.
- Ariffin, D. and A.S. Idris. 1992. The *Ganoderma* selective medium (GSM). *Palm Oil Research Institute of Malaysia (PORIM) Information Series* **8**: 2pp.
- Ariffin, D., A.S. Idris and A. Marzuki. 1996. Spread of *Ganoderma boninense* and vegetative compatibility studies of a single field palm isolates. In: eds. Ariffin D., M.B. Wahid, N. Rahanaidu, T. Dolmat, K. Paranjothy, S.C. Cheah, C.K. Chong and B.H. McCown, *Proceedings of the 1996 PORIM International Palm Oil Congress (Agriculture)*, 23-28 September 1996. Bangi, Selangor, Malaysia, pp. 317-329.
- Ariffin, D., A.S. Idris and G. Singh. 2000. Status of *Ganoderma* in oil palm. In: *Ganoderma Diseases of Perennial Crops*. eds. Flood, J., P.D. Bridge and M. Holderness, pp. 49-68. United Kingdom, Wallingford: CABI Publishing.
- Ariffin, D. and A.S. Idris. 2002. Progress and research on *Ganoderma* basal stem rot of oil palm. In: *Seminar on Recent Progress in the Management of Peat and Ganoderma, May 2002*. Bangi, Selangor, Malaysia. 50pp.
- Armstrong, D.L. 1999. *Better Crops with Plant Food Volume 83 No.1*. Canada: Potash and Phosphate Institute of Canada (PPIC), 40 p.
- Azahar, T.M., J.C. Mustapha, S. Mazliham and B. Patrice. 2011. Temporal analysis of basal stem rot disease in oil palm plantations: An analysis on peat soil. *International Journal of Engineering and Technology* **11**: 96-101.
- Balasundram, S.K., M.H.A. Husni and O.H. Ahmed. 2008. Application of geostatistical tools to quantify spatial variability of selected soil chemical properties from a cultivated tropical peat. *Journal of Agronomy* **7**: 82-87.
- Basiron, Y. 2007. Palm oil production through sustainable plantations. *European Journal of Lipid Science and Technology* **109**: 289-295.
- Benjamin, M. and K.H. Chee. 1995. Basal stem rot of oil palm – a serious problem on inland soils. *MAPPs Newsletter* **19**: 3
- Bivi, M.R., M.S.N. Farhana, A. Khairulmazmi and A.S. Idris. 2010. Control of *Ganoderma boninense*: a casual agent of basal stem rot disease in oil palm with endophyte bacteria *In Vitro*. *International Journal of Agriculture and Biology* **12**: 833-839.
- Blevins, D.G. and K.M. Lukaszewski. 1998. Boron in plant structure and function. *Annual Review of Plant Physiology* **49**: 481–500.

- Bota, C. and C. Deliu. 2011. The effect of copper sulphate on the production of flavonoids in *Digitalis lanata* cell cultures. *Farmacia* **59**: 113-118.
- Box, G.E.P., J.S. Hunter and W.G. Hunter. 2005. *Statistics for Experimenters: Design, Innovation, and Discovery (2ed.)*. University of Michigan, USA: Wiley-Interscience, 664 p.
- Boyhan, G.E., R.L. Torrance and C.D. Hill. 2007. Effects of nitrogen, phosphorus, and potassium rates and fertilizer sources on yield and leaf nutrient status of short-day onions. *Hortscience* **42**: 653-660.
- Bragazza, L. and C. Freeman. 2007. High nitrogen availability reduces polyphenol content in Sphagnum peat. *Science of the Total Environment* **377**: 439-443.
- Bray, R.H. and L.T. Kurtz. 1945. Determination of total organic and available forms of phosphorus in soils. *Soil Science* **11**: 394-402.
- Breton, F., Y. Hasan, Hariadi, Z. Lubis and H. de Franqueville. 2006. Characterization of parameters for the development of an early screening test for basal stem rot tolerance in oil palm progenies. *Journal of Oil Palm Research Special Issue*: 24-36.
- Broeshart, H. 1956. The application of foliar analysis to oil palm cultivation. II. Uptake and distribution of nutrients in mineral deficient oil palm. *Tropical Agriculture* **33**: 238-243.
- Broschat, T.K. 1997. Nutrient distribution, dynamics, and sampling in coconut and canary island date palms. *Journal of the American Society for Horticultural Science* **122**: 884-890.
- Brown, P.H., N. Bellaloui, M.A. Wimmer, E.S. Bassil, J. Ruiz, H. Hu, H. Pfeffer, F. Dannel and V. Romheld. 2002. Boron in plant biology. *Plant Biology* **4**: 205-223.
- Butler, R.A. 2011. First large-scale map of oil palm plantations reveals big environmental toll. Mongabay.com. http://news.mongabay.com/2011/0307-koh_palm_oil.html. Retrieved 24 March 2014.
- Cakmak, I. 2000. Possible roles of zinc in protecting plant cells from damage by reactive oxygen species. *New Phytologist* **146**: 185-205.
- Campanile, G., S.L. Giove and N. Luisi. 2004. Genetic and morphologic variability of *Phellinus torulosus* isolates in some oak woods of Southern Italy. *Journal of Plant Pathology* **86**: 105-115.
- Campbell, C.R. 1992. Determination of total nitrogen in plant tissue by combustion. In: *Plant Analysis References Procedures for the Southern Region of the United States*. eds. Plank, C.O., pp. 20-22. USA: Southern Cooperative Series Bulletin.

- Campbell, N.A. and J.B. Reece. 2005. *Biology* (7th ed.). USA: Pearson Benjamin Cummings, 1312 p.
- Carbajo, J.M., H. Junca, M.C. Terrón, T. González, S. Yagüe, E. Zapico and A.E. González. 2002. Tannic acid induces transcription of laccase gene *cglcc1* in the white-rot fungus *Coriopsis gallica*. *Canadian Journal of Microbiology* **48**: 1041-1047.
- Caruso, C., G. Chilosi, L. Leonardi, L. Bertin, P. Magro, V. Buonocore and C. Caporule. 2001. A basic peroxidase from wheat kernel with antifungal activity. *Phytochemistry* **58**: 43-50.
- Chan, J.J., Z. Latiffah, K.W. Liew and A.S. Idris. 2011. Pathogenicity of monokaryotic and dikaryotic mycelia of *Ganoderma boninense* on oil palm seedlings and germinated seeds in Malaysia. *Australasian Plant Pathology* **40**: 222-227.
- Chapman, G.W. and H.M. Gray. 1949. Leaf analysis and the nutrition of the oil palm. *Annual Botany* **13**: 415-433.
- Chen, E.L., Y.A. Chen, L.M. Chen and Z.H. Liu. 2002. Effect of copper on peroxidase activity and lignin content in *Raphanus sativus*. *Plant Physiology and Biochemistry* **40**: 439-444.
- Chen, H.S., B.B. Jin and T.T. Chang. 2013. Spatial variance characteristics of soil available microelements in tobacco planting fields on slant plain in the east side of Funiushan mountain in central China. *Journal of Food Agriculture and Environment* **11**: 1464-1469.
- Chmielowska, J., J. Veloso, J. Gutierrez, C. Silvar and J. Diaz. 2009. Cross-protection of pepper plants stressed by copper against a vascular pathogen is accompanied by the induction of a defence response. *Plant Science* **178**: 176-182.
- Chong, K.P., M.S. Lum, C.P. Foong, C.M.V.L. Wong, M. Atong and S. Rossall. 2011. First identification of *Ganoderma boninense* isolated from Sabah based on PCR and sequence homology. *African Journal of Biotechnology* **10**: 14718-14723.
- Chong, K.P., A. Markus and S. Rossall. 2012a. The susceptibility of different varieties of oil palm seedlings to *Ganoderma boninense* infection. *Pakistan Journal of Botany* **44**: 2001-2004.
- Chong, K.P., A. Markus and S. Rossall. 2012b. The role of syringic acid in the interaction between oil palm and *Ganoderma boninense*, the causal agent of Basal Stem Rot. *Plant Pathology* **61**: 953-963.
- Chun, J.Y., K.J. Kim, I.T. Hwang, Y.J. Kim, D.H. Lee, I.K. Lee and J.K. Kim. 2007. Dual priming oligonucleotide system for the multiplex detection of respiratory viruses and SNP genotyping of CYP2C19 gene. *Nucleic Acids Research* **35**: e40.

- Chung, G.F. 2011. Management of *Ganoderma* disease in oil palm plantations. *The Planter* **87**: 325-339.
- Cooper, R.M., J. Flood and R.W. Rees. 2011. *Ganoderma boninense* in oil palm plantations: current thinking on epidemiology, resistance and pathology. *The Planter* **87**: 515-526.
- Corley, R.H.V. 1976a. The genus *Elaeis*. In: *Oil Palm Research*, eds. Corley, R.H.V., J.J. Hardon and B.J. Wood, pp. 3-6. Amsterdam: Elsevier Scientific.
- Corley, R.H.V. 1976b. Physiological aspects of nutrition. In: *Oil Palm Research*, eds. Corley, R.H.V., J.J. Hardon and B.J. Wood, pp. 157-164. Amsterdam: Elsevier Scientific.
- Corley, R.H.V. and B.S. Gray. 1976. Growth and morphology. In: *Oil Palm Research*, eds. Corley, R.H.V., J.J. Hardon and B.J. Wood, pp. 7-22. Amsterdam: Elsevier Scientific.
- Corley, R.H.V. and P.B. Tinker. 2003. *The Oil Palm (4th ed.)*. Oxford: Blackwell, 592 p.
- da Fonseca, A.F., A.J. Melfi and C.R. Montes. 2005. Maize growth and changes in soil fertility after irrigation with treated sewage effluent. I. Plant dry matter yield and soil nitrogen and phosphorus availability. *Communications in Soil Science and Plant Analysis* **36**: 1965-1981.
- Department of Irrigation and Drainage Sarawak. 2014. Resource Centre - Peat Swamp Development. http://www.did.sarawak.gov.my/modules/web/page.php?lang=en&id=97&menu_id=0&sub_id=117. Retrieved 24 March 2014.
- Diaz, J., A. Barnal, F. Pomar and F. Merino. 2001. Induction of shikimate dehydrogenase and peroxidase in pepper (*Capsicum annuum* L.) seedlings in response to copper stress and its relation to lignifications. *Plant Science* **161**: 179-188.
- Dobinson, K.F., G.K. Tenuta and G. Lasarovits. 1996. Occurrence of race 2 of *Verticillium dahliae* in processing tomato fields in south-western Ontario. *Canadian Journal of Plant Pathology* **18**: 55-58.
- Dordas, C. 2008. Role of nutrients in controlling plant diseases in sustainable agriculture: A review. *Agronomy for Sustainable Development* **28**: 33-46.
- Elfath, M.E., M.A. Ahmed, J.C. Robert and E.K. Paul. 2000. Multiplex PCR: Optimization and application in diagnostic virology. *Clinical Microbiology Reviews* **13**: 559-570.
- Elliott, M.L. and T.K. Broschat. 2000. *Ganoderma Butt Rot of Palms*. USA: Institute of Food and Agriculture Sciences, University of Florida, 7 p.

- Elliott, M.L. and T.K. Broschat. 2001. Observations and pathogenicity experiments on *Ganoderma zonatum* in Florida. *Palms* **45**: 62-72.
- Fageria, N.K., V.C. Baligar and R.B. Clark. 2002. Micronutrients in crop production. *Advanced in Agronomy* **77**: 185-268.
- Flood, J., Y. Hasan, P.D. Turner and E.B. O'Grady. 2000. The spread of *Ganoderma* from infective sources in the field and its implications for management of the disease in oil palm. In: *Ganoderma Diseases of Perennial Crops*. eds. Flood, J., P.D. Bridge and M. Holderness, pp. 101-112. United Kingdom, Wallingford: CABI Publishing.
- Flood, J., L. Keenan, S. Wayne and Y. Hasan. 2005. Studies on oil palm trunks as sources of infection in the field. *Mycopathologia* **159**: 101-107.
- Foster, H.L. 2003. Assessment of oil palm fertilizer requirements. In: *Oil Palm: Management for Large and Sustainable Yields*, eds. Fairhurst, T. and R. Hardter, pp. 231-257. Canada: Potash and Phosphate Institute of Canada (PPIC).
- Gamble, G.R., D.E. Akin, H.P. Makkar and K. Becker. 1996. Biological degradation of tannins in sericea lespedeza (*Lespedeza cuneata*) by the white rot fungi *Ceriporiopsis subvermispota* and *Cyathus stercoreus* analyzed by solid-state ^{13}C nuclear magnetic resonance spectroscopy. *Applied and Environmental Microbiology* **62**: 3600-3604.
- Gasch, A.P. 2007. Comparative genomics of the environmental stress response in ascomycete fungi. *Yeast* **24**: 961-976.
- Germer, J. and J. Sauerborn. 2008. Estimation of the impact of oil palm plantation establishment on greenhouse gas balance. *Environment, Development and Sustainability* **10**: 697-716.
- Ghosh, M. 2006. Antifungal properties of haem peroxidase from *Acorus calamus*. *Annals of Botany* **98**: 1145-1153.
- Goh, K.J., P.S. Chew and K.K. Kee. 1996. Spatial soil fertility in mature oil palm agroecosystem and its implications on fertiliser management. In: eds. Aminuddin, B.H., A.B. Ismail, A.R. Ahmad and M.Z. Ghazali, *Proceeding of the Soil Science Conference of Malaysia*, 1995. Langkawi, Malaysian Society of Soil Science, Kuala Lumpur, pp. 80-90.
- Goh, K.J., H.H. Gan, K.K. Kee, P.S. Chew and K.C. Teoh. 2007. Boron requirement and distribution in the oil palm (*Elaeis guineensis* Jacq.) and some implications on the manuring practices. In: *Advances in Plant and Animal Boron Nutrition*. eds. Xu, F., H.E. Goldbach, P.H. Brown, R.W. Bell, T. Fujiwara, C.D. Hunt, S. Goldberg and L. Shi, pp. 189-202. Netherlands: Springer.

- Goh, K.J. 2003. General oil palm nutrition. In: *Oil Palm: Management for Large and Sustainable Yields*, eds. Fairhurst, T. and R. Hardter, pp. 191-230. Canada: Potash and Phosphate Institute of Canada (PPIC).
- Gomez, K.A. and A.A. Gomez. 1984. *Statistical Procedures for Agricultural Research* (2nd ed.). USA: Wiley-Interscience, 704 p.
- Graham, D.R. 1983. Effects of nutrients stress on susceptibility of plants to disease with particular reference to the trace elements. *Advances in Botanical Research* **10**: 221–276.
- Graham, D.R. and M.J. Webb. 1991. Micronutrients and disease resistance and tolerance in plants. In: *Micronutrients in Agriculture, 2nd ed.*, eds. Mortvedt, J.J., F.R. Cox, L.M. Shuman and R.M. Welch, pp. 329-370. Madison, Wisconsin, USA: Soil Science Society of America Inc.
- Grewal, H.S., R.D. Graham and Z. Rengel. 1996. Genotypic variation in zinc efficiency and resistance to crown rot disease (*Fusarium graminearum* Schw. Group 1) in wheat. *Plant Soil* **186**: 219-226.
- Gupta, P.K. 2009. *Soil, Plant, water and Fertilizer Analysis* (2nd ed.). Jodhpur, India: Agrobios, 350 p.
- Hanlon, E.A. 2000. Determination of potassium, calcium, and magnesium in plant by atomic absorption techniques. In: *Plant Analysis References Procedures for the Southern Region of the United States*. eds. Plank, C.O., pp. 30-33. USA: Southern Cooperative Series Bulletin.
- Hasan, Y., H.L. Foster and J. Flood. 2005. Investigations on the causes of upper stem rot on standing mature oil palms. *Mycopathologia* **159**: 109-112.
- Hasnol, O., M.D. Farawahida and A.M. Tarmizi. 2009. Technique for mechanically forced unidirectional leaning of oil palm on peat. *Malaysian Palm Oil Board Information Series* 458, MPOB TT No. 417: 4pp.
- Hasnol, O., A.M. Tarmizi, M.D. Farawahida, M.H. Haniff and M.Z. Pilus. 2011. Best management practice for oil palm cultivation on peat: Ground water-table maintenance in relation to peat subsidence and estimation of CO₂ emission at Sessang, *Sarawak. Journal of Oil Palm Research* **23**: 1078-1086.
- Heckman, J.R., B.B. Clarke and J.A. Murphy. 2003. Optimizing manganese fertilization for the suppression of take-all patch disease on creeping bentgrass. *Crop Science* **43**: 1395–1398.
- Henegariu, O., N.A. Heerema, S.R. Dlouhy, G.H. Vance and P.H. Vogt. 1997. Multiplex PCR: critical parameters and step-by-step protocol. *BioTechniques* **23**: 504–511.

- Ho, Y.W. and A. Nawawi. 1985. *Ganoderma boninense* Pat. from basal stem rot of oil palm (*Elaeis guineensis*) in Peninsular Malaysia. *Pertanika* **8**: 425-428.
- Hoe, P.K., C.F.J. Bong, K. Jugah and A. Rajan. 2009. Evaluation of *Metarhizium anisopliae* var. *anisopliae* (Deuteromycotina: Hyphomycete) Isolates and their effects on subterranean termite *Coptotermes curvignathus* (Isoptera: Rhinotermitidae). *American Journal of Agricultural and Biological Sciences* **4**: 289-297.
- Huber, D.M. and R.D. Graham. 1999. The role of nutrition in crop resistance and tolerance to disease. In: *Mineral Nutrition of Crops Fundamental Mechanisms and Implications*, eds. Rengel, Z., pp. 205-226. New York: Food Product Press.
- Huber, D.M. and J.B. Jones. 2013. The role of magnesium in plant disease. *Plant Soil* **368**: 73-85.
- Hue, N.V., R. Uchida and M.C. Ho. 2000. Sampling and analysis of soil and plant tissues. In: *Plant Nutrient Management in Hawaii's Soils, Approaches for Tropical and Subtropical Agriculture*. eds. Silva, J.A. and R.S. Uchida, pp. 23-30. University of Hawaii, Manoa: College of Tropical Agriculture and Human Resources.
- Hushiarian, R., N.A. Yusof and S.W. Dutse. 2013. Detection and control of *Ganoderma boninense*: strategies and perspectives. *SpringerPlus* **2**: 555.
- Ibrahim, M.H., H.Z.E. Jaafar, A. Rahmat and Z.A. Rahman. 2011. The relationship between phenolics and flavonoids production with total non structural carbohydrate and photosynthetic rate in *Labisia pumila* Benth. under high CO₂ and nitrogen fertilization. *Molecules* **16**: 162-174.
- Idris, A.S. 1999. Basal stem rot (BSR) of oil palm (*Elaeis guineensis* Jacq.) in Malaysia: Factors associated with variation in disease severity. Ph.D Thesis. Wye College, University of London, UK.
- Idris, A.S., D. Ariffin, T.R. Swinburne and T.A. Watt. 2001. Distribution of species of *Ganoderma* basal stem rot of oil palm in relation to the environmental conditions in Peninsular Malaysia. In: eds. Ariffin, D. and C.K. Weng, *Proceedings of the 2001 PORIM International Palm Oil Congress (Agriculture)*. Bangi, Selangor, Malaysia, pp. 385-394.
- Idris, A.S, D. Ariffin, T.R. Swinburne and T.A. Watt. 2000a. The identity of *Ganoderma* species responsible for BSR disease of oil palm in Malaysia - Pathogenicity test. *Malaysian Palm Oil Board Information Series* 103, MPOB TT. 77b: 4pp.

- Idris, A.S., D. Ariffin, T.R. Swinburne and T.A. Watt. 2000b. The identity of *Ganoderma* species responsible for basal stem rot (BSR) disease of oil palm in Malaysia – Morphological characteristics. *Malaysian Palm Oil Board Information Series* 102, MPOB TT No. 77a: 4pp.
- Idris, A.S. and D. Ariffin. 2004. Basal stem rot – Biology, detection and control. In: eds. Wahid, M.B., N. Kamaruddin, C.K. Weng, A. Darus, S.R.A. Ali and S. Sundram, *Proceeding of the International Conference on Pest and Diseases of Importance to the Oil Palm Industry*, 18-19 May 2014. Malaysian Palm Oil Board, Kuala Lumpur, pp. 134-162.
- Idris, A.S., A. Kushairi, S. Ismail and D. Ariffin. 2004. Selection for partial resistance in oil palm progenies to *Ganoderma* basal stem rot. *Journal of Oil Palm Research* **16**: 12-18.
- Idris, A.S., S. Kushairi, D. Ariffin and M.W. Basri. 2006. Technique for inoculation of oil palm germinated seeds with *Ganoderma*. *Malaysian Palm Oil Board Information Series* 321, MPOB TT No. 314: 4pp.
- Idris, A.S., S. Rajinder, A.Z. Madiah and M.B. Wahid. 2010. Multiplex PCR-DNA kit for early detection and identification of *Ganoderma* species in oil palm. *Malaysian Palm Oil Board Information Series* 531, MPOB TS No.73: 4pp.
- Isaac, R.A. and W.C. Johnson. 1992. Determination of sulfur in plant tissue by combustion. In: *Plant Analysis References Procedures for the Southern Region of the United States*. eds. Plank, C.O., pp. 42-44. USA: Southern Cooperative Series Bulletin.
- Izzati, M.Z.N.A. and F. Abdullah. 2008. Disease suppression in *Ganoderma*-infected oil palm seedlings treated with *Trichoderma harzianum*. *Plant Protection Science* **44**: 101-107.
- Jaya, J. 2002. *Sarawak: Peat Agricultural Use*. Malaysia: Malaysian Agricultural Research and Development Institute (MARDI), 12 p.
- Jia, S., D. Zhou and D. Xu. 2011. The temporal and spatial variability of soil properties in an agricultural system as affected by farming practices in the past 25 years. *Journal of Food Agriculture and Environment* **9**: 669-676.
- Journel, A.G. and C.H. Huijbregts. 2004. *Mining Geostatistics*. New Jersey, USA: The Blackburn Press, 600 p.
- Karakousis, A., L. Tan, D. Ellis, H. Alexiou and P.J. Wormald. 2006. An assessment of the efficiency of fungal DNA extraction methods for maximizing the detection of medically important fungi using PCR. *Journal of Microbiological Methods* **65**: 38-48.

- Karthikeyan, M., V. Jayakumar, K. Radhika, R. Bhaskaran, R. Velazhahan and D. Alice. 2005. Induction of resistance in host against the infection of leaf blight pathogen (*Alternaria palandui*) in onion (*Allium cepa* var *aggregatum*). *Indian Journal of Biochemistry and Biophysics* **42**: 371-377.
- Kechavarzi, C., Q. Dawson and P.B. Leeds-Harrison. 2010. Physical properties of low-lying agricultural peat soils in England. *Geoderma* **154**: 196-202.
- Khairudin, H., T.K. Ling and A.R.A. Razak. 1991. Pathogenicity of *Ganoderma boninense* Pat. on oil palm seedlings. In: eds. Yusof *et al.*, *Proceedings of the 1993 PORIM International Palm Oil Congress - Agriculture Conference*, 9-14 September 1991. Palm Oil Research Institute of Malaysia, Bangi. Selangor, Malaysia, pp. 418-423.
- Khairudin, H. 1993. Basal stem rot of oil palm caused by *Ganoderma boninense*: An update. In: eds. Jalani *et al.*, *Proceedings of the 1993 PORIM International Palm Oil Congress - Agriculture Conference*, 20-25 September 1993. Palm Oil Research Institute of Malaysia, Bangi. Selangor, Malaysia, pp. 739-749.
- Khairudin, H. and T.C. Chong. 2008. An overview of the current status of *Ganoderma* basal stem rot and its management in a large plantation group in Malaysia. *The Planter* **84**: 469-482.
- Kim, H.K., G.S. Seo and H.G. Kim. 2001. Comparison of characteristics of *Ganoderma lucidum* according to geographical origins: Consideration of morphological characteristics. *Mycobiology* **29**: 80-84.
- Kloepper, J.W., J. Leong, M. Teintze and M.N. Schroth. 1980. *Pseudomonas* siderophores: A mechanism explaining disease-suppressive soils. *Current Microbiology* **4**: 317-320.
- Kok, S.M., Y.K. Goh, H.J. Tung, K.J. Goh, W.C. Wong and Y.K. Goh. 2013. In vitro growth of *Ganoderma boninense* isolates on novel palm extract medium and virulence on oil palm (*Elaeis guineensis*) seedlings. *Malaysian Journal of Microbiology* **9**: 33-42.
- Korn, M.D.G.A., E.S.D.B. Morte, D.C.M.B.D. Santos, J.T. Castro, J.T.P. Barbosa, A.P. Teixeira A.P. Fernandes, B. Welz, W.P.C.D. Santos, E.B.G.N.D. Santos and M. Korn. 2008. Sample preparation for the determination of metals in food samples using spectroanalytical method - A review. *Applied Spectroscopy Reviews* **43**: 67-92.
- Kurki, A., A. Hill and M. Morris. 2006. *Biodiesel: The sustainability dimensions*. National Center of Appropriate Technology, USA: National Sustainable Agriculture Information Service (ATTRA), 12 p.
- Latiffah, Z. and Y.W. Ho. 2005. Morphological characteristics and somatic incompatibility of *Ganoderma* from infected oil palm from three inland estates. *Malaysian Journal of Microbiology* **1**: 46-52.

- Latiffah, Z., K. Harikrishna, S.G. Tan, A. Faridah and Y.W. Ho. 2005. Random amplified polymorphic DNA (RAPD) and random amplified microsatellite (RAMS) of *Ganoderma* from infected oil palm and coconut stumps in Malaysia. *Asia Pacific Journal of Molecular Biology and Biotechnology* **13**: 23-34.
- Lattanzio, V., V.M.T. Lattanzio and A. Cardinali. 2006. Role of phenolics in the resistance mechanisms of plants against fungal pathogens and insects. In: *Phytochemistry: Advances in Research*, eds. Imperato, F., pp. 23-67. Kerala, India: Research Signpost.
- Law, M.C., S.K. Balasundram, M.H.A. Husni, O.H. Ahmed and M.H. Hanif. 2009. Spatial variability of soil organic carbon in oil palm. *International Journal of Soil Science* **4**: 93-103.
- Lewis, N. G., L. B. Davin and S. Sarkanen. 1999. The nature and function of lignins. In: *Comprehensive Natural Products chemistry*, eds. Barton, D.H.R., K. Nakanishi and O. Meth-Cohn, pp. 617-745. Oxford, United Kingdom: Elsevier.
- Lim, K.H. 2005. Soil water and fertilizer management for oil palm cultivation on peat soils. In: *Proceeding of the 2005 International Oil Palm Congress –Agriculture, Biotechnology and Sustainability Conference*, 25-29 September 2005. Malaysian Palm Oil Board, Selangor, Malaysia, pp. 433-455.
- Lim, K.H., S.S. Lim, F. Parish and R. Suharto. 2012. *RSPO Manual on Best Management Practice (BMPs) for Existing Oil Palm Cultivation on Peat*. Kuala Lumpur: RSPO Secretariat Sdn Bhd., 215 p.
- Macrae, M.L., K.J. Devito, M. Strack and J.M. Waddington. 2013. Effect of water table drawdown on peatland nutrient dynamics: implications for climate change. *Biogeochemistry* **112**: 661-676.
- Makinde, E.A., L.S. Ayeni and S.O. Ojeniyi. 2011. Effects of organic, organomineral and NPK fertilizer treatments on the nutrient uptake of *Amaranthus cruentus* L. on two soil types in Lagos, Nigeria. *Journal of Central European Agriculture* **12**: 114-123.
- Maksymiec, W. 1997. Effect of copper on cellular process in higher plants. *Photosynthetica* **34**: 321-342.
- Malaysian Palm Oil Board (MPOB). 2013. Oil palm planted area 2013. Economic and Industry Development Division, MPOB, Selangor, Malaysia, 1pp.
- Malaysian Palm Oil Council (MPOC). 2012. *The Oil Palm Tree*. http://www.mpoc.org.my/The_Oil_Palm_Tree.aspx. Retrieved 17th March 2014
- Marschner, H. 1995. *Mineral Nutrition of Higher Plants (2nd ed.)*. London: Academic Press Ltd., 889 p.

- May, C.Y., M.A. Ngan, C.K. Yoo, R.A. Majid, A.Y.K. Chung, H.L.L. Nang, C.S. Foon, Y.C. Liang, P.C. Wei, N.M. Han and Y. Basiron. 2005. Palm diesel: Green and renewable fuel from palm oil. *Palm Oil Developments* **42**: 1-5.
- McGrath, S.P., J.R. Sanders and M.H. Shalaby. 1988. The effects of soil organic matter level on soil solution concentration and extractabilities of manganese, zinc and copper. *Geoderma* **42**: 177-188.
- Mehlich, A. 1953. *Determination of P, Ca, Mg, K, Na, and NH₄*. Raleigh, North Carolina, USA: Soil Testing Division, Department of Agriculture, 8p.
- Melling, L., J.U. Lau, K.J. Goh, R. Hartono and M. Osaki. 2006. Soils of Loagan Bunut National Park, Sarawak, Malaysia. *Final Report, Peat Swamp Forests Project*, Department of Agriculture, Sarawak, Malaysia.
- Michalak, A. 2006. Phenolic compounds and their antioxidant activity in plants growing under heavy metal stress. *Polish Journal of Environmental Studies* **15**: 523-530.
- Michio, Z., N. Kazuko and T. Kyoji. 1989. Spectrophotometric determination of boron with an azomethine H derivative. *Fresenius Journal of Analytical Chemistry* **334**: 238-241.
- Miller, R.N.G., M. Holderness, P.D. Bridge, G.F. Chung and M.H. Zakaria. 1999. Genetic diversity of *Ganoderma* in oil palm plantings. *Plant Pathology* **48**: 595-603.
- Muniandy, M., O.H. Ahmed, N.M.A. Majid and M.K. Yusop. 2009. Effects of converting secondary forest to oil palm plantation on peat soil carbon and nitrogen and other soil chemical properties. *American Journal of Environmental Science* **5**: 406-412.
- Murphy, J. and J.P. Riley. 1962. A modified single solution method for determination of phosphate in natural water. *Analytica Chimica Acta* **27**: 31-36.
- Murtedza, M., E. Padmanabhan, B.L.H. Mei and W.B. Siong. 2002. *The Peat Soil of Sarawak*. Malaysia: Universiti Malaysia Sarawak, 16 p.
- Mutert, E., T.H. Fairhurst and H.R. von Uexkull. 1999. Agronomic management of oil palms on deep peat. In: *Better Crops International volume 13, No.1*, eds. Armstrong, D.L. and K.P. Griffin, pp. 22-27. Canada: Potash and Phosphate Institute of Canada (PPIC).
- Naher, L., U.K. Yusuf, S.G. Tan and A. Ismail. 2013. Ecological status of *Ganoderma* and basal stem rot disease of oil palms (*Elaeis guineensis* Jacq.). *Australian Journal of Crop Science* **7**: 1723-1727.

- Nathan, M.V., J.A. Stecker and Y. Sun. 2012. *Soil Testing in Missouri - A Guide for Conducting Soil Tests in Missouri*. Missouri, USA: University of Missouri, 48 p.
- Navaratnam, S.J. and K.L. Chee. 1965. Stem rot of oil palms in the Federal Experiment Station, Serdang. *Malaysian Agriculture Journal* **45**: 175-181.
- Ng, S.K. 1972. *The Oil Palm, its Culture, Manuring and Utilisation*. Bern, Switzerland: International Potash Institute (IPI), 145 p.
- Ng, S.K., T.E. Cheah and S. Thamboo. 1969. Nutrient contents of oil palm in Malaya. IV. micronutrients in leaflets. *Malaysian Agriculture Journal* **47**: 41-52.
- Ng, S.K., H. Uexkull and R. Hardter. 2003. Botanical aspects of the oil palm relevant to crop management. In: *Oil Palm: Management for Large and Sustainable Yields*, eds. Fairhurst, T. and R. Hardter, pp. 13-26. Canada: Potash and Phosphate Institute of Canada (PPIC).
- Nudin, N.F.H. and S. Siddiquee. 2012. Comparing interfertility data with random amplified microsatellites DNA (RAMS) studies in *Ganoderma* Karst. Taxonomy. *Molecular Biology Report* **39**: 2861-2866.
- Nur Sabrina, A.A., M. Sariah and A.R. Zaharah. 2012. Suppression of basal stem rot disease progress in oil palm (*Elaeis guineensis*) after copper and calcium supplementation. *Pertanika Journal of Tropical Agriculture Science* **35**: 13-24.
- Nusaibah, S.A., S. Rajinder and A.S. Idris. 2010. Somatic incompatibility and AFLP analysis of four species of *Ganoderma* isolated from oil palm. *Journal of Oil Palm Research* **22**: 814-821.
- Obahiagbon, F.I. 2012. A Review: Aspects of the African Oil Palm (*Elaeis guineensis* Jacq.) and the Implications of its Bioactives in Human Health. *American Journal of Biochemistry and Molecular Biology* **2**: 106-119.
- Ochs, R. and J. Olivin. 1976. Research on mineral nutrition by the IRHO. In: *Oil Palm Research*, eds. Corley, R.H.V., J.J. Hardon and B.J. Wood, pp. 183-214. Amsterdam: Elsevier Scientific.
- Opgenorth, D.C., L. Butler and M. Arciero. 1983. Transmission of *Ceratocystis ulmi* on pruning saws. *Journal of Arboriculture* **9**: 196-197.
- Paavilainen, E. and J. Paivanen. 1995. *Peatland Forestry: Ecology and Principles*, Vol. 111. Berlin: Springer-Verlag, 250 p.
- Paramananthan, S. 2012. *Keys to the Identification of Malaysian Soils Using Parent Materials*, 2nd ed. Selangor, Malaysia: Param Agricultural Soil Surveys (M) Sdn. Bhd., 37 p.

- Parker, I.M. and G.S. Gilbert. 2004. The evolutionary ecology of novel plant-pathogen interactions. *Annual Review of Ecology, Evolution, and Systematics* **5**: 675-700.
- Paterson, R.R.M. 2007. *Ganoderma* disease of oil palm - A white rot perspective necessary for integrated control. *Crop Protection* **26**: 1369-1376.
- Paterson, R.R.M, M. Sariah and L. Nelson. 2009. The feasibility of producing oil palm with altered lignin content to control *Ganoderma* disease. *Journal of Phytopathology* **157**: 649-656.
- Paul, H. 1995. *Fungus Diseases of Tropical Crops*. Cambridge: Cambridge University Press, 624 p.
- Piggott, J. 1990. Growing oil palms: An illustrated guide. Kuala Lumpur, Malaysia: The Incorporated Society of Planters, 152 p.
- Pilon, M., S. E. Abdel-Ghany, C. M. Cohu, K. A. Gogolin and H. Ye. 2006. Copper cofactor delivery in plant cells. *Current Opinion in Plant Biology* **9**: 256-263.
- Pilotti, C.A., F.R. Sanderson and E.A.B. Aitken. 2002. Sexuality and interactions of monokaryotic and dikaryotic mycelia of *Ganoderma boninense*. *Mycological Research of the British Mycological Society* **106**: 1315-1322.
- Pilotti, C.A., F.R. Sanderson and E.A.B. Aitken. 2003. Genetic structure of a population of *Ganoderma boninense* on oil palm. *Plant Pathology* **52**: 455-463.
- Pilotti, C.A., F.R. Sanderson, E.A.B. Aitken and W. Armstrong. 2004. Morphological variation and host range of two *Ganoderma* species from Papua New Guinea. *Mycopathologia* **158**: 251-265.
- Pilotti, C.A. 2005. Stem rots of oil palm caused by *Ganoderma boninense*: Pathogen biology and epidemiology. *Mycopathologia* **159**: 129-137.
- Prabhu, A.S., N.D. Fageria, D.M. Huber and F.A. Rodrigues. 2007. Potassium and plant disease. In: *Mineral Nutrition and Plant Disease*, eds. Datnoff, L.E., W.H. Elmer and D.M. Huber, pp. 57-78. Saint Paul, Minnesota, USA: The American Phytopathological Society.
- Pratt, J.E. 2000. Effect of inoculum density and borate concentration in a stump treatment trial against *Heterobasidion annosum*. *Forest Pathology* **230**: 277-283.
- Prevot, P. and P. de Montbreton. 1958. Etude des gradients en divers elements minéraux selon le rang de la feuille chez le palmier à huile. *Oleagineus* **13**: 317-321.

- Rajaratnam, J.A. and J.B. Lowry. 1974. The role of boron in the oil palm (*Elaeis guineensis*). *Annals of Botany* **38**: 193-200.
- Rankine, I. and T.H. Fairhurst. 1999. Management of phosphorus, potassium and magnesium in mature oil palm. In: *Better Crops International volume 13, No.1*, eds. Armstrong, D.L. and K.P. Griffin, pp. 10-15. Canada: Potash and Phosphate Institute of Canada (PPIC).
- Rao, A.K. 1990. Basal stem rot (*Ganoderma*) in oil palm smallholdings-IADP Johore Barat experience. In: eds. Ariffin, D. and S. Jalani, *Proceedings of the Ganoderma Workshop*, 11 September 1990. Palm Oil Research Institute of Malaysia, Bangi, Selangor, Malaysia, pp.113-131.
- Rees, R.W., J. Flood, Y. Hasan and R.M. Cooper. 2007. Effects of inoculum potential, shading and soil temperature on root infection of oil palm seedlings by the basal stem rot pathogen *Ganoderma boninense*. *Plant Pathology* **56**: 862-870.
- Rees, R.W., J. Flood, Y. Hasan, U. Potter and R.M. Cooper. 2009. Basal stem rot of oil palm (*Elaeis guineensis*); mode of root infection and lower stem invasion by *Ganoderma boninense*. *Plant Pathology* **58**: 982-989.
- Rees, R.W., J. Flood, Y. Hasan, M.A. Wills and R.M. Cooper. 2012. *Ganoderma boninense* basidiospores in oil palm plantations: evaluation of their possible role in stem rots of *Elaeis guineensis*. *Plant Pathology* **61**: 567-578.
- Rengel, Z., R.D. Graham and J.F. Pedler. 1994. Time-course of byosynthesis of phenolics and lignin in roots of wheat genotypes differing in manganese efficiency and resistance to take-all fungus. *Annals of Botany* **74**: 471-477.
- Rezende, B.A., A.F.B. Abreu, M.A.P. Ramalho and E.A. Souza. 2014. Severity evaluation methods in common bean recurrent selection programme for resistance to angular leaf spot. *Journal of Phytopathology* **162**: 643-649.
- Ritchie, F., R.A. Bain and M.P. McQuilken. 2009. Effects of nutrient status, temperature and pH on mycelial growth, sclerotial production and germination of *Rhizoctonia solani* from potato. *Journal of Plant Pathology* **91**: 589-596.
- Robinson, P.W. and C.F. Hodges. 1981. Nitrogen-induced changes in the sugars and amino acids of sequentially senescing leaves of *Poa pratensis* and pathogenesis by *Drechslera sorokiniana*. *Phytopathologische Zeitschrift* **101**: 348-361.
- Romheld, V. and H. Marschner. 1991. Function of micronutrients in plants. In: *Micronutrients in Agriculture, 2nd ed.*, eds. Mortvedt, J.J., F.R. Cox, L.M. Shuman and R.M. Welch, pp. 297-328. Madison, Wisconsin, USA: Soil Science Society of America Inc.

- Sabbe, W.E. 1992. Determination of boron in plants by azomethine-H method. In: *Plant Analysis References Procedures for the Southern Region of the United States*. eds. Plank, C.O., p. 51-53. USA: Southern Cooperative Series Bulletin.
- Saeed, E.M.G., M.M.S. Amin, M.M. Hanafi, A.M.S. Rashid and W. Aimrun. 2003. Spatial variability of total nitrogen, and available phosphorus of large rice field in sawah Sempadan Malaysia. *Science Asia* **29**: 7-12.
- Said, J.M. and S.N.L. Taib. 2009. Peat stabilization with carbide lime. *UNIMAS E-Journal of Civil Engineering* **1**: 1-6.
- Salimin, M.I., S. Gandaseca, O.H. Ahmed and N.M.A. Majid. 2010. Comparison of selected chemical properties of peat swamp soil before and after timber harvesting. *American Journal of Environmental Sciences* **6**: 164-167.
- Sakihama, Y., M.F. Cohen, S.C. Grace and H. Yamasaki. 2002. Plant phenolic antioxidant and prooxidant activities: phenolics-induced oxidative damage mediated by metals in plants. *Toxicology* **177**: 67-80.
- Sanderson, F.R. 2005. An insight into spore dispersal of *Ganoderma boninense* on oil palm. *Mycopathologia* **159**: 139-141.
- Sariah, M. and H. Zakaria. 2000. The use of soil amendments for the control of basal stem rot of oil-palm seedlings. In: *Ganoderma Diseases of Perennial Crops*. eds. Flood, J., P.D. Bridge and M. Holderness, pp. 89-99. United Kingdom, Wallingford: CABI Publishing.
- Sariah, M., M.Z. Hussin, R.N.G. Miller and M. Holderness. 2007. Pathogenicity of *Ganoderma boninense* tested by inoculation of oil palm seedlings. *Plant Pathology* **43**: 507-510.
- SarVision. 2011. *Impact of oil palm plantations on peat land conversion in Sarawak 2005-2010*. Netherlands: Wageningen University, 16 p.
- Seo, G.S. and P.M. Kirk. 2000. Ganodermataceae: Nomenclature and classification. In: *Ganoderma diseases of Perennial Crops*, eds. Flood, J., P.D. Bridge and M. Holderness, pp. 3-20. United Kingdom, Wallingford: CABI Publishing.
- Shamala, S., A. Faridah, A.M.A. Zainal and K.Y. Umi. 2008. Efficacy of single and mixed treatments of *Trichoderma harzianum* as biocontrol agent of *Ganoderma* basal stem rot in oil palm. *Journal of Oil Palm Research* **20**: 470-483.
- Shen, J., L. Yuan, J. Zhang, H. Li, Z. Bai, X. Chen, W. Zhang and F. Zhang. 2011. Phosphorus dynamics: from soil to plant. *American Society of Plant Biologists* **156**: 997-1005.

- Shin, G.C, Y.H. Park, G.S. Seo and D.Y. Cha. 1986. Morphological characters of *Ganoderma lucidum* (Fr) Karsten grown naturally in Korea. *Research Report of Institute of Agriculture Science and Technology* **13**: 44-51.
- Shin, G.C. and G.S. Seo. 1988. Classification of strains of *Ganoderma lucidum*. *Korean Journal of Mycology* **16**: 235-241.
- Singh, G. 1991. *Ganoderma* – the scourge of the oil palm in the coastal areas. *Planter* **67**: 421-444.
- Spann, T.M. and A.W. Schumann. 2009. The role of plant nutrients in disease development with emphasis on citrus and huanglongbing. *Proceedings of the Florida State Horticultural Society* **122**: 169-171.
- Steyaert, R.L. 1980. Study of some *Ganoderma* species. *Bulletin du Jardin Botanique National de Belgique* **50**: 135-186.
- Tarmizi, A.M., H. Othman, M.D. Farawahida, M.H. Haniff, M.Z. Pilus, A.B. Ismail and H. Wosten. 2009. BMP on peat in relation to carbon emission. In: *Proceeding of the PIPOC 2009 International Palm Oil Congress - Balancing Ecologies with Economics*, 9-12 November 2009. Kuala Lumpur, Malaysia, p. 18
- Tayeb, M. D. and A.B. Hamdan. 1999. Relation of fertilizer nutrients to *Ganoderma*. In: eds. Ariffin, D., K.W. Chan and S.R.S.A. Sharifah, *Proceeding of the 1999 International Oil Palm Congress*, 1-6 February 1999. Palm Oil Research Institute of Malaysia, Bangi, Selangor, Malaysia, pp. 422-453.
- Tayeb, M.D. 2005. *Technologies for Planting Oil Palm on Peat*. Bangi, Selangor: Malaysian Palm Oil Board (MPOB), 84 p.
- Thompson, A. 1931. Stem rot of oil palm in Malaysia. *Bulletin Department of Agriculture, Straits Settlements and F.M.S., Science Series* 6: 23pp.
- Thompson, A. 1937. Observations on stem rot of the oil palm. *Bulletin Department of Agriculture, Straits Settlements and F.M.S., Science Series* 21: 20pp.
- Tie, Y.L. 1982. Soil classification in Sarawak. *Technical Paper No. 6*. Department of Agriculture, Sarawak. 86 pp.
- Tohiruddin, L., J. Tandiono, J.S. Abner, N.E. Prabowo and H.L. Foster. 2010. Effects of N, P and K fertilizers on leaf trace element levels of oil palm in Sumatra. *Journal of Oil Palm Research* **22**: 869-877.
- Turner, P.D. 1969. Observations on the incidence, effects and control of upper stem rot in oil palm. In: *Progress in Oil Palm*, ed. Turner, P.D., pp. 143-154. Kuala Lumpur: Incorporated Society of Planters.

- Turner, P.D. and R.A. Gillbanks. 2003. *Oil Palm Cultivation and Management*. Kuala Lumpur, Malaysia: The Incorporated Society of Planters, 915 p.
- Turner, P.D. 1981. *Oil Palm Diseases and Disorders*. Oxford: Oxford University Press, 280 p.
- Tutiempo.net. 2013. Weather Miri and Sib. <http://www.tutiempo.net/en/weather/malaysia/my.html>. Retrieved 1st May 2013.
- United States Department of Agriculture (USDA). 2014. *Oilseeds: World Markets and Trade*. USA: Foreign Agricultural Service, USDA, 34 p.
- Utomo, C., S. Wernrr, F. Niepold and H.B. Deising. 2005. Identification of *Ganoderma*, the causal agent of basal stem rot disease in oil palm using a molecular method. *Mycopathologia* **159**: 159-170.
- Van de Lande, H.L. 1993. Spatio-temporal analysis of spear rot and 'marchitez sorpresiva' in African oil palm in Surinam. *Netherlands Journal of Plant Pathology* **99**: 129-138.
- Van de Lande, H.L. and J.C. Zadoks. 1999. Spatial patterns of spear rot in oil palm plantations in Surinam. *Plant Pathology* **48**: 189-201.
- von Uexkull, H.R. and T.H. Fairhurst. 1991. *Fertilizing for High Yield and Quality - The Oil Palm*. Bern, Switzerland: International Potash Institute (IPI), 79 p.
- von Uexkull, H.R. and T.H. Fairhurst. 1999. Some nutritional disorders in oil palm. In: *Better Crops International volume 13, No.1*, eds. Armstrong, D.L. and K.P. Griffin, pp. 16-21. Canada: Potash and Phosphate Institute of Canada (PPIC).
- Wahid, O., S. Paramanathan, M.H. Harun, N.A. Aziz and A. Kushairi. 2013. Malaysian unified peat classification technique. *Malaysian Palm Oil Board Information Series* 627, MPOB TT No.529: 8pp.
- Webster, J. and R.W.S. Weber. 2007. *Introduction to Fungi*. New York: Cambridge University Press, 841 p.
- Wo, S., X. Guo, X. Zhou, X. Li, Y. Chen and J. Lin. 2009. AFLP analysis of genetic diversity in main cultivated strains of *Ganoderma* species. *African Journal of Biotechnology* **8**: 3448-3454.
- Wong, L.C., C.F.J. Bong and A.S. Idris. 2012. *Ganoderma* species associated with basal stem rot disease of oil palm. *American Journal of Applied Sciences* **9**: 879-885.
- Wong, L.C. 2013. Identification of *Ganoderma* Species Isolated from Basal Stem Rot Infected Oil Palms in Sarawak. M.Sc Thesis. Universiti Putra Malaysia, Selangor, Malaysia.

- Yruela, I. 2005. Copper in plants. *Brazilian Journal of Plant Physiology* **17**: 145-156.
- Yuan, M., Z. Chu, X. Li, C. Xu and S. Wang. 2010. The bacterial pathogen *Xanthomonas oryzae* overcomes rice defences by regulating host copper redistribution. *The Plant Cell* **22**: 3164-3176.
- Zaiton, S., M. Sariah and A.M.A. Zainal. 2008. Effect of endophytic bacteria on growth and suppression of *Ganoderma* infection in oil palm. *International Journal of Agriculture and Biology* **10**: 127-132.
- Zenki, M., K. Nose and K. Toei. 1989. Spectrophotometric determination of boron with an azomethine H derivative. *Fresenius Z Analytical Chemistry* **334**: 238-241.

BIODATA OF STUDENT

Mohd Rakib bin Mohd Rashid was born in 4th December 1985 in Beaufort, Sabah, Malaysia. He received his primary education in Sekolah Rendah Kebangsaan Kundasang Pekan Ranau in 1992, Sekolah Rendah Kebangsaan Tanjung Aru 2 in 1995, and Sekolah Kebangsaan Likas in 1996. In 1998, he pursued his secondary education in Sekolah Menengah Kebangsaan Likas. He completed his Sijil Tinggi Pelajaran Malaysia (STPM) in Sekolah Menengah All Saints, Kota Kinabalu in 2004. In the following year, he continued his higher education in Universiti Putra Malaysia Bintulu Campus and graduated with Bachelor of Science Bioindustry (Honours First Class) in 2009. Afterwards, he was enrolled in degree of Doctor of Philosophy in Plant Pathology since year 2009. During the postgraduate studies, he has published two impact factor journal articles, attended several international conferences inside and outside Malaysia, and enrolled in several academicians part-time jobs.

LIST OF PUBLICATIONS

Journal Articles Published:

1. Rakib, M.R.M., C.F.J. Bong, A. Khairulmazmi and A.S. Idris. 2014. Genetic and morphological diversity of *Ganoderma* species isolated from infected oil palms (*Elaeis guineensis*). *International Journal of Agriculture and Biology* **16**: 691-699.
2. Rakib, M.R.M., C.F.J. Bong, A. Khairulmazmi and A.S. Idris. 2014. Occurrence and spatial distribution of *Ganoderma* species causing upper and basal stem rot in oil palm. *Journal of Food, Agriculture and Environment* **12**: 360-364.

Journal Article Submitted:

1. Rakib, M.R.M., C.F.J. Bong, A. Khairulmazmi and A.S. Idris. Aggressiveness of *Ganoderma boninense* and *G. zonatum* isolated from upper- and basal stem rot of oil palm (*Elaeis guineensis*) in Malaysia. *Journal of Oil Palm Research*.

Conference Papers / Proceedings:

1. Rakib, M.R.M., C.F.J. Bong, A. Khairulmazmi and A.S. Idris. 2013. Genetic and morphological diversity of *Ganoderma* sp. from upper and basal stem rot infected oil palms. In: eds. Nadarajah, K., L.Y. Sze, G. Krishnasamy, F.D.A. Bakar, J. Santanam, M.J. Masarudin, N. Shahab, S.C. Chin, V. Sabaratnam, T.K. Lin and R.A. Rahim, *International Congress of the Malaysian Society for Microbiology*, 12-15 December 2013. Langkawi, Kedah, Malaysia, pp. 80-84.
2. Rakib, M.R.M., C.F.J. Bong, A. Khairulmazmi and A.S. Idris. 2014. Association of copper and zinc to *Ganoderma* spp. spatial distribution in oil palm (*Elaeis guineensis*) plantations on peat. In: *International Conference on Plant Physiology*, 26-28 August 2014. Kuta, Bali, Indonesia.
3. Rakib, M.R.M., C.F.J. Bong, A. Khairulmazmi and A.S. Idris. 2014. Upper stem rot: another threat to oil palm (*Elaeis guineensis*) plantations based on a study in Sarawak. In: *2nd National Postgraduate Symposium on Sustainable Agriculture*, 8-9 October 2014. Universiti Malaysia Sabah (UMS) Sandakan Campus, Sabah, Malaysia.
4. Bong, C.F.J, M.R.M. Rakib and L.C. Wong. 2014. *Ganoderma* diseases in Sarawak. In: *Workshop on Integrated Management of Ganoderma Disease in Oil Palm*, 3-4 December 2014. Kota Kinabalu, Sabah, Malaysia.