




UNIVERSITI PUTRA MALAYSIA

***RELATIONSHIP OF SCROTAL CIRCUMFERENCE AND
TESTICULAR VOLUME TO AGE AND BODY WEIGHT IN DORPER
SHEEP***

NUR AIN SAKINAH BINTI ISHAK

FPV 2017 41

RELATIONSHIP OF SCROTAL CIRCUMFERENCE AND TESTICULAR
VOLUME TO AGE AND BODY WEIGHT IN DORPER SHEEP



SERDANG, SELANGOR

2017

RELATIONSHIP OF SCROTAL CIRCUMFERENCE AND TESTICULAR
VOLUME TO AGE AND BODY WEIGHT IN DORPER SHEEP

by




NUR AIN SAKINAH BINTI ISHAK

A project paper submitted to the
Faculty of Veterinary Medicine, Universiti Putra Malaysia
In partial fulfilment of the requirement for the
DEGREE OF DOCTOR OF VETERINARY MEDICINE
University Putra Malaysia
Serdang, Selangor Darul Ehsan



MARCH 2017

It is hereby certified that I have read this project entitled "Relationship of Scrotal Circumference and Testicular Volume to Age and Body Weight in Dorper Sheep", by Nur Ain SakinahBintiIshak and in my opinion it is satisfactory in terms of scope, quality, and presentation as partial fulfilment of the requirement for the course VPD 4999 – Final Year Project.



**ASSOC. PROF. DR. ROSNINA HJ. YUSOFF
D.V.M (UPM), M.Sc. (UPM), Ph.D. (Guelph)**

Lecturer,

Faculty of Veterinary Malaysia

Universiti Putra Malaysia, Serdang, Selangor

(Supervisor)

DEDICATIONS

In the name of Allah SubhanahuWaTa'ala

I dedicate this thesis to:

My beloved Mummy, Ayah, and my siblings,

lecturers and friends.

Thank you for all the loves and support.



ACKNOWLEDGEMENTS

I am thankful to Allah SWT for the opportunity to be a part of Doctor of Veterinary Medicine society and providing me strength and health throughout the study period.

I submit my heartiest gratitude to my supervisor, Assoc. Prof.Dr.RosninaHj. Yusoff, for her guidance, patience, kindness and motherhood throughout the whole process. I wish to express my appreciation to Prof.Dr. Mohammad Ariff Omar for his guidance, time, and suggestion especially regarding the statistical analysis.

I would like to acknowledge EncikPaizal Jamaluddin, the owner of PJ Livestock SdnBhd and also all personnel for welcoming me there. My deepest thanks and appreciation goes to EncikZulfadhliAizat and EncikMohd. FaizalHamzah for their help in restraining and data collection.

My endless thanks and grateful to my family, especially my father, IshakMdYusoff and my mother, ZalehaMd Dam for their loves, du'a and patience in raising me until today.

Special thanks to my partner during the final year project, NurulLiyana Jamaluddin for her support, company and guidance. My sincere thanks to DVM class 2017, especially my closest friends, Naseeha Sakina Diauudin and Nur Ain SyahiraIshak for their support and laughter. I am deeply indebted to all my respected lecturers for the knowledge.

Lastly, thank you to everyone who has contributed in one way or another in this project.

CONTENTS

	Page
TITLE.....	i
CERTIFICATION	ii
DEDICATIONS	iii
ACKNOWLEDGEMENTS	iv
CONTENTS	v
LIST OF ABBREVIATIONS	vii
LIST OF DIAGRAMS	vii
LIST OF TABLES	viii
LIST OF FIGURES	viii
ABSTRAK	ix
ABSTRACT	xi
1.0 INTRODUCTION	1
2.0 LITERATURE REVIEWS	2
2.1 Dorper sheep	3
2.2 Breeding Soundness Examination (BSE)	3
2.3 Testicular size and sperm production	4
3.0 MATERIALS AND METHODS.....	5
3.1 Study area and sample size	5
3.2 Body condition score	5
3.3 Testicular measurement	6
3.4 Statistical analysis.....	7
4.0 RESULTS	9

5.0	DISCUSSIONS.....	15
6.0	CONCLUSIONS AND RECOMMENDATIONS	18
7.0	REFERENCES	19
8.0	APPENDICES	25
	Appendix A: Reproductive parameters in Dorper sheep	25
	Appendix B: Testicular volume for left and right testes	26

LIST OF ABBREVIATION

Scrotal Circumference	: SC
Body Weight	: Bwt
Body Condition Score	: BCS
Testicular Volume	: TV

LIST OF DIAGRAMS

	Page
Diagram 1: Caudal view for the measurement of height (left) and length (right) of the testis	8
Diagram 2: Lateral view for the measurement of width of the testis.....	8
Diagram 3: Measurement of scrotal circumference	8

LIST OF TABLES

	Page
Table 1: Evaluation of reproductive parameters in Dorper sheep	12
Table 2: Correlation coefficient (r) between age, body weight, body condition score, scrotal circumference and testicular volume in Dorper sheep	13

LIST OF FIGURES

	Page
Figure 1: Comparative testicular volume in left and right testes of Dorper sheep	9
Figure 2: Total testicular volume of two different age group	10
Figure 3: Distribution of mean on all parameters	10
Figure 4: Relationship between scrotal circumference and body weight for two different age groups in Dorper rams.....	14

ABSTRAK

Abstrak daripada kertas projek yang dikemukakan kepada Fakulti Perubatan Veterinar untuk memenuhi sebahagian daripada keperluan kursus VPD 4999 – Projek Tahun Akhir

**HUBUNGKAIT ANTARA LILITAN SKROTAL DAN ISI PADU TESTIS
DENGAN UMUR DAN BERAT BADAN KAMBING BEBIRI DORPER**

oleh

Nur Ain Sakinah Binti Ishak

2017

Penyelia: Prof. Madya Dr. Rosnina Hj. Yusoff

Kajian ini dijalankan untuk menilai parameter pembiakan yang boleh digunakan sebagai petunjuk untuk menganggar kesuburan kambing bebiri Dorper. Lilitan skrotal (SC) dan isipadu testis (TV) adalah parameter pembiakan yang diukur untuk dikolerasikan dengan umur dan berat badan. Sebanyak 32 kambing bebiri jantan Dorper dibahagikan kepada dua kumpulan: Kumpulan A (berusia <12 bulan) dan Kumpulan B (berusia >12 bulan). Terdapat enam ekor kambing bebiri jantan di dalam Kumpulan A dan 26 ekor kambing bebiri jantan di dalam Kumpulan B. SC

kambing bebiri jantan tersebut diukur. Panjang, lebar dan ketinggian testis kiri dan kanan telah diukur untuk mengira TV. Semua data telah dianalisiskan secara statistik. Keputusan terkini menunjukkan bahawa semua parameter berkorelasi secara positif antara satu sama lain. SC dan TV lebih berhubung kait dengan berat badan berbanding dengan usia. SC berkorelasi positif dengan umur ($r=0.605$), berat badan ($r=0.824$) dan TV ($r=0.925$). TV berkorelasi positif dengan umur ($r=0.535$), berat badan ($r=0.873$) dan SC ($r=0.925$). Kajian ini menunjukkan bahawa kambing bebiri jantan berusia lebih daripada 12 bulan dengan berat badan sekurang-kurangnya 45 kg mempunyai SC dan TV yang lebih tinggi. Oleh itu, umur, berat badan, lilitan skrotal dan isipadu testis boleh digunakan sebagai salah satu kriteria dalam pemilihan baka induk.

Kata Kunci: *Kambing bebiri Dorper, lilitan skrotal, isipadu testis*

ABSTRACT

An abstract of the project paper presented to the Faculty of Veterinary Medicine in partial fulfilment of course VPD 4999 – Final Year Project.

RELATIONSHIP OF SCROTAL CIRCUMFERENCE AND TESTICULAR VOLUME TO AGE AND BODY WEIGHT IN DORPER SHEEP

by

Nur Ain SakinahBintiIshak

February, 2017

Supervisor: Assoc. Prof.Dr.RosninaHj. Yusoff

This study was conducted to evaluate reproductive parameters that can be used as indicators to estimate fertility status of Dorper sheep. Scrotal circumference (SC) and testicular volume (TV) were the reproductive parameters that were measured to correlate with age and body weight. A total of 32 Dorper rams were divided into two groups: Group A (< 12 months old) and Group B (> 12 months old). There were six rams in Group A and 26 rams in Group B. The SC of the rams was measured. The length, width and height of left and right testes were measured to calculate TV. All the data were analysed statistically. The present results showed that all parameters are positively correlated to each other. The SC and TV are more correlated to body weight compared with age. The SC is positively correlated with age ($r=0.605$), body

weight ($r=0.824$) and TV ($r=0.925$). The TV is positively correlated with age ($r=0.535$), body weight ($r=0.873$) and SC ($r=0.925$). This study shows that the rams of more than 12 months with body weight of at least 45kg have higher SCand TV. Therefore, age, body weight, scrotal circumference and testicular volume can be used as criteria in selection of sires.

Keywords:*Dorper sheep, scrotal circumference, testicular volume.*

1.0 INTRODUCTION

The livestock industry is important in the agricultural sector of Malaysia. The contribution of livestock production to the Gross Domestic Product (GDP) is inconsistent. Even though there was a decreased in contribution of livestock industry to GDP in the 90's, but in 2005, livestock industry contribution to GDP was slightly increased (0.8%) (Government of Malaysia Economic Report 1990 – 2005, as cited in Mohamed, 2007). The livestock industry includes non-ruminant and ruminant industries. The non-ruminant industry comprises swine and poultry, while the ruminant industry are cattle, buffaloes, goats and sheep. However, goats and sheep play a minor role in livestock industry. The sheep population in Malaysia has increased from 116 387 in 2006 to 132 900 in 2011 (Ministry of Agriculture and Agro-Based Industry Malaysia, 2011). Furthermore, the number of sheep slaughtered from 2006 to 2011 has also increased from 3 554 to 4 171 heads.

Most rams reach puberty at age between 5 and 7 months old. Breeding soundness evaluation is done to predict the potential fertility of the males. It includes physical examination of the animals, assessment of the reproductive organs and semen evaluation. The scrotal circumference is an important measurement as it is related to the current semen production capacity of the rams (Hahn et al., 1969). Studies have shown that age can influence the scrotal circumference and semen characteristics (Toe et al., 1994). This shows that age is directly related to the testicular size and at the same time related to the sperm output. This study was done in hope to enable farmers to identify suitable sires to be used in breeding programme

based on parameters such as age, body weight, scrotal circumference and testicular volume.

Although there is information regarding the relationship of scrotal circumference to age and body weight in sheep, goats and bulls, reports about these in Dorper sheep is inadequate. Therefore, this study was performed to correlate the scrotal circumference and testicular volume to age and body weight in Dorper rams.

7.0 REFERENCES

- Bedford-Guaus, S. J. (2010). Breeding soundness examination of the male. In C. M. Kahn & S. Line (Eds.), *The Merck Veterinary Manual* (pp. 1966-1969). Whitehouse Station, N.J: Merck & Co.
- Boligon, A. A., Silva, J. A. V, Sesana, R. C., Sesana, J. C., Junqueira, J. B., & Albuquerque, L. G. (2010). Estimation of genetic parameters for body weights, scrotal circumference, and testicular volume measured at different ages in Nellore cattle. *Journal of Animal Science*, 88(4), 1215–1219.
<http://doi.org/10.2527/jas.2008-1719>
- Bongso, T. A., Hassan, M. D., & Nordin, W. (1984). Relationship of scrotal circumference and testicular volume to age and body weight in the swamp buffalo (*Bubalus bubalis*). *Theriogenology*, 22(2), 127–134.
[http://doi.org/10.1016/0093-691X\(84\)90425-4](http://doi.org/10.1016/0093-691X(84)90425-4)
- Brito, L. F., Silva, A. E., Unanian, M. M., Dode, M. A., Barbosa, R. T., & Kastelic, J. P. (2004). Sexual development in early-and late-maturing *Bos indicus* and *Bos indicus* × *Bos taurus* crossbred bulls in Brazil. *Theriogenology*, 62(7), 1198–1217. <http://dx.doi.org/10.1016/j.theriogenology.2004.01.006>
- Cloete, S. W. P., Snyman, M. A., & Herselman, M. J. (2000). Productive performance of Dorper sheep. *Small Ruminant Research*, 36(2), 119–135.
[http://doi.org/10.1016/S0921-4488\(99\)00156-X](http://doi.org/10.1016/S0921-4488(99)00156-X)

- Devkota, B., Koseki, T., Matsui, M., Sasaki, M., Kaneko, E., Miyamoto, A., ...& Miyake, Y. I. (2008). Relationships among age, body weight, scrotal circumference, semen quality and peripheral testosterone and estradiol concentrations in pubertal and postpubertal Holstein bulls. *Journal of Veterinary Medical Science*, 70(1), 119-121. <http://doi.org/10.1292/jvms.70.119>
- Fourie, P. J., Neser, F. W. C., Olivier, J. J., & Van Der Westhuizen, C. (2002). Relationship between production performance, visual appraisal and body measurements of young Dorper rams. *South African Journal of Animal Sciences*, 32(4), 256–262.
- Fourie, P. J., Schwalbach, L. M., Neser, F. W. C., & Van Der Westhuizen, C. (2004). Scrotal, testicular and semen characteristics of young Dorper rams managed under intensive and extensive conditions. *Small Ruminant Research*, 54(1–2), 53–59. <http://doi.org/10.1016/j.smallrumres.2003.10.011>
- Gouletsou, P. G., & Fthenakis, G. C. (2010). Clinical evaluation of reproductive ability of rams. *Small Ruminant Research*, 92(1–3), 45–51. <http://doi.org/10.1016/j.smallrumres.2010.04.005>
- Gouletsou, P. G., Galatos, A. D., & Leontides, L. S. (2008). Comparison between ultrasonographic and caliper measurements of testicular volume in the dog. *Animal reproduction science*, 108(1), 1-12. <http://dx.doi.org/10.1016/j.anireprosci.2007.06.020>

- Hahn, J., Foote, R. H., & Seidel, G. E. (1969). Testicular growth and related sperm output in dairy bulls. *Journal of Animal Science*, 29(1), 41–47.
<http://doi.org/10.2527/jas1969.29141x>
- Kimberling, C. V., & Parsons, G. A. (2007). Current Therapy in Large Animal Theriogenology. *Current Therapy in Large Animal Theriogenology*, 620–628.
<http://doi.org/10.1016/B978-072169323-1.50088-X>
- Koyuncu, M., & Altincekic, S. O. (2013). Testicular development and body growth in prepubertal ram lambs of Kivircik, Karacabey Merino and Tahirova in South Marmara region of Turkey. *Bulgarian Journal of Agricultural Science*, 19(1), 152–157.
- Koyuncu, M., Uzun, S. K., Ozis, S., & Duru, S. (2005). Development of testicular dimensions and size, and their relationship to age and body weight in growing Kivircik (western thrace) ram lambs. *Czech Journal of Animal Science*, 50(6), 243–248.
- Loh, T.C. (2002). Livestock production and the feed industry in Malaysia. *Protein sources for the Animal Feed Industry, FAO*. 329 – 339.
- Maurya, V. P., Sejian, V., Kumar, D., & Naqvi, S. M. K. (2015). Impact of heat stress, nutritional restriction and combined stresses (heat and nutritional) on growth and reproductive performance of Malpura rams under semi-arid tropical environment. *Journal of animal physiology and animal nutrition*.
<http://doi.org/10.1111/jpn.12443>

Ministry of Agriculture and Agro-Based Industry Malaysia, (2011). Retrieved on

November 2016 from

[http://www.moa.gov.my/c/document_library/get_file?uuid=851fc0ee-240c-](http://www.moa.gov.my/c/document_library/get_file?uuid=851fc0ee-240c-4cab-b977-d37e0ca37992&groupId=43204)

[4cab-b977-d37e0ca37992&groupId=43204](http://www.moa.gov.my/c/document_library/get_file?uuid=851fc0ee-240c-4cab-b977-d37e0ca37992&groupId=43204)

Mohamed, Z. A. (2007). Chapter 23: The livestock industry. In F. M. Arshad, N. M.

Raja Abdullah, B. Kaur, & A. M. Abdullah (Eds.), *50 years of Malaysian agriculture: transformational issues, challenges and direction* (pp. 553-582).

Penerbit Universiti Putra Malaysia.

Mukasa-Mugerwa, E., & Ezaz, Z. (1992). Relationship of testicular growth and size

to age, body weight and onset of puberty in Menz ram lambs. *Theriogenology*,

38(5), 979–988. [http://dx.doi.org/10.1016/0093-691X\(92\)90172-N](http://dx.doi.org/10.1016/0093-691X(92)90172-N)

Omar, C. (2016). Study of some testicular dimensions and their relationship to body

weight in Karadi ram lambs. *Assiut Veterinary Medical Journal*, 62, 31–38.

Perumal, P. (2014). Scrotal Circumference and Its Relationship with Testicular

Growth, Age, and Body Weight in Tho Tho (*Bos indicus*) Bulls. *International Scholarly Research Notices*, 2014. <http://dx.doi.org/10.1155/2014/249537>

Pezzanite, L., Bridges, D. A., Neary, D. M., & Hutchens, T. (2010). Breeding

Soundness Examination of Rams and Bucks. *Purdue Extension*, 1–4. Retrieved

on January 2017 from <https://www.extension.purdue.edu/extmedia/as/as-599-w.pdf>

- Pukazhenth, B. S., Togna, G. D., Padilla, L., Smith, D., Sanchez, C., Pelican, K., & Sanjur, O. I. (2011). Ejaculate traits and sperm cryopreservation in the endangered Baird's tapir (*Tapirus bairdii*). *Journal of Andrology*, 32(3), 260–270. <http://doi.org/10.2164/jandrol.110.011833>
- Ruttle, J. L., Southward, G. M., Sciences, R., Economics, F., & Cruces, L. (1988). Influence of Age and Scrotal Circumference, 29(4), 945–949.
- Salhab, S. A., Zarkawi, M., Wardeh, M. F., Al-Masri, M. R., & Kassem, R. (2001). Development of testicular dimensions and size, and their relationship to age, body weight and parental size in growing Awassi ram lambs. *Small ruminant research*, 40(2), 187-191. [http://dx.doi.org/10.1016/S0921-4488\(00\)00224-8](http://dx.doi.org/10.1016/S0921-4488(00)00224-8)
- Schoeman, S. J., & Combrink, G. C. (1987). Testicular development in Dorper, Dohne Merino and crossbred rams. *South African journal of animal science*, 17(1), 22-26.
- Thompson, J., & Meyer, H. (1994). Body condition scoring of sheep. Oregon State University, Extension Service. Retrieved from <https://ir.library.oregonstate.edu/xmlui/bitstream/handle/1957/14303/ec1433.pdf>
- Toe, F., Lahou-Kassi, A., & Mukasa-Mugerwa, E. (1994). Semen characteristics of Ile-de-France rams of different age and physical condition. *Theriogenology*, 42(2), 321-326. [http://dx.doi.org/10.1016/0093-691X\(94\)90276-3](http://dx.doi.org/10.1016/0093-691X(94)90276-3)

Wahid, S. A., & Yunus, J. M. (1994). Correlation between testicle measurements and libido and semen quality in rams. *Asian-Australasian journal of animal sciences*, 7(2), 175-178. <https://doi.org/10.5713/ajas.1994.175>

