



UNIVERSITI PUTRA MALAYSIA

PRODUCTION OF RESISTANT STARCH TYPE III AND IV AND THEIR IN VIVO CHARACTERISATION USING BALB/c MICE MODEL

SITI AISYAH MOHD ZAMAN

FSPM 2015 16



PRODUCTION OF RESISTANT STARCH TYPE III AND IV AND THEIR *IN VIVO* CHARACTERISATION USING BALB/c MICE MODEL

By

SITI AISYAH BINTI MOHD ZAMAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

November 2015

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malays



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of
the requirement for the degree of Master of Science

PRODUCTION OF RESISTANT STARCH TYPE III AND IV AND THEIR *IN VIVO* CHARACTERISATION USING BALB/c MICE MODEL

By

SITI AISYAH BINTI MOHD ZAMAN

November 2015

Chair: Shahrul Razid Sarbini, PhD

Faculty: Agricultural and Food Sciences (Bintulu)

Sago is an important agricultural commodity for Sarawak. The potential of sago is highlighted in the Malaysia's 11th plan as part of the poverty eradication programme. Sago starch indigestibility provides a new perspective of sago utilisation as a prebiotic. The indigestible portion of starch is termed as resistant starch (RS). In the present study, RS type III and IV were produced through retrogradation and chemical modifications respectively. Retrogradation was done on different starch paste concentration of 10, 18 and 30%. The resultant RS type III was analysed for solubility, swelling power, amylose and RS content. Resistant starch type IV was produced through hydroxypropylation, acetylation, cross-linking and double modification combining acetylation and cross-linking. The resultant RS was subjected to the same analysis as RS type III. Native sago starch, retrograded starch at 18% initial starch paste concentration, 2% acetylated starch and double modified starch combining 1.5% cross-linking and 2% acetylation were further analysed for prebiotic evaluation through *in vivo* study using BALB/c mice model. All RS was supplemented into a standardised AIN93-M feed formulation. Feeding treatment was conducted for 4 weeks. The daily feed intake and body weight were recorded. Faeces samples were collected on the eighth day and on the final three consecutive days of feeding treatment. All faeces samples were subjected to short chain fatty acid analysis using high-performance liquid chromatography. All starch modifications showed an increase in solubility and swelling power. Native sago showed a high content of RS at 69%. Retrograded starch showed a significantly low amylose and RS content than native starch ($P < 0.05$). Only 1.5 and 2% acetylation, as well as double modification of 1.5% cross-linking and 2% acetylation showed an increase in RS content than native sago starch. Cross-linking and double modified starch showed an increase in amylose content at all level of modification. Mice fed diet supplemented with RS type III and acetylation showed a better overall growth performance with an increase of body weight with a decrease in feed intake. Mice fed diet supplemented with double modified and native sago starch showed a weight loss with a decrease in feed intake. This pattern supported satiety

properties. Short chain fatty acid analysis showed the highest lactate production in mice fed diet supplemented with double modified starch at 43.57%. The highest propionate concentration was shown in mice fed diet supplemented with RS type III and double modified starch at 10.4%. The results showed that double modified sago starch has the potential as a prebiotic candidate ingredient. The weight reduction potential shown in a diet supplemented with double modified starch could also be further implemented in a weight management diet.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Sarjana Sains

PENGHASILAN KANJI RINTANG JENIS III DAN IV DAN PENCIRIAN IN VIVO MENGGUNAKAN MODEL TIKUS BALB/c

Oleh

SITI AISYAH BINTI MOHD ZAMAN

November 2015

Pengerusi: Shahrul Razid Sarbini, PhD

Fakulti: Sains Pertanian dan Makanan (Bintulu)

Sago ialah komoditi pertanian yang penting di Sarawak, kebanyakannya disebabkan oleh potensi sago untuk berkembang subur di tanah gambut. Perladangan sago telah dibesarkan untuk membangunkan sebahagian daripada 1.5 hektar tanah gambut yang tidak mempunyai sistem perparitan yang berkualiti. Potensi sago telah ditekankan di dalam pelan Malaysia ke-11 sebagai sumber untuk program penghapusan kemiskinan. Ketidakhadaman kanji sago memberi perspektif baru dalam penggunaan kanji sago sebagai prebiotik. Bahagian kanji yang tidak dihadam digelar sebagai kanji rintang. Kanji rintang jenis 3 dan 4 dihasilkan melalui proses retrogradasi dan pengubahsuai secara kimia. Proses retrogradasi telah dilakukan ke atas adunan kanji sago dengan kepekatan yang berbeza iaitu 10, 18 dan 30%. Kanji rintang jenis 3 yang terhasil telah dianalisa untuk kelarutan, kuasa pembengkakan, kandungan amilosa dan kanji rintang. Kanji rintang jenis 4 telah dihasilkan melalui proses hidrosipropilasi, pengasetilan, penghubungan silang dan dwi modifikasi yang mencamtmukan proses pengasetilan dan penghubungan silang. Kanji rintang jenis 4 yang terhasil telah menjalani analisa yang sama seperti kanji rintang jenis 3. Kanji sago asli, kanji retrogradasi pada 18% permulaan kepekatan adunan kanji, 2% kanji asetilan dan dwi modifikasi yang menggabungkan 1.5%, penghubungan silang dan 2% pengasetilan telah dianalisa dengan lebih jauh untuk penilaian sebagai prebiotik melalui kajian *in vivo* menggunakan model mencit BALB/c. Kesemua kanji rintang ditambah kedalam rumusan makanan AIN93-M yang telah diseragamkan. Rawatan pemakanan dilaksanakan selama 4 minggu dan pengambilan pemakanan dan berat badan telah direkodkan secara harian. Sampel najis telah dikumpulkan pada hari ke-8 dan 3 hari terakhir rawatan pemakanan. Kesemua sampel najis telah dianalisis melalui kromatografi cair berprestasi tinggi untuk analisa asid lemak rantaian pendek. Kesemua modifikasi telah menunjukkan peningkatan kelarutan dan kuasa kebengkakan. Kanji sago asli menunjukkan kandungan kanji rintang yang tinggi pada 69%. Kanji retrogradasi menunjukkan kandungan amilosa dan kanji rintang yang nyata rendah berbanding kanji asli ($P < 0.05$). Hanya pengasetilan pada tahap 1.5 dan 2% dan juga dwi modifikasi menglibatkan 1.5% penghubungan silang dan 2% pengasetilan menunjukkan peningkatan kandungan kanji rintang berbanding kanji asli. Kanji yang dihubung silang dan di dwi modifikasi menunjukkan peningkatan kandungan amilosa di semua tahap modifikasi. Mencit yang diberi makan diet

pemakanan yang ditambah kanji rintang jenis 3 dan kanji asetilasi menunjukkan prestasi pembesaran keseluruhan yang lebih baik dengan peningkatan berat badan diiringi penurunan kadar pengambilan makanan. Mencit yang diberi makan dengan rumusan pemakanan yang ditambah dengan kanji dwi modifikasi dan kanji sago asli menunjukkan penurunan berat badan dengan diiringi penurunan kadar pengambilan makanan. Corak pemakanan dan berat badan ini menyokong sifat kekenyangan. Analisis asid lemak rantai pendek menunjukkan jumlah penghasilan laktat yang paling tinggi oleh mencit yang diberi makan diet makanan yang ditambah dengan kanji dwi modifikasi pada kadar 43.57%. Kepakatan propionat tertinggi ditunjukkan oleh mencit yang diberi makan diet makanan yang ditambah dengan kanji rintang jenis 3 dan kanji dwi modifikasi pada kadar 10.4%. Data menunjukkan kanji dwi modifikasi mempunyai potensi sebagai calon prebiotik secara *in vivo*. Potensi penurunan berat badan yang ditunjukkan di dalam diet yang ditambah rumusan kanji dwi modifikasi juga boleh selanjutnya digunakan di dalam diet pemakanan untuk pengurusan berat.

ACKNOWLEDGEMENTS



There are so many things to be thankful and subsequently so many people to acknowledge. Throughout the course of my research, across 3 states and 4 cities, I have been fortunate enough to encounter exemplary people. Thus, this author wishes to express her deepest gratitude.

First and foremost, all praise be to Allah S.W.T. for seeing me through it all. "*And whatever of blessings and good things you have, it is from Allah*" [al-Nahl 16:53].

Heartfelt thanks and gratitude to my parents, for the immense emotional support, for all the encouraging words and for always reminding me that I am never alone. Thanks for always believing in me, my dream and my vision even though at times they seems unreachable.

Thanks also to my whole family, especially my siblings for their unending support, both emotionally and financially. Also lots of love to my niece and nephew that often made me forget the stress of being a research student once I reached home.

I would also like to express my heartfelt gratitude and appreciation to my advisor, Dr Shahrul Razid Sarbini, for his excellent supervision, guidance and patience all throughout my research up to the *viva voce* and the completion of this thesis. Thanks for all the motivation and the positive outlook when things look sour. Most of all, thanks for being such a great teacher.

My sincerest thanks also to my co-supervisor, Dr Patricia King Jie Hung, for her guidance and helping hand, particularly through the thesis preparation and submission and the *viva voce*.

Sincerest gratitude also to all staff at Faculty of Agricultural and Food Sciences, UPM Bintulu Campus; especially to Miss Siti Aziah for all her assistance in acquiring lab

necessities and managing the lab altogether. Thanks also to other lab assistants that has helped me throughout the completion of this project.

I would also like to express my thanks to all the staff at the Department of Downstream Processing, CRAUN Research Sdn Bhd for their collaboration in the project. This project would not have had a smooth start if not for the support given by the staff here, from providing the starting material, assay kits, chemicals to constant supervision by the research officers. In particular, I would like to thank En Awang Zulfikar Rizal Bin Awang Seruji and Miss Siti Aminah Ibrahim, and all the lab assistants that has helped throughout the project.

Gratitude are also due to all the staff of Nutrition and Animal House Unit, Institute for Medical Research for their collaboration in the project and all the support given throughout my attachments. In particular, I would like to thank Dr Aswir Rashed, for guiding me in the *in vivo* animal study part of my project, from the acquiring of ethical approval up to necropsy. Also thanks for his assistance in the writing up of the journal paper.

Special thanks also to the veterinarian team of Dr Mohd Shafarin, Dr Azlina and Mdm Ezdiani for all their assistance, professional guidance, monitoring of the *in vivo* study, and for providing me a lot of opportunities not only to grow as a student, but also as a researcher of my own. Also sincerest thanks for the one-on-one training on the handling of laboratory test animal and the protocol of using the *in vivo* facility.

Sincerest gratitude also to all the staff under Animal House Unit, in particular, all the health medical assistant (PPK) for helping me in the physically tiring task, even on weekends.

I would also like to express my thanks to Prof Mohd Salleh Kamaruddin, for allowing me to use the facility in his lab for the preparation of the feed ingredients. Also special thanks to his lab assistant, Mr Salleh for his assistance in the preparation of the feed.

Gratitude are also due to all friends, both new and old. Thanks to all the seniors that has helped me to adapt to the research environment in UPM, and has offer their guidance when I was unsure. Thanks to Miss Phene and Miss Hui Hong for facilitating me with High Performance Liquid Chromatography, Miss Monica and Mr Maru Ali for helping me with the thesis procedure, and also others that I might have missed.

Sincerest gratitude to my friends, especially all the students under Dr Shahrul's supervision; Mdm Kathleen, Mdm Aminah, Miss Mayrylyn, Miss Dayang, Miss Nadiah and Morven, for making my study here less stressful and much more enjoyable. Thanks to Mdm Kathleen for sharing years of her experience as a graduate student and thus providing me with an elevated and a much forward platform to start my own graduate path. Also thanks for sharing with me your mistake to prevent me from doing the same. Thanks to all the "kakak" for making me stay grounded and be mature and to all the "adik" for the crazy, sudden moments.

Thanks also to my core-group of friends, the three musketeers + one dancing queen. Thanks for lending me your ears and shoulders when I need it the most. Thanks also for providing me with a new perspective when I seem too stuck with one. Thanks for being outright honest even when I didn't want to admit it.

Lastly, thanks to Research University Grant Scheme and Science Fund from the Ministry of Science and Innovation for funding this project. Also thanks to Ministry of Higher Education for the scholarship through the MyMaster program.



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Shahrul Razid Sarbini, PhD

Senior Lecturer

Faculty of Agricultural and Food Sciences

Universiti Putra Malaysia

(Chairman)

Patricia King Jie Hung, PhD

Senior Lecturer

Faculty of Agricultural and Food Sciences

Universiti Putra Malaysia

(Member)

BUJANG KIM HUAT, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 10 March 2016

REFERENCES

- Adebawale, K. O., Olu-Owolabi, B. I., Kehinde-Olawumi, E., & Lawal, O. S. (2005). Functional properties of native, physically and chemically modified breadfruit (*Artocarpus altilis*) starch. *Industrial Crops and Products*, 21(3), 343-351.
- Ahmad, F. B., & Williams, P. A. (1998). Rheological properties of sago starch. *Journal of Agricultural and Food Chemistry*, 46(10), 4060-4065.
- Ahmad, F. B., Williams, P. A., Doublier, J.-L., Durand, S., & Buleon, A. (1999). Physico-chemical characterisation of sago starch. *Carbohydrate Polymers*, 38(4), 361-370.
- Al-Tamimi, E. K., Seib, P. A., Snyder, B. S., & Haub, M. D. (2010). Consumption of cross-linked resistant starch (RS4 XL) on glucose and insulin responses in humans. *Journal of Nutrition and Metabolism*.
- Allsopp, P., Possemiers, S., Campbell, D., Ovarzabal, I. S., Gill, C., & Rowland, I. (2013). An exploratory study into the putative prebiotic activity of fructans isolated from *Agave angustifolia* and the associated anticancer activity. *Anaerobe*, 22, 38-44.
- Anderson, G. H., Cho, C. E., Akhavan, T., Mollard, R. C., Luhovyy, B. L., & Finocchiaro, E. T. (2010). Relation between estimates of cornstarch digestibility by the Englyst in vitro method and glycemic response, subjective appetite, and short-term food intake in young men. *The American Journal of Clinical Nutrition*, 91(4), 932-939.
- Anderson, G. H., & Woodend, D. (2003). Consumption of sugars and the regulation of short term satiety and food intake. *American Journal of Clinical Nutrition*, 78(4), 8435-8495.
- Annison, G., Illman, R. J., & Topping, D. L. (2003). Acetylated, propionylated or butyrylated starches raise large bowel short-chain fatty acids preferentially when fed to rats. *The Journal of Nutrition*, 133(11), 3523-3528.
- Arcila, J. A., & Rose, D. J. (2015). Repeated cooking and freezing of whole wheat flour increases resistant starch with beneficial impacts on in vitro fecal fermentation properties. *Journal of Functional Foods*, 12, 230-236.
- Ashogbon, A. O., & Akintayo, E. T. (2014). Recent trend in the physical and chemical modification of starches from different botanical sources: A review. *Starch-Stärke*, 66(1-2), 41-57.
- Ayucitra, A. (2012). Preparation and characterisation of acetylated corn starches. *International Journal of Chemical Engineering and Applications*, 3(3), 156-159.
- Awg-Adeni, D. S., Bujang, K., Hassan, M. A., & Abd-Aziz, S. (2012). Recovery of glucose from residual starch of sago hampas for bioethanol production. *BioMed Research International*.

- Bellisle, F., Drewnowski, A., Anderson, G. H., Westerterp-Plantenga, M., & Martin, C. K. (2012). Sweetness, satiation, and satiety. *The Journal of Nutrition*, 142(6), 1149S-1154S.
- Belobrajdic, D. P., King, R. A., Christophersen, C. T., & Bird, A. R. (2012). Dietary resistant starch dose-dependently reduces adiposity in obesity-prone and obesity resistant male rats. *Nutrition and Metabolism*, 9, 93.
- Ben, X. M., Li, J., Fend, Z. T., Shi, S. Y., Lu, Y. D., Chen, R., & Zhou, X. Y. (2008). Low level of galacto-oligosaccharide in infant formula stimulates growth of intestinal Bifidobacteria and Lactobacilli. *World Journal of Gastroenterology*, 14(42), 6564-6568.
- Benelam, B. (2009). Satiation, satiety and their effects on eating behaviour. *Nutrition Bulletin*, 34(2), 126-173.
- Bergman, E. (1990). Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiological Reviews*, 70(2), 567-590.
- Berry, C. (1986). Resistant starch: formation and measurement of starch that survives exhaustive digestion with amylolytic enzymes during the determination of dietary fibre. *Journal of Cereal Science*, 4(4), 301-314.
- Biliaderis, C. G. (1982). Physical characteristics, enzymic digestibility and structure of chemically modified smooth pea and waxy maize starches. *Journal of Agricultural and Food Chemistry*, 30(5), 925-930.
- Bindels, L. B., Porporato, P., Dewulf, E. M., Verrax, J., Neyrinck, A. M., Martin, J. C., . . . delzenne, N. M. (2012). Gut microbiota derived propionate reduces cancer cell proliferation in the liver. *British Journal of Cancer*, 107(8), 1337-1344.
- Birkett, A., Muir, J., Phillips, J., Jones, G., & O'Dea, K. (1996). Resistant starch lowers fecal concentrations of ammonia and phenols in human. *American Journal of Clinical Nutrition*, 137(2), 830-837.
- Björck, I., Gunnarsson, A., & Østergård, K. (1989). A study of native and chemically modified potato starch. Part II: Digestibility in the rat intestinal tract. *Starch-Stärke*, 41(4), 128-134.
- Bodinham, C. L., Frost, G. S., & Robertson, M. D. (2010). Acute ingestion of resistant starch reduces food intake in healthy adults. *British Journal of Nutrition*, 103(06), 917-922.
- Bodinham, C. L., Smith, L., Wright, J., Frost, G. S., & Robertson, M. D. (2012). Dietary fibre improves first-phase insulin secretion in overweight individuals. *PLoS ONE*, 7(7), e40834.
- Breuer, R. I., Buto, S. K., Christ, M. L., Bean, J., Vernia, P., Paoluzi, P., . . . Caprilli, R. (1991). Rectal irrigation with short-chain fatty acids for distal ulcerative colitis. *Digestive Diseases and Sciences*, 36(2), 185-187.

- Brouns, F., Kettlitz, B., & Arrigoni, E. (2002). Resistant starch and “the butyrate revolution”. *Trends in Food Science & Technology*, 13(8), 251-261.
- Brussow, H. (2013). Microbiota and healthy ageing: observational and nutritional intervention studies. *Microbial Biotechnology*, 6(4), 326-334.
- Bujang, K. (2008). Potentials of Bioenergy from the Sago Industries in Malaysia. *Biotechnology: Encyclopedia of Life Support Systems*.
- Bujang, K. (2014). *Sago: A food and fuel alternative*. Paper presented at the BIOBORNEO 2014:“Sustaining the Bioeconomy Community”.
- Cani, P. D., Holst, J. J., Drucker, D. J., Delzenne, N. M., Thorens, B., Burcelin, R., & Knauf, C. (2007). GLUT2 and the incretin receptors involved in glucose induced incretin secretion. *Molecular and Cell Endocrinology*, 278, 18-23.
- Case, S. E., Capitani, T., Whaley, J. K., Shi, Y. C., Trzasko, P., Jeffcoat, R., & Goldfarb, H. B. (1998). Physical properties and gelation behaviour of a low amylopectin maize starch and other high amylose maize starches. *Journal of Cereal Science*, 27, 301-314.
- Charalampopoulos, D., Wang, R., Pandiella, S. S., & Webb, C. (2002). Application of cereals and cereal components in functional foods: A review. *International Journal of Food Microbiology*, 79, 131-141.
- Charrier, J. A., Martin, R. J., McCutcheon, K. L., Raggio, A. M., Goldsmith, F., Goita, M., . . . Keenan, M. J. (2013). *High fat diet partially attenuates fermentation responses in rats fed resistant starch starch from high amylose maize*: Silver Spring.
- Clark, M. J., Robien, K., & Slavin, J. L. (2012). Effect of prebiotics on biomarkers of colorectal cancer in human: a systematic review. *Nutrition Reviews*, 70(8), 436-443.
- Clausen, M. R., & Mortensen, P. B. (1997). Lactulose, disaccharides and colonic flora. *Drugs*, 53(6), 930-942.
- Cummings, J., & Macfarlane, G. (1991). The control and consequences of bacterial fermentation in the human colon. *Journal of Applied Bacteriology*, 70(6), 443-459.
- Das, A. B., Singh, G., Singh, S., & Riar, C. S. (2010). Effect of acetylation and dual modification on physico-chemical, rheological and morphological characteristics of sweet potato (*Ipomoea batatas*) starch. *Carbohydrate Polymers*, 80(3), 725-732.
- Deckere, De., Kloots, W. J., & Van-Amelsvoort, J. (1993). Resistant starch decreases serum total cholesterol and triacylglycerol concentrations in rats. *The Journal of Nutrition*, 123(12), 2142.

Department of Statistics. (2012). *Metroxylon sagu plantation statistics, Sarawak*. Malaysia: Malaysia.

- Dewulf, E. M., Cani, P. D., Neyrinck, A. M., Possemiers, S., vanHolle, A., Muccioli, G. G., . . . Delzenne, N. M. (2011). Inulin-type fructans with prebiotic properties counteract GPR43 overexpression and PPAR γ -related adipogenesis in the white adipose tissue of high fat diet fed mice. *The Journal of Nutritional Biochemistry*, 22(8), 712-722.
- Duncan, S. H., Louis, P., & Flint, H. J. (2004). Lactate-utilizing bacteria, isolated from human feces, that produce butyrate as a major fermentation product. *Applied and Environmental Microbiology*, 70(10), 5810-5817.
- Eerlingen, R. C., Jacobs, H., & Delcour, J. A. (1994). Enzyme resistant starch. V. Effect of retrogradation of waxy maize starch on enzyme susceptibility. *Cereal Chemistry*, 71, 351-355.
- Enggum, B. O., Juliano, B. O., Perez, C. M., & Acedo, E. F. (1993). The Resistant Starch, Undigestible Energy and Undigestible Protein Contents of Raw and Cooked Milled Rice. *Journal of Cereal Science*, 18(2), 159-170.
- Englyst, H. N., Kingman, S., & Cummings, J. (1992). Classification and measurement of nutritionally important starch fractions. *European Journal of Clinical Nutrition*, 46, S33-50.
- Englyst, K., Liu, S., & Englyst, H. (2007). Nutritional characterization and measurement of dietary carbohydrates. *European Journal of Clinical Nutrition*, 61, S19-S39.
- Escarpa, A., Gonzalez, M. C., Morales, M. D., & Saura-Calixto, F. (1997). An approach to the influence of nutrients and other food constituents on resistant starch formation *Food Chemistry*, 60(4), 527-532.
- Everard, A., Lazarevic, V., Gaia, N., Johansson, M., Stahlman, M., Backhed, F., . . . Cani, P. D. (2014). Microbiome of prebiotic-treated mice reveals novel targets involved in host response during obesity. *ISME J.*
- Faisant, N., Champ, M., Colonna, P., & Buléon, A. (1993). Structural discrepancies in resistant starch obtained in vivo in humans and in vitro. *Carbohydrate Polymers*, 21(2), 205-209.
- Ferguson, L. R., Tasman-Jones, C., Englyst, H., & Harris, P. J. (2000). Comparative effects of three resistant starch preparations on transit time and short-chain fatty acid production in rats. *Nutrition and Cancer*, 36(2), 230-237.
- Flach, M. (1997). Sago palm: Metroxylon sagu Rottb. *Sago palm: Metroxylon Sagu Rottb.*
- Fleming, S., O'Donnell, A., & Perman, J. (1985). Influence of frequent and long-term bean consumption on colonic function and fermentation. *The American Journal of Clinical Nutrition*, 41(5), 909-918.

- Fuentes-Zaragoza, E., Sánchez-Zapata, E., Sendra, E., Sayas, E., Navarro, C., Fernández-López, J., & Pérez-Alvarez, J. A. (2011). Resistant starch as prebiotic: A review. *Starch-Stärke*, 63(7), 406-415.
- Galliard, T., & Bowler, P. (1987). *Morphology and Composition of Starch*. United Kingdom: Wiley, Chichester.
- Ganong, W. F., & Barrett, K. E. (2005). *Review of Medical Physiology* (Vol. 21): McGraw-Hill Medical ^ eNew York New York.
- Gibson, G. R., Probert, H. M., Loo, J. V., Rastall, R. A., & Roberfroid, M. B. (2004). Dietary modulation of the human colonic microbiota: updating the concept of prebiotics. *Nutrition Research Reviews*, 17(02), 259-275. doi: doi:10.1079/NRR200479.
- Gibson, G. R., & Roberfroid, M. B. (1994). Dietary Modulation of the Human Colonie Microbiota: Introducing the Concept of Prebiotics. *J Nutr*, 125(6), 1401-1412.
- Gibson, G. R., & Roberfroid, M. B. (1995). Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *The Journal of Nutrition*, 125(6), 1401-1412.
- Global Industry Analyst, G. (2010). The European Prebiotics Market to Reach \$1.17 Billion by 2015, According to New Report. Retrieved May 1, 2015, from PRWeb.
- Govindasamy, S., Oates, C., & Wong, H. (1992). Characterization of changes of sago starch components during hydrolysis by a thermostable alpha-amylase. *Carbohydrate Polymers*, 18(2), 89-100.
- Gray, G. M. (1975). Carbohydrate digestion and absorption: Role of the small intestine. *New England Journal of Medicine*, 292(23), 1225-1230.
- Haralampu, S. G. (2000). Resistant Starch : A review of the physical properties and biological impact of RS 3. *Carbohydrate Polymer*, 41, 285-292.
- Hashizume, K., Tsukahara, T., Yamada, K., Koyama, H., & Ushida, K. (2003). *Megasphaera elsdenii* JCM1772T normalizes hyperlactate production in the large intestine of fructooligosaccharide-fed rats by stimulating butyrate production. *The Journal of Nutrition*, 133(10), 3187-3190.
- Haska, N., & Ohta, Y. (1991). Glucose production from treated sago starch granules by raw starch digesting amylase from *Penicillium brunneum*. *Starch-Stärke*, 43(3), 102-107.
- Higgins, J. A. (2014). Resistant starch and energy balance: impact on weight loss and maintenance. *Critical Reviews in Food Science and Nutrition*, 54(9), 1158-1166.

- Hollis, J., Hutchison, C., & Hsu, W. (2014). Effect of resistant starch on subjective appetite and food intake in healthy adults. *FASEB Journal*, 28(1).
- Holmes, R. (1971). Carbohydrate digestion and absorption. *Journal of Clinical Pathology. Supplement (Royal College of Pathologists)*. 5, 10.
- Hoover, R., Hannouz, D., & Sosulski, F. (1988). Effects of Hydroxypropylation on Thermal Properties, Starch Digestibility and Freeze-Thaw Stability of Field Pea (*Pisum sativum* cv Trapper) Starch. *Starch-Stärke*, 40(10), 383-387.
- Höverstad, T., & Björneklett, A. (1984). Short-chain fatty acids and bowel functions in man. *Scandinavian Journal of Gastroenterology*, 19(8), 1059-1065.
- Huber, K. C., & BeMiller, J. N. (2001). Location of Sites of Reaction Within Starch Granules 1. *Cereal Chemistry*, 78(2), 173-180.
- Ingredients, M. (2011). Fibersym: Resistant wheat starch. In M. Ingredients (Ed.). Kansas: MGP Ingredients.
- Ingredion, I. (2014). Hi-Maize resistant starch-fibre, carbohydrate and more. In F. Innovation (Ed.). Singapore: Food Innovation.
- Jacobasch, G., Dongowski, G., Schmiedl, D., & Müller-Schmehl, K. (2006). Hydrothermal treatment of Novelose 330 results in high yield of resistant starch type 3 with beneficial prebiotic properties and decreased secondary bile acid formation in rats. *British Journal of Nutrition*, 95(06), 1063-1074.
- Jane, J. (2006). Current understanding on starch granule structures. *Journal of Applied Glycoscience*, 53(3), 205-213.
- Jane, J. L., & Robyt, J. F. (1984). Structure studies of amylose-V complexes and retrograded amylose by action of alpha amylases and a new method for preparing amyloextrins. *Carbohydrate Research*, 132, 105-118.
- Jiang, H., Campbell, M., Blanco, M., & Jane, J. I. (2010). Characterization of maize amylose extender (ae) mutant starches. Part II: Structures and properties of starch residues remaining after enzymatic hydrolysis at boiling water temperature. *Carbohydrate Polymer*, 80(1), 1-12.
- Johnston, K., Thomas, E., Bell, J., Frost, G., & Robertson, M. (2010). Resistant starch improves insulin sensitivity in metabolic syndrome. *Diabetic Medicine*, 27(4), 391-397.
- Kalmokoff, M., Zwicker, B., O'Hara, M., Matias, F., Green, J., Shastri, P., . . . Brooks, S. P. (2013). Temporal change in the gut community of rats fed with high amylose corn starch is driven by endogenous urea rather than strictly on carbohydrate availability. *Journal of Applied Microbiology*, 111(5), 1516-1528.
- Kapelko, M., Zięba, T., Golachowski, A., & Gryszkin, A. (2012). Effect of the production method on the properties of RS3/RS4 type resistant starch. Part 1:

- Properties of retrograded starch (RS3) produced under various conditions and its susceptibility to acetylation. *Food Chemistry*, 135(3), 1494-1504.
- Kapelko, M., Zięba, T., & Michalski, A. (2012). Effect of the production method on the properties of RS3/RS4 type resistant starch. Part 2. Effect of a degree of substitution on the selected properties of acetylated retrograded starch. *Food Chemistry*, 135(3), 2035-2042.
- Karim, A., Tie, A., Manan, D., & Zaidul, I. (2008). Starch from the sago (*Metroxylon sagu*) palm tree—properties, prospects, and challenges as a new industrial source for food and other uses. *Comprehensive Reviews in Food Science and Food Safety*, 7(3), 215-228.
- Karmakar, R., Ban, D., & Ghosh, U. (2013). Comparative study of native and modified starches isolated from conventional and nonconventional sources.
- Kaur, L., Singh, N., & Singh, J. (2004). Factors influencing the properties of hydroxypropylated potato starches. *Carbohydrate Polymers*, 55(2), 211-223.
- Kawabata, A., Sawayama, S., Nagashima, N., & Rosario, R. (1984). Physico-chemical properties of starches from cassava, arrowroot and sago. *Uritani, I.; Reyes, ED (eds.). Tropical Root Crops: Postharvest physiology and processing*.
- Kempermann, G., Kuhn, H. G., & Gage, F. H. (1997). Genetic influence on neurogenesis in the dentate gyrus of adult mice. *Proceedings of the National Academy of Sciences*, 94(19), 10409-10414.
- Kishida, T., Nakai, Y., & Ebihara, K. (2001). Hydroxypropyl-distarch phosphate from tapioca starch reduces zinc and iron absorption, but not calcium and magnesium absorption, in rats. *The Journal of Nutrition*, 131(2), 294-300.
- Kleessen, B., Stoof, G., Proll, J., Schmiedl, D., Noack, J., & Blaut, M. (1997). Feeding resistant starch affects fecal and cecal microflora and short-chain fatty acids in rats. *Journal of Animal Science*, 75(9), 2453-2462.
- Kohyama, K., & Nishikari, K. (1991). Effect of soluble sugars on gelatinization and retrogradation of sweet potato starch. *Journal of Agricultural and Food Chemistry*, 39, 1406-1410.
- Koo, S. H., Lee, K. Y., & Lee, H. G. (2010). Effect of cross-linking on the physicochemical and physiological properties of corn starch. *Food Hydrocolloids*, 24(6), 619-625.
- Lai, H. M., & Cheng, H. H. (2004). Properties of pregelatinized rice flour made by hot air or gum puffing. *International Journal of Food Science & Technology*, 39(2), 201-212. doi: 10.1046/j.0950-5423.2003.00761.
- Lajvardi, A., Mazarin, G. I., Gillespie, M. B., Satchithanandam, S., & Calvert, R. J. (1993). Starches of varied digestibilities differentially modify intestinal function in rats. *The Journal of Nutrition*, 123(12), 2059.

- Laparra, J. M., & Sanz, Y. (2010). Interactions of gut microbiota with functional food components and nutraceuticals. *Pharmacological Research*, 61(3), 219-225.
- Lee, K. Y., Yoo, S. H., & Lee, H. G. (2012). The effect of chemically-modified resistant starch, RS type-4, on body weight and blood lipid profiles of high fat diet-induced obese mice. *Starch-Stärke*, 64(1), 78-85.
- Lee, S. Y., Choi, W. S., & Lim, S. T. (2002). Paste texture of sago starch in comparison with other commercial starches. Paper presented at the International Symposium on Sago, Tokyo, Japan.
- Lehmann, U., & Robin, F. (2007). Slowly digestible starch—its structure and health implication: a review. *Trends in Food Science and Technology*, 18(7), 346-355.
- Leszczyński, W. a. (2004). Resistant starch—classification, structure, production. *Polish Journal of Food and Nutrition Sciences*, 13(54), 37-50.
- Leu, L., R. K., Hu, Y., Brown, I. L., & Young, G. P. (2009). Effect of high amylose maize starches on colonic fermentation and apoptotic response to DNA-damage in the colon of rats. *Nutritional Metabolism (London)*, 6, 11.
- Leszczyński, W. a. (2004). Resistant starch—classification, structure, production. *Polish Journal of Food and Nutrition Sciences*, 13(54), 37-50.
- Li, J. Y., & Yeh, A. I. (2001). Relationships between thermal, rheological characteristics and swelling power for various starches. *Journal of Food Engineering*, 50(3), 141-148. doi: [http://dx.doi.org/10.1016/S0260-8774\(00\)00236-3](http://dx.doi.org/10.1016/S0260-8774(00)00236-3).
- Li, L., Jiang, H., Campbell, M., Blanco, M., & Jane, J.-l. (2008). Characterization of maize amylose-extender mutant starches. Part I: Relationship between resistant starch contents and molecular structures. *Carbohydrate Polymers*, 74(3), 396-404.
- Liu, H., Ramsden, L., & Corke, H. (1999). Physical properties and enzymatic digestibility of hydroxypropylated ae, wx, and normal maize starch. *Carbohydrate Polymers*, 40(3), 175-182. doi: [http://dx.doi.org/10.1016/S0144-8617\(99\)00052-1](http://dx.doi.org/10.1016/S0144-8617(99)00052-1).
- Liu, Z., Lin, X., Huang, G., Zhang, W., Rao, P., & Ni, L. (2014). Prebiotic effects of almonds and almond skins on intestinal microbiota in healthy adult humans. *Anaerobe*, 26(0), 1-6.
- Livingstone, M. B. E., Robson, P. J., Welch, R. W., Burns, A. A., Burrows, M. S., & McCormack, C. (2000). Methodological issues in the assessment of satiety. *Scandinavian Journal of Nutrition/Näringsforskning*, 44(3), 98-103.
- Luoto, R., Ruuskanen, O., Waris, M., Kalliomaki, M., Salminen, S., & Isolauri, E. (2014). Prebiotic and probiotic supplementation prevents rhinovirus infections in preterm infants: a randomized, placebo-controlled trial. *Journal of Allergy Clinical Immunology*, 133(2), 403-413.

- Macfarlane, G. T., Steed, H., & Macfarlane, S. (2008). Bacterial metabolism and health-related effects of galacto-oligosaccharides and other prebiotics. *Journal of Applied Microbiology*, 104(2), 305-344. doi: 10.1111/j.1365-2672.2007.03520.
- Maki, K. C., Pelkman, C. L., Finocchiaro, E. T., Kelley, K. M., Lawless, A. L., Schild, A. L., & Rains, T. M. (2012). Resistant starch from high-amylase maize increases insulin sensitivity in overweight and obese men. *The Journal of Nutrition*, 142(4), 717-723.
- Maki, K. C., Pelkman, C. L., Finocchiaro, E. T., Kelley, K. M., Lawless, A. L., Schild, A. L., & Rains, T. M. (2012). Resistant starch from high amylose maize increases insulin sensitivity in overweight and obese man. *The Journal of Nutrition*, 142(4), 717-723.
- Marlett, J. A., & Longacre, M. J. (1996). Comparison of in vitro and in vivo measures of resistant starch in selected grain products. *Cereal Chemistry*, 73(1), 63-68.
- Matthuis, A. J., vanDenHeuvel, E. G., Schotermann, M. H., & Venema, K. (2012). Galacto-oligosaccharides have prebiotic activity in a dynamic in vitro colon model using a (13) C-labelling technique. *The Journal of Nutrition*, 142(7), 1205-1212.
- Monma, M., Yamamoto, Y., Kagei, N., & Kainuma, K. (1989). Raw Starch Digestion by α -Amylase and Glucoamylase from Chalara paradoxa. *Starch-Stärke*, 41(10), 382-385.
- Morrison, W. R. (1981). Starch lipids: a reappraisal. *Starke*, 33, 408-410.
- Morrison, W. R. (1995). Starch lipids and how they relate to starch granule structure and functionality. *Cereal Foods World*, 40, 437-438, 440-431, 443-436.
- Morrison, W. R. (2000). Starch lipids, starch granule structure and properties. *Special Publication - Royal Society of Chemistry*, 212, 261-270.
- Morrison, W. R., Tester, R. F., Snape, C. E., Law, R., & Gidley, M. J. (1993). Swelling and gelatinization of cereal starches. Some effects of lipid-complexed amylose and free amylose in waxy and normal barley starches. *Cereal Chemistry*, 70, 385-391.
- Muir, J. G., Walker, K. Z., Kaimakamis, M. A., Cameron, M. A., Govers, M., Lu, Z. X., ... O'Dea, K. (1998). Modulation of fecal markers relevant to colon cancer risk: a high-starch Chinese diet did not generate expected beneficial changes relative to a Western-type diet. *The American Journal of Clinical Nutrition*, 68(2), 372-379.
- Munjal, U., Scharlau, D., & Glei, M. (2012). Gut fermentation products of inulin-type fructans modulate the expression of xenobiotic-metabolising enzymes in human colonic tumour cells. *Anticancer Research*, 32(12), 5379-5386.

- Nakayama, T., & Oishi, K. (2013). Influence of coffee (*Coffea arabica*) and the galacto-oligosaccharide consumption of intestinal microbiota and the host responses. *FEMS Microbiology Letters*, 343(2), 161-168.
- Nayak, B., De J. Berrios, J., & Tang, J. (2014). Impact of food processing on the glycemic index (GI) of potato products. *Food Research International*, 56(0), 35-46.
- Nugent, A. P. (2005). Health properties of resistant starch. *British Nutrition Foundation, Nutrition Bulletin*, 30, 27-54.
- Orozco-Martínez, T., & Betancur-Ancona, D. (2004). Indigestible starch of *P. lunatus* obtained by pyroconversion: Changes in physicochemical properties. *Starch-Stärke*, 56(6), 241-247.
- Orzel, D., Bronkowska, M., & Styczynska, M. (2010). Effect of RS4 resistant starch in high-fat diets on magnesium and iron apparent absorption in Wistar rats-a short report. *Polish Journal of Food and Nutrition Sciences*, 60(1).
- Östergård, K., Björck, I., & Gunnarsson, A. (1988). A study of native and chemically modified potato starch. Part I: Analysis and enzymic availability in vitro. *Starch-Stärke*, 40(2), 58-66.
- Panesar, P. S., Kumari, S., & Panesar, R. (2013). Biotechnological approaches for the production of prebiotics and their potential applications. *Critical Reviews in Biotechnology*, 33(4), 345-364.
- Phillips, J. (1995). Effect of resistant starch on fecal bulk and fermentation dependent events in humans. *The American Journal of Clinical Nutrition*, 62(1), 121-130.
- Polesi, L. F., Sarmento, S., & Franco, C. M. (2011). Production and physicochemical properties of resistant starch from hydrolysed wrinkled pea starch. *International Journal of Food Science and Technology*, 46(11), 2257-2264.
- Pongjanta, J., Utaipatanacheep, A., Naivikul, O., & Piyachomkwan, K. (2008). Enzymes-resistant starch (RS III) from pullulanase-debranched high amylose rice starch. *Kasetsart Journal of Natural Science*, 42, 198-205.
- Raina, C., Singh, S., Bawa, A., & Saxena, D. (2006). Some characteristics of acetylated, cross-linked and dual modified Indian rice starches. *European Food Research and Technology*, 223(4), 561-570.
- Raben, A., Andersen, K., Karberg, M., Holst, J., & Astrup, A. (1997). Acetylation of or beta-cyclodextrin addition to potato beneficial effect on glucose metabolism and appetite sensations. *The American Journal of Clinical Nutrition*, 66(2), 304-314.
- Raben, A., Tagliabue, A., Christensen, N. J., Madsen, J., Holst, J. J., & Astrup, A. (1994). Resistant starch: the effect on postprandial glycemia, hormonal response, and satiety. *The American Journal of Clinical Nutrition*, 60(4), 544-551.

- Rauwerdink, J. B. (1986). An essay on Metroxylon, the sago palm. *Principes (USA)*.
- Reeves, P. G., Nielsen, F. H., & Fahey Jr, G. C. (1993). AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-76A rodent diet. *Journal of Nutrition*, 123(11), 1939-1951.
- Ring, S., Akingbala, J., & Rooney, L. (1989). A study of factors affecting amylose content of sorghum determined by an automated method. *Starch-Stärke*, 41(12), 457-461.
- Roberfroid, M., Hoyles, L., McCartney, A. L., Rastall, R. A., Rowland, I., Wolvers, D., . . . Meheust, A. (2010). Prebiotic concept: definition, metabolic and health benefits. *British Journal of Nutrition*, 104, 1-63.
- Robertson, M. D., Currie, J. M., Morgan, L. M., Jewel, D. P., & Frayn, K. N. (2003). Prior short term consumption of resistant starch enhances postprandial insulin sensitivity in healthy subjects. *Diabetologia*, 46(5), 659-665.
- Robinson, D., & Oddy, V. (2004). Genetic parameters for feed efficiency, fatness, muscle area and feeding behaviour of feedlot finished beef cattle. *Livestock Production Science*, 90(2), 255-270.
- Roediger, W., Vernia, P., & Breuer, R. (1989). Fecal anions and lactate in severe ulcerative colitis. *Digestive Diseases and Sciences*, 34(11), 1801-1802.
- Sajilata, M. G., Singhal, R. S., & Kulkarni, P. R. (2006). Resistant starch-A review. *Comprehensive Reviews in Food Science and Food Safety*, 5, 1-17.
- Sarawong, C., Schoenlechner, R., Sekiguchi, K., Berghofer, E., & Ng, P. K. W. (2014). Effect of extrusion cooking on the physicochemical properties, resistant starch, phenolic content and antioxidant capacities of green banana flour. *Food Chemistry*, 143(0), 33-39.
- Sarbini, S. R., Kolida, S., Gibson, G. R., & Rastall, R. A. (2013). *In vitro* fermentation of commercial α -gluco-oligosaccharide by faecal microbiota from lean and obese human subjects. *British Journal of Nutrition*, 109(11), 1980-1989. doi: doi:10.1017/S0007114512004205.
- Sarbini, S. R., & Rastall, R. A. (2011). Prebiotics: Metabolism, structure and function. *Functional Food Reviews*, 3(3), 93-106. doi: 10.2310/6180.2011.00004.
- Sasaki, T., & Matsuki, J. (1998). Effect of wheat starch structure on swelling power. *Cereal Chemistry Journal*, 75(4), 525-529. doi: 10.1094/cchem.1998.75.4.525.
- Scholz-Ahrens, K. E., Ade, P., Marten, B., Weber, P., Timm, W., Açıł, Y., . . . Schrezenmeir, J. (2007). Prebiotics, probiotics, and synbiotics affect mineral absorption, bone mineral content, and bone structure. *The Journal of Nutrition*, 137(3), 838S-846S.

- Scott, K. P., Martin, J. C., Duncan, S. H., & Flint, H. J. (2014). Prebiotic stimulation of human colonic butyrate-producing bacteria and bifidobacteria, *in vitro*. *FEMS Microbiology Ecology*, 87(1), 30-40. doi: 10.1111/1574-6941.12186.
- Segal, I., Hassan, H., Walker, A., Becker, P., & Braganza, J. (1995). Fecal short chain fatty acids in South African urban Africans and whites. *Diseases of The Colon & Rectum*, 38(7), 732-734.
- Shen, R.-L., Zhang, W.-L., Dong, J.-L., Ren, G.-X., & Chen, M. (2014). Sorghum resistant starch reduces adiposity in high-fat diet-induced overweight and obese rats via mechanisms involving adipokines and intestinal flora. *Food and Agricultural Immunology*, 1-11. doi: 10.1080/09540105.2013.876976.
- Shi, Y. C., Capitani, T., Trzasko, P., & Jeffcoat, R. (1998). Molecular structure of a low amylopectin starch and other high amylose maize starch. *Journal of Cereal Science*, 27, 289-299.
- Shukri, R., & Shi, Y.-C. (2015). Physicochemical properties of highly cross-linked maize starches and their enzymatic digestibilities by three analytical methods. *Journal of Cereal Science*, 63, 72-80. doi: <http://dx.doi.org/10.1016/j.jcs.2015.03.001>.
- Sievert, D., & Pomeranz, Y. (1989). Enzyme-resistant starch. I. Characterization and evaluation by enzymatic, thermoanalytical, and microscopic methods. *Cereal Chemistry Journal*, 66(4), 342-347.
- Siew-Wai, L., Zi-Ni, T., Karim, A. A., Hani, N. M., & Rosma, A. (2010). Fermentation of metroxylon sagu resistant starch type III by *Lactobacillus* sp. and *Bifidobacterium bifidum*. *Journal of Agricultural and Food Chemistry*, 58(4), 2274-2278.
- Silva, C. S., Bome, J. J. G. C. v. d., Gerrits, W. J. J., Kemp, B., & Bolhuis, J. E. (2012). Effects of dietary fibers with different physicochemical properties on feeding motivation in an adult female pigs. *Physiology and Behaviour*, 107(2), 218-230.
- Sim, S., Oates, C., & Wong, H. (1991). Studies on sago starch. Part I: characterization and comparison of sago starches obtained from Metroxylon Sagu processed at different times. *Starch-Stärke*, 43(12), 459-466.
- Singh, J., Dartois, A., & Kaur, L. (2010). Starch digestibility in food matrix: a review. *Trends in Food Science and Technology*, 21(4), 168-180.
- Singhal, R. S., Kennedy, J. F., Gopalakrishnan, S. M., Kaczmarek, A., Knill, C. J., & Akmar, P. F. (2008). Industrial production, processing, and utilization of sago palm-derived products. *Carbohydrate Polymers*, 72(1), 1-20.
- Slavin, J. (2004). Whole grains and human health. *Nutrition Research Reviews*, 17(99-110).
- Slavin, J. (2013). Fiber and prebiotics: mechanisms and health benefits. *Nutrients*, 5(4), 1417-1435.

- Soison, B., Jangchud, K., Jangchud, A., Harnsilawat, T., & Piyachomkwan, K. (2015). Characterization of starch in relation to flesh colors of sweet potato varieties. *International Food Research Journal*, 22(6).
- Spinner, J. (2013). Prebiotic markets to hit USD 4.8 billion by 2018. Retrieved May 1, 2015, from William Reed Business Media.
- Srichuwong, S., Sunarti, T. C., Mishima, T., Isono, N., & Hisamatsu, M. (2005a). Starches from different botanical sources I: Contribution of amylopectin fine structure to thermal properties and enzyme digestibility. *Carbohydrate Polymers*, 60(4), 529-538.
- Srichuwong, S., Sunarti, T. C., Mishima, T., Isono, N., & Hisamatsu, M. (2005b). Starches from different botanical sources II: Contribution of starch structure to swelling and pasting properties. *Carbohydrate Polymers*, 62(1), 25-34.
- Stanton, W. R. (1993). Perspectives on, and future prospects for, the sago palm. . *Sago Palm*, 1, 6.
- Tachon, S., Zhou, J., Keenan, M., Martin, M., & Marco, M. L. (2013). The intestinal microbiota in aged mice is modulated by dietary resistant starch and correlated with improvements in host responses. *FEMS Microbial Ecology*, 83(2), 299-309.
- Takahashi, H., Ik Yang, S., Hayashi, C., Kim, M., Yamanaka, J., & Yamamoto, T. (1993). Effect of partially hydrolyzed guar gum on fecal output in human volunteers. *Nutrition research*, 13(6), 649-657.
- Takeda, Y., Takeda, C., Suzuki, A., & Hizukuri, S. (1989). Structures and properties of sago starches with low and high viscosities on amylography. *Journal of Food Science*, 54(1), 177-182.
- Tester, R. F., & Morrison, W. R. (1990a). Swelling and gelatinization of cereal starches. I. Effects of amylopectin, amylose, and lipids. *Cereal Chemistry Journal*, 67(6), 551-557.
- Tester, R. F., & Morrison, W. R. (1990b). Swelling and gelatinization of cereal starches. I. Effects of amylopectin, amylose, and lipids. *Cereal Chemistry Journal*, 67(6), 551-557.
- Tester, R. F., & Karkalas, J. (1996). Swelling and gelatinization of oat starches. *Cereal Chemistry Journal*, 73(2), 271-277.
- Tester, R. F., Qi, X., & Karkalas, J. (2006). Hydrolysis of native starches with amylases. *Animal Feed Science and Technology*, 130, 39-54.
- Thanh-Blicharz, L., J., Anioła, J., Kowalczewski, P., Przygoński, K., Zaborowska, Z., & Lewandowicz, G. (2014). Type IV resistant starch increases cecum short chain fatty acids level in rats. *Acta Biochimica Polonica*, 61(1), 109-114.

- Thompson, L. U., Maningat, C. C., Woo, K., & Seib, P. A. (2011). In vitro digestion of RS4-type resistant wheat and potato starches, and fermentation of indigestible fractions. *Cereal Chemistry*, 88(1), 72-79.
- Thorbecke, G. (2013). *Biology of Aging and Development* (Vol. 3): Springer Science & Business Media.
- Tie, A. P.-L. (2004). *Physico-chemical properties of sago starch in sago palm (*Metroxylon sagu*) at different growth stages.* (Master of Science), University of Science Malaysia.
- Tie, A. P. L., Karim, A. A., & Manan, D. M. A. (2008). Physicochemical properties of starch in sago palms (*Metroxylon sagu*) at different growth stages. *Starch-Stärke*, 60(8), 408-416.
- Toden, S. (2006). Resistant starch prevents colonic DNA damage induced by high dietary cooked red meat or casein in rats. *Cancer Biology and Therapy*, 5(3), 267-272.
- Topping, D. L., & Clifton, P. M. (2001). Short-chain fatty acids and human colonic function: roles of resistant starch and nonstarch polysaccharides. *Physiological reviews*, 81(3), 1031-1064.
- Uggah, M. D. S. D. (2013). Jabu: More focus on sago industry in 11th Malaysia Plan, *The Star*.
- Uthumporn, U., Wahidah, N., & Karim, A. (2014). *Physicochemical Properties Of Starch From Sago (*Metroxylon Sagu*) Palm Grown In Mineral Soil At Different Growth Stages.* Paper presented at the IOP Conference Series: Materials Science and Engineering.
- Waliszewski, K. N., Aparicio, M. A., Bello, L. s. A., & Monroy, J. A. (2003). Changes of banana starch by chemical and physical modification. *Carbohydrate Polymers*, 52(3), 237-242.
- Wang, W., Powell, A., & Oates, C. (1995). Pattern of enzyme hydrolysis in raw sago starch: effects of processing history. *Carbohydrate Polymers*, 26(2), 91-97.
- Wattanachant, S., Muhammad, S. K. S., Hashim, D. M., & Rahman, R. A. (2002). Characterisation of hydroxypropylated crosslinked sago starch as compared to commercial modified starches. *Songklanakarin Journal of Science and Technology*, 24(3), 439-450.
- Wootton, M., & Manatsathit, A. (1984). The influence of molar substitution on the gelatinization of hydroxypropyl maize starches. *Starch - Stärke*, 36(6), 207-208. doi: 10.1002/star.19840360605.
- Woo, K., & Seib, P. A. (1997). Cross-linking of wheat starch and hydroxypropylated wheat starch in alkaline slurry with sodium trimetaphosphate. *Carbohydrate Polymers*, 33(4), 263-271.

- Wootton, M., & Manatsathit, A. (1984). The influence of molar substitution on the gelatinization of hydroxypropyl maize starches. *Starch-Stärke*, 36(6), 207-208.
- Wronkowska, M., Juśkiewicz, J., Zduńczyk, Z., Soral-Śmietana, M., & Krupa-Kozak, U. (2011). Influence of chemically-modified potato starch (RS type 4) on the nutritional and physiological indices of rats. *Polish Journal of Food and Nutrition Sciences*, 61(2), 143-151.
- Yousefi, A., Razavi, S. M., & Norouzy, A. (2015). In vitro gastrointestinal digestibility of native, hydroxypropylated and cross-linked wheat starches. *Food & Function*, 6(9), 3126-3134.
- Zaman, S. A., & Sarbini, S. R. (2015). The potential of resistant starch as a prebiotic. *Critical Reviews in Biotechnology*(0), 1-7.
- Zhang, H., & Jin, Z. (2011). Preparation of resistant starch by hydrolysis of maize starch with pullulanase. *Carbohydrate Polymers*, 83(2), 865-867.
- Zhu, & Zhao. (2013). In vitro fermentation of a retrograded maize starch by healthy adult fecal extract and impacts of exogenous microorganisms on three acids production. *Starke*, 65(3/4), 330-337.
- Zięba, T., Kapelko, M., & Szumny, A. (2013). Effect of preparation method on the properties of potato starch acetates with an equal degree of substitution. *Carbohydrate Polymers*, 94(1), 193-198.