



**UNIVERSITI PUTRA MALAYSIA**

**DIETARY ENERGY AND PROTEIN REQUIREMENTS OF  
LESSER MOUSE DEER (TRAGULUS JA VANICUS)  
IN CAPTIVITY**

**DARLIS**

**IB 2001 1**

**DIETARY ENERGY AND PROTEIN REQUIREMENTS OF  
LESSER MOUSE DEER (*TRAGULUS JAVANICUS*)  
IN CAPTIVITY**

By

**D A R L I S**

**Thesis Submitted in Fulfilment on the Requirement for the Degree of  
Doctor of Philosophy in the Institute of Bioscience  
Universiti Putra Malaysia**

**November 2001**



Abstract of the thesis submitted to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**DIETARY ENERGY AND PROTEIN REQUIREMENTS OF LESSER MOUSE DEER (*TRAGULUS JAVANICUS*) IN CAPTIVITY**

By

**D A R L I S**

**November 2001**

**Chairman : Associate Professor Dr. Norhani Abdullah**  
**Faculty : Institute of Bioscience**

A series of experiments was conducted to study various aspects of nutrition of lesser mouse deer to determine their dietary energy and protein requirements for maintenance and growth in captivity. When given a choice of feeds male mouse deer preferred sweet potato, while the female preferred carrot in terms of fresh weight. However, in terms of dry matter (DM) intake, the rabbit pellet (90% DM) was consumed the most by both male and female mouse deer. The daily intakes of DM, organic matter (OM) and gross energy (GE) were significantly ( $P < 0.05$ ) higher in male than female when given these feeds, but the DM intake was similar when they were fed only rabbit pellet. The crude protein (CP) intake was similar between male and female mouse deer when fed various types of feed or only pellet. The ability of male and female mouse deer to digest DM, OM and GE of feed were also similar.

In the energy expenditure and activity study, it was observed that both male and female mouse deer spent most of their time sitting (926.0 min/d), followed by standing (491.5 min/d) and eating (22.5 min/d). Heat production (HP) for male and



female (0.44 and 0.43 kJ/kg  $W^{0.75}$ /min, respectively) was the highest during eating followed by standing (0.37 and 0.33 kJ/kg  $W^{0.75}$ /min, respectively) and sitting (0.26 and 0.26 kJ/kg  $W^{0.75}$ /min, respectively). The HP for morning to afternoon and afternoon to evening periods were significantly higher than the night and early morning periods. The overall HP of males during morning to afternoon and afternoon to evening periods was significantly ( $P < 0.05$ ) higher than that of females.

When the mouse deer were fed diets containing 5% to 20% of fiber, the digestibility values of DM, OM, CP and GE were significantly ( $P < 0.05$ ) decreased with an increase in fiber level. Positive nitrogen balance and body weight gain was obtained when the levels of fiber were in the range of 5% – 10%. The passage rate constants of solid and fluid were significantly increased when more than 10% fiber was in the diet. On the other hand, the mean retention time of solid and fluid decreased with increasing levels of fiber in the diet.

Metabolisable energy requirements for maintenance in growing and mature mouse deer were in the range of 418.0 kJ/kg  $W^{0.75}$ /d – 424.9 kJ/kg  $W^{0.75}$ /d. Efficiency of utilisation of energy requirement for maintenance was 0.52 for growing mouse deer and 0.44 for mature mouse deer. Maximum nitrogen balance was achieved at 20% metabolisable energy above maintenance. The protein requirement for maintenance of growing male, mature male and mature female was 0.33, 0.25 and 0.28 g N/kg  $W^{0.75}$ /d or 6.4%, 5.4% and 5.3% CP in the diet, respectively and for growth was 0.93, 0.94 and 0.98 g N/kg  $W^{0.75}$ /d or 18.2%, 18.8% and 18.7% CP in the diet, respectively.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan ijazah Doktor Falsafah

**KEPERLUAN TENAGA DAN PROTEIN PELANDUK DALAM  
PELIHARAAN**

**Oleh**

**D A R L I S**

**November 2001**

**Pengerusi : Profesor Madya Dr. Norhani Abdullah**  
**Fakulti : Institut Biosains**

Satu siri eksperimen telah dijalankan untuk mengkaji berbagai aspek pemakanan pelanduk untuk menentukan keperluan tenaga dan protein keperluan hidup asas dan tumbesaran dalam peliharaan. Apabila pelanduk diberi berbagai jenis makanan segar, pelanduk jantan lebih suka ubi kentang sedangkan pelanduk betina lebih suka lobak merah, tetapi dalam bentuk bahan kering (BK), pelanduk jantan dan betina lebih banyak memakan pelet arnab (90% BK). Pengambilan harian BK, bahan organik (BO) dan tenaga (T) pelanduk jantan lebih dari pelanduk betina bila diberi berbagai jenis makanan, tetapi bila hanya diberi pelet arnab, jumlah pengambilan BK pelanduk jantan adalah sama dengan pelanduk betina. Pengambilan protein kasar (PK) pelanduk jantan dan betina adalah sama bila diberi berbagai jenis makanan atau diberi pelet sahaja. Penghadaman BK, BO dan T pelanduk jantan dan pelanduk betina adalah sama.

Dalam kajian aktiviti dan penggunaan tenaga, pelanduk jantan dan betina menghabiskan masa untuk duduk selama 926 min/hari, diikuti dengan berdiri selama



491 min/hari dan makan selama 22.5 min/hari. Penghasilan haba pelanduk jantan dan betina adalah paling tinggi semasa makan (0.44 dan 0.43 kJ/kg  $W^{0.75}$ /min), diikuti semasa berdiri (0.37 dan 0.33 kJ/kg  $W^{0.75}$ /min) dan duduk (0.26 and 0.26 kJ/kg  $W^{0.75}$ /min). Penghasilan haba dari pagi hingga tengahari dan dari tengahari hingga petang adalah lebih tinggi dari malam hingga awal pagi. Penghasilan haba secara keseluruhan pelanduk jantan dari pagi hingga tengahari dan dari tengahari hingga petang adalah lebih tinggi daripada pelanduk betina.

Apabila diberi makanan yang mengandungi 5% - 20% serat kasar, nilai penghadaman BK, BO, PK dan T menurun dengan meningkatnya kandungan serat kasar. Keseimbangan nitrogen dan pertambahan berat badan adalah positif bila kandungan serat kasar dalam lingkungan 5% - 10%. Pengaliran butir-butir kecil digesta dan cecair rumen lebih cepat apabila kandungan serat kasar melebihi 10%. Waktu retensi butir-butir kecil digesta dan cecair rumen menurun dengan meningkatnya kandungan serat kasar.

Keperluan asas tenaga metabolik pelanduk yang membesar dan dewasa adalah 418.0 – 424.9 kJ/kg  $W^{0.75}$ /hari. Kecekapan penggunaan tenaga untuk keperluan asas adalah 0.52 bagi pelanduk yang membesar dan 0.44 bagi pelanduk dewasa. Keseimbangan nitrogen yang maksimum diperolehi bila tenaga metabolik makanan melebihi 20% keperluan asas. Keperluan asas protein untuk pelanduk yang sedang membesar, jantan dewasa dan betina adalah 0.33, 0.25 and 0.28 g N/kg  $W^{0.75}$ /hari atau 6.4%, 5.4% dan 5.3% PK makanan dan untuk pertumbuhan adalah 0.93, 0.94 and 0.98 g N/kg  $W^{0.75}$ /hari atau 18.2%, 18.8% and 18.7% PK makanan.

## ACKNOWLEDGMENTS

I wish to express my sincere appreciation and gratitude to Assoc. Prof. Dr. Norhani Abdullah, chairman of the supervisory committee for her invaluable guidance and untiring assistance throughout the period of my studies. Sincere thanks are also due to Professor. Dr. Ho Yin Wan and Assoc. Prof. Dr. Liang Juan Boo for serving as members of the supervisory committee and for their invaluable advice, suggestion and comments during the course of my work and in the preparation of the thesis.

I would like to thank the Malaysian government for the financial assistance given to me during my study in UPM, the University of Jambi for granting permission for leave of absence to undertake this study, Institute of Bioscience and the Department of Animal Science, Faculty of Agriculture, Universiti Putra Malaysia for providing the research facilities.

I am also grateful to Professor Tan Sri Dr. Syed Jalaludin Syed Salim for his invaluable advice and encouragement, Professor Dr. Dahlan Ismail for allowing me to use a few of his mouse deer, Dr. Bagus Purwanto of the Animal Science Department, Institut Pertanian Bogor, Indonesia for his advice and suggestions in the energy expenditure study and the Department of Wildlife and National Parks of



Malaysia (PERHILITAN) for approval to maintain the mouse deer colony at Universiti Putra Malaysia.

Sincere appreciation is extended to all my colleagues and staff of Digestive Microbiology Laboratory, Latiffah Zakaria, Kalavathy Ramasamy, Lan Gan Qiu, Sieo Chin Chin, Thongsuk Jetana, Vickneswary Dharmadas, Wan Zuhainis, Sideig Adam Abdalla, Khairul Kamar Bakri, Madam Haw Ah Kam, Jivanathan Arumugam, Nagaya Muniandy and Paimon Lugiman for their cooperation and assistance. Sincere appreciation is also extended to all the staff in the Nutrition Laboratory, Ibrahim Mohsin, Bakari Abd. Rahman and Saparin Denim, for their help.

Finally, I would like to express my sincere gratitude and affection to my loving wife Dra. Mardiani for her love, prayers, patience and encouragement throughout the duration of my study and, above all, to Almighty Allah S.W.T., for blessings on me and my family.





## TABLE OF CONTENTS

	<b>Page</b>
ABSTRACT .....	ii
ABSTRAK .....	iv
ACKNOWLEDGEMENTS .....	vi
APPROVAL SHEETS .....	viii
DECLARATION FORM .....	x
LIST OF TABLES .....	xv
LIST OF FIGURES .....	xx
 <b>CHAPTER</b>	
<b>1 INTRODUCTION .....</b>	<b>1</b>
<b>2 LITERATURE REVIEW .....</b>	<b>4</b>
2.1. Mouse Deer .....	4
2.1.1. Classification .....	4
2.1.2. Morphological Characteristics.....	4
2.1.3 Feed Intake Studies in Mouse deer .....	8
2.1.4. Rumen Microorganisms .....	8
2.1.5. Reproduction .....	9
2.2. Feed Intake and Factors Affecting Feed Intake .....	9
2.3. Diet Selection .....	12
2.4. Feed Preference .....	16
2.5. Digestion in Ruminant Animals.....	18
2.6. Passage Rate of Rumen Digesta .....	19
2.7. Energy Requirement and Utilisation .....	22
2.7.1. Energy Available to Animals .....	22
2.7.2. Heat Production .....	24
2.7.3. Energy Requirement for Maintenance .....	26
2.7.4. Utilisation of Energy .....	28
2.7.5. Efficiency of Energy Utilisation .....	30
2.8. Protein Requirement and Utilisation .....	30
2.8.1. Protein Digestion .....	30
2.8.2. Protein Utilisation .....	32
2.8.3. Protein Requirement .....	33
<b>3 PREFERENCE TEST ON FEED AND NUTRIENT INTAKES IN MALE AND FEMALE LESSER MOUSE DEER .....</b>	<b>35</b>
3.1. Introduction .....	35
3.2. Materials and Methods .....	36
3.2.1. Experiment I : Feed Preference and Nutrient Intake .....	36



3.2.1.1.	Animals .....	36
3.2.1.2.	Feeds and Experimental Procedures .....	37
3.2.1.3.	Chemical Analyses .....	38
3.2.1.4.	Statistical Analysis .....	45
3.2.2.	Experiment II : Selection of Feeds with Various Levels of Protein Offered Singly or in Pairs.....	45
3.2.2.1.	Animals .....	45
3.2.2.2.	Feeding .....	46
3.2.2.3.	Single Feed Offer .....	46
3.2.2.4.	Paired-Feed Offer .....	48
3.2.2.5.	Sample Analyses .....	49
3.2.2.6.	Statistical Analysis .....	49
3.3.	Results .....	49
3.3.1.	Feed Preference and Nutrient Intake .....	49
3.3.2.	Selection of Feed with Various Levels of Protein .....	55
3.3.2.1.	Single Feed Offered .....	55
3.3.2.2.	Paired-Feed Offered .....	57
3.4.	Discussion .....	60
<b>4</b>	<b>ENERGY EXPENDITURE IN RELATION TO ACTIVITY OF LESSER MOUSE DEER .....</b>	<b>68</b>
4.1.	Introduction .....	68
4.2.	Materials and Methods .....	68
4.2.1.	Animals and Feed .....	68
4.2.2.	Measurement of Energy Expenditure .....	69
4.2.3.	Statistical Analysis .....	72
4.3.	Results .....	72
4.4.	Discussion .....	80
<b>5</b>	<b>EFFECTS OF FIBRE ON DIGESTIBILITY, NITROGEN BALANCE, FLOW RATE AND HEAT PRODUCTION IN LESSER MOUSE DEER .....</b>	<b>85</b>
5.1.	Introduction .....	85
5.2.	Materials and Methods .....	86
5.2.1.	Animals and Feeds .....	86
5.2.2.	Digestibility Trial .....	88
5.2.3.	Solid and Fluid Flow Rate .....	88
5.2.3.1.	Preparation of Markers .....	89
5.2.3.2.	Sampling .....	90
5.2.3.3.	Sample Analysis .....	90
5.2.3.4.	Calculations .....	91
5.2.4.	Heat Production .....	92
5.2.5.	Statistical Analysis .....	92
5.3.	Results .....	93



5.3.1.	Feed Intake .....	93
5.3.2.	Faecal Output .....	95
5.3.3.	Digestibility .....	95
5.3.4.	Digestible Nutrient Intake .....	99
5.3.5.	Nitrogen Balance .....	101
5.3.6.	Solid Flow Rate .....	103
5.3.7.	Fluid Flow Rate .....	108
5.3.8.	Heat Production .....	112
5.4.	Discussion .....	113
<b>6</b>	<b>ENERGY REQUIREMENT FOR MAINTENANCE OF GROWING AND MATURE MALE LESSER MOUSE DEER ....</b>	<b>119</b>
6.1.	Introduction.....	119
6.2.	Materials and Methods .	120
6.2.1.	Animals .....	120
6.2.2.	Feeds and Feeding Levels .....	120
6.2.3.	Experimental Procedures .....	122
6.2.4.	Sample Analyses .....	123
6.2.5.	Statistical Analysis .....	124
6.3.	Results .....	124
6.4.	Discussion .....	139
<b>7</b>	<b>EFFECTS OF ENERGY AND PROTEIN LEVELS ON FEED INTAKE, NITROGEN BALANCE, NITROGEN REQUIREMENT AND FEED EFFICIENCY OF LESSER MOUSE DEER .....</b>	<b>144</b>
7.1.	Introduction .....	144
7.2.	Materials and Methods .....	145
7.2.1.	Animals .....	145
7.2.2.	Feed .....	145
7.2.3.	Experimental Procedures .....	145
7.2.4.	Statistical Analysis .....	149
7.3.	Results .....	150
7.3.1.	Nutrient Intake .....	150
7.3.2.	Nutrient Output .....	152
7.3.3.	Nutrient Digestibility .....	154
7.3.4.	Digestible Nutrient Intake .....	157
7.3.5.	Nitrogen Balance .....	159
7.3.6.	Efficiency of Feed Utilisation .....	164
7.4.	Discussion .....	169
<b>8</b>	<b>PROTEIN REQUIREMENTS FOR MAINTENANCE AND GROWTH IN GROWING MALE, MATURE MALE AND MATURE FEMALE LESSER MOUSE DEER .....</b>	<b>175</b>
8.1.	Introduction .....	175
8.2.	Materials and Meth	176



8.2.1. Animals .....	176
8.2.2. Feed .....	176
8.2.3. Experimental Procedures .....	176
8.2.4. Endogenous Urinary Nitrogen (EUN) and Metabolic Faecal Nitrogen (MFN) .....	178
8.2.5. Statistical Analysis .....	179
8.3. Results .....	179
8.3.1. Nutrient Intake .....	179
8.3.2. Nutrient Output .....	181
8.3.3. Nutrient Digestibility .....	183
8.3.4. Digestible Nutrient Intake .....	183
8.3.5. Nitrogen Balance .....	186
8.3.6. Efficiency of Feed Utilisation .....	191
8.3.7. Endogenous Urinary Nitrogen and Metabolic Faecal Nitrogen .....	195
8.4. Discussion .....	199
<b>9 GENERAL DISCUSSION AND CONCLUSION .....</b>	<b>203</b>
<b>BIBLIOGRAPHY .....</b>	<b>212</b>
<b>APPENDIX .....</b>	<b>232</b>
<b>VITA .....</b>	<b>233</b>



## LIST OF TABLES

Table	Page
1. Taxonomic relationship of <i>Tragulus javanicus</i> .....	5
2. Mean retention times of digesta in the gut of different animal species.	21
3. Heat production of different species and sizes of animals .....	26
4. Proportion (g/kg) of water, fat, CP and GE contents (MJ/kg) in empty body of lambs at different weights .....	29
5. Ingredient composition of basal feeds L and H (dry weight basis) ....	47
6. Chemical composition of experimental feeds .....	47
7. The chemical composition of feeds offered .....	50
8. Fresh feeds consumed by male and female lesser mouse deer .....	51
9. Dry matter intake (DMI) of feed and their percentage proportions consumed by male and female lesser mouse deer .....	52
10. Feed intake, digestibility, nitrogen balance and body weight gain of male and female lesser mouse deer .....	54
11. Dry matter intake (DMI), crude protein intake (CPI), body weight gain and feed conversion efficiency (FCE) of male and female mouse deer offered single feeds .....	56
12. Feed H intake (FHI), dry matter intake (DMI), crude protein intake (CPI), body weight gain and feed conversion efficiency (FCE) of male and female mouse deer offered paired-feeds .....	58
13. Feed H intake by male and female mouse deer when feeds were placed in different positions .....	59
14. Dry matter (DM), organic matter (OM), crude protein (CP) and gross energy (GE) intakes of male and female mouse deer in the respiration chamber .....	74



15.	Duration (minute) of different activities of male and female mouse deer in the respiration chamber (RC) and in an outdoor enclosure with a small shelter (OD) .....	75
16.	Heat production of mouse deer for different activities in the respiration chamber .....	76
17.	Heat production (kJ/kg W <sup>0.75</sup> ) of mouse deer at different periods in the respiration chamber within 24 h .....	79
18.	Ingredient composition of experimental diets (dry matter basis) .....	87
19.	Chemical composition of experimental diets .....	87
20.	Dry matter (DM), crude protein (CP), organic matter (OM), crude fibre (CF), neutral detergent fibre (ADF), acid detergent fibre (ADF) and gross energy (GE) intakes of mouse deer fed different levels of fibre .....	94
21.	Faecal dry matter (DM), crude protein (CP), organic matter (OM), crude fibre (CF), neutral detergent fibre (ADF), acid detergent fibre (ADF) and energy (E) of mouse deer fed different levels of fibre .....	96
22.	Digestibilities of dry matter (DM), crude protein (CP), organic matter (OM), crude fibre (CF), neutral detergent fibre (ADF), acid detergent fibre (ADF) and gross energy (GE) in mouse deer fed different levels of fibre .....	97
23.	Digestible dry matter (DM), crude protein (CP), organic matter (OM), crude fibre (CF), neutral detergent fibre (ADF), acid detergent fibre (ADF) and energy (E) intakes of mouse deer fed different levels of fibre .....	100
24.	Nitrogen intake (NI), faecal nitrogen (FN), urinary nitrogen (UN), nitrogen balance (NB) and body weight gain (Gain) of mouse deer fed different levels of fibre .....	102
25.	Passage rate constant of particulate matter ( <i>k</i> ), mean retention time (MRT), pool size (PS) and solid flow rate (FR) of mouse deer fed different levels of fibre .....	104
26.	Water intake (WI), dilution rate ( <i>k</i> ), mean retention time (MRT), rumen volume (RV) and fluid flow rate (FR) of mouse deer fed different levels of fibre .....	109



27.	Heat production of mouse deer fed different levels of fibre .....	112
28.	The ingredient composition of pellets (dry matter basis) .....	121
29.	Chemical composition of pellets .....	121
30.	Dry matter (DM), organic matter (OM) and gross energy (GE) intakes of growing and mature male mouse deer fed different levels of feed .....	125
31.	Faecal dry matter (DM), organic matter (OM) and energy (E), urinary energy (UE) and methane energy (M) of growing and mature male mouse deer fed different levels of feed .....	127
32.	Digestibility of dry matter (DM), organic matter (OM) and gross energy (GE) in growing and mature male mouse deer fed different levels of feed .....	129
33.	Digestible dry matter (DM), organic matter (OM) and energy (E) intakes of growing and mature male mouse deer fed different levels of feed .....	131
34.	Metabolisable energy intake (MEI), heat production (HP), energy retention (ER) and body weight gain (gain) of growing and mature male mouse deer fed different levels of feed, calculated using equation I (MEI = 0.82 DE) .....	132
35.	Metabolisable energy intake (MEI), heat production (HP), energy retention (ER) of growing and mature male mouse deer fed different levels of feed, calculated using equation II [MEI = GEI - (FE + UE + M)] .....	133
36.	Linear relationship between metabolisable energy intake (MEI) and energy retention (ER) in mouse deer fed different levels of feed .....	138
37.	Linear relationship between body weight change (BW) and metabolisable energy intake (MEI) in mouse deer fed different levels of feed .....	138
38.	Ingredient composition of pellets .....	146
39.	Chemical composition of pellets .....	147



40. Dry matter (DM), organic matter (OM), crude protein (CP) and gross energy (GE) intakes of mouse deer fed different levels of energy and protein .....	151
41. Faecal dry matter (DM), organic matter (OM), crude protein (CP) and energy (GE) of mouse deer fed different levels of energy and protein .....	153
42. Digestibility of dry matter (DM), organic matter (OM), crude protein (CP) and gross energy (GE) in mouse deer fed different levels of energy and protein .....	155
43. Digestible dry matter (DM), organic matter (OM), crude protein (CP) and energy (E) intakes of mouse deer fed different levels of energy and protein .....	158
44. Nitrogen intake (NI), faecal nitrogen (FN), urinary nitrogen (UN) and nitrogen balance (NB) of mouse deer fed different levels of energy and protein .....	161
45. Linear relationships between nitrogen intake (NI) and nitrogen balance (NB) of mouse deer fed different levels of energy and protein .....	164
46. Energy protein ratio (EP), feed conversion ratio (CFR), feed conversion efficiency (FCE) and body weight gain (Gain) of mouse deer fed different levels of energy and protein .....	165
47. Dry matter (DM), organic matter (OM), crude protein (CP) and gross energy (GE) intakes of growing male (GM), mature male (MM) and mature female (MF) mouse deer fed different levels of protein .....	180
48. Faecal dry matter (DM), organic matter (OM), crude protein (CP) and energy (E) of growing male (GM), mature male (MM) and mature female (MF) mouse deer fed different levels of protein .....	182
49. Digestibility of dry matter (DM), organic matter (OM), crude protein (CP) and gross energy (GE) in growing male (GM), mature male (MM) and mature female (MF) mouse deer fed different levels of protein .....	184
50. Digestible dry matter (DM), organic matter (OM), crude protein (CP) and energy (E) intakes of growing male (GM), mature male (MM) and mature female (MF) mouse deer fed different levels of protein ...	185





51.	Nitrogen intake (NI), faecal nitrogen (FN) urinary nitrogen (UN) and nitrogen balance (NB) of growing male (GM), mature male (MM) and mature female (MF) mouse deer fed different levels of protein ...	187
52.	Linear relationships between nitrogen intake (NI) and nitrogen balance (NB) of growing male, mature male and mature female mouse deer fed different levels of protein .....	190
53.	Nitrogen (N), dietary nitrogen (DN) and dietary crude protein (DCP) requirements for growth of growing male, mature male and mature female mouse deer .....	190
54.	Energy protein ratio (EP), feed conversion ratio (CFR), feed conversion efficiency (FCE) and bodyweight gain (Gain) of growing male (GM), mature male (MM) and mature female (MF) mouse deer fed different levels of protein .....	192
55.	Urinary nitrogen (UN) and faecal nitrogen (FN) of growing and mature male mouse deer fed N-free diet for 7 days .....	196
56.	True digestibility of nitrogen (TD), biological value (BV) and nitrogen requirement for maintenance ( $N_m$ ) of growing male, mature male and mature female mouse deer .....	198



## LIST OF FIGURES

Figure	Page
1. Mature male mouse deer ( <i>Tragulus javanicus</i> ) .....	7
2. Mature female mouse deer ( <i>Tragulus javanicus</i> ) .....	7
3. A model of animal factors affecting feed intake (Source: Lynch <i>et al.</i> , 1992) .....	11
4. Diagrammatic representation of mixtures of feeds require to satisfy animal demands for nutrients. (a) two nutrients; (b) three nutrients; (c) four nutrients. The circled area shows that the animal's requirements upon the nutrients are satisfied. A, B, C, D, E and F indicate types of feeds which content nutrients related to the position in the diagram (Source: Emmans, 1991) .....	13
5. Interrelationships between various forms of energy (Source: McDonald <i>et al.</i> , 1988) .....	23
6. Heat production measurement for mouse deer .....	70
7. The mouse deer keep in an outdoor enclosure with shelter .....	73
8. Energy expenditures of male and female mouse deer in eating, standing and sitting positions .....	77
9. Diurnal pattern of heat production for mouse deer in respiration chamber over 24 hours .....	80
10. Relationships between fibre levels and digestibility of DM (○), OM (▲), CP (●) and GE (□) in mouse deer fed different levels of fibre .....	98
11. Build up curves of Cr in the faeces of mouse deer fed different levels of fibre .....	105
12. Cumulative recovery of Cr in the faeces of mouse deer fed different levels of fibre .....	106
13. Build up curves of Co in the faeces of mouse deer fed different levels of fibre .....	110



14. Cumulative recovery of Co in the faeces of mouse deer fed different levels of fibre .....	111
15. Mean retention times of solid (■) and fluid (□) of mouse deer fed different levels of fibre .....	116
16. Energy retention as a function of metabolisable energy intake for growing (■) and mature (□) male mouse deer fed different levels of feed calculated by using equation I, $MEI = 0.82 DE$ .....	136
17. Energy retention as a function of metabolisable energy intake for growing (■) and mature (□) male mouse deer fed different levels of feed calculated by using equation II, $MEI = GEI - (FE + UE + M)$ ....	136
18. Body weight change as a function of metabolisable energy intake in growing (■) and mature (□) male mouse deer fed different levels of feed calculated by using equation I, $MEI = 0.82 DE$ .....	137
19. Body weight change as a function of metabolisable energy intake in growing (■) and mature (□) male mouse deer fed different levels of feed calculated by using equation II, $MEI = GEI - (FE + UE + M)$ ...	137
20. Changes in CP digestibility in mouse deer fed different levels of protein and energy {1.1 M (□), 1.2 M (■) and 1.3 M (▲)}.....	156
21. Changes in digestible CP intake in mouse deer fed different levels of protein and energy {1.1 M (□), 1.2 M (■) and 1.3 M (▲)}.....	160
22. Relationship between nitrogen intake and nitrogen balance of mouse deer fed different levels of protein and energy {1.1 M (□), 1.2 M (■) and 1.3 M (▲)}.....	163
23. Changes in EP ratio in mouse deer fed different levels of protein and energy {1.1 M (□), 1.2 M (■) and 1.3 M (▲)}.....	167
24. Changes in feed conversion ratio in mouse deer fed different levels of protein and energy {1.1 M (□), 1.2 M (■) and 1.3 M (▲)}.....	168
25. Changes in feed conversion efficiency in mouse deer fed different levels of protein and energy {1.1 M (□), 1.2.M (■) and 1.3 M (▲)}.....	170
26. Relationship between energy protein ratio and body weight gain of mouse deer fed 1.1 M (□), 1.2.M (■) and 1.3 M (▲) energy levels and 8%, 13%, 18% and 23% protein levels.....	174



27. Relationship between nitrogen intake and nitrogen balance in growing male (▲), mature male (●) and mature female mouse deer (□) fed different levels of protein. ....	189
28. Relationship between feed conversion ratio and protein levels of growing male (▲), mature male (●) and mature female (□) mouse deer.....	194
29. Faecal nitrogen (○) and urinary nitrogen (●) of growing male mouse deer fed N-free diet for 7 days .....	197
30. Faecal nitrogen (○) and urinary nitrogen (●) of mature male mouse deer fed N-free diet for 7 days .....	197



## CHAPTER 1

### INTRODUCTION

The lesser mouse deer (*Tragulus javanicus*) is widely distributed throughout the primary and secondary forests of Southeast Asian countries such as Myanmar, Thailand, Kampuchea, Vietnam, Malaysia, Indonesia and the Philippines. It is considered to be one of the most primitive ungulates still in existence (Morat and Nordin, 1978; Kay *et al.*, 1980; Medway, 1983).

Medway (1983) reported that there are two species of mouse deer in Malaysia, namely, the lesser mouse deer (*Tragulus javanicus*) weighing about 1.2 – 1.8 kg and the larger mouse deer (*Tragulus napu*), weighing about 4 – 6 kg. Besides the two species, a number of subspecies such as *Tragulus javanicus pumilus* and *Tragulus javanicus klossi* are also recognized.

The mouse deer is considered a ruminant. However, its stomach consists of only three compartments: the rumen, reticulum and abomasum. Hofmann and Stewards (1972) classified the ruminant into three groups based on their feeding behaviour, i.e. concentrate selector, bulk and roughage eaters and intermediate feeder. The mouse deer can be classified as a concentrate selector (Kay *et al.*, 1980; Agungpriyono *et al.*, 1992). In its natural habitat, the mouse deer feed on

