



***DEVELOPMENT OF GEOSPATIAL MODEL FOR *Metisa plana* (WALKER)
OUTBREAK AND OUTBREAK PREDICTION IN OIL PALM
PLANTATIONS IN MALAYSIA***

SITI AISYAH BINTI RUSLAN

FP 2019 6



**DEVELOPMENT OF GEOSPATIAL MODEL FOR *Metisa plana* (WALKER)
OUTBREAK AND OUTBREAK PREDICTION IN OIL PALM
PLANTATIONS IN MALAYSIA**

By

SITI AISYAH BINTI RUSLAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

November 2018

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

To

The beloved people in my life

For the sacrifices, support, and strength



COPYRIGHT

UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

DEVELOPMENT OF GEOSPATIAL MODEL FOR *Metisa plana* (WALKER) OUTBREAK AND OUTBREAK PREDICTION IN OIL PALM PLANTATION IN MALAYSIA

By

SITI AISYAH BINTI RUSLAN

November 2018

Chairman : Associate Professor Farrah Melissa Muharam, PhD
Faculty : Agriculture

Metisa plana (Walker) is a leaves defoliating insect that is able to cause complete skeletonization and death of the oil palm's fronds. This insect can cause a loss of USD 2.32 billion for two consecutive years given only 10% of the 5 million hectares of oil palms being infested. Hence, efficient, rigorous control methods should be properly planned. In order to do this, the role of environmental factors on the pests' population's fluctuations should be well understood. Nonetheless, the current practices are still leaning towards the conventional approaches that are highly dependent on ineffective, time-consuming in-situ data collection. On the other hand, the utilization of geospatial technologies can be used to obtain data in rapid, harmless, and cost-effective manners. This study utilized the geospatial technologies to i) examine spatial and temporal climatic stresses that cause the outbreak of *Metisa plana*, ii) to construct the relationship between the geospatial data and *Metisa plana* outbreak, and iii) to predict the outbreak of *Metisa plana* in oil palm plantation. *Metisa plana* census data of larvae instar 1, 2, 3, and 4 were collected approximately biweekly over the period of 2014 and 2015. Moderate Resolution Imaging Spectroradiometer (MODIS) and The Tropical Rainfall Measuring Mission (TRMM) satellite images providing values of land surface temperature (LST), rainfall (RF), relative humidity (RH), and Normalized Difference Vegetation Index (NDVI) were extracted and apportioned to 6 time lags; 1 week (T1), 2 weeks (T2), 3 week (T3), 4 weeks (T4), 5 week (T5) and 6 weeks (T6). Linear relationship between *Metisa plana* with LST, RF, RH, and NDVI were carried out using the Pearson's correlation, multiple linear regression (MLR) and multiple polynomial regression analysis (MPR). Artificial neural network (ANN) was then used to develop the best prediction model of *Metisa plana*'s outbreak. Presence of *Metisa plana* was influenced by LST, RF and RH, but not NDVI. The LST between 24°C and 28°C showed a strong relationship with *Metisa plana*, whereby its presence started to decline with LST from 28°C and above. However, the effect of time lag on the presence of *Metisa plana* was not prominent. The best MLR model was obtained

with LST, RF and RH at T4 to T6 with an adjusted $R^2 = 0.29$. The MPR model of LST at T4 to T6 depicted the best fit line with an adjusted $R^2 = 0.50$. The highest accuracy of 95.29% was achieved by models generated by ANN utilizing the relative humidity at T1 to T3. The model generated by combined variables, LST, RF and RH at T4 to T6 was able to predict the presence of *Metisa plana* with the accuracy by up to 89.95%. Based on the result of this study, the elucidation of *Metisa plana*'s landscape ecology was possible with the utilization of geospatial technology.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

PEMBANGUNAN MODEL GEOSPASIAL BAGI LETUSAN *Metisa plana* (WALKER) DAN RAMALAN LETUSANNYA DI PERLADANGAN KELAPA SAWIT MALAYSIA

Oleh

SITI AISYAH BINTI RUSLAN

November 2018

Pengerusi : Profesor Madya Farrah Melissa Muharam, PhD
Fakulti : Pertanian

Metisa plana (Walker) ialah serangga pemakan daun sawit yang berupaya menyumbang kepada kerosakan dan kematian pelepah sawit. Dengan hanya kerosakan minimum sebanyak 10% selama 2 tahun berturut-turut, hasil tanaman daripada 5 juta hektar kawasan lading sawit boleh berkurang sehingga 2.32 bilion USD. Oleh itu, kaedah pengawalan yang efisien bagi menghalang infestasi serangga perosak ini adalah amat penting untuk dirancang. Justeru, kefahaman dan pengetahuan yang mendalam terhadap kesan faktor persekitaran terhadap populasi *Metisa plana* adalah penting. Walau bagaimanapun, pada masa kini, teknik pengumpulan maklumat masih lagi bergantung kepada teknik konservatif yang kurang efisien di mana ianya dilakukan secara *in-situ*. Penggunaan teknologi geospasial mampu untuk membantu pengumpulan maklumat dengan lebih pantas, selamat kepada alam sekitar, dan lebih menjimatkan. Oleh yang demikian, kajian ini memfokuskan kepada penggunaan teknologi geospasial bagi i) mengkaji faktor iklim yang mempengaruhi penularan populasi dan infestasi *Metisa plana* dari segi masa dan spasial, ii) membina hubungan kait di antara data geospasial dan infestasi *Metisa plana*, dan iii) meramal infestasi *Metisa plana* di ladang kelapa sawit. Bancian terhadap bilangan larva *Metisa plana* peringkat satu, dua, tiga, dan empat telah dilakukan pada setiap dua minggu dalam tempoh 2014 dan 2015. Data nilai suhu permukaan (LST), kadar hujan (RF), kelembapan relatif (RH), dan index perbezaan normaliti tumbuhan (NDVI) telah diekstrak daripada imej satelit Moderate Resolution Imaging Spectroradiometer (MODIS) dan The Tropical Rainfall Measuring Mission (TRMM). Data-data yang telah diekstrak daripada imej satelit yang dinyatakan telah dibahagikan kepada enam lag masa iaitu satu sehingga enam minggu sebelum tarikh bancian; satu minggu (T1), dua minggu (T2), tiga minggu (T3), empat minggu (T4), lima minggu (T5), dan enam minggu (T6) bagi tujuan analisa. Analisa korelasi Pearson, regrasi linear berganda (MLR), dan regrasi polynomial berganda (MPR) diantara setiap tarikh bancian dengan 6 lag masa telah digunakan untuk mencari hubungan di antara data geospasial dan

infestasi *Metisa plana*. Setelah itu, rangkaian neural buatan (ANN) telah digunakan untuk menghasilkan model yang terbaik bagi meramal infestasi populasi *Metisa plana*. Hasil kajian telah menunjukkan bahawa terdapat hubungan yang signifikan di antara infestasi *Metisa plana* dengan LST, RF, dan RH, tetapi tidak dengan NDVI. LST di antara 24°C hingga 28°C mempunyai hubungan signifikan yang kuat dengan infestasi *Metisa plana* di mana infestasi *Metisa plana* menurun apabila LST melebihi 28°C. Selain itu, corak lag masa yang tidak konsisten dapat dilihat pada korelasi di antara LST, RF, dan RH dengan *Metisa plana*. Model terbaik telah dihasilkan daripada gabungan LST, RF, dan RH melalui analisa MLR pada T4 hingga T6 dengan R^2 terubah suai = 0.29. Model terbaik daripada MPR menunjukkan LST bagi T4 hingga T6 memberikan garisan fit yang terbaik iaitu pada R^2 terubah suai = 0.5. Model yang dijana oleh ANN menggunakan data kelembapan relatif memberikan ketepatan yang tertinggi iaitu sebanyak 95.29% pada T1 hingga T3. Manakala model yang dijana melalui kombinasi LST, RF, dan RH pada T4 hingga T6 telah memberikan ketepatan sebanyak 89.95%. Hasil kajian ini juga berjaya membuktikan bahawa penggunaan teknologi geospasial mampu meramal infestasi *Metisa plana*.

ACKNOWLEDGEMENTS

Alhamdulillah, the highest gratitude to Allah S.W.T for giving me the opportunity, strength, and wisdom to complete this research project. First and foremost, my deepest thanks are dedicated to the Ministry of Science, Technology, and Innovation (MOSTI) and the Ministry of Higher Education (MOHE) of which without the research funds, this research project would never be executed. My sincerest appreciation to my supervisor, Assoc. Prof. Dr. Farrah Melissa Muharam, who had been my strongest supporter throughout this journey with her love, patience, guidance, understanding, and motivation.

I am nevertheless grateful to my supervisory committee member, Prof. Dr. Dzolkhifli Omar and Dr Zed Diyana Zulkafli, for the help, support, and guidance. Also to the fellow graduate students; Diyana, Zafrullah, Hilmi, Liyana, Ain, Iman, and Adibah for the many kindness and help that was extended to me throughout this journey, without which, the completion of this research would have been impossible.

To my beloved mother, Rihana, my late father, Ruslan, my brother, Asyraf and my sister, Athifah, I deeply owe you for your unconditional love, support and sacrifices. My work is dedicated to you and thank you for being there as a constant reminder that love trumps all.

I would also like to extend my appreciation to the Department of Agriculture Technology academic staff who supported me during my studies, and Tabung Haji Plantation Berhad (THPB), for providing the research area, accommodation, and the necessary support for this study, especially to Haji Pilus Zambri, the General Manager of THPB as well as the staffs of Sungai Mengah estate, your support and guidance will never be forgotten.

Finally, to the Universiti Putra Malaysia, Ministry of Higher Education Malaysia (MOHE) and University Consortium Thesis Grant (SEARCA), I thank the whole organization for supporting me in pursuing my ambition.

This thesis was submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Farrah Melissa Binti Muharam, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Dzolkhifli Bin Omar, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Zed Diyana Binti Zulkafli, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____

Date: _____

Name and Matric No: Siti Aisyah binti Ruslan, GS43879

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Associate Professor Dr. Farrah Melissa Binti Muharam

Signature: _____
Name of Member
of Supervisory
Committee: Professor Dr. Dzolkhifli Bin Omar

Signature: _____
Name of Member
of Supervisory
Committee: Dr. Zed Diyana Binti Zulkafli

TABLE OF CONTENTS

		Page
ABSTRACT		i
ABSTRAK		iii
ACKNOWLEDGEMENTS		v
APPROVAL		vi
DECLARATION		viii
LIST OF TABLES		xii
LIST OF FIGURES		xiv
LIST OF ABBREVIATIONS		xv
CHAPTER		
1	INTRODUCTION	1
	1.1 Study background	1
	1.2 Problem statement	1
	1.3 Justification	2
	1.4 Objective	3
	1.4.1 Specific Objectives	3
2	LITERATURE REVIEW	4
	2.1 Oil palm	4
	2.2 <i>Metisa plana</i>	4
	2.3 Abiotic factors	6
	2.3.1 Temperature	6
	2.3.2 Rainfall	7
	2.3.3 Relative humidity	8
	2.3.4 Normalized difference vegetation index	9
	2.4 Time-lag analysis	10
	2.5 Utilization of remote sensing technology for insect monitoring and outbreak prediction	11
	2.6 Artificial neural network (ANN)	12
	2.7 Conclusion	14
3	MATERIALS AND METHODS	15
	3.1 Study area	15
	3.2 General data collection and processing	17
	3.2.1 Bagworm census data	18
	3.2.2 Geospatial data	18
	3.2.2.1 Land Surface Temperature	18
	3.2.2.2 Rainfall	19
	3.2.2.3 Relative humidity	20
	3.2.2.4 Normalized Difference Vegetation Index	22
	3.2.3 Data extraction	23
	3.3 Data and statistical analysis	25
	3.3.1 Spatio-temporal distribution of <i>Metisa plana</i> 's infestation and weather parameters	25

3.3.2	The determination of the relationship between abiotic factors and <i>Metisa plana</i> 's infestation	25
3.3.3	The development of <i>Metisa plana</i> prediction model using abiotic factors	25
3.3.3.1	Multiple linear regression	25
3.3.3.2	Artificial neural network (ANN)	26
4	RESULTS	28
4.1	Spatio-temporal analysis of <i>Metisa plana</i> 's infestation	28
4.1.1	Description of the seasonal variation of abiotic factors	37
4.2	Relationship between abiotic factors and <i>Metisa plana</i> 's infestation	39
4.2.1	Correlation analysis	39
4.2.1.1	Land surface temperature	39
4.2.1.2	Rainfall	43
4.2.1.3	Relative humidity	47
4.2.1.4	Normalized vegetation index	50
4.2.2	Regression analysis	50
4.3	The development of <i>Metisa plana</i> prediction model	51
5	DISCUSSIONS	53
5.1	Spatio-temporal distribution of <i>Metisa plana</i> 's infestation and weather parameters	53
5.2	The relationship between abiotic factors and <i>Metisa plana</i> 's infestation	55
5.3	Development of <i>Metisa plana</i> 's prediction models using regression analysis and artificial neural network	60
6	CONCLUSIONS	63
	REFERENCES	66
	APPENDICES	79
	BIODATA OF STUDENT	91
	PUBLICATIONS	92

LIST OF TABLES

Table		Page
1	Data source, characteristic, and processing method for independent variables	24
2	Datasets used for regression analysis	26
3	Global Moran 1 analysis of clustering properties of <i>Metisa plana</i> 's infestation in 2014	36
4	Global Moran 1 analysis of clustering properties of <i>Metisa plana</i> 's infestation in 2015	37
5	Comparison of minimum and maximum values of abiotic factors in 2014 and 2015	38
6	Pearson's correlation between 8-days average LST with the average number of <i>Metisa plana</i> in 2014	40
7	Pearson's correlation between 8-days average LST with the average number of <i>Metisa plana</i> in 2015	41
8	Pearson's correlation between 8-days cumulative rainfall with the average number of <i>Metisa plana</i> in 2014	44
9	Pearson's correlation between 8-days cumulative rainfall with the average number of <i>Metisa plana</i> in 2015	45
10	Pearson's correlation between 8-days average relative humidity with the average number of <i>Metisa plana</i> in 2014	47
11	Pearson's correlation between 8-days average relative humidity with the average number of <i>Metisa plana</i> in 2015	48
12	Multiple linear regression analysis between abiotic factors and <i>Metisa plana</i>	50
13	Stepwise multiple polynomial regression analysis between abiotic factors and <i>Metisa plana</i>	51
14	ANN analysis between abiotic parameters and <i>Metisa plana</i>	52
15	t-test analysis to compare abiotic factors of 2014 and 2015	53
16	Comparison of R^2 values obtained from regression analysis and artificial neural network	61

17	Application of remote sensing data for insect monitoring and insect outbreak prediction model	79
18	Application of artificial neural network for insect monitoring and insect outbreak prediction model	86



LIST OF FIGURES

Figure		Page
1	Tabung Haji Plantation, Muadzam Shah	16
2	<i>Metisa plana</i> outbreak prediction method	17
3	<i>Metisa plana</i> 's infestations distribution	29
4	<i>Metisa plana</i> 's infestations distribution	30
5	<i>Metisa plana</i> 's infestations distribution	31
6	<i>Metisa plana</i> 's infestations distribution	32
7	<i>Metisa plana</i> 's infestations distribution	33
8	<i>Metisa plana</i> 's infestations distribution	34
9	Correlation analysis between land surface temperature and number of <i>Metisa plana</i>	42
10	Correlation analysis between rainfall and number of <i>Metisa plana</i>	46
11	Correlation analysis between relative humidity and number of <i>Metisa plana</i>	49
12	<i>Metisa plana</i> 's distribution and LST	54
13	<i>Metisa plana</i> 's distribution and RH	54
14	<i>Metisa plana</i> 's distribution and RF	55

LIST OF ABBREVIATIONS

ANN	Artificial neural network
AWM	Area weighted mean
BBP	Batch back propagation
BP	Back propagation
BSLRC	Broad-spectrum long-residual contact
CGD	Conjugate gradient descent
CMORPH	Climate Prediction Centre morphing method
DAAC	Distributed Active Archive Centre
DBI	David-Bouldin Index
DBM	Diamond backmoth
DD	Degree days
DEM	Digital elevation model
DN	Digital number
ETa	Actual evapotranspiration
ETM+	Enhanced thematic mapper
EVI	Enhanced vegetation index
FFB	Fresh fruit bunch
GES DISC	Goddard Earth Sciences Data and Information Services Center
GIS	Geographic information system
GNI	Gross national income
GPCP-1DD	Global Precipitation Climatology Project 1 Degree Daily
GPS	Global positioning system
ha	Hectare

IPM	Integrated pest management
K	Kelvin
km	Kilometer
LAADS	Level-1 Atmospheric Archive & Distribution System
Landsat TM	Landsat thematic mapper
LM	Levenberg-marquardt
LMQN	Limited memory quasi-newton
LST	Land surface temperature
m	Meter
MAE	Mean Absolute Error
MCI	Moisture condition index
MD	Mean Difference
MLP	Multi-layer perceptron
MLR	Multiple linear regression
mm	Millimetre
MODIS	Moderate Resolution Imaging Spectroradiometer
MPB	Mountain pine beetle
MPR	Multiple polynomial regression
MSE	Mean squared error
MSG	Meteosat second generation
NDIib6	Normalized difference infrared index band 6
NDIib7	Normalized difference infrared index band 7
NDMI	Normalized difference moisture index
NDVI	Normalized vegetation index
NDWI	Normalized difference water index

NetCDF4	Network common data form
NIR	Near infrared
NOAA	National Oceanic and Atmospheric Administration
OBP	Online back propagation
°C	Celsius
p	Calculated probability
PERSIANN-CDR	Precipitation Estimation from Remotely Sensed Information using Artificial Neural Networks Climate Data Record
QN	Quasi- newton
QP	Quick propagation
R	Red
r	Correlation coefficient
R ²	Coefficient of determination
RF	Rainfall
RH	Relative humidity
RMSE	Root Mean Square Error
SARIMA	Seasonal autoregressive integrated moving average
SD	Standart Deviation
SPOT	Satellites Pour l'Observation de la Terre or Earth-observing Satellites
SRTM	Shuttle radar topography mission
SWIR	Short-wavelength infrared
TCI	Temperature condition index
TRMM	The Tropical Rainfall Measuring Mission
USGS	United States Geological Survey

CHAPTER 1

INTRODUCTION

1.1 Study background

The worldwide consumption of palm oil soared from 14.6 million tonnes in 1995 to 61.1 million tonnes in 2015, making it the most consumed oil in the world (European Palm Oil Alliance, 2016). Malaysia ranks second among the largest palm oil global exporter after the neighbouring country Indonesia, supplying by up to 32.6 % of the world's total palm oil exports, which can be translated to approximately USD 9 billion in 2016 (Workman, 2017). Palm oil is also the fourth largest contributor to the national economy, accounting RM 52.7 billion of the Gross National Income (GNI) and is projected to raise to up to RM 178 billion worth of GNI in 2020 (Pemandu.gov.my, 2014). The global dependence of palm oil will continue to rise and its demand is expected to increase to an additional 25 million tonnes per year in the forthcoming 10 years. Consequently, in order to satiate the demand, the production of palm oil will have to be amplified to 85 million tonnes a year (Mielke, 2017). On that account, it is important for diligent efforts and good agriculture practices to take place in order to increase the productivity of oil palm and consequently to achieve the desired yield production. These however, can be compromised by the presence of insect pests such as bagworm.

Bagworm (*Lepidoptera: Psychidae*) is an important defoliating caterpillar that relies on crop leaves as their source of nutrients and shelter. In order to survive, bagworm larvae use 68% of the leaf tissues removed from oil palm for growth and development while the other 32% to construct portable outer cases which act as shelter that they carry around with them as they feed (Basri *et al.*, 1995; Kok *et al.*, 2011). In Malaysia, the most economically devastating species of bagworms for oil palm plantations are *Metisa plana* (Walker), *Pteroma pendula* (Joannis), and *Mahasena corbetti* (Tams). Among these species, *Metisa plana* have caused the most detrimental effect in Peninsular Malaysia, judging by the magnitude of its infestations and damages (Basri *et al.*, 1988) that is resulting from efficient dispersion mechanisms that lead to high reproductive success (Tuck *et al.*, 2011).

1.2 Problem statement

Untreated infestation of *Metisa plana* can lead to a devastating losses in oil palm industry mainly due to the potential of causing a complete skeletonization and an eventual death of oil palm fronds. This will jeopardize the ability of palms to carry out photosynthetic activities that will suppress the growth of the palms and reduce the efficiency of palms to produce optimum yield. Wood *et al.* (1973) earlier captured the threatening nature of this pest by demonstrating that 50% of damage caused by its infestation will bring about 43% or approximately 10 t ha⁻¹ of fresh fruit bunch (FFB) for the next two consecutive years. This was later supported by Basri (1993) which

stated that even at a lower level of damage of 10% to 13%, the loss of yield of oil palm can go up to 40%. These figures can be translated into a loss of approximately USD 2,032 per hectare, which will eventually cause a loss of USD 2.32 billion for two consecutive years, given only 10% of the 5 million hectares of oil palms being infested.

Taking into account the high potential of severe economic losses that could be initiated by *Metisa plana*, rigorous control methods and mitigation systems should be properly planned and executed. Furthermore, the control methods must be efficient in terms of cost, time, and man power, as well as not harmful to oil palms and the environment. The success in doing so is to ensure that necessary steps are to be carried out at the precise times and locations. In order to achieve this, the ecological knowledge of *Metisa plana* especially spatial and temporal domain, should be well understood. Ecological aspects such as weather plays an important role in the lavishness of bagworm's outbreak owing to its adverse influence on insect's behaviour.

1.3 Justification

Currently, the effort of understanding weather aspects affecting the insect pest outbreaks, along with their control practices are still leaning towards the conventional approaches that are highly dependent on in-situ data collection which can be ineffective. The exploitation of modern technology such as geospatial technology have essentially benefited agricultural industry. This technology, which encompasses of remote sensing, global positioning system (GPS), and geographic information system (GIS) have been used in the assessment and monitoring of crop pests and diseases. Unlike the conventional approaches, information on the triggering factors of insect pest outbreaks such as temperature, rainfall, and vegetation's condition can be obtained through this technology in rapid, harmless, and cost-effective manners. Hence, early detection of insect pest outbreaks via geospatial technology can potentially (i) reduce labour time and cost, (ii) limit environmental pollution, and (iii) reduce the potential of a devastating impact of insect pest outbreaks by controlling them before they spread. Furthermore, the monitoring of crops condition within a large area of plantation could be done synchronously with the help of geospatial technology.

1.4 Objective

The overall objective of this study is to develop a model to predict the outbreak of *Metisa plana* in oil palm plantation.

1.4.1 Specific Objectives

The specific objectives of this study are:

- i. To compare the spatio-temporal distribution of *Metisa plana*'s infestation and weather parameters in 2014 and 2015.
- ii. To investigate the remotely-sensed derived factors that influence the outbreak of *Metisa plana* in oil palm plantation.
- iii. To develop model that incorporates the combination of remotely-sensed derived factors influencing the outbreak of *Metisa plana*.

REFERENCES

- Abdullah, R. (2011). World palm oil supply, demand, price, and prospects: Focus on Malaysian and Indonesian palm oil industries. *Oil Palm Industry Economic Journal*, 11, 13–25.
- Ahmadi, P., Muharam, F. M., Ahmad, K., Mansor, S., & Abu Seman, I. (2017). Early detection of ganoderma basal stem rot of oil palms using artificial neural network spectral analysis. *Plant Disease*, 101, 1009–1016. <https://doi.org/10.1094/PDIS-12-16-1699-RE>
- Ameen, M.-U., & Sultana, P. (1977). Biology of the bag-worm moth *Eumeta crameri* Westwood (Lepidoptera: Psychidae) from Dacca, Bangladesh. *Journal of Natural History*, 11, 17–24.
- ArcGIS Pro, (no date). Retrieved from <http://pro.arcgis.com/en/pro-app/tool-reference/spatial-statistics/h-how-spatial-autocorrelation-moran-s-i-spatial-st.htm>
- Assante, S. K., Danthanarayana, W., & Heatwole, H. (1991). Bionomics and population growth statistics of apterous virginoparae of woolly apple aphid, *Eriosoma lanigerum*, at constant temperatures. *Entomologia Experimentalis et Applicata*, 60, 261–270.
- Awmack, C. S., Woodcock, C. M., & Harrington, R. (1997). Climate change may increase vulnerability of aphids to natural enemies. *Ecological Entomology*, 22, 366–368.
- Azevedo de Freitas, F., Zanunciob, T. V., Zanunciob, J. C., Marluci da Conceiçãob, P., Queiroz Fialhob, M. do C., & Bernardinob, A. S. (2005). Effect of plant age, temperature and rainfall on Lepidoptera insect pests collected with light traps in a Eucalyptus grandis plantation in Brazil. *Annals of Forest Science*, 62, 85–90.
- Bale, J. S., Masters, G. J., Hodkinson, I. D., Awmack, C., Bezemer, M., Brown, V. K., ... Whittaker, J. B. (2002). Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Global Change Biology*, 8, 1–16.
- Barbosa, P., Waldvogel, M. G., & Breisch, N. L. (1983). Temperature modification by bags of the bagworm *Thyridopteryx ephemeraeformis* (Lepidoptera: Psychidae). *The Canadian Entomologist*, 115, 855–858.
- Barlow, H. S. (1982). *An Introduction to the Moths of South East Asia*. Kuala Lumpur: Malaysian Nature Society.
- Basri, M. W. (1993). *Life history, ecology, and economic impact of the bagworm, Metisa plana Walker (Lepidoptera: Psychidae) on the oil palm, Elaeis guineensis Jacquon (Palmae), in Malaysia*. University of Guelph.

- Basri, M. W., Abdul Halim, H., & Masijan, Z. (1988). *Bagworms (Lepidoptera: Psychidae) of oil palms in Malaysia*. PORIM Occasional Paper (Vol. 23).
- Basri, M. W., Norman, K., & Hamdan, A. B. (1995). Natural enemies of the bagworm, *Metisa plana* Walker (Lepidoptera Psychidae) and their impact on host population regulation. *Crop Protection*, *14*, 637–645.
- Basri, W., Hassan, A. H., & Zulkifli, M. (1988). *Bagworms (Lepidoptera: Psychidae) of Oil Palms in Malaysia*. Bangi: Institut Penyelidikan Minyak Sawit Malaysia.
- Benali, A., Carvalho, A. C., Nunes, J. P., Carvalhais, N., & Santos, A. (2012). Estimating air surface temperature in Portugal using MODIS LST data. *Remote Sensing of Environment*, *124*, 108–121.
- Benrey, B., & Denno, R. F. (1997). The slow-growth-high mortality hypothesis: a test using the caggabe butterfly. *Ecology*, *78*, 987–999. [https://doi.org/10.1890/0012-9658\(1997\)078\[0987:TSGHMH\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1997)078[0987:TSGHMH]2.0.CO;2)
- Blum, M., & Lensky, Itamar M. Rempoulakis, Polychronis Nestel, D. (2015). Modeling insect population fluctuations with satellite land surface temperature. *Ecological Modelling*, *311*, 39–47.
- Blum, M., Lensky, I. M., & Nestel, D. (2013). Estimation of olive grove canopy temperature from MODIS thermal imagery is more accurate than interpolation from meteorological stations. *Agricultural and Forest Meteorology*, *176*, 90–93. <https://doi.org/10.1016/j.agrformet.2013.03.007>
- Blum, M., Nestel, D., Cohen, Y., Goldshtein, E., Helman, D., & Lensky, I. M. (2018). Predicting *Heliothis (Helicoverpa armigera)* pest population dynamics with an age-structured insect population model driven by satellite data. *Ecological Modelling*, *369*, 1–12.
- Castillo, J., Jacas, J. A., Peña, J. E., Ulmer, B. J., & Hall, D. G. (2006). Effect of temperature on life history of *Quadrastichus haitiensis* (Hymenoptera: Eulophidae), an endoparasitoid of *Diaprepes abbreviatus* (Coleoptera: Curculionidae). *Biological Control*, *36*, 189–196.
- Cheong, Y. L., & Tey, C. C. (2013). Outbreak, Environmental Factors which Influences Bagworm. In *Proceedings of 5th MPOB-IOPRI International Seminar: Sustainable Management of Pests and Disease in Oil Palm* (pp. 1–15). Kuala Lumpur.
- Chon, T.-S., Park, Y.-S., Kim, J.-M., Lee, B.-Y., Yeong-Jin Chung, & Kim, Y. (2000). Use of an artificial neural network to predict population dynamics of the forest-pest pine needle gall midge (Diptera: Cecidomyiida). *Environmental Entomology*, *29*, 1208–1215.
- Chung, G. F., & Sim, S. C. (1991). *Bagworm Census and Control: A Case Study*. Kuala Lumpur: PORIM.

- Colombi, A., Michele, C. De, Pepe, M., & Rampini, A. (2007). Estimation of Daily Mean Air Temperature. *EARSeL EProceedings*, 6, 38–46.
- Corley, R. H. ., & Tinker, P. . (2003a). Disease and Pests of the Oil Palm. In *The Oil Palm* (4th ed., pp. 391–444). Oxford, UK: Blackwell Science Ltd. <https://doi.org/https://doi.org/10.1002/9780470750971.ch3>
- Corley, R. H. V., & Tinker, P. . (2003b). The Climate and Soils of the Oil Palm-growing Regions. In *The Oil Palm* (4th ed., pp. 53–88). Oxford, UK: Blackwell Science Ltd.
- Cowtan, K., & Way, R. G. (2014). Coverage bias in the HadCRUT4 temperature series and its impact on recent temperature trends. *Quarterly Journal of the Royal Meteorological Society*, 140, 1935–1944.
- Cox, D. L., & Potter, D. A. (1986). Aerial dispersal behavior of larval bagworms, *Thyridopteryx ephemeraeformis* (Lepidoptera: Psychidae). *The Canadian Entomologist*, 118, 525–536.
- de Beurs, K. M., & Townsend, P. A. (2008). Estimating the effect of gypsy moth defoliation using MODIS. *Remote Sensing of Environment*, 112, 3983–3990.
- De Conti, B., Bueno, V. H. P., Sampaio, M. V., & Sidney, L. A. (2010). Reproduction and fertility life table of three aphid species (*Macrosiphini*) at different temperatures. *Revista Brasileira de Entomologia*, 54, 654–660.
- Dent, D. R. (1997). Quantifying insect populations: estimates and parameters. In D. R. Dent & M. P. Walton (Eds.), *Methods in ecological and agricultural entomology* (pp. 57–110). New York: Wallingford, Oxon, UK.
- Dury, S. J., Good, J. E. G., Perrins, C. M., Buse, A., & Kaye, T. (1998). The effects of increasing CO₂ and temperature on oak leaf palatability and the implications for herbivorous insects. *Global Change Biology*, 4, 55–62.
- Ekstrand, S. (1996). Landsat TM-Based Forest Damage Assessment: Correction for Topographic Effects. *Photogrammetric Engineering & Remote Sensing*, 62, 151–161.
- European Palm Oil Alliance. (2016). Palm oil consumption. Retrieved August 17, 2017, from <http://www.palmoilandfood.eu/en/palm-oil-consumption>
- Fargues, J., & Luz, C. (2000). Effects of fluctuating moisture and temperature regimes on the infection potential of *Beauveria bassiana* for *Rhodnius prolixus*. *Journal of Invertebrate Pathology*, 75, 202–211.
- Fedor, P., Vaňhara, J., Havel, J., Malenovsky, I., & Spellerberg, I. (2009). Artificial intelligence in pest insect monitoring. *Systematic Entomology*, 34, 398–400.

- Fragoso, D. B., Zanuncio, T. V., Zanuncio, J. C., & Filho, P. J. (2000). Dinâmica populacional de lepidópteros em plantios de *Eucalyptus grandis* em Santa Bárbara, Minas Gerais. *Revista Arvore*, *24*, 253–259.
- Gan, P. Y., & Li, Z. D. (2014). Economic study on Malaysia's oil palm position in the world market to 2035. *Renewable and Sustainable Energy Reviews*, *39*(2014), 740–747. <https://doi.org/https://doi.org/10.1016/j.rser.2014.07.059>
- Gao, B.-C., & Kaufman, Y. J. (2003). Water vapor retrievals using Moderate Resolution Imaging Spectroradiometer (MODIS) near-infrared channels. *Journal of Geophysical Research: Atmospheres*, *108*, 1–9. <https://doi.org/10.1029/2002JD003023>
- Geiger, C. A., & Gutierrez, A. P. (2000). Ecology of *Heteropsylla cubana* (Homoptera: Psyllidae): psyllid damage, tree phenology, thermal relations and parasitism in the field. *Population Ecology*, *29*, 76–86.
- Geng, L., Ma, M., Yu, W., Wang, X., & Jia, S. (2014). Validation of the MODIS NDVI Products in Different Land-Use Types Using In Situ Measurements in the Heihe River Basin. *IEEE Geoscience and Remote Sensing Letters*, *11*, 1649–1653.
- Godfrey, L. D., Holtzer, T. O., & Norman, J. M. (1991). Effects of european corn borer (Lepidoptera: Pyralidae) tunneling and drought stress on field corn gas exchange parameters. *Journal of Economic Entomology*, *84*, 1370–1380.
- Goodwin, Nicholas R. Coops, N. C., Wulder, M. A., Gillanders, S., & Schroeder, Todd A. Nelson, T. (2008). Estimation of insect infestation dynamics using a temporal sequence of Landsat data. *Remote Sensing of Environment*, *112*, 3680–3689.
- Guarneri, A. A., Lazzari, C., Diotaiuti, L., & Lorenzo, M. G. (2002). The effect of relative humidity on the behaviour and development of *Triatoma brasiliensis*. *Physiological Entomology*, *27*, 142–147.
- Gullan, P. J., & Cranston, P. S. (2004). *The Insects: An Outline of Entomology*, 3rd Edition (3rd ed.). Blackwell Publishing Ltd.
- Hachem, S., Duguay, C. R., & Allard, M. (2012). Comparison of MODIS-derived land surface temperatures with ground surface and air temperature measurements in continuous permafrost terrain. *The Cryosphere*, *6*, 51–69.
- Hill, R. L., Gourlay, A. H., & Winks, C. J. (1993). Choosing gorse spider mite strains to improve establishment in different climates. In R. A. Prestidge (Ed.), *Proceedings of the Sixth Australasian Grasslands Invertebrate Ecology Conference* (pp. 377–383). Hamilton, New Zealand: AgResearch.
- Hirose, E., Panizzi, A. R., & Cattelan, A. J. (2006). Effect of relative humidity on emergence and on dispersal and regrouping of first instar *Nezara viridula* (L.) (Hemiptera: Pentatomidae). *Ecology, Behaviour, and Bionomics*, *35*, 757–761.

- Holben, B., & Justice, C. (1980). The topographic effect on spectral response from nadir-pointing. *Photogrammetric Engineering and Remote Sensing*, 46, 1191–1200.
- Hoong, H. W., & Hoh, C. K. Y. (1992). Major pests of oil palm and their occurrence in Sabah. *The Planter*, 68, 193–210.
- Hou, Y., & Weng, Z. (2010). Temperature-dependent development and life table parameters of *Octodonta nipae* (Coleoptera: Chrysomelidae). *Environmental Entomology*, 39, 1676–1684.
- Howard, F. W. (2001). The Animal Class Insecta and the Plant Family Palmae. In F. W. Howard, D. Moore, R. M. Giblin-Davis, & R. G. Abad (Eds.), *Insects on Palms* (p. 414). Wallingford, Oxfordshire: CABI Publishing.
- Huang, Z., Ren, S., & Musa, P. D. (2008). Effects of temperature on the development, survival, longevity, and fecundity of the *Bemisia tabaci gennadius* (Homoptera: Aleyrodidae) predator, *Axinoscymnus cardilobus* (Coleoptera: Coccinellidae). *Biological Control*, 46, 209–215.
- Huete, A., Justice, C., & van Leeuwen, W. (1999). *MODIS vegetation index - ALGORITHM THEORETICAL BASIS*. *Environmental Sciences*.
- Ibrahim, Y., Ho, C. T., & Chong, K. K. (2013). Effects of temperature on the development and survival of the bagworms *Pteroma pendula* and *Metisa plana* (Lepidoptera: Psychidae). *Journal of Oil Palm Research*, 25, 1–8.
- Jandricic, S. E., Wraight, S. P., Bennett, K. C., & Sanderson, J. P. (2010). Developmental times and life table statistics of *Aulacorthum solani* (Hemiptera: Aphididae) at six constant temperatures, with recommendations on the application of temperature-dependent development models. *Environmental Entomology*, 39, 1631–1642.
- Jaworski, T., & Hilszczański, J. (2013). The effect of temperature and humidity changes on insects development and their impact on forest ecosystems in the context of expected climate change. *Forest Research Paper*, 74, 345–355. <https://doi.org/10.2478/frp-2013-0033>
- Ji, R., Xie, B.-Y., Li, D.M., Lia, Z., & Zhang, X. (2004). Use of MODIS data to monitor the oriental migratory locust plague. *Agriculture, Ecosystems and Environment*, 104, 615–620.
- Jiang, L., Shang, Y., & Guan, H. (2016). Mapping interannual variability of maize cover in a large irrigation district using vegetation index-phenological index classifier. *Computers and Electronic in Agriculture*, 123, 351–361.
- Karpakakunjaram, V., Milind, D., & Muralirangan, M.C. (2002) effects of abiotic factors on the population of an acridid grasshopper, *Diabolo-catantops pinguis* (Orthoptera: Acrididae) at two sites in southern India: a three-year study. *Journal of Orthoptera Research*, 11, 55–62.

- Kaufman, Y. J., & Gao, B.C. (1992). Remote sensing of water vapor in the near IR from EOS/MODIS. *IEEE Transactions on Geoscience and Remote Sensing*, 30, 871–884.
- Kaufmann, T. (1968). Observation on the biology and behaviour of the evergreen bagworm moth, *Thyridopteryx ephemeraeformis* (Lepidoptera:Psychidae). *Annals of the Entomological Society of America*, 1, 38–44.
- Kobori, Y., & Amano, H. (2003). Effect of rainfall on a population of the diamondback moth, *Plutella xylostella* (Lepidoptera: Plutellidae). *Applied Entomology and Zoology*, 38, 249–253.
- Kok, C. C., Eng, O. K., Razak, A. R., & Arshad, A. M. (2011). Microstructure and life cycle of *Metisa plana* Walker (Lepidoptera: Psychidae). *Journal of Sustainability Science and Management*, 6, 51–59.
- Kolb, T. E., Fettig, C. J., Bentz, B. J., Stewart, J. E., Weed, A. S., Hicke, J. A., & Ayres, M. P. (2016). Fungal Pathogen Responses to Drought : A Comprehensive Science Synthesis. In T. Vose, James M.; Clark, James S.; Luce, Charles H.; Patel-Weynand (Ed.), *Effects Of Drought on Forests and Rangelands in the United States* (pp. 114–133). Washington, DC: U.S. Department of Agriculture, Forest Service, Washington Office.
- Kollberg, I. (2013). *The Effect of Temperature on Trophic Interactions, Implications for the Population Dynamics of a Forest Pest Insect in a Warmer Climate*. Swedish University of Agricultural Sciences.
- Kuppusamy, A., & Kannan, S. (1993). Life-history of *Eumeta crameri* Westwood (Lepidoptera: Psychidae) and its natural enemy *Sinophorus psycheae* Sonan (Hymenoptera: Ichneumonidae). *Phytophaga*, 5, 9–21.
- Lassau, S. A., & Hochuli, D. F. (2008). Testing predictions of beetle community patterns derived empirically using remote sensing. *Diversity and Distribution*, 14,138–147.
- Lee, K. Y., Chung, N., & Hwang, S. (2016). Application of an artificial neural network (ANN) model for predicting mosquito abundances in urban areas. *Ecological Informatics*, 36, 172–180.
- Lee, T. Y. L. T. Y., & Kaufman, Y. J. (1986). Non-lambertian effects on remote sensing of surface reflectance and vegetation index. *IEEE Transactions on Geoscience and Remote Sensing*, 24, 699–708.
- Levin, D. A. (1973). The role of trichome in plant defense. *The Quarterly Review of Biology*, 48, 3–15.
- Lorenz, C., Ferraudo, A. S., & Suesdeka, L. (2015). Artificial neural network applied as a methodology of mosquito species identification. *Acta Tropica*, 152, 165–169.

- Luoto, M., Kuussaari, M., & Toivonen, T. (2002). Modelling butterfly distribution based on remote sensing data. *Journal of Biogeography*, 29, 1027–1037.
- Luz, C., & Fargues, J. (1997). Temperature and moisture requirements for conidial germination of an isolate of *Beauveria bassiana*, pathogenic to *Rhodnius prolixus*. *Mycopathologia*, 138, 117–125.
- Marques da Silva, J. R., Damásioe, C. V., Sousaa, A. M. O., Bugalhof, L., Pessanhaf, L., & Quaresmaa, P. (2015). Agriculture pest and disease risk maps considering MSG satellite data and land surface temperature. *International Journal of Applied Earth Observation and Geoinformation*, 38, 40–50.
- Mattson, W. J., & Haack, R. A. (1987). Role of drought in outbreaks of plant-eating insects. *Bioscience*, 37, 110–118.
- Merrill, S. C., Holtzer, T. O., Pears, F. B., & Lester, P. J. (2009). Modeling spatial variation of Russian wheat aphid overwintering population densities in Colorado winter wheat. *Journal of Economic Entomology*, 102, 533–541. <https://doi.org/10.1603/029.102.0210>
- Midekisa, A., Senay, G., Henebry, G. M., Semuniguse, P., & Wimberly, M. C. (2012). Remote sensing-based time series models for malaria early warning in the highlands of Ethiopia. *Malaria Journal*, 11, 1–10.
- Mielke, T. (2017). Global Supply, Demand and Price Outlook for Vegetable Oils as well as for Palm Oil. *ISTA Mielke GmbH, OIL WORLD, Hamburg*, (March 2016), 1–5. Retrieved from https://www.oilworld.biz/t/sample/sample_34.pdf
- Mishra, M., Bhavyashree, S., & H. L. Sharma. (2017). Comparative study of the performance of artificial neural network and multiple linear regression techniques for predicting the soybean yield using its attributing characters. *International Quarterly Journal of Life Sciences*, 12, 177–181.
- Mishra, S. . (1978). Observation on the biology of bagworm *Clania crameri* Westwood (Psychidae: Lepidoptera). *Indian Forester*, 104, 35–41.
- Mishra, S., Kumar, P., & Malik, A. (2013). Effect of temperature and humidity on pathogenicity of native *Beauveria bassiana* isolate against *Musca domestica* L. *Journal of Parasitic Diseases*, 39, 697–704.
- Mohd Basri, W., & Kevan, P. (1995). Life history and feeding behaviour of the oil palm bagworm, *Metisa plana* Walker (Lepidoptera: Psychidae). *Elaeis*, 7, 18–34.
- Moran, V. C., Hoffman, J. H., & Basson, N. C. J. (1987). The effects of simulated rainfall on cochineal insects (Homoptera: Dactylopidiidae): colony composition and survival on cactus cladodes. *Ecological Entomology*, 12, 51–60.
- Mullen, B. F., Shelton, H. M., Gutteridge, R. C., & Basford, K. E. (2003). Agronomic evaluation of *Leucaena*. Part 1. Adaptation to environmental challenges in multi-environment trials. *Agroforestry System*, 58, 77–92.

- National Aeronautics & Space Administration (no date). Retrieved from <https://pmm.nasa.gov/trmm/tmi>
- Neal, J., & Santamour, F. (1990). Biotic indicators of host preference by the bagworm (Lepidoptera: Psychidae). *Journal of Economic Entomology*, 83, 93–97.
- Netherer, S., & Schopf, A. (2010). Potential effects of climate change on insect herbivores in European forests—general aspects and the pine processionary moth as specific example. *Forest Ecology and Management*, 259, 831–838.
- Nishikawa, H., Shimada, T., Nakahira, K., & Arakawa, R. (2010). Thermal effect on the development and reproduction of an indigenous mirid bug, *Pilophorus typicus* distant (Heteroptera: Miridae), a potential biological control agent in Japan. *Applied Entomology and Zoology*, 45, 313–318.
- Norhisham, A. R., Abood, F., Rita, M., & Hakeem, K. R. (2013). Effect of humidity on egg hatchability and reproductive biology of the bamboo borer (*Dinoderus minutus* Fabricius). *Springer Plus*, 2, 2–6.
- Norman, K., & Basri, M. W. (2007). Status of common oil palm insect pests in relation to technology adoption. *The Planter*, 83(975), 371–385.
- Norris, R. J., Memmott, J., & Lovell, D. J. (2002). The effect of rainfall on the survivorship and establishment of a biocontrol agent. *Journal of Applied Ecology*, 39, 226–234.
- Ofomata, V. C., Overholt, W. A., Lux, S. A., Huis, A. Van, & Egwuatu, R. I. (2000). Comparative studies on the fecundity, egg survival, larval feeding, and development of *Chilo partellus* and *Chilo orichalcociliellus* (Lepidoptera: Crambidae) on five grasses. *Ecology and Population Biology*, 93, 492–499.
- Parajulee, M. (2006). Influence of constant temperatures on life history parameters of the cotton aphid, *Aphis gossypii*, infesting cotton. *Environmental Entomology*, 36, 666–672.
- Parajulee, M. N., Shrestha, R. B., Leser, J. F., Wester, D. B., & Blanco, C. A. (2006). Evaluation of the functional response of selected arthropod predators on bollworm eggs in the laboratory and effect of temperature on their predation efficiency. *Environmental Entomology*, 35, 379–386.
- Parinussa, R.M., de Jeu, R.A.M., Holmes T.R.H., & Walker, J.P. (2008). Comparison of microwave and infrared land surface temperature products over the NAFE'06 research sites. *IEEE Geoscience and Remote Sensing Letters*, 5, 783-787.
- Parinussa, R.M., Lakshmi, V. Johnson, F., & Sharma, A. (2016). Comparing and combining remotely sensed land surface temperature products for improved hydrological applications. *Remote Sensing*, 8, 1-14.

- Park, Y.-S., Céréghino, R., Compin, A., & Lek, S. (2003). Applications of artificial neural networks for patterning and predicting aquatic insect species richness in running waters. *Ecological Modelling*, *160*, 265–280.
- Paruelo, J. M., & Tomasel, F. (1997). Prediction of functional characteristics of ecosystems: a comparison of artificial neural networks and regression models. *Ecological Modelling*, *97*, 173–186.
- Pellegrino, A. C., Peñaflor, M. F. G. V., Nardi, C., Bezner-Kerr, W., Guglielmo, C. G., Bento, José Maurício Simões, J. N., & McNeil. (2013). Weather forecasting by insects: Modified sexual behaviour in response to atmospheric pressure changes. *Plos One*, *8*, e75004.
- Pemandu.gov.my. (2014). Palm oil and rubber. Retrieved from http://etp.pemandu.gov.my/Palm_Oil-@-Palm_Oil-_-_-Rubber.aspx
- Peng, G., Li, J., Chen, Y., Norizan, A. P., & Tay, L. (2006). High-resolution surface relative humidity computation using MODIS image in Peninsular Malaysia. *Chinese Geographical Science*, *16*, 260–264. <https://doi.org/10.1007/s11769-006-0260-6>
- Pereira, J. M. M., Zanuncio, T. V., Zanuncio, J. C., & Pratisoli, D. (2001). Lepidoptera pests collected in *Eucalyptus urophylla* plantations during five years in Três Marias, State of Minas Gerais, Brazil. *Revista de Biología Tropical*, *49*, 1073–1082.
- Propastin, P. (2012). Multisensor monitoring system for assessment of locust hazard risk in the lake Balkhash drainage basin. *Environmental Management*, *50*, 1234–1246.
- Rhains, M., Davis, D. R., & Price, P. W. (2009). Bionomics of bagworms (Lepidoptera: Psychidae). *The Annual Review of Entomology*, *54*, 209–226. <https://doi.org/https://doi.org/10.1146/annurev.ento.54.110807.090448>
- Rhains, M., Gries, G., & Chew, P. S. (1997). Adaptive significance of density-dependent ballooning by bagworm larvae, *Metisa plana* (Walker) (Lepidoptera: Psychidae). *The Canadian Entomologist*, *129*, 927–931.
- Rhains, M., Gries, G., Ho, C. T., & Chew, P. S. (2002). Dispersal by bagworm, *Metisa plana*: effects of population density, larval sex, and host plant attributes. *Ecological Entomology*, *27*, 204–212.
- Rhains, M., Gries, G., & Mat Min, M. (1999). Size-and density-dependent reproductive success of bagworms *Metisa plana*. *Entomologia Experimentalis et Applicata*, *91*, 375–383.
- Rhains, M., Gries, G., & Min, M. M. (1999). Size- and density-dependent reproductive success of bagworms, *Metisa plana*. *Entomologia Experimentalis et Applicata*, *91*, 375–383.

- Rhains, M., Leather, S. R., & Sadof, C. (2008). Polyphagy, flightlessness, and reproductive output of females: a case study with bagworms (Lepidoptera: Psychidae). *Ecological Entomology*, *33*, 663–672.
- Rouault, G., Candau, J.-N., Lieutier, F., Nageleisen, L.-M., Martin, J.-C., & Warzée, N. (2006). Effects of drought and heat on forest insect populations in relation to the 2003 drought in Western Europe. *Annals of Forest Science*, *63*, 613–624.
- Sahin, M., Kaya, Y., & Uyar, M. (2013). Comparison of ANN and MLR models for estimating solar radiation in Turkey using NOAA/AVHRR data. *Advances in Space Research*, *51*, 891–904.
- Sajap, A. S., & Siburat, S. (1992). Incidence of entomogenous fungi in the bagworm, *Pteroma pendula* (Lepidoptera: Psychidae), a pest of *Acacia mangium*. *Journal of Plant Protection in the Tropics*, *9*, 105–110.
- Scherber, C., Gladbach, D. J., Stevnbak, K., Karsten, R. J., Schmidt, I. K., Michelsen, A., ... Christensen, S. (2013). Multi-factor climate change effects on insect herbivore performance. *Ecology and Evolution*, *3*, 1–12.
- Scriber, J. M. (1977). Limiting effects of low leaf-water content on the nitrogen utilization, energy budget, and larval growth of *Hyalophora cecropia* (Lepidoptera: Saturniidae). *Oecologia*, *28*, 269–287.
- Shipp, J. L., Zhang, Y., Hunt, D. W. A., & Ferguson, G. (2003). Influence of humidity and greenhouse microclimate on the efficacy of *Beauveria bassiana* (Balsamo) for control of greenhouse arthropod pests. *Environmental Entomology*, *32*, 1154–1163.
- Sime Darby Plantation. (2014). *Palm Oil Facts and Figures*.
- Smith, M. P., & Barrows, E. M. (1991). Effects of larval case size and host plant species on case internal temperature in the bagworm, *Thyridopteryx ephemeraeformis* (Haworth) (Lepidoptera: Psychidae). In *Proceedings of the Entomological Society of Washington* (Vol. 93, pp. 834–838). Retrieved from <http://biostor.org/reference/56522>
- Sprintsin, M., Chen, J. M., & Czurylowicz, P. (2011). Combining land surface temperature and shortwave infrared reflectance for early detection of mountain pine beetle infestations in western Canada. *Journal of Applied Remote Sensing*, *5*, 1–13.
- Stephen, C. S. (1962). *Oiketicus kirbyi* (Lepidoptera:Psychidae) : a pest of bananas in Costa Rica. *Journal of Economic Entomology*, *55*, 381–386.
- Syed, R. A., & Shah, S. (1977). Some Important Aspects of Insect Pest Management in Oil Palm Estates in Sabah, Malaysia. In D. A. Earp & W. Newell (Eds.), *International developments in oil palm* (pp. 577–590). Kuala Lumpur: The Incorporated Society of Planters.

- Tailliez, B., & Ballo Koffi, C. (1992). A method for measuring oil palm leaf area. *Oleagineux*, 47, 537–545.
- Tamiru, A., Getu, E., Jembere, B., & Bruce, T. (2012). Effect of temperature and relative humidity on the development and fecundity of *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae). *Bulletin of Entomological Research*, 102, 9–15.
- Tan, M. L., Ibrahim, A. L., Duan, Z., Cracknell, A. P., & Chaplot, V. (2015). Evaluation of six high-resolution satellite and ground-based precipitation products over Malaysia. *Remote Sensing*, 7, 1504–1528. <https://doi.org/10.3390/rs70201504>
- Taveras, R., Hilje, L., Hanson, P., MExzon, R., Carballo, M., & Navarro, C. (2004). Population trends and damage patterns of *Hypsipyla grandella* (Lepidoptera: Pyralidae) in mahogany stand, in Turrialba, Costa Rica. *Agricultural and Forest Entomology*, 6, 89–98.
- Tonnang, H. E. Z., Nedorezov, L. V., Owino, J. O., Ochanda, H., & Löhr, B. (2010). Host-parasitoid population density prediction using artificial neural networks: diamondback moth and its natural enemies. *Agricultural and Forest Entomology*, 12, no-no. <https://doi.org/10.1111/j.1461-9563.2009.00466.x>
- Tuck, H. C. (2002). *Ecological Studies on Pteroma pendula Joannis and Metisa plana Walker (Lepidoptera: Psychidae) Towards Improved Integrated Management of Infestations in Oil Palm*. Universiti Putra Malaysia.
- Tuck, H. C., Ibrahim, Yu., & Chong, K. K. (2011). Infestations by the bagworms *Metisa plana* and *Pteroma pendula* for the period 1986-2000 in major oil palm estates managed by Golden Hope Plantation Berhad in Peninsular Malaysia. *Journal of Oil Palm Research*, 23(August), 1040–1050.
- Ulmer, B. J., Jacas, J. A., Pena, J. E., Duncan, R. E., & Castillo, J. (2006). Effect of temperature on life history of *Aprostocetus vaquitarum* (Hymenoptera: Eulophidae), an egg parasitoid of *Diaprepes abbreviatus* (Coleoptera: Curculionidae). *Biological Control*, 39, 19–25.
- Vancutsem, C., Ceccato, P., Dinku, T., & Connor, S. J. (2010). Evaluation of MODIS land surface temperature data to estimate air temperature in different ecosystems over Africa. *Remote Sensing of Environment*, 114, 449–465.
- Vaňhara, J., Muráriková, N., Malenovský, I., & Havel, J. (2007). Artificial neural networks for fly identification: A case study from the genera *Tachina* and *Ectophasia* (Diptera: Tachinidae). *Biologia, Bratislava*, 62, 462–469.
- Verheye, W. (2010). Growth and production of oil palm. In *Land Use, Land Cover and Soil Sciences. Encyclopedia of Life Support Systems (EOLSS)*. UNESCO-EOLSS Publishers.

- Wang, K., Wan, Z., Wang, P., Sparrow, M., Liu, J., & Haginoya, S. (2007). Evaluation and improvement of the MODIS land surface temperature/emissivity products using ground-based measurements at a semi-desert site on the western Tibetan Plateau. *International Journal of Remote Sensing*, 28, 2549–2565. <https://doi.org/10.1080/01431160600702665>
- Watts, M. J., & Worner, S. P. (2007). Using artificial neural networks to determine the relative contribution of abiotic factors influencing the establishment of insect pest species. *Ecological Informatics*, 3, 64–74.
- Weier, J., & Herring, D. (2000). Measuring Vegetation (NDVI & EVI). Retrieved June 4, 2018, from http://earthobservatory.nasa.gov/Features/MeasuringVegetation/measuring_vegetation_1.php
- Weisser, W. W., Volkl, W., & Hassell, M. P. (1997). The importance of adverse weather conditions for behaviour and population ecology of an aphid parasitoid. *Journal of Animal Ecology*, 66, 386–400.
- White, T. C. R. (1984). The abundance of invertebrate herbivores in relation to the availability of nitrogen in stressed food plants. *Oecologia*, 63, 90–105.
- Wood, B. J. (1971). Development of Integrated control Programs for Pests of Tropical Perennial crops in Malaysia. In C. B. Huffaker (Ed.), *Biological Control* (pp. 422–457). Boston: Springer.
- Wood, B. J., Corley, R. H. V., & Goh, K. H. (1973). Studies on the effect of pest damage on oil palm yield. In R. . Wastie & D. . Earp (Eds.), *Advances in oil palm cultivation : the proceedings* (pp. 360–379). Kuala Lumpur: The Incorporated Society of Planters.
- Wood, B. J., Liao, S. S., & Knecht, C. X. (1974). Trunk injection of systematic insecticides against bagworm, *Metisa plana* (Lepidoptera: Psychidae) on oil palm. *Oléagineux*, 29, 499–505.
- Workman, D. (2017). Palm oil exports by country. Retrieved August 17, 2017, from <http://www.worldstopexports.com/palm-oil-exports-by-country/>
- Wood, F.H., & Foot, M, A., (1981). Graphical analysis of lag in population reaction to environmental change. *New Zealand Journal of Ecology*, 4, 45-51.
- Wylie, F. R., & Speight, M. R. (2012). *Insects pests in tropical forestry* (2nd ed.). Wallingford, Oxfordshire: CAB International.
- Yang, L., Peng, L., Zhang, L., Zhang, L., & Yang, S. (2009). A prediction model for population occurrence of paddy stem borer (*Scirpophaga incertulas*), based on Back Propagation Artificial Neural Network and Principal Components Analysis. *Computers and Electronics in Agriculture*, 68, 200–206.

- Yang, Y. Z., Cai, W. H., & Yang, J. (2017). Evaluation of MODIS land surface temperature data to estimate near-surface air temperature in Northeast China. *Remote Sensing*, 9, 1–19.
- Yones, M. S., Arafat, S., Hadid, A. F. A., Elrahman, H. A. A., & Dahi, H. F. (2012). Determination of the best timing for control application against cotton leaf worm using remote sensing and geographical information techniques. *The Egyptian Journal of Remote Sensing and Space Sciences*, 15, 151–160.
- Yu, X., Guo, X., & Wu, Z. (2014) Land surface temperature retrieval from Landsat 8 TIRS-comparison between radiative transfer equation-based method, split window algorithm and single channel method. *Remote Sensing*, 6, 9829-9852.
- Zad, S.N., Zulkafli, Z. & Muharam, F.M. (2018) Satellite rainfall (TRMM 3B42-V7) performance assessment and adjustment over Pahang river basin, Malaysia. *Remote Sensing*, 10, 1-24.
- Zanuncio, J. C., Zanuncio, T. V., Lopes, E. T., & Ramalho, F. S. (2001). Temporal variations of Lepidoptera collected in an Eucalyptus plantation in the State of Goiás, Brazil. *Netherlands Journal of Zoology*, 50, 435–443.
- Zhang, W., & Zhang, X. (2007). Neural network modeling of survival dynamics of holometabolous insects: A case study. *Ecological Modelling*, 211, 433–443.
- Zhu, W., Lu, A., & Jia, S. (2013). Estimation of daily maximum and minimum air temperature using MODIS land surface temperature products. *Remote Sensing of Environment*, 130, 62–73. <https://doi.org/10.1016/j.rse.2012.10.034>