



UNIVERSITI PUTRA MALAYSIA

**ISOLATION, SCREENING AND MOLECULAR CLONING STUDIES OF
XYLANASE PRODUCING BACILLUS PUMILUS STRAINS**

AINON BINTI HAMZAH

FSMB 1997 3

**ISOLATION, SCREENING AND MOLECULAR CLONING STUDIES OF
XYLANASE PRODUCING *BACILLUS PUMILUS* STRAINS**

By

AINON BINTI HAMZAH

**Dissertation Submitted in Fulfilment of the Requirement for
the Degree of Doctor of Philosophy in the Faculty
of Food Science and Biotechnology,
Universiti Putra Malaysia**

April 1997



In loving memory of my late parents,

Hamzah bin Haji Mahmud and Nyonya binti Abdul Karim

and to

my family

ACKNOWLEDGEMENTS

First and foremost I would like to express my sincere appreciation, gratitude and thanks to Dr. Noor Aini Abdul Rashid Chairperson of the Supervisory Committee for her continual advice, guidance and encouragement in the effort to complete this thesis. I am also grateful to other members of the committee, i.e. Associate Prof. Dr. Che Nyonya A. Razak, Prof. Dr. Ismail Karim, Dr. Junainah A. Hamid and Associate Prof. Dr. Azimahtol Hawariah Lope Pihie for their support and guidance. My indebted appreciation to my former supervisors, Associate Prof. Dr. Abdullah Sipat for introducing me to the field of xylanase and also to Dr. Baharudin Ghani who had given me the full support to finish this study.

I would like to extend my gratitude to the Universiti Pertanian Malaysia for providing the facilities for this work. I am also grateful to the Faculty of Life Sciences, UKM for the facilities in carrying out part of the research work. Thank you also to all the laboratory staff of the Department of Food Science and Biotechnology, UPM; Department of Biochemistry and Microbiology, UPM; Department of Microbiology, UKM, Electron Microscopy Unit Medical Faculty, UKM; and Gene Centre, UKM for their help and cooperation are much appreciated. Special thanks to members in "Wak Dol's Lab" for their friendship and moral support during the hard times.

My special appreciations to all my colleagues in UPM and UKM who had given me all the support that I needed during the critical years and also for their lively discussions in making this venture interesting and successful.

I would like to acknowledge Dr. Warren Wakarchuk, Dr. Saran Narang and other co-workers from School of Biological Science, National Research Council (NRC) Ottawa, Canada for their guidance during my attachment at NRC.

My gratitude is extended to the Universiti Kebangsaan Malaysia for the financial support and study leave for me to pursue this study.

Last but not least is my appreciation and gratefulness to my husband, Anwar Ilyas, my children, Hani, Lan, Nina and Huda for their love that give me the encouragement to accomplish this challenging task.

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
LIST OF TABLES	ix
LIST OF FIGURES	x
ABSTRACT	xiii
ABSTRAK	xv
 CHAPTER	
I INTRODUCTION	1
Usage of xylanase	2
Objective of the research	4
II LITERATURE REVIEW	5
Hemicellulose	5
Xylans	6
Source of xylan	7
Structure of xylan	7
Properties of xylans	11
Xylanolytic enzyme system	13
Occurence of xylanolytic enzymes	14
Mode of action of xylanolytic enzymes	14
Regulation of enzyme synthesis	20
Multiplicity of xylanase	23
Different xylanolytic assays	27
Production of xylanase	29
Purification	32
Properties of xylanase	33
Cloning	37
Gene source	37
Vectors	38
Selection of target gene	38
Expression selection	40
Detection of target gene sequence	41
Xylanase sequence	41
Analysis of hemicellulase sequence	43
Importance of xylanase	45



	Page
III ISOLATION AND IDENTIFICATION OF XYLAN DEGRADING BACTERIA	45
Introduction	48
Materials and Methods	50
Samples	50
Isolation media	50
Preparation of bacterial inoculum	51
Production of xylanase	51
Determination of xylanase activity	52
Characterization	53
Morphological characters	53
Physiological and biochemical tests	54
Result	57
Morphological characters	58
Physiological characteristics	67
Biochemical reactions	73
Discussion	78
Plate assay for primary screening of xylanase activity	78
Isolation and identification of xylanase producing bacteria	81
IV PHYSIOLOGICAL STUDIES OF THE XYLANASE PRODUCING BACTERIA	83
Introduction	83
Materials and Methods	85
Bacterial strains	85
Preparation of bacterial inoculum	85
Cultivation and xylanase production	85
The effect of carbon sources on xylanase production	86
Growth curve of bacteria producing xylanase	87
Results	88
Effect of pH	88
Effect of temperature	88
Effect of agitation	94
The effect of carbon sources	94
The growth curve	101
Discussion	105

	Page
V PURIFICATION AND CHARACTERIZATION OF XYLANASE FROM <i>B. pumilus</i> PJ19	108
Introduction	108
Materials and Methods	110
Strain and growth condition	110
Enzyme assays	110
Protein determination	110
Purification of xylanase	111
Gel electrophoresis	112
Effects of substrate concentration on xylanase activity (K_m determination)	112
Isoelectric point estimation	112
The effect of pH and temperature in enzyme activity	113
To test the effect of prolonged heating on xylanase activity ..	114
Determination of time course of xylan hydrolysis	114
Inhibition by metal	115
The effect of xylose on the activity of xylanase	115
Results	116
Purification of xylanase	116
Gel electrophoresis	116
Effects of substrate concentration on xylanase activity	116
The physicochemical properties xylanase of <i>B. pumilus</i> PJ19	122
Hydrolysis of xylan	122
Effects of metal ions on the activity	133
Effects of xylose on the activity of xylanase	133
Discussion	133
VI CLONING OF THE XYLANASE GENE FROM <i>B. pumilus</i> PJ19	143
Introduction	143
Materials and Methods	145
Chemicals and enzymes	145
Microbial strains	145
General DNA techniques	145
Large scale DNA extraction	146



	Page
Quantitation of DNA	147
Partial digestion and size fractionation of genomic DNA ..	147
Vectors	148
Small scale isolation of plasmid DNA	149
Dephosphorylation of the DNA	150
Ligation of DNA	151
Preparation of competent <i>E. coli</i> cells	151
Transformation of <i>E. coli</i> cells	153
Cloning by PCR	154
Genomic DNA extraction, (method no 2)	154
Purification of plasmid DNA from alkaline lysis with Prep-A-Gene (Bio-Rad)	155
Protein sequencing	155
Oligonucleotide synthesis	156
Polymerase chain reaction (PCR)	156
Amplification of other bacterial xylanase	158
Recovery of PCR products from agarose gel	158
Cloning of PCR products into pCR TM II vector	158
Determination of the molecular size of the insert (pCR2A)	159
Colony blotting	159
Preparation of DNA probe	160
DNA dot blotting	161
Southern transfer	162
Prehybridization and hybridization	163
Antibody treatment	163
Autoradiography	164
DNA sequencing	164
Xylanase production from the pCR2A clone	169
Xylanase activity determination	170
The effect of xylan on the xylanase production from recombinant pCR2A-INV α F'	170
Codon usage, internal restriction site and DNA sequence homology	170
Results	170
Cloning of the xylanase gene from <i>B. pumilus</i> PJ19 into <i>E. coli</i>	170
Cloning by PCR	176
Genomic DNA extraction	176



	Page
N-terminal amino acids sequence of <i>B. pumilus</i> PJ19	
xylanase	176
Construction of primers for PCR amplification	178
Amplification of the xylanase gene	181
Cloning of the PCR products amplified from	
<i>B. pumilus</i> PJ19	182
Analysis of the nucleotide sequence	182
Codon usage	187
DNA sequence homology	187
Internal restriction site	193
Properties of the xylanase from clone pCR2A-INV α F'	193
Discussion	198
Is the <i>B. pumilus</i> PJ19 signal peptide really toxic to <i>E. coli</i> ?	200
Translocation of the enzyme	202
Requirement of a signal peptide	203
Sequence homology	204
VII CONCLUSIONS	205
BILIOGRAPHY	208
APPENDICES	
A Dinitrosalicylic acid (DNS) reagent, Miller et al., (1960).....	228
B Standard materials and methods for	
isolation and identification of bacteria	230
C Media	237
D Protein determination by Lowry et al., (1951)	238
E Polyacrylamide gel electrophoresis (PAGE), (Laemmli, 1970)	239
F Molecular biology techniques	242
G Buffers	244
H The effect of growth temperature on xylanase production of	
<i>B. pumilus</i>	246
I Analysis of variance on the effect of orbital rotation on	
xylanase production of <i>B. pumilus</i> strains	247
J Molecular weight determination of <i>B. pumilus</i> PJ19 xylanase	
using mass spectrometry	248
VITAE	249

LIST OF TABLES

Table	Page
1 Principal structural types found in the xylan family	12
2 Multiplicity of xylanase	25
3 Quantitative comparison of xylanase activities in various microorganisms	30
4 Specific activities of xylanase before and after purification ...	34
5 Characteristics of purified xylanases	36
6 Cloning of xylanase gene from various organisms	39
7 Sources of bacteria producing xylanase	59
8 Assay of xylanase activity from different isolates using the qualitative method	60
9 Assay of xylanase activity of different isolates using the DNS method	62
10 Results for colonial, cellular morphology, growth and biochemical tests of the isolates	65
11 Biochemical test using API 20E	74
12 Results for API 50CHB tests for <i>Bacillus</i> sp.	75
13 Computer analysis by API 50CHB (V2.0)	77
14 Identification of the isolates	79
15 Effect of carbon sources at 1 mg/ml on xylanase production ..	96
16 Effect of carbon sources at 10 mg/ml on xylanase production	96
17 The effect of the combination xylan and xylose on xylanase production by <i>Bacillus pumilus</i> strains PJ19 and P2	100
18 Summary of xylanase purification of <i>B. pumilus</i> PJ19	119
19 Effect of metals on xylanase activity	135
20 The effects of xylose on the degree of inhibition of the purified xylanase activity	136
21 Oligonucleotide primers used in PCR amplification of the <i>B. pumilus</i> PJ19 xylanase genes	157
22 The N-terminal amino acid sequences from the purified xylanase of <i>B. pumilus</i> PJ19 compared with the N-terminal region of <i>B. pumilus</i> IPO determined by DNA nucleotide sequence	177
23 Codon utilization of pre-xylanase <i>B. pumilus</i> PJ19	189
24 Restriction sites found on <i>B. pumilus</i> PJ19 xylanase gene ...	194
25 Effect of xylan on the xylanase production from clone pCR2A-INV α F'	197
26 Cellular distribution of clone pCR2A xylanase in <i>E. coli</i> INV α F'	197



LIST OF FIGURES

Figure		Page
1	Part of a hypothetical xylan molecule	9
2	Composition of (a) a typical hardwood O-acetyl-4-O-methylglucuronoxylan and (b) a typical cereal arabino-4-O-methylglucuronoxylan	10
3	A hypothetical plant heteroxylan fragment and the sites of attack by microbial xylanolytic enzymes	16
4	Hydrolysis mechanism of different xylans proposed for the xylanolytic enzymes of <i>Trichoderma reesei</i>	18
5	The xylanolytic system of <i>Cryptococcus albidus</i> : cellular localization of enzyme components and regulation of their synthesis	22
6	An alignment of the xylanase domains of β -glycanase family F and G xylanases	44
7	Clearing zone of colonies producing xylanase on YTA + RBB-xylan plate after 24 hours incubation	61
8	Zone of hydrolysis on YTA + RBB-xylan plate using the diffusion method at different times of enzyme production	61
9	Electron micrograph showing spore location	63
10	Electron micrograph of a negatively stained <i>Cellulomonas</i> sp. showing polar flagella (X 17,000)	68
11	Colonial morphology on YEA + RBB-xylan plate after 48 hours incubation. (X 20)	69
12	The effect of culture pH on xylanase production of <i>B. pumilus</i> after 24 hours incubation	89
13	The effect of culture pH on xylanase production of <i>B. pumilus</i> after 24 hours incubation	90
14	The effect of culture pH on xylanase production of <i>B. pumilus</i> strains after 48 hours incubation	91
15	Effect of temperature on xylanase production of <i>B. pumilus</i> strains after 24 hours incubation	92
16	Effect of temperature on xylanase production of <i>B. pumilus</i> strains after 48 hours incubation	93
17	The effect of speed rotation on xylanase production of <i>B. pumilus</i> strains	95
18	The effect of different concentrations of carbon sources on the xylanase production from <i>B. pumilus</i>	97



19	The effect of different concentrations of carbon sources on the xylanase production from <i>B. pumilus</i> strains	98
20	Time course of xylanase production by <i>B. pumilus</i> strains	102
21	Time course of xylanase production by <i>B. pumilus</i> strains	103
22	Time course of xylanase production by <i>B. pumilus</i> strain K52A	104
23	Chromatogram of <i>B. pumilus</i> PJ19 xylanase on CM-Sepharose	117
24	Chromatogram of <i>B. pumilus</i> PJ19 xylanase on Sephacryl S-200	118
25	SDS-PAGE of purified xylanase from <i>B. pumilus</i> PJ19	120
26	Determination of molecular weight of xylanase using SDS-PAGE	121
27	Michaelis constant (K_m) determination of xylanase using Eadie Hofstee plot using oat spelt xylan as the substrate	123
28	Michaelis constant (K_m) determination of xylanase using Eadie Hofstee plot using birchwood xylan as the substrate	124
29	Determination of pI using phast gel electrophoresis	125
30	Optimum temperature of purified xylanase	126
31	Optimum pH of purified xylanase	127
32	Effect of pH on the stability of purified xylanase	128
33	Effect of temperature on the stability of the purified xylanase...	129
34	Effect of prolonged heating on xylanase activity (45°C)	130
35	Hydrolysis of oat-spelt xylan by xylanase from <i>B. pumilus</i> PJ19	131
36	Thin-layer chromatogram of xylan hydrolysates	132
37	High-Pressure Liquid Chromatogram of the hydrolysis products of xylanase of <i>B. pumilus</i> PJ19 on oat spelt xylan	134
38	Determination of K_i for xylose inhibitor by the Dixon plot ...	137
39	Partial digestion of genomic DNA from <i>B. pumilus</i> PJ19 incubated with <i>Sau3AI</i> from 5 to 60 minutes at 37°C	171
40	Partial digestion of genomic DNA from <i>B. pumilus</i> PJ19 with <i>Sau3AI</i> after sucrose density gradient, 0-40%	172
41	Circular map of the <i>E. coli</i> expression vector pUC 18/19	173
42	Circular map of <i>E. coli</i> plasmid cloning vector pBR322	175
43	The complete nucleotide sequence and amino acid sequence for xylanase of <i>B. pumilus</i> IPO	179
44	PCR products (289 bp) of <i>B. pumilus</i> PJ19 amplified using primers from the conserved regions (5'C and 3'C) of the xylanase genes	180
45	Map of vector pCR™II	183
46	Comparison of xylanase activity by diffusion method of the supernatant obtained from the wild type <i>B. pumilus</i> PJ19 with recombinant clone pCR2A in <i>E. coli</i> INV α F'	184
47	Agarose-gel electrophoretic analysis of plasmid pCR2A-INV α F'	185
48	The nucleotide sequence of pCR2A-INV α F'	186



49	Complete xylanase sequence of <i>B. pumilus</i> PJ19	188
50	Sequence homology between <i>B. pumilus</i> PJ19 and <i>B. pumilus</i> IPO	190
51	Amino acid sequence maximum homology of xylanase of <i>B. pumilus</i> PJ19 (XYLPJ19) compared with <i>B. pumilus</i> IPO (XYLIPO)	192
52	Genomic DNA of <i>B. pumilus</i> PJ19 was digested with various restriction enzymes, incubated at 37°C for 4 hours (a)	196



Abstract of dissertation submitted to the Senate of Universiti Pertanian Malaysia in fulfillment of the requirement for the Degree of Doctor of Philosophy.

**ISOLATION, SCREENING AND MOLECULAR CLONING STUDIES
OF XYLANASE PRODUCING *BACILLUS PUMILUS* STRAINS**

By

AINON BINTI HAMZAH

April 1997

Chairperson: Dr. Noor Aini Abdul Rashid
Faculty: Food Science and Biotechnology

Cellulosic plant materials are an excellent source of hemicellulolytic microorganisms. Five strains of *Bacillus pumilus* PJ19, P2, K52A, K51 and K5B, two strains of *B. subtilis* B2 and PJ18 and one *Cellulomonas* sp. which produced endoxylanase (1,4- β -D-xylan xylanohydrolase, EC 3.2.1.8) have been isolated locally from plant materials. *B. pumilus* PJ19 produced the highest xylanase activity when grown in shake flask in yeast tryptone broth (YTB) at 200 rpm, 37°C which yielded activity of 265 U/ml. The enzyme was induced in Dubois media by the addition of xylan as carbon source and was repressed by xylose, glucose, fructose, maltose and sucrose. *B. pumilus* strain PJ19 and K5B showed maximum enzyme activity when grown in YTB (pH 7.2), 37°C after 36 hours, P2 (28 hours), while K51 and K52A after 32 hours incubation. The xylanase from *B. pumilus* PJ19 was purified to homogeneity by ammonium sulphate precipitation and gel filtration of CM-Sepharose and Sephacryl S-200. The molecular weight of the purified xylanase was estimated to be 23,000 D by sodium dodecyl sulphate-polyacrylamide gel electrophoresis and



22,515 D by mass spectrometry. The isoelectric point of the enzyme was 9.3. The optimum pH and temperature for hydrolysis of oat spelt xylan were 6.5 and 60°C, respectively. The enzyme was stable at a pH range of 7.5 to 8.5 and at a temperature of up to 45°C but lost 50% of its activity at 58°C after 10 minutes of incubation. The purified enzyme had a K_m of 1.42 mg/ml and V_{max} of 107 $\mu\text{mol}/\text{min}/\text{mg}$ for oat spelt xylan and K_m of 2.15 mg/ml and V_{max} of 29.22 $\mu\text{mol}/\text{min}/\text{mg}$ for birchwood xylan. The major end products of oat spelt xylan hydrolysis were xylobiose, xylotriose and higher oligosaccharides while for birchwood xylan were xylotriose with some xylobiose determined by thin layer chromatography and high performance liquid chromatography. Xylose was not produced as a product of hydrolysis and trans xylosidation was detected. The activity of the enzyme was enhanced in the presence of Mg^{2+} , Ca^{2+} and K^+ but was inhibited by EDTA, Cu^{2+} , Ag^+ , Zn^{2+} , Fe^{2+} and Hg^{2+} . The enzyme was competitively inhibited in the presence of xylose with K_i of 1.98 mM. A complete DNA sequence of the xylanase gene was amplified by polymerase chain reaction and cloned into *E. coli* INV α F' using pCRII cloning vector. The complete DNA sequence was also determined. The structural xylanase gene which started from an ATG initiation codon, consists of an open reading frame of 684 bp, which encoded 202 amino acid residues. The molecular weight of the xylanase was estimated from the amino acid composition to be 22,474 D and is in agreement with the results obtained from SDS-PAGE of the purified xylanase. The xylanase was expressed constitutively by the cloned gene in the absence of xylan. The enzyme was located primarily in the cytoplasm probably because of the incompatibility of the Gram-positive signal peptide in *E. coli* to direct the enzyme extracellularly as in the donor strain.



Abstrak disertasi yang dikemukakan kepada senat Universiti Pertanian Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah.

**PEMENCILAN, PENABIRAN DAN KAJIAN PENGKLONAN MOLEKUL
STRAIN-STRAIN *BACILLUS PUMILUS* PENGHASIL XILANASE**

Oleh

AINON HAMZAH

April 1997

Pengerusi: Dr Noor Aini Abdul Rashid
Fakulti: Sains Makanan dan Bioteknologi

Bahan tumbuhan berselulosa merupakan sumber terbaik bagi mikroorganisma hemiselulolitik. Lima strain *Bacillus pumilus* PJ19, P2, K52A, K51 dan K5B, dua strain *B. subtilis* B2 dan PJ18, dan satu *Cellulomonas* sp. yang menghasilkan endoxilanase (1,4- β -D-xylan xilanohidrolase, EC 3.2.1.8) telah dipencilkan di Universiti Pertanian Malaysia dari bahan tumbuhan tempatan. *B. pumilus* PJ19 menghasilkan aktiviti xilanase tertinggi apabila dihidupkan dalam kelalang goncangan kaldu yis tripton (YTB) pada 200 rpm, 37°C yang menghasilkan aktiviti pada 265 U/ml. Enzim diaruh dalam media Dubois dengan penambahan xilan sebagai sumber karbon dan direncat oleh xilosa, glukosa, fruktosa, maltosa dan sukrosa. *B. pumilus* strain PJ19 dan K5B, menunjukkan aktiviti enzim yang maksimum apabila dihidupkan dalam YTB (pH 7.2), 37°C selepas penderaman selama 36 jam, P2 (28 jam), sementara K51 dan K52A selepas penderaman 32 jam. Xilanase daripada *B. pumilus* PJ19 ditulenkan sehingga homogen dengan pemendakan amonium sulfat dan turasan gel menggunakan CM-Sepharose dan Sephacryl S-200. Berat molekul xilanase tulen dianggarkan 23,000 D dengan



menggunakan elektroforesis gel natrium dodesil sulfat-poliakrilamida, dan 22,515 D dengan kaedah spektrometri jisim. Takat isoelektrik enzim ialah 9.3. Hidrolisis xilan oat spelt adalah optimum pada pH 6.5 dan suhu 60°C. Enzim didapati stabil pada julat pH 7.5-8.5 dan suhu sehingga 45°C, tetapi 50% aktiviti hilang pada suhu 58°C selepas 10 minit pengeraman. Xilanase tulen mempunyai nilai K_m 1.42 mg/ml dan V_{max} 107 $\mu\text{mol}/\text{min}/\text{mg}$ untuk xilan oat spelt dan K_m 2.15 mg/ml dan V_{max} 29.22 $\mu\text{mol}/\text{min}/\text{mg}$ untuk xilan birchwood. Hasil akhir utama hidrolisis xilan oat spelt seperti yang ditentukan dengan kaedah kromatografi lapisan nipis dan kromatografi cecair prestasi tinggi ialah xilobiosa, xilotriosa dan oligosakarida tinggi sementara bagi xilan birchwood pula ialah xilotriosa dengan sedikit xilobiosa. Hidrolisis tidak menghasilkan xilosa kerana berlakunya trans xilosidasi. Aktiviti enzim dapat ditingkatkan dengan kehadiran ion Mg^{2+} , Ca^{2+} dan K^+ tetapi direncat oleh EDTA, Cu^{2+} , Ag^+ , Zn^{2+} , Fe^{2+} dan Hg^{2+} . Aktiviti enzim direncat secara kompetitif oleh kehadiran xilosa pada K_i 1.98 mM. Penjujukan DNA lengkap untuk gen xilanase digandakan secara tindak balas rantai polimerase dan diklonkan ke dalam *E. coli* INV α F' dengan menggunakan vektor pengklonan pCRII. Penjujukan DNA yang lengkap telah juga dilakukan. Gen struktur xilanase yang bermula dari kodon ATG terdiri daripada kerangka pembacaan terbuka 684 bp yang mengekodkan 202 sisa asid amino. Berat molekul xilanase tulen yang dianggarkan melalui komposisi asid amino ialah 22,474 D yang mana bersesuaian dengan nilai yang diperolehi daripada teknik natrium dodesil sulfat dan spektrometri jisim. Gen yang diklonkan menghasilkan xilanase secara konstitutif tanpa kehadiran xilan. Kebanyakan dari xilanase itu dikesan berada di sitoplasma. Ini mungkin disebabkan oleh ketidaksesuaian isyarat peptida gram-positif dalam *E. coli* untuk memberi arahan supaya enzim dibawa keluar sel seperti dalam strain penderma.

CHAPTER I

INTRODUCTION

In recent years, there has been renewed interest in the utilization of plant materials as a source of fuel and chemicals. It is an effort that can partially reduce our total dependence on fossil fuels. An advantage of using plant material is that it is a renewable resource. Total energy content in plant biomass is estimated to be equivalent to 640 billion tonnes of oil (Coughlan, 1985). Furthermore, globally there is an abundance of agricultural residues or agro-wastes such as sugar bagasse, empty fruit bunch from oil palm, coconut husks, paddy straw, wheat straw and forest wastes which accumulate in large quantities every year. From an economic standpoint, it is essential that all of the plant material residues be utilized, or it will cause environmental deterioration and loss of potentially valuable resources.

The main component of plant materials with potential as a source of chemicals is polysaccharides which consists of cellulose (40-60%) with lesser but significant amount of hemicellulose (20-30%) and lignin (15-30%) (Schuerch, 1975). Unlike cellulose, which occurs as homopolymer of glucose, hemicellulose does not have a homogeneous chemical composition. Hemicellulose is a heterogeneous polymer of glucose, mannose, galactose, xylose, arabinose, and uronic acid (Lynch, 1987). Both xylan and cellulose are very difficult to solubilize owing to the β -1,4-glucosidic linkage.

The degradation of cellulosic and hemicellulosic materials occurring in nature is carried out mainly by microorganisms. They include fungi and bacteria, aerobes and anaerobes, mesophiles and thermophiles, and occupy a variety of habitats. One



of the approaches in handling plant biomass residues is to degrade the complex polysaccharides into simple sugars, alcohols, and other useful products.

Technically, there are three methods which can be used for degrading the lignocellulose components or polysaccharides. Firstly is the physical treatments method which involves primarily ball milling (Mandels et al., 1974), high pressure steaming (Saddler et al., 1982) and irradiation (Han & Ciegler, 1983). These processes have their drawbacks; that is they are not cost-effective and unable to remove noncellulosic substances (Andren & Nystrom, 1976). Secondly, is the chemical treatment method which employ alkali and dilute acids such as sulphuric or hydrochloric acids. Likewise, these treatments also have certain disadvantages. These include formation of hard scale on heated surfaces when $\text{Ca}(\text{OH})_2$ is used, the need for expensive corrosion-resistant equipment, contamination of sugars produced by toxic byproducts and often chemicals used are expensive (Marsden & Gray, 1986). Although acid hydrolysis is faster, it can produce toxic compounds which can hinder microbial fermentation. Thirdly, is the employment of biological agents involving enzyme hydrolysis. In fact, this is one of the most important means of converting cellulose or hemicellulose to monosaccharides, which can then be used as fermentation feedstock.

An economic evaluation on the conversion of lignocellulosic materials for ethanol production indicated that the most expensive method is steam explosion coupled with enzyme hydrolysis while the cheapest is acid treatment coupled with enzyme hydrolysis (Parisi & Parisi, 1989). The development of an efficient enzymic hydrolysis offers new prospects for treating hemicellulosic wastes into useful products.

Usage of xylanase

The most widely used enzymes for the hydrolysis of lignocellulose are xylanase and cellulase. Although xylanase enzyme systems for the hydrolysis of xylans have been studied extensively in the past (Dekker & Richards, 1976, Kubicek, 1981), they remain less recognized compared to cellulase systems. This is due to the fact

that cellulose is composed of only D-glucose and can be found in all plants, while the composition of xylan is more complex and it also varies from plant to plant (Biely, 1985; Joseleau et al., 1992). It is also reflected in the literature as there are many reports on enzyme hydrolysis of cellulosic materials to glucose using cellulase from different organisms (Gaden et al., 1976, Gong et al., 1979, Bisaria & Ghose, 1981). However, xylanolytic enzyme systems deserve the same attention as the cellulolytic systems because their biotechnological potential is equally important. This is so as there is a considerable amount of xylan present in agricultural residues, and the conversion of xylan to useful products will strengthen the overall economics. Furthermore, xylan will provide an alternative for energy production from renewable resources.

Enzymes can be derived from various sources, but the use of microbial enzymes for industrial hydrolysis of lignocellulose is advantageous because of the high specificity of the enzyme reactions, the mildness of the reaction conditions, and the absence of substrate loss due to chemical modifications (Wong et al., 1988). That is why this study is focused on the potential use of microbial xylanase. Xylanase is the enzyme for the degradation of xylan, the major component of hemicellulose. According to Gilbert & Hazlewood (1993), xylanase could be exploited in agricultural and industrial processes as listed below:

- i. Pre-treatment of forage crops and other cellulosic biomass with cellulase and xylanase to improve the nutritional quality and digestibility of ruminant feeds or to facilitate composting.
- ii. Enzymic saccharification of agricultural, industrial and municipal wastes to provide sugar syrups for human or animal consumption or for the production of fine chemicals through industrial fermentations.
- iii. Enzymic digestion of industrial wastes as an alternative to landfill deposition.
- vi. Addition of cellulase and xylanase to cattle, pig and poultry cereal-based diets to elicit a significant improvement in nutrient utilization through the hydrolysis of barley β -glucan and arabinoxylans.

- v. Xylanase pretreatment of paper pulps to remove certain xylan components and reducing dependence on chlorine in the brightening process.

Xylanases are also applied to facilitate the bleaching of Kraft pulps or to improve fiber properties (Grabski & Jeffries, 1991). Xylanase can also be used in clarification of juices, preparation of dextrans for use as food thickeners, production of fluids and juices from plant materials, in processes of liquefaction of coffee mucilage for the manufacture of liquid coffee, adjustment of wine characteristics and enhancement of pigment astaxanthin (3,3'-dihydroxy-4,4'-diketo- β -carotene) extraction (Woodward, 1984; Wong et al., 1988; Khasin et al., 1993 and Filho, 1994). The xylanase from *Streptomyces* E86 has been used in the production of xylobiose from commercial hardwood xylan (Kusakabe et al., 1975). Large amounts of hemicellulose which are present in wastes from the pulp and paper industry are in a form that cannot be buried and require expensive cost for disposal. With the advent of the new technology, there is increased interest to use xylan degrading enzymes to reduce these costs. Furthermore, the xylan may be used as raw material to produce xylose that can be used as feedstock or in fermentation of ethanol with or without formation of intermediate xylulose (Gruninger & Fiechter, 1986; Dey et al., 1992 and Filho, 1994).

Objective of the research

With respect to the vast potential in the industrial use of xylanase, this research was done with the ultimate goal of isolating xylanase-producing bacteria from soil. Xylanase is synthesized in large amounts by phytopathogenic soil microorganisms, which are able to degrade plant materials. Bacteria is chosen over fungi due to its fast growth and its ease for scaling up. Physiological and biochemical studies of the bacteria producing xylanase will be carried out to optimize the enzyme production. Through genetic manipulation, superior strains which are able to synthesize enzymes rapidly can be produced. The sequence of the enzyme will show the similarity or difference with other bacterial xylanase.

CHAPTER II

LITERATURE REVIEW

Cellulose, hemicellulose and lignin are the three major components of plant cell wall. Cellulose, the major component of plant cell wall, consists of a linear polymer of anhydroglucose units linked by β -1,4-glucosidic bonds, and do not occur in pure form in any natural resources (Bisaria & Ghose, 1981). In nature, it is always associated with other polysaccharides, such as starch, pectin, lignin and a variety of hemicelluloses. Cellulose has been thoroughly studied by many researchers (Nisizawa, 1973; Bisaria & Ghose, 1981; Coughlan, 1985 and Enari, 1987) and the mechanisms of hydrolysis has been well understood.

As this study is concerned with xylan and the xylanolytic enzyme systems, a comprehensive literature survey on the subjects is presented in this chapter. A brief literature review as background information on specific topics will be presented in each subsequent chapter.

Hemicellulose

Hemicelluloses are heteropolymers composed of various pentoses and hexoses of varying proportions, depending on their botanical origin. Introduced in 1891 by Schulze, they refer to the easily hydrolyzable parts of the cell wall. According to Timell (1964), hemicelluloses are polysaccharides of low molecular weights that occur in plant tissues together with cellulose, and which can be extracted from the original or the delignified material by using alkali or water. Hemicelluloses may

also be defined as polysaccharides present in the cell wall and intercellularly, that can be extracted from higher land-plant lignified tissues by alkali treatment (Wilkie, 1979). Certain carbohydrates in cereal endosperm, namely, non-starch polysaccharides that are described as cereal gums or pentosans are also classified as hemicellulose (Wilkie, 1979). However, they were later redefined to include only plant cell wall polysaccharides that bind noncovalently to cellulose. This definition is based on the chemical properties of these polysaccharides that are relatively easy to measure and are related to their proposed biological formation (Kennedy & White, 1988).

The hemicelluloses are one of the major constituents of lignocellulosic materials, and may comprise 30-40% of the total plant cell carbohydrate. They are important structural components and occur in close association with lignin and cellulose (Zimmermann, 1989; Williams, 1989).

Hemicellulose does not have a homogeneous chemical composition and is a heterogeneous polymer of different types of sugars in the backbone chain and in the side chain or appendages. These may be D-glucose, D-galactose, D-mannose, D-xylose, L-arabinose, D-glucuronic acid, D-galacturonic acid, or sometimes esters of O-acetyl, feruloyl coumaryl linked via L-arabinose residues to the backbone (Woodward, 1984; Puls & Poutanen, 1989). The composition of hemicelluloses in plants can be influenced by various factors such as growth, maturation, nature of soil, climate, length of the day, geographical location, and type of fertilizer used (Wilkie, 1979). However, the main component of hemicellulose is xylan.

Xylans

Xylans form a major component of hemicellulose comprising β -1,4-linked polymer of D-xylose as a backbone. They are the second biopolymers after cellulose, present in nature in large amounts. β -1,4-xylans are mainly found in secondary walls of plants, the major component of woody tissue (Timmel, 1967). The amount of xylan varies in different plants, from 35% of the dry weight of birchwood to as little as 7% in some gymnosperms (Whistler & Richards, 1970). Since agricultural

wastes and plant residues are generated on a massive scale each year, xylan represents a considerable potential source of fuel, single-cell protein, solvent and other chemicals by the selective use of specific fermentative microorganisms (Biely, 1985). This also can contribute to the elimination of agricultural wastes. The efficient exploitation of xylans as a source of industrial raw materials requires understanding of the enzyme systems that affect their conversion.

Source of xylan

Xylans are present in many types of terrestrial plants ranging from monocots to dicots and are mainly located in secondary plant cell walls (Timell, 1967). They are the major and minor components of mature cell walls of woody tissue and the primary wall in dicots, respectively (Timell, 1967; McNeil et al., 1984) and are also present as major components of monocot primary cell walls (Dey & Brinson, 1984). Xylan is also present in the primary walls of growing cells (Joseleau & Barnound, 1974), cell walls of the aleurone layer (McNeil et al., 1975) and sometimes in seeds and bulbs (Shaw & Stephen, 1966). In general, xylan is the major hemicellulose in wood from angiosperms but is less abundant in wood from gymnosperms; it accounts for approximately 15% to 30% and 7% to 12% of the total dry weight, respectively (Whistler & Richards, 1970).

In perennial or cereals, xylans are present in wheat straw, wheat leaf, oat straw, corn cobs, wheat bran, maize fiber and rice husk. Xylan is also present in woody plants, such as beechwood, pine, larch, oak, hemlock and spruce (Aspinall, 1959). As such, Malaysian woody plants are an equally good source of xylans.

Structure of xylan

Xylans from land plants are heteropolysaccharides that constitute a large group of related hemicellulose with a great variability in their structures. There is a relationship between the chemical structure of the xylan to their botanical origin and their cytological localization. Therefore, this results in the formation of several different polymers of xylans of related structures.

Most xylans from terrestrial plants have molecular structural which are branched and the degree of branching varies depending on the source (Biely, 1985) and methods of extraction (Wilkie, 1979). The main chain of xylan is a linear backbone structure composed of D-xylose units linked by β -1,4-xylopyranosyl which may be substituted, depending on the origin of xylan (Dey & Brinson, 1984; Wong et al., 1988; Puls & Poutanen, 1989). Substitutions include: acetylation at C-2 or C-3 of the xylose units, α -1,2-linked glucuronic or 4-O-methylglucuronic acid groups, α -1,3-linked arabinofuranosyl units, and ferulic or coumaric acids esterified to O-5 of arabinose (Coughlan, 1992) as shown in Figure 1.

Very few unbranched linear xylan homopolysaccharides from land plants have been isolated. The best known example is xylan from esparto grass (Aspinall, 1959), which is normally used as a model for chemical and physical studies. It is composed of a straight chain of β -1,4-linked to D-xylopyranosyl residues. Water soluble, linear xylan, has also been isolated from tobacco stalks which consisting of almost pure β -1,4-D-xylan without any other sugar components (Eda et al., 1976).

All xylans from terrestrial plants (hardwoods, softwoods and grasses) are linked by β -1,4-D-xylopyranosyl residues (Timell, 1967). However, those isolated from the marine algae are β -1,3-linked (Dekker & Richards, 1976), while xylans from sea-weeds, such as *Rhodymenia palmata*, contain unbranched chains of about 80% of β -1,4-linkages and 20% of β -1,3-linkages (Kato & Nevins, 1984). In some species of the Chlorophyceae and the Rhodophyceae where cellulose is absent, the xylans appeared to be in the form of only β -1,3-D-xylopyranose residues which form a highly crystalline fibrillar material (Aspinall, 1959; Joseleau, et al., 1992).

Hardwoods contain acetylated xylan, which made up 10-35% of the total dry weight with a degree of polymerization (DP) of 150 to 200 (Puls & Shuseil, 1993). Seven out of ten xylose units are acetylated on C-2 and/or C-3 position (Puls & Poutanen, 1989; Gilbert & Hazlewood, 1993) and the typical structure is shown in Figure 2.2a. For example, birch xylan contains more than 1 mole of acetic acid per 2 mole of D-xylose. The presence of acetyl groups makes the xylan