



UNIVERSITI PUTRA MALAYSIA

**CAROTENE EXTRACTION FROM CRUDE PALM OIL
USING SYNTHETIC ABSORBENTS**

RAZAM ABD. LATIP

FSMB 2000 7

**CAROTENE EXTRACTION FROM CRUDE PALM OIL
USING SYNTHETIC ABSORBENTS**

By

RAZAM ABD. LATIP

**Degree of Master of Science in the Faculty of
Food Science and Biotechnology
Universiti Putra Malaysia**

September 2000



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Master of Science

**CAROTENE EXTRACTION FROM CRUDE PALM OIL USING
SYNTHETIC ADSORBENTS**

By

RAZAM ABD. LATIP

September 2000

Chairman : Assoc. Prof. Badlishah Sham Baharin, M.Sc.

Faculty : Food Science and Biotechnology

Crude palm oil (CPO) has the highest content of carotenoids (500 - 700 ppm), a precursor of vitamin A. The only commercially viable method, so far, is transesterification followed by phase separation. However, the edible oil used as raw material has to be converted to methyl ester, therefore destroying the oil into non-edible product.

The present study on carotene extraction from CPO focussed on adsorption using synthetic adsorbent followed by solvent extraction. By this method, the carotene can be recovered without destroying the oil therefore it can be used for food applications. The

objectives of this study were mainly to find out the suitable adsorption process that selectively extracts the carotene from CPO and to determine the effect on CPO quality after going through this process.

Based on the studies conducted, it was found that the synthetic adsorbent SP850, SP825, HP20, Relite Exa 32 and Relite Exa 50 were capable of adsorbing carotene from CPO. The percentage of carotene extracted varied from 10 to 80% with the carotene concentration ranging from 1000 to 20,000 ppm depending on the process conditions. Combinations of adsorbent HP 20 and SP 850 slightly increased the percentage of carotene extracted. Adsorbent/CPO ratio of 4 was most suitable for this process for optimum recovery and concentration of carotene. The minimum adsorption time required was 0.5 hr. The IPA extraction time was determined based on the final carotene concentration required. The suitable temperature for adsorption and solvent extraction process was at 40°C. There is no significant difference on the percentage of carotene extracted and carotene concentration between with and without agitation during IPA extraction process.

The quality of CPO after going through the carotene extraction process slightly deteriorated in terms of moisture content, impurities, peroxide value (PV), anisidine value (AV), discriminant function (DF) and deterioration of bleaching index (DOBI). However, changes in the chemical properties of the oil such as triglyceride (TG)

carbon number and fatty acid composition (FAC) and it can be refined to produce refined bleached deodorized palm oil (RBDPO) that is able to meet Palm Oil Refinery Association of Malaysia (PORAM) standard specifications.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGEKSTRAKAN KEROTIN DARIPADA
MINYAK SAWIT MENTAH**

OLEH

RAZAM ABD. LATIP

September 2000

Pengerusi : Prof. Madya Badlishah Sham Baharin, M.Sc.

Fakulti : Sains Makanan dan Bioteknologi

Minyak sawit mentah (CPO) mengandungi *carotenoids* pemula untuk Vitamin A yang paling tinggi (500-700 ppm). Setakat ini hanya kaedah *transesterification* diikuti dengan pengasingan berperingkat telah dimajukan secara komersial. Walaubagaimanapun minyak yang digunakan sebagai bahan mentah akan diubah kepada *methyl ester*, mengakibatkan pemusnahan minyak kepada bahan bukan makanan.

Kajian pengekstrakan kerotin dari CPO ini menumpukan penggunaan bahan penjerap sintetik melalui proses penjerapan diikuti dengan pengekstrakan pelarut. Kaedah ini akan mengekstrak kerotin tanpa merosakkan minyak, oleh itu ianya boleh digunakan untuk

penggunaan bahan makanan. Tujuan utama kajian ini dijalankan adalah untuk menentukan sistem penjerapan yang sesuai dimana ianya akan menjerap hanya kerotin daripada CPO dan menentukan kesan ke atas kualiti CPO selepas menjalani proses tersebut.

Berdasarkan daripada kajian yang dijalankan, didapati bahan penjerap SP850, SP825, HP20, Relite Exa 32 dan Relite Exa 50 berkemampuan untuk menjerap kerotin dari CPO. Peratus pengekstrakan kerotin adalah diantara 10% hingga 80% dimana kepekataannya adalah diantara 1,000 hingga 20,000 ppm bergantung kepada keadaan proses. Peratus pengekstrakan kerotin bertambah sedikit dengan menggunakan gabungan bahan penjerap HP 20 dan SP 850. Nisbah bahan penjerap/CPO pada kadar 4 adalah keadaan proses yang paling sesuai untuk mendapatkan kepekatan dan kadar peratusan pengekstrakan kerotin yang optimum. Masa penjerapan minima yang diperlukan adalah pada 0.5 jam. Masa pengekstrakan IPA ditentukan berdasarkan kepada kepekatan kerotin yang dikehendaki. Suhu yang sesuai bagi proses penjerapan dan pengekstrakan pelarut adalah pada 40°C. Peratus pengekstrakan kerotin dan kepekatan kerotin tidak berubah samada dengan menggunakan pengacau atau tidak semasa proses pengekstrakan IPA.

Kualiti CPO selepas menjalani proses pengekstrakan kerotin didapati mengalami sedikit penurunan terutamanya kandungan kelembapan, bendasing, nilai pengoksidaan (PV), nilai anisidine (AV),

fungsi diskriminen (DF) dan indeks penurunan pelunturan (DOBI). Walaubagaimanapun tiada perubahan di dalam bahan-bahan kimia minyak itu seperti TG dan FAC, dan ianya boleh menjalani proses penjernihan untuk menghasilkan minyak tulin, terluntur dan ternyabau (RBDPO) yang memenuhi spesifikasi Piawaian Persatuan Pengilang-Pengilang Kelapa Sawit Malaysia (PORAM).

ACKNOWLEDGEMENTS

With great thanks to Almighty Allah S. W. T and His mercy for giving me life and the opportunity to write this thesis as required.

I would like to take this opportunity to express my most sincere appreciation and deepest gratitude to Assoc. Professor Badlishah Sham Bin Baharin, the Chairman of my Supervisory Committee. Without his valuable guidance, advice, suggestions and constructive criticisms, I would not have been able to present my work and in the preparation of this thesis.

I offer,

Committee: Professor Dr. Yaakob Bin Che Man and Assoc. Prof. Dr. Russly Bin Abd. Rahman for their supervision, support and comments.

Special thanks and appreciation to my friends, staff, Pn. Sadikom and En. Radhakrishnan, for their assistance in preparing this thesis. I would like to acknowledge Golden Hope Plantations Berhad for approving my M.Sc studies and using the laboratory facilities in Golden Jomalina Food Industries Sdn. Bhd.

I also owe my special thanks to my beloved wife Aziah Lajis for her inspiration, moral encouragement, support, patience and sacrifices which had helped me in undertakings and completing this research study.

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL SHEETS	x
DECLARATION FORM	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xx
LIST OF PLATES	xxiii
CHAPTER	
I. GENERAL INTRODUCTION	1
II. REVIEW OF LITERATURE	4
Palm Oil	4
Palm Oil Production and Trade	4
Composition of Palm Oil	6
Palm Oil Refining	11
The Uses of Palm Oil	19
Carotenoids	19
Other	21
Nomenclature and Structure	21
Applications of Carotenes	28
Various Processes of Carotene Recovery from Palm Oil	34
Saponification	34
Extraction Using Absorbents	34
Solvent Extraction	41
Transesterification	41
Supercritical Fluid Technology	50
III. EVALUATION OF DIFFERENT TYPES OF SYNTHETIC ADSORBENTS FOR CAROTENE EXTRACTION FROM CRUDE PALM OIL	53
Introduction	53
Materials and Methods	54
Materials	54
Adsorption	55
Solvent Extraction	55
Analysis	55

Calculations for the Percentage of Carotene Extracted	56
Statistical analysis	56
Results and Discussion	57
Evaluation of Different Types of Adsorbents	57
Combination of Adsorbents (HP 20 and SP 850)	60
Adsorbent/CPO Ratio	62
Summary	66
IV. THE EFFECT OF ADSORPTION AND SOLVENT EXTRACTION PROCESS ON THE PERCENTAGE OF CAROTENE EXTRACTED FROM CRUDE PALM OIL	67
Introduction	67
Materials and Methods	67
Materials	67
Adsorption	68
Solvent Extraction	68
Adsorption Time	68
IPA Extraction Time	68
Temperature of Adsorption and Solvent Extraction Process	69
Agitation during IPA Extraction	69
Adsorbent Lifespan	70
Analysis	70
Calculations for the Percentage of Carotene Extracted	70
Statistical analysis	70
Results and Discussion	70
Adsorption Time	70
IPA Extraction Time	71
Temperature of Adsorption and Solvent Extraction Process	74
Agitation during IPA Extraction	76
Adsorbent Lifespan	77
Summary	80
V. THE EFFECT OF CAROTENE EXTRACTION PROCESS ON CRUDE PALM OIL QUALITY, CAROTENE COMPOSITION AND CAROTENE STABILITY DURING STORAGE	82
Introduction	82
Materials and Methods	82
Materials	82
Adsorption	82
Solvent Extraction	83
Degumming	83
Bleaching	83
Deodorization	83
Determination of Total Carotene	84
Triglyceride (TG) Carbon Number Composition	84
Fatty Acid Composition (FAC)	84

High Performance Liquid Chromatography (HPLC)	
Analysis of Carotene	85
Results and Discussion	85
Analysis on CPO Quality	85
Pilot Plant Refining	89
Triglyceride (TG) Carbon Number Composition	90
Analysis on Fatty Acid Composition (FAC)	92
Analysis on Carotene Compositions by HPLC	93
Stability of Carotene during Storage	93
Summary	96
VI. SUMMARY, CONCLUSION AND RECOMMENDATION	97
Summary	97
Conclusion and Recommendation	99
BIBLIOGRAPHY	101
APPENDICES	108
BIOGRAPHICAL SKETCH	137

LIST OF TABLES

Table	Page
2.1 World Palm Oil Production 1960 – 1997 (1,000 metric ton)	6
2.2 World Major Exporters of Palm Oil 1960 – 1997 (1,000 metric tonnes)	7
2.3 Fatty Acid Composition of Palm Oil	8
2.4 Triglyceride Composition of Crude Palm Oil	9
2.5 Solid Fat Content of Crude Palm Oil	10
2.6 Refining of Crude Palm Oil: Unit Processes	13
2.7 Desirable Quality of Pretreated and RBD Palm Oil from the Factory	16
2.8 Desirable Quality of Freshly Produced Intermediate and Final Products in Alkaline Refining of Palm Oil	18
2.9(a) Typical Carotenoids Content of Selected Fruits and Vegetables (mcg/100 g portion)	23
2.9(b) Typical Carotenoid Content of Selected Fruits and Vegetables (mcg/100 g portion)	24
2.10 End Group Designation of Carotenes	25
2.11 Aromatic Synthetic Adsorbents – HP Series	36

2.12	Aromatic Synthetic Adsorbents – SP 800 Series	38
2.13	Aromatic Synthetic adsorbents – SP 200 Series	40
3.1	Physical Properties of the Synthetic Adsorbents	54
3.2	Effect of Different Types of Adsorbents on the Percentage of Carotene Extracted	58
3.3	Effect of Combination of Adsorbents on the Percentage of Carotene Extracted	61
3.4	The Effect of Adsorbent/CPO Ratio on the Percentage of Carotene Extracted	63
3.5	Selection Matrix for Adsorbent/CPO Ratio	65
3.6	The Effect of Combination of Adsorbents on The Percentage of Carotene Extracted (First Run)	110
3.7	The Effect of Combination of Adsorbents on the Percentage of Carotene Extracted (Second Run)	111
3.8	The Effect of Combination of Adsorbents on the Percentage of Carotene Extracted (Third Run)	112
3.9	The Effect of Adsorbent/CPO Ratio on the Percentage of Carotene Extracted (First Run)	113
3.10	The Effect of Adsorbent/CPO Ratio on the Percentage of Carotene Extracted (Second Run)	114
3.11	The Effect of Adsorbent/CPO Ratio on the Percentage of Carotene Extracted (Third Run)	115
4.1	The Effect of Adsorption Time on the Percentage of Carotene Extracted	72
4.2	The Effect of IPA Extraction Time on the Percentage of Carotene Extracted	73

4.3	The Effect of Temperature of Adsorption and IPA Extraction Process on Percentage of Carotene Extracted	75
4.4	The Effect of Agitation during IPA Extraction on Percentage of Carotene Extracted	78
4.5	The Adsorbent Lifespan and the Effect on the Percentage of Carotene Extracted	79
4.6	The Effect of Adsorption Times on the Percentage of Carotene Extraction (First Run)	116
4.7	The Effect of Adsorption Times on the Percentage of Carotene Extraction (Second Run)	116
4.8	The Effect of Adsorption Times on the Percentage of Carotene Extraction (Third Run)	117
4.9	The Effect of IPA Extraction Time on the Percentage ... of Carotene Extraction (Adsorbent/CPO Ratio : 1.5)	117
4.10	The Effect of IPA Extraction Time on the Percentage ... of Carotene Extraction (Adsorbent/CPO Ratio : 2.0)	118
4.11	The Effect of IPA Extraction Time on the Percentage ... of Carotene Extraction (Adsorbent/CPO Ratio : 3.0)	118
4.12	The Effect of IPA Extraction Time on the Percentage ... of Carotene Extraction (Adsorbent/CPO Ratio : 5.0)	119
4.13	The Effect of IPA Extraction Time on the Percentage ... of Carotene Extraction (Adsorbent/CPO Ratio : 6.0)	119
4.14	The Effect of IPA Extraction Time on the Percentage ... of Carotene Extraction (Adsorbent/CPO Ratio : 7.5)	120
4.15	The Effect of Temperature during Adsorption Process and IPA Extraction of Percentage of Carotene Extraction (First Run)	120

4.16	The Effect of Temperature during Adsorption Process and IPA Extraction of Percentage of Carotene Extraction (Second Run)	121
4.17	The Effect of Temperature during Adsorption Process and IPA Extraction of Percentage of Carotene Extraction (Third Run)	121
5.1	Quality of Feed Crude Palm Oil (CPO) and CPO from Isopropanol (IPA) Fractions	87
5.2	Crude Palm Oil (CPO) Grade Based on Deterioration of Bleachability Index (DOBI) and Discriminant Function (DF) Values	89
5.3	Quality of CPO from IPA Fractions and RBD Palm Oil using 1%, 2% and 3% of Bleaching Earth	91
5.4	Triglyceride Carbon Number Composition of Feed CPO, CPO from IPA Fractions and RBD Palm Oil	92
5.5	Fatty Acid Composition of Feed CPO, CPO from IPA Fractions and RBD Palm Oil	93

LIST OF FIGURES

Figure		Page
2.1	Triglyceride	7
2.2	Flow Diagram of Alkali-Refining and Physical Refining Processes for Palm Oil	12
2.3	Basic Structure of Carotenoids	22
2.4	Carotenoid Structure and Numbering of the Parent Carotene (Steam Name)	22
2.5	End Group Designation of Carotenes	25
2.6	Example of Carotenes	26
2.7	Example of Carotene after changes in Hydrogenation Level	27
2.8	Example of Oxygenated Carotenoids	29
2.9	Major Commercial Uses of β -carotene	30
2.10	Structural Formulate of Some Common Carotenoids and Vitamin A	32
2.11	Chemical Structure of the Synthetic Adsorbent HP and SP Series	36
2.12	Chemical Structure of the Chemically Modified Synthetic Adsorbent SP200 Series	37

2.13	Types and Categories of Synthetic Adsorbents	39
2.14	Carotenoids Extraction from CPO PORIM Process	44
2.15	Carotenoids and Tocotrienols Extraction from CPO Carotech Associates, USA	48
2.16	Carotenoids Extraction from CPO, Lion Corporation, Japan	51
2.17	Schematic Diagram of Continuous Supercritical Carbon Dioxide Processing of Palm Oil	52
3.1	The Effect of Adsorbent/CPO Ratio on the Percentage of Carotene Extracted	70
4.1	The Adsorption Lifespan and the Effect on Percentage of Carotene Extracted	81
5.1	High Performance Liquid Chromatography (HPLC) Chromatogram of Carotenes in Feed CPO (A) and Hexane Fraction (B)	94
5.2	Carotene Stability during Storage	95
5.3	Gas Chromatogram of Triglyceride Carbon Number Composition of Crude Palm Oil	123
5.4	Gas Chromatogram of Triglyceride Carbon Number Composition of Oil from IPA Fraction	124
5.5	Gas Chromatogram of Triglyceride Carbon Number Composition of RBDPO from Plant Production (Golden Jomalina Food Industries Sdn. Bhd.)	125
5.6	Gas Chromatogram of Triglyceride Carbon Number Composition of RBDPO from Pilot Plant Refining	126

5.7	Gas Chromatogram of Fatty Acids Composition of Crude Palm Oil	127
5.8	Gas Chromatogram of Fatty Acids Composition of Oil from IPA Fraction	128
5.9	Gas Chromatogram of Fatty Acids Composition of RBDPO from Plant Production (Golden Jomalina Food Industries Sdn. Bhd.)	129
5.10	Gas Chromatogram of Fatty Acids Composition of RBDPO from Pilot Plant Refining	130

LIST OF PLATES

Plate		Page
3.1	Adsorption Process	132
3.2	Soxhlet Extractor	132
3.3	Vacuum Evaporator	133
3.4	Shidmadzu UV-1601 Spectrophotometer	133
4.1	Jacketed Flask Extractor	134
5.1	Degumming and Bleaching Process	134
5.2	Filtration Process	135
5.3	Deodorization Process	135
5.4	Gas-Liquid Chromatography (GLC)	136
5.5	High-Performance Liquid Chromatography	136
	(HPLC)	

CHAPTER I

GENERAL INTRODUCTION

According to Goh (1991) palm oil is composed predominantly of triglycerides (TG) (and partial glycerides), with minor constituents of carotenoids (mainly α - and β -carotene), tocopherols, tocotrienols, sterols (e.g. cholesterol), phospholipids, triterpenes phospholipids and aliphatic hydrocarbons. Carotenoids and vitamin E are the principal minor components (Ooi, 1995) and they are known to possess important nutritional and physiological properties.

According to Choo et al., (1989) the concentration of carotene in palm oil can range from 400 ppm to 4600 ppm, depending on the species of the palm fruit from which the oil is obtained. The commercial Malaysian crude palm oil (CPO) contains about 500 ppm to 700 ppm (Jacobsberg, 1974 and Goh et al., 1985) of carotenoids with α - and β -carotene formed up to 90% of the total carotenoids.

Carotenes, in particular β -carotene, are known for their pro-vitamins A activities as they can be transformed into Vitamin A *in vivo*. The vitamin A equivalents of α -, β - and γ -carotenes and β -zeacarotenes are 0.9, 1.67,