

UNIVERSITI PUTRA MALAYSIA

KINETICS AND MECHANISM OF CADMIUM, COPPER AND LEAD ION BIOSORPTION USING ASPERGILLUS FLAVUS 44-1 LIVE BIOMASS

KOK KEAN HIN

FSMB 2002 3

KINETICS AND MECHANISM OF CADMIUM, COPPER AND LEAD ION BIOSORPTION USING ASPERGILLUS FLAVUS 44-1 LIVE BIOMASS

By

KOK KEAN HIN

Thesis Submitted in Fulfilment of the Requirement for the Degree of Master of Science in the Faculty of Food Science and Biotechnology Universiti Putra Malaysia

June 2002



DEDICATION

Specially dedicated to.

My parents, F.O., Kok and P.H., Yeap My siblings, S.N., Kok, S.F., Kok and S.Y., Kok All of my family members All of my friends

My supervisor, Professor Dr. Mohamed Ismail Abdul Karım My committee members, Associate Professor Dr. Arbakarıya Arıff Dr. Suraını Abdul Azız

Thank you for everything.



Abstract of thesis presented to the senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Master of Science

KINETICS AND MECHANISM OF CADMIUM, COPPER AND LEAD ION BIOSORPTION USING *ASPERGILLUS FLAVUS* 44-1 LIVE BIOMASS

By

KOK KEAN HIN

June 2002

Chairman: Professor Mohamed Ismail Abdul Karim, Ph.D.

Faculty: Food Science and Biotechnology

Study on the feasibility of using live biomass of *Aspergillus flavus* as biosorbent to remove heavy metals, such as cadmium (Cd), copper (Cu) and lead (Pb) from solution was carried out in batch sorption isotherm experiments using 500.0 mL shake flask and 2 L stirred tank reactor. The effect of metal concentration (0 - 480.0 mg/L), biosorbent concentration (0 - 5.0 g/L), pH (pH 1.0 - pH 5.0) and temperature ($10.0^{\circ}C - 60.0^{\circ}C$) were investigated in single (Cd, Cu, Pb) and multimetals (CdCu, CdPb, CuPb, CdCuPb) system. Preliminary study on the biosorption heat of metal ions (Cd²⁺, Cu²⁺, Pb²⁺) on *Aspergillus flavus* was also conducted. Microscopic study using Scanning and Transmission Electron Microscope and X-ray Energy Dispersive analysis were also performed.

Results obtained from single cadmium (Cd), copper (Cu) and lead (Pb) system in the shake flask experiments revealed that biomass of *Aspergillus flavus*



was a potential biosorbent for the removal of Cd, Cu and Pb from solution. Optimum pH for the maximum removal of Cd, Cu and Pb was at pH 2.0, pH 5.0 and pH 4.0, respectively. Optimum temperature for the maximum removal of Cd, Cu and Pb was occurred at 30.0° C, 30.0° C and 40.0° C, respectively. Results from this study also showed that a small amount of *Aspergillus flavus* biosorbent, less than 0.6g, was sufficient to remove a significant large amount of metal ions, almost 23.33 mg/L (Cd²⁺, Cu²⁺, Pb²⁺) from 100.0 mg/L solution.

In the dual (CdCu, CdPb, CuPb) and tri-metals (CdCuPb) system, removal of cadmium (Cd), copper (Cu) and lead (Pb) from solution were interfered by the presence of inhibiting cations. The presence of competing ions have altered the equilibrium state and stability of solution chemistry of the system. The system would shift to another equilibrium in favour of the more influenced ion in the system.

Performance of biosorption in a more controlled surrounding in enclosed contactor, such as stirred tank reactor was preferred by the industry. Maximum lead uptake (59.70 mg/g) by *Aspergillus flavus* in stirred tank reactor could be achieved at pH 5.0, 30.0°C and biosorbent (*Aspergillus flavus*) concentration of 2.0 g/L.

Biosorption data of cadmium (Cd), copper (Cu) and lead (Pb) adsorption in single Cd, Cu and Pb system revealed that the Langmuir, Scatchard and Freundlich models were applicable to the biosorption system. However, the applicability of these adsorption models in the dual(CdCu, CdPb, CuPb) and trimetals (CdCuPb) system were not encouraging.

Desorption with appropriate eluant (HCl, HNO₃, H₂SO₄) was able to recover the metal ion (Cd²⁺, Cu²⁺, Pb²⁺) from solution and prevent secondary



pollution to our environment. The possibility of regenerating the adsorbeat (Aspergillus flavus) have led to the development of this promising technology.



KINETIK DAN MEKANISMA BIO-PENJERAPAN ION KADMIUM, KUPRUM DAN PLUMBUM DENGAN MENGGUNAKAN BIOJISIM HIDUP ASPERGILLUS FLAVUS 44-1

Oleh

KOK KEAN HIN

Jun 2002

Pengerusi: Profesor Mohamed Ismail Abdul Karim, Ph.D.

Fakulti: Sains Makanan dan Bioteknologi

Kebolehan biojisim hidup *Aspergillus flavus* sebagai biopenjerap dalam pengasingan logam-logam berat seperti kadmium (Cd), kuprum (Cu) dan plumbum (Pb) daripada larutan telah dikaji menggunakan ujikaji penjerapan isoterma tidak selanjar dalam kelalang kon 500.0 mL dan tangki pengaduk 2 liter. Kesan kepekatan logam (0 – 480.0mg/L), kepekatan biopenjerap (0 – 5.0g/L), pH (pH 1.0 – pH 5.0) dan suhu (10.0°C – 60.0° C) telah diselidik dalam ekperimen sistem satu logam (Cd, Cu, Pb) dan sistem pelbagai (CdCu, CdPb, CuPb, CdCuPb) logam. Penyelidikan awal pada haba biopenjerapan oleh ion logam (Cd²⁺, Cu²⁺, Pb²⁺) terhadap *Aspergillus flavus* telah diselidik. Kajian mikroskopik dengan menggunakan mikroskop elektron pengimbas dan penembus serta kajian pengagihan tenaga sinar X telah juga dijalankan. Kajian kinetik dengan penggunaan pelbagai model penjerapan isoterma seperti Langmuir, Scatchard and Freundlich telah dikaji.





Keputusan diperolehi daripada sistem satu logam (Cd, Cu, Pb) dalam eksperimen kelalang kon menunjukkan biojisim *Aspergillus flavus* berpotensi digunakan sebagai biopenjerap dalam pengasingan kadmium (Cd), kuprum (Cu) dan plumbum (Pb) daripada larutan berbanding dengan penjerap yang lain. pH optima dalam pengasingan maksima Cd, Cu dan Pb daripada larutan adalah pada pH 2.0, pH 5.0 dan pH 4.0, manakala suhu optima dalam pengasingan maksima Cd, Cu dan Pb daripada larutan berlaku pada suhu 30.0°C, 30.0°C dan 40.0°C. Kajian juga menunjukkan penggunaan biopenjerap (*Aspergillus flavus*) dalam kuantiti yang rendah, kurang daripada 0.6g, sudah memadai dalam pengasingan maksima logam-logam berat (Cd, Cu, Pb), hampir 23.33mg/L daripada 100.0mg/L larutan dan penambahan biopenjerap (*Aspergillus flavus*) tidak diperlukan.

Bagi pengasingan sistem dua (CdCu, CdPb, CuPb) dan tiga logam (CdCuPb), pengasingan kadmium (Cd), kuprum (Cu) dan plumbum (Pb) daripada larutan telah dipengaruhi oleh kehadiran kation pengganggu. Kehadiran ion penyaing ini telah mengubah keadaan keseimbangan dan kimia larutan sistem tersebut. Sistem tersebut akan berubah ke satu keadaan keseimbangan yang bersesuaian di mana ia lebih berpihak kepada ion penyaing yang lebih berpengaruh.

Keberkesanan proses biopenjerapan di dalam keadaan terkawal, seperti di dalam pengaduk tertutup dan secara amnya merujuk kepada tangki pengaduk, lebih diberi perhatian oleh pihak industri. Maksima penjerapan plumbum (59.70mg/g) oleh *Aspergillus flavus* di dalam tangki pengaduk boleh diperolehi pada pH 5.0, 30.0°C dan pada kepekatan biopenjerap (*Aspergillus flavus*) sebanyak 2.0 g/L.



Data biopenjerapan untuk penjerapan kadmium (Cd), kuprum (Cu) dan plumbum (Pb) di dalam sistem satu logam (Cd, Cu, Pb) menunjukkan model penjerapan isoterma Langmuir, Scatchard dan Freundlich boleh digunakan untuk menjelaskan kinetik proses biopenjerapan ini. Walau bagaimanapun, penggunaan model-model penjerapan isoterma ini dalam sistem dua (CdCu, CdPb, CuPb) dan tiga logam (CdCuPb) tidak bersesuaian.

Proses penyahjerapan dapat dilakukan dengan agen penyahjerap yang sesuai (HCl, HNO₃, H₂SO₄) dan boleh mengelakkan pencemaran sekunder terhadap alam sekeliling serta kitar semula bahan penjerap(*Aspergillus flavus*) dalam proses yang seterusnya. Keupayaan bagi penggunaan semula biopenjerap telah membuka peluang baru dalam perkembangan seterusnya dalam bidang teknologi ini.



ACKNOWLEDGEMENTS

For the hard time that have been history yesterdays,

.....Daylight I must wait for the sunrise

I must think of a new life

And I mustn't give in

When the dawn comes

Tonight will be a memory too

And a new day will begin.....T.S. Eliot & Trevor Nunn

First of all, I wish to express my deepest gratitude to my supervisor, Professor Dr. Mohamed Ismail Abdul Karim for his invaluable guidance, constant encouragement and constructive ideas throughout the course of this study. His patience and willingness to look into all of my problems, when I held in the bottlenecks of my study, really assist me lots. I wish to express my thankfulness to Professor Ismail again for his advice and moral support during my days of up and down. I really appreciate it !

My appreciation and gratitude also go to other members of my supervisory committee, Associate Professor Dr. Arbakariya Ariff and Dr. Suraini Abdul Aziz for their guidance and valuable comments during my study. Sincere thanks are also extended to Associate Professor Badlishah Sham Baharin, Dr. Lee Kong Hung, Dr. Foo Hooi Ling, Associate Professor Dr. Mohamed Ali Hassan, Dr. Rosfarizan Mohamed, Associate Professor Dr. Yazid Bin Abdul Manap and Dr. Kamariah Long for their not much, but valuable advice and encouragement.

I also wish to express my appreciation to Encik Rosli Alim, Encik Azman, Puan Aluyah and the staffs in the Fermentation and Bioprocess Engineering laboratory, Enzyme and Bioprocess Engineering laboratory, Waste



and Bioprocess Engineering laboratory and Biochemistry laboratory for their support and assistantship throughout my study. My appreciation also extended to Mr. Ooi and Mr. Chan of Food Technology Center, Malaysian Agricultural Research and Development Institute for their guidance to assist me in better understanding of the Flame Atomic Absorption Spectrophotometer. Thanks also extended to Encik Karim of Department of Soil Science, Faculty of Agriculture, Universiti Putra Malaysia for his guidance to enhance my skill in performing the Flame Atomic Absorption Spectrophotometer. My appreciation also extended to Associate Professor Dr. Fauziah Othman and the staffs, especially Mr. Ho, Puan Faridah, Cik Azilah and Puan Siti Selena of Electron Microscopic Unit, Enzyme and Microbial Technology Laboratory, Institute of Bioscience, Universiti Putra Malaysia for their guidance during my practice on the Environmental and Variable Pressure Scanning Electron Microscope, Transmission Electron Microscope and X-ray Energy Dispersive analysis. Not forgetting the other staffs, such as Mr. Ooi, Encik Sobri, Encik Rezal, Encik Bazli and others in Fermentation Technology Unit, Enzyme and Microbial Technology Laboratory, Institute of Bioscience, Universiti Putra Malaysia for their assistantship in carry out the bioreactor system. I would also like to express my gratitude to the Ministry of Science, Technology and Environment, Malaysia for funding this study under the Intensification Research of Priority Area Scheme (RM7) for 1 1/2 years.

Heartfelt appreciation is also due to all of the members of the faculty, staffs, fellow postgraduate and undergraduate students of the Department of Biotechnology, Faculty of Food Science and Biotechnology and Institute of Bioscience for their co-operation and assistance during my study.



Finally, 1 owe my family a debt of gratitude for what's they have sacrifice for me. Thanks for their understanding, care and invaluable support. Not forgetting my friends and colleagues who are always be helpful and blessed me with solutions during our communication.



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Thesis Supervisory Committee are as follows:

MOHAMED ISMAIL ABDUL KARIM, Ph.D., Professor, Institute of Bioscience, Universiti Putra Malaysia. (Chairman)

ARBAKARIYA ARIFF, Ph.D., Associate Professor, Fermentation Technology Unit, Enzyme and Microbial Technology Laboratory, Institute of Bioscience, Universiti Putra Malaysia. (Member)

SURAINI ABDUL AZIZ, Ph.D., Department of Biotechnology, Faculty of Food Science and Biotechnology, Universiti Putra Malaysia. (Member)

> AINI IDERIS, Ph.D, Professor/Dean, School of Graduate Studies, Universiti Putra Malaysia

Date:



TABLE OF CONTENTS

DEDICATION	2
ABSTRACT	3
ABSTRAK	6
ACKNOWLEDGEMENTS	9
APPROVAL SHEETS	12
DECLARATION FORM	14
LIST OF TABLES	20
LIST OF FIGURES	22
LIST OF PLATES	28
LIST OF ABBREAVIATIONS	30

CHAPTER

Ι	GENERAL INTRODUCTION	34
II	LITERATURE REVIEW	37
	Environmental Pollution by Disposal	
	of Metals in Malaysia	37
	Cadmium, Copper and Lead	38
	Cadmium	38
	Copper	38
	Lead	39
	Conventional Physico-chemical	
	Treatment of Metal Bearing Wastes	39
	Flocculation and Coagulation	40
	Ion Exchange	40
	Photodegradation	40
	Incineration	41
	Membrane Filtration System	41
	Bioremediation and Biosorption	41
	The Application of Microorganism	
	in Bioremediation Technology	42



Algae	43
Bacteria	44
Yeast	44
Fungi	45
Metal Elution and Regeneration of	
Biosorbent	46
Microbiology of Aspergillus flavus	46
General Background	46
Factors that Influence the Growth	
and Survival of Aspergillus flavus	47
The Mechanisms of Biosorption	47
Factors that Influence the Process	
of Biosorption	50
Effect of Metal Concentration	50
Effect of Biosorbent Concentration	50
Effect of pH	51
Effect of the Presence of Interfering	
Ions	51
Effect of Temperature	52
Application of Various Adsorption	
Isotherm Models	52
Langmuir Adsorption Isotherm	
Model	53
Freundlich Adsorption Isotherm	
Model	54
Scatchard plot	55

III	GENERAL MATERIALS AND	
	METHODS	56
	Microorganism	56
	Media Composition	56
	Preparation of Biosorbent	56
	Free Live Cell as Biosorbent	56
	Non-viable Powderized Biomass as	
	Biosorbent	57
	Reagents	57
	Experimental Layout	57
	Analytical Procedures	59
	Determination of Cd ²⁺ , Cu ²⁺ and	
	Pb ²⁺ in Solution using Atomic	
	Absorption Spectrophotometer	59
	Measurement of Cell Dry Weight	61
	Microscopic Study using	



Scanning Electron Microscope-	
X-ray Energy Dispersion analysis	
(SEM-EDAX) and Transmission	
Electron Microscope (TEM)	62

IVEFFECT OF CADMIUM, COPPER AND
LEAD IN GROWTH MEDIUM OF
Aspergillus flavus64Introduction64Materials and Methods64Results66Discussions77Conclusion79

AND [
<i>ivus</i> 80
80
81
82
93
98

R AND
М
flavus 100
100
101
102
117
122

VII	UPTAKE OF CADMIUM, COPPER AND	
	LEAD IN TRI-METALS SYSTEM	
	USING BIOMASS OF Aspergillus flavus	124
	Introduction	124
	Materials and Methods	125
	Results	125
		120



Discussions	133
Conclusion	137

VIII	PRELIMINARY STUDY ON THE	
	DETERMINATION OF THE	
	BIOSORPTION HEAT (Δ Hads)	
	OF CADMIUM, COPPER AND	
	LEAD ON Aspergillus flavus	138
	Introduction	138
	Materials and Methods	140
	Results	143
	Discussions	149
	Conclusion	151

IX	APPLICATION OF DIFFERENT ADSORPTION ISOTHERM MODELS ON CADMIUM, COPPER AND LEAD	
	UPTAKE USING BIOMASS OF	
	Aspergillus flavus	152
	Introduction	152
	Materials and Methods	155
	Results	156
	Discussions	163
	Conclusion	167

Х	THE PERFORMANCE OF LEAD BIOSORPTION USING BIOMASS OF <i>Aspergillus flavus</i> IN BATCH STIRRED	
	TANK REACTOR	168
	Introduction	168
	Materials and Methods	169
	Results	171
	Discussions	178
	Conclusion	183

XI GENERAL DISCUSSIONS, CONCLUSIONS AND RECOMMENDATIONS 184



BIBLIOGRAPHY	187
APPENDICES	198
BIODATA OF THE AUTHOR	201



LIST OF TABLES

Table		Page
1	Metal accumulation by algae	44
2	Metal accumulation by bacteria	44
3	Metal accumulation by yeast	45
4	Metal accumulation by fungi	45
5	Capability of cadmium adsorption between Aspergillus flavus with other adsorbents	94
6	Capability of copper adsorption between Aspergillus flavus with other adsorbents	94
7	Capability of lead adsorption between Aspergillus flavus with other adsorbents	95
8	General characterization of cadmium, copper and lead	118
9	Cadmium, copper and lead biosorption data of Langmuir linearized equation using dead cell of <i>Aspergillus flavus</i> at different temperatures	148
10	Cadmium, copper and lead biosorption data of Langmuir linearized equation in single Cd, Cu and Pb system, dual metals system of CdCu, CdPb and CuPb and tri-metals system of CdCuPb	158
11	Scatchard analysis for cadmium, copper and lead biosorption data of <i>Aspergillus flavus</i> in single Cd, Cu and Pb system, dual metals system of CdCu, CdPb and CuPb and tri-metals system of CdCuPb	140
		160



12	Cadmium, copper and lead biosorption data of Freundlich equation by <i>Aspergillus</i> <i>flavus</i> in single Cd, Cu and Pb system, dual metals system of CdCu, CdPb and CuPb and tri-metals system of CdCuPb	162
13	Effect of temperature on lead adsorption by Aspergillus flavus under controlled conditions in batch stirred tank reactor	176



LIST OF FIGURES

Figure		Page
1	Experimental layout for Cd ²⁺ , Cu ²⁺ and Pb ²⁺ biosorption in single, dual and tri-metals system	58
2	Experimental layout for study of biosorption heat on <i>Aspergillus flavus</i> and lead biosorption in batch stirred tank reactor	59
3	Standard curve for cadmium	60
4	Standard curve for copper	61
5	Standard curve for lead	61
6	Effect of cadmium, copper and lead presence in the growth medium of Aspergillus flavus	66
7	Effect of biosorbent concentration on cadmium biosorption in single cadmium system by Aspergillus flavus	82
8	Effect of biosorbent concentration on copper biosorption in single copper system by Aspergillus flavus	83
9	Effect of biosorbent concentration on lead biosorption in single lead system by <i>Aspergillus flavus</i>	83
10	Effect of [Initial cadmium/biosorbent] ratio on cadmium uptake capacity in single cadmium system by Aspergillus flavus	84
11	Effect of [Initial copper/biosorbent] ratio on copper uptake capacity in single copper system by Aspergillus flavus	85
12	Effect of [Initial lead/biosorbent] ratio on lead uptake capacity in single lead system by Aspergillus flavus	85
13	Effect of cadmium concentration on cadmium biosorption in single cadmium system by Aspergillus flavus	86



14	Effect of copper concentration on copper biosorption in single copper system by Aspergillus flavus	87
15	Effect of lead concentration on lead biosorption in single lead system by Aspergillus flavus	87
16	Effect of pH on cadmium uptake in single cadmium system by Aspergillus flavus	88
17	Effect of pH on copper uptake in single copper system by Aspergillus flavus	89
18	Effect of pH on lead uptake in single lead system by Aspergillus flavus	90
19	Effect of temperature on cadmium biosorption in single cadmium system by Aspergillus flavus	91
20	Effect of temperature on copper biosorption in single copper system by Aspergillus flavus	92
21	Effect of temperature on lead biosorption in single lead system by Aspergillus flavus	93
22	Effect of biosorbent concentration on cadmium and copper biosorption in binary system of CdCu by Aspergillus flavus	103
23	Effect of biosorbent concentration on cadmium and lead biosorption in binary system of CdPb by Aspergillus flavus	103
24	Effect of biosorbent concentration on copper and lead biosorption in binary system of CuPb by Aspergillus flavus	104
25	Effect of metal concentration on cadmium and copper biosorption in binary system of CdCu by <i>As pergillus</i>	



	flavus	105
26	Effect of metal concentration on cadmium and lead biosorption in binary system of CdPb by <i>Aspergillus flavus</i>	106
27	Effect of metal concentration on copper and lead biosorption in binary system of CuPb by Aspergillus flavus	107
28	Effect of pH on cadmium biosorption in binary system of CdCu by Aspergillus flavus	108
29	Effect of pH on copper biosorption in binary system of CdCu by Aspergillus flavus	109
30	Effect of pH on cadmium biosorption in binary system of CdPb by <i>Aspergillus flavus</i>	110
31	Effect of pH on lead biosorption in binary system of CdPb by Aspergillus flavus	110
32	Effect of pH on copper biosorption in binary system of CuPb by Aspergillus flavus	111
33	Effect of pH on lead biosorption in binary system of CuPb by Aspergillus flavus	112
34	Effect of temperature on cadmium biosorption in binary system of CdCu by Aspergillus flavus	113
35	Effect of temperature on copper biosorption in binary system of CdCu by Aspergillus flavus	114
36	Effect of temperature on lead biosorption in binary system of CdPb by Aspergillus flavus	114
37	Effect of temperature on cadmium biosorption in binary system of CdPb	

