



UNIVERSITI PUTRA MALAYSIA

**SAMPLING TECHNIQUE FOR DETERMINING EGG DISTRIBUTION
COCOA POD BORER (*CONOPORMORPHA CRAMERELLA*) (SNELLEN)
(LEPIDOPTERA: GRACILLARIIDAE) AND COCOA POD PHENOLOGY**

NOORAZLIN MOHD ALI

FS 2005 48

**SAMPLING TECHNIQUE FOR DETERMINING EGG DISTRIBUTION
COCOA POD BORER (*CONOPORMORPHA CRAMERELLA*) (SNELLEN)
(LEPIDOPTERA: GRACILLARIIDAE) AND COCOA POD PHENOLOGY**

By

NOORAZLIN MOHD ALI

**Thesis Submitted to the School of Graduate Studies, University Putra Malaysia,
in Fulfilment of the Requirement for the
Degree of Master of Science**

2005



DEDICATION

To my beloved husband, Zamran Haji Hari thanks for helping me making my dreams a reality. Your sacrifices, patience, support and encouragement are truly helpful.

I am truly indebted to my father, Haji Mohd Ali Mohd Yusof and mother, Hajah Azizah Tajuddin. I really appreciate your support, inspiration and motivation. Its hard for me without your love and sacrifice. There is no other compatible parent such both of you.

My heartfelt appreciation to every member of my family; Alang, Fadzli, Firdaus and Faiz, thanks for your understandable, assistance and encouragement during my studies.

And to my inspiring little kid, Nurin Aqilah Zamran; you understand the hardship of me and believe me dreams will be reality with Allah S.W.T permitted.



Abstract of thesis presented to the Senate of University Putra Malaysia in fulfillment of the requirement for the degree of Master Science

**SAMPLING TECHNIQUE FOR DETERMINING EGG DISTRIBUTION
COCOA POD BORER (*CONOPORMORPHA CRAMERELLA*) (SNELLEN)
(LEPIDOPTERA: GRACILLARIIDAE) AND COCOA POD PHENOLOGY**

By

NOORAZLIN BT MOHD ALI

June 2005

Chairman: Professor Syed Tajuddin Syed Hassan, PhD

Faculty: Science

An in-depth investigation on sampling of cocoa pod bore (CPB) was carried out over a 12 month period in 2000 at Malaysia Cocoa Board Research station at Hilir Perak, Sg Sumur Perak conducted over a period of 12 months. Several components were examined. The components examined were the most reliable and efficient sampling technique, determining distribution of CPB with respect to pod phenology, determining optimum sample size to produce sampling parameters for CPB. Generally, during low population levels, the mean population densities and their relative variations were similar for all sampling techniques. However, during high population densities, mean densities and relative variations increased. This is especially for random sampling which has been found as the most reliable sampling technique. Using two aggregation models which are widely used; Taylor's Power Law (b coefficient) CPB's egg were found to be highly aggregated. In determining optimum sample sizes, two parameters were calculated; b of Green's model and β for



Kuno's model. The Green model was suitable to produce the most efficient sampling sizes during low and high population densities. During high population densities, random sampling was the most reliable with Taylor's Power Law providing the optimum coefficients for developing optimum sample sizes.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**TEKNIK PENSAMPELAN UNTUK MENENTUKAN TABURAN ULAT
PENGOREK BUAH KOKO (UPBK) (*CONOPORMORPHA CRAMERELLA*)
(SNELLEN) (LEPIDOPTERA: GRACILLARIDAE DAN FENOLOGI BUAH**

Oleh

NOORAZLIN BT MOHD ALI

Jun 2005

Pengerusi: Professor Syed Tajuddin Syed Hassan, PhD

Fakulti: Sains

Pada tahun 2000, satu kajian terperinci mengenai pensampelan ulat pengorek buah koko (UPBK) telah dijalankan selama 12 bulan. Terdapat beberapa komponen utama yang dikaji meliputi kaedah yang paling bersesuaian dan efektif, mengenal pasti bentuk taburan ulat pengorek buah koko (UPBK) berdasarkan fenologi buah koko, dan penentuan saiz sampel optimum untuk menghasilkan parameter pensampelan yang paling optimum. Secara umum, apabila populasi UPBK rendah, nilai min populasi dan relatif variasi yang diperolehi menyerupai bagi kaedah-keadah yang digunakan. Namun begitu, apabila populasi bertaburan tinggi kaedah rawak merupakan kaedah yang paling persis. Dengan menggunakan dua model pengelompokan yang terkenal, Hukum Kuasa Taylor dan Indeks Pengelompokan Iwao, UPBK ditemui secara meluas dalam keadaan berkelompok. Bagi penentuan saiz sampel optimum, dua bentuk model iaitu Green, b dan Kuno, β digunakan. Model Green didapati paling sesuai menghasilkan kaedah pensampelan paling

effektif pada populasi UPBK tinggi dan rendah. Apabila populasi UPBK tinggi, kaedah pensampelan secara rawak memberikan hasil yang paling berpadanan dimana Hukum Kuasa Taylor memberikan pekali optimum untuk pembentukan saiz sample paling optimum.

ACKNOWLEDGEMENT

My sincere praises and thanksgiving to Allah S.W.T for permitting me to complete my Master of Science project successfully.

My deep appreciation extends foremost to the Chairman supervisory committee, Prof. Dr. Syed Tajuddin Syed Hassan for his valuable encouragement, guidance, criticisms and support enabling me to complete this project. Special thanks are also expressed to members of the supervisory committee, Dato' Dr. Azhar Ismail from Malaysian Cocoa Board (MCB), and to Professor Mohd Yusof Husien and Prof. Dr. Ahmad Said Sajap for their guidance, and assistance in preparation and completion of this thesis. I would like to thank the former member whom largely responsible for the privilege of researching with full accommodations at the MCB field station at Hilir Perak, Entomological Laboratory, Malaysian Cocoa Board of Hilir Perak, En. Roslan, En. Shukor, En. Zaki and Puan Hamimah for their assistance during the field study and observation. Assistance from Cik Aina of Faculty Science for help in solving technical problems regarding registration is gratefully acknowledged.

To all members of Entomology Laboratory of the Department of Biology Universiti Putra Malaysia; Ruzila, Saripah, Navies, Mai Shihah, Rysalls and student project in batch 2002-2003 and 2004-2005, Nur, Rosni, Wan, Rizal, Rafie, Izwan, Kamal, Najwa, Nur, Niza, Iz, Zul, Ami, Nik and Amran, I sincerely express heartfelt gratitude for continual friendship, support and assistance.



To my parents, husband and family; their sincere most love, patience and encouragement light the beacon of perseverance for me to complete this project.

Last but not least, the I gratefully acknowledge MARA sponsorship which enables me to complete this project.

I certify that an Examination Committee has on 18 June 2005 to conduct the final examination of Noorazlin Mohd Ali on her Master of Science thesis entitled “Sampling Techniques for Determining Egg Distribution of Cocoa Pod Borer, *Canopomorpha cramerella* (Snellen) (Lepidoptera: Gracillariidae) and Cocoa Pod Phenology” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the degree of Master of Science.

Members of the Examination Committee were as follows:

Umi Kalsum Yusof, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Nor Azwady Abdul Aziz, PhD


Lecturer
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Abdul Rahim Ismail, PhD

Lecturer
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Idris Abdul Ghani, PhD

Associate Professor
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(External Examiner)



HASANA H. GHAZALI, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 08 DEC 2005

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement of the degree of Master of Science. The members of the supervisory Committee are as follows:

Syed Tajuddin Syed Hassan, PhD

Professor
Faculty of Science
University Putra Malaysia
(Chairman)

Ahmad Said Sajap, PhD

Professor
Faculty of Forestry
University Putra Malaysia
(Member)

Dato' Dr. Azhar Ismail, PhD

Director General
Malaysian Cocoa Board
(Member)



AINI IDERIS, PhD

Professor/Dean
School of Graduate Studies
University Putra Malaysia
Date: 8 DEC 2005

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



NOORAZLIN MOHD ALI

Date: 2007

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xii
LIST OF FIGURES	
LIST OF ABBREVIATIONS	xviii
CHAPTER	
1 INTRODUCTION	1
1.1 Problem statement	2
1.2 Integrated Pest Management	2
1.3 Research Objectives	3
2 LITERATURE REVIEW	4
2.1 The purpose of sampling	4
2.2 Sampling concepts	5
2.2.1 Sampling unit and samples	5
2.2.2 Sampling patterns	6
2.2.3 Sampling technique	7
2.2.4 Sampling program	8
2.3 Population distribution	9
2.4 Mathematical distribution for describing population	11
2.4.1 Poisson distribution	12
2.4.2 Negative binomial	13
2.5 Indices for classifying dispersion	14
2.5.1 Variance to Mean Ratio	14
2.5.2 Lylod's Mean Crowding	15
2.6 Regression techniques for evaluating dispersion	15
2.6.1 Taylor's Power Law	15
2.6.2 Iwao's Patchiness regression	17
2.7 Comparison of sampling technique	18
2.8 Biology of Cocoa Pod Borer, <i>Conopomorpha cramerella</i> (CPB)	18
2.8.1 Eggs	19
2.8.2 Larva	19
2.8.3 Pupa	19
2.8.4 Adult	19
3 GENERAL MATERIALS AND METHODOLOGY	21
3.1 Study Site	21
3.1.1 Block 12D	23
3.1.2 Block 3A	25
3.1.3 Block 10A	27



3.2	Sampling Eggs	29
3.2.1	Sample units	29
3.3	Sampling Technique	30
3.3.1	Random	30
3.3.2	Systematic	30
3.3.3	Stratified	31
3.4	Timing of sampling	32
3.5	Recording	32
3.6	Statistical Analysis	33
4	COMPARISON OF SAMPLING TECHNIQUES	35
4.1	Introduction	35
4.2	Materials and Methodology	36
4.2.1	Data source	36
4.2.2	Sampling Techniques	36
4.2.3	Analysis	36
4.3	Results	38
4.3.1	Mean population (\bar{x})	40
4.3.2	Relative variation (RV)	43
4.3.3	Cost reliability and relative net precision (RNP)	46
4.4	Discussion	49
5	TEMPORAL VARIATION ON CPB'S EGG POPULATION	51
5.1	Introduction	51
5.2	Materials and Methods	52
5.2.1	Data source	52
5.2.2	Dispersion analysis	52
5.2.3	Ecological dispersion analysis	52
5.2.4	Regression technique analysis	53
5.3	Results	53
5.3.1	Distribution for overall sampling location	63
5.3.2	Distribution parameter based on the sampling methods	63
5.3.2.1	Random	63
5.3.2.2	Systematic	65
5.3.2.3	Stratified	67
5.3.3	k parameters	70
5.4	Discussion	72
5.4.1	Distribution of the CPB	72
5.4.2	Pod phenology	76
6	OPTIMUM SAMPLE SIZE	80
6.1	Introduction	80
6.2	Materials and Methodology	81
6.2.1	Data sampling	81
6.2.2	Analysis	81
6.2.2.1	Optimum sample size determination	82
6.3	Results	83
6.3.1	Optimum sampling size	83
6.4	Discussion	90

7	SEQUENTIAL SAMPLING PLAN	92
7.1	Introduction	92
7.2	Materials and Methodology	93
7.2.1	Data sampling	93
7.2.2	Sampling Plans	93
7.2.3	Analyses	93
7.3	Result	94
7.4	Discussion	100
8	GENERAL DISCUSSION AND CONCLUSION	102
	REFERENCES	107
	APPENDICES	120
	BIODATA OF THE AUTHOR	172



LIST OF TABLES

Table		Page
4.1	Result of Normality Test as one component in Exploratory data analysis vertical	38
4.2	Result of Kruskal –Wallis Test based on every categories.	39
4.3	Result of Friedman-test on the difference of the sampling block, duration and technique	39
4.4	Monthly mean of three different sampling technique random, systematic and stratified.	41
4.5	Chisquare test for Relative Variation (RV) on the sampling duration and sampling technique.	44
4.6	Monthly RV of three different sampling technique random, systematic and stratified	44
4.7	Average time for sampling technique	46
4.8	Chisquare test for Relative Net Precision (RNP) on the sampling duration and sampling technique.	47
4.9	Monthly RNP of three different sampling technique random, systematic and stratified	47
5.1	Distribution parameter for overall sampling location based on Taylor Power Law using random sampling technique	64
5.2	Distribution parameter for overall sampling location based on Iwao Mean Crowding regression using random sampling technique	64
5.3	Distribution parameter for overall sampling location based on Taylor Power Law using systematic sampling technique	65
5.4	Distribution parameter for overall sampling location based on Iwao Mean Crowding regression using systematic sampling technique	66
5.5	Distribution parameter for overall sampling location based on Taylor Power Law using stratified sampling technique	67

5.6	Distribution parameter for overall sampling location based on Iwao Mean Crowding regression using systematic sampling Technique	68
5.7	K values for each sampling technique	70
5.8	R^2 of the regression between mean densities of each technique	70
6.1	Distribution parameter obtained using Taylor Power Law	85
6.2	Distribution parameter obtained using Iwao's Regression Indices	85
6.3	Optimum sample sizes using Taylor Power Law model and Iwao Patchiness Regression model for random sampling	86
6.4	Optimum sample sizes using Taylor Power Law model and Iwao Patchiness Regression model for systematic sampling	87
6.5	Optimum sample sizes using Taylor Power Law model and Iwao Patchiness Regression model for stratified sampling	88
7.1	Sequential sampling plan for random sampling technique	96
7.2	Sequential sampling plan for systematic sampling technique	96
7.3	Sequential sampling plan for stratified sampling technique	97

LIST OF FIGURES

Figure	Page
2.2 The cocoa pod borer, <i>Conopomorpha cramerella</i>	20
3.1 Sampling location	22
3.2 Block 12D	23
3.3 Map of Block 12D	24
3.4 Block 3A	25
3.5 Map of Block 3A	26
3.6 Block 10A	27
3.7 Map of Block 10A	28
3.8 Random	30
3.9 Systematic	31
3.6 Stratified	31
3.7 Eggs on the pod	32
3.8 Various cocoa pod sizes	33
3.9 Block 12D	30
3.10 Block 10A	31
3.11 Block 3A	32
4.1 Comparison of monthly mean densities of CPB's egg population using three sampling techniques, random, systematic and stratified.	42
4.2 Comparison of monthly relative variation of CPB's egg population three sampling techniques, random, systematic and stratified.	45
4.3 Comparison of monthly relative net precision values of CPB's egg population using three sampling techniques, random, systematic and stratified.	48
5.1 Frequency of pods with CPB eggs sampled using random, Systematic and stratified sampling technique at block 12D.	54-55

5.2	Frequency of pods with CPB eggs sampled using random, systematic and stratified sampling technique at block 10A.	56-57
5.3	Frequency of pods with CPB eggs sampled using random, systematic and stratified sampling technique at block 3A.	58-59
5.4	Comparison of temporal variation of b and β value based on three different methods of sampling from August 2000 till September 2001	60
5.5	Temporal variation of b value between three different locks, bock 12D, block 10A and block 3A.	61
5.6	Temporal variation of β value between three different locks, bock 12D, block 10A and block 3A.	62
5.7	Comparison of temporal variation of b and β value based on three different technique of sampling from August 2000 till September 20001	69
5.8	The changes pattern of the k value using three sampling techniques.	71
6.1	Optimum sample size for overall sampling location based on Taylor's model.	77
6.2	Optimum sample size for overall sampling location based on Iwao's model.	77
6.3	Comparison of Taylor's model and Iwao's model optimum sample Size at various precision levels for three different method.	89
7.1	Stop line for random techniques for model Green at $D_0 = 0.25, 0.25$ and 0.35 .	98
7.2	Stop line for random, systematic and stratified techniques for both model Green and Kuno at $D_0 = 0.25, 0.25$ and 0.35 .	99

LIST OF ABBREVIATIONS

b	Taylor's index
β	Iwao's index
$^{\circ}\text{C}$	Celcius
<i>et al.</i>	others
i.e	that is
m^2	meter square
ml	mililiter
r^2	correlation coefficient of regression
RV	relative variation
RNP	relative net precision
SE	standard error
TPL	Taylor Power Law
\bar{x}	mean density



CHAPTER 1

INTRODUCTION

Cocoa is the third most important primary crop in Malaysia which provide RM 0.66 billion revenue to the country (MCB, 2001). The Malaysian cocoa industry has developed rapidly since late 70's with production reaching 240, 000 tons in 1990. It is the most productive crop in the world with a yield of 1 tons/ha/year (MCB, 2001). However, in 1980's the industry was beset with the problems of prolonged depressed price and the infestation of a serious pest, the cocoa pod borer, *Conopomorpha cramerella* (Snellen) (CPB).

The cocoa pod borer was first discovered in late 1970 in Tawau and subsequently found in Peninsular Malaysia in 1986. It is indigenous to South East Asia as it is known to attack Leguminosae species such as *Nephelium lappaceum* (rambutan) pulasan and *Nephelium mutabile* (pulasan) and other members of Sapindicae such as *Pometia pinnata* (kasai). The life cycle of CPB begins with the laying of eggs on pod surface especially within the primary furrow (Azhar and Long, 1993). The larva hatches, and burrows directly into the pod to feed within the placenta. The penetration into the placenta of the young pods resulted in retarded pod development and malformed beans. This poses persistent problems to the growers, which subsequent increase in production cost and reduction yield.

1.1 Problem statement

Several methods are recommended to manage CPB including frequent harvesting, bagging or sleeving, chemical treatment and the use of natural enemies and pheromones. However, the use of chemical treatment or insecticides is the most common approach to control the CPB infestation. Many estate managers make treatment decisions typically based on infestation level (Sidhu *et al.* 1987), thus setting a time lag in between the time of decision and the actual implementation of control. Assessment made during observations does not reflect the actual situation when decisions are made. In addition, the continuous use of insecticides poses environmental problems and increase in production cost (Azhar *et al.* 2001). The severity of pest infestation too has increased since many pests have become more resistant to existing insecticides (Voss, 1988, Senn *et al.* 1994). Hence, an effective timing of insecticide applications based on the population level and crop phenology is needed.

1.2 Integrated Pest Management

The Integrated Pest Management (IPM) programs in agroecosystems have shown a positive impact on rice (Hassan *et al.* 1996). Since the primary objective of IPM is to reduce losses resulting from pests in a very effective, economically and ecologically compatible way, various pest management approaches have been recommended to reduce the reliance on chemical pesticides to improve pest management decision making at the farm level (MacKay *et al.* 1993).

Sampling is an important IPM component, especially in the development and execution of quantitative studies, estimation assignments, monitoring and surveillance strategies, pest population modeling and simulation, and forecasting protocols (Hassan, 1999). The major purposes of sampling are; determining statistics of samples or parameters of population, and secondly accessing the status of the population with respect to action thresholds (Hassan, 1999). Many factors should be considered such as within plant distribution of the pest (Maisin *et al.* 1997) in order to represent population attributes for decision-making analysis whether to treat or to classify the density of the population. Besides, the choice of accurate and relevant sampling plan and dynamics should be scrutinized (Wilson and Room, 1982 ; Hassan *et al.* 1992). This helps to produce a commercial procedure with the fastest method for estimating the yield of uninfested pods based on the distribution pattern of eggs study (Azhar and Long, 1993). The contribution of phenology growth of the cocoa pod is essential in most CPB's management program (Day, 1956).

1.3 Research Objectives

The main objectives of this research were 1) to determine the most reliable sampling technique for CPB egg distribution pattern, 2) to analyse the distribution pattern of CPB with respect to the phenology of growth of the cocoa pod, 3) to determine the optimum sample sizes for population management purposes.

CHAPTER 2

LITERATURE REVIEW

2.1 The Purpose of Sampling

Sampling is an essential activity in entomological and ecological research in order to get accurate and precise estimates of species abundance where the understanding of the population's dynamics is desired (Buntin, 1994). It provides current information on the pest status to make timely decisions for management activities in integrated pest management (Gonzalez, 1971) and form a baseline population research database (Morris, 1955).

In integrated pest management (IPM), there are two main purposes in sampling i.e., sampling for density estimation and for decision-making purposes (Binns and Nyrop, 1992). Ruesink (1980) reported that the reasons for sampling are to determine whether the economic threshold has been exceeded and hence the need for control measure and secondly, to estimate the population density with some predetermined degree of reliability. An example of management purpose sampling is sequential sampling by Azhar and Long (1991, 1993), who developed the fastest method for estimating cocoa yield.

An example of the second purpose sampling is assessment of insect abundance in a particular habitat (Azhar and Hassan, 2000), where reasonably accurate determination of population density is important. Estimation of density sampling involve technical and logistic constraints which need to be addressed to provide reliable data estimates.

2.2 Sampling Concepts

2.2.1 Sample Unit and Samples

Sampling universe in statistics refers as a whole population in which samples are taken (Pedigo, 1989). In arthropod sampling, the sampling universe represents the habitat in which the population occurs (Lee, 1997). For instance, the cocoa and rice fields are the sampling universe.

The sampling unit and samples are taken within the sampling universe. The measurable entity is based on the sampling unit such as a single leaf, single fruit or whole plant. In rice ecosystem, Hassan and Rashid (1997a; 1997b) a hill was the sampling unit while Azhar and Long (1991; 1993) used a cocoa pod as the sampling unit for cocoa ecosystem, and a cocoa tree as another alternative

According to Azhar and Hassan (2000) a sampling unit should follow several criteria as listed below;