



UNIVERSITI PUTRA MALAYSIA

**CHEMICAL CONSTITUENTS OF *MICROMELUM MINUTUM*
AND *ACRONYCHIA LAURIFOLIA* (RUTACEAE)**

YAP YUN HIN @ MD. TAUFIQ

FSAS 1994 3

**CHEMICAL CONSTITUENTS OF *MICROMELUM MINUTUM*
AND *ACRONYCHIA LAURIFOLIA* (RUTACEAE)**

By

YAP YUN HIN @ MD. TAUFIQ

Thesis Presented in Fulfilment of the
Requirements for the Degree of Master of Science
in the Faculty of Science and Environmental Studies,
Universiti Pertanian Malaysia

1994



DEDICATIONS

To my beloved father and mother,

THE LATE YAP YOON SANG
ARWAH ABDULLAH LIM HENG FUI
SITI AISHAH CHIN SU KIAW

To my brothers and sister,

LIM YIT HIANG
NORAINI LIM FUNG KUN
ABDUL HALIM LIM YIT VUI

To my dearest friends

SAIDAH BTE MAMAT
NORIZAN BIN REMBUN

*and the members of *Nahwan Nur**

WITHOUT WHOSE LOVE AND
CONTINUED SUPPORT THIS THESIS
WOULD NOT HAVE BEEN POSSIBLE.



ACKNOWLEDGEMENTS

I would like to express my sincere and deepest appreciation to Assoc. Prof. Dr. Mawardi Rahmani for the invaluable advises, guidances, encouragement, patient and support throughout the course of the work, Dr. Hazar B.M. Ismail, Dr. Mohd Aspollah Hj. Sukari, Assoc. Prof. Dr. Jambari Hj. Ali, Assoc. Prof. Dr. Karen Badri and Dr. Mohd. Niyadz Khan for their constant support.

I am also very grateful to my colleagues, friends and staff of the Chemistry Department of UPM, Dr. Faujan Hj. Ahmad, Dr. Mahiran Basri, En. Zainal Abidin Kasim, En. Zainal Zahari Zakaria, En. Nordin Ismail and En. Zainuddin Samadi, whose help, advice and encouragement were invaluable during this research project.

My special thanks also goes to: Prof. P.G. Waterman of University of Strachlyde for obtaining the ^1H - and ^{13}C -NMR spectra of microminutinin, 6-methoxymicrominutinin and MM5; Prof. T.W. Sam of USM and Dr. Hiroshi Sugiyama of Kyoto University, for obtaining of ^1H - and ^{13}C -NMR spectra of microminutinin and 6-methoxymicrominutinin; Prof. Latif Muhammad and En. A. Zainuddin Ibrahim of UKM, for identifying and collecting the plant materials; Assoc. Prof. Dr. S.C. Quah of UPM, for kindly supplying the *Drosophila melanogaster* culture; Puan Noraini Abu Bakar, for typing and



doing some editing work on the thesis; and Universiti Pertanian Malaysia for providing facilities and space for completion of the project.

Financial support from Malaysian Government under the IRPA programme is gratefully acknowledged



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	ii
LIST OF TABLES	vii
LIST OF FIGURES	ix
LIST OF ABBREVIATIONS	xi
ABSTRACT	xii
ABSTRAK	xiv
PART I	
EXTRACTION AND ISOLATION OF CHEMICAL COMPONENTS FROM <i>MICROMELUM MINUTUM</i> AND <i>ACRONYCHIA LAURIFOLIA</i> (RUTACEAE)	
INTRODUCTION AND LITERATURE REVIEW OF <i>MICROMELUM MINUTUM</i> (FROST. f.) SEEM AND <i>ACRONYCHIA</i> <i>LAURIFOLIA</i> BLUME.....	2
OBJECTIVES OF STUDY.....	13
RESULTS AND DISCUSSION.....	14
<i>Micromelum minutum</i> (Frost. f.) Seem.....	14
<i>Acronychia laurifolia</i> Blume.....	70
CONCLUSION.....	79



EXPERIMENTAL.....	79
EXTRACTION OF <i>MICROMELUM MINUTUM</i> (KELANTAN).....	81
Isolation of Microminutinin (32).....	82
Isolation of 6-Methoxymicrominutinin (35).....	83
Isolation of Scopoletin (13).....	84
Isolation of CHCl ₃ extract	86
EXTRACTION OF <i>MICROMELUM MINUTUM</i> (PAHANG).....	86
Isolation of Squalene (36).....	86
Isolation of Micromelin (1).....	87
Isolation of Scopoletin (13).....	88
Isolation of MM5 (37).....	89
EXTRACTION OF <i>ACRONYCHIA LAURIFOLIA</i>	90
Isolation of Sitosterol (38).....	91
Isolation of Seselin (39).....	92
Isolation of AP6.....	93

PART II PRELIMINARY BIOASSAYS AGAINST BRINE SHRIMP (*Artemia salina*) AND VINEGAR FLY (*Drosophila melanogaster*)

INTRODUCTION	95
--------------------	----



OBJECTIVES OF STUDY.....	100
RESULTS AND DISCUSSION.....	100
CONCLUSION.....	111
EXPERIMENTAL.....	111
Hatching of Brine Shrimp Cysts.....	111
Preparation of Assay Solution.....	112
Bioassay Procedure.....	112
Analysis of Results.....	113
Preparation of Culture Medium for <i>Drosophila</i>	113
Preparation of <i>D. melanogaster</i> Larvae.....	114
Preparation of Assay Media.....	114
Bioassay Procedure.....	115
BIBLIOGRAPHY.....	116
APPENDIX.....	122
VITA.....	123



LIST OF TABLES

TABLE		Page
1	Compounds from <i>Micromelum sp.</i>	9
2	¹ H-NMR Chemical Shifts (δ) and Coupling Patterns of the Protons in Various ¹ H-NMR Techniques of (32).....	25
3	¹³ C-NMR Chemical Shift Data (ppm) of (32), (33) and (34).....	30
4	¹³ C-NMR Chemical Shift Data (ppm) of (32) and (35).....	40
5	¹³ C-NMR Chemical Shift Data (ppm) of Micromelin (1)	56
6	Coupling of Protons of MM5 (37).....	66
7	¹³ C-NMR Chemical Shift Data (ppm) and ¹³ C- ¹ H Coupling of MM5.....	69
8	¹ H-NMR Spectral Data (ppm) of Seselin and Norbraylin with the Observed Data.....	76
9	Plant Extract Concentrations used in Brine Shrimp Bioassay.....	101
10	Brine Shrimp Bioassays Results of Extracts (Pahang) and Compounds of <i>M. minutum</i>	102
11	Brine Shrimp Bioassays Results of CHCl ₃ Extracts of <i>A. laurifolia</i>	103
12	Diet Formulas for <i>D. melanogaster</i>	105
13	Effect of <i>M. minutum</i> (Kelantan) Extract on <i>D. melanogaster</i> Larvae.....	106
14	Effect of <i>M. minutum</i> (Kelantan) MeOH Extract on <i>D. melanogaster</i>	107



15	Effect of CHCl ₃ Extract (leaves) of <i>M. minutum</i> (Pahang) Extract on <i>D. melanogaster</i> Larvae.....	108
16	Effect of Pure Compounds on <i>D. melanogaster</i> Larvae....	109



LIST OF FIGURES

FIGURE		Page
1	Mass Spectrum of (32).....	16
2	IR Spectrum of (32).....	17
3	¹ H-NMR Spectrum of (32).....	19
4	COSY-45 Spectrum of (32).....	20
5	COSY-LR Spectrum of (32).....	21
6	Long-range Coupling of Various Protons Observed from COSY-LR Spectrum.....	22
7	NOESY Spectrum of (32).....	23
8	DEPT ¹³ C-NMR Spectrum of (32).....	24
9	HC-COBI Spectrum of (32).....	26
10	HMBC Spectrum of (32).....	27
11	HMBC Spectrum of (32).....	28
12	¹³ C- ¹ H Coupling Pattern Observed in HMBC Spectra of (32).....	29
13	Mass Spectrum of (35).....	32
14	IR Spectrum of (35).....	33
15	¹ H-NMR Spectrum of (35).....	35
16	COSY-45 Spectrum of (35).....	36
17	¹³ C-NMR Spectrum of (35).....	37
18	DEPT ¹³ C-NMR Spectrum of (35).....	39
19	Mass Spectrum of (13).....	42



20	IR Spectrum of (13).....	43
21	¹ H-NMR Spectrum of (13).....	45
22	¹³ C-NMR Spectrum of (13).....	46
23	Mass Spectrum of (36).....	47
24	IR Spectrum of (36).....	48
25	¹ H-NMR Spectrum of (36).....	50
26	Mass Spectrum of (1).....	52
27	IR Spectrum of (1).....	53
28	¹ H-NMR Spectrum of (1).....	54
29	MS Spectrum of (37).....	58
30	IR Spectrum of (37).....	59
31	¹ H-NMR Spectrum of (37).....	61
32	COSY-45 Spectrum of (37).....	62
33	NOESY Spectrum of (37).....	63
34	DEPT ¹³ C-NMR Spectrum of (37).....	64
35	HC-COBI Spectrum of (37).....	65
36	¹³ C- ¹ H Coupling Pattern of MM5 (37).....	67
37	HMBC Spectrum of (37).....	68
38	IR Spectrum of (38).....	71
39	Mass Spectrum of (38).....	73
40	Mass Spectrum of (39).....	74
41	¹ H-NMR Spectrum of (39).....	77
42	DEPT ¹³ C-NMR Spectrum of (39).....	78



LIST OF ABBREVIATIONS

br.	broad
CHCl ₃	chloroform
CDCl ₃	deuterated chloroform
d	doublet
dt	doublet of triplet
dq	double of quartet
g	gram
IR	Infra Red
l	litre
LC	Lethal Concentration
m	medium
ml	mili litre
mg	mili gram
Me	methyl
MeOH	methanol
m.p.	melting point
MS	Mass Spectrum
NMR	Nucleur Magnetic Resonance
PE	petroleum ether
PLC	Preparative Thin Layer Chromatography
ppm	part per million
q	quartet
s	strong
TLC	Thin Layer Chromatography
UV	Ultra Violet
w	weak



Abstract of the thesis presented to the Senate of Universiti Pertanian Malaysia in fulfilment of requirements for the degree of Master of Science.

**CHEMICAL CONSTITUENTS OF *MICROMELUM MINUTUM*
AND *ACRONYCHIA LAURIFOLIA* (RUTACEAE)**

By

YAP YUN HIN @ MD. TAUFIQ

April 1994

Chairman : Associate Professor Dr. Mawardi Rahmani
Faculty : Science and Environmental Studies

A chemotaxonomic investigation on the leaves and stems of *Micromelum minutum* (Rutaceae), has resulted in the isolation of six components. The structure of these compounds was elucidated by means of spectroscopic methods including by the extensive use of various NMR techniques and also comparison with previous studies. The use of High Field NMR is essential in structural determination of these complex molecules. With the aid of various NMR experimental techniques and other spectroscopic methods such as MS, UV and IR, the correct structures of the pure isolated compounds were established.

M. minutum collected from Kelantan yielded two novel coumarins, microminutinin and 6-methoxymicrominutinin, and a known coumarin,



scopoletin. From another separate collection of the plant from Pahang, a novel compound, MM5, squalene, and two known coumarin, micromelin and scopoletin have also been isolated. A similar study on the leaves and stems of *Acronychia laurifolia* collected from Pulau Singa Besar, Langkawi Island afforded three compounds, sitosterol, seselin and an unidentified compound.

The presence of bioactive compounds in these plants was detected by the use of brine shrimp (*Artemia salina*). With the aid of this bioassay system the activity of either the crude plant extracts or pure isolated compounds could be determined. The LC₅₀ value of CHCl₃ and MeOH extracts of *M. minutum* (Pahang) gave no significant activity while CHCl₃ extracts of leaves and stems of *A. laurifolia* had LC₅₀ of 2.4 and 22.6 ppm respectively. Microminutinin gave strong test result with LC₅₀ of 34.7 ppm and MM5 with LC₅₀ of 165.5 ppm.

A more specific bioassay was also carried out on the crude and pure compounds for development inhibition activity by using fruit fly, *Drosophila melanogaster* larvae. The MeOH extract of *M. minutum* (Kelantan), at concentration of 200 mg/2 g diet, completely retarded the larval growth. For the pure compounds, microminutinin has been shown to have a strong development inhibition activity. At only 4 mg/2 g diet, it caused 100% mortality against first/second instar larvae of the insect, whereas no activities was observed from MM5.



Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia
bagi memenuhi keperluan Ijazah Master Sains.

**KANDUNGAN KIMIA DARI *MICROMELUM MINUTUM* DAN
ACRONYCHIA LAURIFOLIA (RUTACEAE)**

Oleh

YAP YUN HIN @ MD. TAUFIQ

April 1994

Pengerusi : Professor Madya Dr. Mawardi Rahmani
Fakulti : Sains dan Pengajian Alam Sekitar

Kajian kimotaksonomi ke atas ekstrak mentah daun dan ranting *Micromelum minutum* (Rutaceae) telah memberikan enam komponen. Struktur sebatian-sebatian ini telah dapat dikenal pasti dengan menggunakan kaedah spektroskopi termasuk penggunaan yang meluas pelbagai teknik NMR dan juga perbandingannya dengan kajian-kajian yang lepas. Penggunaan NMR berkuasa tinggi adalah amat diperlukan untuk menentukan struktur yang rumit. Dengan bantuan berbagai teknik ini dan juga kaedah spektroskopi yang lain seperti MS, UV dan IR, struktur yang betul bagi sebatian tulen yang dipencilkan dapat ditentukan.

M. minutum yang dikumpulkan dari Kelantan telah menghasilkan dua kumarin baru, mikrominutinin dan 6-metoksikrominutinin dan satu



koumarin yang telah dikenali, skopoletin. Dari pengumpulan berasingan (Pahang) pokok yang sama telah memberikan satu sebatian baru, MM5, skualena dan dua koumarin yang telah dikenali, mikromelin dan skopoletin. Kajian yang serupa ke atas daun dan ranting *Acronychia laurifolia* yang dikumpulkan dari Pulau Singa Besar, Pulau Langkawi telah menghasilkan tiga sebatian, sitosterol, seselin dan satu sebatian yang belum dikenalpasti.

Kehadiran sebatian bioaktif dalam tumbuhan-tumbuhan ini dapat dikesan dengan menggunakan udang pepai (*Artemia salina*). Dengan bantuan sistem biocerakanan ini, aktiviti sama ada dari ekstrak mentah tumbuhan atau sebatian tulen dapat ditentukan. Nilai LC_{50} dari ekstrak $CHCl_3$ dan MeOH dari *M. minutum* (Pahang) tidak memberikan keputusan yang bermakna sementara ekstrak $CHCl_3$ bagi daun dan ranting *A. laurifolia* masing-masing memberikan LC_{50} 2.4 dan 22.6 ppm. Mikrominutinin memberikan keputusan yang kuat dengan LC_{50} 34.7 ppm dan MM5 dengan LC_{50} 165.5 ppm.

Kaedah biocerakanan yang lebih spesifik juga telah dijalankan ke atas ekstrak mentah dan sebatian tulen bagi aktiviti penghalang perkembangan dengan menggunakan larva *Drosophilamelanogaster*. Ekstrak MeOH bagi *M. minutum* (Kelantan) pada kepekatan 200 mg/2 g diet telah sepenuhnya menghalang perkembangan larva ini. Bagi sebatian tulen pula, mikrominutinin telah menunjukkan aktiviti penghalang perkembangan. Pada kepekatan hanya 4 mg/2 g diet, mikrominutinin telah menyebabkan 100% kematian terhadap larva

peringkat pertama dan kedua bagi serangga ini, sementara itu tiada aktiviti yang dikesan bagi MM5.

PART 1

**EXTRACTION AND ISOLATION OF
CHEMICAL COMPONENTS FROM
MICROMELUM MINUTUM
AND
ACRONYCHIA LAURIFOLIA
(RUTACEAE)**





**INTRODUCTION AND LITERATURE REVIEW OF
MICROMELUM MINUTUM (FROST. f.) SEEM AND
ACRONYCHIA LAURIFOLIA BLUME**

Ten species of *Micromelum* have been found throughout South-Eastern Asia and the Pacific of which only two species occur in Peninsular Malaysia, *M. minutum* and *M. hirsutum*. *M. minutum* (synonymous with *M. pubescens* Blume) is a shrub or small tree of the family Rutaceae, which can grow up to 25 feet high. In Peninsular Malaysia, *M. minutum* which is locally known as "Chemama", "Cherek-cherek" or "Secherek" normally occurs in the hilly parts of the northern half and in the north-west area.

The plants are used medicinally. The leaves are normally pounded with tamarinds and salt and applied to the skin to relieve pain. In Peninsular Malaysia, it is traditionally used in the treatment of fever and giddiness and a poultice of the boiled roots is applied to treat ague (Burkill, 1966). The plant is also found widely throughout the Fiji Islands, where it is termed "Qiqila" and is also used by the Fijians in traditional folk medicine (Craft and Toia, 1989).

Previous phytochemical studies of *Micromelum* species revealed a number of prenylated coumarins, a pyranoquinoline alkaloid and polyoxygenated flavanoids. In 1966, Lamberton *et al.* isolated a neutral constituent, micromelin (**1**), C₁₅H₁₂O₆, m.p. 218-219.5°C from the MeOH extract of *M. minutum*. Another coumarin, osthol (**2**), m.p. 82-83.5°C was

also found in the methylene chloride extract. Later Chatterjee *et al.* (1967) reported the isolation of coumarins, micropubescin, $C_{15}H_{14}O_4$ (**3**), m.p. 130-131°C and (**1**) from the stem bark of *M. pubescens*. In 1974, Joshi *et al.* isolated a novel coumarin, 6-(2,3-dihydroxy-3-methylbutyl)-7-methoxycoumarin or ulopterol (**4**), $C_{15}H_{18}O_5$, m.p. 139-140°C from *M. pubescens* Blume.




(1) R =


(4) R = $-\text{CH}_2\text{CH}(\text{OH})\text{C}(\text{OH})(\text{CH}_3)_2$

(2) R = $-\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$


(3) R = $-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2\text{C}(\text{CH}_3)=\text{CH}_2$



(5) R =







An investigation of *M. minutum* by Tantivatana *et al.* (1982) yielded a cytotoxic compound microminutin (**5**), and a pyranoquinoline alkaloid, flindersine (**6**). Microminutin, m.p. 154-155°C was found to be inactive in KB cytotoxicity test but was reported to show weak activity (ED_{50} 3.7 $\mu\text{g/mL}$) against the P-388 lymphocytic leukemia test system *in vitro*.



(6)

Further study on *M. minutum* collected in Diphu area of the Karbi Anglong District, Assam, India by Das *et al.* (1984) afforded four known coumarins, (1), (2), murralongin (7) and murrangatin (8); and four new coumarins, dihydromicromelin A and B, (9a and 9b), acetyl dihydromicromelin A (9c), minumicrolin/mupanidin (10) and a new flavone, 7,12-ether of 5,7-dihydroxy-3,6,8,4'-tetramethoxyflavone (11).






(9a) R = α -OH

(9b) R = β -OH

(9c) R = α -OAc




(11)


Tantishaiyakul *et al.* (1986) undertook the examination of the stem bark extract of *M. minutum* from which they reported the isolation of phebalosin (12), m.p. 120.5-121.5°C. Phebalosin (14) was reported to be significantly toxic to brine shrimp (LC₅₀ 47 ppm,) and to significantly inhibit the development of crown gall tumors on potato disc but had insignificant cytotoxic activity.

From the stem-leaf of *M. integerrimum*, Cassady *et al.* (1979) discovered the occurrence of two known coumarins, (1) and scopoletin (13), and both were demonstrated to have antitumor activities. 7-[4'-(4"-methyl-5"-


oxo-2'',5''-dihydro-2''-furyl)-3'-methyl-2'-butenyloxy]coumarin (**14**), has been isolated by De Silva *et al.* (1980) from the PE extract of the leaves of *M. zeylanicum* Swingle.



(12)




(13)




(14)


In another investigation by Bowen and Perera (1981), the leaves and stems of *M. zeylanicum* have yielded a new oxazole alkaloid, *o*-methylhalfordinol (**15**), coumarin (**1**) and 5-hydroxy-3,3',4',7,8-pentamethoxyflavone (**16**). In addition, the stems also contained 6-formyl-7-methoxycoumarin, or crenulatin (**17**) and the leaves contained the carbazole alkaloid koenigine (**18**) and β -sitosterol.




(15)



(16)



(17)



(18)

Another investigation by Kong *et al.* (1988) on the roots of *M. falcatum* has revealed the occurrence of the indole alkaloid, yuehchukene (19), the novel carbazole alkaloid, 5,6-pyranoglycozoline (20), as well as (1) and (10). The various chemical components present in *Micromelum sp.* are listed in Table 1.