



UNIVERSITI PUTRA MALAYSIA

**MICROEMULSION WTH ESTER IN MIXED SURFACTANTS AND
ITS ASSOCIATION WITH BETULINIC ACID**

WAN RUSMAWATI BT WAN MAHAMOD

FSAS 2000 26

**MICROEMULSION WITH ESTER IN MIXED SURFACTANTS AND
ITS ASSOCIATION WITH BETULINIC ACID**

By

WAN RUSMAWATI BT WAN MAHAMOD

Thesis Submitted in Fulfilment of the Requirements for
the Degree of Master of Science in the Faculty of
Science and Environmental Studies
Universiti Putra Malaysia

February 2000



CHAPTER 1

INTRODUCTION

A surface active agent, or ‘surfactant’, is a chemical compound with a polar or ionic head and a hydrocarbon tail (Figure 1) (Ottewil, 1983; Benson, 1983; Lucassen, 1981). The hydrocarbon tail is hydrophobic and can be either linear or branched. The polar, or ionic, head is hydrophilic, interacting strongly with the water molecules with hydrogen bonding and dispersing them via dipole-dipole or ion-dipole interaction. A structure having both a hydrophobic and hydrophilic moieties is said to be ‘amphiphatic’.

Surfactants are classified by their hydrophilic heads into four types - ionic (either cationic or anionic) and nonionic (either zwitterionic or amphoteric) (Figure 2).

The balance between the hydrophobic and hydrophilic portions of a surfactant confers it its unique properties in use (Karsa, 1987; Myers, 1990). The applications of surfactants are many, in both everyday household products and personnel care, and in the industrial production and processing of materials (Figure 3).

*Dedicated to
In memory of my father*

*"Thank you for your everlasting love and support.
Although you are not able to share my success
I know that you will be very happy for me."*

**Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Master of Science.**

**MICROEMULSION WITH ESTER IN MIXED SURFACTANTS AND
ITS ASSOCIATION WITH BETULINIC ACID**

By

WAN RUSMAWATI BT WAN MAHAMOD

February 2000

Chairman : Assoc. Prof. Dr. Faujan Bin Haji Ahmad

Faculty : Science and Environmental Studies

The phase behaviour of Tween 80/Brij 30/H₂O and Tween 80/Brij 30/glycerol was constructed to determine the isotropic region produced by the mixed surfactants (MS) – Tween 80:Brij 30 - in aqueous and nonaqueous systems. The weight ratios of 90:10, 80:20 and 60:40 Tween 80:Brij 30 were selected for further construction of MS systems of MA or EB/Tween 80:Brij 30/water and MA or EB/Tween 80:Brij 30/glycerol. MS in aqueous systems produced a larger isotropic region than the single phase MA or EB/Tween 80/water and MA or EB/Brij 30/water systems. The system of MA/Tween 80:Brij 30 (90:10, 80:20 and 60:40)/water was then studied to determine the surfactants' aggregation and their association with BA.

The conductivity was measured to assess the charge carrier and the movement of ions in the system. The structure of the isotropic region was also determined. A reverse micelle (w/o micromulsion), bicontinuous structure

and micelle (o/w microemulsion) formed in the range of 20 – 40wt%, 40 – 60wt% and >60wt% water, respectively. However, the MA/Tween 80:Brij 30 (80:20)/water system did not form a micelle. The results were confirmed by light scattering measurement in which the particle size in each region was determined.

BA was more soluble in the system with higher Tween 80. The solubility of BA increased with the MA content and decreased with the water content.

In the antimicrobial study, 0.01 wt% BA in the microemulsion MA/Tween 80:Brij 30 (80:20)/water was more active against *Candida lipolytica* ATCC 2075 than the pure microemulsion itself. A higher mixed surfactant content (20:80 weight ratio of MA and MS) gave better activity than a lower one (50:50 weight ratio). BA in the reverse micelle region gave the highest activity compared to in the monomer and bicontinuous regions.

The cytotoxicity study produced unexpected results. The microemulsion systems were very active and killed all the cells (CEM-SS) leaving no control data with which to assess the cytotoxicity.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk syarat ijazah Master Sains

MIKROEMULSI DENGAN ESTER DI DALAM CAMPURAN SURFAKTAN DAN GABUNGANNYA DENGAN ASID BETULINIK

Oleh

WAN RUSMAWATI BT WAN MAHAMOD

Februari 2000

Chairman : Assoc. Prof. Dr. Faujan Bin Haji Ahmad

Fakulti : Sains dan Pengajian Alam Sekitar

Gambarajah fasa Tween 80/Brij 30/air dan Tween 80/Brij 30/gliserol dibina bagi menentukan kawasan isotropik yang terhasil dari campuran Tween 80:Brij 30 dalam akuas dan nonakuas. Nisbah berat antara Tween 80 dan Brij 30 pada 90:10, 80:20 and 60:40 dipilih bagi pembinaan gambarajah fasa sistem MA atau EB/Tween 80:Brij 30/air and MA atau EB/Tween 80:Brij 30/gliserol. Campuran surfaktan dalam system akuas menghasilkan kawasan isotropik yang lebih luas berbanding dengan gambarajah fasa tunggal, MA atau EB/Tween 80/air and MA atau EB/Brij 30/air. Sistem MA/Tween 80:Brij 30 (90:10, 80:20 and 60:40)/air kemudiannya digunakan dalam kajian lanjutan agregasi surfaktan yang terhasil serta gabungannya dengan BA.

Konduktiviti diukur bagi menentukan pembawaan cas dan pergerakan ion di dalam sistem. Struktur bagi kawasan isotropik juga telah ditentukan. Misel songsang, misel campuran dan misel normal masing-masing wujud

dalam julat 20 – 40%, 40 – 60% dan >60% berat air. Walau bagaimanapun, sistem MA/Tween 80 :Brij 30 (80:20)/air tiada pembentukan misel. Keputusan diperkuuhkan lagi melalui pengukuran sebaran cahaya di mana saiz partikel di dalam setiap kawasan ditentukan.

BA lebih larut di dalam sistem yang mengandungi peratusan Tween 80 yang tinggi. Kelarutan BA meningkat dengan peningkatan komposisi MA dan menurun dengan peningkatan komposisi air.

Di dalam kajian antimikrob, 0.01% berat BA di dalam sistem mikroemulsi MA/Tween 80:Brij 30 (80:20)/air paling aktif terhadap *candida lipolytica* ATCC 2075 berbanding dengan sistem mikroemulsi tersebut. Sistem yang mengandungi campuran surfaktan yang tinggi (nisbah berat MA dan campuran surfaktan pada 20:80) adalah lebih aktif berbanding dengan sistem yang rendah komposisi campuran surfaktan (nisbah berat MA dan campuran surfaktan pada 50:50). BA di dalam misel songsang memberikan aktiviti yang paling baik berbanding dengan kawasan monomer dan misel campuran.

Kajian sitotoksik menghasilkan keputusan diluar jangkaan. Sistem mikroemulsi sangat aktif dan telah membunuh semua sel CEM-SS tanpa meninggalkan kontrol bagi penilaian kesan sitotoksik.

ACKNOWLEDGEMENTS

I would like to express my most sincere gratitude and deep appreciation to the committee Chairman, Assoc. Prof. Dr. Faujan B.H. Ahmad for his invaluable contribution, guidance, ideas and time towards my project. I would also like to thank the committee members, Prof. Dr. Hamdan Suhaimi and Assoc. Prof. Dr. Anuar Kassim.

My gratitude goes to all the lecturers in the Chemistry Department for their noteworthy assistance, not forgetting a thousand thanks to Assoc. Prof. Dr. Manaf Ali, Mr. Majid Eshagi and Mr. Sahak for their help, guidance and ideas.

Furthermore my caring love goes to my family - Ma, abang, Shila, Ridzwan, Rosni and Mira - for their support and encouragement.

TABLE OF CONTENTS

	Page
DEDICATION.....	2
ABSTRACT.....	3
ABSTRAK.....	5
ACKNOWLEDGEMENTS.....	7
APPROVAL SHEETS.....	8
DECLARATION FORM.....	10
LIST OF TABLES.....	13
LIST OF FIGURES.....	14
LIST OF ABBREVIATIONS.....	20

CHAPTER

I INTRODUCTION.....	1
Microemulsion.....	10
Microemulsion with esters.....	15
Microemulsion with nonionic surfactants.....	17
Application of microemulsions.....	18
Microemulsion in pharmaceuticals.....	18
Phase equilibria.....	22
Betulinic acid	23
Solubilisation.....	27
Instrumentations.....	28
Objectives of this study	30
II MATERIALS AND METHODS.....	32
Materials.....	32
Maintenance of the microorganisms.....	33
Target microbes.....	33
Methods.....	33
Determination of phase equilibria.....	35
Conductivity measurements.....	39
Light scattering measurements.....	39
Solubility of betulinic acid (BA).....	41
Antimicrobial assay of betulinic acid (BA) in microemulsion system.....	42
Screening test for BA in DMSO.....	43
Assay for cytotoxicity.....	43
<i>In vitro</i> anti-tumour assay system.....	43
Cell counting.....	43
Protocol 1: Screening for cytotoxicity in natural products.....	45
Protocol 2: Micro-culture tetrazolium salt (MTT) assay	45

Screening test for cytotoxicity.....	47
III RESULTS AND DISCUSSION.....	48
Aqueous Systems.....	48
MA/Surfactant/Water system.....	48
MA/Mixed Surfactants/Water system.....	53
EB/Surfactant/Water system.....	60
EB/Mixed Surfactants/Water system.....	60
Nonaqueous Systems.....	70
MA/Surfactant/Glycerol system.....	70
MA/Mixed Surfactants/Glycerol system.....	70
EB/Surfactant/Glycerol system.....	75
EB/Mixed Surfactants/Glycerol system.....	82
Conductivity.....	89
Particle Size.....	98
Solubility Studies	108
Solubility of betulinic acid (BA) in organic solvents....	108
Solubility of betulinic acid (BA) in microemulsion system.....	110
Toxicity Studies.....	120
Antimicrobial assay of BA in a microemulsion system..	120
Cytotoxicity assay of BA in microemulsion system.....	124
IV CONCLUSION	127
BIBLIOGRAPHY.....	129
APPENDIX A.....	136
APPENDIX B.....	146
APPENDIX C.....	157
BIODATA OF AUTHOR.....	158

LIST OF TABLES

Table	Page
1 Some medicinal values of betulinic acid.....	26
2 Chemicals used, their sources and purity.....	33
3 Conductivity of binary mixtures of MA and MS.....	90
4 Solubility of betulinic acid in some aliphatic alcohol.....	109
5 Solubility of betulinic acid in organic solvents and surfactants.....	109
6 Activity of betulinic acid (BA) in DMSO against various microbes.....	121
7 Activity of microemulsion samples with 0.01 wt% betulinic acid on microbial.....	122
8 Viability of microemulsion samples against cell CEM-SS.....	124

LIST OF FIGURES

Figure	Page
1 Structure of a surfactant.....	2
2 Classification of surfactants.....	3
3 Some uses of surfactants (Karsa, 1987).....	4
4 Behaviour of a surfactants in solution (Benson, 1983).....	6
5 Structures of normal and reversed micelles (Gruen, 1983).....	7
6 Possible states of an amphiphile in solution (Myer, 1988).....	8
7 Structures of (a) inverse micelle (b) w/o microemulsion and (c) w/o emulsion.....	13
8 Structures of (a) normal micelle (b) o/w microemulsion and (c) o/w emulsion.....	14
9 Approximate size ranges for surfactant assemblies and related structures (Tadros, 1984).....	16
10 Some applications of microemulsions.....	19
11 Quaternary phase diagram.....	24
12 Chemical structure of betulinic acid (BA).....	25
13 Possibles sites of incorporation of solubilisate in a micelle; (a) in the hydrocarbon core; (b) short penetration of the palisade layer; (c) deep penetration of the palisade layer; (d) adsorption on the surface of the micelle; (e) in the polyoxyethylene shell of the micelle of a nonionic detergent (a to d) (Attwodd and Florence, 1983).....	29
14 Diagram of Experimental Design.....	34
15 Construction of both w/o microemulsion region in the presence of esters, namely, methyl acetate (MA) and ethyl benzoate (EB in: a) aqueous (water; b) nonaqueous (glycerol) systems at 25.0 ± 0.5°C.....	38

16	Preparation of sample for antimicrobial and antifungus testing by using ELISA.....	44
17	Flow chart of preparation samples for cytotoxicity screening and table of calculation the amount of betulinic acid.....	46
18	Phase diagram of MA/Tween 80/water system at $25.0 \pm 0.5^{\circ}\text{C}$. L ₁ = o/w microemulsion, L ₂ = w/o microemulsion; D = lamellar liquid crystal region; and T- two-phase region.....	49
19	Phase diagram of MA/Brij 30/ water system at $25.0 \pm 0.5^{\circ}\text{C}$. L ₂ = w/o microemulsion; D = lamellar liquid crystal region; and T- two-phase region.....	50
20	Phase diagrams of a)—— -MA/Tween 80/ water system; b)----- - MA/Brij 30/water system at $25.0 \pm 0.5^{\circ}\text{C}$. L ₁ = o/w microemulsion, L ₂ = w/o microemulsion; D = lamellar liquid crystal region; and T- two-phase region.....	52
21	Phase diagrams of Tween 80/Brij 30/ water system at $25.0 \pm 0.5^{\circ}\text{C}$. L = isotropic region; D = lamellar liquid crystal region; E = hexagonal liquid crystal region; and T =two-phase region.....	54
22	Phase diagrams of MA/Tween 80:Brij 30 (90:10)/water system at $25.0 \pm 0.5^{\circ}\text{C}$. L = isotropic region; and T =two-phase region...	55
23	Phase diagrams of MA/Tween 80:Brij 30 (80:20)/water system at $25.0 \pm 0.5^{\circ}\text{C}$. L = isotropic region; and T =two-phase region...	57
24	Phase diagrams of MA/Tween 80:Brij 30(60:40)/ water system at $25.0 \pm 0.5^{\circ}\text{C}$. L = isotropic region; D = lamellar liquid crystal region; and T =two-phase region.....	58
25	Phase diagrams of a)—— - MA/Tween80:Brij 30(90:10)/ water system b) ----- - MA/Tween80:Brij 30(80:20)/ water system and c) --- MA/Tween 80:Brij 30(60:40)/ water system at $25.0 \pm 0.5^{\circ}\text{C}$. L = isotropic region; D = lamellar liquid crystal region; and T =two-phase region.....	59
26	Phase diagrams of a)—— - EB/Tween 80/ water and b)----- - EB/Brij 30/water system at $25.0 \pm 0.5^{\circ}\text{C}$. L ₂ = w/o microemulsion; and T = Two-phase region.....	61
27	Phase diagrams of EB/Tween 80:Brij 30(90:10)/ water system at $25.0 \pm 0.5^{\circ}\text{C}$. L ₂ = w/o microemulsion; E = hexagonal liquid crystal region; and T = two-phase region.....	62

28	Phase diagrams of EB/Tween 80:Brij 30(80:20)/ water system at $25.0 \pm 0.5^\circ\text{C}$. L_1 = o/w microemulsion, L_2 = w/o microemulsion; D = lamellar liquid crystal region; and T = two-phase region.....	63
29	Phase diagrams of EB/Tween 80:Brij 30(60:40)/water system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = w/o microemulsion; D = lamellar liquid crystal region; and T = two-phase region.....	65
30	Phase diagrams of a) — - EB/Tween 80:Brij 30(90:10)/ water system; b) -EB/Tween 80:Brij 30(80:20)/ water system; and c) --- - EB/Tween 80:Brij 30(60:40)/ water system at $25.0 \pm 0.5^\circ\text{C}$. L_1 = o/w microemulsion, L_2 = w/o microemulsion; D = lamellar liquid crystal; E= hexagonal liquid crystal region; and T = two-phase region.....	67
31	Phase diagrams of a) ____ - MA/Tween 80/water system; EB/Tween 80/water system; b)____ - MA/Brij 30/water system; and -EB/Brij 30/ water system at $25.0 \pm 0.5^\circ\text{C}$. L_1 = w/o microemulsion; L_2 = o/w microemulsion; D – lamellar liquid crystal region; and T- two-phase region.....	68
32	Phase diagrams of — - MA/Tween 80:Brij 30/ water system; -EB/Tween 80:Brij 30/ water system at $25.0 \pm 0.5^\circ\text{C}$. L_1 = w/o microemulsion; L_2 = o/w microemulsion; D = lamellar liquid crystal region; E = hexagonal liquid crystal region; and T = two-phase region.....	69
33	Phase diagram of MA/Tween 80/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = Two-phase region.....	71
34	Phase diagram of MA/Brij 30/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L = Isotropic region; and T = two-phase region.....	72
35	Phase diagrams of — - MA/Tween 80/glycerol system; and - MA/Brij 30/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region.....	73
36	Phase diagram of Tween 80/Brij 30/glycerol system at $25.0 \pm 0.5^\circ\text{C}$ where L – Isotropic region; and T- Two-phase region.....	74
37	Phase diagram of MA/Tween 80:Brij 30(90:10)/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region.....	76

38	Phase diagram of MA/Tween 80:Brij 30(80:20)/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region.....	77
39	Phase diagram of MA/Tween 80:Brij 30(60:40)/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region.....	78
40	Phase diagrams of a)—— MA/Tween 80:Brij 30(90:10)/glycerol system; b)..... - MA/Tween 80:Brij 30(80:20)/glycerol system; and c) --- - MA/Tween 80:Brij 30(60:40)/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region.....	79
41	Phase diagram of EB/Tween 80/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = Two-phase region.....	80
42	Phase diagram of EB/Brij 30/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region.....	81
43	Phase diagram of EB/Tween 80:Brij 30(90:10)/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region.....	83
44	Phase diagrams of EB/Tween 80:Brij 30(80:20)/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region.....	84
45	Phase diagrams of EB/Tween 80:Brij 30(60:40)/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region	85
46	Phase diagrams of a)—— - EB/Tween 80:Brij 30(90:10)/glycerol system; b)..... - EB/Tween 80:Brij 30(80:20)/glycerol system; and c)--- - EB/Tween 80:Brij 30(60:40)/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L_2 = o/w microemulsion; and T = two-phase region.....	86
47	Phase diagrams of a)—— - MA/Tween 80/glycerol system;- EB/Tween 80/glycerol system; and b) —— MA/Brij 30/glycerol system; - EB/Brij 30/glycerol system at $25.0 \pm 0.5^\circ\text{C}$. L = Isotropic region; and T = two-phase region.....	87
48	Phase diagrams of ____ - MA/Tween 80:Brij 30/glycerol system; - EB/Tween 80:Brij 30/glycerol system at $25.0 \pm 0.5^\circ\text{C}$ where L – Isotropic region; and T- Two-phase region.....	88

49	Conductivity of MA/Tween 80:Brij 30 (90:10)/water system as a function weight percentage of water at constant MA-to-MS weight ratio, r at $25.0 \pm 0.5^\circ\text{C}$	91
50	Variation of w_o (water, wt%) and conductivity, K with weight ratio with MA:Mixed Surfactants (MA/MS) of MA/Tween 80:Brij 30(90:10)/water system at $25.0 \pm 0.5^\circ\text{C}$ at a) Percolation Threshold; b) Bicontinuous Structure and c) Micellar Region.....	93
51	Conductivity of MA/Tween 80:Brij 30(80:20)/water system as a function weight percentage of water at constant MA-to-MS weight ratio, r at $25.0 \pm 0.5^\circ\text{C}$	95
52	Variation of w_o (water, wt%) and conductivity, K with weight ratio with MA:Mixed Surfactants (MA/MS) of MA/Tween 80:Brij 30(80:20)/water system at $25.0 \pm 0.5^\circ\text{C}$ at a) Percolation Threshold; b) Bicontinuous Structure and c) Micellar Region.....	96
53	Conductivity of MA/Tween 80:Brij 30(60:40)/water system as a function weight percentage of water at constant MA-to-MS weight ratio, r at $25.0 \pm 0.5^\circ\text{C}$	97
54	Variation of w_o (water, wt%) and conductivity, K with weight ratio with MA:Mixed Surfactants (MA/MS) of MA/Tween 80:Brij 30(60:40)/water system at $25.0 \pm 0.5^\circ\text{C}$ at a) Percolation Threshold; b) Bicontinuous Structure and c) Micellar Region.....	99
55	Conductivity, K at Percolation Threshold with weight ratio with MA:Mixed Surfactants (MA/MS) of a) MA/Tween 80:Brij 30(90:10)/water system; b) MA/Tween 80:Brij 30(80:20)/water system; and c) MA/Tween 80:Brij 30(60:40)/ water system at $25.0 \pm 0.5^\circ\text{C}$	100
56	Conductivity, K at Bicontinuous structure with weight ratio with MA:Mixed Surfactants (MA/MS) of a) MA/Tween 80:Brij 30(90:10)/water system; b) MA/Tween 80:Brij 30(80:20)/water system; and c) MA/Tween 80:Brij 30(60:40)/water system at $25.0 \pm 0.5^\circ\text{C}$	101
57	Conductivity, K at Micellar region with weight ratio with MA:Mixed Surfactants (MA/MS) of a) MA/Tween 80 : Brij 30 (90:10)/water system; b) MA/Tween 80:Brij 30(80:20)/water system; and c) MA/Tween 80:Brij 30(60:40)/water system at $25.0 \pm 0.5^\circ\text{C}$	102

58	Plots of Z_{ave} against water weight percentage (w) at different MA and MS weight ratios of MA/Tween 80:Brij 30(80:20)/water system at $25.0 \pm 0.5^{\circ}\text{C}$	104
59	Plots of Z_{ave} against water weight percentage (w) at different MA and MS weight ratios of MA/Tween 80:Brij 30(60:40)/water system at $25.0 \pm 0.5^{\circ}\text{C}$	106
60	Solubility of BA in MA/Tween 80:Brij 30(90:10)/water system at different methyl acetate, MA and mixed surfactants, MS weight ratios as water weight percentage at $25.0 \pm 0.5^{\circ}\text{C}$	111
61	Solubility of BA in MA/Tween 80:Brij 30(80:20)/water system at different methyl acetate, MA and mixed surfactants, MS weight ratios as water weight percentage at $25.0 \pm 0.5^{\circ}\text{C}$	113
62	Solubility of BA in MA/Tween 80:Brij 30(60:40)/water system at different methyl acetate, MA and mixed surfactants, MS weight ratios as water weight percentage at $25.0 \pm 0.5^{\circ}\text{C}$	114
63	Solubility of BA in MA/Tween 80:Brij 30/water [60/40/water] at a) 90:10; b) 80:20; and c) 60:40 weight ratio of Tween 80 and Brij 30 system at different MA and MS weight ratios as water weight percentage at $25.0 \pm 0.5^{\circ}\text{C}$	116
64	Solubility of BA in MA/Tween 80:Brij 30/water [50/50/ water] at a) 90:10; b) 80:20; and c) 60:40 weight ratio of Tween 80 and Brij 30 system at different MA and MS weight ratios as water weight percentage at $25.0 \pm 0.5^{\circ}\text{C}$	117
65	Solubility of BA in MA/Tween 80:Brij 30/water [35/65/ water] at a) 90:10; b) 80:20; and c) 60:40 weight ratio of Tween 80 and Brij 30 system at different MA and MS weight ratios as water weight percentage at $25.0 \pm 0.5^{\circ}\text{C}$	118
66	Solubility of BA in MA/Tween 80:Brij 30/water [20/80/ water] at a) 90:10; b) 80:20; and c) 60:40 weight ratio of Tween 80 and Brij 30 system at different MA and MS weight ratios as water weight percentage at $25.0 \pm 0.5^{\circ}\text{C}$	119
67	Screening BA cytotoxicity in DMSO against CEM-SS cells	126

LIST OF ABBREVIATIONS

BA	Betulinic acid
MA	Methyl acetate
H ₂ O	Deionised water
MS	Mixed surfactants
DMSO	Dimethyl sulphoxide
ATCC	American Type Culture Collection
NRRL	Northern Regional Research
EB	Ethyl benzoate
PDA	Potato Dextrose Agar
PDB	Potato Dextrose Broth
NA	Nutrient Agar
NB	Nutrient Broth
MTT	Micro-culture Tetrazolium Salt
W/O	water-in-oil
O/W	oil-in-water
OD	Optical Density
CEM-SS	T-Lymphoblastic Leukemia cell
IC ₅₀	Inhibitory Concentration at 50% cells reduction
L	Lamellar liquid crystal
D	Cubic liquid crystal
E	Hexagonal liquid crystal
L ₁	Normal micelle (o/w microemulsion)

L ₂	Reverse micelle (w/o microemulsion)
HLB	Hydrophilic Lipophilic Balance
T	Two-phase region

CHAPTER 1

INTRODUCTION

A surface active agent, or ‘surfactant’, is a chemical compound with a polar or ionic head and a hydrocarbon tail (Figure 1) (Ottewil, 1983; Benson, 1983; Lucassen, 1981). The hydrocarbon tail is hydrophobic and can be either linear or branched. The polar, or ionic, head is hydrophilic, interacting strongly with the water molecules with hydrogen bonding and dispersing them via dipole-dipole or ion-dipole interaction. A structure having both a hydrophobic and hydrophilic moieties is said to be ‘amphiphatic’.

Surfactants are classified by their hydrophilic heads into four types - ionic (either cationic or anionic) and nonionic (either zwitterionic or amphoteric) (Figure 2).

The balance between the hydrophobic and hydrophilic portions of a surfactant confers it its unique properties in use (Karsa, 1987; Myers, 1990). The applications of surfactants are many, in both everyday household products and personnel care, and in the industrial production and processing of materials (Figure 3).




Figure 1: Structure of a surfactant