



UNIVERSITI PUTRA MALAYSIA

**THE KINETICS OF UPTAKE AND RELEASE OF
POLYCYCLIC AROMATIC HYDROCARBONS IN THE GREEN
MUSSELS *Perna viridis* FOR BIOMONITORING OF MARINE
POLLUTION**

ALI MASHINCHIAN MORADI

FSAS 2001 16

**THE KINETICS OF UPTAKE AND RELEASE OF
POLYCYCLIC AROMATIC HYDROCARBONS IN THE GREEN MUSSELS
Perna viridis FOR BIOMONITORING OF MARINE POLLUTION**

By

ALI MASHINCHIAN MORADI

**Thesis Submitted in Fulfilment of the Requirement for the Degree of
Doctor of Philosophy in the Graduate School
Universiti Putra Malaysia**

December 2001



DEDICATION

To my dear family, my parents, wife and sisters who have been my source of inspiration, wisdom and strength through the most difficult times of my life.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

**THE KINETICS OF UPTAKE AND RELEASE OF
POLYCYCLIC AROMATIC HYDROCARBONS IN THE GREEN MUSSELS
Perna viridis FOR BIOMONITORING OF MARINE POLLUTION**

By

ALI MASHINCHIAN MORADI

December 2001

Chairman: Associate Professor Jambari Hj. Ali, Ph.D.

Faculty: Science and Environmental Studies

The kinetics of accumulation and depuration of polycyclic aromatic hydrocarbons (PAHs) in green mussel *Perna viridis* has never been investigated under field conditions. In this study, green mussels were used to assess the spatial distribution of PAH pollution along the west coast of Peninsular Malaysia, and to determine PAHs uptake and release kinetics in this species under field conditions. In addition, the effect of sex and exposure duration on the uptake and release kinetics of these contaminants were investigated. Overall more than 1500 individual mussels were collected, soft tissues were homogenized, extracted, cleaned up, fractionated and concentration of 17 individual PAH compounds were analysed using Gas Chromatography-Mass Spectrometry (GC-MS). Among the eleven stations sampled in 1997 and 1998 along the Straits of Malacca, mussels from Pasir Puteh (01° 26.05' N and 103° 55.94' E) had the highest (2855-3450 ng/g lipid weight) and from Pasir Panjang (02° 24.96' N and 101° 56.54' E) the lowest (278-366 ng/g lipid weight) levels of PAHs. As a result, Pasir Puteh was selected as the "hot-spot" and Pasir



Panjang as the "clean site" to further conduct the kinetics experiments. The PAH levels in mussels from Malaysian coastlines in comparison with the polluted areas worldwide were found to be relatively low. At the end of both uptake and release experiments (28 and 36 days, respectively), PAH levels in transplanted and back-transplanted mussels were not significantly different ($p>0.05$) from those found in natives (controls in Pasir Puteh and Pasir Panjang, respectively). It was found that sex did not make a significant difference ($p>0.05$) in the uptake and release rates. Similarly, duration of exposure did not significantly affect the release rates. However, significant differences ($p<0.05$) were found both in uptake and release rates when target PAHs were divided into three, four and five to seven-ring compounds. In general, rapid rates of uptake and release of relatively less lipophilic PAHs were observed, indicating that the primary mechanism of bioaccumulation of these lipophilic pollutants in *Perna viridis* complies with the concept of lipid-water partitioning equilibrium. The biological half-life of individual PAHs ranged from 7 to 29 days. This study illustrated that green mussels are a useful tool in monitoring of PAHs. However, due to relatively rapid rates of uptake and release of these compounds, to get an average level of PAHs over a long period, "time-bulking" should be considered for future biomonitoring programmes using this species.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**KINETIK PENGAMBILAN DAN
PELEPASAN POLISIKLIK AROMATIK HIDROKARBON (PAHs)
DI DALAM KUPANG *Perna viridis*
BAGI PEMANTAUAN BIOLOGI PENCEMARAN MARIN**

Oleh

ALI MASHINCHIAN MORADI

Disember 2001

Pengerusi: Profesor Madaya Jambari Hj. Ali, Ph.D.

Fakulti: Sains dan Pengajian Alam Sekitar

Kinetik pengumpulan dan depurasi polisiklik aromatik hidrokarbon (PAHs) di dalam kupang *Perna viridis* tidak pernah dikaji di bawah keadaan lapangan. Dalam kajian ini, kupang telah digunakan untuk mengenalpasti ruang taburan pencemaran PAHs di sepanjang persisiran pantai barat Semenanjung Malaysia, dan menentukan kinetik pengambilan dan pelepasan bagi PAHs di dalam spesies ini di bawah keadaan lapangan. Disamping itu, kesan jantina dan jangkamasa pendedahan kepada kinetik pengambilan dan pelepasan terhadap pencemar ini telah dilakukan. Keseluruhannya, lebih daripada 1500 individu kupang telah disampel, tisu kupang telah dihomogenat, diekstrak, dicuci, dipecah dan kepekatan bagi 17 individu sebatian PAHs telah dianalisis dengan menggunakan gas kromatografi spektrometri jisim (GC-MS). Keputusan bagi fasa satu menunjukkan antara 11 stesen yang disampel di Selat Melaka, kupang dari Kg. Pasir Puteh (01° 26.05' N dan 103° 55.94' E) mempunyai nilai tertinggi (2855-3450 ng/g berat lipid) dan Pasir Panjang (02° 24.96' N dan

101° 56.54' E) mempunyai nilai terendah (278-366 ng/g berat lipid) bagi tahap PAHs yang dikaji. Oleh itu, Kg. Pasir Puteh telah dipilih sebagai "stesen tercemar" dan Pasir Panjang sebagai "stesen bersih" untuk eksperimen kinetik. Keputusan kajian menunjukkan tahap PAHs dalam tisu kupang di persisiran pantai Malaysia adalah rendah jika dibandingkan dengan tahap di kawasan tercemar lain di dunia. Pada akhir eksperimen biopenimbunan dan depurasi (28 hari, dan 36 hari) tahap PAHs bagi kupang yang dipindah dan yang dikembalikan ke tempat asal tidak mencapai tahap yang signifikan ($p > 0.05$) berbanding dengan kupang asal (kawalan - Kg. Pasir Puteh dan Pasir Panjang). Keputusan kajian telah menunjukkan jantina tidak menunjukkan perbezaan yang signifikan ($p > 0.05$) dalam kadar pengambilan dan pelepasan. Pada masa yang sama, masa pendedahan tidak memberi kesan ke atas kadar pelepasan. Bagaimanapun, perbezaan yang signifikan ($p < 0.05$) telah didapati bagi kadar pengambilan apabila PAHs sasaran dibahagikan kepada sebatian gegelang tiga, empat, lima hingga tujuh. Secara amnya, kadar pengambilan dan pelepasan yang cepat bagi PAHs telah berlaku bagi PAHs yang lebih lipofilik. Ini telah menunjukkan bahawa mekanisme primer untuk pencemar lipofilik didalam *Perna viridis* menepati konsep keseimbangan pembahagian "lipid-air" Separuh hayat biologi individu PAHs berjulat 7 hingga 29 hari. Kajian in menunjukkan kupang adalah bahan yang sangat berguna untuk pemantauan pemantauan. Bagaimanapun, oleh kerana kadar pengumpulan dan pelepasan yang cepat bagi sebatian tersebut demi mendapatkan tahap hitung panjang PAHs "time-bulking" haruslah diambilkira dalam kajian pemantauan PAHs dimasa hadapan.

ACKNOWLEDGEMENTS

I am most indebted and grateful to Mr. Mohamad Pauzi Zakaria who has guided and supported me along the way and given me a lot of constructive comments, advice, and ideas, and finally for his kind friendship. He is certainly the person who is responsible for the successful completion of this study.

I would like to extend my gratitude to my supervisor, Associate Professor Dr. Jambari Hj. Ali, for his advice, kind and constant support, encouragement and omnipresence at all my difficulties. My appreciation also goes to the members of my Supervisory Committee, Professor Dr. Fatimah Md. Yusoff and Associate Professor, Dr. Mohamad Ismail Yaziz for their kind support.

This research was made possible through the financial support of the Ministry of Science, Technology and Environment, Malaysia (IRPA - Intensification of Research Priority Areas, project no. 0802040013) which is highly appreciated.

Special thanks to my spouse for all her patience, kind help and morale support. Many thanks to Yap Chee Kong and Hamid Assilzadeh and all my other friends who contributed so much indirectly towards this achievement.



TABLE OF CONTENTS

DEDICATION.....	ii
ABSTRACT.....	iii
ABSTRAK.....	v
ACKNOWLEDGEMENTS.....	vii
APPROVAL.....	viii
DECLARATION.....	x
LIST OF TABLES.....	xiii
LIST OF FIGURES.....	xviii
LIST OF PLATES.....	xxii
LIST OF ABBREVIATIONS.....	xxiii

CHAPTER

1 INTRODUCTION.....	1
1.1 Background of Study.....	1
1.2 Problem to be Addressed.....	2
1.3 Significance of the Study.....	3
1.4 Purpose of Study.....	5
1.5 Study Objectives.....	6
2 LITRATURE REVIEW.....	8
2.1 Polycyclic Aromatic Hydrocarbons (PAHs).....	8
2.2.1 Sources of PAHs in the Aquatic Environment.....	9
2.2.2 Distribution and Fate of PAHs in the Aquatic Environment.....	10
2.2.3 Adverse Biological Effects of PAHs.....	13
2.2 Bioaccumulation.....	15
2.3 Biomonitoring of the Aquatic Environment.....	16
2.3.1 Bioindicators.....	17
2.3.2 Mussel Watch Concept.....	20
2.3.3 Transplantation of Biomonitors.....	21
2.4 <i>Perna viridis</i> as a Bioindicator Organism.....	22
2.4.1 Taxonomy.....	23
2.4.2 Morphology.....	23
2.4.3 Habitat and Distribution.....	24
2.4.4 Food.....	25
2.4.5 Reproduction.....	25
2.5 Uptake, Retention and Release of Organic Pollutants in Bivalves.....	27
3 MATERIALS AND METHODS.....	32
3.1 Analytical Method Development.....	32
3.1.1 GC-MS SIM-MODE (Selective Ion Monitoring).....	32
3.1.2 Calibration Curves and Linearity of the Response Factors.....	36
3.1.3 Determination of Elusion Volume for Silica-Gel Column Chromatography.....	40
3.1.3.1 Clean-up Step.....	40



3.1.3.2 Fractionation Step.....	42
3.2 Biomonitoring of the Coastal Waters of the Straits of Malacca.....	44
3.2.1 Sampling.....	44
3.2.1.1 Sampling Locations.....	44
3.2.1.2 Individual Mussel Size.....	47
3.2.1.3 Sample Size.....	47
3.2.1.4 Collection Time.....	48
3.2.3 Sample Storage and Transport.....	49
3.3 Transplantation and Sample Collection for the Kinetics Studies.....	49
3.3.1 Transplantation of Mussels.....	49
3.3.2 Back-transplantation.....	54
3.4 Analytical Procedure.....	56
3.4.1 Sample Preparation.....	56
3.4.1.1 Apparatus Cleaning.....	56
3.4.1.2 Homogenization.....	56
3.4.1.3 Extraction.....	57
3.4.1.4 Clean-up of the Sample Extracts.....	58
3.4.1.5 Fractionation.....	59
3.4.2 GC-MS Analysis.....	60
3.4.3 Calculation of Concentrations in the Samples.....	62
4 RESULTS AND DISCUSSION.....	65
4.1 Biomonitoring of the Coastal Waters of the Straits of Malacca.....	65
4.1.1 The First Stage of Biomonitoring.....	65
4.1.2 The Second Stage of Biomonitoring.....	72
4.1.3 Comparison of Results of the First and Second Stages of Biomonitoring.....	78
4.1.4 Identification of PAH Sources in the Sampling Sites.....	81
4.1.5 Comparison of the Results with Worldwide PAH concentrations in mussels.....	87
4.2 The Kinetics of Uptake of PAHs in the Mussel <i>Perna viridis</i>	90
4.2.1 The Uptake Kinetics of PAHs in Male Mussels.....	92
4.2.2 The Uptake Kinetics in Female Mussels.....	110
4.2.3 Comparison of Uptake Rates in Male and Female Mussels.....	121
4.2.4 Effect of Environmental Factors on the Uptake Kinetics.....	123
4.3 The Kinetics of Release of PAHs in the Mussel <i>Perna viridis</i>	130
4.3.1 The Release Kinetics of PAHs after a Short-term Exposure.....	133
4.3.2 The Release Kinetics of PAHs after a Long-term Exposure.....	151
4.3.3 Effect of Sex and Duration of Exposure on the Release Kinetics.....	165
4.3.4 Effect of Environmental Factors on the Release Kinetics.....	170
5 CONCLUSION AND RECOMMENDATIONS.....	175
REFERENCES.....	182
VITA.....	203



LIST OF TABLES

Table		Page
2.1	Results of different hydrocarbon uptake/depuration studies using bivalves.....	31
3.1	Target PAH compounds and their molecular weight, log <i>K_{ow}</i> (octanol-water partition coefficient) and water solubility.....	34
3.2	Selected character ions and time intervals for developing SIM-MODE for PAH target compounds analysis	35
3.3	Response factor (peak area) of individual PAH compounds with different concentrations.....	38
3.4	Description of sampling sites along the west coast of Peninsular Malaysia.....	46
3.5	Retention time, peak area and peak height of PAH compounds in one of the mussel tissue samples analysed in the SIM-MODE.....	61a
3.6	Relative standard deviations(RSD) of concentrations of PAH compounds in four replications of mussel samples from Pasir Puteh.....	61b
4.1	Biological details of green mussels collected in the first stage of biomonitoring along the Straits of Malacca.....	66
4.2	Concentration of 17 individual PAH compounds (ng/g lipid weight) in mussels from different stations in the first stage of biomonitoring.....	69
4.3	Concentration of 17 individual PAH compounds (ng/g wet weight) in mussels from different stations in the first stage of biomonitoring.....	70
4.4	Biological details of green mussels collected in the second stage of biomonitoring along the Straits of Malacca.....	72



4.5	Concentration of 17 individual PAH compounds (ng/g lipid weight) in mussels from different stations in the second stage of biomonitoring.....	74
4.6	Concentration of 17 individual PAH compounds (ng/g wet weight) in mussels from different stations in the second stage of biomonitoring.....	75
4.7	Concentration of individual PAHs in mussel samples (ng/g wet weight.) collected in the first and second stages of biomonitoring.....	79
4.8	PAHs composition for mussels, crude oil, lubricating oil, used crankcase oil, street dust and atmospheric aerosol.....	82
4.9	Concentrations of PAHs in the soft tissues of mussels from different parts of the world.....	88
4.10	Biological details of mussels <i>P. viridis</i> collected during the uptake experiment.....	91
4.11	Concentrations of individual PAHs (ng/g wet wt.) in the transplanted male mussels during the uptake experiment.....	93
4.12	Concentrations of individual PAHs (ng/g lipid wt.) in the transplanted male mussels during the uptake experiment.....	94
4.13	Concentrations of individual PAHs (ng/g wet wt.) in native male mussels (control-Pasir Puteh) during the uptake experiment.....	95
4.14	Concentrations of individual PAHs (ng/g lipid wt.) in native male mussels (control-Pasir Puteh) during the uptake experiment.....	95
4.15	The uptake rate constants (k_1) of the male mussels transplanted in Pasir Puteh.....	105
4.16	Bioconcentration Factor values in male mussels for individual PAH compounds both from field measurements [BCF (Cm/Cw)] and from the kinetics model [BCF (k_1/k_2)]......	106



4.17	Concentrations of individual PAHs (ng/g wet wt.) in the transplanted female mussels during the uptake experiment.....	111
4.18	Concentrations of individual PAHs (ng/g lipid wt.) in the transplanted female mussels during the uptake experiment.....	112
4.19	Concentrations of individual PAHs (ng/g wet wt.) in the native female mussels (control-Pasir Puteh) during the uptake experiment.....	113
4.20	Concentrations of individual PAHs (ng/g lipid wt.) in native female mussels (control-Pasir Puteh) during the uptake experiment.....	113
4.21	The uptake rate constants (k_1) in the female mussels transplanted in Pasir Puteh.....	117
4.22	Bioconcentration Factor values in female mussels for individual PAH compounds from field measurements [BCF (Cm/Cw)] and from the kinetics model [BCF (k_1/k_2)]......	119
4.23	Physico-chemical parameters of seawater (mean \pm S.D., n = 3) in Pasir Puteh during the uptake experiment.....	124
4.24	Correlation coefficients (r) between the variation of total PAH concentrations and physico-chemical parameters during the uptake experiment in Pasir Puteh.....	126
4.25	Concentrations of PAHs in the seawater (ng/ml), and sediments (ng/g dry wt.) from Pasir Puteh.....	127
4.26	Biological details of short-term exposed mussel samples collected during the release experiment.....	131
4.27	Biological details of long-term exposed mussel samples collected during the release experiment.....	132
4.28	Concentrations of individual PAHs (ng/g wet wt.) in the back-transplanted (short-term exposed) male mussels during the release experiment.....	134



4.29	Concentrations of individual PAHs (ng/g wet wt.) in the back-transplanted (short-term exposed) female mussels during the release experiment.....	134
4.30	Concentrations of individual PAHs (ng/g lipid wt.) in the back-transplanted (short-term exposed) male mussels during the release experiment.....	135
4.31	Concentrations of individual PAHs (ng/g lipid wt.) in the back-transplanted (short-term exposed) female mussels during the release experiment.....	135
4.32	Concentrations of individual PAHs (ng/g wet wt.) in native mussels (control-Pasir Panjang) during the release experiment.....	137
4.33	Concentrations of individual PAHs (ng/g lipid wt.) in native mussels (control-Pasir Panjang) during the release experiment.....	137
4.34	The release rate constants (k_2), biological half-lives (BHL), days to 90% uptake equilibrium ($t_{(90\%)}$) and correlation coefficients (r) in the equation used to obtain k_2 in male mussels, after a short-term exposure..	145
4.35	The release rate constants (k_2), biological half-lives (BHL), days to 90% uptake equilibrium ($t_{(90\%)}$) and correlation coefficients (r) in the equation used to obtain k_2 in female mussels, after a short-term exposure.....	146
4.36	Concentrations of individual PAHs (ng/g wet wt.) in the long-term exposed transplanted male mussels during the release experiment.....	152
4.37	Concentrations of individual PAHs (ng/g wet wt.) in the long-term exposed transplanted female mussels during the release experiment.....	152
4.38	Concentrations of individual PAHs (ng/g lipid wt.) in the long-term exposed transplanted male mussels during the release experiment.....	153
4.39	Concentrations of individual PAHs (ng/g lipid wt.) in the long-term exposed transplanted female mussels during the release experiment.....	153



4.40	The release rate constants (k_2), biological half-lives (BHL), days to reach 90% uptake equilibrium ($t_{(90\%)}$) and correlation coefficients (r) in the equation used to obtain k_2 in long-term exposed male mussels.....	161
4.41	The release rate constants (k_2), biological half-lives (BHL), days to 90% uptake equilibrium ($t_{(90\%)}$) and correlation coefficient (r) in the equation used to obtain k_2 in long-term exposed female mussels.....	162
4.42	Physico-chemical parameters of seawater (mean \pm S.D., $n = 3$) in Pasir Panjang during the release experiment.....	170
4.43	Correlation coefficients (r) between the variation of total PAH concentrations in mussels and physico-chemical parameters during the release experiment in Pasir Panjang.....	172
4.44	Concentrations of PAHs in the seawater (ng/ml), and sediments (ng/g dry wt.) from Pasir Panjang.....	173
4.45	Correlation coefficients (r) between individual PAH concentrations in long-term and short-term exposed male and female mussels and their concentrations in seawater and sediments from Pasir Panjang.....	173



LIST OF FIGURES

Figure		Page
3.1	Molecular structure of analysed PAH compounds.....	33
3.2	Chromatograms of 17 PAH standard mixtures with different concentrations (0.1, 0.25, 0.5 and 0.75 ppm) and constant concentration of internal injection standard -IIS (1 ppm).....	37
3.3	Calibration curves for the individual PAH compounds: Relative response of each compound versus its concentration.....	39
3.4	Sampling stations along the coastline of West Peninsular Malaysia.....	46
3.5	Experimental design for mussel transplantation and sampling for the kinetics studies.....	55
3.6	The analytical procedure of PAHs measurement in mussel samples.....	61
3.7	An example of GC-MS chromatogram of PAH compounds in one of the mussel tissue samples analysed in the SIM MODE.....	61a
3.8	Typical GC-MS chromatogram of a PAH standard mixture.....	62
3.8	Schematic chromatograms and peaks in the standard mixture solution (a) and sample solution (b), (D is peak of deuterated internal injection standard (IIS) and S is peak of one of the target compounds).....	63
4.1	Concentration of total PAHs in soft tissue of mussels from different stations in the first stage of biomonitoring.....	67
4.2	Profiles of PAH concentrations (ng/g lipid wt.) in mussel samples from different sampling sites in the first stage of biomonitoring.....	71
4.3	Concentration of total PAHs in soft tissue of mussels from different stations in the second stage of biomonitoring.....	73



4.5	Concentrations of total PAHs in mussels from the first and second stages of biomonitoring.....	78
4.6	Average relative abundance of individual PAH compounds in mussels from all sampling sites (first and second stages of biomonitoring).....	80
4.7	PAHs profiles for sediments, crude oil, fresh lubricating oil, used crankcase oil, street dust and aerosols in Malaysia (data from Zakaria <i>et al.</i> , in preparation).....	84
4.8	Total PAHs concentration (sum of concentrations of 17 individual target compounds) in the transplanted and native male mussels during the uptake experiment.	96
4.9	The uptake trends of three-ring, four-ring and five to seven ring PAHs in male mussels during the exposure period.....	97
4.10	Concentration of individual PAHs (ng/g lipid wt.) in the transplanted and native male mussels in Pasir Puteh during the uptake experiment.....	99
4.11	Correlation of BCF values from field measurements (C_m/C_w) versus the kinetics model (k_1/k_2) in male mussels.....	107
4.12	Correlation between octanol-water partition coefficients (K_{ow}) and (a) biological concentration factors from field measurements (C_m/C_w), (b) biological concentration factors from the kinetics model (k_1/k_2), and (c) uptake rate constants (k_1) in male mussels.....	108
4.13	Total PAHs concentration (sum of concentrations of 17 individual target compounds) in the transplanted and native female mussels during the uptake experiment.....	110
4.14	The uptake trends of three-ring, four- ring and five to seven ring PAHs in female mussels during the exposure period.....	114



4.15	Concentrations of individual PAHs (ng/g lipid wt.) in the transplanted and native female mussels in Pasir Puteh during the uptake experiment.....	116
4.16	Correlation of BCF values from field measurements (C_m/C_w) versus values resulted from the kinetics model (k_1/k_2) in female mussels.....	119
4.17	Correlation between octanol-water partition coefficients (K_{ow}) and (a) biological concentration factors from field measurements (C_m/C_w), (b) biological concentration factors from the kinetics model (k_1/k_2), and (c) uptake rate constants (k_1) in female mussels.....	120
4.18	Comparison of the uptake rate values (k_1) in male and female mussels: (a) individual PAH compounds, and (b) grouped PAH compounds.....	122
4.19	Fluctuation of physico-chemical parameters in the seawater during the uptake period in Pasir Puteh.....	125
4.20	Correlation between concentration of individual PAH compounds in seawater (ng/ml) and in (a) male mussels and (b) female mussels (ng/g lipid wt.) at the end of the uptake period.....	128
4.21	Time trend of total PAHs concentration during the release experiment in female (a), and male (b), back-transplanted (short-term exposed) and native mussels (control-Pasir Panjang).....	138
4.22	The release trends of three-ring, four-ring and five to seven-ring PAHs in short-term exposed male mussels (a), and female mussels (b), during the release period.	140
4.23	Time trend of the concentration of individual PAHs (ng/g lipid wt.) in back-transplanted (short-term exposed) and native (control) male mussels during the release experiment.....	142
4.24	Time trend of the concentration of individual PAHs (ng/g lipid wt.) in back-transplanted (short-term exposed) and native (control) female mussels during the release experiment.....	143



4.25	Release rate constant (k_2) versus octanol-water partition coefficient (K_{ow}) for individual PAH compounds in male (a), and female mussels (b), after a short-term exposure.....	147
4.26	Time trend of total PAHs concentration during the release experiment in female (a) and male (b) transplanted and native mussels, after a long-term exposure in Pasir Panjang.....	155
4.27	The release trends of three-ring, four-ring and five to seven-ring PAHs in long-term exposed male mussels (a), and female mussels (b), during the release period.....	156
4.28	Time trend of the individual PAHs concentration (ng/g lipid wt.) in long-term exposed transplanted and native (control) male mussels, during the release experiment.	158
4.29	Time trend of the individual PAHs concentration (ng/g lipid wt.) in long-term exposed transplanted and native (control) female mussels, during the release experiment.....	159
4.30	Release rate constants (k_2) versus octanol-water partition coefficient (K_{ow}) for individual PAH compounds in male (a), and female mussels (b), after a long-term exposure.....	163
4.31	Comparison of release rate constants (k_2) in different sexes of mussels: Individual PAHs in long-term exposed (a), grouped PAHs in long term exposed (b), individual PAHs in short-term exposed (c), grouped PAHs in short-term exposed (d), male and female mussels.....	166
4.32	Comparison of release rate constants (k_2) after different exposure durations: Individual PAHs in male mussels (a), grouped PAHs in male mussels(b), individual PAHs in female mussels (c), grouped PAHs in female mussels (d).....	167
4.33	Fluctuation of physico-chemical parameters in the seawater during the release period in Pasir Panjang.....	171



LIST OF PLATES

Plate		Page
3.1	Pasir Puteh sampling site (the hot-spot) and its industrial hinterland.....	51
3.2	A cluster of native (long-term exposed) mussels from Pasir Puteh.	51
3.3	Pasir Panjang sampling site (the clean site).....	52
3.4	Native mussels at a mussel culture site in Pasir Panjang.....	52
3.5	Cages used for transplantation and back-transplantation of mussels (0.2 m × 1.0 m, mesh size = 2 cm).....	53
3.6	The tissue color of different sexes of mussel <i>Perna viridis</i> : Female in orange (left) and male in creamy color (right).....	53

LIST OF ABBREVIATIONS

PAHs:	Polycyclic aromatic hydrocarbons
BCF:	Biological concentration factor
BHL:	Biological half-life
$t_{90\%}$:	Time required to reach 90% uptake equilibrium
K_{ow} :	Octanol-water partition coefficient
k_1 :	Uptake rate constant
k_2 :	Release rate constant
GC-MS:	Gas chromatography mass spectrometry
IIS:	Internal injection standard
SIS:	Surrogate internal standard
Lipid wt.:	Lipid weight basis
Wet wt.:	Wet weight basis
DBT:	Dibenzothiophene
PHN:	Phenanthrene
ANT:	Anthracene
2MANT:	2-methyl anthracene
2MPHN:	2-methyl phenanthrene
FLU:	Fluoranthene
PYR:	Pyrene
RET:	Retene
1MPYR:	1-methyl pyrene
BaNT:	Benzo(a)anthracene
CHR:	Chrysene
BkFLU:	Benzo(k)fluoranthene
BeP:	Benzo(e)pyrene
BaP:	Benzo(a)pyrene
BeACPH:	Benzo(e)acephenanthrylene
DBahANT:	Dibenz(a,h)anthracene
COR:	Coronene



CHAPTER 1

INTRODUCTION

1.1 Background of Study

The coastal zones of West Peninsular Malaysia are heavily populated and lined by urban, industrial, agricultural areas and shipping ports (DOE, 1995). As a result, the coastal waters of this region receive a broad range of anthropogenic organic micropollutants from various land-based and marine-based sources (Law, 1994). Organic micropollutants are also transported from other regions via atmospheric and global ocean transport systems (Zubir, 1992; Iwata *et al.*, 1993). The Straits of Malacca is one of the busiest waterways in the world and subject to various types of pollution (Zubir, 1992; Law, 1994; DOE, 1995). Oil spills in the Straits of Malacca have been occurring very frequently in the past and have been affecting coastal ecosystems (Law, 1994; Zakaria *et al.*, 1999). Monitoring reports have shown that the coastal waters of West Peninsular Malaysia are polluted by organic contaminants (Sittig, 1985; WHO, 1989; Law and Ravinthar, 1989; Law *et al.*, 1990; Denker *et al.*, 1994; WHO, 1994). These substances may include polychlorinated biphenyls (PCBs), DDT insecticide and its derivatives, and polycyclic aromatic hydrocarbons (PAHs) (Iwata *et al.*, 1993; Iwata *et al.*, 1994; Dominic, 1994). The spatial distribution, levels and sources of PAHs in the Straits of Malacca are not well known.